Demographic consequences of climate variation along an elevational gradient for a montane

terrestrial salamander

Nicholas M. Caruso[1,2,3]

Leslie J. Rissler[1,4]

[1]Department of Biological Sciences, Box 870345 MHB Hall, University of Alabama,

Tuscaloosa, Alabama 35487, USA

[2]Author for correspondence: carusonm@gmail.com

[3]Current address: Directorate for Biological Sciences, National Science Foundation, 4201 Wilson

Blvd. Arlington, VA 22230, USA

[4]Current address: Department of Fish and Wildlife Conservation, Virginia Polytechnic Institute

and State University College of Natural Resources, Blacksburg, VA 24061

NMC ORCID ID# 000-0002-1059-6907

Abstract

Climate change represents a significant threat to amphibians, which are already imperiled.

However, for many species, the relationship between demographic vital rates (survival and

growth) and climate is unknown, which limits predictive models. Here we describe the life

history variation of *Plethodon montanus* using capture-recapture data over a period of four years,

at five sites along an elevational gradient to determine how survival and growth vary with

temperature, precipitation, and how these relationships vary with elevation. We used a

hierarchical model to estimate asymptotic size and growth rate, and used a spatial Cormack-

Jolly-Seber model to estimate probability of capture and survival, as well as dispersal variance.

Our results show that during the active season, growth and survival rates are both positively

affected by precipitation, while survival was positively affected by temperature at all elevations,

the relationship between growth rates and temperature varied along the elevational gradient.

Generally at lower elevations, higher temperatures led to a decrease in growth while at higher

elevations the opposite was true. During the inactive season we found elevational variation in the

relationship between survival and the amount of snow; at low elevations snowfall was low but

survival decreased with increasing snowfall while at higher elevations increasing snowfall lead

to higher survival. Our results demonstrate that understanding how the environment can affect

salamander demography to develop mechanistic models, will require knowledge of the actual

environmental conditions experienced by a given population as well as an understanding of the

overall differences in climate at a given site.

Key words: Capture-Recapture; Growth; Life history; MCMC; *Plethodon*; Survival

Introduction

Amphibians are one of the most endangered vertebrate taxa (McCallum 2007; Hoffman et al. 2010; IUCN 2016) and face multiple onslaughts including emerging infectious diseases, habitat loss, invasive species, and climate change (Hoffman et al. 2010; Blaustein et al. 2011; Grant et al. 2016). Currently, at least 41% of the approximately 6,500 recognized amphibian species are considered threatened (Hoffman et al. 2010; IUCN 2016) and at least 50% of all salamander species are currently listed as "critically endangered", "endangered", or "vulnerable" (IUCN 2016). Trends in declining salamander populations have recently become both taxonomically and geographically widespread (e.g., Rovito et al. 2009; Adams et al. 2013; Spitzen-van der Sluijs et al. 2013). These declines are especially concerning because salamanders represent a significant portion of the total forest biomass and function as keystone predators (Burton and Likens 1975; Milanovich and Peterman 2016).

Given that many populations are already experiencing declines, future changes in climate, represent a compounding threat to amphibian populations (Milanovich et al. 2010; Sutton et al. 2015; Caruso et al. 2017). Recent evidence suggests that contemporary changes in climate have already affected amphibian life history traits (e.g., Reading 2007; Caruso et al. 2014; but see Connette et al. 2015). In addition, warmer temperatures result in metabolic depression (Catenazzi 2016) and slower growth rates of salamanders (Muñoz et al. 2016), which can negatively affect fitness. Under future climate change, populations may become further isolated to higher, cooler elevations (Bernardo and Spotila 2006; Bernardo et al. 2007; Gifford and Kozak 2012; Lyons et al. 2016). Current model predictions of how changes in climate may affect salamander distributions are generally limited to correlative models (e.g., Milanovich et al. 2010; Sutton et al. 2015; Caruso et al. 2017), which do not take into account metrics of

demographic vital rates (i.e., survival, growth, and reproduction) as they are lacking for many

species. Therefore, current models likely underestimate the effects of future changes in climate

(Buckley et al. 2010, Urban et al. 2016).

Demographic vital rates can vary across spatial gradients, and these rates are driven by

the biotic (e.g., competition) and the abiotic (e.g., temperature) environment. Lower quality

environmental conditions can limit a species' distribution, while higher quality environments

allow for persistence (Gaston 2003). In general, pole-ward range limits are thought to be set

primarily by abiotic factors and equator-ward limits by biotic interactions (Schemske et al.

2009). Contrastingly, correlative niche models suggest that amphibian ranges may be more

limited at the warmer range edges by the abiotic environment (Cunningham et al. 2016).

Although data for montane salamander species are sparse, physiological constraints (Bernardo

and Spotila 2006; Gifford and Kozak 2012; Lyons et al. 2016) and results of reciprocal transplant

experiments (Cunningham et al. 2009; Caruso et al. 2017) support this trend. However, detailed

sampling of vital rates across multiple populations distributed across a species' range is time-

consuming and labor-intensive; therefore, few studies on salamanders have used this information

to inform models of range limits and shifts.

As global climates continue to shift, demographic vital rates have become increasingly

important to characterize the health of natural populations and to develop informed population

models (Pauly 1995; Caswell 2000; Sarrazin and Legendre 2000; Tenhumberg et al. 2004;

Coulson et al. 2005; Buckley et al. 2010; Urban et al. 2016). Unfortunately, vital rates and life

history traits are unknown for many plethodontid species, adding further uncertainty to their

potentially bleak future (e.g., Milanovich et al. 2010). Even when such studies are done,

sampling biases such as unobservable ecological states, imperfect and variable detection, and

measurement error can distort vital rate estimates of natural populations (Leberg et al. 1989;

Royle and Dorazio 2008; Schwarz and Runge 2009; Eaton and Link 2011; Kéry and Schaub

2012; Connette and Semlitsch 2015; Connette et al. 2015; Kéry and Royle 2016). Capture-

recapture (CR) methods offer a solution for accounting for these biases; individual observable

ecological states can be tracked, while uncertainty in these states can be modeled (Kéry and

Schaub 2012; Kéry and Royle 2016). Survival is often a focus of CR studies, as understanding

survival, its variation (both temporal and spatial), and the abiotic and biotic factors that drive this

variation, are necessary to understanding the underlying spatial and temporal variation in

population growth (Lebreton et al. 1992; Saether and Bakke 2000). Similarly, growth is useful

for understanding population demographics since larger body size in many species, especially

amphibians, is associated with higher fitness (e.g., Petranka 1998). Both survival and growth

estimates can be improved using CR methods. Survival can be modeled by accounting for

capture probability and dispersal (Lebreton et al. 1992; Schaub et al. 2004; Schaub and Royle

2014), while estimating measurement error and the variation in growth within and among

individuals can improve growth estimates (Eaton and Link 2011; Link and Hesed 2015).

In this study, we collected four years of capture-recapture data for *Plethodon montanus* at

five sites along an elevational gradient and used hierarchical models to explore the relationship

between demography and climate. The objectives of our study were to 1) determine how

demographic vital rates (growth and survival) vary along an elevational gradient and among

seasons, 2) determine how environmental conditions (temperature and precipitation) affect

variation in growth and survival, and 3) determine how the relationship between both survival

and growth and environmental conditions varies along the elevational gradient and among

seasons. We hypothesized that *P. montanus* vital rates would be driven by climate, whereby

warmer and drier conditions would decrease both survival and growth and that lower elevation

populations, by virtue of being warmer and drier, will show reduced survival and growth

compared to those at higher elevations.

Methods

Locations and Sampling

We established five sites along an elevational gradient within the range of *P. montanus* in

Pisgah National Forest in 2013: SPG (Spivey Gap; 996m), IMG (Iron Mountain Gap; 1,134m),

HG, (Hughes Gap; 1,231m), BBT (Big Butt Trail; 1,300m), and CG (Carver's Gap; 1,464m).

These sites were chosen to minimize the differences among sites in leaf litter depth, aspect,

canopy coverage, and number of surface retreats while establishing an elevational gradient

within distribution of *P. montanus*.

Within each site, we delineated one-150 m^2 plot (10 x 15 m). Starting in 2014, we

established a grid (25-2 x 3 m sections) within the plot to determine the location of each

individual salamander within 0.5 m. Although surveys differed in number of people, effort, and

type (i.e., diurnal and nocturnal), salamanders were processed similarly regardless of survey

type, amount of effort, or number of surveyors. We captured all salamanders encountered and

measured their body size from the snout to the posterior margin of the vent (SVL). We marked

P. montanus using Visual Implant Elastomer (VIE; Northwest Technology Inc., Shaw Island,

Washington) tags, which have minimal effects on fitness and low incidence of tag loss (Bailey

2004). After all salamanders were processed on a particular sampling occasion, we released all

individuals back to the original point of capture. See appendix A for additional detail about

sampling and site characteristics.

Site- and Survey-Specific Climate

First, we defined two seasons based on our sampling; the active season (i.e., when

salamanders were typically active and when we conducted surveys – 27 May to 13 October), and

the inactive season (14 Oct to 26 May). Next, for the length of our study, we obtained daily

temperature and precipitation data from the DAYMET database (http://www.daymet.org;

Thornton et al. 1997) as covariates in our growth and survival models. We defined temperature

for both the active and inactive season as the mean maximum temperature (^{o}C) during the

sampling interval, while we defined precipitation for the active season as mean precipitation

(mm) and for the inactive season as the mean snow water equivalent (mm) during the sampling

interval.

Asymptotic Size and Growth

To test the degree to which growth is driven by climate, we used a hierarchical model to

estimate the effects of season, elevation, temperature, and precipitation on growth rates. Our

model estimates the expected size of the ith individual at the tth time (ES_{it}) using the von

Bertalanffy growth curve, parameterized for unknown ages (Fabens 1965) as a function of its

expected size at the previous measurement time (TS_{it-1}), elevation-specific asymptotic size (a_{elev}),

active [$k(A)_{it}$] and inactive [$k(IA)_{it}$] season growth rates scaled for 365 day increments, and the

interval between captures (number of days) during the active [$\Delta t(A)_{it}$] or inactive [$\Delta t(IA)_{it}$]

seasons (Equation 1).

$$ES_{it} = ES_{it-1} + (a_{elev} - ES_{it-1}) \times (1 - e^{-k(a)_{it} \times \frac{\Delta t(A)_{it}}{365} - k(i)_{it} \times \frac{\Delta t(IA)_{it}}{365}})$$ (Eq. 1)

The logarithm of active and inactive season growth rates were subsequently defined by season-

specific intercepts [$\alpha_{k(A)}, \alpha_{k(IA)}$] , as well as covariates for elevation [$\beta elev_{k(A)}, \beta elev_{k(IA)}$],

temperature [$\beta temp_{k(A)}, \beta temp_{k(IA)}$], precipitation [$\beta precip_{k(A)}$], SWE [$\beta swe_{k(IA)}$] and the

interaction between elevation and temperature [$\beta et_{k(A)}, \beta et_{k(IA)}$], precipitation [$\beta ep_{k(A)}$], or

SWE [$\beta es_{k(IA)}$; Equation 2].

157
$$\log(k(A)_{it}) = \alpha_{k(A)} + \beta elev_{k(A)} + \beta temp_{k(A)} + \beta precip_{k(A)} + \beta et_{k(A)} + \beta ep_{k(A)}$$
$$\log(k(IA)_{it}) = \alpha_{k(IA)} + \beta elev_{k(IA)} + \beta temp_{k(IA)} + \beta swe_{k(IA)} + \beta et_{k(IA)} + \beta es_{k(IA)}$$
(Eq. 2)

Lastly, our measurements of a given individual ($SVL_{i,t}$) are described by independent normal

random variables with a mean of the expected size ($ES_{i,t}$) and an estimated standard deviation

(σ_{SVL}; i.e., measurement error). Therefore, using this hierarchical model we estimated 18

parameters. For further model details and code, see appendix B.

Bayesian Growth Analysis

To evaluate our growth models, we assigned vague normal priors (mean = 0; variance =

100) to all growth rate covariates, uniform priors for elevation specific asymptotic size (min =

50, max = 80), and a vague Gamma prior (shape and rate = 0.001) to the parameter $\dfrac{1}{\sigma^2_{SVL}}$ (i.e.,

precision). All continuous covariates were first scaled and centered prior to model fitting. We fit

the model using Markov chain Monte Carlo (MCMC), generating three chains, each with

400,000 iterations. We used an adaptation phase of 1,000, discarded 250,000 burn-in iterations,

and used a thinning rate of 50, retaining 3,000 iterations from each chain to estimate posterior

distributions (9,000 total samples).

Capture, Dispersal, and Survival

To test the degree to which survival is driven by climate, we used a spatial Cormack-

Jolly-Seber (s-CJS) model (Schaub and Royle 2014) to estimate the effects of season, elevation,

temperature, and precipitation on survival while accounting for variation in capture probability,

and dispersal. For each individual, we modeled survival to each primary period after its initial

capture. Therefore, an individual's ecological state during the primary period where it is first

captured and marked is known (i.e., equal to one). For subsequent primary periods, the

ecological state of the *i*th individual at the *t*th primary period ($z_{i,t}$) is described by a Bernoulli

distribution where the probability of success (i.e., the individual is alive, given that it was alive

previously) is the product of the probability of survival of the *i*th individual to the *t*th primary

period ($\phi_{i,t}$) and the ecological state of the *i*th individual at the previous primary period ($z_{i,t-1}$).

Our observation process is likewise described by a Bernoulli distribution where the probability

of success (i.e., finding the *i*th, individual, at the *t*th primary period, and *tt*th secondary period,

given that it is alive and within the bounds of the study area) is the product of the capture

probability ($p_{i,tt,t}$), ecological state ($z_{i,t}$), and spatial state ($r_{i,tt,t}$) of the *i*th individual, at the *t*th

primary period, and *tt*th secondary period (Schaub and Royle 2014; Kéry and Schaub 2012).

To account for the fact that some individuals emigrated and thus represent apparent

survival (Schaub and Royle 2014), we included each individual's spatial location within each

study site and estimated dispersal from subsequent recaptures. The spatial state ($r_{i,tt,t}$) of the *i*th

individual, at each *tt*th secondary period, and *t*th primary period, is given a value of one if the

location in the x- and y-axes ($Gx_{i,tt,t}$, $Gy_{i,tt,t}$ respectively) of that individual, at that time is within

the study area, while the spatial state receives a value of zero if that individual, at that time is

outside the study area. Because we sampled secondary periods within primary periods (2013-

2014), we first describe the primary period center of activity in the x- and y-axes ($Gx_{i,t}$, $Gy_{i,t}$

respectively). The initial primary period center of activity of the *i*th individual, at the *t*th primary

period is described by a uniform distribution, which is bounded by the lower and upper bounds

of the x- and y-axes of the plot area (i.e., an individual must be within the study area to be

marked). For subsequent primary periods (*t+1*), the center of activity of the *i*th individual is

normally distributed where the mean is the center of activity at the previous primary period in the

x- and y-axes and two estimated precision parameters ($\frac{1}{\sigma^2_{Gxt}}$, $\frac{1}{\sigma^2_{Gyt}}$). Lastly, the spatial location

of the *i*th individual at the *tt*th secondary period and *t*th primary period are also normally

distributed where the mean is the center of activity during the *t*th primary period and two

estimated precision parameters ($\frac{1}{\sigma^2_{Gxtt}}$, $\frac{1}{\sigma^2_{Gytt}}$). We therefore estimated four parameters for the

dispersal portion of our model.

We modeled the logit of the capture probability ($p_{i,tt,t}$) for the *i*th individual, at the *t*th

primary period, and *tt*th secondary period as a function of an intercept (α_p), survey type (diurnal

or nocturnal; $\beta surv_p$), effort (1 or 2; βeff_p), number of people (1 or 2; $\beta pers_p$), linear and

quadratic terms for Julian day ($\beta jday_p$, $\beta jday_p^2$ respectively), site ($\beta site_p$) and random

intercepts for individuals (ε_i) and primary period ($\gamma_{i,t}$; Equation 3).

$\text{logit}(p_{i,tt,t}) = \alpha_p + \beta surv_p + \beta eff_p + \beta pers_p + \beta jday_p + \beta jday_p^2 + \beta site_p + \varepsilon_i + \gamma_{i,t}$ (Eq. 3)

Therefore, for capture probability we estimated twelve total parameters (i.e., ten for the fixed

effects and one for the precision component for each of the two random intercepts). Lastly, we

modeled the logit of survival ($\phi_{i,t}$) of the *i*th individual at the *t*th primary period as a function of

an intercept (α_ϕ), size (last SVL measurement; $\beta size_\phi$), elevation ($\beta elev_\phi$), season ($\beta season_\phi$),

temperature, precipitation, and SWE during the active [$\beta temp_{\phi(A)}$, $\beta precip_{\phi(IA)}$] and inactive [

$\beta temp_{\phi(A)}$, $\beta swe_{\phi(IA)}$] seasons, as well as interactions between elevation with season (βese_ϕ),

active season temperature and precipitation [$\beta et_{\phi(A)}$, $\beta et_{\phi(IA)}$] and inactive temperature and

precipitation and SWE [$\beta ep_{\phi(A)}$, $\beta esw_{\phi(IA)}$; Equation 4].

221
$$\begin{aligned} \text{logit}(\phi_{i,t}) = {} & \alpha_\phi + \beta size_\phi + \beta elev_\phi + \beta season_\phi + \beta temp_{\phi(A)} + \beta temp_{\phi(IA)} + \beta precip_{\phi(A)} \\ & + \beta swe_{\phi(IA)} + \beta ese_\phi + \beta et_{\phi(A)} + \beta et_{\phi(IA)} + \beta ep_{\phi(A)} + \beta esw_{\phi(IA)} \end{aligned}$$
(Eq. 4)

We therefore estimated a total of 13 parameters for survival. For further model details and code,

see appendix B.

Bayesian Survival Analysis

To evaluate our survival models, we first scaled and centered all continuous fixed. We

assigned uniform priors (min = 0, max = 10) to all spatial variance estimates. For fixed

parameters, we assumed vague normal priors (mean = 0; variance = 100), and random intercepts

(γ_t and ε_i) were given normal priors, which had estimated precision parameters from a uniform

distribution (min = 0; max = 10). We fit the model using MCMC, generating three chains, each

with 600,000 iterations. We used an adaptation phase of 1,000, discarded 300,000 burn-in

iterations, and used a thinning rate of 50, retaining 6,000 iterations from each chain to estimate

posterior distributions (18,000 total samples).

All analyses were performed in program R version 3.3.1 (R Core Team 2016). We used

the *jagsUI* package (Kellner 2017) to call JAGS (Plummer 2003), from Program R for MCMC

analyses. We examined traceplots of parameters for adequate mixing among chains and the $\hat{R}$

statistic (Gelman et al. 2004) to evaluate model convergence (see appendix C), and we evaluated

parameter significance based on the overlap of 95% highest posterior density (HPD) with zero.

Results

Over 190 diurnal (n = 58; 31%) and nocturnal (n = 132; 69%) surveys, we captured 2,962

salamanders representing nine species (*P. montanus* = 2,413, 81%; non-target species = 549,

19%). We marked a total of 1,343 individuals, and recapture events constituted 1,070 (44%) of

our total captures of *P. montanus* captures; we recaptured 559 (42%) individuals at least once

(range = 1 – 15 times). Recapture rates generally increased throughout the duration of this study

and average recapture rates were at least 40% during the final year of this study at all sites

(Appendix A7).

Growth

We used animals that were captured at least twice for all growth analyses, which included

1,586 total measurements (544 unique individuals), with a range of 92 – 728 measurements per

site (36 – 215 unique individuals per site). Although the highest elevation site (CG) had the

largest asymptotic size estimate (62.2 mm) this size was similar to the asymptotic size estimate

for IMG (62.1 mm), which is approximately 300 meters lower in elevation, while the remaining

three sites had smaller asymptotic sizes (Fig. 1A). During the active season, we found significant

effects of average precipitation ($\beta precip_{k(A)}$; 95% HPD = 0.15 – 0.45) and the interaction

between average temperature and elevation ($\beta et_{k(A)}$; 95% HPD = 0.10 – 0.32; Fig 1B). At all

sites, increased precipitation was associated with higher growth rates at all sites, whereas the

relationship between temperature and growth rate varied along the elevational gradient; higher

temperatures at lower elevations resulted in lower growth rates while the opposite was observed

at higher elevations (Fig 2A). Average growth rates during the inactive season were considerably

lower ($\alpha_{k(IA)}$; 95% HPD = -28.37 – -6.26) than those during the active season ($\alpha_{k(A)}$; 95% HPD

= 0.11 – 0.37) and inactive season growth rates were not significantly influenced by climate (Fig

1C).

Dispersal and Capture

Dispersal variance estimates from s-CJS models were similar for primary and secondary

seasons with values of approximately one in both the x- and y-axes (Fig. 3A). Therefore, we

would expect that 95% of an individual's movements in the x- and y-axes would be found within

~2 meters from their previous point of capture. Individual variation in capture probability (σ_ε)

was lower (95% HPD = 0.01 – 0.20) compared to primary period variation (σ_γ; HPD 95% = 0.67

– 1.04) and we found that survey type ($\beta surv_p$; 95% HPD = 2.01 – 2.42), the amount of effort (

βeff_p; 95% HPD = 0.81 – 1.95), the number of people ($\beta pers_p$; 95% HPD = 0.41 – 1.18)

significantly explained capture probability (Fig. 3B). Capture probability was higher during

nocturnal surveys and those that had two people and an increased effort (Fig. 3B).

Survival

We found that survival was significantly affect by size ($\beta size_\phi$; 95% HPD = 1.22 – 1.56),

elevation ($\beta elev_\phi$; 95% HPD = -0.58 – -0.02), and season ($\beta season_\phi$; 95% HPD = -2.48 – -1.25;

Fig. 3C). Survival increased with increased SVL, was lower at high elevations compared to low

elevations, and was higher during the active season compared to the inactive season (Figs. 3C,

4). During the active season both temperature ($\beta temp_{\phi(A)}$; 95% HPD = 0.36 – 0.94) and

precipitation ($\beta precip_{\phi(A)}$; 95% HPD = 0.21 – 0.97) positively influenced survival, increased

precipitation and higher temperatures were associated with higher survival (Fig. 4A, B).

However, during the inactive season we found that survival was significantly influenced by the

interaction between average SWE and elevation ($\beta esw_{\phi(IA)}$; 95% HPD = 0.22 – 2.51; Fig. 3C).

At higher elevations, a greater amount of SWE was associated with higher survival while the

opposite was observed at lower elevations (Fig. 4C).

Discussion

We present four years of capture-recapture data to provide a detailed account of how

survival and growth of *Plethodon montanus* varies along an elevational gradient and with

climate. We found that survival and growth of *P. montanus* is influenced by climate and

importantly that the relationship between either survival or growth and climate can vary along an

elevational gradient. Our results suggest that *P. montanus* exhibits variation in life history along

this elevational gradient, likely resulting from the differences in abiotic environment experienced

by those populations. Therefore, understanding how the environment can affect salamander

populations, via survival and growth of individuals, may require more than just knowledge of the

actual environmental conditions experienced by a given population but also understanding the

overall differences in climate at a given site.

Dispersal

Previous studies have found small home ranges and low dispersal in other species of

terrestrial plethodontids (Kleeberger and Werner 1982; Mathis 1991; Merchant 1972; Muñoz et

al. 2016). Our observations for *P. montanus* are consistent with these patterns. Dispersal and

subsequent immigration can buffer sink populations from declines even when climate negatively

affects demographic vital rates and population growth (Brown and Kodric-Brown 1977; Dias

1996; Pulliam 1988; Tavecchia et al. 2016). Unfortunately, montane salamanders, like *P.*

montanus, are also physiologically restricted at lower elevations, and tracking suitable climate

would likely be limited through warmer valleys and across latitudes, which may exacerbate

population isolation and range contractions (Bernardo and Spotila 2006; Kozak and Wiens

2006). The low dispersal observed for *P. montanus* and other terrestrial plethodontids (Cabe et

al. 2007; Liebgold et al. 2011; Marsh et al. 2004; Ousterhout and Liebgold 2010; Peterman and

Semlitsch 2013) likely further increases their risk of population decline under future climate

change.

Active Season

We found that increased precipitation is associated with both higher growth rates across

the elevational gradient (Fig 2B). It has been well-established that rainfall can influence surface

activity of plethodontid salamanders (Petranka and Murray 2001; Bendik and Gluesenkamp

2013; Connette et al. 2015), which require moist conditions due to their high rates of evaporative

water loss (Spight 1968; Spotila 1972; Spotila and Berman 1976). Conditions, such as increased

precipitation, allow for a greater surface activity window and subsequently increased foraging

time (Feder and Londos 1984) or may increase the encounter rates of salamanders with their prey

(Jaeger 1978, 1980); both of which would likely increase the total number of prey captured and

would lead to increases in growth rates, assuming other environmental conditions are equal (e.g.,

temperature; Catenazzi 2016). Higher precipitation was also associated with an increase in

survival across the elevational gradient (Fig 4B). Direct mortality from reduced rainfall is

unlikely as salamanders are able to find and take advantage of moist microhabitats in seemingly

unfavorable habitats (e.g., Yanev and Wake 1981); however, dehydration can cause a reduction

in locomotor performance (Feder and Londos 1984), which may in turn increase predation and

decrease survival.

Temperature-sensitivity in various aspects of growth rates (e.g., rates of assimilation and

digestion) has been previously demonstrated in plethodontids (e.g., Fitzpatrick 1973; Bobka et al.

1981; Muñoz et al. 2016; Clay and Gifford 2017). In our study, the effect of temperature on

growth rates of *P. montanus* varied along the elevational gradient; higher temperatures resulted

in lower growth rates at low elevations and higher growth rates at high elevations (Fig. 2A). This

pattern may be explained, at least in part, by the range of average temperatures experienced by

these populations; high elevation populations generally experienced temperatures lower than

those experienced by low elevations (Appendix A4; Fig. 2A). Indeed, Clay and Gifford (2017)

found a similar pattern, in which energy assimilation under laboratory conditions in *P. montanus*

steadily increased with increasing temperatures but rapidly dropped after reaching an optimum

(22.8^oC for high elevations and 22.6^oC for low elevations) similar to the average temperature at

which the growth rate temperature relationship in our study (~$23.5\ ^o$C) changed from positive

(lower temperatures) to negative (higher temperatures; Fig. 2A). Survival, on the other hand, was

affected by temperature similarly across our elevational gradient; higher temperatures were

associated with higher rates of survival (Fig. 4A). While high temperatures can invariably lead to

death in plethodontids, temperatures experienced by our populations never reached higher than

30^oC, which is lower than the critical thermal maximum of similar species (Spotila 1972).

Increased temperature can have a positive influence on terrestrial salamander endurance and

speed (Else and Bennett 1987; Johnson et al. 2010), which may allow them to more readily

escape predation. Moreover, under extended warmer conditions, plethodontids may reduce

surface activity (Spotila 1972), which would result in a reduction in encounters with potential

surface predators. If higher temperatures increased survival via a reduction in surface activity,

we would also expect populations to show a reduction in growth rates due to reduced foraging

opportunities, which is supported in our growth analyses (Fig. 2A).

Inactive Season

Our results suggest a disparity in both growth and survival along elevations during the

inactive season compared to the active season (Figs. 1 and 3). Low growth rates during the

inactive season is likely due to the extended periods of inactivity and lower prey availability,

although salamanders have been found on the surface during the winter when temperatures are

above freezing (e.g., Catenazzi 2016). Reduced food intake due to the decreased foraging time

and/or lower prey availability may also explain the lower survival that we observed during the

inactive season compared to the active season. Salamanders that are active during warmer winter

conditions may not be able to find the necessary food sources to compensate for this increased

activity, leading to decreased body condition and lower survival (Reading 2007; Sinclair et al.

2013; Catenazzi 2016).

During the inactive season, we found that the interaction between SWE and elevation was

a significant predictor of survival for *P. montanus* (Fig. 4C). Increased SWE lead to an increase

in survival at higher elevations but a decrease in survival at lower elevations; although SWE at

the lowest elevations was low and was less variable (Fig. 4C). We suggest two mechanisms by

which increases in the amount of snow at higher elevations could increase survival. First,

salamanders may have increased survival in areas with more snowpack throughout the winter

because of their reduced surface activity, which leads to a reduction in the number of encounters

with surface predators (Turbill et al. 2011). Alternatively survival may increase with an increase

in snowfall because snow acts as a soil insulator (Decker et al. 2003) and less snowpack can lead

to more variable and colder soil temperatures (Groffman et al. 2001; Henry 2008; Bale and

Hayward 2010; Brown and DeGaetano 2011). Therefore, hibernating salamanders in areas with

more snowpack would have a greater buffer from subzero temperatures (Decker et al. 2003).

Importantly, predictions of salamander population growth under future climate change

that only account for responses to the active season conditions may underestimate losses.

Warming is predicted to be unequal among seasons, winter months will likely see a greater

increase in temperatures than the other seasons (Xia et al. 2014). For logistic reasons, studies of

terrestrial plethodontids have typically focused on the active season (i.e., when individuals are

available for capture). Therefore, determining the effect of winter conditions on salamander

demography (e.g., survival) through the experimental manipulation of temperature or snowpack

would improve mechanistic predictive population models (Sanders-DeMott and Templer 2017).

Conclusions

Future warming is predicted to be a major challenge for Appalachian salamanders

(Milanovich et al. 2010; Sutton et al. 2015; Catenazzi 2016; Caruso et al. 2017). Yet mechanistic

population growth models are lacking, due, in part, to the paucity of demographic data for many

species. Through surveying multiple populations along an elevational gradient, this study was

not only able to estimate survival and growth and their variation with relevant climatic factors,

but also we were able to demonstrate that the relationships between salamander demography and

climate also vary with elevations. This spatial variation in demographic vital rates, and their

relationship with relevant climatic factors, is necessary to model population growth and develop

conservation strategies (Caswell 2001; Buckley et al. 2010; McLean et al. 2016; Urban et al.

2016). Furthermore, our study demonstrates the importance of climate variation in life history

strategies in *P. montanus* likely resulting from both the overall environmental differences in our

elevational gradient as well as the variation in climate experience by these populations over the

course of this study.

Acknowledgements

Thanks to C. Houser, J. Jacobs, and E. Kabay for assistance in the field and to D. Adams, C.

Staudhammer, G. Starr, S. Wiesner, S. Kunwor, S. George, and two anonymous reviewers for

comments that improved this manuscript. This research was funded through Graduate Research

Fellowship, E.O. Wilson Fellowship, and the Herpetologists' League E.E. Williams Research

Grant awarded to NMC.

References

Adams MJ, Miller DAW, Muths E, Corn PS, Grant EHC, Bailey LL, Fellers GM, Fisher RN,

Sadinski WJ, Waddle H, Walls SC (2013) Trends in amphibian occupancy in the United

States. PLoS ONE 8:e64347. doi:10.1371/journal.pone.0064347.

Bailey LL (2004) Evaluating elastomer marking and photo identification methods for terrestrial

salamanders; marking effects and observer bias. Herpetological Review 35:38-41.

Bale JS, Hayward SAL (2010) Insect overwintering in a changing climate. Journal of

Experimental Biology 213:980-994.

Bendik NF, Gluesenkamp AG (2013) Body length shrinkage in an endangered amphibian is

associated with drought. Journal of Zoology 290:35-41.

Bernardo J, Spotila JR (2006) Physiological constraints on organismal response to global

warming: mechanistic insights from clinally varying populations and implications for

assessing endangerment. Biology Letters 2:135-139.

Bernardo J, Ossola RJ, Spotila J, Crandall KA (2007) Interspecific physiological variation as a

tool for cross-species assessments of global warming-induced endangerment: validation

of an intrinsic determinant of macroecological and phylogeographic structure. Biology

Letters 3:695-698.

Blaustein AR, Walls SC, Bancroft BA, Lawler JJ, Searle CL, Gervasi SS (2010) Direct and

indirect effects of climate change on amphibian populations. Diversity 2:281-313.

Bobka MS, Jaeger RG, McNaught DC (1981) Temperature dependent assimilation efficiencies

of two species of terrestrial salamanders. Copeia 1981:417-421.

Brown PJ, DeGaetano AT (2011) A paradox of cooling winter soil surface temperatures in a

warming northeastern United States. Agricultural and Forest Meterology 151:947-956.

Brown JH, Kodric-Brown A (1977) Turnover rates in insular biogeography: effect on

immigration on extinction. Ecology 58:445-449.

Buckley LB, Urban MC, Angilletta MJ, Crozier LG, Rissler LJ, Sears MW (2010) Can

mechanism inform species' distribution models? Ecology Letters 13:1041-1054.

Burton TM, Likens GE (1975) Energy flow and nutrient cycling in salamander populations in the

Hubbard Brook Experimental Forest, New Hampshire. Ecology 56:1068-1080.

Cabe PR, Page RB, Hanlon TJ, Aldrich ME, Marsh DM (2007) Fine-scale population

differentiation and gene flow in a terrestrial salamander (*Plethodon cinereus*) living in

continuous habitat. Heredity 98:53-60.

Caruso NM, Jacobs JF, Rissler LJ (2017). An experimental approach to understanding elevation

limits in a montane terrestrial salamander, *Plethodon montanus*. Submitted to Copeia

bioRxiv doi: https://doi.org/10.1101/131573

Caruso NM, Sears MW, Adams DC, Lips KR (2014) Widespread rapid reductions in body size

of adult salamanders in response to climate change. Global Change Biology 20:1751-

1759.

Caswell H (2000) Prospective and retrospective perturbation analyses: their roles in conservation

biology. Ecology 81:619-627.

Caswell H (2001) Matrix Population Models: Construction, Analysis, and Interpretation. 2nd

edn. Sinauer Associates, Sunderland

Catenazzi A (2016) Ecological implications of metabolic compensation at low temperatures in

salamanders. PeerJ 14:e2072 10.7717/peerj.2072.

Clay TA, Gifford ME (2017) Population level differences in thermal sensitivity of energy

assimilation in terrestrial salamanders. Journal of Thermal Biology 64:1-6.

Connette GM, Crawford JA, Peterman WE (2015) Climate change and shrinking salamanders:

alternative mechanisms for changes in plethodontid salamander body size. Global

Change Biology 21:2834-2843.

Connette GM, Semlitsch RD (2015) A multistate mark-recapture approach to estimating survival

of PIT-tagged salamanders following timber harvest. Journal of Applied Ecology

52:1316-1324.

Coulson T, Gaillard J-M, Festa-Bianchet M (2005) Decomposing the variation in population

growth into contributions from multiple demographic rates. Journal of Animal Ecology

74:789-801.

Cunningham HR, Rissler LJ, Apodaca JJ (2009) Competition at the range boundary in the slimy

salamander: using reciprocal transplants for studies on the role of biotic interactions in

spatial distributions. Journal of Animal Ecology 78:52-62.

Cunningham HR, Rissler LJ, Buckley LB, Urban MC (2016) Abiotic and biotic constraints

across reptile and amphibian ranges. Ecography 39:1-8.

Decker KL, Wang MD, Waite C, Scherbatskoy T (2003) Snow removal and ambient air

temperature effects on forest soil temperatures in northern Vermont. Soil Science Society

of America Journal 67:1234-1243.

Dias PC (1996) Sources and sinks in population biology. Trends in Ecology and Evolution

11:326-330.

Eaton MJ, Link WA (2011) Estimating age from recapture data: integrating incremental growth

measures with ancillary data to infer age-at-length. Ecological Applications 21:2487-

2497.

Else, PL Bennett AF (1987) The thermal-dependence of locomotor performance and muscle

contractile function in the salamander *Ambystoma tigrinum nebulosum*. Journal of

Experimental Biology 128:219-233.

Fabens AJ (1965) Properties and fitting of von Bertalanffy growth curve. Growth 29:265-289.

Feder ME, Londos PL (1984) Hydric constraints upon foraging in a terrestrial salamander

Desmognathus ochrophaeus (Amphibia: Plethodontidae). Oecologia 64:413-418.

Fitzpatrick LC (1973) Influence of seasonal temperatures on the energy budget and metabolic

rates of the northern two-lined salamander *Eurycea bislineata bislineata*. Comparative

Biochemistry and Physiology 45A:807-818.

Gaston KJ (2003) The Structure and Dynamics of Geographic Ranges. Oxford University Press.

Gelman A, Carlin JB, Stern HS, Rubin DB (2004) Bayesian data analysis, 2nd edn.

CRC/Capman and Hall, Boca Raton, FL.

Gifford ME, Kozak KH (2012) Islands in the sky or squeezed at the top? Ecological causes of

elevational range limits in montane salamanders. Ecography 35:193-203.

Grant EHC, Miller DAW, Schmidt BR, Adams MJ, Amburgey SM, Chambert T, Cruickshank

SS, Fisher RN, Green DM, Hossack BR, Johnson PTJ, Joseph MB, Rittenhouse TAG,

Ryan ME, Waddle JH, Walls SC, Bailey LL, Fellers GM, Gorman TA, Ray AM, Pilliod

DS, Price SJ, Saenz D, Sadinski W, Muths E (2016) Quantitative evidence for the effects

of multiple drivers on continental-scale amphibian declines. Scientific Reports 6:25625

doi:10. 1038/srep25625.

Groffman PM, Rustad LE, Templer PH, Campbell JL, Christenson LM, Lany N, Socci AM,

Vadebncoeur MA, Schaberg PG, Wilson G, Driscoll C, Fahey TJ, Fisk MC, Goodale CL,

Green MB, Hamburg SP, Johnson CE, Mitchell MJ, Morse JL, Pardo LH, Rodenhouse

NL (2012) Long-term integrated studies show complex and surprising effects of climate

change in the northern hardwood forest. BioScience 62:1056-1066.

Henry HAL (2008) Climate change and soil freezing dynamics: historical trends and projected

changes. Climatic Change 87:421-434.

Hoffman M, et al (2010) The impact of conservation on the status of the world's vertebrates.

Science 330:1503-1509.

IUCN (2016) The IUCN red list of threatened species. Version 2016-3. <www.iucnredlist.org>.

Downloaded on 22 March 2017.

Jaeger RG (1978) Plant climbing by salamanders: periodic availability of plant-dwelling prey.

Copeia 1978:686-691.

Jaeger RG (1980) Fluctuations in prey availability and food limitation for a terrestrial

salamander. Oecologia 44:335-341.

Johnson JR, Johnson BB, Shaffer HB (2010) Genotype and temperature affect locomotor

performance in a tiger salamander hybrid swarm. Functional Ecology 24:1073-1080.

Kéry M, Royle JA (2016) Applied hierarchical modeling in ecology: analysis of distribution,

abundance and species richness in R and BUGS (volume 1 – prelude and static models).

Academic Press, San Diego, CA.

Kéry M, Schaub M (2012) Bayesian population analysis using WinBUGS – a hierarchical

perspective. Academic Press, San Diego, CA.

Kellner K (2017) jagsUI: a wrapper around 'rjags' to streamline 'JAGS' analyses. R package

version 1.4.9.

Kleeberger SR, Werner JK (1982) Home range and homing behavior of *Plethodon cinereus* in

Northern Michigan. Copeia 1982:409-415.

525 Kozak KH, Wiens JJ (2006) Does niche conservatism drive speciation? A case study in North

526   American salamanders. Evolution 60:2585-2603.

527 Leberg PL, Brisbin IL, Smith MH, White GC (1989) Factors affecting the analysis of growth-

528   patterns of large mammals. Journal of Mammalogy 70:275-283.

529 Lebreton J-D, Burnham KP, Clobert J, Anderson DR (1992) Modeling survival and testing

530   biological hypotheses using marked animals: a unified approach with case studies.

531   Ecological Monographys 62:67-118.

532 Liebgold EB, Brodie III ED, Cabe PR (2011) Female philopatry and male-biased dispersal in a

533   direct-developing salamander, *Plethodon cinereus*. Molecular Ecology 20:249-257.

534 Link WA, Hesed KM (2015) Individual heterogeneity in growth and age at sexual maturity: a

535   gamma process analysis of capture-mark-recapture data. Journal of Agricultural,

536   Biological, and Environmental Statistics 20:343-352.

537 Lyons MP, Shepard DB, Kozak KH (2016) Determinants of range limits in montane woodland

538   salamanders (genus *Plethodon*). Copeia 104:101-110.

539 Marsh DM, Thakur KA, Bulka KC, Clarke LB (2004) Dispersal and colonization through open

540   fields by a terrestrial, woodland salamander. Ecology 85:3396-3405.

541 Mathis A (1991) Territories of male and female terrestrial salamanders: costs, benefits, and

542   intersexual spatial associations. Oecologia 86:433-440.

543 McLean N, Lawson CR, Leech DI, van de Pol M (2016) Predicting when climate-driven

544   phenotypic change affects population dynamics. Ecology Letters 19:595-608.

545 McCallum ML (2007) Amphibian decline or extinction? Current declines dwarf background

546   extinction rate. Journal of Herpetology 41:483-491.

Merchant H (1972) Estimated population size and home range of the salamanders *Plethodon*

jordani and *Plethodon glutinosus*. Journal of the Washington Academy of Sciences

62:248-257.

Milanovich JR, Peterman WE (2016) Revisiting Burton and Likens (1975): nutrient standing

stock and biomass of a terrestrial salamander in the Midwestern United States. Copeia

104:165-171.

Milanovich JR, Peterman WE, Nibbelink NP, Maerz JC (2010) Projected loss of a salamander

diversity hotspot as a consequence of projected global climate change. PLoS ONE

5:e12189 DOI 10.1371/journal.pone.0012189.

Muñoz DJ, Hesed KM, Grant EHC, Miller DAW (2016) Evaluating within-population variability

in behavior and demography for the adaptive potential of a dispersal-limited species to

climate change. Ecology and Evolution 6:8740-8755.

Ousterhout B, Liebgold EB (2010) Dispersal versus site tenacity of adult and juvenile red-backed

salamanders (*Plethodon cinereus*). Herpetologica 63:269-275.

Pauly D (1995) Anecdotes and shifting baseline syndrome of fisheries. Trends in Ecology and

Evolution 10:430. doi:10.1016/S0169-5347(00)89171-5

Peterman WE, Semlitsch RD (2013) Fine-scale habitat associations of a terrestrial salamander:

the role of environmental gradients and implications for population dynamics. PLoS ONE

8:e62184. doi:10.1371/journal.pone.0062184.

Petranka JW (1998) Salamanders of the United States and Canada. Smithsonian Institution Press,

Washington, DC.

Petranka JW, Murray SS (2001) Effectiveness of removal sampling for determining salamander

density and biomass: a case study in an Appalachian streamside community. Journal of

Herpetology 35:36-44.

Plummer M (2003) JAGS: A program for analysis of Bayesian graphical models using Gibbs

sampling. Proceedings of the 3rd International Workshop on Distributed Statistical

Computing (DSC2003), pp 20-22.

Pullimam HR (1988) Sources, sinks, and population regulation. The American Naturalist

132:652-661.

R Core Team (2016) R: a language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Reading CJ (2007) Linking global warming to amphibian declines through its effects on female

body condition and survivorship. Oecologia 151:125-131.

Rovito SM, Parra-Olea G, Vásquez-Almazán CR, Papenfuss TJ, Wake DB (2009) Dramatic

declines in neotropical salamander populations are an important part of the global

amphibian crisis. Proceedings of the National Academy of Sciences USA 106:3231-3236.

Royle JA, Dorazio RM (2008) Hierarchical modeling and inference in ecology. Academic Press,

San Diego, CA.

Saether B-E, Oyvind B (2000) Avian life history variation and contribution of demographic traits

to the population growth rate. Ecology 81:642-653.

Sanders-DeMott R, Templer PH (2017) What about winter? Integrating the missing season into

climate change experiments in seasonally snow covered ecosystems. Methods in Ecology

and Evolution doi: 10.1111/2041-210X.12780

Sarrazin F, Legendre S (2000) Demographic approach to releasing adults versus young in

reintroductions. Conservation Biology 14:488-500.

Schaub M, Gimenez O, Schmidt BR, Pradel R (2004) Estimating survival and temporary

emigration in the multistate capture-recapture framework. Ecology 85:2107-2113.

Schaub M, Royle JA (2014) Estimating true instead of apparent survival using spatial Cormack-

Jolly-Seber models. Methods in Ecology and Evolution 4:1316-1326.

Schemske DW, Mittelbach GG, Cornell HV, Sobel JM, Roy K (2009) Is there a latitudinal

gradient in the importance of biotic interactions? Annual Review of Ecology, Evolution,

and Systematics 40:245-269.

Schwarz LK, Runge MC (2009) Hierarchical Bayesian analysis to incorporate age uncertainty in

growth curve analysis and estimates of age from length: Florida manatee (*Trichechus*

manatus) carcasses. Canadian Journal of Fisheries and Aquatic Sciences 66:1775-1789.

Sinclair BJ, Stinziano JR, Williams CM, MacMillan HA, Marshall KE, Storey KB (2013) Real-

time measurement of metabolic rate during freezing and thawing of the wood frog, *Rana*

sylvatica: implications for overwinter energy use. Journal of Experimental Biology

216:292-302.

Spight TM (1968) The water economy of salamanders: evaporative water loss. Physiological

Zoology 41:195-203.

Spitzen-van der Sluijs AM, Spikmans F, Bosman W, de Zeeuw M, van der Meij T, Goverse E,

Kik M, Pasmans F, Martel A (2013) Rapid enigmatic decline drives the fire salamander

(*Salamandra salamandra*) to the edge of extinction in the Netherlands. Amphibia-

Reptilia 34:233-239.

Spotila JR (1972) Role of temperature and water in the ecology of lungless salamanders.

Ecological Monographs 42:95-125.

Spotila JR, Berman EN (1976) Determination of skin resistance and the role of the skin in

controlling water loss in amphibians and reptiles. Comparative Biochemistry and

Physiology 55A:407-411.

Sutton WB, Barrett K, Moody AT, Loftin CS, deMaynadier PG, Nanjappa P (2015) Predicted

changes in climatic niche and climate refugia of conservation priority salamander species

in the northeastern United States. Forests 6:1-26. doi:10.3390/f6010001.

Tavecchia G, Tenan S, Pradel R, Igual J-M, Genovart M, Oro D (2016) Climate-driven vital

rates do not always mean climate-driven population. Global Change Biology 22:3960-

3966.

Tenhumberg B, Tyre AJ, Shea K, Possingham HP (2004) Linking wild and captive populations

to maximise species persistence: optimal translocation strategies. Conservation Biology

18:1304-1314.

Thornton PE, Running SW, White MA (1997) Generating surfaces of daily meteorological

variables over large regions of complex terrain. Journal of Hydrology 190:214-251.

Turbill C, Bieber C, Ruf T (2011) Hibernation is associated with increased survival and the

evolution of slow life histories among mammals. Proceedings of the Royal Society B:

Biological Sciences 278:3355-3363. doi:10.1098/rspb.2011.0190.

Urban MC, Bocedi G, Hendry AP, Mihoub J-B, Pe'er G, Singer A, Bridle JR, Crozier LG, De

Meester L, Godsoe W, Gonzalez A, Hellmann JJ, Holt RD, Huth A, Johst K, Krug CB,

Leadley PW, Palmer SCF, Pantel JH, Schmitz A, Zollner PA, Travis JMJ (2016)

634   Improving the forecast for biodiversity under climate change. Science

635   353**:**aad8466(2016) DOI: 10.1126/science.aad8466.

636 Xia J, Chen J, Piao S, Ciais P, Luo Y, Wan S (2014) Terrestrial carbon cycle affected by non-

637   uniform climate warming. Nature Geoscience 7:173-180.

638 Yanev KP, Wake DB (1981) Genic differentiation in a relict desert salamander, *Batrachoseps*

639   *campi*. Herpetologica 37:16-28.

Fig. 1 Highest posterior density for growth model with asymptotic size (**a**), active season

parameters and measurement error (**b**), and inactive season parameters (**c**). Points represent mean

estimates; thick lines show 90% HPD while thin lines show 95% HPD. Colors denote the

proportion of the posterior sample that has the same sign as the mean estimate (f).

Fig. 2 Relationship between the logarithm of growth rate with temperature and elevation (**a**), and

precipitation (**b**). Line colors in show site elevation (**a**) while the gray shaded ribbon shows the

95% HPD (**b**). Predicted relationships were limited to the actual range of climate experience by

each site (**a**) or at all sites (**b**).

Fig. 3 Highest posterior density for s-CJS model with dispersal parameters (**a**), capture

parameters (**b**), and survival parameters (**c**). Points represent mean estimates; thick lines show

90% HPD while thin lines show 95% HPD. Colors denote the proportion of the posterior sample

that has the same sign as the mean estimate (f).

Fig. 4 Relationship between survival with temperature (**a**), and precipitation (**b**) and snow water

equivalent (SWE; **c**). In all plots, line colors in show site elevation, while the plot facets

represent the 10, 30, 50, 70, 90% quantiles of SVL (mm) to show the variation in survival over

the range of sizes. Predicted relationships were limited to the actual range of climate experience

by each site (**a, b, c**).