Structure-based analysis of CysZ-mediated cellular uptake of sulfate

Authors and Affiliations:

Zahra Assur Sanghai[1,2], Qun Liu[3], Oliver B. Clarke[2], Edgar Leal Pinto[4], Brian Kloss[5], Shantelle Tabuso[5], James Love[5], Marco Punta[6], Surajit Banerjee[7], Kanagalaghatta R. Rajashankar[7], Burkhard Rost[8], Diomedes Logothetis[4], Matthias Quick[9], Wayne A. Hendrickson[1,2,6] and Filippo Mancia[1*]

[1] Department of Physiology and Cellular Biophysics, [2] Department of Biochemistry and Molecular Biophysics, and [9] Department of Psychiatry, Center for Molecular Recognition, Columbia University, New York, NY 10032, USA.

[3] Biology Department, Brookhaven National Laboratory, Upton, NY 11973, USA.

[4] Department of Physiology and Biophysics, Virginia Commonwealth University School of Medicine, Richmond, VA 23298, USA.

[5] New York Structural Biology Center, New York, NY 10027, USA.

[6] Centre for Evolution and Cancer, The Institute of Cancer Research, London, UK.

[7] Department of Chemistry and Chemical Biology, Cornell University, NE-CAT, Advanced Photon Source, Argonne, IL 60439, USA.

[8] Department of Informatics, Technical University of Munich, Munich 85748, Germany

* To whom correspondence should be addressed: E-mail fm123@cumc.columbia.edu (F.M.)

ABSTRACT

Sulfur, most abundantly found in the environment as sulfate (SO_4^{2-}), is an essential element in metabolites required by all living cells, including amino acids, co-factors and vitamins. Current understanding of the cellular delivery of SO_4^{2-} at the molecular level is limited however. CysZ has been described as a SO_4^{2-} permease, but its sequence family is without known structural precedent. Based on crystallographic structure information, SO_4^{2-} binding and uptake experiments in cells and proteoliposomes, and single-channel conductance measurements, we provide insight into the molecular mechanism of CysZ-mediated translocation of SO_4^{2-} across membranes. CysZ properties differ markedly from those of known transporters and ion channels. The structures display a hitherto unknown fold with dual topology, assembling in CysZ from *Pseudomonas denitrificans* as a trimer of antiparallel dimers in the membrane. CysZ structures from two other species recapitulate dimers from this assembly. Mutational studies highlight the functional relevance of conserved CysZ residues.

Sulfur has a central role in many cellular processes across all kingdoms of life. It is a vital component of several essential compounds, including the sulfur-containing amino acids cysteine and methionine, in prosthetic groups such as the Fe-S clusters, as well as vitamins and micronutrients such as biotin (vitamin H), thiamine (vitamin B1) and lipoic acid, and in coenzymes A and M (Barton, 2005). Whilst mammals obtain the majority of the necessary sulfur-containing metabolites directly from the diet, plants, fungi, and bacteria are able to assimilate and utilize sulfur from organic and inorganic sources (Barton, 2005). Sulfate (SO_4^{2-}) is the most abundant source of sulfur in the environment and its utilization is contingent upon its

entry into the cell (Kertesz, 2000). In certain fungi, and prokaryotes, once internalized, SO_4^{2-} is

first reduced to sulfite (SO_3^{2-}), and then further to sulfide (S^{2-}), a form that can be used by the

cell (Kredich, Hulanicka, & Hallquist, 1979) (Fig. 1-S1). In *Escherichia coli* and other gram-

negative bacteria, the culmination of the aforementioned sulfate assimilatory (also known as

reductive) pathway is the formation of cysteine by the addition of S^{2-} to O-acetylserine by

cysteine synthase, followed by the synthesis of methionine from homocysteine (Kredich, 1971)

(Fig. 1-S1).

In prokaryotes, the entry of SO_4^{2-} into the cell is mediated by four known families of dedicated

transport systems: the ABC sulfate transporter complexes SulT or CysTWA, the SulP family of

putative SLC13 sodium:sulfate or proton:sulfate symporters or SLC26 solute:sulfate exchangers,

the phosphate transporter-like CysP/PitA family, and the CysZ family classified as SO_4^{2-}

permeases (Aguilar-Barajas, Diaz-Perez, Ramirez-Diaz, Riveros-Rosas, & Cervantes, 2011;

Hryniewicz, Sirko, Palucha, Bock, & Hulanicka, 1990; Kertesz, 2001; Loughlin, Shelden,

Tierney, & Howitt, 2002; Mansilla & de Mendoza, 2000; Sirko, Zatyka, Sadowy, & Hulanicka,

1995). CysZ family members are 28-30 kDa bacterial inner-membrane proteins found

exclusively in prokaryotes with no apparent homology to any of the established channel or

transporter folds, and are scarcely studied in the literature (Zhang, Jiang, Nan, Almqvist, &

Huang, 2014). The *cysZ* gene owes its name to its presence in the cysteine biosynthesis regulon.

In two reports from thirty years ago, an *E. coli* K12 strain with a *cysZ* deletion showed a severe

impairment in its ability to accumulate SO_4^{2-} and was not viable in sulfate-free media without an

alternate sulfur source such as thiosulfate ($S_2O_3^{2-}$) (Britton et al., 1983; Parra, Britton, Castle,

Jones-Mortimer, & Kornberg, 1983). Recently, a third report studying the functional properties

of CysZ, concluded that the protein from *E. coli* functions as a high affinity, highly specific pH-

dependent SO_4^{2-} transporter, directly regulated by the toxic, assimilatory pathway intermediate,

SO_3^{2-} (Zhang et al., 2014).

To investigate the role of CysZ in cellular sulfate uptake at a molecular level, we have

undertaken an approach that combines structural and functional studies. To this end, we

determined the crystal structures of CysZ from three species, *Idiomarina loihiensis (Il; IlCysZ)*,

Pseudomonas fragi (Pf; PfCysZ), and *Pseudomonas denitrificans (Pd; PdCysZ)*, and

characterized CysZ function in purified form, in reconstituted proteoliposomes, in planar lipid

bilayers, and in cells. Combining the structural information from the three orthologs reveals that

CysZ features a novel protein fold that assembles as oligomers with a dual topology. This

arrangement can be understood as being derived from trimers of dimers akin to the hexameric

assembly captured in one of the structures. Interpreting the functional data in a structural context

has allowed us to formulate a mechanistic model for CysZ-mediated SO_4^{2-} translocation across

the bacterial cytoplasm membrane.

Both the structures and the functional properties of CysZ proteins are distinct from those of any

known membrane transporter or ion channel. Besides not resembling other transporter structures,

CysZ mediates sulfate uptake into cells or proteoliposomes without coupling to ion gradients,

partner proteins, or exogenous energy sources such as ATP. Besides differing from other ion

channel structures, CysZ generates sulfate currents across lipid bilayers with unusual

characteristics. These distinctive properties make CysZ appealing as a model system for studies

of biophysical principles of membrane protein biogenesis and transmembrane ion passage.

RESULTS

Structure determination of CysZ

Following a structural genomics approach aimed at crystallization for structural analysis, we

cloned and screened a total of 63 different bacterial homologs of CysZ for high-level expression

and stability in detergents (Love et al., 2010; Mancia & Love, 2011). Crystal structures were

determined for CysZ from three organisms. Chronologically, the structure of *Il*CysZ was the first

solved, to 2.3 Å resolution in space group *C2* by SAD, initially based on a single selenate ion

bound to the protein and subsequently also by selenomethione derivatization (SeMet) SAD, and

multi-crystal native SAD (Liu et al., 2012). The structure of *Pf*CysZ was solved second, to 3.5 Å

resolution, with crystals also belonging to space group *C2*. Although *Pf*CysZ and *Il*CysZ share

42% sequence identity, molecular replacement failed to find a convincing solution, and we

instead used SeMet-derivatized *Pf*CysZ to obtain phase information by multi-crystal SeMet

SAD. Third, we determined the structure of *Pd*CysZ, which crystallized in multiple forms

belonging to space groups $P6_3$, $P4_122$ and $P2_12_12_1$, revealing the same architecture and

oligomeric assembly each time (Fig. 1-S2b). *Pd*CysZ structures in the $P6_3$ and $P2_12_12_1$ lattices

each contain an entire hexamer in their asymmetric units, whereas a molecular diad coincides

with a crystallographic axis in the $P4_122$ lattice. We focused our analysis on the best of these

(3.4 Å resolution in $P6_3$). The location of the selenium sites was obtained from a SeMet SAD

data set, and the structure was solved by combining the resulting SAD phases with those from a

molecular replacement solution obtained by positioning the *Pf*CysZ model (75% sequence

identity) onto SeMet fiducials in the initial electron density map (Table 1, Fig. 1-S2a).

The hexameric structure of *Pd*CysZ

The refined structure of *Pd*CysZ comprises an entire hexamer of near-perfect D3 symmetry. Antiparallel pairs of protomers arrange together as a trimer of dimers (Fig. 1a), with the three-fold axis oriented perpendicular to the plane of the putative membrane and three two-fold axes between dimers of the hexamer. Both the periplasmic and cytoplasmic faces of the hexamer are essentially identical by symmetry, resulting in a dual-topology assembly for *Pd*CysZ. The hexamer has a triangular face of equal sides measuring approximately 75 Å, with the perpendicular span of about 65 Å. The interaction of the six protomers results in a total buried surface area (Krissinel & Henrick, 2007) of 5,700 Å^2. A surface electrostatic representation reveals a hydrophobic belt along the mid-section of the hexamer when viewed from its side, outlining the orientation of CysZ in the lipid bilayer (Czodrowski, Dramburg, Sotriffer, & Klebe, 2006; Dolinsky et al., 2007; Dolinsky, Nielsen, McCammon, & Baker, 2004) (Fig. 1b). A surface representation of the sequence conservation, calculated by analysis of multiple sequence alignments (MSA) (Ashkenazy et al., 2016; Glaser et al., 2003) highlights the regions of invariance in the sequence and in turn, the areas on the molecule that are most likely to have structural and functional importance (Fig. 1c).

The CysZ protomer is an alpha-helical integral membrane protein with two long transmembrane (TM) helices (H2b and H3a) and two pairs of shorter helices (H4b-H5a and H7-H8) that insert only partially into the membrane (hemi-penetrating), forming a funnel or tripod-like shape within the membrane (Fig. 2a, b). The protein has an extra-membranous hydrophilic 'head', comprising an iris-like arrangement of the two short helices, H1 and H6, and kinked helices H3b, H4a, and H5b. The amino and carboxyl termini are also located in this region (Fig. 2a, b). H4b and H5a lie

partly inserted in the membrane, with the turn between the two helices pointing in towards the

three-fold axis at the center of the hexamer. CysZ amino-acid sequences from different

organisms are very similar; for example, those of *Pd*CysZ and *E. coli* CysZ (*Ec*CysZ) are 40.5%

identical and those of *Ec*CysZ and our three structures have 30.0% of their residues exactly in

common (Fig. 2c). Helix boundaries are also essentially the same in the structures of *Pd*CysZ,

*Il*CysZ and *Pf*CysZ (Fig. 2c).

The dimeric assembly of *Il*CysZ and *Pf*CysZ

Unlike *Pd*CysZ, both *Il*CysZ and *Pf*CysZ crystallize as dimers (Fig. 3a, b), in agreement with

their different behavior in detergent-containing solution. Indeed, size-exclusion chromatography

runs of the three species of CysZ in the same buffer and detergent conditions, show mono-

disperse peaks eluting at 13.42 ml (*Il*CysZ), 13.95 ml (*Pf*CysZ) and 12.56 ml (*Pd*CysZ) (Fig. 3-

S1). This result is consistent with a different and smaller oligomeric state of *Il*CysZ and *Pf*CysZ

(similar retention volume) compared to *Pd*CysZ (eluting 1 ml ahead).

The protomers of the *Il*CysZ dimer are arranged in a head-to-tail antiparallel association, with

H4b-H5a protruding at an angle that is nearly parallel to the putative plane of the membrane

(Fig. 3a, 3-S2a, b). The *Pf*CysZ dimer is also arranged in an antiparallel orientation but with a

different dimer interface (Fig. 3b). In the *Pf*CysZ structure, H4b-H5a together form a narrower

angle with H2 and H3, and tuck-in closer to the rest of the molecule (Fig. 3b). The resulting

dumbbell-shaped *Pf*CysZ dimer is predicted, by OPM/PPM (Orientation of Proteins in

Membranes) (Lomize, Pogozheva, Joo, Mosberg, & Lomize, 2012), to lie in the membrane at a

31° tilt to the perpendicular, in agreement with the position of its central hydrophobic belt, as

revealed by surface-electrostatics calculations (Fig. 3- S2e, f). The dimer interfaces of *Il*CysZ and

*Pf*CysZ bury 780 Å^2 and 1136 Å^2 of surface area respectively (Krissinel & Henrick, 2007)

The individual protomers of CysZ from all three species adopt the same topology and fold, and

superpose well with an overall pairwise root mean squared deviation (r.m.s.d) of ~2.5 Å (Fig.

3c). The greatest variation between the protomers of each structure is seen in the orientation of

H4b-H5a with respect to the TM helices, H2 and H3 (Fig. 3c), which seem to be the most

conformationally flexible with respect to the rest of the molecule.

Comparison of the structures of *Il*CysZ and *Pf*CysZ with that of *Pd*CysZ revealed that these two

distinctive dimeric structures are both represented in the hexameric one. Indeed, the *Il*CysZ

dimer (Fig. 3a) resembles the vertically arranged pair of protomers in the *Pd*CysZ hexamer (Fig.

3d) at each of the vertices of the triangular structure. On the other hand, the *Pf*CysZ dimer (Fig.

3b) resembles the transverse pair of protomers lying at a tilt, just as was predicted by OPM

(Lomize et al., 2012), along the side of the *Pd*CysZ hexamer when viewed from inside the plane

of the membrane (Fig. 3e). In essence, the *Pd*CysZ hexamer can be seen as a trimer of either

*Il*CysZ or *Pf*CysZ dimers (Fig. 3f).

Dual-topology assembly of CysZ

To validate the dual-topology assembly of CysZ observed in all our structures, we performed

disulfide crosslinking assays on engineered cysteine mutants of CysZ designed to capture the

antiparallel dimer, utilizing isolated membranes. Disulfide-trapping experiments of the

transverse dimer, performed by crosslinking a pair of mutants (L161C-A164C) on H5 of *Pf*CysZ,

as well as the corresponding pair in *Il*CysZ (V157C-Q163C) confirm the dimer interface

observed in the structures of *Pf*CysZ and *Il*CysZ, and, as a consequence, the dual-topology

assembly of the two proteins (Fig. 3-S2f, g, h).

To further validate the dual-topology orientation of CysZ in the membrane, we performed a

cysteine accessibility scan experiment, mapping residues expected to be located outside the lipid

bilayer, by fluorescence labeling of the thiol groups of various single cysteine mutants with a

membrane-impermeable dye. Membrane fractions isolated from recombinant cultures expressing

*Il*CysZ cysteine mutants introduced at positions along the edge of helix H4 predicted to be

solvent accessible based on our structure, were indeed labeled with a membrane-impermeable

fluorescent thiol-specific maleimide dye (Fig. 3-S2c, d).

Functional characterization of CysZ

To characterize the functional properties of CysZ, we used a four-pronged approach: (i)

radiolabeled $[^{35}S]O_4^{2-}$ uptake experiments in cells expressing plasmid-encoded CysZ, (ii)

radiolabeled ($[^{35}S]O_4^{2-}$) binding to purified CysZ in detergent solution, (iii) $[^{35}S]O_4^{2-}$ uptake in

proteoliposomes reconstituted with purified CysZ, and (iv) single-channel electrophysiological

recordings in a planar lipid bilayer reconstituted with CysZ.

First, we compared the time course of SO_4^{2-} accumulation in an *E. coli cysZ* knockout strain (*E.*

coli K12 JW2406-1, CysZ⁻) (Baba et al., 2006) with that in the wild-type (WT, *E. coli* K12

BW25113, *cysZ*⁺) strain. CysZ⁻ cells, after growth in minimal media and 12 hours of sulfate

starvation, showed significantly diminished SO_4^{2-} uptake when compared to the WT strain,

consistent with previous results (Fig. 4a) (Parra et al., 1983). The CysZ⁻ strain still showed a low

level of sulfate accumulation, which could be attributed to the other endogenous sulfate transport

systems present in the bacteria (for example ABC transporter and SulP). This uptake-deficient

phenotype of the CysZ⁻ cells could be rescued by plasmid-driven expression of PdCysZ (Fig.

4b), as well as IlCysZ and PfCysZ (data not shown). Consistent with previously reported data

(Zhang et al., 2014), we observed that SO_3^{2-} ions severely impede or block the accumulation of

$[^{35}S]O_4^{2-}$ by CysZ-expressing cells (Fig. 4b).

To further characterize the interaction of CysZ with both SO_4^{2-} and SO_3^{2-}, we performed binding

experiments on purified PdCysZ using the scintillation proximity assay (SPA) (Quick & Javitch,

2007). To determine the concentration of half-maximal binding (EC_{50}), we isotopically diluted

$[^{35}S]O_4^{2-}$ with non-labeled SO_4^{2-}, obtaining an EC_{50} of 0.81 ± 0.045 µM (Fig. 4c). Competing

binding of $[^{35}S]O_4^{2-}$ with SO_3^{2-} revealed that PdCysZ binds SO_3^{2-} with greater affinity as reflected

by a half-maximum inhibition constant (IC_{50}) of 0.12 ± 0.034 µM.

Next, we conducted $[^{35}S]O_4^{2-}$ uptake experiments on detergent solubilized and purified CysZ

reconstituted in proteoliposomes. Proteoliposomes provide a means for assessing the activity of a

membrane protein independent of other proteins found natively in cellular expression systems.

Such binding partners could be involved in SO_4^{2-} uptake as measured in intact $E.\ coli$ cells (Fig.

4a). Furthermore, in addition to the SPA measurements, these experiments allowed us to confirm

that the detergent-based extraction from the membrane and purification of CysZ used for our

crystallization approaches did not compromise its activity. Acting alone, PdCysZ in

proteoliposomes mediated the accumulation of SO_4^{2-} in a time-dependent manner; the uptake

occurred rapidly, peaking at about 30 seconds after incubation with the radioligand (Fig. 4d).

Consistent with our observations on cells, the influx of sulfate was inhibited by the presence of

SO_3^{2-} at micromolar concentrations. *Il*CysZ and *Pf*CysZ exhibited similar uptake profiles in

liposomes, suggesting similar functional properties (Fig. 4-S1a).

To determine whether sulfate uptake by CysZ was driven by the concentration gradient of sulfate

itself, or if it was coupled to a secondary driving force such as an ion (e.g., H^+, Na^+) gradient, we

tested the effect of the uncouplers CCCP (carbonyl cyanide m-chlorophenyl hydrazone), and

oligomycin, as well as that of the ionophores valinomycin and gramicidin on CysZ-mediated

$[^{35}S]O_4^{2-}$ accumulation in our cell-based assay (Buckler & Vaughan-Jones, 1998). Since none of

these uncoupling agents had a significant effect on the accumulation of $[^{35}S]O_4^{2-}$, these

experiments suggest that CysZ-mediated $[^{35}S]O_4^{2-}$ flux is not dependent on the 'classical' H^+ or

Na^+ electrochemical transmembrane gradient, thus indicating that the observed uptake solely

depends on the SO_4^{2-} transmembrane concentration gradient (Fig. 4-S1b), a notion that is

consistent with a channel-like mechanism for CysZ-mediated SO_4^{2-} translocation.

Finally, we employed the use of single-channel electrophysiological current recordings on CysZ

(*Pd*CysZ and *Il*CysZ) reconstituted in a planar lipid bilayer (Fig. 4-S2). Consistent with our

$[^{35}S]O_4^{2-}$-based flux assays, these electrophysiological measurements showed that reconstituted

CysZ exhibits channel-like properties when a membrane potential is applied in symmetrical

Na_2SO_4 solutions on both sides of the bilayer. The *Il*CysZ protein exhibits an average open

probability (P_{OPEN}) of 0.8 and conductance estimated as 92 ± 12 pS. However, unitary

conductance levels varied, showing a dependence on the amount of protein reconstituted in the

planar lipid bilayer. The *Pd*CysZ protein passes a broad distribution of currents at fixed voltages.

For both proteins, the SO_4^{2-} currents are completely abolished by the presence of low micromolar

SO_3^{2-} concentrations.

Sulfate Binding Site

A bound SO_4^{2-} ion was observed in the structure of *Il*CysZ. This binding site was confirmed by

purifying and crystallizing CysZ in the presence of the heavier SO_4^{2-}-analog selenate (SeO_4^{2-}). In

the SO_4^{2-} bound structure, each protomer in the dimer showed electron density consistent with a

bound sulfate, but only one of the two refined to full occupancy, hence only one is shown in the

refined model of *Il*CysZ (Fig. 5a). The location of the bound SO_4^{2-} is close to the putative

membrane interface where it is coordinated by two arginine residues (R27 and R28 in *Il*CysZ)

and the backbone amides of residues G25 and L26. This SO_4^{2-}-binding site is in the loop between

H1 and H2, with the motif GLR(R) being well conserved among the CysZ family. Consistent

with this observation, crystals of *Il*CysZ in which R27 and R28 were replaced with alanines did

not show interaction with SO_4^{2-} in that site based on i) the lack of any density in the refined

mutant structure at 2.3 Å and ii) functional studies (see below). We could not confirm the

presence of SO_4^{2-} in either of the other structures of *Pf*CysZ and *Pd*CysZ, owing, at least in part,

to their lower resolution limit (~3.5 Å). Soaking and co-crystallization attempts on both *Pf*CysZ

and *Pd*CysZ with SeO_4^{2-} resulted in cracking and destabilization of crystals, and loss of

measurable diffraction. In this vein, *Il*CysZ R27A/R28A, and the corresponding *Pf*CysZ R25A

and *Pd*CysZ R23A mutants all have severely impaired sulfate uptake capability, as demonstrated

in cells and proteoliposomes (Fig. 5b, 4-S1a), consistent with a conserved role for these targeted

arginine residues in all CysZ variants.

Conserved core in the hydrophilic head

Each apex of the triangular faces of the *Pd*CysZ hexamer comprises an extra-membranous hydrophilic head of a CysZ protomer (Fig. 1a and 5c). The central core of this hydrophilic head consists of the ends of H3 and H5 and the start of H4, with their residues forming an intricate network of hydrogen bonds and salt bridges that hold this helical bundle together (Fig. 5c). Two highly conserved motifs in this region, ExVE and QYxDYPxDNHK (Fig. 2c), likely play critical roles in this network of interactions. The two aspartates E103 and E106 (*Pd*CysZ) in the first motif interact with R129, N184 and R110, H185, respectively (Fig. 5c). The conserved R129 and R133 on H4b in turn also interact with D183 and N184 in the DNHK sequence stretch of H5b in the second motif. In the second motif, a conserved tyrosine (Y177) lies below the membrane interface towards the core of the molecule, along H5.

Mutational studies on a subset of these conserved hydrophilic-head residues highlight their functional relevance (Fig. 5d). R129 mutants (R129E or Q) exhibited a severe loss of SO_4^{2-} binding and uptake, and mutations made to Q176 and Y177 also showed loss in function, albeit to a lesser extent. Charge reversal mutations made to E103 and E106 (to K or R) or the equivalent *Il*CysZ E107, E110 resulted in very poor expression and stability levels, suggesting that the disruption of the interaction network in this region may destroy the structural integrity of the protein.

Putative pore and sulfate translocation pathway of CysZ

It was already evident from the initial *Il*CysZ structure that the hydrophilic head presented an incipient opening into the membrane, as seen in the center of the Fig. 5c inset, but it was quite

unclear how this opening might relate to transmembrane sulfate translocation. The 'transverse'

dimer structure of *Pf*CysZ clarified the possibility for ion permeation by showing that openings

in its two protomers aligned across the putative membrane; however, the prospective

transduction pathway would then be open on one side to the bilayer. The *Pd*CysZ structure

showed that the open sides of transverse dimers line a central cavity in the *Pd*CysZ hexamers.

Thus, plausible transduction pathways became more evident.

The view into a protomer surface along the direct pathway between transverse dimer apices (e.g.

green to orange in Fig. 3e) displays the pattern of exceptionally high conservation associated

with the entrance to a putative pore for sulfate translocation (Fig. 6a, b). This putative entrance

or 'pore' lies in the midst of a tight network of interacting residues, which is seen detailed in the

inset of Fig. 5c in the very same view as for Fig. 6a. These interactions close the incipient pore

from ion conduction in this conformation. The electrostatic potential surface of PdCysZ (Fig. 6c)

shows striking and puzzling electronegativity at the conserved pore entrance (Fig. 6d compared

with Fig. 6b). Because of the conservations, similar electrostatics pertain to the two other

homologs. Moreover, we observe the same helix dispositions, pore shape and charge interactions

in all three structures, evident when the protomers are superimposed (Fig. 3c, top view).

Pathway prediction algorithms (namely, PoreWalker (Pellegrini-Calace, Maiwald, & Thornton,

2009)) performed on the CysZ protomer revealed a putative ion translocation pathway that

begins at the entrance to the putative pore and ends in the large central cavity of the CysZ

hexamer that lies within the membrane (Fig. 7a, b). In the structure of *Pd*CysZ, the entrance of

the putative pore appears to be closed by the network of charge-interactions by conserved

residues, R110, E106, N184, H185 and W235 (Fig. 6a, 7c-1). These residues interact to tightly

restrict access to the putative translocation pathway and constrict the entry of ions, likely

selected for size and charge (Fig. 7-S1a). After the narrow entrance, the putative pathway

broadens, surrounded by a ring of conserved asparagine and glutamine residues, N33, N91, N98,

Q176, and N220 (Fig. 7c-2). Mutations to N98 (N98A, N98D) led to impaired sulfate uptake

(Fig. 5d). Following this polar environment, the putative pathway widens even further leading

into a large, primarily hydrophobic, internal cavity encapsulated by the TM helices of the

*Pd*CysZ hexamer, located in the plane of the lipid bilayer (Fig. 7b, 7-S1c, d). The hydrophobic

cavity is enclosed on both the top and bottom by pairs of helices H4-H5, three on each side from

the six protomers, pointing in towards the three-fold axis, with each side of the cavity, at its mid-

section, measuring ~50 Å, when viewed from above or outside the membrane (Fig. 7-S1c, d).

Given the symmetric, dual topology nature of the CysZ assembly, the ions could then exit the

cavity via the same pathway as they entered, but traveling through CysZ protomers located on

the opposite side of the membrane.

DISCUSSION

There are four known families of dedicated SO_4^{2-} transport systems in prokaryotes[6], of which

CysZs are the least studied (Zhang et al., 2014). The sequence of CysZ, coding for an integral

membrane protein with four predicted TM segments, shows no resemblance to any other known

protein. This, and the quest to set the basis for a mechanistic understanding of function,

prompted us to investigate the structure of CysZ.

We present here three structures of CysZ from different species, all determined by x-ray

crystallography. These structures are all similar, showing a novel fold comprising two extended

TM helices and two hemi-penetrating helical hairpins, giving rise to a tripod-like shape within

the membrane, and a hydrophilic head (Fig. 2a, b, 3c). Two of the structures (*Il*CysZ and

*Pf*CysZ) show dimers in the crystals, albeit with different interfaces (Fig. 3a, b), and the third

structure (*Pd*CysZ) displays a hexameric assembly in multiple crystal forms (Fig. 1a).

All three structures have a dimeric component, each of which is present in the hexameric

arrangement present in all the crystal forms of *Pd*CysZ (Fig. 3a, b), showing how they are all

related. Furthermore, *Il*CysZ can be crosslinked in membranes to the alternative *Pf*CysZ dimer

by engineering of disulfide cysteine mutants (Fig. 3-S2), which suggests that *Il*CysZ can adopt

both of the dimer assemblies observed in the *Pd*CysZ hexamer. Thus, we hypothesize that CysZ

is a hexamer in nature, as observed in *Pd*CysZ, and that in the case of *Il*CysZ and *Pf*CysZ, this

assembly may have come apart into the subsequently crystallized dimeric components during the

process of detergent extraction from the membrane and purification. This could be explained by

the unusually labile, and conformationally-flexible, nature of the H4b-H5a helical hairpin of

CysZ, with only one pair of stabilizing TM helices per protomer (H2-H3). We presume that the

hexamer is maintained with the support and scaffolding of the lipid bilayer.

CysZ shows a clear dual topology arrangement, which we confirmed with cross-linking

experiments on membranes (Fig. 3-S2). Dual-topology insertion is fairly uncommon in

membrane proteins; however, reported cases include the well-documented EmrE, a multi-drug

resistant export protein that inserts into the membrane as an anti-parallel dimer, and the more

recently discovered and studied family of double-barreled 'Fluc' fluoride channels (Amadi,

Koteiche, Mishra, & McHaourab, 2010; Korkhov & Tate, 2009; Rapp, Granseth, Seppala, & von

Heijne, 2006; Stockbridge et al., 2015; Stockbridge, Robertson, Kolmakova-Partensky, & Miller,

2013).

Functional characterizations of the three CysZs for which we have obtained structural

information show that all three mediate SO_4^{2-} uptake and that this uptake is inhibited by SO_3^{2-},

for which the protein has higher affinity. Comparable results were obtained in cell-based as well

as in proteoliposome-based uptake assays (Fig. 4b, d, 4-S1a). Affinities for SO_4^{2-} and SO_3^{2-} were

measured by SPA (Fig. 4c). These results are consistent with previously reported data on *E. coli*

CysZ (Zhang et al., 2014). Experiments performed on cells with uncoupling agents (Fig. 4-S1b)

showed that CysZ-mediated SO_4^{2-} uptake was independent of a classical electrochemical

gradient, thus suggestive of the driving force being a concentration gradient of SO_4^{2-} across the

membrane. This hypothesis is supported by our single-channel bilayer experiments performed

with *Pd*CysZ and *Il*CysZ, which show that CysZ has channel-like properties (Fig. 4-S2).

Building on our functional and structural discoveries, our results suggest a fascinating hypothesis

for mechanisms of SO_4^{2-} transfer and regulation. There is a conserved central core (Fig. 6a),

which is likely to have a structural role. Close to this lies a sulfate-binding site, with functional

implications (Fig. 5a), and the entrance to a putative pore (Fig. 5c). The entrance of this putative

pore is delineated by a hydrophilic network of conserved residues that form a tight constriction

in our observed conformation. Following this hypothetical route, sulfate ions that might enter

through the three separate pores, one per CysZ dimer pair, would converge into a central

hydrophobic cavity (Fig. 7b, 7-S1c, d). Once sulfate ions enter this central cavity, due to the

unfavorable environment, these are likely to exit it rather rapidly through one of the three

available exit pores on the cytoplasmic side of the hexamer. Hydrophobic cavities and pores are

seen commonly in ion channels, with examples ranging from the well documented hydrophobic

inner pores of the various potassium channels (Doyle et al., 1998) to the SLAC1 (Chen et al.,

2010) and bestrophin anion channels (Yang et al., 2014), and the MscS and MscL

mechanosensitive channels (Anishkin, Akitake, Kamaraju, Chiang, & Sukharev, 2010; Bass,

Strop, Barclay, & Rees, 2002; Birkner, Poolman, & Kocer, 2012; Chen et al., 2010; Doyle et al.,

1998), facilitating the rapid passage of ions due to the unfavorable environment, as well as

potentially providing a means of 'hydrophobic gating' (Aryal, Sansom, & Tucker, 2015).

The surface electrostatics of CysZ (Fig. 1b and Fig. 6c, d) draw attention to the negative

potential of the conserved core of each protomer, which is then surrounded by a more neutral

annulus. While this feature seems contradictory to admission of SO_4^{2-} ions at the extracellular

side, it could be advantageous for expulsion into the cytoplasm on the opposite side. In any case,

the structures that we have determined are evidently in a closed state, implying that a

conformational change would have to occur to allow passage of SO_4^{2-}, likely modifying the

surface electrostatics of the protein. It is tempting to speculate that the regulation of the opening

of the pore could be modulated by the binding of sulfate ions to the identified sulfate-binding

site, as it is near the entrance of the putative pore. L22 lies in proximity of the entrance of the

putative pore, and its backbone amide (along with G21) coordinates the sulfate ion in the GLR

motif of the sulfate-binding site. Thus, the binding of sulfate to the GLR motif could trigger a

conformational change needed to displace L22, allowing for a wider opening for the sulfate ions

to enter the pore. Sulfite could hypothetically exert its inhibitory effect on CysZ function by

binding to this site.

There is an overall electropositive region in the center of the hexameric molecule, when viewed

from the top (Fig. 1b, 6c, d). This central region is lined by conserved residues along helices

H4b-H5a, namely, R129, R133 and K137. The hydrophobic tips of helices H4b-H5a of the three

protomers on each side of the membrane then converge in the center of the hexamer. A pore

through the three-fold axis of the hexamer could provide an alternative passageway for SO_4^{2-}

ions through this assembly. Although we cannot exclude it, this possibility seems less plausible

because of the poor conservation at the tips of helices H4b-H5a and the strictly hydrophobic

nature of this region.

We observe, in agreement with previous data, that SO_3^{2-} inhibits CysZ-mediated SO_4^{2-} flux (Fig.

4b, d, 4-S2a, f) (Zhang et al., 2014). The dual topology nature of CysZ could provide a means for

internal as well as external regulation. However, experiments in solutions – such as

crystallizations – where all molecules are exposed to the same chemical environment, makes

capturing such a state in an open or SO_3^{2-} blocked conformation challenging. Despite this

limitation, our CysZ structures and associated functional experiments have allowed us to make

substantial progress in the understanding SO_4^{2-} uptake by these membrane permeases. This work

sets the framework for future experiments aimed at unraveling the molecular details of how

SO_4^{2-} is translocated across the membrane by CysZ and how this process is regulated.

PDB Accession codes: *Idiomarina loiheinsis* CysZ: **3TX3**, *Pseudomonas fragi* CysZ: **TBD,**

Pseudomonas denitrificans CysZ: **TBD**.

Materials and Methods

Ortholog selection and cloning

A total of 63 *cysZ* candidate genes were selected by a bioinformatics approach implemented by

the New York Consortium of Membrane Protein Structure (NYCOMPS), as previously described

(Punta et al., 2009). The majority of the genes (including *Il*CysZ, uniprot ID: Q5QUJ8) were

PCR-amplified from fully sequenced prokaryotic genomic DNA (obtained from ATCC®) (Love

et al., 2010). CysZ genes from certain species, such as *Pf*CysZ (uniprot ID: A0A0X8F058) and

*Pd*CysZ (uniprot ID: M4XKU7) were chemically synthesized by GenScript®, with codon-

optimization for protein expression. All genes were cloned by ligation-independent cloning

(LIC) (Aslanidis & de Jong, 1990) into an IPTG (isopropyl β-D-1-thiogalactopyranoside)

inducible, kanamycin-resistant pET derived plasmid (Novagen), with an N-terminal deca-

histidine tag (His10) and a TEV (tobacco etch virus) protease site to allow for tag cleavage upon

purification.

Protein expression and purification

Expression plasmids bearing the *cysZ* genes were transformed into BL21(DE3)pLysS cells using

standard protocols, and grown at 37 °C in 2XYT media supplemented with 50 μg/ml kanamycin

and 50 μg/ml chloramphenicol in an orbital shaker at 250 rpm. Protein expression was induced

for ~16 hours at 22 °C with 0.2 mM IPTG once an absorbance ($A_{600\ nm}$) of 0.8-1.0 was reached.

Selenomethione (Se-Met)-incorporated proteins were expressed in BL21(DE3)pLysS cells

grown using an M9 minimal media kit (Shanghai Medicilon Inc.) supplemented with the

necessary minerals, vitamins and non-inhibitory amino acids. Se-Met was added prior to IPTG

induction at an $A_{600\ nm}$ of 1.2. The Se-Met-incorporated protein was purified using the same

procedures as the native protein. Once harvested, the cells were resuspended at 0.2 g/ml in lysis

buffer containing 20 mM Na-Hepes pH 7.5, 200 mM NaCl, 20 mM MgSO4, DNase I and RNase

A, 0.5 mM PMSF (phenylmethylsulfonyl fluoride), EDTA-free Complete protease inhibitor

cocktail (Roche) and 1 mM TCEP-HCl (Tris (2-carboxyethyl) phosphine hydrochloride) as a

reducing agent. Initial small-scale expression and detergent screening was performed on 80 mg

of pelleted cells (wet weight), and 7-10 g of cells for large-scale protein purification. Cells were

lysed using an Avestin[®] EmusiFlex-C3 homogenizer, followed by protein solubilization with 1%

(w/v) decyl maltopyranoside (DM) (Anatrace® Affymetrix) for 1 hour at 4 °C, after-which

insoluble material was removed by ultra-centrifugation at 100,000 x g. The solubilized protein

was applied to Ni-NTA Sepharose (Qiagen) in batch, washed with lysis buffer containing 0.2%

DM and 40 mM imidazole and eluted in buffer containing 250 mM imidazole. Upon elution,

CysZ was dialyzed overnight with His-tagged TEV protease at 4°C, against a buffer containing

20 mM Na-Hepes pH 7.0, 200 mM NaCl, 0.2% DM, 1 mM TCEP-HCl and 20 mM Na2SO4,

allowing for the cleavage of the His10 tag and removal of the imidazole. Tagless CysZ was then

re-passaged over Ni-NTA sepharose to re-bind of any uncleaved CysZ, TEV protease and the

cleaved His10 tag. The protein was then subjected to size-exclusion chromatography (Superdex

200 10/30 HR, GE Healthcare) in 20 mM Na-Hepes pH 7.0, 200 mM NaCl, 1 mM TCEP-HCl,

20 mM Na2SO4 and appropriate detergent for crystallization, for *Il*CysZ: 0.06% Lauryl

dimethylamine oxide (LDAO) and for *Pf*CysZ and *Pd*CysZ: 1% β-octyl glucopyranoside (β-

OG). The choice of detergent was made based on protein yield, stability and mono-disperse gel-

filtration peaks obtained in the initial small-scale detergent screening. A yield of ~1.5 mg of

purified CysZ was typically obtained from a cell pellet of 7-8 grams (1 liter of culture).

Protein Crystallization

_Il_CysZ: Crystals of _Il_CysZ in LDAO were obtained by vapor diffusion at a protein concentration

of 6-8 mg/ml at 4°C, in a 1:1 v/v ratio against a precipitant of 28-32% PEG400, 0.1M Tris-HCl

pH 8.0, with salt additive of 0.1 M NaCl or 0.1 M $MgCl_2$. The crystals appeared overnight,

continued to grow in size over the course of 2-4 days after set-up. After optimization, the crystals

grew to a maximum size of ~200 μm x 100 μm x 50 μm with a rhomboid or cuboid shape. The

crystals were harvested directly without the addition of a cryo-protectant and flash-frozen into

liquid nitrogen, for data collection on the X4A/X4C beamlines at National Synchrotron Light

Source (NSLS), Brookhaven National Labs (Upton, NY). SeMet derivatized and selenate co-

crystals were obtained from the same conditions as the native crystals.

_Pf_CysZ: Crystals of _Pf_CysZ in β-OG at 5 mg/ml were initially obtained at 4 °C by vapor

diffusion, in a 1:1 protein to precipitant ratio, after 1-2 days against 28% PEG400, 0.1 M MES

pH 6.0. They were cuboid in shape and grew in clusters of multiple crystals originating from a

common locus. The crystals were optimized to a maximal size of ~150-200 μm x 50 μm x 50

μm, with the best diffracting crystals grown under silicone oil (visc. 500) in microbatch Terazaki

plates. Crystals were directly flash-frozen into liquid nitrogen without the use of a cryo-

protectant and were exposed to X-rays at the NE-CAT (241DC and E) beamlines at APS,

Argonne National Lab (Argonne, IL) for data collection. SeMet crystals were obtained from the

same conditions as the native protein.

*Pd*CysZ: Crystals of *Pd*CysZ in β-OG at 5-8 mg/ml were initially obtained at 4°C by vapor

diffusion, in a 1:1 protein to precipitant ratio, after 2-3 days against 22-30% PEG550MME, 0.1

M Na-Hepes pH 7.0. The rod-like crystals were hexagonal on one face, and grew in clusters

originating from a common locus as well as on the edge of the drop. The multiple crystal forms

observed were all obtained in the same crystallization conditions. The crystals were optimized to

a maximal size of ~200-250 μm x 25 μm x 50 μm. With the addition of 20% glycerol (w/v) as a

cryo-protectant, the crystals were flash-frozen into liquid nitrogen and were exposed to X-rays at

the NE-CAT (24-IDC and 24-IDE) beamlines at APS, Argonne National Lab (Argonne, IL) for

data collection.

Data collection and structure determination

*Il*CysZ: The structure of CysZ was determined by the single-wavelength anomalous diffraction

(SAD) method from anomalous diffraction of a selenate (SeO_4^{2-}) derivative crystal. The

anomalous signals were measured at the Se K-edge peak wavelength, which was determined

experimentally from fluorescence scanning of the crystal prior to data collection. All diffraction

data were recorded at 100K using an ADSC Q4R CCD detector at the NSLS X4 beamline.

Diffraction data were indexed, integrated, scaled, and merged by HKL2000 (Otwinowski &

Minor, 1997). Selenate substructure determination was performed with the SHELXD program

through HKL2MAP (Pape & Schneider, 2004). A resolution cut-off at 2.6 Å was used for

finding Se sites by SHELXD. A strong peak found by SHELXD was used to calculate initial

SAD phases, which were improved by density modification by SHELXE (Sheldrick, 2010). With

a solvent content of 65% corresponding to 2 molecules in the asymmetric unit, 50 cycles of

density modification resulted in an electron density map of sufficient quality for model building.

The initial polypeptide chain was built by Arp/Warp (Langer, Cohen, Lamzin, & Perrakis, 2008),

at 2.1 Å by using experimental phases. Further cycles of model building were performed

manually using COOT (Emsley, Lohkamp, Scott, & Cowtan, 2010) and all rounds of

refinements were performed with PHENIX (Adams et al., 2010). The native structure of CysZ

with bound sulfate was determined both by multi-crystal native SAD (Liu et al., 2012) (final

resolution of 2.3 Å) and by molecular replacement with the selenate bound model (final

resolution of 2.1 Å). In addition, phase information obtained from Se-Met derivatized protein

with 9 Se sites per CysZ molecule, verified our model obtained from the selenate data.

*Pf*CysZ: Multi-crystal SeMet-SAD data sets were collected at APS beamline 24-IDC with a

Pilatus 6M pixel array detector under a cryogenic temperature of 100 K. To enhance anomalous

signals from Se atoms for phasing (Liu, Zhang, & Hendrickson, 2011), the X-ray wavelength

was tuned to the Se-K edge (λ=0.9789 Å). The orientation of crystals was random without

special consideration of crystal alignment, and beam size was adjusted to match the crystal size.

A total of 22 data sets were collected, each from a single crystal. An oscillation angle of 1° was

used for data collection with a total of 360 frames for each data set. The beam size was adjusted

to match the crystal size. The 22 single-crystal data sets were processed individually by using

XDS (Kabsch, 2010) and CCP4 packages (Winn et al., 2011). For phasing purposes, the low-

resolution anomalous signals were enhanced by increased multiplicity. By rejection of 7 outlier

crystals (Liu et al., 2012), anomalous diffraction data from 15 statistically-compatible crystals

were scaled and merged for phasing. For outlier rejection, a unit-cell variation of 1.0σ was used.

CCP4 program POINTLESS and SCALA (Evans, 2006) were used for data combining; and

Bijvoet pairs were kept separately throughout the data flow. For refinement purposes, keeping

the high angle data was important, and done by limiting radiation damage as well as by

increasing multiplicity. Although most *Pf*CysZ crystals diffracted to only about 3.5 Å spacings

or poorer, we intentionally set the detector distance to include higher spacings. Higher resolution

data were retained through a data merging procedure that is described as follows: 1) The 22

individually processed data sets were analyzed by diffraction dissimilarity analysis by using only

high angle data between 3.5-3.0 Å, resulting in three subsets. 2) The data statistics of members in

each subset were checked manually and the subset that contained the highest angle data set, e.g.

data set 6, was selected for further analyses and data combination. 3) Each data set within the

selected subset was compared with data set 6 by high-angle intensity correlation. Six of the

highest resolution data sets were statistically comparable and therefore were selected for

merging. For phasing, substructure solutions were found by SHELXD (Sheldrick, 2010) and

were further refined and completed by PHASER (McCoy et al., 2007) and then used to compute

initial SAD phases at the data limit by SAD phasing with PHENIX (Adams et al., 2010). Phases

were density modified with solvent flattening and histogram matching as implemented in CCP4

program DM (Cowtan & Zhang, 1999) to improve phases and also to break phase ambiguity.

The estimated solvent contents of 71% were used for density modification. The model was

initially built into the experimental electron density map by COOT (Emsley et al., 2010),

followed by iterative refinement by PHENIX and model building in COOT. The refined model

does not contain solvent molecules at this resolution.

*Pd*CysZ:

Native crystal data were collected at the APS beamline 24-IDC with a Pilatus 6M pixel array

detector at a cryogenic temperature of 100 K at an X-ray wavelength of $\lambda=1.023$ Å. The sample-

to-detector distance was set to 500 mm. An oscillation angle of 0.5° was used for data collection.

The beam size was adjusted to match the crystal size. Molecular replacement was attempted

using a variety of search models (*Pf*CysZ and *Il*CysZ monomer/dimer models, with various

degrees of truncation). Success was achieved by searching for six copies of a search model

consisting of the *Pf*CysZ monomer, with residues 36-53 deleted and the sequence adjusted using

CHAINSAW (Stein, 2008) (pruning non-conserved residues to the gamma carbon). Density

modification of the initial map was performed in PARROT, incorporating solvent flattening,

histogram matching and NCS-averaging. An initial round of model building was performed in

COOT (Emsley et al., 2010) into this map, followed by further phase improvement and bias-

removal using *phenix.prime_and_switch* (Adams et al., 2010*)*. The improved map was used for a

second round of model building, followed by iterative cycles of reciprocal space refinement

using *phenix.refine*, and real-space refinement and correction in COOT.

All graphical representations and figures of our structural models were made in either PyMOL

(Schrodinger, 2010) or Chimera (Pettersen et al., 2004).

Radioligand Binding by Scintillation Proximity Assay (SPA)

CysZ was purified by standard protocols as described above, with the exception of leaving the

histidine-tag intact without cleavage by the TEV protease. The imidazole was removed by

dialysis and the purified protein was not run over the size exclusion column, and instead was

directly concentrated to 2 mg/ml for the SPA experiment. $[^{35}S]O_4^{2-}$ obtained in the form of

sulfuric acid (American Radiolabeled Chemicals (ARC)) was used as the radioligand. 100 ng –

250 ng of CysZ was used per assay point, diluted in 100 μl of assay buffer containing 20 mM

Hepes pH 7.0, 200 mM NaCl, 0.2% DeM, 20% glycerol, 0.5 mM TCEP, with 12.5 μl of Copper

YSi beads (Perkin Elmer). A trace amount of $[^{35}S]O_4^{2-}$ (~10-30 nM) mixed with cold ligand

(Na$_2$SO$_4$) was added to each well of a 96 well clear bottom plate along with the mix of protein,

buffer and beads. The plate was agitated for a minimum of 30 minutes to as long as overnight,

and measured in a scintillation counter the following morning (MicroBeta, Perkin Elmer).

Competitive binding experiments with a gradient of a competing cold ligand ranging from (10

nM − 100 mM) were performed in triplicate, with the control measured at each concentration by

adding 1 M Imidazole to the reaction buffer, to prevent binding of the protein to the scintillation

beads. All data were analyzed and graphically represented with GraphPad™ Prism6 software.

[^{35}S]O$_4^{2-}$ Uptake Experiments

In whole cells: The *cysZ* gene knockout strain in *E. coli* K-12 was obtained from the Coli

Genetic Stock Center at Yale University (http://cgsc.biology.yale.edu/), originally found in the

Keio knockout collection (Baba et al., 2006). Genotype: F-, Δ(araD-araB)567, Δ

lacZ4787(::rrnB-3), λ-, ΔcysZ742::kan,rph-1, Δ(rhaD- rhaB)568, hsdR514. The knockout

strain was made competent by standard protocols (Hanahan, 1983) to allow for the

transformation and expression of *cysZ*-containing plasmids, for rescue experiments. Wild-type

parental *E. coli* K-12 was used as a control strain. The strains were grown in LB (Luria Broth)

without any antibiotic (for the WT cells) and with 50 μg/ml of kanamycin for the knockout cells

at 37 °C overnight to saturation. The culture was spun down at 3000 x g the next morning and

the cells were resuspended in Davis-Mingioli (DM) minimal media without sulfate (MgSO$_4$ was

replaced with MgCl$_2$ and (NH$_4$)$_2$SO$_4$ was replaced by NH$_4$Cl), supplemented with 0.63 mM L-

cysteine to allow the cells to grow in the absence of sulfate (Davis & Mingioli, 1950). The cells

were incubated in DM media for a minimum of 9 hours, maximum overnight, at 37°C after

which they were spun down and washed 3 times in 5 mM Na-Hepes pH 7.0. This ensured that

the cells were starved of SO_4, to deplete the sulfate stores in the cell, enhancing sulfate uptake

measured (F. Parra, personal communication). The cells were resuspended in the same buffer at

0.7 g/ml at room temperature, and 10 μl of the cell suspension was used for each point in the

uptake experiment. A final concentration of 320 μM of SO_4 (Na_2SO_4 with $[^{35}S]O_4^{2-}$ as a tracer)

was used outside. The uptake experiments were performed in triplicate, measured over a time

course of typically 0 - 300 seconds. The reaction was stopped by dilution, by adding 1.5 ml of

ice-cold buffer (5 mM Na-Hepes pH 7.0) to the cells in the tube. The reaction mix was

immediately poured onto a vacuum filtration device, with an individual glass-fiber filter (0.75

μm, GF/F) per assay point, after which the filters are washed once more with 1.5 ml of buffer.

The dry filters, with the cells attached to their surface, were then moved to scintillation vials,

containing 4 ml of EconoSafe® scintillation cocktail, to be counted the next morning. For the

CysZ cell rescue experiments, a similar protocol was used, transformed with an ampicillin

resistant expression vector containing the CysZ gene, mutant or empty vector (as the control).

Upon transformation, the cells were grown overnight in a 4 ml starter culture of LB with 50

μg/ml kanamycin and 100 μg/ml ampicillin. The next morning the entire 4 ml was used to

inoculate 75 ml of LB (Kan, Amp), and grown at 37 °C for 2.5-3 hours until an OD_{600} of 0.6-0.8

was reached. Protein expression was then induced at 37 °C for 4 hrs with 0.2 mM IPTG. After

4hrs, the cells were spun down and resuspended in the DM minimal media without sulfate, 0.63

mM cysteine, Amp, Kan and 0.2 mM IPTG, and incubated overnight at 22 °C. The next morning

the cells were spun down, washed 3 times with 5 mM Na-Hepes 7.0 and resuspended at final

concentration of 0.7 g/ml for uptake.

In proteoliposomes: CysZ was purified by the procedure described above, and upon elution from

the size-exclusion column concentrated to 1 mg/ml for reconstitution into liposomes at a protein

to lipid ratio of 1:100. The liposomes were comprised of a 3:1 ratio of *E. coli* polar lipids and

phosphatidylcholine (PC) (Avanti Polar Lipids), prepared by previously described methods

(Rigaud, Pitard, & Levy, 1995). Typically, 10 μl of CysZ-proteoliposomes and control 'empty'

liposomes at a concentration of 5-10 mg/ml, were used per assay point and diluted into a reaction

buffer of 100 μl, containing $[^{35}S]O_4^{2-}$ (at trace amounts of ~10-15 nM) supplemented with

Na_2SO_4 at the desired concentration. The experiment was performed in triplicate and sulfate

accumulation was measured either over a time-course or at a fixed time with different amounts

of SO_4^{2-} or SO_3^{2-} present. The reaction was stopped by dilution, by adding 1.5 ml of ice-cold

buffer (5 mM Na-Hepes pH 7.0) to the cells in the tube. The reaction mix was immediately

poured onto a vacuum filtration device, with an individual nitrocellulose filter (0.22 μm) per

assay point, after which the filters are washed once more with 1.5 ml of buffer. The dry filters,

with the cells attached to their surface, were then moved to scintillation vials, containing 4 ml of

EconoSafe® scintillation cocktail, to be counted the next morning.

Measurement of CysZ single-channel activity in the planar lipid bilayer

A previously described method (Mueller, Rudin, Tien, & Wescott, 1962) to insert ion channels

incorporated in liposomes into "painted" planar lipid bilayers by vesicle fusion was used to

incorporate CysZ into a lipid bilayer created on a small aperture between two aqueous

compartments, called the cis and trans compartments (Morera, Vargas, Gonzalez, Rosenmann, &

Latorre, 2007). Since this system is very sensitive to contaminants, CysZ was expressed in and

purified from a porin- deficient strain of *E. coli* cells to prevent any carry through of

contaminating porins that could create large conductances and artifacts in the single-channel

recordings. Phosphatidylethanolamine (PE) and phosphatidylserine (PS) (Avanti Polar Lipids),

in a ratio of 1:1 were dissolved in chloroform to mix, and dried completely under an argon

stream. The mixed and dried lipids were then dissolved in n-decane to a final concentration of 50

μg/ml, and kept at 4 °C. The lipids are always prepared fresh, on the same day of the experiment.

The purified CysZ protein were incorporated into PE:PS (1:1) liposomes by brief sonication at

80kHz for 1 minute at 4 °C. The experimental apparatus consisted of two 1-ml buffer chambers

separated by a Teflon film that contains a single 20- to 50-μm hole. A lipid bilayer was formed

by "painting" the hole with the 1:1 mixture of PE:PS, this results in a seal between the two cups

formed by the lipids (Leal-Pinto, London, Knorr, & Abramson, 1995). For these studies, the cis

side was defined as the chamber connected to the voltage-holding electrode and all voltages are

referenced to the trans (ground) chamber. Stability of the bilayer was determined by clamping

voltage at various levels. If a resistance > 100 Ω and noise < 0.2 pA were maintained in the

patch, the proteoliposomes containing CysZ were added to the trans side of the chamber and

stirred for 1 min. The fusion event or insertion of a channel into the bilayer was assessed by the

presence of clear transitions from 0 current to an open state.

679 Site-specific cysteine labeling experiments

All functional and cysteine mutants of CysZ were generated by site-directed mutagenesis, using

the QuikChange® site-directed mutagenesis kit (Strategene). The sequence-verified mutants

were then tested for expression in comparison to the WT CysZ. To address the membrane

topology of *Il*CysZ, single cysteine mutants were designed to perform site-directed fluorescence

labeling based on the accessibility of the cysteine to the membrane impermeable thiol-directed

fluorescent probe (Ye, Jia, Jung, & Maloney, 2001). A set of surface-exposed residues at

different positions on the CysZ molecule were selected to be mutated to cysteines, based on the

*Il*CysZ structure. The cysteine mutants were expressed by standard protocols that were used for

the WT protein. The membrane fraction of each mutant was pelleted after cell lysis by

ultracentrifugation at 100,000 x g and resuspended at 20 mg/ml (Bradford assay) in fresh buffer

containing 20 mM Na-Hepes pH 7.0, 200 mM NaCl, protease inhibitors: 0.5 mM PMSF and

Complete protease inhibitor cocktail EDTA-free and 1 mM TCEP-HCl. 1 ml of membranes were

then incubated with 30 μM membrane-impermeant fluorescein-5-maleimide dye (Invitrogen, 2

mM stock freshly prepared in water, protected from light) for 30 minutes in the dark at room

temperature. The labeling reaction was stopped by the addition of 6 mM β- mercaptoethanol, and

the membranes were spun down and resuspended in fresh buffer to remove any remaining

unreacted fluorescent dye. The fluorescently labeled protein was then purified from the

membranes by solubilization with 1% DM, using a standard Ni-NTA purification protocol,

qualitatively analyzed on an SDS-PAGE, and quantitatively measured by a Tecan® fluorescence

plate reader at an excitation of 495 nm, emission of 535 nm.

Crosslinking of Cysteine Mutant Pairs in CysZ

Cysteine mutants of *Pf*CysZ and *Il*CysZ were designed based on pairs of residues that were in

close proximity (within 3-7 Å) of each other in our structure that could have the ability to

covalently join the 2 protomers of the dimer. The single and double cysteine mutants were made

by site-directed mutagenesis using the QuikChange® Site-Directed Mutagenesis Kit

(Strategene). Mutants and WT were expressed using the standard protocols CysZ expression.

Isolated membrane fraction was resuspended by homogenization in fresh lysis buffer at a

membrane protein concentration of ~25 mg/ml (measured by Bradford Assay). Bis-

methanethiosulfonate (Bis-MTS) crosslinkers (Santa Cruz Biotechnolgies) of different spacer

lengths were used at 0.5mM (dissolved in DMSO) added to 1ml of resuspended membranes at

RT for 1 hour. Bis-MTS crosslinkers are membrane permeable, highly reactive and specific to sulfhydryl groups, and the covalent linkage is resistant to reducing agents like β-mercaptoethanol (Akabas, Stauffer, Xu, & Karlin, 1992). The crosslinking lengths used were: 1,1-Methanediyl Bismethanethiosulfonate (3.6 Å), 1,2- Ethanediyl Bismethanethiosulfonate (5.2 Å), 1,4-Butanediyl Bismethanethiosulfonate (7.8 Å) and 1,6-Hexanediyl Bismethanethiosulfonate (10.4 Å). The reaction was quenched by the addition of 10 mM free cysteine, and the protein was then extracted and purified from the membranes with 1% DM followed by Ni-NTA resin. The imidazole is then diluted out in the Ni-elute, and the His-tag is cut with TEV protease in small scale. The cleaved protein is passaged over Ni-NTA resin to remove contaminants and uncleaved protein, and flowthrough is run on a reducing SDS- PAGE and stained with Coomassie blue to analyze dimer formation. The mutants designed were L161C (single mutant), L161C-A164C and N160C-L168C (double mutants) for *Pf*CysZ; and L156C-Q163C and V157C-Q163C (double mutants) for *Il*CysZ.

Acknowledgments:

Crystallographic data for this study were collected on the NSLS beamline X4A at Brookhaven National Laboratory and on the NE-CAT beamlines 24ID-C and E (supported by NIH-NIGMS grant P41 GM103403) at the Advanced Photon Source. The Pilatus 6M detector on 24-ID-C beam line is funded by a NIH-ORIP HEI grant (S10 RR029205). This work was supported by NIH-NIGMS grants R01 GM098617 (F.M.) and R01 GM107462 (W.A.H.), and by grants through the New York Structural Biology Center for the New York Consortium on Membrane Protein Structure (NYCOMPS; U54 GM095315) and for the Center on Membrane Protein Production and Analysis (COMPPÅ; P41 GM116799). O.B.C. was supported by a Charles H.

Revson Senior Fellowship. We thank John Schwanof for assistance during data collection at

NSLS, Joe Mindell, Francisco Parra, Albano Meli, Giuliano Sciara and Carlos A Villalba Galea

for helpful advice and contributions at the early stages of the project, and Leora Hamberger for

her assistance managing the Mancia Lab.

Declaration of Competing Interests:

The authors declare that they have no financial or non-financial competing interests.

Figures and Tables:

Table 1: Crystallographic data and refinement statistics.

Data collection	Native *IlCysZ*	*IlCysZ* w/SeO$_4^{2-}$ (SAD)	SeMet *PfCysZ* (6 crystals, refinement)	SeMet *PfCysZ* (15 crystals, SAD)	Native *PdCysZ* (MR-SAD)
Beamline	NSLS X4A	NSLS X4A/C	APS 24-ID-E	APS 24-ID-C	APS 24-ID-C
Space group	C2	C2	C2	C2	P6$_3$
Cell dimensions:					
a, b, c (Å)	128.9, 82.0, 100.4	128.9, 81.9, 100.3	172.29, 56.9, 96.17	172.35, 56.9, 96.31	225.13, 225.13, 96.62
α, β, γ (°)	90, 125.1, 90	90, 125.1, 90	90, 91.43, 90	90, 91.33, 90	90, 90, 120
Z$_a$	2	2	2	2	6
Wavelength	1.7432	0.96789	0.97890	0.97890	1.0230
Bragg spacings (Å)	30-2.30	50-2.10	40-3.20	40-3.50	86-3.40
R$_{merge}$	0.047 (0.257)	0.046 (0.417)	0.136 (6.506)	0.135 (2.13)	0.095 (2.89)
I / σ$_I$	29.4 (7.9)	25.0 (1.9)	12.3 (0.7)	24.2 (2.6)	22.4 (1.2)
Completeness (%)	99.9 (99.9)	99.9 (100.0)	99.8 (99.3)	99.9 (100.0)	100.0 (100.0)
Multiplicity	7.3 (7.2)	7.6 (7.6)	27.2	72.8	19.7
Refinement:					
Resolution (Å)	2.30		3.50		3.40
No. of reflections	38075		20741		37221
R$_{work}$ / R$_{free}$	0.200/0.238		0.297/0.339		0.239/0.277
No. of atoms:					
Protein	3679		3428		11415
Ligand/ion	249		0		560
Water	162		0		0
Average B-factors (Å^2)	49.4		198.9		204.18
Protein	48.4		198.9		204.18
Ligand/Ion	77.3		-		-
Water	45.3		-		-
Bond Ideality (r.m.s.d.):					
Bond lengths (Å)	0.006		0.006		0.004
Bond angles (°)	0.894		1.133		0.96
Ramachandran Analysis:					
Favored (%)	99.0		99.2		98.5
PDB accession code	3TX3		TBD		TBD

744
Values in parentheses are from the highest resolution shell. R$_{free}$ was calculated using 5% of data

excluded from refinement.

Figure 1.

Fig. 1. Overall structure of the *P. denitrificans* CysZ (*Pd*CysZ) hexamer. a. Side and top

views of the hexamer as a ribbon diagram with each protomer chain colored differently. The

approximate dimensions of the hexamer marked in Å. b. Side and top views represented by

surface electrostatics as calculated by APBS, with negative and positive surface potential

represented in red and blue respectively. c. Side and top views representing conservation of

residues as calculated by ConSurf, with maroon being most conserved to cyan being least

conserved.

Figure 1 – Figure Supplement 1.

Fig. 1. S1. Schematic of assimilatory sulfate reduction in bacteria for cysteine biosynthesis.

Assimilatory reduction is energetically dispendious as sulfate is extremely stable. In this

pathway, sulfate ions enter the cell via a transporter or channel, like CysZ; this is followed by its

activation by ATP-sulfurylase, which utilizes ATP to form adensosine phosphosulfate (APS) and

inorganic pyrophosphate (PPi) as a by-product. APS is then further activated by the addition of a

second phosphate forming phosphoadenosine phosphosulfate (PAPS) and this high-energy

intermediate product is poised for sulfur removal to form sulfite ($SO3^{2-}$) by the thioredoxin

enzyme, which is then further reduced to sulfide ions (S^-). Sulfide, the final reduced sulfur

product, can then be used in a variety of ways by the cell, such as the incorporation into O-

acetylserine to synthesize cysteine.

Figure 1 – Figure Supplement 2.

774
775

Fig. 1. S2. Representative electron density of the crystal structures of *Il*CysZ, *Pf*CysZ and

***Pd*CysZ, and the different crystal forms observed for *Pd*CysZ.** a. Representative electron

density in blue mesh for: *Il*CysZ, contoured at 1 x r.m.s.d., *Pf*CysZ, contoured at 2 x r.m.s.d, with

a negative B-factor of -100 $Å^2$ applied to map; *Pd*CysZ, contoured at 1 x r.m.s.d, with a negative

B-factor of -100 $Å^2$ applied to map. Final atomic models (Cα trace with side chains depicted) are

shown as yellow stick representation. b. Preliminary molecular replacement analyses of the 3

different crystal forms obtained for *Pd*CysZ, showing the hexameric assembly of the molecule,

seen in each case. The Cα trace of an asymmetric unit in each case is depicted in grey, with their

symmetry mates are shown in yellow. Space group and unit cell dimensions are listed below.

Figure 2.

Fig. 2. Structure and topology diagram of *Pd*CysZ protomer, with sequence alignment and

conservation a. Ribbon diagram of the *Pd*CysZ protomer colored in rainbow colors from N

(blue) to C terminus (red), viewed from within the plane of the membrane, shown in the same

orientation as the protomer drawn in green in Fig. 1a. b. Topology diagram of the *Pd*CysZ

protomer with helices marked from 1 to 8. Helices H2b and H3a are transmembrane helices,

whereas H4-H5 and H7-H8 are hemi-penetrating helical hairpins, only partially inserted into the

membrane. c. Sequence alignment of *E. coli* CysZ (*Ec*CysZ), *P. denitrificans* CysZ (*Pd*CysZ), *P. fragi* CysZ (*Pf*CysZ), and *I. loihiensis* CysZ (*Il*CysZ). Residues are colored based on conservation, with maroon being most conserved and cyan least conserved, as calculated by ConSurf using a sequence alignment of 150 non-redundant sequences from the CysZ family as input. Spirals above residues mark the extent of the helical segments based on the atomic structure of *Pd*CysZ with helices numbered H1-H8; letters mark residue identities; black dots above identify every tenth residue (modulo 10) in the *Pd*CysZ sequence and black underlines mark functionally relevant motifs discussed in the text.

Figure 3.

Fig. 3. Structures of *Il*CysZ and *Pf*CysZ, with comparison to *Pd*CysZ a. Ribbon diagram of

the structure of CysZ from *I. loihiensis (Il*CysZ) at 2.3 Å. Protomers of the dimer are colored in

salmon pink and teal blue, arranged in a head-to-tail conformation in the membrane, with helical

hairpins H2-3 and H4-5 labeled for clarity. The dimer interface of *Il*CysZ involves H2-3. b.

Ribbon diagram of the structure of CysZ from *P. fragi (Pf*CysZ*)* at 3.2 Å. The protomers of the

dual topology dimer in the membrane are colored gold and purple, with helical hairpins H2-3 and

H4-5 again labeled. The dimer interface here involves the interaction of helices H4-5 of each

protomer. c. Side and top views of the superposition of the three different protomers from

*Pd*CysZ (green), *Il*CysZ (pink) and *Pf*CysZ (yellow) after aligning H1-H3. d, e. The same dimer

interfaces observed in *Il*CysZ (a) and *Pf*CysZ (b) observed in the hexameric assembly of

*Pd*CysZ, as highlighted in green and blue (d) and green and orange (e). f. Schematic

representation showing how three copies of the dimeric protomers of *Il*CysZ (green and blue,

left) and of *Pf*CysZ (orange and green, center), can coexist in and each recapitulate the

hexameric assembly of *PdCysZ*.

Figure 3 – Figure Supplement 1.

Fig. 3. S1. Size-exclusion chromatography of CysZ shows a mono-disperse elution profile for each of the three species purified - *Pd*CysZ, *Pf*CyZ and *Il*CysZ. All three proteins were solubilized from isolated membrane fractions purified in the presence of 0.2% decyl maltopyranoside (DeM) and exchanged into buffer containing 1% β-octyl glucopyranoside (β-OG) on the size-exclusion column (Superdex 200 10/30 HR). A schematic of the oligomeric states observed in the three crystal structures of CysZ are shown below the elution profiles. Crystallization trials were set-up directly after the size-exclusion chromatography step. A shift of

830   ~1 ml was observed for the elution of *Pd*CysZ (peak at 12.56 ml), with respect to *Pf*CysZ and

831   *Il*CysZ, indicating that the size (volume) and shape of *Pd*CysZ is significantly larger than the

832   other two species. *Il*CysZ elutes at 13.42 ml, 0.5 ml ahead of *Pf*CysZ, which could be explained

833   by the shape of *Il*CysZ, occupying more volume as compared to *Pf*CysZ, with its helices (H4-

834   H5) pointing outward, away from the body of the dimer.

  Figure 3 – Figure Supplement 2.

Fig. 3. S2. Surface electrostatics, hydrophobicity and site-directed fluorescence labeling of cysteine mutants located on H4 of *Il*CysZ. a. A depiction of the surface electrostatic potentials of the *Il*CysZ dimer with negative surface potential represented in red, and positive potential in blue as calculated by APBS. b. Surface of *Il*CysZ rendered by level of hydrophicity, using the Kyte-Doolitle hydrophobicity scale with cyan being most polar, to orange being most hydrophobic, reveals a hydrophobic belt along the center of the dimer, suggesting the orientation of the dimer in the lipid bilayer. c, d. Single cysteine mutants were designed to be located along

the length of helix 4 of CysZ, to gauge the extent of its membrane insertion. The labeled protein

was extracted from the membrane and purified after quenching the labeling reaction.

Fluorescence intensity was measured and quantified by a Tecan fluorescence plate reader at an

excitation of 485nm and emission of 535nm the results of which were plotted on the left, with

error bars representing the standard deviation from the mean for n=3. The results show that of all

the residues on the helix, only R132C (the top-most residue) was accessible to the fluorophore

and hence exposed out of the membrane. d. Locations of the mutated residues on helix H4 are

marked on the model of *Il*CysZ in green. Crosslinking of L161C-A164C cysteine mutant of

*Pf*CysZ and *Il*CysZ exhibits dimer formation. e. A depiction of the surface electrostatic

potentials of the *Pf*CysZ dimer with negative surface potential represented in red, and positive

potential in blue as calculated by APBS, highlights hydrophobic belt marking the orientation of

the *Pf*CysZ dimer in the membrane. f. Membrane orientation of *Pf*CysZ predicted by OPM/PPM

server shows its 31-degree tilt to the perpendicular, with a zoomed in view of the location of the

cysteine mutants at the dimer interface, used for crosslinking of the dimer. g. h. Dimer formation

by the crosslinking of *Pf*CysZ (g) and *Il*CysZ (h) with introduced structure-based cysteine

substitution mutations in the H4-H5 dimer interface. Analogous pairs of cysteine mutants from

each species are shown here: *Pf*CysZ L161C-A164C and *Il*CysZ L156C-Q163C and V157C-

Q163C. Sulfydryl specific Bis-MTS crosslinkers of a certain spacer length (in this case 5.2Å)

were used at 0.5mM for 1hour at room temperature to crosslink the protein. Experiment was

performed on protein in the membrane, as well as on solubilized protein. WT protein (cysteine-

less) was used as a control. Proteins were purified after stopping the reaction, and run on an SDS

PAGE with reducing dye, and stained with Coomassie blue.

    Figure 4.

Fig. 4. Functional characterization of CysZ. a. Time course of $[^{35}S]O_4^{2-}$ uptake (320 μM) by

cells of WT *E. coli K12* (strain BW25113) and by CysZ⁻ (strain JW2406-1) cells (n=3). b. Sulfite

inhibition of sulfate uptake. Radiolabeled sulfate (320 μM $[^{35}S]O_4^{2-}$) uptake is rescued in the

same *cysZ* knockout strain by transformation with an expression vector for *Pd*CysZ whereas the

control cells were transformed with an empty vector. Sulfate uptake is inhibited by the addition

of 10 mM SO_3^{2-} (n=3). c. Competitive binding of labeled sulfate in presence of unlabeled

ligands. The inhibition of labeled sulfate (7 nM $[^{35}S]O_4^{2-}$) binding by non-labeled sulfate or

sulfite is measured by the scintillation proximity assay using purified PdCysZ (n=3). PdCysZ

binds SO_4^{2-} with an EC_{50} of 0.81 ± 0.045 μM and exhibits an IC_{50} of 0.12 ± 0.034 μM for SO_3^{-2}.

d. Radiolabeled sulfate (10 μM $[^{35}S]O_4^{2-}$) accumulation into proteoliposomes reconstituted with

detergent-purified PdCysZ. Uptake into reconstituted proteoliposomes (open squares) is shown

in comparison with control (empty) liposomes (solid discs). Uptake is inhibited by the presence

of 0.1 mM SO_3^{2-} (n=3).

Figure 4 – Figure Supplement 1.

Fig. 4. S1. a. Radiolabeled sulfate (10 μM [^{35}S]O$_4^{2-}$) accumulation is measured from

detergent purified *Il*CysZ and *Pf*CysZ reconstituted in proteoliposomes, in comparison to

the control (empty) liposomes. Sulfate-binding site alanine mutants (*Il*CysZ R27A/R28A and

*Pf*CysZ R25A) display a significantly diminished sulfate uptake capability. Error bars represent

the standard deviation from the mean (SEM), for n=3. b. Effect of uncouplers and ionophores on

the sulfate accumulation by CysZ. *Il*CysZ expressed in the *E. coli* knockout strain shows

unchanged sulfate uptake levels in the presence of the uncouplers CCCP and oligomycin or the

ionophores gramicidin and valinomycin. Sulfate accumulation was measured in triplicate at pH 7

and total accumulation was plotted after a 10 s incubation. Error bars were calculated by standard

deviation from the mean, for n=3.

Figure 4 – Figure Supplement 2.

Fig. 4. S2.   Reconstitution of CysZ in planar lipid bilayers yields sulfite-sensitive

conductances. a. Sample traces of *Il*CysZ under 150 mM symmetrical Na_2SO_4 solutions at the

indicated membrane potentials and pH = 5.4. The lowest trace shows inhibition of activity by 15

μM Na_2SO_3. b. Unitary conductance estimated at 92 ± 12 pS with a reversal potential (E_{rev}) at 0

mV. c. Unitary open probability does not change as a function of membrane potential, and is

stable at an average P_{OPEN} = 0.8. d. Stable bilayer recording at a membrane potential V_m = +50

mV, with no protein reconstituted. An all point current amplitude histogram is plotted on the left

of the trace.   The peak of the normal distribution denotes the baseline zero current. e.

Reconstitution of *Pd*CysZ under 150 mM symmetrical Na_2SO_4 solutions at +50 mV. An all point

current amplitude histogram is plotted on the left of the trace. The high frequency activity shows

a poorly resolved distribution of unitary currents above baseline. f. Inhibition of activity by 5 μM

Na_2SO_3. An all point current amplitude histogram is plotted on the left of the trace. Sulfite

inhibited activity back to baseline levels.

Figure 5.

Fig. 5. Functionally relevant features of the CysZ molecule a. Side-view of a ribbon diagram

of the *Pd*CysZ hexamer, with one chain rainbow-colored from N- (blue) to C-termini (red).

Insets show the sulfate-binding site located at the start of H2a in *Pd*CysZ (left), with conserved

residues G21, L22, R23 and L24 labeled, and the corresponding site in *Il*CysZ (right), with

residues G25, L26, R27, and R28 labeled and showing the SO_4^{-2} ion as bound in the crystal

structure. b. Sulfate uptake by sulfate-binding site mutants of *Pd*CysZ. *E. coli K12* CysZ⁻ cells

transformed with the listed *Pd*CysZ R23 mutants were used to measure $[^{35}S]O_4^{-2}$ uptake (n=3).

Sulfate uptake was abolished for R23A, and rescued to 50% and 40% of the wild type levels for

the R23K and R23Q mutants respectively. c. A top-view of the *Pd*CysZ hexamer, colored as in a.

The inset magnifies the central core of CysZ to show the associated network of hydrogen bonds

(R129-N184, E106-H185), van der Waals interactions (E103-N184, E106-N184, Q176-E134)

and salt bridges (R129-D183, R133-D183, R110-E106) between pairs of highly conserved

residues. Interatomic contacts are shown as purple dotted lines with distances (in Å) marked in

blue. d. Sulfate uptake by central-core mutants of *Pd*CysZ. *E. coli K12* CysZ⁻ cells transformed

with the listed *Pd*CysZ mutants were used to measure $[^{35}S]O_4^{-2}$ uptake (n=3). N98A and R129E

and R129Q showed severely impaired sulfate uptake, whereas more conservative substitutions

such as N98D, Q176E and Q176N had less of a negative effect on function. Y177A and Y177F

do not show any impaired function.

Figure 6.

Fig. 6. Conservation and entrance of putative pore of *Pd*CysZ. a. Ribbon diagram colored by

conservation with residues in maroon being most conserved to cyan being least conserved

(calculated by ConSurf) to highlight the entrance to the putative pore; an asterisk (*) marks the

location of the sulfate-binding site (GLR motif) at top of H2a. b. Same view and coloring scheme

as in a, but now shown in surface representation. c. Electrostatic representation of hexameric

*Pd*CysZ as viewed from the top, with negative surface potential represented in red, and positive

potential in blue as calculated by APBS, with location of the putative pore marked by a dashed

circle. d. Close-up view in electrostatic representation of the putative pore within a *Pd*CysZ

protomer, surface and orientation as in b.

Figure 7.

Fig. 7. Putative ion conductance pathway of CysZ. a. A *Pd*CysZ protomer looking into the

incipient pore entrance. The polypeptide ribbon is oriented as in 6a and colored by level of

sequence conservation (calculated by ConSurf, with maroon being most conserved and cyan

least). The putative pore and ion conduction pathway is shown as a yellow tube of 1 Å diameter,

calculated by PoreWalker. b. Surface representation of the *Pd*CysZ hexamer, as side views, both

semi-transparent (left) and with surface clipped (right, cut surfaces colored in grey) to allow for

the internal visualization of the pathway leading to the central cavity. c. Side view of a *Pd*CysZ

protomer left, viewed as in b and rotated 90° from a. Insets show magnified views of cross-

sections along the pathway: At level 1, the entrance to the pore consists of a narrow constriction

created by a network of highly conserved polar and charged residues (E106, R110, E134, N184,

H185) tightly interacting with one another. At level 2, the pathway broadens and becomes less

charged, but yet polar in nature as lined by conserved asparagine, tyrosine and threonine residues

(N33, T87, N91, N98, Q176, Y177). From level 3, the pathway widens further and ultimately

leads into the large central hydrophobic cavity.

Figure 7 – Figure Supplement 1.

a

b

c

d

974
975

Fig. 7. S1. Plot of the pore diameter of the putative ion conduction pathway. a. *Pd*CysZ b.

*Pf*CysZ, showing a very narrow constriction of under 2 Å in diameter near the entrance of the

pore, which then widens as the ions move past the core of the protomer to the inner central cavity

of the hexameric channel (as calculated by PoreWalker, measured at 3 Å steps along the x-axis)

c. Central hydrophobic cavity of *Pd*CysZ hexamer, cross-section of central cavity of the

hexameric *Pd*CysZ (top view) and d. side view, shows a triangular-shaped cavity with each side

measuring 49.6 Å in length. The cavity is primarily hydrophobic, as depicted and colored with

the Kyte-Doolitle hydrophobicity scale from most polar colored in cyan to most hydrophobic

colored in orange.

References:

Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Davis, I. W., Echols, N., Headd, J. J., Hung, L. W., Kapral, G. J., Grosse-Kunstleve, R. W., McCoy, A. J., Moriarty, N. W., Oeffner, R., Read, R. J., Richardson, D. C., Richardson, J. S., Terwilliger, T. C., & Zwart, P. H. (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 2), 213-221. doi:S0907444909052925 [pii]
10.1107/S0907444909052925

Aguilar-Barajas, E., Diaz-Perez, C., Ramirez-Diaz, M. I., Riveros-Rosas, H., & Cervantes, C. (2011). Bacterial transport of sulfate, molybdate, and related oxyanions. *Biometals, 24*(4), 687-707. doi:10.1007/s10534-011-9421-x

Akabas, M. H., Stauffer, D. A., Xu, M., & Karlin, A. (1992). Acetylcholine receptor channel structure probed in cysteine-substitution mutants. *Science, 258*(5080), 307-310.

Amadi, S. T., Koteiche, H. A., Mishra, S., & McHaourab, H. S. (2010). Structure, dynamics, and substrate-induced conformational changes of the multidrug transporter EmrE in liposomes. *J Biol Chem, 285*(34), 26710-26718. doi:10.1074/jbc.M110.132621

Anishkin, A., Akitake, B., Kamaraju, K., Chiang, C. S., & Sukharev, S. (2010). Hydration properties of mechanosensitive channel pores define the energetics of gating. *J Phys Condens Matter, 22*(45), 454120. doi:10.1088/0953-8984/22/45/454120

Aryal, P., Sansom, M. S., & Tucker, S. J. (2015). Hydrophobic Gating in Ion Channels. *Journal of Molecular Biology, 427*(1), 121-130. doi:10.1016/j.jmb.2014.07.030

Ashkenazy, H., Abadi, S., Martz, E., Chay, O., Mayrose, I., Pupko, T., & Ben-Tal, N. (2016). ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation in macromolecules. *Nucleic Acids Res, 44*(W1), W344-350. doi:10.1093/nar/gkw408

Aslanidis, C., & de Jong, P. J. (1990). Ligation-independent cloning of PCR products (LIC-PCR). *Nucleic Acids Res, 18*(20), 6069-6074.

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K. A., Tomita, M., Wanner, B. L., & Mori, H. (2006). Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol, 2*, 2006 0008. doi:10.1038/msb4100050

Barton, L. L. (2005) *Structural and Functional Relationships in Prokaryotes* (pp. 680-720): Springer.

Bass, R. B., Strop, P., Barclay, M., & Rees, D. C. (2002). Crystal structure of Escherichia coli MscS, a voltage-modulated and mechanosensitive channel. *Science, 298*(5598), 1582-1587. doi:10.1126/science.1077945

Birkner, J. P., Poolman, B., & Kocer, A. (2012). Hydrophobic gating of mechanosensitive channel of large conductance evidenced by single-subunit resolution. *Proc Natl Acad Sci U S A, 109*(32), 12944-12949. doi:10.1073/pnas.1205270109

Britton, P., Boronat, A., Hartley, D. A., Jones-Mortimer, M. C., Kornberg, H. L., & Parra, F. (1983). Phosphotransferase-mediated regulation of carbohydrate utilization in Escherichia coli K12: location of the gsr (tgs) and iex (crr) genes by specialized transduction. *J Gen Microbiol, 129*(2), 349-356.

Buckler, K. J., & Vaughan-Jones, R. D. (1998). Effects of mitochondrial uncouplers on
intracellular calcium, pH and membrane potential in rat carotid body type I cells. *J
Physiol, 513 (Pt 3)*, 819-833.
Chen, Y. H., Hu, L., Punta, M., Bruni, R., Hillerich, B., Kloss, B., Rost, B., Love, J.,
Siegelbaum, S. A., & Hendrickson, W. A. (2010). Homologue structure of the SLAC1
anion channel for closing stomata in leaves. *Nature, 467*(7319), 1074-1080.
doi:10.1038/nature09487
nature09487 [pii]
Cowtan, K. D., & Zhang, K. Y. (1999). Density modification for macromolecular phase
improvement. *Prog Biophys Mol Biol, 72*(3), 245-270.
Czodrowski, P., Dramburg, I., Sotriffer, C. A., & Klebe, G. (2006). Development, validation, and
application of adapted PEOE charges to estimate pKa values of functional groups in
protein-ligand complexes. *Proteins, 65*(2), 424-437. doi:10.1002/prot.21110
Davis, B. D., & Mingioli, E. S. (1950). Mutants of Escherichia coli requiring methionine or
vitamin B12. *J Bacteriol, 60*(1), 17-28.
Dolinsky, T. J., Czodrowski, P., Li, H., Nielsen, J. E., Jensen, J. H., Klebe, G., & Baker, N. A.
(2007). PDB2PQR: expanding and upgrading automated preparation of biomolecular
structures for molecular simulations. *Nucleic Acids Res, 35*(Web Server issue), W522-
525. doi:10.1093/nar/gkm276
Dolinsky, T. J., Nielsen, J. E., McCammon, J. A., & Baker, N. A. (2004). PDB2PQR: an
automated pipeline for the setup of Poisson-Boltzmann electrostatics calculations.
Nucleic Acids Res, 32(Web Server issue), W665-667. doi:10.1093/nar/gkh381
Doyle, D. A., Morais Cabral, J., Pfuetzner, R. A., Kuo, A., Gulbis, J. M., Cohen, S. L., Chait, B.
T., & MacKinnon, R. (1998). The structure of the potassium channel: molecular basis of
K+ conduction and selectivity. *Science, 280*(5360), 69-77.
Emsley, P., Lohkamp, B., Scott, W. G., & Cowtan, K. (2010). Features and development of
Coot. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 4), 486-501. doi:S0907444910007493
[pii]
10.1107/S0907444910007493
Evans, P. (2006). Scaling and assessment of data quality. *Acta Crystallogr D Biol Crystallogr,
62*(Pt 1), 72-82. doi:10.1107/S0907444905036693
Glaser, F., Pupko, T., Paz, I., Bell, R. E., Bechor-Shental, D., Martz, E., & Ben-Tal, N. (2003).
ConSurf: identification of functional regions in proteins by surface-mapping of
phylogenetic information. *Bioinformatics, 19*(1), 163-164.
Hanahan, D. (1983). Studies on transformation of Escherichia coli with plasmids. *Journal of
Molecular Biology, 166*(4), 557-580.
Hryniewicz, M., Sirko, A., Palucha, A., Bock, A., & Hulanicka, D. (1990). Sulfate and
thiosulfate transport in Escherichia coli K-12: identification of a gene encoding a novel
protein involved in thiosulfate binding. *J Bacteriol, 172*(6), 3358-3366.
Kabsch, W. (2010). Xds. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 2), 125-132.
doi:10.1107/S0907444909047337
Kertesz, M. A. (2000). Riding the sulfur cycle--metabolism of sulfonates and sulfate esters in
gram-negative bacteria. *FEMS Microbiol Rev, 24*(2), 135-175.
Kertesz, M. A. (2001). Bacterial transporters for sulfate and organosulfur compounds. *Res
Microbiol, 152*(3-4), 279-290.

Korkhov, V. M., & Tate, C. G. (2009). An emerging consensus for the structure of EmrE. *Acta Crystallogr D Biol Crystallogr, 65*(Pt 2), 186-192. doi:10.1107/S0907444908036640

Kredich, N. M. (1971). Regulation of L-cysteine biosynthesis in Salmonella typhimurium. I. Effects of growth of varying sulfur sources and O-acetyl-L-serine on gene expression. *J Biol Chem, 246*(11), 3474-3484.

Kredich, N. M., Hulanicka, M. D., & Hallquist, S. G. (1979). Synthesis of L-cysteine in Salmonella typhimurium. *Ciba Found Symp*(72), 87-99.

Krissinel, E., & Henrick, K. (2007). Inference of macromolecular assemblies from crystalline state. *Journal of Molecular Biology, 372*(3), 774-797 'Protein interfaces, surfaces and assemblies' service PISA at the European Bioinformatics Institute. (http://www.ebi.ac.uk/pdbe/prot_int/pistart.html). doi:10.1016/j.jmb.2007.05.022

Langer, G., Cohen, S. X., Lamzin, V. S., & Perrakis, A. (2008). Automated macromolecular model building for X-ray crystallography using ARP/wARP version 7. *Nat Protoc, 3*(7), 1171-1179. doi:nprot.2008.91 [pii]

10.1038/nprot.2008.91

Leal-Pinto, E., London, R. D., Knorr, B. A., & Abramson, R. G. (1995). Reconstitution of hepatic uricase in planar lipid bilayer reveals a functional organic anion channel. *J Membr Biol, 146*(2), 123-132.

Liu, Q., Dahmane, T., Zhang, Z., Assur, Z., Brasch, J., Shapiro, L., Mancia, F., & Hendrickson, W. A. (2012). Structures from anomalous diffraction of native biological macromolecules. *Science, 336*(6084), 1033-1037. doi:10.1126/science.1218753

Liu, Q., Zhang, Z., & Hendrickson, W. A. (2011). Multi-crystal anomalous diffraction for low-resolution macromolecular phasing. *Acta Crystallogr D Biol Crystallogr, 67*(Pt 1), 45-59. doi:10.1107/S0907444910046573

Lomize, M. A., Pogozheva, I. D., Joo, H., Mosberg, H. I., & Lomize, A. L. (2012). OPM database and PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res, 40*(Database issue), D370-376. doi:10.1093/nar/gkr703

Loughlin, P., Shelden, M. C., Tierney, M. L., & Howitt, S. M. (2002). Structure and function of a model member of the SulP transporter family. *Cell Biochem Biophys, 36*(2-3), 183-190. doi:CBB:36:2-3:183 [pii]

10.1385/CBB:36:2-3:183

Love, J., Mancia, F., Shapiro, L., Punta, M., Rost, B., Girvin, M., Wang, D. N., Zhou, M., Hunt, J. F., Szyperski, T., Gouaux, E., MacKinnon, R., McDermott, A., Honig, B., Inouye, M., Montelione, G., & Hendrickson, W. A. (2010). The New York Consortium on Membrane Protein Structure (NYCOMPS): a high-throughput platform for structural genomics of integral membrane proteins. *J Struct Funct Genomics, 11*(3), 191-199. doi:10.1007/s10969-010-9094-7

Mancia, F., & Love, J. (2011). High throughput platforms for structural genomics of integral membrane proteins. *Curr Opin Struct Biol, 21*(4), 517-522. doi:10.1016/j.sbi.2011.07.001

Mansilla, M. C., & de Mendoza, D. (2000). The Bacillus subtilis cysP gene encodes a novel sulphate permease related to the inorganic phosphate transporter (Pit) family. *Microbiology, 146 (Pt 4)*, 815-821.

McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., & Read, R. J. (2007). Phaser crystallographic software. *Journal of Applied Crystallography, 40*(Pt 4), 658-674. doi:10.1107/S0021889807021206

Morera, F. J., Vargas, G., Gonzalez, C., Rosenmann, E., & Latorre, R. (2007). Ion-channel reconstitution. *Methods Mol Biol, 400*, 571-585. doi:10.1007/978-1-59745-519-0_38

Mueller, P., Rudin, D. O., Tien, H. T., & Wescott, W. C. (1962). Reconstitution of cell membrane structure in vitro and its transformation into an excitable system. *Nature, 194*, 979-980.

Otwinowski, Z., & Minor, W. (1997). Processing of X-ray Diffraction Data Collected in Oscillation Mode. *Methods in Enzymology, 276*, 307-326.

Pape, T., & Schneider, T. R. (2004). HKL2MAP: a graphical user interface for macromolecular phasing with SHELX programs. *J. Appl. Cryst., 37*, 843-844.

Parra, F., Britton, P., Castle, C., Jones-Mortimer, M. C., & Kornberg, H. L. (1983). Two separate genes involved in sulphate transport in Escherichia coli K12. *J Gen Microbiol, 129*(2), 357-358.

Pellegrini-Calace, M., Maiwald, T., & Thornton, J. M. (2009). PoreWalker: a novel tool for the identification and characterization of channels in transmembrane proteins from their three-dimensional structure. *PLoS Comput Biol, 5*(7), e1000440. doi:10.1371/journal.pcbi.1000440

Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., & Ferrin, T. E. (2004). UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem, 25*(13), 1605-1612. doi:10.1002/jcc.20084

Punta, M., Love, J., Handelman, S., Hunt, J. F., Shapiro, L., Hendrickson, W. A., & Rost, B. (2009). Structural genomics target selection for the New York consortium on membrane protein structure. *J Struct Funct Genomics, 10*(4), 255-268. doi:10.1007/s10969-009-9071-1

Quick, M., & Javitch, J. A. (2007). Monitoring the function of membrane transport proteins in detergent-solubilized form. *Proc Natl Acad Sci U S A, 104*(9), 3603-3608. doi:0609573104 [pii]

10.1073/pnas.0609573104

Rapp, M., Granseth, E., Seppala, S., & von Heijne, G. (2006). Identification and evolution of dual-topology membrane proteins. *Nat Struct Mol Biol, 13*(2), 112-116. doi:10.1038/nsmb1057

Rigaud, J. L., Pitard, B., & Levy, D. (1995). Reconstitution of membrane proteins into liposomes: application to energy-transducing membrane proteins. *Biochim Biophys Acta, 1231*(3), 223-246.

Schrodinger, LLC. (2010). *The PyMOL Molecular Graphics System, Version 1.3r1*.

Sheldrick, G. M. (2010). Experimental phasing with SHELXC/D/E: combining chain tracing with density modification. *Acta Crystallogr D Biol Crystallogr, 66*(Pt 4), 479-485. doi:S0907444909038360 [pii]

10.1107/S0907444909038360

Sirko, A., Zatyka, M., Sadowy, E., & Hulanicka, D. (1995). Sulfate and thiosulfate transport in Escherichia coli K-12: evidence for a functional overlapping of sulfate- and thiosulfate-binding proteins. *J Bacteriol, 177*(14), 4134-4136.

Stein, N. (2008). CHAINSAW: a program for mutating pdb files used as templates in molecular replacement. *Journal of Applied Crystallography, 41*, 641-643. doi:10.1107/S0021889808006985

Stockbridge, R. B., Kolmakova-Partensky, L., Shane, T., Koide, A., Koide, S., Miller, C., &
Newstead, S. (2015). Crystal structures of a double-barrelled fluoride ion channel.
Nature, 525(7570), 548-551. doi:10.1038/nature14981
Stockbridge, R. B., Robertson, J. L., Kolmakova-Partensky, L., & Miller, C. (2013). A family of
fluoride-specific ion channels with dual-topology architecture. *Elife, 2*, e01084.
doi:10.7554/eLife.01084
Winn, M. D., Ballard, C. C., Cowtan, K. D., Dodson, E. J., Emsley, P., Evans, P. R., Keegan, R.
M., Krissinel, E. B., Leslie, A. G., McCoy, A., McNicholas, S. J., Murshudov, G. N.,
Pannu, N. S., Potterton, E. A., Powell, H. R., Read, R. J., Vagin, A., & Wilson, K. S.
(2011). Overview of the CCP4 suite and current developments. *Acta Crystallogr D Biol*
Crystallogr, 67(Pt 4), 235-242. doi:10.1107/S0907444910045749
Yang, T., Liu, Q., Kloss, B., Bruni, R., Kalathur, R. C., Guo, Y., Kloppmann, E., Rost, B.,
Colecraft, H. M., & Hendrickson, W. A. (2014). Structure and selectivity in bestrophin
ion channels. *Science, 346*(6207), 355-359. doi:10.1126/science.1259723
Ye, L., Jia, Z., Jung, T., & Maloney, P. C. (2001). Topology of OxlT, the oxalate transporter of
Oxalobacter formigenes, determined by site-directed fluorescence labeling. *J Bacteriol,*
183(8), 2490-2496. doi:10.1128/JB.183.8.2490-2496.2001
Zhang, L., Jiang, W., Nan, J., Almqvist, J., & Huang, Y. (2014). The Escherichia coli CysZ is a
pH dependent sulfate transporter that can be inhibited by sulfite. *Biochim Biophys Acta,*
1838(7), 1809-1816. doi:10.1016/j.bbamem.2014.03.003
1184