Distinct effects of tubulin isotype mutations on neurite growth in *Caenorhabditis elegans*

Chaogu Zheng[*,†], Margarete Diaz-Cuadros[*,†], Ken C.Q. Nguyen[‡], David H. Hall[‡], and Martin Chalfie[*]

[*] Department of Biological Sciences, Columbia University, New York, New York, 10027, U.S.A.
[‡] Department of Neuroscience, Albert Einstein College of Medicine, Bronx, NY 10461, U.S.A.
[†] These authors contributed equally to the work.

Correspondence:

Martin Chalfie
Department of Biological Sciences
1012 Fairchild, MC#2446
Columbia University
1212 Amsterdam Avenue
New York, NY 10027
Phone: 212-854-8870
Fax: 212-865-8246
Email: mc21@columbia.edu.

Table of Content (TOC) Highlight Summary

Different tubulin isotypes perform different functions in the regulation of MT structure and

neurite growth, and missense mutations of tubulin genes have three types of distinct effects on

MT stability and neurite growth. One α-tubulin isotype appears to induce relative instability due

to the lack of potential post-translational modification sites.

Abstract

Tubulins, the building block of microtubules (MTs), play a critical role in both supporting and regulating neurite growth. Eukaryotic genomes contain multiple tubulin isotypes, and their missense mutations cause a range of neurodevelopmental defects. Using the *C. elegans* touch receptor neurons, we analyzed the effects of 67 tubulin missense mutations on neurite growth. Three types of mutations emerged: 1) loss-of-function mutations, which cause mild defects in neurite growth; 2) antimorphic mutations, which map to the GTP binding site and intradimer and interdimer interfaces, significantly reduce MT stability, and cause severe neurite growth defects; and 3) neomorphic mutations, which map to the exterior surface, increase MT stability, and cause ectopic neurite growth. Structure-function analysis reveals a causal relationship between tubulin structure and MT stability. This stability affects neuronal morphogenesis. As part of this analysis, we engineered several disease-associated human tubulin mutations into *C. elegans* genes and examined their impact on neuronal development at the cellular level. We also discovered an α-tubulin (TBA-7) that appears to destabilize MTs. Loss of TBA-7 led to the formation of hyperstable MTs and the generation of ectopic neurites; the lack of potential sites for polyamination and polyglutamination on TBA-7 may be responsible for this destabilization.

Introduction

Microtubules (MTs) play important roles in many aspects of neurite development, being involved in the formation, extension, guidance, and maintenance of neurites (reviewed in Dent *et al.*, 2011; Prokop, 2013; Sainath and Gallo, 2015). MTs constantly explore the growth cone periphery until they are captured by stabilized actin filaments or membrane receptors enriched at the side of the growth cone responding to a guidance cue (Tanaka *et al.*, 1995; Challacombe *et al.*, 1996; Schaefer *et al.*, 2008; Qu *et al.*, 2013). This capture transiently stabilizes MTs and enables MT elongation in the direction that the growth cone has turned. In addition to providing physical support for the neurite growth that follows the changes in actin dynamics, MTs also play an instructive role in neurite guidance. Since local application of the MT-stabilizing drug paclitaxel (also known as taxol) induced growth cone attraction, of the MT-destabilizing drug nocodazole induced repulsion, and of the MT-depolymerizing drug colchicine resulted in branching (Bray *et al.*, 1978; Buck and Zheng, 2002), signals that act by altering MT stability appear to directly initiate growth cone turning. Indeed, the guidance molecule Wnt can induce growth cone remodeling by changing the organization of MT structure through the inactivation of MT-plus end binding protein Adenomatous Polyposis Coli (Purro *et al.*, 2008). These results indicate that the regulation of MT dynamics is crucial for neurite growth and guidance, but how MT stability is controlled locally at specific sites of the growth cone and globally in the entire neuron is not well understood.

As the building blocks of MTs, α- and β-tubulins are crucial determinants of MT stability. In fact, eukaryotic genomes contain multiple tubulin genes encoding different isotypes (Sullivan, 1988; McKean *et al.*, 2001) that are expressed in spatially and temporally distinct patterns (Leandro-Garcia *et al.*, 2010) and that confer specific dynamic properties on MTs in tubulin

polymerization assays *in vitro* (Hoyle and Raff, 1990; Panda *et al.*, 1994; Pucciarelli *et al.*, 2012).

These observations support the "multi-tubulin hypothesis," which proposes that distinct tubulin

isotypes impart specific properties onto MTs, so they can perform particular cellular functions

(Fulton and Simpson, 1976; Cleveland, 1987). Moreover, tubulins also undergo a range of

complex post-translational modification (reviewed by Song and Brady, 2015), which affect the

dynamics of MTs and their interaction with other proteins and can add to the complexity of the

tubulin code. In neurons specifically, stable axonal MTs are more detyrosinated, acetylated, and

glutamylated, whereas the dynamic MTs in the growth cone are more tyrosinated (Liao and

Gundersen, 1998; Konishi and Setou, 2009). Tubulin isotypes and modifications at the C-

terminal tail also control the velocity and processivity of MT motor proteins and microtubule

depolymerization rates *in vitro* (Sirajuddin *et al.*, 2014). In turn MT motors can also regulate the

organization of MTs (Verhey and Gaertig, 2007; Yu *et al.*, 2015).

The clinical importance of tubulin genes in the development of the nervous system is

seen in the finding that α- and β- tubulin mutations in people are associated with microcephaly,

lissencephaly, pachygyria, and other cortical malformations, as well as a range of axon guidance

defects, including agenesis or hypoplasia of the corpus callosum, internal capsule, commissural

fibers, and corticospinal tracts (Tischfield *et al.*, 2011). Three summaries of human mutations

(Bahi-Buisson *et al.*, 2014; Liu and Dwyer, 2014; Chakraborti *et al.*, 2016) identify 60 point

mutations in α-tubulin genes (51 in TUBA1A, 1 in TUBA3E, and 8 in TUBA4A), one splicing-

affecting intronic deletion in TUBA8, 48 point mutations in β-tubulin genes (2 in TUBB2A, 24

in TUBB2B, 19 in TUBB3, and 3 in TUBB5), and one exonic deletion in TUBB2B that produce

neurological disorders in heterozygous carriers (Table S1); all mutations, except for the TUBA8

deletion, appear to act as dominant-negative mutations. Although mutated residues are found

throughout the molecules, many are found in region predicted to mediate either GTP binding,

heterodimer stability, inter-dimer interaction, or association with motor proteins and other MAPs

(Tischfield *et al.*, 2011). Despite some *in vitro* studies on the effects of the mutations on tubulin

folding, heterodimer assembly, and MT growth (Jaglin *et al.*, 2009; Tian *et al.*, 2010; Tischfield

et al., 2010), very few *in vivo* studies can systematically test how these different tubulin

mutations impact axon guidance and extension in living organisms. Moreover, the complexity of

the mammalian nerve system makes the analysis of the developmental consequences of these

tubulin mutations very difficult.

Here we use the morphologically simple and well-defined touch receptor neurons (TRNs)

in the nematode *Caenorhabditis elegans* to model the effects of tubulin mutations on neurite

growth. By analyzing a large collection of missense mutations in several tubulin genes, we found

that these mutations caused three morphologically distinct defects in TRN neurite outgrowth: 1)

the shortening of all TRN neurites; 2) the specific shortening of posteriorly directed neurites; and

3) the production of ectopic posteriorly directed neurites. The structural location of the mutated

residue correlated with the resulting phenotype. Many tubulin mutations characterized in our

study affect the same amino acid residue or region as the disease-causing mutations in humans.

We generated several such human mutations in *C. elegans* tubulin genes through genome editing

and found that they also caused distinct neurite growth defects that fall into the above categories.

Thus, our system may be used to understand the different effects of the clinically identified

tubulin mutations and to facilitate their classification. Moreover, we found that null mutations in

two α-tubulin genes led to very different phenotypes, supporting the hypothesis that tubulin

isotypes perform specific and often non-overlapping roles in neurite development.

Results

Neurite morphology and MT organization in the TRNs

The six mechanosensory TRNs (ALML/R, PLML/R, AVM, and PVM) in *C. elegans* are a useful model to study axonal outgrowth and guidance because of their well-defined morphology (Chalfie and Sulston, 1981). The ALM and PLM neurons are two pairs of embryonically derived, bilaterally symmetric cells, whereas the AVM and PVM neurons arise from postembryonic lineages. All six neurons have a long anteriorly-directed neurite (AN); in addition, the two PLM neurons have a posteriorly-directed neurite (PN), making them bipolar. Except in PVM, the ANs branch at their distal ends; we refer to this branch as the synaptic branch.

Unlike most of *C. elegans* the cells, which have only a few (~5 per cross section in ventral cord neurons) 11-protofilament (11-p) MTs, TRNs contain a great number (~31) of large diameter (15-protofilament, 15-p) MTs that assemble into bundles that fill the neurites (Chalfie and Thomson, 1979; Savage *et al.*, 1994). The abundance and prominence of the MTs in the TRNs permits easy observation of alterations in MT structure and organization by electron microscopy.

The *C. elegans* genome contains nine α-tubulin genes (*mec-12*, *tba-1*, *tba-2*, and *tba-4* through *tba-9*) and six β-tubulin genes (*ben-1*, *mec-7*, *tbb-1*, *tbb-2*, *tbb-4*, and *tbb-6*). Previous studies found that the α-tubulin MEC-12 and the β-tubulin MEC-7 are expressed specifically at high levels in the TRNs (Hamelin *et al.*, 1992; Mitani *et al.*, 1993) and are required for the mechanosensory functions of the TRNs (Chalfie and Au, 1989; Savage *et al.*, 1994). TRNs also express the ubiquitously present α-tubulin, TBA-1 and TBA-2, and β-tubulin, TBB-1 and TBB-2, but their loss had no effect on the development or function of TRNs (Fukushige *et al.*, 1993,

1995; Lockhead *et al.*, 2016). Here, we identify an additional α-tubulin TBA-6, which is also

expressed in TRNs and functions to prevent excessive neurite growth.

In the following sections, we focus on MEC-7, MEC-12, and TBA-7. We first describe

three types of *mec-7* missense mutations and their distinct effects on MT structure, neurite

growth, and neuronal function, and then discuss the phenotypes of similar mutations in *mec-12*.

These latter phenotypes are generally weaker than the corresponding *mec-7* mutations. To allow

for a comparison of the *mec-7* and *mec-12* phenotypes, however, we have organized the first four

figures by the type of data (Figure 1 for electron microscopy, Figure 2 for process outgrowth,

Figure 3 for structural analysis, and Figure 4 for TRN activity). Finally, we describe the effect of

the loss of *tba-7*.

Deletion of *mec-7*/β-tubulin led to the loss of 15-p MTs and defects in posteriorly directed

neurite growth

Using electron microscopy (EM), we found that *mec-7(ok2152)* knockout animals had on

average 6 MTs in a cross section of the ALM neurite compared to 31 MTs in the wild type

animals (Figure 1). The *ok2152* mutation deletes the first four exons (Figure S1A) and is thus is a

likely null allele. The diameter of MTs in *mec-7(ok2152)* animals was much smaller (16.6 nm)

than that of the wild type (21.5 nm), and tannic acid staining indicated that the 15-p MTs in

TRNs were replaced by smaller 11-p MTs in the absence of MEC-7 (Figure 1). These results

suggest that the abundance of MTs and the formation of large diameter 15-p MTs requires MEC-

7, which is consistent with our previous observations on *mec-7(e1506)* animals (the *e1506*

mutation alters the start codon and results in no detectable *mec-7* mRNA; Chalfie and Thomson,

1982; Savage *et al.*, 1994).

We next examined how *mec-7(ok2152)* mutation affected TRN morphogenesis. The *mec-*

7 deletion allele *mec-7* only slightly affected the growth of anteriorly-directed neurites of PLM

neurons but markedly shortened the posteriorly-directed neurites (Figure 2B). This result

suggests that PLM-AN and PLM-PN, which arise at the same time in the embryo, have different

tubulin requirements. Normally the PLM-AN extends anteriorly past the vulva to within 50 μm

of ALM cell body. Lockhead *et al.* (2016) reported that the gap between the end of PLM and

ALM cell body was wider in animals with the presumed null *mec-7(u241)* due to the shortening

of the PLM-AN. We found a similar modest widening of the gap in *mec-7(ok2152)* animals, but

92% of the PLM-ANs still extended beyond the vulva (Figure 2B and D). Moreover, no

shortening of the ALM-AN was seen. A more penetrant and striking defect, however, was the

loss of the synaptic branch in 78% of PLM-ANs in *mec-7(ok2152)* animals (Figure 2A and S2;

the synaptic branch normally arises posterior to the vulva). The ALM synaptic branch was not

affected. *mec-7* nonsense alleles *u156* and *u440* produced similar morphological defects (Figure

S1B and data not shown). These results suggest that the smaller, generic 11-p MTs in *mec-7*

knockout mutants are, in general, capable of supporting the growth of most TRN neurites, but the

normal extension of PLM-PN and the elaboration of the posterior synaptic branch require the

special 15-p MTs.

The neurites of wild-type TRNs have many 15-p MTs associated into a bundle. This

bundle may provide stronger support for neurite growth than the few 11-p MTs found in *mec-7*

deletion mutants. If so, the requirement for the 15-p MTs in the PLM-PN may be due to a higher

sensitivity to changes in microtubule stability there than in the PLM-AN. In fact, treatment with

the MT-destabilizing drug colchicine (1mM), which specifically removes most of the TRN MTs

(Chalfie and Thomson, 1982), led to a similar shortening of the PLM-PN, phenocopying the

mec-7 null mutants (Figure S3A).

Antimorphic mutations in *mec-7* led to severe neurite growth defects

We next examined 55 *mec-7* alleles that represent 48 different missense mutations (Table

1) and found three categories of TRN morphological defects. First, 12 mutations caused neurite

growth defects similar to the knockout allele and were thus classified as loss-of-function (*lf*)

mutations. Data from *u1020* (G34S) animals is shown in Figure 2 as an additional *lf* example.

When mapped to the structure of a bovine αβ tubulin dimer (Figure 3; Nogales *et al.*, 1998), *lf*

mutations were found throughout the molecule and presumably affected protein folding. Second,

19 mutations caused severe shortening of all TRN neurites; the ALM-AN did not reach the

pharynx, the PLM-AN terminated before reaching the PVM cell body and the PLM-PN was

significantly shortened (Figure 2B and D). Because of the dominant-negative nature of these

alleles (all, except *u430*, were either dominant or semi-dominant), we classified them as

antimorphic (*anti*) gain-of-function mutations. We also identified 4 weak *mec-7(anti)* mutants

(Table 1), in which ALM-AN did not extend beyond the nerve ring and PLM-AN did not reach

the vulva. Third, 6 mutations led to the growth of an extra posteriorly-directed neurite in ALM

neurons, as well as the overextension of the PLM-PN. Among the six alleles, *u278* (C303Y)

produced the strongest phenotype; the ectopic posterior neurite of ALM neurons often extended

posteriorly to the PLM cell body (Figure 2C). Because this phenotype was novel and different

from the phenotype of either *mec-7 lf* or *anti* mutations, we classified these alleles as neomorphic

(*neo*) gain-of-function mutations. Except two semi-dominant ones (*u278* and *ky852*), most of the

neo alleles were recessive. Because of the antagonism between anteriorly and posteriorly

directed outgrowth (Zheng *et al.*, 2016), some *neo* alleles, especially *u1017*, caused mild

shortening of the ANs, i.e. PLM-AN terminated slightly posterior to the vulva (Figure 3C).

EM studies of the *mec-7(anti)* allele, *e1343* (P171L), found that the mutants contained

very few (2.8 ± 0.5 in a cross section) MTs (Chalfie and Thomson, 1982), which suggests that

the *anti* mutations block MT polymerization and thereby cause the severe neurite outgrowth

defects. In fact, when mapped to the tubulin structure, amino acid residues mutated in the *anti*

alleles are clustered in three regions: 1) the GTP/GDP binding pocket, 2) the intradimer interface,

and 3) the lateral or longitudinal interdimer interface. All of which are important for tubulin

polymerization. For example, *u430* (A97V), *u911* (P171S), *u957* (P171L), and *u262* (N226Y)

mutations alter amino acids in direct contact with the GTP molecule, and *u48* (S176F) and *u449*

(V179A) change amino acids in the B5 (the fifth β-strand)-to-H5 (the fifth α-helix) loop, which

is crucial for forming the GTP/GDP binding pocket (Figure 3C). Two other mutations *u445*

(M300V) and *u98* (M300T) substitute M300, which is located in a loop near the GTP binding

site and may participate in positioning H7 (the seventh α-helix) with its critical N226, and thus

indirectly affect GTP binding.

The second group of mutated residues in *mec-7(anti)* alleles is located in the intradimer

interface, where β-tubulin makes contact with α-tubulin to form the heterodimer (red residues in

Figure 3D). *u283* (P243L), *u958* (G244S), *n434* (N247I), and *u162* (D249N) mutations all affect

the residues on the H7-to-H8 loop, which interacts extensively with residues on H1 and H2 of α-

tubulin. Two other dominant mutations *gk286000* (A314V) and *u955* (A352T) changed residues

that located on B8 and B9, respectively, which are physically adjacent to the H7-to-H8 loop at

the interface. In addition, two prolines mutations, *u910* (P357L) and *u956* (P358L) possibly

disrupted the B9-to-B10 loop (a.a. 356-361) critical for the positioning of B9.

The third group of *mec-7(anti)* alleles include *e1527* (V286D), *u18* (A393T), and

gk285997 (A393V), which affect residues involved in the lateral (V286) and longitudinal (A393)

interdimer interaction between one α/β-tubulin heterodimer and the neighboring one on the MTs.

These mutations may interfere with the interaction between tubulin dimers.

Overall, the MEC-7/β-tubulin *anti* mutants likely act in a dominant-negative manner by

forming poisonous α/β dimers, whose incorporation into MTs could terminate MT

polymerization and induce instability. The mutated β-tubulin either cannot properly bind to GTP

(the first group), form misshaped α/β heterodimers that block the growing end of MTs (the

second group) or disrupt the stacking of tubulin dimers (the third group). Therefore, changes in

tubulin structure led to compromised MT elongation, which caused severe defects in neurite

outgrowth, highlighting the importance of functional MTs in neuronal morphogenesis.

Neomorphic mutations in *mec-7* led to ectopic neurite growth by increasing MT stability

The *mec-7(neo)* mutations appear to induce the growth of ectopic ALM-PNs by forming

hyperstable MTs, as first suggested by Kirszenblat *et al.* (2013). Supporting this hypothesis,

pharmacological stabilization of MTs with paclitaxel induced the growth of ALM-PN, although

at a low frequency, and destabilizing MTs with colchicine can partially suppress this ectopic

growth in *mec-7(ky852 neo)* mutants (Kirszenblat *et al.*, 2013) and in *mec-7(u278 neo)* and *mec-*

7(u1017 neo) animals (Figure S3A). We found that *mec-7(neo)* animals had increased resistance

to colchicine compared to the wild-type animals. More of this MT-destabilizing drug was needed

to produce the same level of reduction in both TRN function (touch sensitivity) and PLM-PN

length in *neo* mutants than in wild-type animals (Figure S3B-C). Moreover, these mutants also

showed increased tubulin acetylation (Figure 4C), an indication of stable MTs (Song and Brady,

2015).

The greater MT stability and higher resistance to destabilizing reagents in the *neo*

mutants could be the result of either reduced dynamics of individual MTs or more stable bundles.

MTs in wild-type TRNs showed very little dynamics, since Hsu *et al.* (2014) found and we

confirmed that the GFP labeled MT plus-end binding protein EBP-2/EB2 barely moved in the

wild-type TRNs. Thus, the already very high stability of individual MTs is unlikely to increase

greatly in the *neo* mutants. However, the MT bundles in the TRNs of *neo* mutants have changed

and so may be the cause of the drug-resistance in these animals. By quantifying EM data

previously obtained (Savage *et al.*, 1994), we found that *mec-7(u278 neo)* and *mec-7(u170 neo)*

mutants kept the large diameter MTs (slightly larger than in wild-type cells), but had fewer of

them (Figure 1B). Importantly, the MT bundles in the mutants are more closely spaced than the

wild type and are not surrounded by the electron-lucent region seen in the wild type (Savage *et*

al., 1994). We quantified these differences by showing that the distance between two closest

center points of MTs is smaller in *mec-7(neo)* mutants compared to the wild type and that MTs

occupied a bigger proportion of the cross-sectional area of the neurite in the mutants (Figure 1B).

In fact, the TRN neurite in these mutants is about 50% thinner than the wild type animals. The

observation that tightly packed 15-p MT bundles filled up most of the space in the TRN neurites

is consistent with increased MT stability, which presumably allowed the excessive neurite

growth towards the posterior, overcoming the normal inhibition of this growth in the wild-type

ALM neurons.

The majority of the altered residues in the *mec-7(neo)* alleles are located on the exterior

surface of MTs, where they may interfere with the binding of motor proteins or MAPs to the

MTs and therefore cause changes in MT stability. Examples include *u1017* (L377F) and *u170*

(E407K), which are located on the H11 and H12 helices, respectively, *gk286003* (R162Q) on the

H4-to-B5 loop, and *u278* (C303Y) on the H9-to-B8 loop are also exposed on the MT surface

(Figure 3E). Interestingly, although *u278* (C303Y) produced the strongest *neo* phenotype, C303S

mutation in allele *gk286001* did not lead to similar excessive growth (Table 1), suggesting that

tyrosine's bulky phenolic ring may be responsible for the phenotype. In contrast, the *neo* alleles

gk895768 (T149I) and *ky852* (P220S) affect residues on the inside the MTs and may be

important for the folding of the protein.

Loss-of-function, antimophic, and neomorphic mutations of *mec-12*/α-tubulin caused

similar but weaker phenotypes than the *mec-7* mutations

We next examined the contribution of MEC-12/α-tubulin to MT organization and neurite

growth in the TRNs. *mec-12(tm5083)* allele deletes part of exon 3 and intron 3 (Figure S1B) and

is likely to be a *lf* mutation. Those deletion mutants had much fewer (~7) MTs in a cross section

of ALM neurites than the wild type animals, but the MTs had the same diameter as the 15-p MTs

in the wild type (Figure 1B). Moreover, we observed no defects in TRN outgrowth in the *mec-*

12(tm5083 lf) mutants (Figure 2) and confirmed the results using three CRISPR/Cas9-induced

frameshift mutations (*u1026*, *u1027*, and *u1028*; Figure S1B and data not shown). Therefore,

although the abundance of MTs requires MEC-12, the formation of 15-p MTs does not require

MEC-12, and the reduction of MT numbers in *mec-12(lf)* mutants did not affect neurite

development. Because *C. elegans* genome contains nine α-tubulin genes, other tubulin isotypes

may compensate for the loss of MEC-12. Although TBA-1 and TBA-2 are expressed in the

TRNs, *tba-1(lf)*; *mec-12(lf)* and *tba-2(lf); mec-12(lf)* double mutants had very mild defects in

neurite growth, only the slight shortening of PLM-AN (Lockhead *et al.*, 2016). In this study, we

found another α-tubulin TBA-7, which is also expressed and functions in TRNs (see below);

however, *mec-12; tba-7* double mutants were still capable of forming and growing normal TRN

neurites, suggesting further genetic redundancy.

The EM results of *mec-12(lf)* mutants were unexpected, because the *mec-12[(e1607*

(G144S)] allele, which had been regarded as the null allele (Bounoutas *et al.*, 2009a; Bounoutas

et al., 2011; Hsu *et al.*, 2014), had very few (~2) MTs, all of which had the small diameter

(Chalfie and Au, 1989). We repeated the EM studies using the newly isolated *mec-12(u1021)*

allele that carries the same G144S mutation and found the ALM neurites had ~2.7 MTs in a cross

section and 15% of the 72 sections examined had no MTs; the diameter of MTs was 17.9 nm

(Figure 1B). We found that *e1607* and several other missense alleles (Table 1) were in fact

antimorphic (*anti*) gain-of-function alleles, which caused the significant shortening of PLM-PN

(Figure 2B and E). Like the *mec-7(anti)*, the five *mec-12(anti)* mutations also mapped to the

GTP binding pocket [*u76* (D69N), *u950* (S140F), and *u1021* (G144S); labeled in red in Figure

3B], the intradimer interface [*u1016* (E97K); Figure 3D], or the interdimer interface [*u1019*

(G354E)]. However, unlike the *mec-7(anti)* alleles, *mec-12(anti)* are mostly recessive at 20 °C,

except for *u1019*, and did not cause strong, general defects in TRN neurite outgrowth, but rather

specific shortening of PLM-PN similar to the *mec-7(lf)* mutants.

One possible explanation for this phenotype is that MEC-12/α-tubulin preferentially or

exclusively binds to MEC-7/β-tubulin, and the MEC-12(*anti*) proteins sequester and disable

MEC-7 but no other β-tubulin isotypes. As a result, both *mec-12(anti)* and *mec-7(lf)* removes

MEC-7 from the pool of free tubulins, leading to the loss of the 15-p MTs that require MEC-7

and the shortening of PLM-PN. Consistent with this hypothesis, *mec-7(lf); mec-12(anti)* double

mutants did not show additive effects with regards to the neurite growth defects compared to

either single mutant (Figure 2E).

Four of the 19 *mec-12* missense mutations (Table 1) were *neo* alleles [*gk170196* (P32S),

gk915672 (E196K), *u917* (V260I), *gk515972* (V323I)]. These alleles produced a similar but less

severe phenotypes than the *mec-7(neo)* alleles. Specifically, the ectopic ALM-PN was shorter in

the *mec-12(neo)* mutants than in the *mec-7(neo)* mutants (Figure 2F). This observation supports

the notion that MEC-12 may be less important than MEC-7 in regulating MT stability and

subsequent neurite growth. E196 and V260 are located on loops on the exterior surface, whereas

P32 and V323 may be involved in protein folding (Figure 3F).

The production of the ectopic ALM-PN requires both *mec-7* and *mec-12*. The deletion of

mec-12 suppressed the growth of ALM-PN in *mec-7(neo)* mutants, and the deletion of *mec-7*

similarly suppressed the effect of *mec-12(neo)* alleles (Figure S3A). These results suggest that

the tubulin *neo* mutations induce excessive neurite growth by altering the properties of mostly

MEC-7/MEC-12 heterodimers and not other α/β heterodimers.

mec-7 and mec-12 mutations also affect TRN functions, vesicle transport, and protein

expression

In addition to neurite growth, MTs are important for many other functions in TRNs

(Bounoutas *et al.*, 2009a; Hsu *et al.*, 2014). Deletion of either *mec-12* or *mec-7* resulted in touch

insensitivity, defects in the localization of pre-synaptic vesicles, and severe loss of MT

acetylation, a mark for stable MTs (Figure 4 and S4A). Since MEC-12 is the only α-tubulin

isotype that contains the acetylation site (lysine 40) in *C. elegans*, the TRN-specific 15-p MTs

are likely the only MTs that can be acetylated. The loss of *mec-7* and *mec-12* also induced a

global reduction in protein levels through a mechanism dependent on the MAPK pathway dual

leucine zipper-bearing kinase DLK-1 (Bounoutas *et al.*, 2011). Interestingly, mutations in *dlk-1*

restored the normal protein expression in *mec-7* and *mec-12* null mutants but failed to rescue the

PLM-PN growth defects in *mec-7(lf)* animals (Figure 4D and S4B). Therefore, this PLM-PN

defect is most likely a direct effect of the loss of MEC-7 and not the result of secondary changes

in protein levels.

Compared to the *mec-7(lf)* mutants, the *mec-7(anti)* mutations caused stronger defects not

only in neurite outgrowth, but also in mechanosensation, synaptic vesicle transport, tubulin

acetylation, and protein expression levels (Figure 4 and S4B). Those phenotypes in *mec-12(anti)*

mutants were more severe than the phenotypes in the *mec-12(lf)* alleles and were comparable to

those produced by *mec-7(lf)* mutations but not as strong as the *mec-7(anti)* mutations (Figure 4).

The *mec-7* and *mec-12 neo* alleles also caused touch insensitivity, synaptic vesicle transport

defects, and reduced TRN protein levels, as found in the *lf* mutants; but the *neo* mutants had

markedly increased MT acetylation (Figure 4). Moreover, the fact that both impaired (in *anti*

mutants) and excessive (in *neo* mutants) neurite growth can occur in conjunction with other TRN

defects suggests that the role of MTs in regulating neurite development is genetically separable

from other MT functions.

Furthermore, several *mec-12* partial *lf* mutations affected only a subset of MT functions.

mec-12(e1605) animals carrying the H192Y missense mutation were defective in

mechanosensation but had retained normal 15-p MT structures, tubulin acetylation, axonal

transport, protein levels, and neurite growth patterns (Figure S5; Bounoutas *et al.*, 2009a).

Similarly, *mec-12* mutants carrying the *u63* (E415K) or *gm379* (G416E) alleles were partially

touch-insensitive but kept the large dimeter MTs; however, these animals showed mistargeting

of synaptic vesicles presumably because alterations in the EEGE (amino acid 414-417) motif

increased affinity with the motor dynein (Hsu *et al.*, 2014). *u63* and *gm379* mutations also

caused a slight decrease in tubulin acetylation and a partial reduction in protein production;

gm379 allele led to very mild defects in the growth of PLM-PN (Figure S5). These mutations

may be useful in understanding some specific aspects of MT functions.

Modeling the effects of tubulin mutations using the TRN neurites

Our genetic analysis shows that missense mutations in tubulin genes can have distinct

effects on MT stability and neurite growth pattern, and those different phenotypes appear to

correlate with the positions of the altered residues in the tubulin structure. Such structure-

function analyses have been carried out for the yeast α-tubulin TUB1 and the Drosophila testis-

specific β-tubulin βTub85D (Fackenthal *et al.*, 1995; Richards *et al.*, 2000) but not for tubulins

expressed in the nervous system. Moreover, recent discovery of over a hundred missense

mutations in tubulin genes in patients with a range of neurological disorders prompted us to

examine the cellular impact of specific tubulin mutations on neurons. Combining the

convenience of genome editing in *C. elegans* and the ease of observing TRN morphology as the

readout of MT stability and neurite growth, we could systematically study the effects of those

human tubulin mutations. To provide a proof of concept, we generated five clinically observed

β-tubulin mutations (S172P, P173L, A302T, P380S, and E410K) in *mec-7* through

CRISPR/Cas9-mediated gene editing (Materials and Methods; Figure 5A) and found that these

mutations indeed caused distinct phenotypes on TRN morphogenesis.

Heterozygous S172P and P173L mutations in human TUBB3 result in microcephaly,

polymicrogyria, cortical dysplasia, and agenesis of the corpus callosum, all of which are defects

in neuronal migration and axon growth (Jaglin *et al.*, 2009; Poirier *et al.*, 2010; Bahi-Buisson *et*

al., 2014). *In vitro* studies found that tubulin heterodimer containing the β-tubulin S172P mutant

could not be incorporated into MTs (Jaglin *et al.*, 2009), indicating that the mutated protein is

nonfunctional. Consistent with these findings, the *u1056* (S172P) mutation of *mec-7* led to a

recessive phenotype similar to *mec-7(lf)* in the TRNs, which showed significantly shortened

PLM-PN and slightly shortened PLM-AN with branching defects (Figure 5B). Surprisingly, the

u1057 (P173L) allele did not cause any TRN morphological or functional defects (Figure 5B).

The difference between the mutant phenotypes in humans and worms is puzzling, since both

S172 and P173 are located on the GTP binding B5-to-H5 loop (Figure 3C), which is identical in

C. elegans MEC-7 and human TUBB3 (Figure S6). Perhaps the differences arise from

differences in MT structure and organization: the *C. elegans* TRNs have 15-p MTs that are

bundled, whereas human neurons have 13-p MTs that are not bundled. In fact, when 15-p MTs

were converted to 13-p MTs and the bundle was disrupted in *mec-17* (α-tubulin acetyl-

transferase) mutants (Topalidou *et al.*, 2012), both MEC-7 S172P and P173L mutations led to

decreased MT stability and the loss of ALM-PN (Figure 5B and S7). This result suggests that

P173L mutation indeed compromises MT stability in TRNs under sensitized conditions.

Heterozygous A302T and R380C mutations in human TUBB3, which affect residues on

the external surface of the MTs, were associated with moderate congenital fibrosis of the

extraocular muscles 3 (CEFOM3), anterior commissure hypoplasia, and corpus callosum

hypoplasia. Mutation of A302 and R380 in yeast led to the formation of highly stable, benomyl-

resistant MTs (Tischfield *et al.*, 2010). Consistent with these observations, we found that the

same mutations of MEC-7 produced a *neo* phenotype (the generation of an ectopic ALM-PN;

Figure 5C).

TUBB3 E410K mutation was also found in patients with severe CEFOM3 and hypoplasia

of anterior commissure and corpus callosum, but these patients also suffer from facial weakness

and progressive axonal sensorimotor polyneuropathy (Tischfield *et al.*, 2010). In yeast, MTs

containing β-tubulin with the E410K substitution were less stable and less resistant to benomyl

and had markedly decreased plus-end accumulation of kinesin-like motor proteins, compared to

MTs with A302T and R380C mutations (Tischfield *et al.*, 2010). In *C. elegans*, MEC-7(E410K)

produced a distinct neurite growth phenotype: ALM-AN and PLM-AN, but not PLM-PN were

significantly shortened, which indicates reduced MT stability; but the PLM-PN was not affected,

making the phenotype also different from that of the *mec-7(anti)* alleles, which cause general

defects in MT polymerization (Figure 5D). Since E410 is also exposed to the exterior of MTs

like A302 and R380 (Figure 3E), these results suggest that the cellular impact of the missense

mutations would depend on the specific amino acid change instead of the location of the affected

residues alone.

The loss of *tba-7/α*-tubulin also caused excessive posterior neurite growth

In addition to MEC-7 and MEC-12, we identified a second α-tubulin gene *tba-7* that also

regulates neurite growth in the TRNs. In the course of a mutagenesis for mutants with abnormal

TRNs (Supplemental Results), we isolated a missense *tba-7* allele *u1015* (G92D), which caused

the growth of an ectopic ALM posterior neurite, resembling the phenotype of *mec-7* and *mec-12*

neo alleles (Figure 6A and B). This *tba-7* allele is likely a *lf* allele, because it failed to

complement with *gk787939* (Q230*), which is presumably a null and produced a similar

phenotype as *u1015* (Figure S1C). We also examined mutants of other tubulin isotypes, and did

not find morphological defects in the TRNs (we did not test for redundancy among the genes;

Table S2).

tba-7 was expressed in the TRNs, and the excessive growth phenotype of *tba-7(u1015)*

could be rescued by expressing *tba-7(+)* from the TRN-specific *mec-17* promoter, indicating that

TBA-7 acts cell-autonomously in the TRNs (Figure S8). *tba-7*(*lf)* mutants did not exhibit touch

insensitivity, synaptic vesicle mistargeting, or reduced protein levels, suggesting that the general

function of TRN MTs was not affected and that the primary function of TBA-7 is to regulate

neurite growth (Figure S9). We hypothesized that the loss of TBA-7 led to the formation of

hyperstable MTs, similar to the *mec-7* and *mec-12 neo* mutations. Supporting this hypothesis,

electron microscopy studies revealed that, like *mec-7(neo)* mutants, *tba-7(u1015)* animals

retained the large diameter 15-p MTs and have closely packed MT bundles, which occupied a

larger-than-normal area of a neurite cross-section (Figure 1). *tba-7(lf)* mutants also retained

normal tubulin acetylation levels (Figure S9C) and had increased resistance to colchicine (Figure

S3), which confirmed the presence of stable MTs. Treatment with 1 mM colchicine fully

suppressed in the growth of ALM-PN in *tba-7(u1015)* animals, as did either *lf* or *anti* mutations

in *mec-7* or *mec-12* (Figure 6B).

The amino acids sequences of α-tubulin TBA-7 and MEC-12 are 82% identical and 94%

similar; they mainly differ in a N-terminal region and the C-terminal tail (Figure 6C). Domain

swapping experiments showed that changing L31 of TBA-7 to Q found in MEC-12 or replacing

the C-terminal DANDNGD (a.a. 435-441 in TBA-7) sequence with MEDNGEEG (a.a. 440-447

in MEC-12) partially impaired the function of TBA-7 proteins to restrict excessive neurite

growth; and TBA-7 proteins with both L31Q mutation and the C-terminal replacement

completely lost the ability to rescue the *tba-7(lf)* phenotype (Figure 6D). Q31 in MEC-12 is a

potential site for polyamination (Song *et al.*, 2013) and E445 of the GEE motif (a.a. 444-446 in

MEC-12, but absent in TBA-7) is a target for polyglutamination (Edde *et al.*, 1990). Post-

translational modifications at these two sites could increase the stability of neuronal MTs (Song

and Brady, 2015). Interestingly, replacing the a.a. 35- 47 region of TBA-7 with the sequence

from MEC-12 that contains lysine 40 (the site for α-tubulin acetylation) did not affect the TBA-7

function (Figure 6D), suggesting that the absence of this MT acetylation site in TBA-7 was not

responsible for its activity in preventing ectopic neurite growth.

Since TBA-7 lacks some modification sites (e.g. Q31 and E445) that could stabilize MTs,

our results suggest that TBA-7 may be a MT-destabilizing tubulin isotype, as compared to MEC-

12. In wild-type animals, TBA-7 incorporation into the 15-p MTs with MEC-7 and MEC-12 may

reduce MT stability; when TBA-7 is not available, more MEC-12 and possibly other α-tubulin

isotypes replace TBA-7 and this replacement led to the formation of hyperstable MTs.

The *tba-7(u1015 lf); mec-7(u1017 neo)* double mutant had an ALM-PN that was similar

in length to that in *mec-7(u1017)* single mutants (Figure 6A) but additionally produced up to 4

and 5 short ectopic neurites sprouting from the ALM and PLM cell bodies, respectively; this

phenotype was rarely observed in *mec-7(u1017)* animals (Figure 6E and F). The additive effect

in the double mutant suggest that the loss of TBA-7 and the MEC-7 *neo* mutations act in parallel

to elevate MT stability.

Mutations in MT-destabilizing Kinesin 13 affect TRN neurite growth

Since motor proteins and MAPs regulate MT dynamics (Akhmanova and Steinmetz,

2015), we expected mutations in MAP genes to cause similar phenotype to those of some of the

tubulin mutations. Indeed, Ghosh-Roy *et al.* (2012) found and we have confirmed that the loss of

the MT-depolymerizing kinesin *klp-7* induced the growth of an ectopic posterior neurite in ALM

neurons (Figure 6G). KLP-7 belongs to the kinesin-13 family of catastrophe factors that bind to

MT plus-ends and promotes depolymerization, thus generating dynamic microtubules (Han *et al.*,

2015). Importantly, we found that, similar to the tubulin *neo* alleles, the growth of the ectopic

ALM-PN in *klp-7(lf)* mutants was suppressed by the loss of *mec-7* or *mec-12*, but the *klp-7*

phenotype was not enhanced by *mec-7(neo)* mutations. These results suggest that the mutated

MEC-7 proteins may render MTs hyperstable by reducing their interaction with KLP-7 or by

making the MTs insensitive to the action of KLP-7.

Discussion

Neuronal morphogenesis requires optimal MT stability

MTs both support and regulate neurite growth. As the building block of MTs, α- and β-

tubulins determine the structural properties of MTs and mediate their interaction with motor

proteins and MAPs. As a result, any alteration in the tubulin proteins can potentially lead to

changes in MT stability, which then affects neurite formation, guidance, and extension. In this

study, we analyzed the effects of 67 tubulin missense mutations on neurite development in *C.*

elegans touch receptor neurons. Based on the phenotypes, we categorized these mutations into

three classes: loss-of-function (*lf*), antimorphic (*anti*), and neomorphic (*neo*) mutations. 1) *lf*

mutations cause only mild neurite growth defects by reducing MT stability moderately; 2) *anti*

mutations block MT polymerization and cause severe neurite growth defects; 3) *neo* mutations

cause excessive neurite growth by inducing hyperstable MTs (Figure 7). The directionality of the

excessive neurite in ALM reflects its potential to grow towards the posterior. In fact, about 30%

of the adult ALM neurons have a very short (less than one cell body length) posterior protrusion,

and manipulation of cellular signaling (e.g. activating the small GTPase Rac) can induce the

production of ALM-PN (Zheng *et al.*, 2016). Thus, the ectopic growth of ALM-PN may serve as

an indication for altered cytoskeletal organization and dynamics.

The correlation between the amount of neurite growth and the stability of MTs suggests

that MT structure and dynamics are major determinants of neurite growth. Given the great

number of proteins involved in both positively and negatively regulating MT assembly and

disassembly (Mimori-Kiyosue, 2011), our results support the hypothesis that an optimal level of

MT stability is required for proper neuronal morphogenesis. The finding that negative regulators

of MT stability, such as the α-tubulin isotype TBA-7 and the MT-destabilizing kinesin-13 KLP-

7, prevent excessive TRN neurite growth further supports this hypothesis.

Structure-function analysis of tubulin mutations

The large number of missense mutations analyzed in this study allowed us to perform a

structure-function analysis of neuronal tubulin proteins in living animals. Mapping the altered

amino acid residues onto the *Bos taurus* tubulin α/β heterodimer structure 1JFF (Nogales *et al.*,

1998), we found that *lf* mutations mostly affect residues located in the interior of the structure

and possibly disrupt protein folding. *anti* mutations affected residues that participate in GTP

binding, intradimer interaction, or interdimer interactions. *neo* mutations mostly altered residues

on the exterior surface and probably perturb the association of motor proteins and MAPs with

MTs (Figure 3 and 7).

Our findings are consistent with similar structure-function relationship studies conducted

in yeast α-tubulin *TUB1* and β-tubulin *TUB2* and Drosophila testis-specific *β2*-tubulin (Reijo *et*

al., 1994; Fackenthal *et al.*, 1995; Richards *et al.*, 2000) and provide another model to study the

properties of tubulin proteins. For example, yeast *TUB1* mutations that caused supersensitivity to

the MT-destabilizing drug benomyl mainly altered amino acids involved in GTP binding (e.g.

D70) and intradimer (e.g. E98) and interdimer interaction; the change of equivalent residues in

MEC-12 (D69 and E97) were identified in *mec-12(anti)* mutants. Similarly, some Drosophila *β2*-

tubulin mutations (S25L and G96E) that made MTs less stable (Fackenthal *et al.*, 1995) affected

the same or the adjacent residue in *mec-7(anti)* alleles [*u319* (S25F) and *u430* (A97V)].

Moreover, yeast *TUB1* mutations conferring benomyl-resistance mostly affected the residues on

the outer surface of the MTs (Richards *et al.*, 2000) as did the *mec-7(neo)* mutations similarly

that were resistant to colchicine. Thus, comparative studies using the three different organisms

can help identify key residues in the tubulin structure responsible for various MT functions.

Modeling the effects of neuronal tubulin mutations

We have used the *C. elegans* TRNs as a model to study neuronal MTs, which are much

more stable and have different types of post-translational modification than the MTs in dividing

cells (Baas *et al.*, 2016). Clinical studies in the past five years identified more than 100 missense

mutations in tubulin genes that cause a wide spectrum of neurodevelopmental disorders, such as

microcephaly, lissencephaly, polymicrogyria, and cortical malformation, along with severe

defects in axon guidance and growth, including the complete or partial absence of corpus

callosum, defects in commissural fiber tracts, and degeneration of motor and sensory axons

(Tischfield *et al.*, 2011; Chakraborti *et al.*, 2016). Patients carrying different tubulin mutations

often display distinct symptoms, but the complexity of the human nervous system makes it

difficult to analyze the impact of specific mutations at the cellular level.

Combining CRISPR/Cas9-mediated genome editing with *in vivo* examination of TRN

morphology, we were able to create several disease-causing human β-tubulin mutations in the *C.*

elegans mec-7 gene and examined their effects on neurite growth at a single neuron resolution.

Our phenotypic evaluation was largely consistent with previous characterization of those

mutations, suggesting that the TRN system could be instrumental to model the effects of human

tubulin mutations identified in the clinic.

In fact, many disease-causing tubulin mutations alter amino acids that are physically

adjacent to or located in the same region as the ones affected in our mutants, suggesting that they

may affect MT stability in similar ways. By mapping 51 TUBA1A, 24 TUBB2B, and 19

TUBB3B mutations that were clinically identified onto the tubulin structure, we could correlate

the location of some of the affected residues with the resulting clinical manifestation (Table S1).

For example, 7/8 (87.5%) mutations that alter residues involved in GTP binding caused complete

agenesis of corpus callosum, an indication of severe defects in axonal growth, whereas only 5/23

(21.7%) changes of residues in the interior of the structure and 10/32 (31.2%) mutations of

residues involved in MAP binding did so (Table S1).

Position of the mutated residues alone, however, could not determine the effects of the

missense mutation. For example, although the β-tubulin B5-to-H5 loop (from V170 to V179)

contacts GTP, and four strong *mec-7(anti)* mutations (P171S, P171L, S176F, and V179A)

mapped to this loop, the S172P mutation led to a *lf* phenotype and P173L mutation caused no

defects in TRNs. One possible explanation is that S172 and P173 are further away from the GTP

binding site, thus their mutations interfere less with microtubule function.

The identity of the residue replacing the wild type one is also important. For instance,

u278 (C303Y) was the strongest *mec-7(neo)* mutant, but *gk286001* (C303S) had no effect at all;

since both tyrosine and serine have the polar hydroxyl group, the bulky phenyl group of the

tyrosine may be responsible for disrupting some MT functions. Moreover, the *u1058* (A302T)

mutation also caused a strong *mec-7(neo)* phenotype. A302 and C303 on the H9-to-B8 loop are

exposed to the exterior of MTs but are not located on the major landing surface (H11 and H12)

for motor protein and MAPs (Nogales *et al.*, 1998). Our results suggest that A302 and C303 may

represent a previously unrecognized site of interaction between the MTs and their associated

proteins. Even for residues located on H11 and H12, the effects of their mutation can be different.

u1059 (R380S) is a *mec-7(neo)* mutation, whereas *u1060* (E410K) allele produced an *anti*-like

phenotype, suggesting that the impact of the missense mutation depends on how the interaction

of MTs with the motor proteins or MAP is affected by the amino acid change.

MT-destabilizing tubulin isotype and "multi-tubulin hypothesis"

The presence of multiple tubulin genes in eukaryotic genomes, their subtle differences in

the protein sequences, and their functional differences led to the "multi-tubulin hypothesis." In

multicellular organism, the tubulins may have distinct expression patterns, *e.g.* Drosophila *β2*-

tubulin is only expressed in the male germline (Kemphues *et al.*, 1980); human TUBB3B is

highly enriched in the nervous system, whereas TUBB2B is expressed in many tissues (Sullivan

and Cleveland, 1986). In single-cell organisms like *Tetrahymena*, different β-tubulin isotypes are

spatially separated; some are only used to make the nuclear MTs of the mitotic apparatus, and

some are only detected in the MTs of somatic cilia and basal bodies (Pucciarelli *et al.*, 2012).

Our studies, however, found that two distinct α-tubulin isotypes (MEC-12 and TBA-7) could

have different functions when both incorporated into the same MTs in the same neurons; MEC-

12 promotes stability and TBA-7 promotes instability.

The abundance of large diameter 15-p MTs in the TRNs requires MEC-12, but the role of

TBA-7 is to increase MT dynamics and to prevent the ectopic growth of posteriorly directed

neurites in ALMs. Thus, TBA-7 is a MT-destabilizing tubulin isotype, and this destabilizing

activity depends on the absence of potential polyamination and polyglutamination sites that

could increase MT stability. Our results are consistent with early *in vitro* experiments suggesting

that TUBB3 (also known as β_{III}) also promotes MT dynamics; removal of TUBB3 from the brain

extract resulted in a tubulin mixture that assembled much more rapidly than the unfractionated

control (Banerjee *et al.*, 1990), and MTs assembled from the purified $\alpha\beta_{III}$ heterodimers were

considerably more dynamic than MTs made from the $\alpha\beta_{II}$ and $\alpha\beta_{IV}$ dimers (Panda *et al.*, 1994).

However, the long-standing question is whether some tubulin isotypes do destabilize MTs *in vivo*.

Our findings suggest the answer is yes. A balanced incorporation of multiple tubulin isotypes

into the same MT structure is critical to generate MTs with the optimal stability.

The absence of post-translational modification sites in tubulin isotypes may lead to

functional diversity. In addition to TBA-7, among the other eight *C. elegans* α-tubulin isotypes,

TBA-8 and TBA-5 also lack the Q31 for polyamination, and TBA-6 lacks the E445 for

polyglutamination; all nine human α-tubulins have Q31, but TUBAL3 has a short C-terminal tail

and does not contain potential polyglutamination sites (Figure S10). The incorporation of those

tubulin isotypes may be a general regulatory mechanism to control MT stability. Despite the

functional differences between MEC-12 and TBA-7, the fact that both *mec-12(lf)* mutants and

mec-12(lf); tba-7(lf) double mutants retained the ability to grow out a normal amount of neurites

suggests also genetic redundancy among the α-tubulins in supporting neurite growth.

Different contributions of tubulin isotypes to MT dynamics were also observed in *C.*

elegans embryogenesis (Honda *et al.*, 2017). The spindle MTs required two β-tubulins (TBB-1

and TBB-2), but TBB-2 was incorporated into the MTs twice as much as TBB-1; the loss of

TBB-2 caused a dramatic decrease in MT growth rate and an increase in catastrophe frequency,

leading to highly unstable MTs, whereas the loss of TBB-1 only slightly reduced growth rate.

These data support the "multi-tubulin concept" that different tubulin isotypes have distinct

functions in the same cells.

Materials and Methods

Strains and genetic screens

C. elegans wild type (N2) and mutant strains were maintained at 20 °C as previously described

(Brenner, 1974). *mec-7* alleles *u910, u911, u955, u956, u957, u958, u1017,* and *u1020, mec-12*

alleles *u917, u950, u1016, u1019,* and *u1021,* and *tba-7(u1015)* were isolated by visually

screening for mutants with TRN differentiation defects using TU4069, which carries *uIs134*

[mec-17p::RFP] for the visualization of the TRNs, as the starter strain and ethyl

methanesulfonate as the mutagen (Brenner, 1974). Mutants were outcrossed with wild type, and

the phenotype-causing mutations were identified as previously described by whole-genome

resequencing (Zheng *et al.*, 2013) or by complementation tests with reference alleles. A detailed

description of the screen can be found in the supplemental results.

mec-7 alleles *e1343*, *e1505*, *e1527*, *e1522*, *n434*, *u10*, *u18*, *u22*, *u48*, *u58*, *u98*, *u127*,

u129, *u162*, *u170*, *u223*, *u225*, *u278*, *u234*, *u249*, *u262*, *u275*, *u283*, *u305*, *u319*, *u428*, *u429*,

u430, *u433*, *u449*, and *u445*, and *mec-12* alleles *e1605*, *e1607*, *u50*, *u63*, *u76*, and *u241* were

previously isolated (Chalfie and Sulston, 1981; Chalfie and Au, 1989; Savage *et al.*, 1994) and

re-examined in this study.

u1056 to *u1060* alleles were created through CRISPR/Cas9-mediated homologous

recombination (Dickinson *et al.*, 2013). Guide RNAs were designed according to the target

sequence closest to the desired codon change (Figure 5A). Recombination templates were

created by cloning the *mec-7* coding region and 466 bp 3'UTR sequence into the SalI and

BamHI sites of pBlueScript II SK (+) vector and then introducing the desired missense mutation

(red in Figure 5A), along with synonymous mutations (blue in Figure 5A) that change the guide

RNA target sites, using the Q5 Site-Directed Mutagenesis Kit from New England Biolabs (NEB;

Ipswich, MA). pDD162 constructs expressing Cas9 and specific guide RNAs were injected

together with the corresponding recombination template and marker *myo-2p::RFP* into TU4069

animals. F1s expressing RFP in the muscle were singled out and genotyped to identify

heterozygotes with successful edits, and F2s carrying homozygous mutations were then isolated

and examined for TRN morphology.

The *gk* alleles of *mec-7* and *mec-12* listed in Table 1 and *tba-7(gk787939)* were generated

in the million mutation project (Thompson *et al.*, 2013). These *gk* alleles, as well as *mec-*

17(ok2019), *dlk-1(ju476)*, and *dlk-1(km12)* were obtained from the Caenorhabditis Genetics

Center, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

klp-7(tm2143), and *mec-12(tm5083)* were generated by the National Bioresource Project of

Japan, and *mec-12(gm379)* was kindly provided by Dr. Chun-Liang Pan at the National Taiwan

University.

Additional *mec-12* null alleles (*u1026*, *u1027*, and *u1028*) were generated by

CRISPR/Cas9-mediated genome editing targeting 5'-GAAGTAATTTCGATTCACATCGG-3'

in exon 2 of *mec-12* as described above (Dickinson *et al.*, 2013). Frameshift-causing mutations

were identified by sequencing the *mec-12* locus (Figure S2).

Constructs and Transgenes

A *tba-7::GFP* reporter (TU#1632) was made by cloning a 1.2 kb *tba-7* promoter and the

entire coding region from the genomic DNA to wild-type genomic DNA into the Gateway

pDONR221 P4-P1r vector; the resulting entry vector, together with the pENTR-GFP, pENTR-

unc-54-3'UTR, and destination vector pDEST-R4-R3 were used in the LR reaction to create the

final expression vectors. *mec-17p::tba-7(+)* (TU#1629) was created by cloning a 1.9 kb *mec-17*

promoter and the *tba-7* coding sequence into pDONR221 P4-P1r and pDONR221 vectors,

respectively, and assembling these entry vectors with pENTR-*unc-54*-3'UTR and the destination

vector. Details about the Gateway cloning method by Life Technologies can be found at

www.invitrogen.com/site/us/en/home/Products-andServices/Applications/Cloning/Gateway-

Cloning.html. *tba-7(+)* locus, including the 1.2 kb promoter, coding region, and a 911-bp 3'UTR,

was cloned using SalI and NotI into pBlueScript II SK(+) vector; NEB Q5 site-directed

mutagenesis kit was then used to generate constructs expressing the TBA-7 variants shown in

Figure 6E.

Transgenes *uIs31[mec-17p::GFP]* III, *uIs115[mec-17p::RFP]* IV, and *uIs134[mec-*

17p::RFP] V were used to visualize TRN morphology (Zheng *et al.*, 2015). *jsIs821[mec-*

7p::GFP::RAB-3] was used to assess synaptic vesicle localization (Bounoutas *et al.*, 2009b).

Electron microscopy

We re-analyzed previously collected cross-section images of ALM neurites from wild

type animals (N501, N631, N933, and N934 from the Hall lab collection) and from *mec-7(neo)*

mutants *u170* and *u278* (Savage *et al.*, 1994). The wild type animals had been fixed by either

chemical immersion fixation without tannic acid (samples N501, N631; Hall, 1995) or by high-

pressure freezing/freeze substitution (HPF/FS) including tannic acid (samples N933, N934;

Topalidou *et al.*, 2012). The *mec-7(neo)* alleles had also been fixed by chemical immersion,

without tannic acid.

For this study, we fixed *mec-7(ok2152 lf)*, *mec-12(tm5083 lf)*, and *tba-7(u1015 lf)* adults

using an HPF/FS protocol that included a first fixation in 0.5% glutaraldehyde + 0.1% tannic

acid and a second fix in 2% osmium tetroxide + 0.1% uranyl acetate (UAc), both at -90 °C,

followed by staining in 1% UAc in acetone at 0 °C. We also fixed adults of *mec-12(anti)* mutants,

u950, *u76*, and *u1021*, using HPF/FS using a similar protocol, but without tannic acid. Eighty-

nanometer transverse sections were collected at multiple positions along the ALM anterior

neurite and post-stained with uranium acetate and lead citrate. A Philips CM10 electron

microscope with an attached Morada digital camera (Olympus) was used to acquire the images.

For each animal, we counted the number of MTs in the ALMs and measured the MT

diameter on sections collected from at least five different positions along the ALM-AN; data

from ALML and ALMR were combined. For each strain, we examined sections from at three

different animals. For wild-type animals, *tba-7(lf)*, and *mec-7(neo)* mutants, we used at least ten

sections to measure the distance between two closest MT centers and the proportion of MT-

occupied area to the cross-sectional area of the neurite.

We noticed that our earlier studies found that wild-type TRN MTs had much larger

diameters (29.6 ± 0.4 nm; Chalfie and Thomson, 1979) than we find here (21.5 ± 1.6 nm).

Fixation method could not explain the difference, because we obtained similar results from N501

and N631 (fixed by chemical immersion) and N933 and N934 (fixed by HPF/FS); data obtained

using both methods showed smaller MT diameters than the measurements from earlier studies,

which used chemical immersion fixation. Although we could not explain this discrepancy, we

are very confident in the current calibration of the electron microscope.

Phenotype Scoring and Statistical Analysis

We measured the length of TRN neurites in at least 30 fourth stage larvae or young adults

grown at 20° C, except where otherwise stated. Relative length of posteriorly directed neurites

was calculated by dividing the neurite length by the diameter of the cell body. Defects in TRN

anterior neurite length were assessed by counting the percentage of cells whose neurites failed to

reach the vulva (for PLM) or the posterior pharyngeal bulb (for ALM); at least 50 cells were

examined for each strain.

Fluorescence intensity of the RFP expressed from the *uIs134* transgene in the ALM cell

body was used to measure TRN protein levels; intensity was calculated after manual background

subtraction using ImageJ as previously described (Chen and Chalfie, 2015). Synaptic vesicle

localization was quantified by calculating the percentage of animals in each of the four

phenotypical categories: 1) normal localization to the PLM synaptic branch; 2) reduced intensity

of the synaptic marker at the synapses as partial defect; 3) complete loss of normal localization to

the synapse as complete defect; 4) localization to the PLM-PN as mistargeting. At least 50 adult

animals were examined.

Acetylated α-tubulin staining using antibody [6-11B-1] (Abcam, Cambridge, MA) was

performed and analyzed as previously described (Topalidou *et al.*, 2012). For colchicine

treatment, animals were grown in standard NGM agar plates containing different concentrations

(from 0.06 mM to 2 mM) of colchicine before phenotypic analysis as described before

(Bounoutas *et al.*, 2009a). Similarly, 100 nM paclitaxel was added to the NGM agar plates, L1

animals were placed onto the plates, and young adults were examined for TRN morphology.

PyMOL (Schrodinger, 2015) was used to view the structure of α/β tubulin dimer (1jff.pdb;

Nogales *et al.*, 1998) and to label affected residues in *mec-7* and *mec-12* mutants,.

For statistical analysis, ANOVA and the post hoc Dunnett's test (comparison of each of

many treatments with a single control) or Tukey-Kramer test (studentized range test for all

pairwise comparison) were performed using the GraphPad Prism (version 5.00 for Windows,

GraphPad Software, La Jolla California USA, www.graphpad.com) to identify significant

difference between the mutants and wild type animals. Student's *t* test was used to find

significant difference between two samples in paired comparisons. Single and double asterisks

indicated $p < 0.05$ and $p < 0.01$, respectively. The χ^2 test was used for the categorical data to find

significant difference between different strains.

Acknowledgements
792
793 We thank Dan Dickinson, Chun-Liang Pan, and Andrew Chisholm for sharing reagents.

794 This work was supported by NIH Grant GM30997 (to M.C.) and OD 010943 (to D.H.H.). Core

795   facilities for electron microscopy were supported by NICHD P30 HD71593 for the RFK-IDDRC

796   at Albert Einstein College of Medicine.

References
Akhmanova, A., and Steinmetz, M.O. (2015). Control of microtubule organization and dynamics: two
ends in the limelight. Nat Rev Mol Cell Biol *16*, 711-726.
Baas, P.W., Rao, A.N., Matamoros, A.J., and Leo, L. (2016). Stability properties of neuronal microtubules.
Cytoskeleton (Hoboken) *73*, 442-460.
Bahi-Buisson, N., Poirier, K., Fourniol, F., Saillour, Y., Valence, S., Lebrun, N., Hully, M., Bianco, C.F.,
Boddaert, N., Elie, C., Lascelles, K., Souville, I., Consortium, L.I.-T., Beldjord, C., and Chelly, J. (2014). The
wide spectrum of tubulinopathies: what are the key features for the diagnosis? Brain *137*, 1676-1700.
Banerjee, A., Roach, M.C., Trcka, P., and Luduena, R.F. (1990). Increased microtubule assembly in bovine
brain tubulin lacking the type III isotype of beta-tubulin. The Journal of biological chemistry *265*, 1794-
1799.
Bounoutas, A., Kratz, J., Emtage, L., Ma, C., Nguyen, K.C., and Chalfie, M. (2011). Microtubule
depolymerization in Caenorhabditis elegans touch receptor neurons reduces gene expression through a
p38 MAPK pathway. Proceedings of the National Academy of Sciences of the United States of America
108, 3982-3987.
Bounoutas, A., O'Hagan, R., and Chalfie, M. (2009a). The multipurpose 15-protofilament microtubules in
C. elegans have specific roles in mechanosensation. Current biology : CB *19*, 1362-1367.
Bounoutas, A., Zheng, Q., Nonet, M.L., and Chalfie, M. (2009b). mec-15 encodes an F-box protein
required for touch receptor neuron mechanosensation, synapse formation and development. Genetics
183, 607-617, 601SI-604SI.
Bray, D., Thomas, C., and Shaw, G. (1978). Growth cone formation in cultures of sensory neurons.
Proceedings of the National Academy of Sciences of the United States of America *75*, 5226-5229.
Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-94.
Buck, K.B., and Zheng, J.Q. (2002). Growth cone turning induced by direct local modification of
microtubule dynamics. The Journal of neuroscience : the official journal of the Society for Neuroscience
22, 9358-9367.
Chakraborti, S., Natarajan, K., Curiel, J., Janke, C., and Liu, J. (2016). The emerging role of the tubulin
code: From the tubulin molecule to neuronal function and disease. Cytoskeleton (Hoboken) *73*, 521-550.
Chalfie, M., and Au, M. (1989). Genetic control of differentiation of the Caenorhabditis elegans touch
receptor neurons. Science *243*, 1027-1033.
Chalfie, M., and Sulston, J. (1981). Developmental genetics of the mechanosensory neurons of
Caenorhabditis elegans. Developmental biology *82*, 358-370.
Chalfie, M., and Thomson, J.N. (1979). Organization of neuronal microtubules in the nematode
Caenorhabditis elegans. J Cell Biol *82*, 278-289.
Chalfie, M., and Thomson, J.N. (1982). Structural and functional diversity in the neuronal microtubules
of Caenorhabditis elegans. J Cell Biol *93*, 15-23.
Challacombe, J.F., Snow, D.M., and Letourneau, P.C. (1996). Actin filament bundles are required for
microtubule reorientation during growth cone turning to avoid an inhibitory guidance cue. J Cell Sci *109*
(Pt 8), 2031-2040.
Chen, X., and Chalfie, M. (2015). Regulation of mechanosensation in C. elegans through ubiquitination of
the MEC-4 mechanotransduction channel. The Journal of neuroscience : the official journal of the
Society for Neuroscience *35*, 2200-2212.
Cleveland, D.W. (1987). The multitubulin hypothesis revisited: what have we learned? J Cell Biol *104*,
381-383.

Dent, E.W., Gupton, S.L., and Gertler, F.B. (2011). The growth cone cytoskeleton in axon outgrowth and
guidance. Cold Spring Harb Perspect Biol *3*.

Dickinson, D.J., Ward, J.D., Reiner, D.J., and Goldstein, B. (2013). Engineering the Caenorhabditis elegans
genome using Cas9-triggered homologous recombination. Nature methods *10*, 1028-1034.

Edde, B., Rossier, J., Le Caer, J.P., Desbruyeres, E., Gros, F., and Denoulet, P. (1990). Posttranslational
glutamylation of alpha-tubulin. Science *247*, 83-85.

Fackenthal, J.D., Hutchens, J.A., Turner, F.R., and Raff, E.C. (1995). Structural analysis of mutations in the
Drosophila beta 2-tubulin isoform reveals regions in the beta-tubulin molecular required for general and
for tissue-specific microtubule functions. Genetics *139*, 267-286.

Fukushige, T., Yasuda, H., and Siddiqui, S.S. (1993). Molecular cloning and developmental expression of
the alpha-2 tubulin gene of Caenorhabditis elegans. J Mol Biol *234*, 1290-1300.

Fukushige, T., Yasuda, H., and Siddiqui, S.S. (1995). Selective expression of the tba-1 alpha tubulin gene
in a set of mechanosensory and motor neurons during the development of Caenorhabditis elegans.
Biochim Biophys Acta *1261*, 401-416.

Fulton, C., and Simpson, F.A. (1976). Selective synthesis and utilization of flagellar tubulin: the multi-
tubulin hypothesis. In: Cell Motility, vol. 3, eds. R. Goldman, T. Pollard, and J. Rosenbaum, New York:
Cold Spring Harbor Publications, 987-1005.

Ghosh-Roy, A., Goncharov, A., Jin, Y., and Chisholm, A.D. (2012). Kinesin-13 and tubulin posttranslational
modifications regulate microtubule growth in axon regeneration. Developmental cell *23*, 716-728.

Hall, D.H. (1995). Electron microscopy and three-dimensional image reconstruction. Methods Cell Biol
48, 395-436.

Hamelin, M., Scott, I.M., Way, J.C., and Culotti, J.G. (1992). The mec-7 beta-tubulin gene of
Caenorhabditis elegans is expressed primarily in the touch receptor neurons. The EMBO journal *11*,
2885-2893.

Han, X., Adames, K., Sykes, E.M., and Srayko, M. (2015). The KLP-7 Residue S546 Is a Putative Aurora
Kinase Site Required for Microtubule Regulation at the Centrosome in C. elegans. PLoS One *10*,
e0132593.

Honda, Y., Tsuchiya, K., Sumiyoshi, E., Haruta, N., and Sugimoto, A. (2017). Tubulin isotype substitution
revealed that isotype composition modulates microtubule dynamics in C. elegans embryos. J Cell Sci.

Hoyle, H.D., and Raff, E.C. (1990). Two Drosophila beta tubulin isoforms are not functionally equivalent. J
Cell Biol *111*, 1009-1026.

Hsu, J.M., Chen, C.H., Chen, Y.C., McDonald, K.L., Gurling, M., Lee, A., Garriga, G., and Pan, C.L. (2014).
Genetic analysis of a novel tubulin mutation that redirects synaptic vesicle targeting and causes neurite
degeneration in C. elegans. PLoS genetics *10*, e1004715.

Jaglin, X.H., Poirier, K., Saillour, Y., Buhler, E., Tian, G., Bahi-Buisson, N., Fallet-Bianco, C., Phan-Dinh-Tuy,
F., Kong, X.P., Bomont, P., Castelnau-Ptakhine, L., Odent, S., Loget, P., Kossorotoff, M., Snoeck, I., Plessis,
G., Parent, P., Beldjord, C., Cardoso, C., Represa, A., Flint, J., Keays, D.A., Cowan, N.J., and Chelly, J.
(2009). Mutations in the beta-tubulin gene TUBB2B result in asymmetrical polymicrogyria. Nat Genet *41*,
746-752.

Kemphues, K.J., Raff, E.C., Raff, R.A., and Kaufman, T.C. (1980). Mutation in a testis-specific beta-tubulin
in Drosophila: analysis of its effects on meiosis and map location of the gene. Cell *21*, 445-451.

Kirszenblat, L., Neumann, B., Coakley, S., and Hilliard, M.A. (2013). A dominant mutation in mec-7/beta-
tubulin affects axon development and regeneration in Caenorhabditis elegans neurons. Mol Biol Cell *24*,
285-296.

Konishi, Y., and Setou, M. (2009). Tubulin tyrosination navigates the kinesin-1 motor domain to axons.
Nature neuroscience *12*, 559-567.

Leandro-Garcia, L.J., Leskela, S., Landa, I., Montero-Conde, C., Lopez-Jimenez, E., Leton, R., Cascon, A.,
Robledo, M., and Rodriguez-Antona, C. (2010). Tumoral and tissue-specific expression of the major
human beta-tubulin isotypes. Cytoskeleton (Hoboken) *67*, 214-223.
Liao, G., and Gundersen, G.G. (1998). Kinesin is a candidate for cross-bridging microtubules and
intermediate filaments. Selective binding of kinesin to detyrosinated tubulin and vimentin. The Journal
of biological chemistry *273*, 9797-9803.
Liu, G., and Dwyer, T. (2014). Microtubule dynamics in axon guidance. Neurosci Bull *30*, 569-583.
Lockhead, D., Schwarz, E.M., O'Hagan, R., Bellotti, S., Krieg, M., Barr, M.M., Dunn, A.R., Sternberg, P.W.,
and Goodman, M.B. (2016). The tubulin repertoire of C. elegans sensory neurons and its context-
dependent role in process outgrowth. Mol Biol Cell.
McKean, P.G., Vaughan, S., and Gull, K. (2001). The extended tubulin superfamily. J Cell Sci *114*, 2723-
2733.
Mimori-Kiyosue, Y. (2011). Shaping microtubules into diverse patterns: molecular connections for
setting up both ends. Cytoskeleton (Hoboken) *68*, 603-618.
Mitani, S., Du, H., Hall, D.H., Driscoll, M., and Chalfie, M. (1993). Combinatorial control of touch receptor
neuron expression in Caenorhabditis elegans. Development *119*, 773-783.
Nogales, E., Wolf, S.G., and Downing, K.H. (1998). Structure of the alpha beta tubulin dimer by electron
crystallography. Nature *391*, 199-203.
Panda, D., Miller, H.P., Banerjee, A., Luduena, R.F., and Wilson, L. (1994). Microtubule dynamics in vitro
are regulated by the tubulin isotype composition. Proceedings of the National Academy of Sciences of
the United States of America *91*, 11358-11362.
Poirier, K., Saillour, Y., Bahi-Buisson, N., Jaglin, X.H., Fallet-Bianco, C., Nabbout, R., Castelnau-Ptakhine, L.,
Roubertie, A., Attie-Bitach, T., Desguerre, I., Genevieve, D., Barnerias, C., Keren, B., Lebrun, N., Boddaert,
N., Encha-Razavi, F., and Chelly, J. (2010). Mutations in the neuronal ss-tubulin subunit TUBB3 result in
malformation of cortical development and neuronal migration defects. Hum Mol Genet *19*, 4462-4473.
Prokop, A. (2013). The intricate relationship between microtubules and their associated motor proteins
during axon growth and maintenance. Neural Dev *8*, 17.
Pucciarelli, S., Ballarini, P., Sparvoli, D., Barchetta, S., Yu, T., Detrich, H.W., 3rd, and Miceli, C. (2012).
Distinct functional roles of beta-tubulin isotypes in microtubule arrays of Tetrahymena thermophila, a
model single-celled organism. PLoS One *7*, e39694.
Purro, S.A., Ciani, L., Hoyos-Flight, M., Stamatakou, E., Siomou, E., and Salinas, P.C. (2008). Wnt regulates
axon behavior through changes in microtubule growth directionality: a new role for adenomatous
polyposis coli. The Journal of neuroscience : the official journal of the Society for Neuroscience *28*, 8644-
8654.
Qu, C., Dwyer, T., Shao, Q., Yang, T., Huang, H., and Liu, G. (2013). Direct binding of TUBB3 with DCC
couples netrin-1 signaling to intracellular microtubule dynamics in axon outgrowth and guidance. J Cell
Sci *126*, 3070-3081.
Reijo, R.A., Cooper, E.M., Beagle, G.J., and Huffaker, T.C. (1994). Systematic mutational analysis of the
yeast beta-tubulin gene. Mol Biol Cell *5*, 29-43.
Richards, K.L., Anders, K.R., Nogales, E., Schwartz, K., Downing, K.H., and Botstein, D. (2000). Structure-
function relationships in yeast tubulins. Mol Biol Cell *11*, 1887-1903.
Sainath, R., and Gallo, G. (2015). Cytoskeletal and signaling mechanisms of neurite formation. Cell Tissue
Res *359*, 267-278.
Savage, C., Xue, Y., Mitani, S., Hall, D., Zakhary, R., and Chalfie, M. (1994). Mutations in the
Caenorhabditis elegans beta-tubulin gene mec-7: effects on microtubule assembly and stability and on
tubulin autoregulation. J Cell Sci *107 (Pt 8)*, 2165-2175.

Schaefer, A.W., Schoonderwoert, V.T., Ji, L., Mederios, N., Danuser, G., and Forscher, P. (2008).
Coordination of actin filament and microtubule dynamics during neurite outgrowth. Developmental cell
15, 146-162.
Schrodinger, LLC. (2015). The PyMOL Molecular Graphics System, Version 1.8.
Sirajuddin, M., Rice, L.M., and Vale, R.D. (2014). Regulation of microtubule motors by tubulin isotypes
and post-translational modifications. Nature cell biology *16*, 335-344.
Song, Y., and Brady, S.T. (2015). Post-translational modifications of tubulin: pathways to functional
diversity of microtubules. Trends Cell Biol *25*, 125-136.
Song, Y., Kirkpatrick, L.L., Schilling, A.B., Helseth, D.L., Chabot, N., Keillor, J.W., Johnson, G.V., and Brady,
S.T. (2013). Transglutaminase and polyamination of tubulin: posttranslational modification for stabilizing
axonal microtubules. Neuron *78*, 109-123.
Sullivan, K.F. (1988). Structure and utilization of tubulin isotypes. Annu Rev Cell Biol *4*, 687-716.
Sullivan, K.F., and Cleveland, D.W. (1986). Identification of conserved isotype-defining variable region
sequences for four vertebrate beta tubulin polypeptide classes. Proceedings of the National Academy of
Sciences of the United States of America *83*, 4327-4331.
Tanaka, E., Ho, T., and Kirschner, M.W. (1995). The role of microtubule dynamics in growth cone motility
and axonal growth. J Cell Biol *128*, 139-155.
Thompson, O., Edgley, M., Strasbourger, P., Flibotte, S., Ewing, B., Adair, R., Au, V., Chaudhry, I.,
Fernando, L., Hutter, H., Kieffer, A., Lau, J., Lee, N., Miller, A., Raymant, G., Shen, B., Shendure, J., Taylor,
J., Turner, E.H., Hillier, L.W., Moerman, D.G., and Waterston, R.H. (2013). The million mutation project: a
new approach to genetics in Caenorhabditis elegans. Genome Res *23*, 1749-1762.
Tian, G., Jaglin, X.H., Keays, D.A., Francis, F., Chelly, J., and Cowan, N.J. (2010). Disease-associated
mutations in TUBA1A result in a spectrum of defects in the tubulin folding and heterodimer assembly
pathway. Hum Mol Genet *19*, 3599-3613.
Tischfield, M.A., Baris, H.N., Wu, C., Rudolph, G., Van Maldergem, L., He, W., Chan, W.M., Andrews, C.,
Demer, J.L., Robertson, R.L., Mackey, D.A., Ruddle, J.B., Bird, T.D., Gottlob, I., Pieh, C., Traboulsi, E.I.,
Pomeroy, S.L., Hunter, D.G., Soul, J.S., Newlin, A., Sabol, L.J., Doherty, E.J., de Uzcategui, C.E., de
Uzcategui, N., Collins, M.L., Sener, E.C., Wabbels, B., Hellebrand, H., Meitinger, T., de Berardinis, T.,
Magli, A., Schiavi, C., Pastore-Trossello, M., Koc, F., Wong, A.M., Levin, A.V., Geraghty, M.T., Descartes,
M., Flaherty, M., Jamieson, R.V., Moller, H.U., Meuthen, I., Callen, D.F., Kerwin, J., Lindsay, S., Meindl, A.,
Gupta, M.L., Jr., Pellman, D., and Engle, E.C. (2010). Human TUBB3 mutations perturb microtubule
dynamics, kinesin interactions, and axon guidance. Cell *140*, 74-87.
Tischfield, M.A., Cederquist, G.Y., Gupta, M.L., Jr., and Engle, E.C. (2011). Phenotypic spectrum of the
tubulin-related disorders and functional implications of disease-causing mutations. Curr Opin Genet Dev
21, 286-294.
Topalidou, I., Keller, C., Kalebic, N., Nguyen, K.C., Somhegyi, H., Politi, K.A., Heppenstall, P., Hall, D.H.,
and Chalfie, M. (2012). Genetically separable functions of the MEC-17 tubulin acetyltransferase affect
microtubule organization. Current biology : CB *22*, 1057-1065.
Verhey, K.J., and Gaertig, J. (2007). The tubulin code. Cell Cycle *6*, 2152-2160.
Yu, I., Garnham, C.P., and Roll-Mecak, A. (2015). Writing and Reading the Tubulin Code. The Journal of
biological chemistry *290*, 17163-17172.
Zheng, C., Diaz-Cuadros, M., and Chalfie, M. (2015). Dishevelled attenuates the repelling activity of Wnt
signaling during neurite outgrowth in Caenorhabditis elegans. Proceedings of the National Academy of
Sciences of the United States of America *112*, 13243-13248.
Zheng, C., Diaz-Cuadros, M., and Chalfie, M. (2016). GEFs and Rac GTPases control directional specificity
of neurite extension along the anterior-posterior axis. Proceedings of the National Academy of Sciences
of the United States of America *113*, 6973-6978.

Zheng, C., Karimzadegan, S., Chiang, V., and Chalfie, M. (2013). Histone methylation restrains the
expression of subtype-specific genes during terminal neuronal differentiation in Caenorhabditis elegans.
PLoS genetics *9*, e1004017.

983
984 **Table legend**
985
Table 1. The *mec-7* and *mec-12* mutations analyzed in this study. Several mutations are

represented by multiple alleles, whose phenotypes were found to be similar. For touch sensitivity,

+ indicate the average response to 5 anterior stimuli is above 4; ± indicate the average is between

4 and 1; – indicate the average is below 1. Partial *lf* alleles of *mec-12* showed some but not all of

the *lf* phenotypes (see the text). Asterisks indicate that the mutation was originally found in

humans and was created in *mec-7* gene through CRISPR/Cas9-mediated genome editing.

Mapping of the amino acid residues to the structural domains was done according to Tischfield

et al. (2011).

994
995 **Figure legends**
996
Figure 1. MT structures in tubulin mutants. (A) Cross sectional images of ALM-AN in wild type

and *mec-12(tm5083 lf)*, *mec-7(ok2152 lf)*, *tba-7(u1015 lf) mec-12[u950* (S140F) *anti]* and *mec-*

12[u1021 (G144S) *anti]* mutants. Images for *mec-7(anti)* and *mec-7(neo)* can be found in

previous publications (Chalfie and Thomson, 1982; Savage *et al.*, 1994). Insets in the lower right

corner show the 15-p, 11-p, and 15-p structure of tannic acid stained MTs of *mec-12(lf)*, *mec-*

7(lf), and *tba-7(lf)* animals, respectively (4-fold enlarged). Scale bar = 100 nm. (B) Measures of

MT structure and organization. Mean ± SD are shown, and numbers of observations are in

parentheses. A Dunnett's test was performed to compare the mutants with the wild type.

Throughout the figures, one asterisk represents a statistical significance of $p < 0.05$ and two

asterisks indicate $p < 0.01$. We did not measure center-center distance and MT-occupied area in

mec-7(lf), *mec-12(lf)*, and *mec-12(anti)* mutants, because they contained very few MTs, which

did not form bundles.

Figure 2. *mec-7* and *mec-12* mutations affect TRN neurite length. (A) Schematic diagram of

ALM and PLM morphology in *mec-7* and *mec-12 lf*, *anti*, and *neo* mutants. (B) On the left panel,

compared to the wild-type animals, *mec-7(ok2152 lf)* animals have an increased gap between

PLM-AN and ALM cell body (dashed line) and significantly shortened PLM-PN (arrows). PLM-

AN still extends beyond the position of the vulva indicated by the triangle. The middle panel

shows PLM posterior neurites in the various animals. Arrows point to the shortened PLM-PN.

The right panel displays the TRN morphologies in *mec-7[u957* (P171L) *anti]*, *mec-7[u958*

(G244S) *anti]*, and *mec-12[u1019* (G354E) *anti]* animals. (C) TRN morphology of *mec-7[u278*

(C303Y) *neo]*, *mec-7[u1017* (C377F) *neo]*, and *mec-12[u917* (V260I) *neo]* animals. Arrows

point to the ectopic ALM-PN. (D) The percentage of ALM and PLM cells that had significantly

shortened anterior neurites (ALM-AN not reaching the posterior pharyngeal bulb and PLM-AN

not extending beyond the vulva). Amino acid changes in the alleles can be found in Table 1. A χ^2

test for categorical data was used to compare mutants with the wild type. (E-F) The relative

length (mean ± SEM) of PLM-PN or ALM-PN in *mec-7* and *mec-12 lf*, *anti*, and *neo* mutant

animals, as well as the double mutants of *mec-7(ok2151 lf)* with *mec-12 neo* alleles. Dunnett's

tests were used to analyze the data.

Figure 3. Position of amino acid residues changed in *mec-7* and *mec-12* mutants. (A) The

structure of the α/β tubulin dimer (1jff.pdb) visualized using PyMOL. To be consistent with the

original structure (Nogales *et al.*, 1998) and the mapping of human tubulin mutations (Tischfield

et al., 2011), α-tubulin (wheat) is shown on top of β-tubulin (cyan, magenta, and pink, which

labels α-helices, β-sheets, and loops, respectively). GTP is labeled in green. (B-C) Amino acid

changes around the GTP binding pocket in *mec-7* (red) and *mec-12* (pink) antimorphs. P173L

and S172P (blue) are disease-associated mutations found in human TUBB3 that were engineered

in *mec-7* to test their effects (see Figure 5). (D) *mec-7* (red) and *mec-12* (pink) *anti* mutations

mapped to intradimer interface. (E-F) Amino acid alterations in *mec-7* (E, yellow) and *mec-12* (F,

yellow) *neo* alleles. A302T, R380S, and E410K (orange) were clinically identified TUBB3

mutations. E415 and G416 (green), mutated in *mec-12* partial *lf* alleles, are located at the exterior

surface of α-tubulin.

Figure 4. Effects of *mec-7* and *mec-12* mutations on TRN function and activity. (A) The average

number of touch responses out of five for *mec-7* and *mec-12 lf*, *anti*, and *neo* mutants when they

are touch anteriorly. (B) Defective targeting of the synaptic vesicle marker RAB-3::GFP in *mec-*

7 and *mec-12* mutant PLM neurons. RAB-3::GFP signal is found in patches where PLM-AN

branches synapse onto target neurons in the wild-type ventral nerve cord. This targeting was

partially (reduced GFP signal at the synapse) or completely (no GFP signal at the synapse) lost in

many of the mutants; this loss in *mec-7(lf)* and *mec-12(anti)* mutants could be partly caused by

the lack of PLM synaptic branch. In *mec-7(u278*; C303Y*)* mutants, however, the marker was

mistargeted to the distal end of PLM-PN (arrow). (C) Immunofluorescent intensity of antibody

labeling of acetylated α-tubulin. (D) Fluorescent intensity of the TRN marker *uIs134 [mec-*

17p::RFP] in various *mec-7* and *mec-12* mutants and their doubles with the *dlk-1 lf (ju476)* allele.

For (C-D), representative images are shown in Figure S3. Dunnett's tests were performed to

compare the mutants with the wild type animals, and *t* tests with Bonferroni correction were used

to identify significant difference between the tubulin single mutants and *dlk-1; mec-7* or *dlk-1;*

mec-12 double mutants.

Figure 5. Modeling human tubulin mutations in *C. elegans* TRNs. (A) Sequence changes

designed to introduce TUBB3 missense mutations in *mec-7* gene. For each mutation, the line

with an arrow underscores the 20 bp DNA sequence for the guide RNA target, and the

directionality of the arrow indicates the strand of the target. The bottom DNA sequence contains

the desired nonsynonymous (red) and synonymous (blue) mutations on the homologous repair

template. Amino acids sequence with the positional information was shown on the top. (B-D)

TRN morphologies of animals carrying the engineered missense mutations. For *u1056*, the

dashed line, triangle, and arrow indicate the gap from PLM-AN to ALM cell body, the position

of the vulva, and the shortened PLM-PN, respectively. *u1057* showed normal TRN morphology.

The growth of ALM-PN (arrow) was suppressed in *mec-17(ok2109 lf); mec-7(u1056)* and *mec-*

17(ok2109 lf); mec-7(u1057) double mutants. For *u1058* and *u1059* (C), arrows point to the

ectopic ALM-PN. For *u1060* (D), the asterisk demarcates the end of the severely shortened

PLM-AN.

Figure 6. Mutations in *tba-7*, and *klp-7* result in the growth of ectopic ALM posterior neurites.

(A) A representative image of the ectopic ALM-PN in *tba-7(u1015 lf)* mutants. (B) The length of

ALM-PN in *tba-7(u1015 lf)* and the effect of 1 mM colchicine and *mec-7(ok2152 lf)*, *mec-*

7(u957 anti), *mec-12(tm5083 lf)*, and *mec-12(u1016 anti)* mutations on the *tba-7(lf)* phenotype.

(C) The differences between the TBA-7 and MEC-12 amino acid sequences in the alignment

were labeled in color. (D) The length of ALM-PN in *tba-7(u1015 lf)* carrying the transgene

expressing the wild type or mutant TBA-7 proteins. TBA-7 (NT SWAP) had

TSMEPDGNSGSLG (a.a. 35-47) replaced by QMPSDKSLGGSDDSFS, TBA-7 (CT SWAP)

had DANDNGD (a.a. 435-441) replaced by MEDNGEEG, and TBA-7 (L31Q + CT SWAP)

carried both mutations. (E) Sprouting of neurites from the ALM and PLM cell bodies in *tba-*

7(u1015 lf); *mec-7(u1017 neo)* double mutants. (F) The number of neurites emanating from the

cell bodies in *tba-7(u1015 lf)* and *mec-7(u1017 neo)* single mutants and their double mutants.

Box plot shows the minimum, first quartile, median, third quartile, and maximum; the mean is

indicated by a + sign. (G) The length of ALM-PN in *klp-7(tm2143 lf)* mutants and their double

mutants with *mec-7(ok2152 lf)*, *mec-7(u957 anti)*, *mec-12(tm5083 lf)*, and *mec-7(u1017 neo)*.

Figure 7. A model for the effect of MT dynamics on neurite development. (A) In wild-type

animals, 15-p MTs are mostly made of MEC-12/α-tubulin (brown) and MEC-7/β-tubulin (cyan).

Another α-tubulin TBA-7 (yellow) is also incorporated into the MTs, probably to decrease the

stability. For simplicity, other tubulin isotypes that may function in TRNs, including TBA-1,

TBA-2, TBB-1, and TBB-2 (Lockhead *et al.*, 2016), were shown in blue. Arrows towards the left

and right represent depolymerization and polymerization, respectively. Stars represent mutations

on the tubulin protein. (B) In the absence of MEC-7, the abundant 15-p MTs are replaced by a

few 11-p MTs, which use other β-tubulins for polymerization and can still support neurite

growth. Whether those β-tubulins are normally incorporated into the MTs is unclear. (C) *mec-*

7(anti) mutations produce dominant-negative MEC-7 proteins with defects in the GTP binding

pocket or the intradimer or interdimer interfaces. Such MEC-7 mutants can either sequester all

α-tubulin or form toxic heterodimers that can terminate MT elongation upon incorporation. Thus,

MT polymerization is blocked and neurite growth is disrupted. (D) *mec-7(neo)* mutations mostly

map to the exterior facing surface of the tubulin structure. When incorporated into the MTs,

those MEC-7 mutant proteins may affect the binding of MAPs or motor proteins (e.g. Kinesin 13

family protein KLP-7) by either binding poorer to destabilizing proteins or better to stabilizing

proteins. The net result would be hyperstable MTs. The loss of TBA-7 presumably allowed more

MEC-12 to be incorporated into the MTs, which led to increased MT stability. Abnormally

elevated MT stability caused excessive neurite growth.

Gene	Allele	Mutation	Structural function	Classification	Morphological defects	Touch sensitivity	Expression
mec-7	*u319*	S25F	Tubulin folding	weak antimorph	moderately shortened neurites	-	semidominant
	u305, u1020	G34S	Lumen-facing loop	lf	short PLM-PN	±	recessive
	gk906464	G38E	Lumen-facing loop	N/A	no defects	+	N/A
	u58, u223	P61L	Lateral interaction	weak antimorph	moderately shortened neurites	−	semidominant
	u249	P61S	Lateral interaction	weak antimorph	moderately shortened neurites	−	semidominant
	gk337318	V64I	Tubulin folding	N/A	no defects	+	N/A
	u430	A97V	GTP binding	antimorph	short TRN neurites	−	recessive
	u222	G109E	Lateral interaction	weak antimorph	moderately shortened neurites	−	semidominant
	u429, u433	G141E	Tubulin folding	lf	short PLM-PN	−	recessive
	gk595364	G141R	Tubulin folding	lf	short PLM-PN	−	recessive
	u275	G148R	Tubulin folding	lf	short PLM-PN	−	recessive
	gk895768	T149I	Tubulin folding	neomorph	ectopic ALM-PN	±	recessive
	gk286003	R162Q	MAP binding	neomorph	ectopic ALM-PN	±	recessive
	u911	P171S	GTP binding	antimorph	short TRN neurites	−	dominant
	u957, u127, e1343	P171L	GTP binding	antimorph	short TRN neurites	−	semidominant
	u1056	S172P*	GTP binding	lf	short PLM-PN	−	recessive
	u1057	P173L*	GTP binding	N/A	no defects	+	N/A
	u48	S176F	GTP binding	antimorph	short TRN neurites	−	semidominant
	u449	V179A	GTP binding	antimorph	short TRN neurites	−	semidominant
	u10	S188F	Tubulin folding	lf	short PLM-PN	±	recessive
	u225	T214P	Tubulin folding	lf	short PLM-PN	±	recessive
	ky852	P220S	Tubulin folding	neomorph	ectopic ALM-PN	−	semidominant
	u262	N226Y	GTP binding	antimorph	short TRN neurites	−	semidominant
	gk286002	P243S	Intradimer interaction	lf	short PLM-PN	±	recessive
	u283	P243L	Intradimer interaction	antimorph	short TRN neurites	−	dominant
	u129, u958	G244S	Intradimer interaction	antimorph	short TRN neurites	−	dominant
	n434	N247I	Intradimer interaction	antimorph	short TRN neurites	−	dominant
	u162	D249N	Intradimer interaction	antimorph	short TRN neurites	−	dominant
	e1505	G269D	Tubulin folding	lf	short PLM-PN	±	recessive
	e1527	V286D	Lateral interaction	antimorph	short TRN neurites	−	dominant
	u445	M300V	Tubulin folding	antimorph	short TRN neurites	−	semidominant
	u98	M300T	Tubulin folding	antimorph	short TRN neurites	−	semidominant
	u1058	A302T*	MAP binding	neomorph	ectopic ALM-PN	±	semidominant
	u278	C303Y	MAP binding	neomorph	ectopic ALM-PN	−	semidominant
	gk286001	C303S	MAP binding	N/A	no defects	+	N/A
	gk286000	A314V	Tubulin folding	antimorph	short TRN neurites	−	semidominant
	e1522	F317I	Tubulin folding	lf	short PLM-PN	±	recessive
	u234	R318Q	Tubulin folding	lf	short PLM-PN	±	recessive
	u955	A352T	Intradimer interaction	antimorph	short TRN neurites	−	dominant
	u910, gk373602	P357L	Tubulin folding	antimorph	short TRN neurites	−	dominant

	Allele	Mutation	Function	Morph	Phenotype		Inheritance
	u956	P358L	Tubulin folding	antimorph	short TRN neurites	−	dominant
	u428	G369E	Tubulin folding	lf	short PLM-PN	−	recessive
	u1017	L377F	MAP binding	neomorph	the growth of ectopic ALM-PN	±	recessive
	u1059	R380S*	MAP binding	neomorph	the growth of ectopic ALM-PN	±	recessive
	u18	A393T	Longitudinal interaction	antimorph	short TRN neurites	−	dominant
	gk285997	A393V	Longitudinal interaction	antimorph	short TRN neurites	−	dominant
	u170	E407L	MAP binding	neomorph	the growth of ectopic ALM-PN	±	recessive
	u1060	E410K*	MAP binding	antimorph	shortened TRN-ANs	−	semidominant
mec-12	*gk170196*	P32S	Lumen-facing loop	neomorph	the growth of ectopic ALM-PN	±	recessive
	gk170195	S50N	Lumen-facing loop	N/A	no defects	+	N/A
	gk636747	R60H	Lumen-facing loop	N/A	no defects	+	N/A
	u76	D69N	GTP binding	antimorph	short PLM-PN	−	recessive
	u1016	E97K	Intradimer interaction	antimorph	short PLM-PN	−	recessive
	u950, gk672907	S140F	GTP binding	antimorph	short PLM-PN	−	recessive
	gk600523	G142E	GTP binding	lf	no defects	±	recessive
	u1021, e1607	G144S	GTP binding	antimorph	short PLM-PN	−	recessive
	gk583647	L152F	Tubulin folding	N/A	no defects	+	N/A
	u50, e1605	H192Y	MAP binding	partial lf	no defects	−	recessive
	gk915672	E196K	MAP binding	neomorph	the growth of ectopic ALM-PN	±	recessive
	u1041	G246E	Tubulin folding	lf	no defects	±	recessive
	u917	V260I	MAP binding	neomorph	the growth of ectopic ALM-PN	±	recessive
	gk854211	P307L	MAP binding	N/A	no defects	+	N/A
	gk515972	V323I	Longitudinal interaction	neomorph	the growth of ectopic ALM-PN	±	recessive
	u241, u1019	G354E	Longitudinal interaction	antimorph	short PLM-PN	−	recessive
	gk341552	G365E	Lumen-facing loop	N/A	no defects	+	N/A
	u63	E415K	MAP binding	partial lf	no defects	±	recessive
	gm379	G416E	MAP binding	partial lf	no defects	±	recessive

A

WT | *mec-12(lf)* | *mec-7(lf)*

tba-7(lf) | *mec-12(u950 anti)* | *mec-12(u1021 anti)*

B

Genotype	Number of MTs	MT diameters (nm)	Center-center distance (nm)	MT-occupied area (%)
wild type	31.1 ± 7.3 (79)	21.5 ± 1.6 (126)	34.4 ± 1.0 (320)	46.7 ± 5.5 (38)
mec-7(ok2152 lf)	6.0 ± 2.2 (52)**	16.6 ± 1.4 (123)**	–	–
mec-12(tm5083 lf)	7.4 ± 2.4 (36)**	21.3 ± 2.0 (103)	–	–
tba-7(u1015 lf)	27.0 ± 10.1 (109)	22.4 ± 1.8 (248)**	32.3 ± 2.1 (278)**	57.2 ± 10.1 (74)**
mec-7(u170 neo)	18.0 ± 5.8 (25)**	22.7 ± 2.2 (192)**	31.7 ± 2.3 (148)**	58.6 ± 5.2 (20)**
mec-7(u278 neo)	20.6 ± 6.8 (23)**	22.5 ± 1.9 (176)**	32.9 ± 1.5 (164)**	59.0 ± 8.1 (18)**
mec-12(u76 anti)	1.6 ± 1.5 (52)**	17.6 ± 2.6 (53)**	–	–
mec-12(u950 anti)	2.5 ± 2.1 (40)**	18.0 ± 2.0 (75)**	–	–
mec-12(u1021 anti)	2.7 ± 2.4 (72)**	17.9 ± 1.8 (114)**	–	–

A

S172P
```
169  V  V  P  S  P  K  V  S  D  T  V  V  E  P  Y  183
     GTAGTTCCAAGTCCAAAAGTATCCGACACTGTAGTCGAACCCTAC
     GTAGTTCCACCTCCAAAGGTCTCCGACACTGTAGTGCTTGGGATC
```

P173L
```
169  V  V  P  S  P  K  V  S  D  T  V  V  E  P  Y  183
     GTAGTTCCAAGTCCAAAAGTATCCGACACTGTAGTCGAACCCTAC
     GTAGTTCCAAGTCTGAAGGTCTCCGACACTGTAGTGCTTGGGATC
```

A302T
```
296  A  K  N  M  M  A  A  C  D  P  R  H  G  R  Y  310
     GCAAAGAACATGATGGCTGCATGCGATCCAAGACATGGACGTTAT
     GCAAAGAACATGATGGCTACTTGTGACCCTAGACATGGACGTTAT
```

R380S
```
369  G  N  S  T  A  I  Q  E  L  F  K  R  I  S  E  383
     GGAAACTCCACCGCCATTCAAGAGCTCTTTAAGAGAATCTCAGAG
     GGAAACTCCACCGCCATACAGGAACTGTTCAAGAGTATCTCAGAG
```

E410K
```
402  G  M  D  E  M  E  F  T  E  A  E  S  N  M  N  416
     GGAATGGACGAGATGGAATTCACTGAGGCGGAGAGCAACATGAAC
     GGAATGGACGAGATGGAGTTTACCGAAAGCGGAGAGCAACATGAAC
```

B *u1056* (S172P)   ALMs   PLMs
100μm

u1057 (P173L)   PLMs   ALMs
100μm

mec-17(ok2109 lf)   ALM   20μm
mec-17(lf); mec-7(u1056)
mec-17(lf); mec-7(u1057)

C *u1059* (R380S)   ALM   20μm

u1058 (A302T)   ALMs   PLMs   100μm

D *u1060* (E410K)   ALMs   PLMs   100μm

A wild type; *mec-17p::RFP*   *tba-7(u1015 lf)*
ALM   ALM
20μm   20μm

B ALM-PN

Neurite length/ Cell body length

Colchicine − + − + − − − − − −
mec-7 + + + + *lf* *anti* + + − −
mec-12 + + + + − − *lf* *anti* + +
tba-7 + + + *lf* *neo* *neo*

C

TBA-7   26 LEHGI**L**PDG**TSMEP---DGNSGSLG**TFF 50
MEC-12 26 LEHGI**Q**PDG**QMPSDKSLGGSDDSFS**TFF 53

TBA-7 418 EDLAALEKDYEEVGADS**DANDN-GD**DEY 444
MEC-12 423 EDLAALEKDYEEVGVDS**MEDNGEEG**DEY 450

D ALM-PN

Neurite length/ Cell body length

tba-7 (+) +
tba-7 (L31Q) +
tba-7 (NT SWAP) +
tba-7 (CT SWAP) +
tba-7 lf

E *tba-7(u1015 lf); mec-7(u1017)* ALM

tba-7(u1015 lf); mec-7(u1017) PLM

20μm

F

Number of neurites

ALM   PLM

mec-7 + + *u1017* + + *u1017*
tba-7 + *lf* + *lf* + *lf*

G ALM-PN

Neurite length/ Cell body length

mec-7 + *lf* *lf* *anti* *neo*
mec-12 + + *lf* +
klp-7 *lf*