TITLE PAGE

Title: Population temporal structure supplements the rate code during sensorimotor transformations

Abbreviated title: Population temporal dynamics in gaze control

Authors: Uday K. Jagadisan[1,3,4] and Neeraj J. Gandhi[1,2,3]

Author affiliations:
[1]Department of Bioengineering
[2]Department of Neuroscience
[3]Center for the Neural Basis of Cognition
University of Pittsburgh, Pittsburgh, PA 15213
[4]Present Address: Department of Molecular and Cell Biology
University of California, Berkeley, Berkeley, CA 94720

Corresponding author: Correspondence and requests for more information should be addressed to
U.K.J (kj.udayakiran@gmail.com) or N.J.G (neg8@pitt.edu).
14

Author Contributions: U.K.J and N.J.G designed the study. U.K.J. performed the experiments, analyzed
the data, and performed model simulations. U.K.J and N.J.G wrote the manuscript.

Number of pages: 50

Number of figures: 6 + 9 supplementary figures, 1 supplementary table

Number of words: Abstract - 174, Main text - 5819, Supplementary sections - 2, Methods - 5512

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: We thank Drs. A. Batista, C. Olson, M. Smith, and B. Yu for scientific discussions and
critical feedback on previous versions of the manuscript and J. McFerron for programming assistance.
The study was funded by the following NIH R01 grants: EY022854 and EY024831 awarded to N.J.G.

Summary

Sensorimotor transformations are mediated by premotor brain networks where individual neurons represent sensory, cognitive, and movement-related information. Such multiplexing poses a conundrum – how does a decoder know precisely when to initiate a movement if its inputs are active at times when a movement is not desired (e.g., in response to sensory stimulation)? Here, we propose a novel hypothesis: movement is triggered not only by an increase in firing rate, but critically by a reliable temporal pattern in the population response. Laminar recordings in the macaque superior colliculus (SC), a midbrain hub of orienting control, and pseudo-population analyses in SC and cortical frontal eye fields (FEF) corroborated this hypothesis. Importantly, we used spatiotemporally patterned microstimulation to causally verify the importance of temporal structure and demonstrate its role in gating movement initiation. We also offer a spiking neuron model with dendritic integration as a putative mechanism to decode this temporal information. These findings offer new insights into the long-standing debate on movement generation and highlight the importance of short-term population history in neuronal communication and behavior.

Abstract

Sensorimotor transformations are mediated by premotor brain networks where individual neurons represent sensory, cognitive, and movement-related information. Such multiplexing poses a conundrum – how does a decoder know precisely when to initiate a movement if its inputs are active at times when a movement is not desired (e.g., in response to sensory stimulation)? Here, we propose a novel hypothesis: movement is triggered not only by an increase in firing rate, but critically by a reliable temporal pattern in the population response. Laminar recordings in the superior colliculus (SC), a midbrain region that plays an essential role in orienting eye movements, indicate that the temporal structure across neurons is a factor governing movement initiation. Specifically, using a measure that captures the fidelity of the population code - here called temporal stability - we show that the temporal structure fluctuates during the visual response but becomes increasingly stable during the movement command, even when the mean population activity is similar between the two epochs. Analyses of pseudo-populations in SC and cortical frontal eye fields (FEF) corroborated this model. We also used spatiotemporally patterned microstimulation to causally test the contribution of population temporal stability to movement initiation and found that stable stimulation patterns were more likely to evoke a movement, even when other features of the patterns such as mean pulse rates and population state subspaces were matched. Finally, a spiking neuron model was able to discriminate between stable and unstable input patterns, providing a putative biophysical mechanism for decoding temporal structure. These findings offer an alternative perspective on the relationship between movement preparation and generation by situating the correlates of movement initiation in the temporal features of activity in shared neural substrates. They also suggest a need to look beyond the instantaneous rate code at the single neuron or population level and consider the effects of short-term population history on neuronal communication and behaviour.

INTRODUCTION

In order to successfully interact with the environment, the brain must funnel down the sensory inputs it receives to specific movements at specific times. Such sensory-to-motor transformations are critically mediated by premotor brain networks where evolving activities in individual neurons represent sensory, cognitive, and movement-related information (*1-3*). For example, in brain regions involved in the control of gaze, including the superior colliculus (SC) and the frontal eye fields (FEF), a population of visual neurons processes the stimulus, while another group of motor or movement neurons discharges a volley of spikes to produce the saccade to the location of the stimulus. So-called visuomovement neurons, which constitute the majority, burst for both stimulus presentation and saccade generation (*1*). In reality, visuomovement and motor neurons span a continuum that exhibits different amounts of visual and movement-related activities. Such neurons from both SC and FEF project to the saccade generation circuitry in the brainstem (*4, 5*). How the activity patterns of visuomovement and motor neuron populations collectively drive saccade initiation remains an unresolved question in motor systems neuroscience.

Canonically, accumulation to threshold mechanisms have been thought to underpin saccade initiation (*6, 7*). The accumulator model posits that population activity in SC and FEF increases stochastically toward a constant threshold activation level, leading to the generation of a high-frequency burst and execution of a saccade. While support for the constant threshold criterion has been reported for FEF motor neurons (*6*), visuomovement neurons in FEF as well as visuomovement and motor cells in SC do not seem to obey the criterion (*8*). Moreover, in many visuomovement neurons, the threshold activation level estimated from the premotor burst is also crossed by the sensory response, yet a movement is not triggered early. It has also been suggested that the threshold criterion applies to the population response, not necessarily to every neuron in the ensemble (*9*). Even then, it remains possible that the population activity could exceed the threshold level during the sensory response. Thus, the accumulator model has limitations.

Variability of neural population activity has also been proposed as a potential mechanism of movement initiation, notably in the skeletomotor system. For simultaneously recorded neural populations, e.g., in premotor or motor cortex, the ensemble response for each trial can be represented as a neural trajectory in multi-dimensional population activity space. These trajectories have been shown to traverse through "optimal subspaces", defined as regions of reduced variability computed across trials (*10*). In the delayed reach task, for example, such optimal subspaces can be defined for stimulus onset, motor preparation and movement execution stages (*11*). A closely related model posits that muscles are recruited and a movement is initiated when neural activity traverses an optimal or "motor-potent" region of the population state space (*12, 13*) and is inhibited otherwise, e.g., during movement preparation (*13, 14*). A major strength of these approaches is that the firing rate need not reach a threshold to trigger the movement (but see (*15*)), yet this framework also has limitations. Since variability measures are computed by combining information across trials, they are not available to be used by downstream structures on individual trials. More importantly, neural variability models do not provide direct mechanistic insights into how neurons decoding their inputs map the presence of a trajectory in a particular subspace into a specific function, nor have they been validated using causal manipulations.

In this study, we propose a novel mechanism for movement generation based on the temporal structure
of neural population activity. Our central hypothesis is that saccade initiation occurs when high firing
rate (burst) is coupled with stable temporal structure in population activity. We specifically propose that
the visual burst is unstable and therefore cannot trigger a saccade, while the premotor burst is stable
and does initiate the movement when activity crosses threshold. We demonstrate the presence of this
code using a combination of laminar population recordings in the SC and pseudo-population analyses in
SC and FEF, suggesting that temporal stability features permeate through the entire visuomotor
neuraxis, encompassing both cortex and subcortex (*1*). We also show that the code is robust across tasks
and sub-regions of SC. Crucially, we provide causal validation of this mechanism by delivering temporally
structured or unstructured microstimulation patterns into the SC with a linear electrode array and
comparing their movement-inducing efficacy. Finally, we show how population temporal structure could
be decoded by a spiking neuron model with dendritic integration. We foresee this computation as a
pervasive feature of neuronal communication, and we apply it specifically to understand the
mechanisms of saccade initiation.

RESULTS

Joint representation of visual stimulus and premotor processing in visuomovement neurons

The aforementioned, dual nature of visuomovement neurons is best illustrated by examining their
activity in the delayed response paradigm (left panel in Figure 1A), which requires the subject to
withhold a saccade to a stimulus in the visual periphery until after the disappearance of a central
fixation cue. We recorded population activity from the SC using laminar microelectrode arrays (right
panel in Figure 1A) in two monkeys (*Macaca mulatta*). Figures 1B shows responses on individual
channels (top row) and population activity averaged across channels (bottom row) aligned on target
(left) and saccade (right) onsets for three example trials in one session. The population exhibits a high-
frequency visual burst following target onset and a subsequent premotor burst prior to a saccade. The
peak magnitude of the visual burst is lower than that of the premotor burst on some trials (light traces
in Figure 1B), but it is not uncommon for the peak visual response to match or exceed the premotor
activity (medium and dark traces in Figure 1B), especially when accounting for the 10-20 ms efferent
delay from neural initiation to movement onset (*6, 16, 17*) (vertical dashed line in bottom panel of
Figure 1B, also see Figure 1G), consistent with previous reports that tectoreticular neurons involved in
saccade generation show strong visual and premotor transients (*4*). Yet, on these trials, the visual burst
does not trigger a movement, an observation that casts doubt on thresholding (*6, 8*) as a singular
mechanism governing movement initiation.

We also tested whether saccade initiation could be controlled solely by "motor" neurons, thought to be
active only during saccades and not during the visual epoch. This hypothesis requires motor neurons to
exclusively have a high motor potential, i.e., be most correlated with saccade kinematics (*18*). While
motor neurons indeed have a high motor potential (Supplementary Fig. 1A and B), we found that
neurons with high visual activity also were among the strongest drivers of the saccade (Supplementary
Fig. 1C and D), disputing the categorical motor neuron hypothesis. Given that visuomovement and
motor neurons project directly to the brainstem saccade burst generator that initiates and guides
saccadic gaze shifts (*4, 5, 19*), we asked how downstream structures are able to differentiate between
the two bursts.

Population temporal structure is scrambled during the visual response and stable during the premotor response

We reasoned that if static features of population activity such as the mean firing rate are insufficient to discriminate between visual and premotor bursts, the answer likely resides in the spatiotemporal structure of activity during the two bursts. Indeed, a growing body of work has revealed that precise coordination in the timing of input spikes is more efficient at driving downstream cortical and subcortical neurons (20-22). Specifically, we hypothesized that a critical criterion for a decoder to generate a movement should be a high level of certainty in the instructed movement and its metrics. Since the exact decoding scheme is unknown, we considered whether certainty is provided by consistency in the population pattern over the course of a burst, under any generalized weighted pooling scheme. To quantify this consistency, we first constructed a population firing rate trajectory as a function of time. Next, we normalized the population vector at each time point, constraining it to a unit hypersphere in state space (Figure 1C; see Methods for details). This step factors out global changes in firing across the population and focuses on the relative activity pattern. We then computed the dot product between two of these unit vectors separated in time (parametrized by τ) - we call this measure the temporal stability of the population code.

Figure 1D shows the evolution of temporal stability averaged across all sessions (n = 280 channels/16 sessions, mean +/- s.e.m., colored traces; also see Supplementary Fig. 2A, top row). The stability of the population pattern decreased relative to baseline during the visual burst and increased during the premotor burst. Moreover, this property was preserved when considering only the subset of trials where the peak visual response matched or exceeded the premotor activity 15 ms before saccade onset (gray trace in Figure 1D) and was significantly different from the trend in the null condition when the saccade was directed to a stimulus in the opposite hemifield (black trace in Figure 1D). Scrambling the population code by either shuffling the activations of individual neurons at each time point or shuffling across time points lowered the stability profile for the entire trial significantly (Supplementary Fig. 2A, bottom row). The anti-phase relationship of stability between the visual and premotor bursts was also present across a range of separation times between the two population vectors used for the dot product (Supplementary Fig. 2B). We also computed temporal stability by realigning activity on the peak population visual response, to discount any effect of visual response onset latency differences across the population and found similar (if not stronger) effects (Figure 1E, also see Supplementary Fig. 2C). Finally, temporal stability afforded more robust discrimination between the visual and premotor bursts even on individual trials and sessions (Figure 1F), compared to the population firing rate (Figure 1G). These results were unaffected by spike sorting (Supplementary Fig. 3, also see Methods), in line with recent evidence that spike sorting has a negligible effect on population analyses (23). Thus, population temporal stability seems to impose a constraint that prevents movement initiation at an undesirable time (visual epoch), allowing the animal to successfully perform the task at hand. Once cued to execute a gaze shift, the activity in the same population rises in a stable manner allowing saccade initiation.

The distinction between visual and premotor temporal structure is robust across brain regions and tasks

Next, we explored the robustness of the temporal stability hypothesis. We tested whether the properties of stability observed in other populations of neurons or conditions were consistent with its

predicted role in movement initiation. For this part of the study we used data obtained from single-
electrode recordings in SC and the frontal eye fields (FEF), the latter of which plays a major role in the
cortical control of saccade initiation (*6, 24*). First, we constructed pseudo-trials by random sampling of
trials from individual neurons/single-unit recording sessions (see Methods for details) and used this
pseudo-population to construct surrogate neural trajectories (Figure 2A, top row, also see Methods and
Supplementary Fig. 4). Since SC and FEF both project to saccade-generating brainstem structures (*25*),
we also analyzed a combined SC+FEF pseudo-population. The middle row of Figure 2A shows the
average normalized pseudo-population activity of 60 SC neurons, 30 FEF neurons, and the combined 90-
neuron population, reaffirming the visuomovement pattern of activity discussed earlier. Importantly,
the temporal structure profiles (Figure 2A, bottom row) were remarkably similar to those observed with
simultaneous population recordings – the dramatic reduction in the stability during the visual burst and
an increase in stability during the premotor burst – in both SC and FEF individually (transparent and
opaque colored traces in Figure 2A, bottom row), as well as in the combined population (black traces in
Figure 2A, bottom row).

We then used pseudo-population analyses to test whether the temporal stability framework was
obeyed by SC neurons beyond those that are involved in the generation of large or non-fixational
saccades. Neurons in the rostral SC are tonically active during fixation and reduce their activity during
larger movements (*26*) (Figure 2B, middle row). Importantly, they also project to the saccade burst
generator (*27*). Assuming population stability modulates the input drive to downstream structures, we
hypothesized that the temporal structure in rostral SC neurons must decrease during large saccades but
remain elevated during fixation, even when a visual burst occurs in other parts of SC and FEF. In other
words, we expected the evolution of temporal stability across rostral SC neurons to be the inverse of
what occurs in caudal SC - Figure 2B (bottom row) confirms this prediction, both when rostral SC is
considered separately, as well as in the combined SC population. In addition to the antiphase
relationship with caudal SC during large saccades, rostral SC is also known to play a causal role in the
generation of microsaccades (*28*) during fixation. Therefore, we asked whether the rostral SC population
exhibits stable temporal structure during the burst that produces microsaccades. This was indeed the
case, where the same rostral SC population that showed a reduction in temporal stability during large
saccades showed an increase during microsaccades (Figure 2C). Thus, the population activity of neurons
in rostral SC, including its temporal structure, complements the pattern in other parts of SC in both
suppressing and initiating movements.

We also examined whether population temporal structure was related to the propensity to trigger
saccades off the visual burst. In a modified task paradigm called the gap task (top row, Figure 2D),
"express saccades" occur under certain conditions, and are reflected in the initial peak of a bimodal
reaction time distribution (inset in middle row, Figure 2D). These low-latency saccades are thought to be
triggered directly by the initial visual burst in SC (middle row, Figure 2D). If temporal structure of SC
activity is related to the saccade generation process, then temporal stability should not decrease at all or
as much during the visual burst when express saccades are triggered. The bottom row of Figure 2D
shows that this was indeed the case – the temporal stability in the visual epoch is higher for express
saccades (it drops slightly but quickly recovers) compared to the notable trough for regular saccades. In
sum, temporal stability seems robustly correlated with saccade generation across brain regions (rostral
SC, caudal SC, and FEF), tasks (delayed, regular, and express saccades), experimental techniques

(laminar and single electrode recordings), and analytical approaches (spike sorted, unsorted, and
pseudo-population analyses).

Microstimulation patterns with stable temporal structure are more likely to evoke movements

Thus far, we have shown that population temporal structure is a candidate measure that can be used to
discriminate between visual and premotor bursts, using purely correlational analyses. To test whether
this measure is actively used by the brain in a causal manner, we performed multi-site patterned
microstimulation in SC. We first identified individual contacts of the laminar probe that evoked low-
latency saccades with suprathreshold stimulation, to verify their placement in the intermediate layers of
SC. We then designed temporally stable and unstable stimulation patterns restricted to these contacts,
limiting stimulation parameters for individual sites to the sub-threshold regime and verifying that the
overall stimulation was near-threshold (for details, see Methods). Stable patterns were created with a
linearly decreasing inter-pulse interval (IPI) sequence for one contact (29) (to simulate a burst) and
scaling the IPIs for other contacts by a uniformly spaced factor. For each stable pattern, we also created
a *paired* unstable pattern by jittering the pulse times within a window and shuffling spikes between
contacts, preserving both the pulse count for each site and the average pulse rate across the
"population".

An example pair of stable-unstable pulse trains is shown in Figure 3A (top row). The pulse rates
determined from these trains (second row) illustrate the scaled rates for the stable pattern and the
fluctuating rates for the unstable pattern for individual sites, despite the comparable population rates
on individual trials (thick traces; see Supplementary Fig. 5B,C for more examples). Note that the
assignment of high and low rates to different contacts was randomized across trials, resulting in similar
trial-averaged pulse rates for any given channel, for either type of pattern (third row). The temporal
stability for stable and unstable patterns for all trials in the example session (bottom row) confirm the
impression provided by the pulse trains and rates, i.e., the scaled patterns are highly stable compared to
the relative instability of the jittered patterns.

We delivered these stimulation patterns during the "gap period" in a gap saccade task (Supplementary
Fig. 5A). Figure 3B shows a scatter plot of the stimulation-aligned saccade latencies observed using
stable versus unstable stimulation patterns for all pairs from one session. For the majority of stable-
unstable trial pairs, a saccade was evoked with the stable but not the unstable pattern, henceforth
referred to as S+US- trial pairs. These are the points in the blue shaded window spanning the stimulation
duration in Figure 3B. Points in the unshaded region refer to trials in which neither stable or unstable
pair evoked a saccade, and these trials were assigned the latency relative to stimulation onset of the
saccade directed to a target presented after the gap period. The relative likelihood of evoking a
movement with the stable stimulation pattern only, as computed in the graphic on the right, is shown as
a function of session number in Figures 3C (blue points). For most sessions, the observed relative
likelihood values were significantly higher (i.e., biased towards the stable pattern; p < 0.01, permutation
tests) than the null distribution (gray points) obtained by shuffling trials with randomly assigned stable-
unstable identities, reinforcing the observation that the stable pattern was more likely to evoke a
movement compared to a *state-matched* unstable pattern. We also analyzed the movement vectors
evoked by stable and unstable patterns – both sets of saccades, when they occurred, were similar to
each other in both amplitude and direction and to the site-specific vectors evoked by constant

frequency multi-channel stimulation (Supplementary Fig. 5D and 5E; p > 0.01 for most sessions,
Wilcoxon rank-sum tests; for details, see Methods).

Additionally, we explored whether differences in pulse rates on subsets of trials could explain the
across-sessions range of movement likelihood effects seen in Figure 3C (Supplementary Fig. 6). Sessions
in which the stable pattern was more likely to evoke movements were also the ones in which the S+US-
trials (Supplementary Fig. 6A) were closest to the intended uniform sampling of pulse rates across
channels, as reflected in the low relative dispersion index for those sessions (Supplementary Fig. 6B and
6C). Sessions with no significant (or negative) effect on movement likelihood were the ones which had a
high across-channel pulse rate sampling asymmetry, potentially allowing extremes in pulse rates to
mask any underlying effect of temporal structure (Supplementary Fig. 6B and 6C; also see Methods).

Linear decoders using temporal information outperform static models of movement generation

Having demonstrated the role of population temporal structure in saccade initiation using both
correlative and causal approaches, we sought to disambiguate the stability framework from extant
models of movement initiation. We earlier argued for the implausibility of threshold-based gating based
on the fact that the population activity during the visual burst can sometimes exceed premotor activity
(Figure 1B & 1G). Could other population activity-based mechanisms, such as weighted linear readout
models that seem to govern movement generation in the skeletomotor system (e.g., the optimal or
movement-potent subspace frameworks) (*12-14*), play a role here? To test this, we first used factor
analysis (FA) to visualize the low-dimensional neural states (*30*) in the visual and premotor bursts (see
Methods). An example 3-dimensional FA projection of the two bursts is shown in Figure 4A – the two
sets of states are clearly separable, likely an effect of the subtle yet distinct trends in visual versus
premotor activity levels along the dorso-ventral extent of the SC (e.g., Figure 1B) (*31*). Indeed, a linear
discriminant analysis (LDA) classifier was able to easily discriminate between the visual and premotor
states for most sessions (histogram inset in Figure 4A). The properties of neural activity in SC therefore
seem to be consistent with a static state space code as well.

Next, we used data from the patterned microstimulation experiments to disambiguate between the
optimal subspace and temporal stability frameworks. Using a representative low-dimensional FA
projection to differentially visualize stable and unstable stimulation patterns was infeasible because all
the electrode sites contributed equally to the patterns, due to the random assignment of pulse rates
across sites on different trials. From the example 3-dimensional projection in Figure 4B, it is evident that
the stable and unstable patterns are indiscriminable in state space. Indeed, an LDA classifier failed to
discriminate between stable and unstable patterns based on population pulse states alone, never
yielding above-chance classification (Figure 4B, orange bar in histogram inset). However, an LDA model
that also included stability information was able to classify perfectly, reaffirming the difference in
temporal structure (by design) between the two sets of patterns (Figure 4B, green bar in inset; also see
right matrix in Figure 4C). These results indirectly provide a key piece of evidence in support of the
notion that the brain in fact uses temporal information, since the evoked behavior reflected the
difference between stable and unstable patterns while a static linear readout of population states did
not.

The next set of analyses examined the effect of stimulation pattern on behavior. We focused on the trial
pairs in which only one of stable or unstable pattern evoked a movement (left matrix in Figure 4C), since
these are likely to provide insight into the discriminatory role played by temporal structure. We trained
another LDA classifier to discriminate between trials in which stimulation evoked a movement and those
where no movement was evoked. When using the pulse states alone, the linear decoder was unable to
determine whether a movement was evoked on a given trial with above-chance accuracy (orange points
in Figure 4D). Since the stable-unstable pairs were matched in terms of whether a movement was
evoked or not for this subset, this readout of the population pattern was independent of its temporal
structure. Crucially, and in contrast, when the population pulse states were supplemented with stability
information (added as a native dimension in the input to the decoder, see green matrix in Figure 4C),
classifier performance increased to reflect the broad trend in relative movement likelihood (compare
blue points in Figure 3C with green points in Figure 4D). The correlation between performance of the full
LDA classifier and the behavioral effects of stimulation is better highlighted in Figure 4E, i.e., the brain's
ability to extract temporal structure to generate movements is reflected by a model that utilizes this
temporal information. We also independently evaluated whether temporal stability was more
informative than each and all pulse count dimensions by performing model selection analysis with a
logistic regression fit to the movement occurrence data used for the previous analysis. For every
stimulation session, the best model (one with the lowest Bayesian Information Criterion or BIC) was one
that only had the temporal stability dimension (blue asterisks indicating minima in BIC curves in Figure
4F). Together with the effect of dispersion seen earlier (Supplementary Fig. 6), these analyses reveal two
independent contributors to movement initiation in SC – a static "laminar" code, presumably related to
the optimal or potent subspace, that may be a function of the distribution of preferred stimulation sites,
and a dynamic "temporal" code, where stability of the population pattern controls movement initiation
even under matched state space conditions.

We also tested how the temporal stability hypothesis was related to modified threshold-based
mechanisms that rely on pooling the activity of a correlated ensemble of neurons (*9, 32*). In this
framework, correlations in the accumulation rates of neurons influence movement initiation (reaction
times) under a given pooling scheme, with higher correlations leading to shorter reaction times. Spike
count correlations between SC neurons in the visual and premotor epochs (for details, see Methods)
were inconsistent with this notion – the correlations during the visual epoch were slightly, but
significantly, higher compared to the premotor epoch (Supplementary Fig. 7A, mean difference between
visual and premotor epoch correlations = 0.0167 > 0, p = 2.5E-6, one-tailed t-test). We also computed
pulse count correlations for the stable and unstable stimulation patterns and found no difference
(Supplementary Fig. 7B, median difference in correlations between conditions = 4.11E-4, not
significantly different from 0, p = 0.279, Wilcoxon signed-rank test), despite the significant difference in
their effects on behavior. Thus, the temporal stability framework is, to a large extent, independent of
extant models of movement initiation.

Biophysical models with synaptic facilitation can decode population temporal structure

Finally, we sought to identify a mechanism by which downstream neurons could discriminate between
stable and unstable population codes. We modelled the decoder as a spiking neuron that receives
population inputs through its network of dendrites (Figure 5A). The decoder can be thought to represent

neurons in the pons that receive and integrate SC inputs and burst for saccades (*27*). To mimic the
potent inhibitory gating (and disinhibition during saccades) provided by the omnipause neurons (OPNs)
on the burst neurons, we also included a spiking disinhibitor unit with reciprocal inhibitory connections
with the decoder. The disinhibitor also received both excitatory (*33*) and inhibitory (*34*) inputs from the
same population as the decoder, creating a balance between excitation and inhibition. This balance
must be overcome by some property of the input activity in order to produce a saccade. How might the
population pattern be converted into a signal that initiates a movement only when the inputs are
stable? Conceptually, the decoder should have a mechanism to keep track of the short-term history of
population activity, use this history to evaluate temporal structure, and respond selectively when the
activity pattern is deemed stable over the time scale of integration. We incorporated these heuristic
requirements by using state-dependent modulation of input-evoked excitatory post-synaptic potentials
in the decoder, i.e., inputs arriving when the local post-synaptic potential was depolarized caused a
higher subsequent change in the potential (*35, 36*) (Figure 5B).

We simulated the model with the stable and unstable patterns from the microstimulation experiments
as inputs, since they readily offered mean-matched input sequences differing only in temporal structure.
We characterized the efficacy of integration by the decoder as the time of first spike in the burst, a
proxy for movement initiation latency. Although there were several pairs of trials for which both or
neither the stable and unstable inputs caused the decoder to spike (Figure 5C), there were a number of
pairs for which only the stable input pattern was decoded (arrow in Figure 5C). To facilitate visualization
and comparison with experimental data (Figure 3B), latency values greater than the stimulation duration
were randomly assigned when the decoder wasn't recruited; this provides a visualization of the number
of trials where no movement was evoked by either pulse train of the pair. Figure 5D shows the
membrane potential output of the disinhibitor (cyan and magenta traces) and decoder (blue and red
traces) units for a pair of stable (top row) and unstable (bottom row) input patterns from this subset of
trial pairs. The disinhibitor maintained a tonic firing rate throughout the trial when the inputs were
unstable, and sharply reduced its activity at some point during stable stimulation (Figure 5E - magenta
and cyan traces, respectively). This latter disinhibition on stable trials was in anti-phase relationship with
the bursting exhibited by the decoder, which was absent on unstable trials (Figure 5E - blue and red
traces, respectively). See Supplementary Fig. 8 for membrane potential waveforms when both or neither
the stable and unstable patterns recruit the decoder unit. The firing rate profiles of the decoder and
disinhibitor units share a striking resemblance to medium-lead burst neurons and OPNs, respectively, in
the pons (*37*).

Thus, a relatively simple, yet biophysically realistic, module seems capable of discriminating between
population inputs based solely on their temporal characteristics, offering a putative mechanism by
which downstream networks could use temporal information to make decisions about movement
generation. The model simulations predict that temporal structure will impact integration near
activation threshold, but a surplus of spikes in an unstable input may still be integrated well enough to
produce a movement, consistent with the observations in the patterned microstimulation experiment.
Another testable prediction is that disinhibition of the gaze control circuitry by silencing the OPNs
should diminish the effect of temporal structure in determining movement initiation, producing
movements for both stable and unstable inputs. We also found that a firing rate-based accumulator
model with short-term synaptic plasticity produces comparable results (Supplementary Fig. 9),

demonstrating the flexibility of various biophysical mechanisms in their ability to decode temporal
structure and hence, the model-independence of this result.

DISCUSSION

Neurons in premotor structures are constantly bombarded with information from thousands of
presynaptic neurons that are active during sensorimotor processing. It is unclear how activity relevant
for movement initiation is discriminated from activity related to other processing. Extant models of
movement generation that rely on firing rate, including rise-to-threshold (6), inhibitory gating (37), and
dynamical switches at the population level (12, 13), leave certain explanatory gaps unfilled. A canonical
model of movement initiation, especially in the oculomotor system, is threshold-based gating(6) (Figure
6A-B). Current knowledge points to a role by the OPNs in defining the threshold and controlling saccade
initiation (6, 38). However, thresholds vary across behavioral paradigms (8, 39), raising the question of
how the threshold is set in a particular condition. Furthermore, evidence that the threshold changes
during the course of a trial purely based on OPN activity is limited; OPN activity increases transiently
during the visual burst but only by an additional spike or two and only in a small subset of neurons (38).
Critically, the existence of trials in which the population activity during the visual response exceeds
premotor activity strongly reduces the likelihood of thresholding operating as a singular mechanism.

An extension of simple thresholding is a mechanism based on the pooled activity of neurons with
varying degrees of correlation in the population (9). However, we show that population correlation
statistics are not necessarily related to temporal structure or saccade initiation (Supplementary Fig. 7). A
related alternative explanation is that a movement is initiated specifically when movement-only neurons
are active, but most neurons in sensorimotor structures like SC and FEF span a continuum between
visuomovement activity and movement activity (1, 31, 40). In fact, antidromically identified
tectoreticular neurons that presumably drive saccade generation exhibit both visual and premotor
bursts(4). Moreover, as seen earlier, SC neurons with the highest correlation to movement velocity tend
to also have strong visual responses (Supplementary Fig. 1). Finally, the FEF also projects to the
premotor circuitry in the brainstem (25). Our pseudo-population analysis revealed similar temporal
stability waveforms for both regions considered individually (Figure 2A, colored traces) as well as
collectively (Figure 2A, black trace). Incorporating the result that impairment of either region alters
saccade kinematics produced by the other (41, 42) together with these results suggests that both
regions operate cooperatively to initiate saccades.

Other mechanisms, primarily based on static weighted linear readouts of population activity, have been
proposed for how movement generation unfolds in the skeletomotor system. These models posit that
muscles are recruited and a movement is initiated when neural activity traverses an optimal or motor-
potent region of the population state space (12, 13) and is inhibited otherwise, e.g., during movement
preparation (13, 14) (Figure 6C). While the potent- and null- subspaces hypothesis certainly seem to be
consistent with recorded neural activity in SC (Figure 4A), they cannot readily account for the
observation that neck (43) and upper limb (44) muscles are recruited time-locked to the visual target, as
proxied through electromyography. We instead show that temporal stability plays an independent role
in determining movement initiation, even when the putative potent subspace is matched in a causal
manipulation (Figure 4D, green symbols). We reason that for a decoder downstream, it is important to
ensure the stability - or consistency over time - of the input code while processing it in order to

influence the motor output. Population activity that creates a high firing rate drive but is unstable is also
likely unreliable and must have a reduced efficacy of triggering a movement. This requirement is critical
especially in the case of ballistic movements such as saccades, where the ability to reverse the decision
once the movement has been initiated is limited.

Thus, we present a novel mechanism, wherein both high firing rate and consistent temporal structure
are necessary conditions for movement initiation, allowing for population trajectories that traverse
away (increasing overall rate) but within hypersectors that fan out (stable temporal pattern) from the
state space origin (Figure 6D). Note that this model does not preclude population activity from
traversing a localized "optimal" region of state space, possibly due to hardwired network constraints.
Crucially moreover, the proposed mechanism is also consistent with classic population vector decoding
schemes (45, 46), since any weighted population vector readout during the sensory and movement
epochs will produce decoded movement vectors that are similar given a sufficiently large integration
window. This allows for the same physical movement to be planned and executed by a given population
of neurons in the labelled line sense, a mechanism that is precluded by weighted neural readout
mechanisms of movement preparation and generation, such as the potent/null space models.

Our findings are closely related to the premotor theory of attention (47) and offer a way to reconcile the
attention- intention debate. They could also account for the mirror-like activity recorded during both
action observation and execution in neurons known to project directly to motoneurons in the spinal
cord (48). In both cases, the same neuronal population represents two distinct signals that serve
different functional roles. The results presented here suggest that this multiplexing ability may be
provided by the distinct temporal structures of population activity patterns. Indeed, intracellular
recordings have demonstrated that visual stimulation drives cortical networks into an asynchronous
state (49), which may be a critical requirement for sensory processing.

Additionally, the differential effects of stable and unstable population stimulation patterns are
reminiscent of the desynchronizing effects of patterned deep brain stimulation recently developed for
the treatment of Parkinson's disease (50). In this so-called "coordinated reset" approach, sequential,
non-overlapping spatiotemporal sequences are more effective at resetting the affected neuronal
population from the pathological synchronized state to an asynchronous state, enabling better
movement control (51).

Previous studies have looked at the role of precise coordination in the timing of incoming spikes in the
transmission of information and the efficacy of driving the recipient neuron (20, 21). However, input
firing rates may vary greatly across the population, limiting the ability to compare to spike times. Our
study proposes a mean-field equivalent to the spike-based temporal or correlation code (22, 52) by
looking at the temporal structure of population firing rates, thus tying together the notion of rate,
temporal, and population codes. We suggest that temporal structure of population activity is critical to
understanding movement generation as well as, more broadly, neuronal communication and its
relationship to behavior.

Figure 1. **Temporal structure of population activity during sensorimotor multiplexing. A.** Left - Sequence of
events in the delayed saccade task. The top row shows a typical display sequence, and the bottom rows show the
timeline. The fixation target is shown as a plus symbol for illustration purposes. Dotted lines depict line of sight of
the animal. Right – Linear electrode arrays with 16-24 contacts were used to penetrate the SC normal to the
surface. Thus, all neurons encountered along the dorso-ventral axis of the SC prefer similar optimal vectors. **B.** Top:

Example recording session showing activity from 16 channels aligned on target (left column) and saccade (right
column) onsets for 3 different trials (traces with different color saturations). Bottom: Population activity averaged
across the 16 channels for the 3 example trials. Note the considerable variation in the amplitude of the population
visual burst across trials. **C.** Schematic depicting computation of temporal stability. The population vector (center)
was used to construct a single-trial neural trajectory, which was then normalized at each instance to represent a
unit length population vector traversing along a hypersphere. The left and right panels highlight unstable and
stable parts of a schematic trajectory. Note that in both cases, the two population vectors (thin arrows in green
and purple) are meant to be separated by a similar length of time. In the unstable case, the normalized vectors
(thick arrows) move around on the surface of the hypersphere (green and purple vectors are spatially separate),
whereas in the stable case, they stay pointed roughly in the same direction (the two vectors overlap in space). **D.**
Temporal stability is computed as a dot product between two normalized population vectors separated by
$2\tau = 20\ ms$ aligned on target (left) and saccade (right) onsets. The stability of population activity, averaged across
sessions, drastically decreases during the visual burst (red traces, mean +/- s.e.m.; p < 0.01 for Wilcoxon signed-
rank test with respect to baseline) and increases before and during movement onset (blue traces; p < 0.01 for
Wilcoxon signed-rank test with respect to baseline). The gray trace is the stability computed only for those trials
where the peak visual burst matched or exceeded activity during the premotor burst at saccade initiation (after
accounting for an efferent delay of 15 ms). It is effectively identical to the red and blue waveforms. The black trace
shows population stability on trials in which the target and saccade were directed to the opposite hemifield, far
away from the response field (RF) of these neurons. **E.** Temporal stability aligned to peak visual burst. The peak
time was computed on the average population response of each trial, and the peak-aligned trace was averaged
across trials and sessions. Note the sharper dip in the stability of the visual response when aligned to the peak. Like
in D, the gray trace shows peak-aligned stability for trials in which the peak visual activity matched or exceeded
premotor activity at saccade initiation. The gray and colored traces are nearly identical. F. Scatter plot of temporal
stability in a 20 ms window preceding the peak of visual and premotor bursts for individual sessions. Each point is
the median of the stability values computed on individual trials for one session (error bars are +/- 95% confidence
intervals). Filled circles indicate significant difference between the visual and premotor stability (p < 0.01, Wilcoxon
signed-rank test). The inset shows the scatter plot for individual trials for an example session (encircled in green).
Thus, temporal stability of visual and premotor epochs on individual trials also showed the same effect. **G.** Similar
to F but for the peak population firing rate during the visual and premotor bursts. The inset is for the same
example session as in F. For F and G, the premotor burst is truncated at 15 ms before saccade onset, the putative
efferent delay between SC and the extraocular muscles.

Figure 2. **Temporal stability in SC and FEF pseudo-populations. A.** Top row – Pseudo-population schematic. We combined single-neuron recording trials from different active populations on the SC map (left) into one pseudo-population (middle) that would be active for an average saccade, enabling the construction of a pseudo-population trajectory. A similar construction (not depicted) was performed for FEF. Middle row – Mean (+/- s.e.m.) normalized pseudo-population activity of visuomovement and motor neurons in SC (light red and blue traces) and FEF (dark red and blue traces) during the delayed saccade task. The black traces show the combined SC+FEF population putatively jointly projecting to the brainstem. The left and right panels show data aligned on target and saccade onsets, respectively. Bottom row – Temporal stability of the SC, FEF, and combined SC+FEF populations (colors as in the middle row). Pseudo-population stability shows a drastic reduction at the time of the visual burst and an increase prior to the onset of movement, consistent with the simultaneously recorded laminar population data in the previous figure. **B.** Top row – Schematic depicting active population regions in rostral and caudal SC. Middle row – Mean normalized pseudo-population activity in rostral SC neurons during the delayed saccade task (dark red and blue traces), overlaid with caudal SC activity (light red and blue traces as in panel A) and the combined rostral+caudal SC pseudo-population (black traces). Bottom row – Temporal stability of the rostral, caudal, and combined rostral+caudal pseudo-populations. Note that activity suppression in the rostral SC for large saccades is associated with a concomitant reduction in temporal stability. **C.** Population activity of rostral SC neurons in stable during microsaccades. Top row – Normalized population activity of microsaccade-related neurons in rostral SC (black trace). Note the strong burst compared to the suppression for large saccades in the same population (blue trace). Bottom row – Temporal stability of the rostral SC population for microsaccades and large saccades. Neurons which burst for microsaccades remain stable during these fixational eye movements, consistent with the hypothesis that both and increase in firing rate and high stability is required for movement generation. **D.** Top row – Sequence of events in the Gap task, designed to elicit express saccades. Middle row – Mean normalized pseudo-population activity on regular (red traces) and express (blue traces) saccade trials, as denoted in the inset of reaction time distribution. Bottom row – Population temporal stability decreases during the visual burst on regular saccade trials (red traces) like in the delayed saccade task but remains relatively high on express saccade trials in the same time period.

Figure 3. **Patterned microstimulation supports temporal stability as a model of movement initiation. A.** Example
pair of stable-unstable stimulation patterns. Top row – Pulse trains for 14 of 16 channels that were stimulation-
viable with suprathreshold parameters for this session. Red and blue trains indicate an example pair of unstable
and stable patterns, respectively. Second row – Pulse rates for the 14 channels in the stable-unstable pulse
patterns, with the population rates averaged across channels overlaid as thick, dashed traces. Note that the rates
in the unstable pattern are highly fluctuating despite the matched population average. Third row -- Average pulse
rates for each channel across all trials for the stable (blue) and unstable (red) patterns used in this session. The
clustering of trial-averaged rates in a narrow band is the intended result after randomization of pulse rate
assignment to individual channels on different trials. Bottom row -- Temporal stability of the stable and unstable
patterns (blue and pink, respectively) for all trials in the example session. Across-trials averages are shown as thick
traces. **B.** Scatter plot of saccade latencies (relative to stimulation onset) for an example session. Each point
reflects the outcome of stimulation with stable and unstable pulse patterns constituting each pair. The points in
shaded regions are stimulation-evoked for at least one condition. N_{s+us-}, points in the light blue shaded box, counts
trial pairs for which a stable stimulation pattern evoked a saccade but its unstable counterpart did not. N_{us+s-},
points in the light red shaded box, denotes the trial pairs for which an unstable stimulation pattern evoked a
saccade but its stable counterpart did not. Points in the purple box show subset of trials in which both stable and
unstable pairs evoked a saccade. Latency values are comparable for near-threshold stimulation parameters(*53,
54*). Points in the unshaded (white) region refer to trials in which neither stable nor unstable pattern of the pair

evoked a saccade. For these pairs, the trial was assigned the latency of the saccade (relative to stimulation onset)
directed to a target presented after the gap period. The graphic on the right shows the calculation of relative
likelihood of the stable pattern evoking a stimulation-evoked movement for a given session. **C.** Relative
proportions of the stable pattern evoking a movement for each session (blue points, error bars represent 95% CI of
the bootstrapped distribution). The sessions are sorted in descending order of proportion for viewing clarity. Note
that this sorting order is used in all subsequent figures depicting individual sessions (Figure 4D and Supplementary
Fig. 6B). Gray points and error bars are computed from a surrogate dataset in which stable/unstable trial identities
are completely shuffled (mean +/- 95% CI). Asterisks denote a significant effect for that session ($p < 0.01$ on the
permutation test).

Figure 4. Linear discriminability of population activity and microstimulation patterns. A. Three-dimensional factor analysis (FA) projection of population activity states during the visual and premotor bursts (red and blue points, respectively; 50 ms around peak of visual burst, 50 ms before saccade initiation in premotor burst. The dimensions were chosen arbitrarily to illustrate the separation of the two states. The solid lines and ellipses show the axes of maximum variance and covariance ellipses for each cluster, respectively. The histogram in the inset shows the performance of a linear discriminant analysis (LDA) classifier on labelling visual and premotor states on individual sessions (perfect classification is 1, chance is 0). **B.** FA projection of population pulse states for stable and unstable microstimulation patterns (blue and red points, respectively). The projection dimensions were chosen arbitrarily, but no view showed meaningful separation between the two clusters. The histogram in the inset shows LDA performance on classifying stable and unstable stimulation patterns, based on population pulse states alone (orange bars), or with a combination of population states and temporal stability (green bars). Note that perfect classification is expected by definition in the latter case. **C.** Schematic of the trials and data matrix used for classification of stimulation-evoked movement occurrence. Left - Trial pairs in which only one of the stable or unstable pattern evoked a movement. Right - The classifier was first trained on pulse counts alone (part of the matrix highlighted by the orange boundary – match color to the next panel). Separately, the classifier was also trained on an additional dimension of temporal stability values (full matrix highlighted by the green

boundary). **D.** LDA classification performance on discriminating trials in which stimulation evoked a movement
from those in which it did not. The orange points are with the classifier trained solely on the population pulse
states. The green points are with the addition of temporal stability. For most sessions, addition of temporal
information increased classification performance ($p < 0.01$, one-tailed t-test of a difference between the cross-
validated performance distributions with and without addition of temporal stability). Sessions are sorted by the
order determined in Figure 3C based on relative likelihood of evoking a movement. **E.** Scatter plot showing
correlation between classifier performance and behavior (values of blue points in Figure 3C rectified around 0.5).
The classifier's discriminative ability predicted the relative behavioral effect size between stable and unstable
patterns. **F.** Model selection for discriminating between trials in which a movement occurred versus those in which
it did not occur. Bayesian information criterion (BIC) traces as a function of model type for individual stimulation
sessions show that the lowest BIC (blue asterisks) was attained for the stability-only model for every session.

Figure 5. Spiking neuron model with dendritic processes can discriminate population temporal structure. A.
Schematic of the model architecture. The decoder (green neuron), representing high frequency burst neurons in
the brainstem, receives population inputs (output of purple neurons, or, in this case, microstimulation pulses)
through its dendritic network (excitatory inputs, represented by the arrowhead terminals). The disinhibitor (orange
neuron), representing pontine omnipause neurons, also receives inputs from the same population (both excitatory
and inhibitory(*33, 34*), represented by the triangular terminals), in addition to a constant current that produces
tonic firing activity (not shown). The decoder and disinhibitor are both modelled as leaky integrate-and-fire spiking
neurons and mutually inhibit each other (flat terminals). **B.** Schematic of the model heuristic. Each input neuron's
activation level is represented by its size and thickness. The effect of an input spike at the local dendritic site (i.e.,
excitatory post-synaptic conductance changes) is represented by the thickness of the arrow. Two time points are
shown for illustration (left column – arbitrary initial point, right column – subsequent time point). In the top row,

unstable inputs lead to a scenario where the post-synaptic conductances are no longer aligned to the strong inputs
that initially created them, whereas in the bottom, stable inputs lead to matched strong input and post-synaptic
conductances, resulting in stronger accumulation and firing. **C.** Scatter plot of first spike latencies in the decoder
(putatively representing saccade initiation) for stable versus unstable inputs from matched pairs. The input
duration was 150 ms and thus only latency values <150 ms correspond to actual first spikes produced by the
decoder. To facilitate comparison with Figure 3B, latency values were assigned randomly for trials in which the
decoder did not generate any spikes, sampled from a distribution with an arbitrarily selected mean of 350 ms.
Note the occurrence of a number of trial pairs for which only the stable input causes the decoder emit spikes (blue
arrow). **D.** Simulated membrane potential of the decoder (blue and red traces) and disinhibitor (cyan and magenta
traces) for an example matched trial pair with stable (top row) and unstable (bottom row) input patterns. **E.**
Average activity of the disinhibitor (cyan and magenta traces, stable and unstable inputs, respectively) and
decoder (blue and red traces, stable and unstable inputs, respectively) for all trial pairs in which only the stable
input produced a spike/movement (i.e., trial pairs indicated by the arrow in panel D).

Figure 6. Summary of models of movement preparation in a state space framework. A and **B.** Multi-dimensional
representations of the threshold hypotheses. Panel A depicts single neuron thresholds, i.e., the activity of each
neuron must rise to a fixed threshold at the same time in order to initiate a movement. Fixed thresholds for each
neuron are equivalent to a point (or a small region, represented by the gray sphere) in neural state space. Activity
profiles that meet this criterion (e.g., blue trajectory) and beyond (blue region) lead to movement generation while
those that don't meet this criterion (red traces) do not result in a movement. Alternatively, activity may need to
cross a fixed threshold at the population level (sum of neurons' activities = constant, i.e., an n-1 dimensional
hyperplane, dark gray plane), depicted in panel B. **C.** The optimal subspace hypothesis offers more leeway by
allowing neuronal activity to reach a relatively larger region of population state space (blue trajectory and gray
ellipsoid) over a period of time in order to signal the initiation of a movement. Activity trajectories that evolve
outside this "optimal subspace" (red traces) do not lead to a movement. **D.** The temporal stability hypothesis
suggests that a burst of neural activity that is consistent over time in state space, i.e., an activity trajectory that
points in the same direction (blue trajectories and vectors) is likely to be interpreted as a movement command by
a decoder, while high-frequency activity that is inconsistent (fluctuating directions, red trajectories and arrows) is
not. It does not matter where in state space the activity happens to evolve – a different subpopulation of neurons
could be active, but as long as they are pointing in the same direction (i.e., evolving within one of the blue sectors),
it will lead to a movement. The gray sphere around the origin represents a unit hypersphere for visualization of
back-projected unit vectors.

METHODS

Subjects and Surgical Procedures

All experimental and surgical procedures were approved by the Institutional Animal Care and Use Committee at the University of Pittsburgh and were in compliance with the US Public Health Service policy on the humane care and use of laboratory animals. We used three adult rhesus monkeys (*Macaca mulatta*, 2 male and 1 female, ages 6 - 12) for our experiments. Both SC and FEF were recorded in monkeys BB and BL whereas only SC was recorded in monkey WM. Under isoflourane anaesthesia, recording chambers were secured to the skull over craniotomies that allowed access to the SC and FEF. In addition, posts for head restraint and scleral search coils to track gaze were implanted. Analgesic and antibiotic treatment protocols were followed after surgery. Post-recovery, the animal was trained to perform standard eye movement tasks for a liquid reward.

Visual stimuli and behavior

Visual stimuli were displayed either by back-projection onto a hemispherical dome or on a LED-backlit flat screen monitor. Stimuli were white squares on a dark grey background, 4x4 pixels in size and subtended approximately 0.5° of visual angle. Eye position was recorded using the scleral search coil technique or using an EyeLink 1000 eye tracker, both sampled at 1 kHz. Stimulus presentation and the animal's behavior were under real-time control with a Labview-based controller interface(55). All monkeys were trained to perform standard oculomotor tasks. In the delayed saccade task (Figure 1A), the monkey was required to initiate the trial by aligning gaze on a central fixation target. Next, a target appeared in the periphery, but the fixation point remained illuminated for a variable 500-1200 ms, and the animal was required to delay saccade onset until the fixation point was extinguished (GO cue). The gap task (Figure 2D and Supplementary Fig. 5A) was used for express saccade analysis and for the patterned microstimulation experiments. In this task, initial fixation on a central target was followed by a gap period (200-300 ms) during which the fixation point disappeared, while the animal was required to maintain fixation at the now vacant location. On stimulation trials, microstimulation pulses were delivered 100 ms into the gap period and window constraints were relaxed to allow for the stimulation-evoked movements. This was followed by the appearance of a saccade target in the periphery which was also the GO cue for the animal to make a target-directed saccade. All animals performed these tasks with >95% accuracy. Incorrectly performed trials were removed from further analyses. The tasks were occasionally interleaved with visual search paradigms used for a different study.

Experimental sessions

We used data from several different types of sessions in this study. Each session was either a laminar recording session (n = 20 sessions), a single-unit recording session (n = 108 sessions), or a patterned microstimulation session (n = 16 sessions). The laminar and single-unit recording sessions employed mostly the delayed saccade task with a few microstimulation trials to verify electrode presence in SC (or FEF) and estimate the location on the topographic map. The patterned microstimulation sessions were restricted to the gap task, with stimulation trials and non-stimulation trials interleaved.

Laminar recordings and single-electrode electrophysiology

During each laminar recording session, a linear microelectrode array (LMA, AlphaOmega, Inc.) or a Plexon V-probe (Plexon, Inc.) was inserted into the SC chamber using a hydraulic microdrive. The arrays had either 16 or 24 contacts, with inter-contact spacing of 150 μm or 175 μm, respectively. Neural activity was amplified, digitized and recorded using the Grapevine Neural Interface Processor (Ripple, Inc.) and visualized using the associated Trellis interface. Neural activity was band-pass filtered between 500 Hz and 5 kHz to record spiking activity and between 0.1-250 Hz to record local field potentials for another study. Approach towards the SC surface was identified by luminance-based visual modulation of activity in the lowermost channels, after which the electrode was driven down a further 2-3 mm until known hallmarks of SC activity were observed in a majority of the channels. The presence of several contacts in the intermediate layers was further confirmed by the ability to evoke movements with single-channel microstimulation at these contacts. For the initial subset of experiments, a tungsten microelectrode was lowered into the FEF or SC. Neural activity was amplified and band-pass filtered between 200 Hz and 5 kHz and fed to a digital oscilloscope for visualization and spike discrimination. A window discriminator was used to threshold and trigger spikes online, and the corresponding spike times were recorded. Both SC and FEF were confirmed by the presence of visual and movement-related activity as well as the ability to evoke fixed vector saccadic eye movements at low stimulation currents (20-40 μA, 400 Hz, 100 ms). Before beginning data collection for a single or laminar recording site, the neural response field was roughly estimated. Since the electrode track was directed orthogonal to the SC surface, most neurons preferred similar visual response fields and/or movement vectors. In most data collection sessions with either electrode, the saccade target was placed either in the neurons' response field or at the diametrically opposite location in a randomly interleaved manner. In addition, stimulation-evoked saccades were recorded to identify the response field centers (or "hotspots") for the cells recorded during that session. For recordings in rostral SC, stimuli were presented at one of four locations at an eccentricity sufficient to induce a reduction in activity during the large amplitude saccade.

Data analysis and pre-processing

Data were analyzed using a combination of in-house software and Matlab. Eye position signals were smoothed with a phase-neutral filter and differentiated to obtain velocity traces. Saccades were detected using standard velocity criteria (30-50°/s). The animal was considered to be maintaining fixation if the gaze remained within a 2-3° window around the fixation target. We also detected any microsaccades that occurred during the delay period in each trial by using a velocity criterion based on the noise in the velocity signal for that trial. Only one of the two monkeys (WM) in whom we recorded neural activity in the rostral SC made sufficient number of microsaccades to permit further analysis.

Raw voltage signals filtered to spiking frequencies (500 Hz – 5 kHz, see above) were passed through a threshold criterion to obtain spike times corresponding to multi-unit activity (MUA). Spike density waveforms were computed for each neuron (or multi-unit activity cluster) and each trial by convolving the spike trains with the causal portion of a Gaussian kernel (width = 5 ms; in some instances, we used 10 ms for display purposes). Data from the patterned microstimulation experiments (n = 16 sessions)

were also subjected to this procedure to compute pulse density waveforms. For the laminar recordings,
we analyzed channel activity on single trials independently. Because direct kernel-based estimation of
firing rates from binary spike trains on single trials can be noisy, we first computed the inverse of the
inter-spike intervals as a measure of the firing rate prior to convolution with the smoothing kernel. All
units that crossed the spiking threshold of 6σ were considered for further analyses, regardless of their
characteristics (i.e., number of neurons/multi-unit clusters matched number of channels for these
datasets). Only sessions which had at least 6 channels with typical visuomovement activity patterns
were considered for further analysis, resulting in 16 laminar recording sessions (monkey BB – 104
channels/6 sessions, monkey BL – 176 channels/10 sessions). For single-electrode recordings, the spike
densities from condition-matched trials were randomly combined to form a pseudo-population dataset
(see next section). Neurons were classified as visuomovement neurons if the spike density was
significantly elevated above baseline during the visual epoch (50-200 ms following target onset) and
during the premotor epoch (50 ms before and after saccade onset). This resulted in 60 caudal SC
neurons and 30 FEF neurons. In addition, rostral SC neurons were defined as fixation-related if the
activity during the premotor epoch of large saccades was significantly reduced below baseline (36
neurons). A subset of these neurons also elevated their discharge around the onset of microsaccades
(13 neurons). To minimize the effect of noise in the spike density waveforms due to insufficient number
of trials in our analysis, we used only neurons which had at least 10 trials for a given condition. This was
not a factor in most cases (we typically had 50-100 trials). For the gap task analysis, we used neurons
from sessions that had at least 10 express saccade trials, defined as the set of trials in the first peak of
the saccade reaction time (SRT) histogram (SRT < 150ms). Regular saccade trials comprised the second
peak (SRT > 150ms).

To verify that spike isolation quality did not have an undue effect on our analyses, we also repeated the
main analysis from Figure 1 on spike sorted data. Spike sorting was performed using Kilosort2, an
algorithm developed for dense laminar probe recordings such as those with Neuropixels probes
(https://www.ucl.ac.uk/neuropixels/spike-sorting). The results of sorting were further manually
inspected to remove obvious misclassification errors. Sorted spike unit times were converted to spike
density waveforms using the same procedure as for MUA laid out above. When multiple units were
detected in a single laminar electrode channel, they were treated as distinct components of the
population vector corresponding to that session. The sorting produced 391 units over 280 channels
across 16 sessions (session counts – 12, 22, 23, 18, 24, 13, 40, 20, 27, 40, 23, 15, 29, 32, 29, 24 units).
The results of the analysis on spike sorted data strongly resembled the main temporal stability results
from MUA (compare Supplementary Fig. 3 and Fig. 1). Therefore, we restricted all analyses in the rest of
the paper to MUA, especially since spike sorting has been recently also found to have a negligible effect
on population-level analyses(23), and unsorted MUA may be more representative of the overall input
processed by downstream brain networks.

Estimating population dynamics from laminar and single-unit recordings

We define $\mathbf{R}(t)$ as a population activity vector:

$$\mathbf{R}(t) = \begin{bmatrix} R_1(t) \\ R_2(t) \\ ... \\ R_n(t) \end{bmatrix}$$

where $\mathbf{R}(t)$ represents the instantaneous activity at time t as a point in an n-dimensional space, n is the number of neurons, and $R_i(t)$ is the spike density function of the i^{th} neuron. The curve connecting successive points over time is the neural trajectory that describes the evolution of population activity. For laminar recording data, a neural trajectory was determined directly for each trial. Analysis of single electrode data relied on pseudo-population analysis, for which a surrogate dataset was created with individual trials drawn from each single neuron session. To ensure that these "pseudo-trials" were matched in relevant metrics, we first binned trials from each session into quartiles based on saccade reaction time and length of the delay period (the latter only for the delayed saccade task). A pseudo-trial was then created by drawing randomly from matched quartiles for each session, and this process was repeated 1000 times to create a full surrogate dataset from which pseudo-population trajectories were computed for individual pseudo-trials. Since SC and FEF neurons have fairly broad visual and movement fields, many neurons in each region contribute to the active population for any given visual stimulus or saccade(*56*), respectively. Our neurons were sampled roughly (but not exactly) around the hotspot of the active population for a given session. Therefore, the pooled data from individual sessions can be thought of as an approximation of the population mound active for an arbitrary location/RF in the visual field on any given trial. Many recent studies have reconstructed such pseudo-populations from sequentially recorded neurons and found comparable properties from simultaneous and serial recordings and found comparable properties from simultaneous and serial recordings(*13, 57, 58*). Indeed, this is also expected of our pseudo-population under the assumption of isotropy – that each neuron's contribution to its respective local active population is similar regardless of the locus of the population, and consistency between the results we observed in the laminar dataset and pseudo-population confirms this. To better demonstrate this, we estimated the location of a given neuron in the active pseudo-population as follows (we use SC for illustrative purposes because of its convenient topography). We mapped the target location onto the SC map and used this locus as a representation of the center of the active population for that session, and used the stimulation-evoked saccade vector to identify the location of that neuron on the SC. We referenced the mapped locus of each target location to a single location to create an active pseudo-population and translated the neuron locations relative to this population center (Supplementary Fig. 4A). All mathematical equations for transforming between visual and SC tissue coordinates have been defined previously(*59*). Inclusion of stimuli/saccades in the anti-preferred RF of the neurons allowed us to also estimate a pseudo-population of neurons in the ipsilateral SC. To complete the representation of activity across the SC topography, we also recorded from and included in our analyses neurons in the rostral portion of SC, which are active during fixation(*26*), burst during microsaccades(*28*), and are suppressed during large saccades(*26*).

812 **Temporal stability analysis**

813 To assess temporal stability, we first normalized the population trajectory $\mathbf{R}(t)$ by its Euclidean norm
814 ($\|\mathbf{R}(t)\|$), equivalently its magnitude, at each time point to yield $\widehat{\mathbf{R}}(t)$:

$$\widehat{\mathbf{R}}(t) = \frac{\mathbf{R}(t)}{\|\mathbf{R}(t)\|}$$

815 The normalized trajectory can be visualized as a unity length population vector that points in an n-
816 dimensional direction at each instant in time. That is, while $\mathbf{R}(t)$ is free to traverse the n-dimensional
817 activity space, $\widehat{\mathbf{R}}(t)$ is constrained to the surface of an n-dimensional hypersphere (Figure 1C). Temporal
818 stability or consistency of the evolving population was then quantified by the dot product of two time-
819 shifted unity length vectors:
820

$$S(t) = \widehat{\mathbf{R}}(t - \tau) \cdot \widehat{\mathbf{R}}(t + \tau)$$

The stability metric ($S(t)$) tracks the running similarity of the normalized trajectory separated in time by
2τ ms. Crucially, the normalization constrains $S(t)$ between 0 and 1. Thus, if $S(t) \rightarrow 1$, the population
activity is considered stable since the relative contribution of each neuron is consistent. If $S(t) \ll 1$, the
population activity is deemed unstable because the relative contribution is variable. If the neural
trajectory is not normalized, the dot product quantifies similarity across the vectors' magnitude and
direction. It roughly mimics the quadratic of the firing rate (exactly so for $\tau = 0$). When the vector
direction remains constant, the dot product yields no additional information than that already present in
the firing rate. In contrast, the normalization scales the neural trajectory and thus always has unity
magnitude. It neither alters the relative contributions of the neurons nor compromises the vector
direction. The dot product therefore performs an unbiased evaluation of stability based only on vector
direction and is an estimate of the fidelity of the population code modulo a multiplicative gain factor.
Intuitively, the stability measure is analogous to the correlation between the neurons' activities at two
different time points. We chose the dot product, however, because of its interpretability as a measure
of pattern similarity in n-dimensional activity space.

We assessed the significance of the stability profiles in two ways. First, we compared the across-trial
distribution of stability values in appropriate windows around the visual burst and during the premotor
epoch for each session (Figure 1F). For this analysis, the stability values were averaged in a 20 ms
window preceding the peak of the visual burst for the visual epoch and in a 20 ms window preceding the
saccade initiation time (15 ms before saccade onset) in the premotor burst for the premotor epoch.
Second, for each trial, we randomly shuffled the activities of various neurons (channels for laminar
recordings) at each time point (dark gray trace, Supplementary Fig. 2A, bottom row), or randomly
shuffled the time points at which the population patterns occurred during the trial (light gray trace,
Supplementary Fig. 2A, bottom row). The former shuffle retains the average firing rate at each instant
and the latter shuffle retains the overall population correlation structure but both shuffles remove any
temporal correlation in the firing rate across neurons. We performed multiple such shuffles and re-
computed temporal stability for each instance, followed by across-trial averaging, baseline correction,
and across-session averaging to obtain the distribution of stability profiles expected to occur by chance
if the neurons' activities were uncoordinated.

To mitigate the effect of potentially variable visual response onset latencies (on different trials) on the
trial- and session-averaged mean temporal stability profiles in the visual epoch, we also performed the
averaging after realigning the data to the peak visual response on individual trials (Figure 1E). In order to
do this, the time of peak visual burst was identified from the population mean (average across channels)
firing rate on individual trials and was used to align the traces across trials before computing the trial
average.

For the mean-matched control in Figure 1D&E, we performed the temporal stability analysis only on the
subset of trials in which the peak of the population visual response was greater than or equal to the
population activity at the time of saccade initiation, after accounting for an efferent delay. We chose 15
ms before saccade onset as the latest time at which SC activity could influence saccade initiation,
because it is consistent with a range of putative efferent delays reported using various approaches(*6, 16,*
17).

Patterned microstimulation

The patterned microstimulation experiments allowed us to causally evaluate and compare the temporal
stability model to other models of movement initiation. Given the ability to control the delivery of
individual stimulation pulses to each contact on the laminar electrode, we used stimulation patterns
with specific spatiotemporal features to evaluate their relative efficacy in evoking movements. We
designed these patterns as follows. All individual stimulation pulses were biphasic with a leading
cathodic phase, monophasic pulse widths of 200-267 µs, and an interphase interval of 100 µs. Our first
step was to determine the appropriate range of current intensities and frequencies for each experiment.
During each session (i.e., for each electrode penetration into SC, total n = 16 sessions), we first
stimulated from individual contacts with standard suprathreshold parameters (40 µA, 400 Hz) to identify
the set of contacts that elicited a movement at a sub-50 ms latency. We restricted the experiment to
these contacts for the rest of that session. We then stimulated simultaneously across all these contacts
at the same current and frequency, starting from 4 µA, 100 Hz, stepping up by 1 µA and 50 Hz, to
determine the threshold for evoking a movement with multi-channel stimulation. The threshold, defined
as the current/frequency combination for which stimulation generated movements on approximately
half the trials, ranged from 5-12 µA and 150-200 Hz across sessions. For each session, we used the
metrics of the saccades evoked at these threshold parameters from constant frequency stimulation in
lieu of "fixed vector saccades" as a control against which to compare the patterned microstimulation
results (Supplementary Fig. 5D&E).

Once we identified the threshold parameters, we designed stimulation patterns at that current intensity
but with time-varying frequencies intended to simulate a burst of activity. Each stable pattern was
created with a linearly decreasing inter-pulse interval (IPI) sequence for one contact (to simulate a burst)
and scaling the IPIs for other contacts by a uniformly spaced factor (see below). The 150 ms duration
burst was composed of a flat baseline (40 ms), a rising phase (80 ms), and a falling phase (30 ms). The
peak frequency for individual channels ranged uniformly between 20 Hz and 400 Hz with the mean peak
frequency across contacts pegged to the threshold frequency determined from the constant frequency
multi-channel stimulation in an earlier step. At these parameters, stimulation at no individual channel
evoked movements. Each stable stimulation pattern (e.g., blue pulse pattern in Figure 3A, top row) can
be envisioned as a $n \times m$ matrix, when n is the contact number and m is stimulation duration. The

matrix contains ones and zeros, identifying the time of pulse delivered to each contact. To create
additional realizations of the stable stimulation patterns (e.g., blue pulse patterns in Supplementary Fig.
5B & C), the assignment of specific peak frequencies to individual channels was randomly shuffled across
trials.

For each stable stimulation pattern, we created a corresponding unstable pattern/pair by jittering
stimulation pulses within a 20 ms window for each channel and randomly shuffling pulses across
channels (e.g., red pulse patterns in Figure 3A, top row and Supplementary Fig. 5B & C). This step
created instability in the population pulse pattern by destroying the relative scaling of pulse rates across
neurons while also ensuring that both total pulse counts per channel and the mean instantaneous pulse
rates across channels were preserved between the stable and unstable patterns. Thus, all subsequent
analyses were performed on pairs of trials with stable and unstable stimulation patterns matched in
these aspects. We also ensured that the inter-pulse interval was never less than 2 ms (i.e., the peak
frequency never exceeded 500 Hz) for any stimulation train. The stimulation patterns were generated
offline and the pattern trains corresponding to each trial were introduced 100 ms into the gap period in
a gap task. Stable-unstable pairs were randomly interleaved within a block in which roughly 80% of all
trials were stimulation trials.

We used the latency of the first saccade after stimulation onset but before stimulation offset as an
indicator of the occurrence of a stimulation-evoked saccade (Figure 3B); such saccades exhibited
latencies <160 ms. If the microstimulation was ineffective at evoking a saccade, the first movement was
typically directed to the target presented after the gap period ended; in such cases, the saccade was
produced >400 ms after stimulation onset. We estimated the relative likelihood of evoking a movement
by examining trials in which only one of the stable or unstable pulse patterns in a pair yielded a
stimulation-evoked saccade (also see Figure 3C and next section). This was quantified based on the
number of trials pairs in which only the stable pattern evoked a movement during the window of
stimulation (= N_{s+us-}, number of points in the blue shaded region in Figure 3B) and the number of trial
pairs in which only the unstable pattern evoked a movement (= N_{us+s-}, number of points in the red
shaded region in Figure 3B). The relative likelihood of evoking a movement with the stable pattern was
defined as

919
$$RL_s = \frac{N_{s+us-}}{N_{s+us-}+N_{us+s-}} .$$

RL_s ranges from 0 to 1 and is symmetric with a neutral value of 0.5. To estimate the significance of this
estimate, we performed permutation tests with respect to a surrogate dataset in which trials were
randomly assigned stable/unstable labels. This lowered the expected estimate of relative likelihood to
purely chance levels (Figure 3C).

Discriminability of population patterns – neural activity and microstimulation

To determine whether the visual and premotor bursts are discriminable using static, non-temporal
features of population activity, we performed factor analysis (FA) on 50 ms snippets of activity from
these two epochs (visual epoch: 50 ms centered around peak response, motor epoch: 50 ms before
saccade onset). We used DataHigh(*30*), a dimensionality reduction and visualization toolbox written in
Matlab, to perform FA on the 16- or 24-dimensional laminar recordings and visualize the projections

onto the top latent dimensions. The top 3 dimensions accounted for >95% of the variance in neural
states for all our datasets (eigenspectrum not shown), and a 3-dimensional projection is shown for an
example dataset in Figure 4A, with the visual and premotor snippets coded in different colors. We also
performed FA on the microstimulation patterns (the "states" were computed across the entire 150-ms
stimulation duration). Almost all dimensions were needed to account for >95% of the variance in the
stimulation patterns, owing to the randomized assignment of frequencies across contacts on different
trials. Thus, we chose an arbitrary 3-dimensional projection for visualization in Figure 4B; however, note
that the qualitative result did not depend on the projection. Since we used this step purely for
visualization, we did not perform any subsequent analysis on the estimated FA dimensions.

Next, we used a linear discriminant analysis (Fisher's LDA)(*60*) to assess a decoder's ability to discern
whether its inputs signify a movement command at the population level, based on static population
features alone, and how this discriminability relates to temporal stability. We first trained a binary LDA
classifier on the visual and premotor 50 ms snippets described above in the native neural space. LDA
finds a projection defined by a hyperplane that maximizes the separation between the two classes:

$$\boldsymbol{w} \propto \Sigma^{-1}(\boldsymbol{\mu_1} - \boldsymbol{\mu_0}) \,,$$

where 0 and 1 are the two class labels, the vector $\boldsymbol{w}$ is normal to the hyperplane defining the class
boundary, $\Sigma = \frac{n_0 \Sigma_0 + n_1 \Sigma_1}{n_0 + n_1}$, the pooled covariance matrix derived from the within-class covariance
matrices Σ_0 and Σ_1, n_0 and n_1 are class occupancies, and $\boldsymbol{\mu_0}$ and $\boldsymbol{\mu_1}$ are the class means. We subjected
the linear scores

$$s_i = \boldsymbol{w}. (x - \frac{\mu_i}{2})$$

obtained from the LDA projections to a softmax transformation to compute class probabilities:

$$P(x \in i) = \frac{e^{s_i}}{\sum_i e^{s_i}}$$

In all cases, we used k-fold crossvalidation to test LDA performance (on the classification of visual and
premotor population states, and on stimulation patterns described below), with the choice of k dictated
by the total number of trials in the dataset such that each fold (test set) had at least 10 trials. We
quantified classifier performance using the Matthews Correlation Coefficient (MCC)(*61*), since accuracy
can be significantly biased and have high variance for classification problems with unequal number of
training or test points(*62*), which was the case with the stimulation patterns below. Instead, MCC takes
the complete confusion matrix into account to quantify binary classifier performance as:

$$\text{MCC} = \frac{N_{00} N_{11} - N_{01} N_{10}}{\sqrt{(N_{00} + N_{01})(N_{00} + N_{10})(N_{11} + N_{01})(N_{11} + N_{10})}} \,,$$

where N_{ij} represents the number of instances where a trial in class i is classified to be in class j in the
classification confusion matrix. Similar to a standard correlation coefficient, MCC values range from -1 to
1 (-1, 0, and 1 respectively indicate poor, average, and perfect classification).

To evaluate the discriminability of microstimulation patterns, we split each dataset in 2 different ways.
The first split was based on the predefined stable-unstable pattern pairs and was used to classify pulse
patterns as stable or unstable based on their population states alone (inset in Figure 4B). The second

split was used to classify stimulation patterns as movement-evoking or non-evoking, in order to identify
other features of the stimulation patterns that may have impacted movement occurrence. For this
analysis, we focused on stable-unstable trial pairs where only one of the stable-unstable pair evoked a
movement (Figure 4C). The rationale was that a classifier trained on this subset may allow for the
identification of temporal stability as a predictor because of the differential (i.e., not one-to-one)
relationship between stability and movement occurrence. We further evaluated whether a dynamic
feature of population activity such as temporal stability can be exploited for this classification by re-
training the classifier with explicit addition of temporal stability as an input dimension and comparing it
with the performance of one without this addition.

We also performed model selection analysis using nested logistic regression models to identify the
predictor variable(s) in the microstimulation patterns that best explained the observed behavior. Logistic
regression models the log odds of an outcome (i.e., movement versus no movement) as a linear function
of desired predictors (population pulse states and temporal stability) as

979
$$\ln \frac{p}{1-p} = \beta_0 + \sum_{i=1}^{n} \boldsymbol{\beta}_i \boldsymbol{x}_i \, ,$$

where p is the probability of a movement occurring, i.e., P(stimulation evokes movement), $\boldsymbol{x}$ are the
predictors, and $\boldsymbol{\beta}$ are the model coefficients. We used a generalized linear model with a logit link
function and obtained the maximum likelihood estimate of the coefficients $\boldsymbol{\beta}$. We did this for models
with one or more predictors chosen from the set of population pulse counts and temporal stability.
Since the number of samples and/or predictors can have an effect on the log-likelihood of a given
model, we used the Bayesian information criterion (BIC) to estimate model performance, which
normalizes log-likelihood by a factor related to the number of samples and predictors. This analysis is
shown in Figure 4F.

To explain the variability in behavioral effects of stable/unstable microstimulation patterns across
sessions, we also considered whether certain sessions had idiosyncratic patterns of activation across
channels for trial subsets in which a movement was evoked. That is, we explored whether differences in
pulse rates on subsets of trials could explain the across-sessions range of movement likelihood effects
seen in Figure 3C. The rationale was that over many trials, individual channel pulse rates should average
out to a narrow distribution of rates given the random uniform sampling on individual trials. Thus,
discrepancies from this expected narrow distribution may reflect the effect of pulse rates alone for trial
subsets parsed by behavioral effect. We computed relative dispersion for a particular subset of trials as
the difference between the dispersion (i.e., pulse count variance across the population) for that subset
and dispersion on all trials for that session. That is, $RD_{subset} = var(x_{subset}) - var(x_{all})$, where x is
the distribution across channels of trial-averaged pulse counts for a set of trials, and subset refers to
S+US- or US+S- trials. The relative dispersion index was then defined as the relative dispersion for S+US-
trials divided by the sum of the two RDs:

1001
$$RDI = \frac{RD_{s+us-}}{RD_{s+us-} + RD_{us+s-}} \, .$$

1002 Thus, on sessions where the pulse rates for S+US- trials were closer to the distribution of rates on all
1003 trials (closer to the intended uniform sampling of pulse rates across channels), more so than the pulse
1004 rates for US+S- trials, the RDI is lower than 0.5 and vice versa. This is an indicator of pulse rate bias in

1005   one set of trials versus the other, with the unbiased subset (one with lower RDI) more likely to minimize
1006   the influence of pulse rates on the behavioral outcome. This analysis is shown in Supplementary Fig.5.

1007

Correlation analysis

In order to compare against pooled accumulator mechanisms(9), we estimated the correlation structure
in our population of neurons and microstimulation patterns. We first computed the spike count
correlation across trials for every pair of neurons (or multi-unit clusters, n = 1963 pairs) during the visual
and motor epochs as defined above. For the microstimulation patterns, we computed the pulse count
correlations for every pair of stimulation channels (n = 899 pairs) during the stimulation window for the
stable and unstable patterns separately. Note that in this case, pulse count correlation is equivalent to
computing the correlation between accumulation rates (and therefore directly comparable to the
accumulator mechanism mentioned above), since the duration of stimulation was fixed and the baseline
and peak rates were matched between the stable and unstable conditions.

1018

Spiking neuron model

We developed a biophysically realistic computational model with spiking neurons to discriminate and
read out temporal structure in population activity. The core component of the model was a recurrently
connected module comprising a decoder element and a disinhibitor element, putatively representing
pontine high-frequency burst neurons and OPNs, respectively(37). The decoder and disinhibitor were
both modelled as leaky integrate-and-fire (LIF) spiking neurons:

$$\tau_m \frac{dV_{neu}}{dt} = -(V_{neu}(t) - V_{rest}) + R_m I_{neu}(t)$$

where $V_{neu}(t)$ is the membrane potential of the neuron, V_{rest} is the resting potential, τ_m is the
membrane time constant, R_m is the membrane resistance, $I_{neu}(t)$ is the net time-varying input to the
neuron, and *neu* represents the decoder (*dec*) or disinhibitor (*dis*). The spiking followed standard
threshold crossing rules, i.e., if $V_{neu}(t = t^k) > V_{th}$, where V_{th} represents the action potential
threshold, the membrane potential was reset to $V_{neu}(t = t^k) \equiv V_{reset}$, and the neuron's spike times
$[t^i]$ were updated with the k^{th} spike time t^k.

The current input to both the decoder and disinhibitor was composed of excitatory and inhibitory
components:

$$I_{neu}(t) = G_{neu}^{exc}(t)\big(E_{exc} - V_{neu}(t)\big) + G_{neu}^{inh}(t)\big(E_{inh} - V_{neu}(t)\big) + I_{neu}^0$$

where $G_{neu}^{exc}(t)$ and $G_{neu}^{inh}(t)$ are the respective net excitatory and inhibitory post-synaptic conductances,
and E_{exc} and E_{inh} are the excitatory and inhibitory reversal potentials representing the contribution of
AMPA and GABA channels, respectively. I_{neu}^0 is a constant current term that was set to zero for the
decoder and to a fixed value of 4 for the disinhibitor in order to simulate tonic firing without external
input.

The decoder received excitatory inputs from the SC in the form of stimulation pulse trains, and
inhibitory input from the disinhibitor. The disinhibitor received both excitatory and inhibitory inputs
from the SC(*33, 34*) and reciprocal inhibition from the decoder. The net time-varying conductances were
modelled as follows:

$$1042 \quad G_{dec}^{exc}(t) = \sum_{i=1}^{N} w_i^{exc-dec} g_i(t) \ , \ \text{ and } \ g_i(t) = \sum_{k=1}^{n_i} \epsilon\left(t - t^k\right)\left(1 + g_i\left(t^k\right)\right) \ ,$$

$$1043 \quad G_{dec}^{inh}(t) = w^{dis-dec} \sum_{k=1}^{n_{dis}} \epsilon\left(t - t^k\right) \ ,$$

$$1044 \quad G_{dis}^{exc}(t) = \sum_{i=1}^{N} w_i^{exc-dis} g_i(t) \ , \ \text{ and } \ g_i(t) = \sum_{k=1}^{n_i} \epsilon\left(t - t^k\right)\left(1 + g_i\left(t^k\right)\right) \ ,$$

$$1045 \quad G_{dis}^{inh}(t) = w^{dec-dis} \sum_{k=1}^{n_{dec}} \epsilon\left(t - t^k\right) + \sum_{i=1}^{N} w_i^{inh-dis} g_i(t) \ , \ \text{ and } \ g_i(t) = \sum_{k=1}^{n_i} \epsilon\left(t - t^k\right)\left(1 + g_i\left(t^k\right)\right).$$
1046

In each equation they appear, N is the number of input units (i.e., number of channels with pulse trains),
$w_i^{exc/inh-neu}$ is the excitatory/inhibitory weight from the i^{th} input unit to the spiking neuron, n_i is the
number of pulses in the i^{th} input unit, $w^{dis-dec}$ is the weight from the disinhibitor to the decoder,
$w^{dec-dis}$ is the weight from the decoder to the disinhibitor, and t^k is the time of the k^{th} input pulse or
output spike (depending on which summation term it appears in). The g_i term, when it appears in the
sum that produces g_i (as in all but the second conductance equation), represents the state-dependent
influence of each incoming pulse input. $\epsilon(t)$ is the post-synaptic conductance kernel defined as:

$$\epsilon\left(t - t^k\right) = \begin{cases} -e^{-\frac{t}{\tau_1}} + e^{-\frac{t}{\tau_2}} \ if \ t > t^k \\ 0 \ if \ t < t^k \end{cases},$$

1054 The conductance and input current terms that were dependent solely on the pulse input patterns were
1055 pre-computed and fed into a Eulerian solver (time step = 0.1 ms) for the differential equations governing
1056 the membrane potential of the spiking decoder and disinhibitor units. Supplementary Table 1 shows the
1057 values of the parameters and constants used for the model simulations.

1058

1059

1060

Acknowledgements: We thank Drs. A. Batista, C. Olson, M. Smith, and B. Yu for scientific discussions and
critical feedback on previous versions of the manuscript and J. McFerron for programming assistance.

Funding: The study was funded by the following NIH R01 grants: EY022854 and EY024831.

Author Contributions: U.K.J and N.J.G conceived and designed the study. U.K.J. performed the
experiments, analyzed the data, and performed model simulations. U.K.J and N.J.G wrote the
manuscript.

Competing Interests: The authors declare no competing financial interests.

Data and code availability: The data that support the findings of this study are available from the
corresponding author(s) upon reasonable request. The custom MATLAB code is available from the
authors upon reasonable request.

1071

Supplementary Figure 1. Strong saccade drivers also have high visual activity. A. Scatter plot of the motor potential, calculated as the peak correlation between saccade velocity and preceding premotor activity (see Methods), as a function of peak premotor burst for individual neurons. Each point represents a neuron/MUA recorded on an individual channel, and each line is the linear regression fit for the points from an individual, laminar probe recording session (bold lines indicate significant trends, p < 0.01). **B.** Neurometric plot of the proportion of neurons in each of 4 quartiles split based on peak premotor activity that have a motor potential (as defined above) higher than the abscissa. The traces are colored based on their quartile as indicated in the legend. The vertical line and arrowhead indicate the median motor potential. For example, roughly 80% of the neurons in the highest premotor activity quartile (yellow trace) show an activity-velocity correlation above the median and are thus considered strong saccade drivers. **C.** Scatter plot similar to **A** except the abscissa shows peak visual activity. A strong correlation between the neurons' peak visual activity and its motor potential, which is still computed using its premotor activity, is observed for several sessions and in the overall distribution. **D.** Neurometric plot similar to **B** of the proportion of neurons in each of 4 quartiles split based on peak *visual* activity that have a motor potential higher than the abscissa. The majority of neurons with high visual activity (>60% in top 2 quartiles, green and yellow traces) exhibit motor potential above the median (vertical line and arrowhead), indicating that neurons with high visual activity also strongly influence saccade kinematics and thus are candidates for causal drivers of the movement.

1089

1090

Supplementary Figure 2. Temporal stability profiles within and across laminar recording sessions. A. Top row: Trial-averaged temporal stability traces for individual sessions (thin traces) have different baselines, possibly due to varying noise levels, spike isolation differences, etc. The baselines were shifted to the across-session mean level before session-averaging in Figure 1D. Note that the raw and not baseline-shifted values were used for statistical analyses. Bottom row: Temporal stability averaged across trials and sessions for two types of shuffled datasets. The light gray traces (mean +/- s.e.m.) are for time-shuffled data where relative activity levels between neurons are preserved but time points during the trial are randomly re-ordered. The dark gray traces are for neuron-shuffled data where the instantaneous population firing rate is preserved but the firing rate distribution is shuffled across neurons at each time point. The colored traces show the session-averaged temporal stability (as in Figure 1D) for comparison. **B.** In order to ensure that the choice of τ did not have an undue effect on our results, we computed stability for several values of τ. The panels show temporal stability profiles for (from top to bottom), τ = 1, 2.5, 5, 10, and 20, respectively. The colored traces for τ =10 are the ones shown in the Figure 1D. As evident, the absolute magnitude of stability was inversely related to τ, a consequence of the smooth and continuous nature of the trajectory. In other words, since the state of the neural population evolves smoothly, it must traverse intermediate states in order to move from one state to another, resulting in greater similarity between state vectors close together in time than those further apart. However, the relative shape of stability profile was largely preserved for $\tau > 5$. **C.** Similar to top row of panel A but for temporal stability aligned to peak of visual burst.

1108

Supplementary Figure 3. Spike sorting does not affect temporal structure findings. A. Similar to Figure 1B but for
spike sorted data. Top left: Example laminar recording trial showing activity from 16 channels aligned on target
(left column) and saccade (right column) onsets. Multiple spike sorted units in a channel (if any) are shown as
different traces. Bottom: Population activity averaged across the 16 channels for the example trial. Top right:
Corresponding mean waveforms of spike sorted units in each channel, with multiple units from a channel shown as
different traces. **B.** Temporal stability for spike sorted laminar recordings shows a similar trend to the unsorted
multi-unit activity data in the main figures (dip during the visual epoch and increase in the premotor epoch, p <
0.01 for Wilcoxon signed-rank test with respect to baseline , cf. Figure 1D). **C.** Temporal stability for spike sorted
data aligned to peak visual burst, similar to Figure 1E. **D**. Scatter plot of mean peak population firing rate from
spike sorted data for individual sessions, with trials from an example session shown in the inset (cf. Figure 1G). **E.**
Similar to D but for temporal stability in a 20 ms window preceding the peak visual and premotor bursts (cf. Figure
1F). In D and E, filled circles indicate significant difference between the corresponding values computed during the
visual burst and the premotor burst (p < 0.01, Wilcoxon signed-rank test).

1122

1123

1124

1125

Supplementary Figure 4. Reconstructed locations of neurons in pseudo-population on SC map. *Top row* - Target
loci (red points) mapped from visual space to SC coordinates across all experimental sessions. The locations on the
SC were computed using traditionally established transformations between visual space and SC tissue coordinates
(see Methods). *Middle row* - Endpoints of the stimulation-evoked saccade vector at the recording site (blue points)
shown with the target loci from above. The stimulation vector provides a proxy for the RF center of the recorded
neuron. Neuron-target pairs (blue-red) from individual sessions are connected using black lines (unconnected
points did not have the corresponding stimulation/target data). *Bottom row* - The active pseudo-population on the
SC map. The red locations from the previous panels were referenced to one arbitrarily selected location on the SC
map (here, R=15, theta = 0) and the blue locations were translated accordingly. Points from both colliculi are
shown on a single SC for the sake of clarity.

1136

Supplementary Figure 5. Microstimulation experiment - task, examples, and evoked saccade metrics. A.
Schematic of the gap task used in the patterned microstimulation experiments. **B** and **C.** Two more examples of
pulse rasters and pulse rates for individual channels for stable (blue) and unstable (red) pairs of stimulation
patterns (cf. Figure 3A). **D.** Mean stimulation-evoked saccade amplitude for stable (blue points) and unstable (red
points) stimulation patterns plotted against the fixed vector saccade amplitude obtained from near-threshold
constant frequency stimulation (see Methods). The fixed-vector saccade is also known as the site-specific saccade.
Each point represents one session for that condition. The diagonal represents the unity relationship. Filled circles
denote a significant difference of the stable/unstable saccade amplitude from the fixed vector saccade (p < 0.01,
Wilcoxon rank-sum test). Points circumscribed by black circles denote sessions where the saccade amplitudes on
stable and unstable trials were significantly different from each other (p < 0.01, Wilcoxon signed-rank test). **E.**
Similar to D (including criteria for statistical significance), but for stimulation-evoked saccade direction.

1148

1149

Supplementary Figure 6. Variability in behavior is partly explained by variability in stimulation pulse rates across subsets of trials. A. Individual channel pulse rates averaged across specific subsets of trials for an example session. *Left* - Average stable pattern for the set of trials in which, only the stable pattern evoked a movement out of the stable-unstable pair (S+US- pairs, cyan traces), only the unstable pattern evoked a movement (S-US+ pairs, magenta traces), and all trials (gray traces). *Middle* - Schematic of which trial pairs are plotted in the left and right columns, highlighted in the appropriate color. Relative dispersion is calculated as the across-channel dispersion (variance) of the pulse counts for a particular subset (e.g., S+US-) minus the dispersion of the pulse counts for all trials. *Right* – Average unstable pattern for the three subsets of trials as in the left column. **B.** Relative dispersion index computed as in the formula in the inset, plotted for each session (green points). Values *less than* 0.5 indicate that there was less dispersion in the S+US- trials, and trials with lower dispersion are less dependent on pulse rate effects. The behavioral effect (from Figure 3C) is overlaid for comparison (blue points and error bars). The sessions are ordered by behavioral effect size as in Figure 3C. **C.** Scatter plot showing correlation between behavioral effect size and relative dispersion index. The sessions with negative (ordinate < 0.5) or non-existent effect sizes in behavior are also the ones with the highest dispersion index, indicating that those sessions were dominated by pulse rate effects and less likely to reveal an effect of temporal structure.

1165

1166   38

1167

1168

Supplementary Figure 7. Across-trial spike and pulse count correlations. A. Left: Histograms of spike count
correlations for all pairs of channels (n = 1963) for the visual and premotor epochs (red and blue histograms,
respectively). One of the histograms is shown inverted for the sake of clarity. Right: Distribution of differences
between visual and premotor epoch correlations computed for each pair. Positive values indicate larger spike
count correlations for spikes in the visual epoch. The two vertical lines represent zero and the mean of the
distribution (left and right lines, respectively). **B.** Same as A for pulse counts correlations for all pairs of stimulation
channels (n = 899) for the stable (blue) and unstable (red) patterns, with the corresponding difference histogram
shown on the right. Positive values indicate higher pulse count correlations in the unstable stimulation pattern.

1177

1178

Supplementary Figure 8. Performance of spiking neuron model for various simulation outcomes with stable and unstable pulse patterns. A. Scatter plot of first spike latencies in the decoder (putatively representing saccade initiation) for stable versus unstable inputs from matched pairs (same as Figure 5C). The next two panels are based on the trial pairs highlighted by the arrow (in which both stable and unstable patterns produced a saccade). **B.** Simulated membrane potential of the decoder (blue and red traces) and disinhibitor (cyan and magenta traces) for an example matched trial pair from the subset in panel A, for stable (top row) and unstable (bottom row) input patterns. **E.** Average activity of the disinhibitor (cyan and magenta traces, stable and unstable inputs, respectively) and decoder (blue and red traces, stable and unstable inputs, respectively) for all trial pairs in which both stable and unstable input patterns produced a movement. **D.** Same as panel A, except for the focus on trial pairs (arrow). The next two panels are based on the trial pairs in which neither stable nor unstable patterns produced a saccade. **E.** Simulated membrane potential of the decoder and disinhibitor for an example matched trial pair from the subset in panel D. Layout and color scheme as in panel B. **F.** Average activity of the disinhibitor and decoder for all trial pairs in which neither stable nor unstable input patterns produced a movement. Color scheme as in panel C. Note the small increase in disinhibitor activity around the time when a movement would have normally occurred, reminiscent of the slight increase in OPN activity during visual input(*38*).

1194

Supplementary Figure 9. Leaky accumulator with facilitation (LAF) model can discriminate population temporal structure. A. Inputs to the model from SC visuomovement neurons. Raw unstable (red) and stable (blue) input profiles (inset – population means). The two sets of inputs are 180 ms snippets taken from the visual and premotor bursts, respectively, in the spike density profiles shown in Figure 2A. **B.** Mean-matched input profiles and population means. **C.** Temporal stability of the two mean-matched populations. **D.** Output of the LAF accumulator in response to the stable and unstable model inputs. **E.** Evolution of synaptic weights for the two conditions. **F.** Correlation between instantaneous weights and input rates for the two conditions. **G.** Pattern sensitivity index of the model's ability to discriminate between the two types of inputs. Values in the top half of the plot indicate higher sensitivity (faster accumulation) to stable population input.

1204

1205   **Supplementary Table 1. Values of the constants and parameters used for the spiking neuron model.**

Biophysical constants	Values
τ_m	10 ms
R_m	10 Mohms
V_{th}	-50 mV
V_{rest}	-65 mV
V_{reset}	-70 mV
E_{exc}	50 mV
E_{inh}	-80 mV
Model parameters	
τ_1	2 ms
τ_2	10 ms
$\boldsymbol{w}^{exc-dec}$	$5 \times 10^{-6} \; uniform[0,2]$
$w^{dis-dec}$	5
$\boldsymbol{w}^{exc-dis}$	$0.2 \, \boldsymbol{w}^{exc-dec}$
$\boldsymbol{w}^{inh-dis}$	$0.8 \, \boldsymbol{w}^{exc-dec}$
$w^{dec-dis}$	0.1
Numerical parameters	
dt (time step)	0.1

1206

S1. Inferring population dynamics from single-unit recordings

The main text and methods describe how we reconstructed a pseudo-population from unit recordings. Using knowledge of the RF center of a given session's local population obtained from microstimulation, and known transformation from visual space to SC tissue coordinates, here we reconstruct the pseudo-population on the SC map.

We first transferred target locations (R_T, θ_T) inside the RF for each neuron onto the SC map using the following transformations(59) –

$x_T = \frac{B_x}{2} \ln \frac{(H+A)^2 + V^2}{A^2}$, $y_T = B_y \tan^{-1} \frac{V}{H+A}$, where,

$H = R_T \cos \theta_T$, $= R_T \sin \theta_T$, and $A = 3$, $B_x = 1.4$, $B_y = 1.8$.

The transformed locations (x_T, y_T) on the bilateral SC map are shown in Supplementary Fig. 3A, top row. In order to move these target locations to one pseudo-population, we need to identify where these points reside in the RF of each neuron or local population. We used the endpoints of the site-specific microstimulation-evoked saccades as a proxy for the respective RF centers. The transformed endpoints, with their locations relative to the corresponding target locations, are shown in Supplementary Fig. 3A, middle row. We picked an arbitrary location in the visual field as the RF center of the pseudo-population, $(R_c, \theta_c) = (15,0)$. We used the fact that the size of the active SC population is invariant regardless of the encoded saccade to preserve relative distances in tissue space between a site's target location and RF centre, and translated the transformed microstimulation endpoints to the common pseudo-population center, along with the respective target locations. To construct one active population from sites gleaned from both hemifields (and therefore both colliculi), we reflected the coordinates from one colliculus onto the other. The resulting pseudo-population (Supplementary Fig. 3A, bottom row) shows a fairly representative sampling of neurons from the pseudo-population, with its extent consistent with a large (25% of SC tissue) active population, albeit one that is biased to one side of the population.

S2. Leaky accumulator with facilitation (LAF) model

We developed an alternative computational model to demonstrate how the temporal structure of
population activity could be used by neural circuits to gate decisions such as movement initiation.
Similar to the spiking neuron model presented in the main text, a core principle behind this model is the
hypothesis that signal integration is stronger when the temporal structure in input activity is stable,
which allows the decoder neuron to reach threshold during the high frequency burst that triggers the
movement. We constructed an accumulator as an abstraction of a neuron receiving population inputs
through its network of dendrites. The total synaptic current $I(t)$ and the firing rate $v(t)$ of the
accumulator were defined as

$$\tau_s \dot{I} = -I + \sum_1^n w_i u_i$$

1241 and

$$v = F(I)$$

where u_i and w_i are the instantaneous firing rate and synaptic gain of the i^{th} input neuron, and τ_s is the
time constant of the synaptic current. The output firing rate of the single decoder neuron $v(t)$ can be
described by a standard monotonic function (e.g., linear or sigmoid) applied to the net current(*63*). The
family of stochastic, leaky accumulator models that integrate the firing rate of neurons toward a
threshold criterion has been commonly used to explain reaction times, decision making, and
perception(*9, 64-66*). We used the following heuristic in order to extend this framework to incorporate
temporal structure. In order to assess stability, the decoder neuron must keep track of the short term
history of the population activity, use this "memory" to evaluate stability, and respond selectively when
the activity pattern is deemed stable over some time scale. We implemented these requirements by
introducing short-term plasticity in the form of facilitatory connections(*67, 68*) from the input
population onto the output unit (accumulator). The change in connection strength or gain of each
neuron on the decoder neuron w_i can be defined by a differential equation that incorporates a Hebbian-
like learning rule and leak current:

$$\tau_w \dot{w}_i = -w_i + \frac{u_i v}{wnf_i}$$

The Hebbian-learning component, $u_i v$ describes the change in weight coupled to the firing rates of the
i^{th} pre-synaptic neuron and the post-synaptic accumulator neuron. τ_w is the time constant of the weight
parameter. Since this version of the model contains only excitatory connections, the weight parameter
can exhibit unbounded accumulation, which can be controlled by incorporating normalization. We
normalized the Hebbian-learning component $u_i v$ with the contribution of that neuron to the total
resource pool(*69*). The resource pool available for facilitation at any given time was defined as the sum
of the Hebbian-learning component across all input units $\sum_1^n u_i v$. The contribution of a neuron to the
output rate $v(t)$ then determines its contribution to the overall pool, giving rise to the weight
normalization factor:

$$wnf_i = \sum_1^n u_i w_i u_i$$

For this model simulation, we used visuomovement neuron activity drawn from the SC pseudo-
population as inputs (Supplementary Fig. 8A). We created two sets of input snippets from the visual and
premotor bursts, each 180 ms in length. For the visual burst, we used the activity upto the peak in the
population response. For the premotor burst, we used activity upto 35 ms before saccade onset (which
was the time when the response magnitude reached the same level as the peak of the visual burst). In
order to control for the effect of average firing rate on the accumulation, we mean-matched the input
profiles as follows. We divided the activations in the premotor input set at each instant by the mean
activation of the visual inputs at the corresponding instant. That is,

$$u_i^{pre}(t)_{mm} = \frac{u_i^{pre}(t)}{\frac{1}{n}\sum_1^n u_i^{vis}(t)}$$

where $u_i^{vis}(t)$ and $u_i^{pre}(t)$ are the activity of neuron i at time t in the visual and premotor input sets,
respectively, and $u_i^{pre}(t)_{mm}$ is the premotor input instantaneously rescaled to match the mean of the
visual input (Supplementary Fig. 8B). This ensured that we isolated the effect of temporal structure
(Supplementary Fig. 8C), independent of firing rate, on the model's response.

The accumulator increased its activity at a faster rate in response to the stable pattern compared to the
unstable pattern (Supplementary Fig. 8D), suggesting that the network was able to discriminate
between two types of population patterns even though the net input drive was identical. How critical is
facilitation to this function? Like the inputs themselves, the weights also showed greater fluctuation
during the visual burst (Supplementary Fig. 8E). Consistent with our heuristic, the weights tracked the
input rates when the input was stable, but this correlation dropped away when the input was unstable
(Supplementary Fig. 8F). Note that the shape of the correlation profile is not unlike the stability profile in
Supplementary Fig. 8C. Therefore, facilitation allows the weights to retain the memory of a pattern over
the time scale of tens of milliseconds, but the memory can be scrambled by a fluctuating input pattern.

We also quantified the ability of the synaptic weights to track the inputs by computing the Pearson's
correlation between the weight and input vectors at each time point (Supplementary Fig. 8G). To
quantify the accumulator's ability to discriminate temporal pattern in population input, we computed a
pattern sensitivity index (*psi*) as

$$psi(t) = \frac{v^{pre}(t) - v^{vis}(t)}{v^{pre}(t) + v^{vis}(t)}$$

1289　where $v(t)$ is the output of the accumulation for the corresponding inputs.

1290　The results here suggest that the accumulator is driven more strongly by the stable premotor burst,
1291　even when other features of population activity remain the same, providing yet another biophysical
1292　mechanism by which temporal structure could be read out.

1293

1294

1295

1296

References

1.   R. H. Wurtz, M. A. Sommer, M. Paré, S. Ferraina, Signal transformations from cerebral cortex to
superior colliculus for the generation of saccades. *Vision Res* **41**, 3399-3412 (2001).
2.   C. A. Buneo, M. R. Jarvis, A. P. Batista, R. A. Andersen, Direct visuomotor transformations for
reaching. *Nature* **416**, 632-636 (2002).
3.   V. Gallese, L. Fadiga, L. Fogassi, G. Rizzolatti, Action recognition in the premotor cortex. *Brain : a*
journal of neurology **119 (Pt 2)**, 593-609 (1996).
4.   C. K. Rodgers, D. P. Munoz, S. H. Scott, M. Pare, Discharge properties of monkey tectoreticular
neurons. *J Neurophysiol* **95**, 3502-3511 (2006).
5.   M. A. Segraves, Activity of monkey frontal eye field neurons projecting to oculomotor regions of
the pons. *J Neurophysiol* **68**, 1967-1985 (1992).
6.   D. P. Hanes, J. D. Schall, Neural control of voluntary movement initiation. *Science* **274**, 427-430
(1996).
7.   R. Ratcliff, J. N. Rouder, Modeling response times for two-choice decisions. *Psychol Rev* **9**, 347-
356 (1998).
8.   J. J. Jantz, M. Watanabe, S. Everling, D. P. Munoz, Threshold mechanism for saccade initiation in
the frontal eye field and superior colliculus. *J Neurophysiol* **109**, 2767-2780 (2013).
9.   B. Zandbelt, B. A. Purcell, T. J. Palmeri, G. D. Logan, J. D. Schall, Response times from ensembles
of accumulators. *Proceedings of the National Academy of Sciences of the United States of*
America **111**, 2848-2853 (2014).
10.   B. M. Yu *et al.*, Gaussian-process factor analysis for low-dimensional single-trial analysis of
neural population activity. *J Neurophysiol* **102**, 614-635 (2009).
11.   K. V. Shenoy, M. Sahani, M. M. Churchland, Cortical control of arm movements: a dynamical
systems perspective. *Annu Rev Neurosci* **36**, 337-359 (2013).
12.   M. M. Churchland *et al.*, Neural population dynamics during reaching. *Nature* **487**, 51-56 (2012).
13.   M. T. Kaufman, M. M. Churchland, S. I. Ryu, K. V. Shenoy, Cortical activity in the null space:
permitting preparation without movement. *Nature neuroscience* **17**, 440-448 (2014).
14.   G. F. Elsayed, A. H. Lara, M. T. Kaufman, M. M. Churchland, J. P. Cunningham, Reorganization
between preparatory and movement population responses in motor cortex. *Nat Commun* **7**,
13239 (2016).
15.   A. Afshar *et al.*, Single-trial neural correlates of arm movement preparation. *Neuron* **71**, 555-564
(2011).
16.   N. Miyashita, O. Hikosaka, Minimal synaptic delay in the saccadic output pathway of the
superior colliculus studied in awake monkey. *Exp Brain Res* **112**, 187-196 (1996).
17.   U. K. Jagadisan, N. J. Gandhi, Removal of inhibition uncovers latent movement potential during
preparation. *Elife* **6**, (2017).
18.   I. Smalianchuk, U. K. Jagadisan, N. J. Gandhi, Instantaneous Midbrain Control of Saccade
Velocity. *J Neurosci* **38**, 10156-10167 (2018).
19.   P. J. May, J. D. Porter, The laminar distribution of macaque tectobulbar and tectospinal neurons.
Vis Neurosci **8**, 257-276 (1992).
20.   T. Branco, B. A. Clark, M. Hausser, Dendritic discrimination of temporal input sequences in
cortical neurons. *Science* **329**, 1671-1675 (2010).
21.   A. G. Carter, G. J. Soler-Llavina, B. L. Sabatini, Timing and location of synaptic inputs determine
modes of subthreshold integration in striatal medium spiny neurons. *J Neurosci* **27**, 8967-8977
(2007).

22. M. A. Harvey, H. P. Saal, J. F. Dammann, 3rd, S. J. Bensmaia, Multiplexing stimulus information
through rate and temporal codes in primate somatosensory cortex. *PLoS biology* **11**, e1001558
(2013).

23. E. M. Trautmann *et al.*, Accurate Estimation of Neural Population Dynamics without Spike
Sorting. *Neuron* **103**, 292-308 e294 (2019).

24. C. J. Bruce, M. E. Goldberg, Primate frontal eye fields. I. Single neurons discharging before
saccades. *J Neurophysiol* **53**, 603-635 (1985).

25. G. B. Stanton, M. E. Goldberg, C. J. Bruce, Frontal eye field efferents in the macaque monkey: II.
Topography of terminal fields in midbrain and pons. *J Comp Neurol* **271**, 493-506 (1988).

26. D. P. Munoz, R. H. Wurtz, Fixation cells in monkey superior colliculus. I. Characteristics of cell
discharge. *J Neurophysiol* **70**, 559-575 (1993).

27. A. K. Moschovakis *et al.*, An anatomical substrate for the spatiotemporal transformation. *J
Neurosci* **18**, 10219-10229 (1998).

28. Z. M. Hafed, L. Goffart, R. J. Krauzlis, A neural mechanism for microsaccade generation in the
primate superior colliculus. *Science* **323**, 940-943 (2009).

29. D. L. Kimmel, T. Moore, Temporal patterning of saccadic eye movement signals. *J Neurosci* **27**,
7619-7630 (2007).

30. B. R. Cowley *et al.*, DataHigh: graphical user interface for visualizing and interacting with high-
dimensional neural activity. *Journal of neural engineering* **10**, 066012 (2013).

31. C. Massot, U. K. Jagadisan, N. J. Gandhi, Sensorimotor transformation elicits systematic patterns
of activity along the dorsoventral extent of the superior colliculus in the macaque monkey.
Commun Biol **2**, 287 (2019).

32. R. P. Heitz, J. D. Schall, Neural mechanisms of speed-accuracy tradeoff. *Neuron* **76**, 616-628
(2012).

33. N. J. Gandhi, E. L. Keller, Spatial distribution and discharge characteristics of superior colliculus
neurons antidromically activated from the omnipause region in monkey. *J Neurophysiol* **78**,
2221-2225 (1997).

34. K. Yoshida, Y. Iwamoto, S. Chimoto, H. Shimazu, Saccade-related inhibitory input to pontine
omnipause neurons: an intracellular study in alert cats. *J Neurophysiol* **82**, 1198-1208 (1999).

35. R. A. Deisz, G. Fortin, W. Zieglgansberger, Voltage dependence of excitatory postsynaptic
potentials of rat neocortical neurons. *J Neurophysiol* **65**, 371-382 (1991).

36. G. Stuart, B. Sakmann, Amplification of EPSPs by axosomatic sodium channels in neocortical
pyramidal neurons. *Neuron* **15**, 1065-1076 (1995).

37. E. L. Keller, Participation of medial pontine reticular formation in eye movement generation in
monkey. *J Neurophysiol* **37**, 316-332 (1974).

38. S. Everling, M. Paré, M. C. Dorris, D. P. Munoz, Comparison of the discharge characteristics of
brain stem omnipause neurons and superior colliculus fixation neurons in monkey: implications
for control of fixation and saccade behavior. *J Neurophysiol* **79**, 511-528 (1998).

39. C. K. Hauser, D. Zhu, T. R. Stanford, E. Salinas, Motor selection dynamics in FEF explain the
reaction time variance of saccades to single targets. *Elife* **7**, (2018).

40. U. K. Jagadisan, N. J. Gandhi, Disruption of Fixation Reveals Latent Sensorimotor Processes in the
Superior Colliculus. *J Neurosci* **36**, 6129-6140 (2016).

41. D. P. Hanes, R. H. Wurtz, Interaction of the frontal eye field and superior colliculus for saccade
generation. *J Neurophysiol* **85**, 804-815 (2001).

42. T. R. Peel, S. Dash, S. G. Lomber, B. D. Corneil, Frontal eye field inactivation alters the readout of
superior colliculus activity for saccade generation in a task-dependent manner. *J Comput
Neurosci*, (2020).

43.   B. D. Corneil, E. Olivier, D. P. Munoz, Visual responses on neck muscles reveal selective gating
1390   that prevents express saccades. *Neuron* **42**, 831-841 (2004).

1391   44.   C. Gu, D. K. Wood, P. L. Gribble, B. D. Corneil, A Trial-by-Trial Window into Sensorimotor
1392   Transformations in the Human Motor Periphery. *J Neurosci* **36**, 8273-8282 (2016).

1393   45.   A. P. Georgopoulos, A. B. Schwartz, R. E. Kettner, Neuronal population coding of movement
1394   direction. *Science* **233**, 1416-1419 (1986).

1395   46.   C. Lee, W. H. Rohrer, D. L. Sparks, Population coding of saccadic eye movements by neurons in
1396   the superior colliculus. *Nature* **332**, 357-360 (1988).

1397   47.   G. Rizzolatti, L. Riggio, I. Dascola, C. Umilta, Reorienting attention across the horizontal and
1398   vertical meridians: evidence in favor of a premotor theory of attention. *Neuropsychologia* **25**,
1399   31-40 (1987).

1400   48.   G. Vigneswaran, R. Philipp, R. N. Lemon, A. Kraskov, M1 corticospinal mirror neurons and their
1401   role in movement suppression during action observation. *Curr Biol* **23**, 236-243 (2013).

1402   49.   A. Y. Tan, Y. Chen, B. Scholl, E. Seidemann, N. J. Priebe, Sensory stimulation shifts visual cortex
1403   from synchronous to asynchronous states. *Nature* **509**, 226-229 (2014).

1404   50.   C. Hauptmann, P. A. Tass, Restoration of segregated, physiological neuronal connectivity by
1405   desynchronizing stimulation. *Journal of neural engineering* **7**, 056008 (2010).

1406   51.   O. V. Popovych, P. A. Tass, Control of abnormal synchronization in neurological disorders. *Front
1407   Neurol* **5**, 268 (2014).

1408   52.   W. Softky, Sub-millisecond coincidence detection in active dendritic trees. *Neuroscience* **58**, 13-
1409   41 (1994).

1410   53.   T. R. Stanford, E. G. Freedman, D. L. Sparks, Site and parameters of microstimulation: evidence
1411   for independent effects on the properties of saccades evoked from the primate superior
1412   colliculus. *J Neurophysiol* **76**, 3360-3381 (1996).

1413   54.   H. A. Katnani, N. J. Gandhi, The relative impact of microstimulation parameters on movement
1414   generation. *J Neurophysiol* **108**, 528-538 (2012).

1415   55.   C. L. Bryant, N. J. Gandhi, Real-time data acquisition and control system for the measurement of
1416   motor and neural data. *J Neurosci Methods* **142**, 193-200 (2005).

1417   56.   D. L. Sparks, R. Holland, B. L. Guthrie, Size and distribution of movement fields in the monkey
1418   superior colliculus. *Brain Res* **113**, 21-34 (1976).

1419   57.   V. Mante, D. Sussillo, K. V. Shenoy, W. T. Newsome, Context-dependent computation by
1420   recurrent dynamics in prefrontal cortex. *Nature* **503**, 78-84 (2013).

1421   58.   M. Rigotti *et al.*, The importance of mixed selectivity in complex cognitive tasks. *Nature* **497**,
1422   585-590 (2013).

1423   59.   F. P. Ottes, J. A. Van Gisbergen, J. J. Eggermont, Visuomotor fields of the superior colliculus: a
1424   quantitative model. *Vision Res* **26**, 857-873 (1986).

1425   60.   R. A. Fisher, The use of multiple measurements in taxonomic problems. *Ann Eugenic* **7**, 179-188
1426   (1936).

1427   61.   B. W. Matthews, Comparison of the predicted and observed secondary structure of T4 phage
1428   lysozyme. *Biochim Biophys Acta* **405**, 442-451 (1975).

1429   62.   S. Boughorbel, F. Jarray, M. El-Anbari, Optimal classifier for imbalanced data using Matthews
1430   Correlation Coefficient metric. *PLoS One* **12**, e0177678 (2017).

1431   63.   A. Rauch, G. La Camera, H. R. Luscher, W. Senn, S. Fusi, Neocortical pyramidal cells respond as
1432   integrate-and-fire neurons to in vivo-like input currents. *J Neurophysiol* **90**, 1598-1612 (2003).

1433   64.   M. Usher, J. L. McClelland, The time course of perceptual choice: the leaky, competing
1434   accumulator model. *Psychol Rev* **108**, 550-592 (2001).

1435   65.   B. A. Purcell *et al.*, Neurally constrained modeling of perceptual decision making. *Psychol Rev*
1436   **117**, 1113-1143 (2010).

1437   66.   K. F. Wong, A. C. Huk, M. N. Shadlen, X. J. Wang, Neural circuit dynamics underlying
1438   accumulation of time-varying evidence during perceptual decision making. *Front Comput*
1439   *Neurosci* **1**, 6 (2007).
1440   67.   T. Tsujimoto, A. Jeromin, N. Saitoh, J. C. Roder, T. Takahashi, Neuronal calcium sensor 1 and
1441   activity-dependent facilitation of P/Q-type calcium currents at presynaptic nerve terminals.
1442   *Science* **295**, 2276-2279 (2002).
1443   68.   J. G. Borst, B. Sakmann, Facilitation of presynaptic calcium currents in the rat brainstem. *The*
1444   *Journal of physiology* **513 (Pt 1)**, 149-155 (1998).
1445   69.   G. G. Turrigiano, K. R. Leslie, N. S. Desai, L. C. Rutherford, S. B. Nelson, Activity-dependent scaling
1446   of quantal amplitude in neocortical neurons. *Nature* **391**, 892-896 (1998).

1447