Geographical co-occurrence of butterfly species: The importance of niche filtering

by host plant species

Ryosuke Nakadai[1,2,*,†], Koya Hashimoto[1,*], Takaya Iwasaki[3], Yasuhiro Sato[1,4]

[1]Center for Ecological Research, Kyoto University, Hirano 2-509-3, Otsu, Shiga,

520-2113 Japan

[2]Current address: Faculty of Science, University of the Ryukyus, Senbaru 1, Nishihara,

Okinawa, 903-0213 Japan.

[3]Department of Biological Sciences, Faculty of Science, Kanagawa University,

Tsuchiya 2946, Hiratsuka, Kanagawa 259-1293 Japan

[4]Current address: Department of Plant Life Sciences, Faculty of Agriculture, Ryukoku

University, Yokotani 1-5, Otsu, Shiga 520-2194 Japan.

[*]Equal contribution

[†]*Author Correspondence:* R. Nakadai

Faculty of Science, University of the Ryukyus, Senbaru 1, Nishihara, Okinawa 903-0213,

Japan.

E-mail: r.nakadai66@gmail.co.jp

Author contributions: RN and KH conceived and designed the study; KH and RN

collected the data from the literature; TI conducted the analysis of ecological niche

modeling; YS and RN performed the statistical analyses; and RN, KH, YS, and TI

wrote the manuscript.

Conflict of Interest: The authors declare that they have no conflict of interest.

Abstract

The relevance of interspecific resource competition in the context of community assembly by herbivorous insects is a well-known topic in ecology. Most previous studies focused on local species assemblies, that shared host plants. Few studies evaluated species pairs within a single taxon when investigating the effects of host plant sharing at the regional scale. Herein, we explore the effect of plant sharing on the geographical co-occurrence patterns of 232 butterflies distributed across the Japanese archipelago; we use two spatial scales (10×10 km and 1×1 km grids) to this end. We considered that we might encounter one of two predictable patterns in terms of the relationship between co-occurrence and host sharing among butterflies. On the one hand, host sharing might promote distributional exclusivity attributable to interspecific resource competition. On the other hand, sharing of host plants may promote co-occurrence attributable to filtering by resource niche. At both grid scales, we found significant negative correlations between host use similarity and distributional exclusivity. Our results support the hypothesis that the butterfly co-occurrence pattern across the Japanese archipelago is better explained by filtering via resource niche rather than interspecific resource competition.

Keywords: Climatic niche, dispersal ability, herbivorous insect, Japanese archipelago, taxonomic relatedness

Introduction

Efforts to understand community assembly processes are of major importance in ecological research for obtaining past, current, and future biodiversity information. Dispersal limitations, environmental filtering via both abiotic and biotic niches, and interspecific interactions are thought to sequentially determine local community structures (reviewed by Cavender-Bares et al. 2009). Of the various relevant factors, the significance of interspecific interaction has often been assessed by examining how different species co-occur spatially even when they share similar resource niches (Diamond 1975; Gotelli and McCabe 2002). As different species with similar niches likely prefer similar environmental habitats, but may compete strongly, recent studies have often compared the significance of interspecific interactions (in terms of community assembly patterns) from the viewpoint of niche filtering, which indicates species sorting via performance and survival rates of each species against both abiotic (e.g., climate) and biotic (e.g., resource) properties from the species pool (Webb et al. 2002; Mayfield and Levine 2010).

In terrestrial ecosystems, many plant species are used by various herbivorous insects as both food resources and habitats; however, the importance of interspecific resource competition among herbivorous insects was once largely dismissed (reviewed by Kaplan and Denno 2007). Although some earlier studies described niche partitioning between co-occurring herbivores and considered that partitioning was attributable to interspecific competition (Ueckert and Hansen 1971; Benson 1978, Waloff 1979), other studies have observed frequent co-occurrence of multiple herbivorous insect species on shared host plants despite the fact that their niches extensively overlapped (Ross 1957; Rathcke 1976; Strong 1982; Bultman and Faeth 1985). Several authors have even argued that interspecific competition among herbivorous insects is too rare to structure herbivore communities (Lawton and Strong 1981; Strong et al. 1984). In subsequent decades however, many experimental ecological studies revealed that herbivorous

insects do compete with one another, often mediated by plastic changes in plant defense

traits following herbivory (Faeth 1986; Harrison and Karban 1986; Brown and Weis

1995; Agrawal 1999; Agrawal 2000; Viswanathan et al. 2005). These findings have

prompted insect ecologists to revisit exploring the prevalence of interspecific resource

competition in structuring herbivore communities (Kaplan and Denno 2007). Also,

evolutionary studies have pointed out that interspecific resource competition may have

regulated speciation and species diversification of herbivorous insects (Ehrlich and

Raven 1964; Rabosky 2009; Nosil 2012; Thompson 2013). For example, rapid

diversification after host shift to distant relative plants (Fordyce 2010) was recognized

as indirect evidence of diversity regulation by interspecific competition (Ehrlich and

Raven 1964; Rabosky 2009; Nakadai 2017), because the release from diversity

regulation by host shift (i.e., a key innovation and ecological release; Yoder et al. 2010)

may cause following specialization and diversification of herbivorous insects. In this

context, the co-occurrence pattern, especially on a large spatial scale and not within the

local community, is an important indicator for revealing the effects of interspecific

resource competition in herbivore diversification, which eventually feeds back to their

species pool as a bottom-up process (Loreau 2000). However, very little is known about

the extent to which the results of the cited studies can be extrapolated to describe

patterns, at the regional scale, of the distribution of herbivorous insects within a single

taxon.

The effects of interspecific competition and filtering via resource niches on the

co-occurrence patterns of herbivorous insects may be obscured by other potential

factors. For example, climatic niches (often associated with differences in potential

geographical distributions in the absence of interspecific interactions; Warren et al.

2008; Takami and Osawa 2016) may drive niche filtering, which may in turn mediate

the impact of host use on assembly patterns, because the distributions of host plants are

also strongly affected by climate niches (Hawkins et al. 2014; Kubota et al. 2017). In

addition, taxonomic relatedness should reflect niche similarity; the niche of any

organism should be partly determined by its phylogenetic history (Webb et al. 2002).

For example, phylogenetic conservatisms of host use (i.e., resource niche) in

herbivorous insects were often reported (Lopez-Vaamonde et al. 2003; Nyman et al.

2010; Jousselin et al. 2013; Doorenweerd et al. 2015). Thus, the outcomes of

competition within shared niches should reflect both resource and climatic factors

(Mayfield and Levine 2010: Connor et al. 2013). As such factors may correlate with

host use by individual species; any focus on host use alone may yield misleading results.

Furthermore, dispersal ability may complicate assembly patterns, often being associated

with both the extent of the geographical range and other factors (Kneitel and Chase

2004). A high dispersal ability may enhance the extent of co-occurrence. Thus, when

attempting to explore the effects of competition and filtering via resource niches on

distribution patterns, it is important to consider all of taxonomic relatedness, climatic

niche preferences, and dispersal ability.

In the present study, we explored whether interspecific competition or niche

filtering better explained the geographical co-occurrence of a group of herbivorous

butterflies. Given that the host specificity of butterflies is relatively high among

herbivorous insects (Novotny et al. 2010), butterflies that share a host plant are more

likely to be potential competitors for each other than other less specialized herbivores

are. Butterfly larvae often consume huge amounts of leaves in the larval stage, and even

eat whole material of plants in some cases (Inouye and Johnson 2005; Fei et al. 2016;

Hashimoto and Ohgushi 2017). Moreover, it is suggested that not all plants or parts of a

plant are suitable for butterfly larvae due to within- and among-individual variations in

plant quality (e.g., nutritional status and defensive traits such as secondary metabolites)

(Damman 1989; Dempster 1997; Van Zandt and Agrawal 2004). Hence, the observation

that most parts of terrestrial plants seem to remain unconsumed (Hairston et al. 1960;

Polis 1999) does not necessary mean that butterflies are not facing resource competition.

Several studies have examined butterfly–butterfly competition (Brower 1962; Millan et

al. 2013), and many have reported interspecific resource competition among

herbivorous insects that involved butterfly species (Agrawal 1999; Redman and Scriber

2000; Agrawal 2000; Van Zandt and Agrawal 2004; Prior and Hellman 2010). Thus,

resource-mediated competition is expected to be prevalent in herbivore butterflies. In

particular, we focused on herbivorous butterflies of the Japanese archipelago for the

following reasons. First, the Biodiversity Center of Japan has extensive records of

butterfly distribution. Second, forewing length (easily measured on photographs) is a

useful index of butterfly dispersal ability (Chai and Srygley 1990; Shirôzu 2006).

Finally, and most importantly, a great deal is known about the hosts used by butterfly

species (Saito et al. 2016). Thus, data on Japanese butterflies can be used to explore the

effects of host plant sharing and other factors on the co-occurrence patterns at regional

scales.

We expected to discern one of two patterns when evaluating the significance

of interspecific competition in terms of the geographical co-occurrence of herbivore

butterfly. If interspecific resource competition is sufficiently intense to cause

competitive exclusion with the precondition that intraspecific competition is

comparably low, sharing of host plants would be positively associated with exclusive

distribution (Hypothesis 1 in Table 1). Alternatively, if filtering via a resource niche is

relatively stronger than resource competition, sharing of host plants would promote

species co-occurrence (Hypothesis 2 in Table 1). To distinguish between these two

patterns, we examined the correlations between host use similarity (i.e., the extent of

sharing of host species) and the exclusiveness of geographical distribution among pairs

of entirely herbivorous Japanese butterflies. The former pattern predicts that a positive

correlation would be evident between host use similarity and the exclusiveness of

geographic distribution. The latter pattern predicts that the correlation would be

negative. Furthermore, we also considered taxonomic relatedness, climatic niche

similarities, and overall dispersal abilities in the course of our work as factors affecting

the association between host sharing and geographical co-occurrence (major hypotheses

are summarized in Table 1).

Materials and methods

Study area

The Japanese archipelago, including the Ryukyu Islands, forms a long chain of

continental islands lying off the eastern coast of Asia. The Japanese archipelago was

recognized as a hotspot of biodiversity (Gerardo and Brown 1995; Mittermeier et al.

2011) and insect diversity, with about 32,000 insect species recorded (Clausnitzer et al.

2009; Tojo et al. 2017). The latitudinal range of the archipelago (22°N to 45°N)

embraces hemi-boreal, temperate, and subtropical zones. The mean temperatures in the

coldest and warmest months are −19.0°C and 31.5°C, respectively; the annual

precipitation ranges from 867 to 3,908 mm (Kubota et al. 2014).

Study organisms

The Japanese archipelago hosts over 280 species of butterflies of five families (the

Papilionidae, Pieridae, Lycaenidae, Nymphalidae, and Hesperiidae) (Shirôzu 2006).

Over 95% of the larvae of Japanese butterflies feed on plants (Honda 2005). The host

plants are diverse and include both dicots and monocots (Saito et al. 2016). Lycaenid

butterflies include two non-herbivorous species, but all species of all other families are

exclusively herbivorous.

Metadata compilation

Butterfly census data are available on the website of the Biodiversity Center of Japan,

Ministry of the Environment (http://www.biodic.go.jp/index_e.html). We used the

results of the fourth and fifth censuses (1988 to 1991 and 1997 to 1998, respectively) of

the National Survey of the Natural Environment in Japan

(http://www.biodic.go.jp/kiso/15/do_kiso4.html). This database includes records of 273

species/subspecies of butterflies from the entire Japanese archipelago, in grid cells of

latitude 5 min and longitude 7.5 min (the Japanese Standard Second Mesh). These grid

dimensions are about 10 km × 10 km, and this grid is described below as the "10-km

grid." Furthermore, the Biodiversity Center also contains records from grid cells of

latitude 30 s and longitude 45 s (the Japanese Standard Third Mesh). These grid

dimensions are about 1 km × 1 km, and this grid is described below as the "1-km grid."

As processes driving community assemblies may vary between spatial scales

(Cavender-Bares et al. 2009), we evaluated data from both grids. We summarized data

at the species level, and converted all records into the presence or absence (1/0) of a

species in each grid. Finally, the number of records used in the study after summarizing

the duplicate presence records was 79,003 and 165,102 for each 10-km grid and 1-km

grid for targeted species, respectively. We used the taxonomy of Shirôzu (2006).

Data on host plants and forewing length were evaluated as possible variables

explaining, respectively, host use and dispersal ability. The host plants of 278 butterfly

species/subspecies (3,573 records in total, after excluding duplications) were obtained

from data by Saito *et al.* (2016), and the host records were described in 8 major

illustrated reference books, 2 checklists, and 14 other pieces of literature; the presence

of larvae on plants, and the observations of larvae eating plants or insects in the field

were considered host records. We used 2,939 records of targeted species in this study

(Table S1). The datasets of Saito *et al.* (2016) included some incomplete records (e.g.,

Asarum sp. of *Luehdorfia japonica*, *Cotoneaster* spp. of *Acytolepis puspa*) including 34

records (about 1.14% of whole records) on targeted species; those incomplete records

were excluded from our analyses. Dispersal ability was evaluated by reference to adult

wing length. We compiled wing data on 284 species using published illustrations

(Shirôzu 2006). We used Image J software (Abramoff et al. 2004) to extract forewing

lengths (in cm) from plates that included centimeter scale bars. Multiple forewing

lengths were extracted when individual, sexual, and/or geographical variations were

evident.

We assembled data on the distributions, host plants, climatic niches (described

below), and forewing lengths of 232 butterfly species. Twenty-four species were

excluded from the analysis, for the following reasons. First, the taxonomic status of

three species (*Papilio dehaanii*, *Pieris dulcinea*, and *Eurema hecabe*) changed in the

interval between the fourth and fifth biodiversity censuses (Inomata 1990; Shirôzu

2006). Thus, we excluded these species because identifications were unreliable. Second,

we excluded three species of non-herbivorous butterflies (*Taraka hamada*, *Spindasis*

takanonis, and *Niphanda fusca*). Finally, we excluded a further 18 species because the

models used to evaluate their ecological niches failed to satisfy the criteria that we

imposed (i.e., shortage of occurrence records for the niche modeling program or less

than 0.7 of the values of area under the curve [AUC], the details are in Supplementary

File 1).

Data analysis

Species distribution exclusiveness. We used the checkerboard scores (C-scores; Stone

and Roberts 1990) to evaluate the exclusivity of distributions between each species pair.

We set r_i and r_j as the numbers of grids in which species *i* and *j*, respectively, were

present; the checker unit C_{ij} associated with the two species was defined as: $C_{ij} = (r_i -$

$S_{ij}) \times (r_j - S_{ij})$, where S_{ij} indicates the extent of co-presence (i.e., the number of grid

cells shared by the two species). Thus, the checker unit became larger as the two species

occurred more commonly in different grid cells. We simulated null models to allow the

observed checker units to be compared with stochastic distributions. We used the

method of Jonsson et al. (2001) to maintain the observed frequencies of species

occurrence and randomized the presence/absence matrices for each pair of butterfly

species. The null models were run 999 times for each species pair. C_{obs} and C_{null} were

the checker units of the observed and null distributions, respectively; the checker unit

was standardized as: $C_{std} = (C_{obs} - C_{null})/SD_{null}$, where SD_{null} indicates the standard

deviation of all checker units of the null models. The checker unit of the null model,

C_{null}, was the average checker unit of all null models. Thus, positive and negative values

of C_{std} indicate that two species are allopatrically and sympatrically distributed,

respectively, to extents greater than indicated by the null models. We also attempted to

construct a more constrained null model by weighting each grid by its species richness

to consider the potential size of the butterfly community as well as the observed

frequency of each species. However, this weighted index was very highly correlated

with C_{std} ($r = 0.97$, $n = 300$ randomly selected pairs); thus, we adopted Jonsson's

method in this study. All statistical analyses were performed with the aid of R software

version 3.2.0 (R Core Team 2015).

Climatic niche similarities. We used ecological niche modeling (ENM)

(Franklin 2010) to evaluate climatic niche similarities among butterfly species (Warren

et al. 2008). ENM associates distributional data with environmental characteristics, thus

estimating the response functions of, and the contributions made by, environmental

variables. Furthermore, potential distributional ranges may be obtained by projecting

model data onto geographical space. In the present study, potential distributional ranges

estimated by ENM should be influenced by abiotic environmental variables alone

(climate and altitude); we did not consider interspecific interactions among butterflies or

dispersal abilities in this context. Thus, comparisons of potential distribution patterns

estimated by ENM allow evaluation of climatic niche similarities among butterfly

species. The maximum entropy algorithm implemented in MaxEnt ver. 3.3.3e software

(Phillips et al. 2006; Phillips and Dudík 2008) was employed in ENM analyses

(Supplementary File 1 contains the details). The logistic outputs of MaxEnt analyses

can be regarded as presence probabilities (Phillips and Dudík 2008). Finally, we then

used Schoener's (1968) D statistic to calculate climatic niche similarities between pairs

of butterfly species, based on the MaxEnt outputs. $P_{x,i}$ and $P_{y,i}$ were the probabilities

(assigned by MaxEnt) that species x and y would mesh to the extent of i on a geographic

scale; the climatic niche similarity between the two species was defined as: $D_{env} = 1 -$

$0.5 \times \Sigma|P_{x,i} - P_{y,i}|$, where D_{env} ranged from 0 (no niche overlap) to 1 (completely

identical niches). The probability assigned to the presence of species x in grid i was $P_{x,i}$

$= p_{x,i} / \Sigma p_{x,i}$, where $p_{x,i}$ was the logistic Maxent output for species x in grid i.

Explanatory variables. We evaluated both host use similarity and other factors

that might explain exclusive species distributions (i.e., taxonomic relatedness). We

calculated the total dispersal abilities of species pairs and climatic niche similarities (as

explained above). Host use similarity was calculated as 1 minus Jaccard's dissimilarity

index (Koleff et al. 2003) when host plant species were shared by two butterflies. The

taxonomic relatedness of each species pair was classified as: 2: in the same genus; 1: in

the same family; and 0: in different families. The total dispersal ability was calculated

as the sum of the ln (forewing lengths) of each species pair.

Statistical tests. We used the Mantel test, in which the response matrix yielded

pairwise C_{std} data, and which included explanatory matrices, to examine the effects of

host use similarity and other factors (i.e., taxonomic relatedness, climatic niche

similarity, and total dispersal ability) on the exclusivity of butterfly distribution. We

calculated Spearman's correlations because one of the explanatory variables (taxonomic

relatedness) was a rank variable. P-values were determined by running 9,999

permutations. In addition, when analyzing correlations between host use similarity and

other explanatory matrices, we ran Mantel tests with 9,999 permutations, and calculated

Spearman's correlations. We used partial Mantel tests, in which the response matrix

yielded pairwise C_{std} data, and in which a matrix of host use similarity including the

other three explanatory matrices served as a co-variable, to evaluate the effects of

confounding factors associated with host use similarity on the exclusivity of butterfly

distribution. We ran 9,999 permutations and calculated Spearman's correlations. We

employed the vegdist function of the vegan package (Oksanen et al. 2015) and the

mantel function of the ecodist package (Goslee and Urban 2007) implemented in R. We

performed all analyses using data from both the 10-km and the 1-km grids. The same

analysis was conducted for the datasets after excluding the species (the bottom 10% in

the 10-km grid and 1-km grid scales, to reduce the bias of rare species in the analysis.

The histogram of presence records for each butterfly species is shown in Figure S1.

Results

The standardized checkerboard scores (C_{std} values) of most species pairs were negative,

indicating that, in general, Japanese butterflies were more likely to co-occur than

expected by chance (Fig. 1). The Mantel test showed that host use similarity was

significantly and negatively correlated with the C_{std} values at both geographical scales

(Fig. 1, Table 2a); all three explanatory variables exhibited significant negative

correlations with the C_{std} values at both scales (Table 2a). Host use similarity was

significantly and positively correlated with both taxonomic relatedness and climatic

niche similarity, but we found no significant correlation with total dispersal ability

(Table 2b). The partial Mantel tests showed negative correlations between the C_{std}

values and host use similarity at both geographical scales when controlling both

taxonomic relatedness and dispersal ability. The differences of the correlation

coefficients from the results of Mantel tests were small, specifically less than 1% (Table

2c). In contrast, we found significant positive correlations when controlling for climatic

niche similarity and all three factors in the 10-km grid dataset, in which the correlation

coefficients between host use similarity and C_{std} score largely changed in the positive

direction from the results of Mantel tests, but no significant correlation in the 1-km grid

dataset (Table 2c). When we excluded rare species from the analyses, most of the

results were consistent with the results of original datasets, except for the correlation

between the C_{std} values and host use similarity under the control of climatic niche

similarity in the 10-km grid dataset, which was not significant (Tables S5 and 6).

Discussion

Significant negative correlations were clearly evident between the C_{std} scores and host

use similarities at both grid scales (Fig. 1, Table 2a), indicating that a pair of Japanese

butterflies sharing host plants is more likely to co-occur (supporting Hypothesis 2).

Significant negative correlations between C_{std} scores and host use similarities were

evident after controlling for each of the other potentially confounding factors,

taxonomic relatedness, and total dispersal ability (Table 2c). The predicted pattern from

interspecific resource competition (i.e., positive correlations between the C_{std} score and

host use similarity) were confirmed only after controlling for the effects of climatic

niche similarity and all three factors in the 10-km grid data (supporting Hypothesis 6).

However, the correlation coefficient was low (Table 2c), and we could not confirm the

pattern in the analysis excluding rare species (Table S5c). Note that we need to interpret

the results carefully, as interspecific resource competition is not the only process that

generates negative correlations between C_{std} score and host use similarity (e.g.,

speciation, Warren et al. 2014). Our results are consistent with the idea that interspecific

resource competition is too weak to organize communities of herbivorous insects

effectively (Lawton and Strong 1981; Strong et al. 1984; Nakadai and Kawakita 2017).

Rather, our results suggest that the geographic pattern of species co-occurrence among

Japanese butterflies is better explained by niche filtering.

The most likely explanation of our data is that the relative strength of

structuring via resource competition may be weaker than that associated with niche

filtering. As the geographical distributions of host plants would be expected to be

strongly associated with the local climatic environment, the impacts of resource and

climatic niche filtering may combine to ensure that butterfly species sharing host plants

assemble in the same places. In addition, the dispersal of adult butterflies from the

patches in which they were born may counteract the structuring force imposed by

interspecific competition. However, although co-occurrence was facilitated by the

overall total dispersal ability (Table 2a), the negative correlations between the C_{std}

scores and host use similarity were evident even when we controlled for the effects of

total dispersal ability (Table 2c). This means that dispersal alone may not explain the

weak impact of resource competition on the co-occurrence patterns of Japanese

butterflies. Other potential factors reducing the effects of interspecific resource

competition may also be in play, although we did not address these topics in this study.

For example, co-occurring butterfly species, which share their host plants, are also more

likely to share their evolutionary histories. The accumulated time in which species

co-occur (i.e., historical co-occurrence) may cause character displacement and reduce

the effects of interspecific resource competition on the co-occurrence pattern after

convergence (e.g., Germain et al. 2016). In the case of Japanese butterflies, there are

differences in historical dispersal events, especially the timing and route to the Japanese

archipelago, due to the repeated connections and disconnections between Eurasia

continent and the Japanese archipelago (reviewed by Tojo et al. 2017). In addition,

butterflies sharing host plants may experience similar historical dispersal events

corresponding to the distribution changes of plant species. Such a historical process can

generate faunal similarity along distance and climate (Nekola and White 1999; Kubota

et al. 2017) and indirectly affect the pattern of herbivorous insects, which was also

detected in our study (see the correlation between C_{std} and climate niche similarity in

Table 2a). Therefore, the effects of historical co-occurrence on the patterns of current

geographical co-occurrence may also occur, although we did not discuss the details

here.

The negative correlation evident between taxonomic relatedness and the C_{std}

scores (Table 2a) suggests that niche filtering is in play among Japanese butterflies,

given that taxonomic relatedness serves as a proxy of niche similarity including host use.

Indeed, we found significant (positive) correlations between host use similarity and the

taxonomic relatedness of Japanese butterflies (Table 2b), as has often been shown for

other herbivorous insects (e.g., Nyman et al. 2010; Jousselin et al. 2013). These results

reconfirm niche conservatisms associated with host use, which were also reported in

previous studies (Lopez-Vaamonde et al. 2003; Nyman et al. 2010; Jousselin et al.

2013; Doorenweerd et al. 2015; Nakadai and Kawakita 2016). Moreover, when host use

similarity was controlled using the partial Mantel test, taxonomic relatedness did not

significantly affect co-occurrence at the 10-km grid scale (Table S4). These results

suggest that, at least at the 10-km grid scale, the effects of taxonomic relatedness largely

reflect host use similarity. Although we mainly discussed taxonomic relatedness as a

niche similarity between species pairs throughout the manuscript, the correlation

between geographical co-occurrence and taxonomic relatedness could also be

interpreted as the imprint of allopatric speciation (Hypothesis 3-2, Barraclough et al.

1998; Warren et al. 2014). In that context, we can also interpret our results as the

imprint of allopatric speciation not being strong enough to construct the co-occurrence

pattern at the regional scale.

In the present study, we used ENM to evaluate the effects of climatic niche

similarity on co-occurrence patterns. When we controlled for the effects of such niche

similarity, the negative correlations between the C_{std} scores and host plant similarities

disappeared at both spatial scales (Table 2c). This suggests that the explanatory power

of climatic niche filtering is stronger than that of resource niche filtering. It is

reasonable that host plants are also distributed in relation to climatic niches (Hawkins et

al. 2014). Moreover, we also only found a positive correlation when controlling for the

effects of climatic niche similarity in the 10-km grid data, which supports the

hypothesis of interspecific resource competition (Hypothesis 6). This result suggests the

possibility that the competitive exclusions of herbivorous insects do exist but usually

cannot be observed due to masking by climatic niche filtering and other effects (Table

2c). It should be noted that, although ENM has been widely used to quantify climatic

niches (e.g., Kozak et al. 2008; Warren et al. 2008; Takami and Osawa 2016) when we

interpret this result, ENM data should be treated with caution (Peterson et al. 2011;

Warren 2012; Warren et al. 2014). For example, Warren et al. (2014) noted that ENM

always includes non-targeted factors that limit real distributions if those distributions

correlate spatially with environmental predictors. Such confounding effects may cause

overestimation of any positive correlation between climatic niche similarity and

butterfly co-occurrence, and may also cause the effects of host use similarity to be

underestimated when controlling for the effects of climatic niche similarity. The use of

ENM to study species co-occurrence patterns requires careful consideration; more case

studies evaluating the relative importance of climatic niches are required.

Although a number of studies have reported that resource competition

involving butterfly species occur (e.g., Redman and Scriber 2000; Prior and Hellman

2010), few studies have indicated competitive exclusion or resource partitioning due to

resource competition involving butterflies (Blakley and Dingle 1978; Millan et al. 2013).

Some authors have argued that reproductive interference among butterflies sharing

similar niches is a more plausible mechanism causing resource or habitat niche

displacement than resource competition (Jones et al. 1998; Friberg et al. 2008, 2013:

reviewed in Noriyuki et al. 2015; Shuker and Burdfield-Steel 2017). Friberg *et al.*

(2013) suggested that habitat separation of two congeners of wood white butterflies

Leptidea sinapis and *L. juvernica* is caused by reproductive interference. Because

habitat displacement by reproductive interference is more likely to work at the local

scale (Friberg et al. 2013), its effects may be difficult to observe at the regional or

broader spatial scale. This does not contradict our results in which robust evidence of

negative co-occurrence among host-sharing species was not detected.

We focused on butterfly species, which is one of the best-studied herbivore

taxa in both ecology and evolution (Ehrlich and Raven 1964; Hanski and Singer 2001;

Fordyce et al. 2010), as well as data on fundamental information (e.g., occurrence

information and host use of each species) that are abundant in butterfly species. Our

analysis relied on this advantage, enabling the use of huge datasets of butterfly species,

and attempted to clarify the geographical co-occurrence on a larger scale. However, our

study had several limitations. First, there was underrepresentation of rare species. The

abundance and the range of distribution are different among butterfly species, and the

records of rare species had some bias in our analysis using huge datasets. We

additionally conducted analyses after excluding the species, which were confirmed in

the lower number of grids, to eliminate the bias due to the unfounded presence records

of rare species. As a result, most of the results showed the same trends as the results of

the original datasets, so the bias of rare species may not be very large (Tables S5 and 6).

The second issue was regarding the temporal fidelity of observations. We compiled

datasets of field surveys over multiple years, but using single-year datasets, which are

more appropriate for assessing the effects of interspecific resource competition on the

patterns of geographical co-occurrence. If butterfly distributions largely change over the

years, this approach would obscure the real patterns of resource-driven butterfly

co-occurrence. Finally, we only used occurrence data with the accuracy of a large grid

(10-km and 1-km squares); combining datasets of deferent resolution (grid-level and

community-level data) may help to resolve co-occurrence patterns among herbivore

butterflies (Cardillo and Warren 2016).

Conclusions

The significance of interspecific resource competition in terms of the structuring of

herbivorous insect communities is a source of long-standing controversy. Many

researchers have sought to explain the general patterns of relationships between the

co-occurrence of, and the use of different niches by, herbivorous insects. However, the

data remain limited because most previous studies employed narrow taxonomic and

spatial scales. In this context, our study is the first to provide a comprehensive picture of

the co-occurrence patterns among a single taxonomic group over a large region.

Although co-occurrence of Japanese butterflies is more likely to be driven by niche

filtering than interspecific resource competition, it is essential to employ broad

taxonomic and spatial scales when attempting to reveal general patterns of community

assembly among herbivorous insects. Future studies should explore the relative

importance of each assembly stage not only ecologically but also over evolutionary time

(Rabosky 2009). Such work would answer the important question: "Why have

herbivorous insects become one of the most diverse groups of the natural world?"

Acknowledgements

We thank K. Kadowaki for his comments and advice on the original version of our

manuscript, M. U. Saito for giving us the detail information of collecting datasets of

host plants, the Biodiversity Center of Japan for allowing access to butterfly data at the

1-km grid scale, and the associate editor and three reviewers for their comments, which

improved our manuscript. We are particularly grateful to the entomologists and

naturalists who accumulated the information on butterflies used in this study. This work

was supported by a grant from the Grant-in-Aid Program for JSPS Fellows (Grant No.

15J00601).

References

Agrawal AA (1999) Induced responses to herbivory in wild radish: effects on several

herbivores and plant fitness. Ecology 80:1713–1723

Agrawal AA (2000) Specificity of induced resistance in wild radish: causes and

consequences for two specialist and two generalist caterpillars. Oikos 89:493–500

Abramoff MD, Magalhães PJ, Ram SJ (2004) Image processing with ImageJ. Biophot

inter 11:36–42

Barraclough TG, Vogler AP, Harvey PH (1998) Revealing the factors that promote

speciation. Philos Trans R Soc Lond B Biol Sci 353:241–249

Benson WW (1978) Resource Partitioning in Passion Vine Butterflies. Evolution

32:493–518

Blakley NR, Dingle H (1978) Competition: butterflies eliminate milkweed bugs from a

Caribbean Island. Oecologia 37:133–136

Brower LP (1962) Evidence for interspecific competition in natural populations of the

Monarch and Queen butterflies, *Danaus plexippus* and *D. gilippus berenice* in

south central Florida. Ecology 43:549–552

Brown DG, Weis AE (1995) Direct and indirect effects of prior grazing of goldenrod

upon the performance of a leaf beetle. Ecology 76:426–436

Bultman TL Faeth SH (1985) Patterns of intra- and inter- specific association in

leaf-mining insects on three oak host species. Ecol Entomol 10:121–129

Cantor SB, Sun CC, Tortolero-Luna G, Richards-Kortum R, Follen M (1999) A

Comparison of C/B Ratios from studies using receiver operating characteristic

curve analysis. J Clin Epidemiol 52:885–892

Cardillo M, Warren DL (2016) Analysing patterns of spatial and niche overlap among

species at multiple resolutions. Global Ecol Biogeogr 25: 951-963

Cavender-Bares J, Kozak KH, Fine PVA, Kembel SW (2009) The merging of

community ecology and phylogenetic biology. Ecol Lett 12:693–715

Chai P, Srygley RB (1990) Predation and the flight, morphology, and temperature of

neotropical rainforest butterflies. Am Nat 135:748–765

Clausnitzer V, Kalkman VJ, Ram M, Collen B, Baillie JEM, Bedjanič M, Darwell WRT,

Dijkstra K-DB, Dow R, Hawking J, Karube H, Malikova E, Paulson D, Schütte K,

Suhling F, Villanueva RJ, Ellenrieder N, Wilson K (2009) Odonata enter the

biodiversity crisis debate: the first global assessment of an insect group. Biol

Conserv 142:1864–1869

Connor EF, Collins MD, Simberloff D (2013) The checkered history of checkerboard

distributions. Ecology 94:2403–2414.

Damman H (1989) Facilitative interactions between two lepidopteran herbivores of

Asimina. Oecologia 78:214–219

Dempster JP (1997) The role of larval food resources and adult movement in the

population dynamics of the orange-tip butterfly (*Anthocharis cardamines*).

Oecologia 111:549–556

Diamond J (1975) Assembly of species communities. In: Cody ML, Diamond JM (eds)

Ecology and evolution of communities. Harvard University Press, Cambridge, pp

342–444

Doorenweerd C, van Nieukerken EJ, Menken SBJ (2015) A global phylogeny of

leafmining *Ectoedemia* moths (Lepidoptera: Nepticulidae): exploring host plant

family shifts and allopatry as drivers of speciation. PLoS One 10:e0119586

Ehrlich PR, Raven PH (1964) Butterflies and plants: a study in coevolution. Evolution

18:586–608

Fordyce JA (2010) Host shifts and evolutionary radiations of butterflies. Proc R Soc

Lond B Biol Sci 277:3735–3743

Faeth SH (1986) Indirect interactions between temporally separated herbivores

mediated by the host plant. Ecology 67:479–494

Fei M, Gols R, Zhu F, Harvey JA (2016) Plant quantity affects development and

survival of a gregarious insect herbivore and its endoparasitoid wasp. PLoS One

11:e0149539

Franklin J (2010) Mapping species distributions: spatial inference and prediction.

Cambridge University Press.

Friberg M, Bergman M, Kullberg J, Wahlberg (2008) Niche separation in space and

time between two sympatric sister species—a case of ecological pleiotropy. Evol

Ecol 22:1–18

Friberg M, Leimar O, Wiklund C (2013) Heterospecific courtship, minority effects and

niche separation between cryptic butterfly species. J Evol Biol 26:971–979

Gerardo C, Brown JH (1995) Global patterns of mammalian diversity, endemism, and

endangerment. Conservation Biology 9:559–568

Germain RM, Weir JT, Gilbert B (2016) Species coexistence: macroevolutionary

relationships and the contingency of historical interactions. Proc R Soc B

283:20160047

Goslee SC, Urban DL (2007) The ecodist package for dissimilarity-based analysis of

ecological data. Journal of Statistical Software 22:1–19

Gotelli NJ, McCabe DJ (2002) Species co-occurrence: A meta-analysis of J. M.

Diamond's assembly rules model. Ecology 83:2091–2096

Hairston NG, Smith FE, Slobodkin LB (1960) Community structure, population control,

and competition. Am Nat 94:421–425

Harrison S, Karban R (1986) Effects of an early-season folivorous moth on the success

of a later-season species, mediated by a change in the quality of the shared host,

Lupinus arboreus Sims. Oecologia 69:354–359

Hashimoto K, Ohgushi T (2017) How do two specialist butterflies determine growth

and biomass of a shared host plant? Popul Ecol 59:17–27

Hawkins BA, Rueda M, Rangel TF, Field R,   Diniz-Filho JAF (2014) Community

phylogenetics at the biogeographical scale: cold tolerance, niche conservatism and

the structure of North American forests. J Biogeogr 41:23–38

Honda K (2005) Larval feeding habit and host selection. In: Honda K, Kato Y (eds)

Biology of butterflies. University of Tokyo Press, Tokyo, pp 255–301 (in

Japanese).

Inomata T (1990) Keys to the Japanese butterflies in natural color. Hokuryukan.

Inouye BD, Johnson DM (2005) Larval aggregation affects feeding rate in *Chlosyne*

poecile (Lepidoptera: Nymphalidae). Florida Entomol 88:247–252

Jones MJ, Lace LA, Harrison EC, Stevens-Wood B (1998) Territorial behavior in the

speckled wood butteries *Pararge xiphia* and *P. aegeria* of Madeira: a mechanism

for interspecific competition. Ecography 21: 297–305

Jonsson BG (2001) A null model for randomization tests of nestedness in species

assemblages. Oecologia 127:309–313

Jousselin E, Cruaud A, Genson G, Chevenet F, Foottit RG, Cœur d'acier A (2013) Is

ecological speciation a major trend in aphids? Insights from a molecular phylogeny

of the conifer-feeding genus *Cinara*. Front Zool 10:56

Kaplan I. Denno RF (2007) Interspecific interactions in phytophagous insects revisited:

a quantitative assessment of competition theory. Ecol Lett 10:977–994

Kneitel JM, Chase JM (2004) Trade-offs in community ecology: Linking spatial scales

and species coexistence. Ecol Lett 7:69–80

Koleff P, Gaston KJ, Lennon JJ (2003) Measuring beta diversity for presence-absence

data. J Ani Ecol 72:367–382

Kozak KH, Graham CH, Wiens JJ (2008) Integrating GIS-based environmental data

into evolutionary biology. Trends Ecol Evol 23:141–148

Kubota Y, Hirao T, Fujii S, Shiono T, Kusumoto B (2014) Beta diversity of woody

plants in the Japanese archipelago: the roles of geohistorical and ecological

processes. J Biogeogr 41:1267–1276

Kubota Y, Kusumoto B, Shiono T, Tanaka T (2017). Phylogenetic properties of Tertiary

relict flora in the East Asian continental islands: imprint of climatic niche

conservatism and in situ diversification. Ecography, 40:436–447

Lawton JH, Strong DR (1981) Community patterns and competition in folivorous

insects. Am Nat 118:317–338

Liu C, Berry PM, Dawson TP, Pearson RG (2005) Selecting thresholds of occurrence in

the prediction of species distributions. Ecography 28:385–393

Lopez-Vaamonde, C, Godfray HCJ, Cook JM (2003) Evolutionary dynamics of

host-plant use in a genus of leaf- mining moths. Evolution 57:1804–1821

Loreau M (2000). Are communities saturated? On the relationship between α, β and γ

diversity. Ecol Lett 3:73-76

Mayfield MM, Levine JM (2010) Opposing effects of competitive exclusion on the

phylogenetic structure of communities. Ecol lett 13:1085–1093

Millan C, Borges SS, Rodrigues D, Moreira GRP (2013) Behavioral and life-history

evidence for interspecific competition in the larvae of two heliconian butterflies.

Naturwissenschaften 100:901–911

Mittermeier RA, Turner WR, Larsen FW, Brooks TM, Gascon C (2011) Global

biodiversity conservation: the critical role of hotspots. In: Zachos FE, Habel JC

(eds) Biodiversity hotspots. Springer, Berlin, Heidelberg, pp 3–22

Nakadai R (2017) Species diversity of herbivorous insects: a brief review to bridge the

gap between theories focusing on the generation and maintenance of diversity.

Ecol Res

Nakadai R, Kawakita A (2016) Phylogenetic test of speciation by host shift in leaf cone

moths (*Caloptilia*) feeding on maples (*Acer*). Ecol Evol 6:4958–4970

Nakadai R, Kawakita A (2017) Patterns of temporal and enemy niche use by a

community of leaf cone moths (*Caloptilia*) coexisting on maples (*Acer*) as revealed

by metabarcoding. Mol Ecol 26:3309–3319

Nekola JC, White PS (1999) The distance decay of similarity in biogeography and

ecology. J Biogeogr 26:867-878

Noriyuki S (2015) Host selection in insects: reproductive interference shapes behavior

of ovipositing females. Popul Ecol 57:293-305

Nosil P (2012) Ecological speciation. Oxford University Press, New York

Novotny V, Miller SE, Baje L, Balagawi S, Basset Y, Cizek L, Craft KJ, Dem F, Drew

RAI, Hulcr J, Leps J, Lewis OT, Pokon R, Stewart AJA, Samuelson GA, Weiblen

GD (2010) Guild-specific patterns of species richness and host specialization in

plant-herbivore food webs from a tropical forest. J Ani Ecol 79:1193–1203

Nyman T, Vikberg V, Smith DR, Boevé J-L (2010) How common is ecological

speciation in plant-feeding insects? A "Higher" Nematinae perspective. BMC Evol

Biol 10:266

Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB (2015) Vegan:

community ecology package

[https://cran.r-project.org/web/packages/vegan/index.html].

Peterson AT, Soberon J, Pearson RG, Anderson RP, Martinez-Meyer E, Nakamura M,

Araújo MB (2011) Ecological niches and geographic distributions. Monographs in

Population Biology. Princeton University Press. Princeton. New Jersey, USA

Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species

geographic distributions. Ecol Model 190:231–259

Phillips SJ, Dudík M (2008) Modeling of species distributions with Maxent: new

extensions and a comprehensive evaluation. Ecography 31:161-175

Prior KM, Hellmann JJ (2010) Impact of an invasive oak gall wasp on a native

butterfly: a test of plant-mediated competition. Ecology 91:3284–3293

Polis GA (1999) Why are parts of the world green? Multiple factors control productivity

and the distribution of biomass. Oikos 86:3–15

Rabosky DL (2009) Ecological limits and diversification rate: alternative paradigms to

explain the variation in species richness among clades and regions. Ecol Lett 12:

735– 743

Rathcke BJ (1976) Competition and co-existence within a guild of herbivorous insects.

Ecology 57:76–87

R Core Team (2015). R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/

Redman A, Scriber J (2000) Competition between the gypsy moth, *Lymantria dispar*,

and the northern tiger swallowtail, *Papilio canadensis*: interactions mediated by

host plant chemistry, pathogens, and parasitoids. Oecologia 125:218–228

Ross HH (1957). Principles of natural coexistence indicated by leafhopper populations.

Evolution 11:113–129

Saito MU, Jinbo U, Yago M, Kurashima O, Ito M (2016) Larval host records of

butterflies in Japan. Ecol Res 31:491–491

Schoener TW (1968) The anolis lizards of bimini: Resource partitioning in a complex

fauna. Ecology 49:704–726

Shirôzu T (2006) The butterflies of Japan in color. In Japanese. Gakken Holdings,

Tokyo, Japan.

Shuker DM, Burdfield-Steel ER (2017) Reproductive interference in insects. Ecol

Entomol 42:65−75

Stone L, Roberts A (1990) The checkerboard score and species distributions. Oecologia

85:74–79

Strong DR (1982) Harmonious coexistence of *Hispine* beetles on *Heliconia* In

experimental and natural communities. Ecology 63:1039–1049

Strong DR, Lawton JH, Southwood SR (1984) Insects on plants. Community patterns

and mechanisms. Harvard University Press, Cambridge, MA.

Takami Y, Osawa T (2016) Ecological differentiation and habitat unsuitability

maintaining a ground beetle hybrid zone. Ecol Evol 6:113–124

Thompson JN (2013) Relentless Evolution. University of Chicago Press.

Tojo K, Sekiné K, Takenaka M, Isaka Y, Komaki S, Suzuki T, Schoville SD (2017)

Species diversity of insects in Japan: Their origins and diversification processes.

Entomol Sci 20:357–381

Ueckert DN, Hansen RM (1971) Dietary overlap of grasshoppers on sandhill rangeland

in northeastern Colorado. Oecologia, 8:276-295

Van Zandt PA, Agrawal AA (2004) Specificity of induced plant responses to specialist

herbivores of the common milkweed *Asclepias syriaca*. Oikos 104:401–409

Viswanathan D V, Narwani AJT, Thaler JS (2005) Specificity in induced plant

responses shapes patterns of herbivore occurrence on *Solanum dulcamara*.

Ecology 86:886–896

Waloff N (1979) Partitioning of resources by grassland leafhoppers (Auchenorrhyncha,

Homoptera). Ecol Entomol 4:379-385

Warren DL, Cardillo M, Rosauer DF, Bolnick DI (2014) Mistaking geography for

biology: inferring processes from species distributions. Trends Ecol Evol

29:572–580

Warren DL, Glor RE, Turelli M (2008) Environmental niche equivalency versus

conservatism: Quantitative approaches to niche evolution. Evolution

62:2868–2883

Warren DL (2012) In defense of 'niche modeling'. Trends Ecol Evol 27:497–500

Webb CO, Ackerly DD, McPeek MA, Donoghue MJ (2002) Phylogenies and

community ecology. Annu Rev Ecol Syst 33:475–505

Yoder JB, Clancey E, Des Roches S, Eastman JM, Gentry L, Godsoe W, Hagey TJ,

Jochimsen D Oswarld BP, Robertson J, Sarver BAJ, Schenk JJ, Spear SF Harmon

LJ (2010) Ecological opportunity and the origin of adaptive radiations. J Evol Biol

23:1581–1596

Supporting Materials

Table S1. Japanese butterfly species analyzed in this study.

Table S2. Summary of the MaxEnt data at the 10-km grid scale.

Table S3. Summary of the MaxEnt data at the 1-km grid scale.

Table S4. Correlations among host use similarity (Host), taxonomic relatedness

(Taxon), climate niche similarity (Climate), and total dispersal ability (Dispersal), at

two spatial scales. (a) Summary of Mantel test data on pairwise correlations between the

explanatory matrices. (b) Summary of partial Mantel test data on standardized C_{std}

scores between pairs of butterfly species. The "Taxon-Dispersal" data (b) were obtained

using the datasets at either grid mesh scale.

Table S5. Results using the datasets that excluded rare species. This table corresponds

to Table 2.

Table S6. Results using the datasets that excluded rare species. This table corresponds

to Table S4.

Figure S1 Histogram of the number of observed grids for each butterfly species at both

10-km and 1-km grid scales: (a) all targeted butterfly species at the 10-km grid scale,

(b) butterfly species at the 10-km grid scale, with an observed grid number less than 50,

(c) all targeted butterfly species at the 1-km grid scale, and (d) butterfly species at the

1-km grid scale, with an observed grid number less than 50.

Supplementary File 1. Details of the methods used for ecological niche modeling.

Figure Legends

Figure 1. The relationships between C_{std} scores and host use similarities (10-km grid

scale: Spearman $\rho = -0.128$, $P = 0.0001$; 1-km grid scale: Spearman $\rho = -0.130$, $P =$

0.0001; Mantel test).

Figure 1.

(a) 10km grid scale

(b) 1km grid scale

Table 1 Major hypotheses and predictions tested in this study, associated with host use similarity (Host), taxonomic relatedness (Taxon), climate niche similarity (Climate), and total dispersal ability (Dispersal).

Variables	Hypotheses	Predictions
Host	Resource competition is sufficiently intense to cause competitive exclusion (1).	Positive correlation between host use similarity and C_{std} score (exclusiveness of distribution).
Host	Niche filtering through sharing host plants is relatively stronger than resource competition (2).	Negative correlation between host use similarity and C_{std} score.
Host-Taxon	Taxonomic relatedness reinforces resource competition, increasing competitive exclusion (3-1). Imprint of allopatric speciation reinforces exclusive distribution among close relatives, which share host plants because of niche conservatism (3-2)*.	Correlation between host use similarity and C_{std} score would be changed in the negative direction when controlling for taxonomic relatedness.
Host-Taxon	Taxonomic relatedness, which indicates a niche filter due to niche conservatism including host use, acts counteracting competitive exclusion (4).	Correlation between host use similarity and C_{std} score would be changed in the positive direction when controlling for taxonomic relatedness.
Host-Climate	Filtering through climate niche reinforces resource competition, increasing competitive exclusion (5).	Correlation between host use similarity and C_{std} score would be changed in the negative direction when controlling for climate niche similarity.

Host-Climate	Filtering through climate niche acts counteracting competitive exclusion or jointly with the host plant niche filter (6).	Correlation between host use similarity and C_{std} score would be change in the positive direction when controlling for climate niche similarity.
Host-Dispersal	Higher dispersal ability promotes distribution overlapping, thereby counteracting local assembling processes such as competitive exclusion (7).	Correlation between host use similarity and C_{std} score would change in the negative direction when controlling for total dispersal ability.
Host-Dispersal	Higher dispersal ability promotes distribution overlapping, thereby niche filtering through sharing host plants (8).	Correlation between host use similarity and C_{std} score would change in the positive direction when controlling for total dispersal ability.

The bracketed number following each hypothesis only indicates the number of each hypothesis for descriptive purposes.

*Hypothesis 3-2 has a slightly different perspective from the others, because it is mainly based on the background of speciation history.

Table 2. Correlations among host use similarity (Host), taxonomic relatedness (Taxon), climate niche similarity (Climate), and total dispersal ability (Dispersal), at two spatial scales. (a) Summary of Mantel test data on standardized C_{std} scores between pairs of butterfly species. (b) Summary of Mantel test data on pairwise correlations between host use similarity and the data of other explanatory matrices. (c) Summary of partial Mantel test data on standardized C_{std} scores between pairs of butterfly species. The "Host-Taxon" and "Host-Dispersal" data (b) were analyzed using the same datasets between two grid scales.

(a)

Factor	10-km grid		1-km grid	
	Mantel ρ	P-values	Mantel ρ	P-values
Host	−0.128	**0.0001**	−0.130	**0.0001**
Taxon	−0.025	**0.0093**	−0.036	**0.0009**
Climate	−0.718	**0.0001**	−0.631	**0.0001**
Dispersal	−0.045	**0.0125**	−0.073	**0.0008**

(b)

Factor	10-km grid		1-km grid	
	Mantel ρ	P-values	Mantel ρ	P-values
Host-Taxon	0.098	**0.0001**	0.098	**0.0001**
Host-Climate	0.209	**0.0001**	0.225	**0.0001**
Host-Dispersal	0.061	<u>0.0947</u>	0.061	<u>0.0947</u>

(c)

Factor	Covariate	10-km grid		1-km grid	
		Mantel ρ	P-values	Mantel ρ	P-values
Host	Taxon	−0.126	**0.0001**	−0.127	**0.0001**
Host	Climate	0.033	**0.0044**	0.016	0.1889
Host	Dispersal	−0.125	**0.0001**	−0.126	**0.0001**
Host	All three	0.036	**0.0021**	0.021	<u>0.0890</u>

Spearman's correlation coefficients (ρ values) are shown for all four factors.

Bold: $P < 0.05$; Underlined: $0.05 < P < 0.1$ after 9,999 permutations.

(a) 10km grid scale

(b) 1km grid scale