Disparity, Diversity, and Duplications in the Caryophyllales

Stephen A. Smith[*,1], Caroline Parins-Fukuchi[1], Ya Yang[2], Michael J. Moore[3], Samuel F.

Brockington[4], Riva Bruenn[3,5], Chloe P. Drummond[3,6], Joseph W. Brown[1], Joseph F. Walker[1],

Noah Last[2], Norman A. Douglas[3,7]

[1]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, 48103

[2] Department of Plant Biology, University of Minnesota-Twin Cities, 1445 Gortner Avenue, St.

Paul, MN 55108

[3]Department of Biology, Oberlin College, 119 Woodland St., Oberlin, Ohio 44074-1097

[4]Department of Plant Sciences, University of Cambridge, Cambridge CB2 3EA, United

Kingdom

[5]Current Address: Department of Plant & Microbial Biology, University of California, Berkeley,

Berkeley, California 94720-3102

[6]Current Address: Department of Botany, University of Wisconsin-Madison, 430 Lincoln Drive,

Madison, WI 53706

[7]Current address: Department of Biology, University of Florida, 618A Carr Hall, Gainesville, FL

32605

CORRESPONDING AUTHOR: Stephen A. Smith; 734-615-5510; eebsmith@umich.edu

WORD COUNT

Total: 5285

Summary: 197

Introduction: 1177

Materials and Methods: 1674

Results and Discussion: 2428

This manuscript also contains 3 Main Figures (two in color), 4 Main Tables, and 5

Supplementary Figures.

Summary

- The role whole genome duplication (WGD) plays in the history of lineages is actively debated. WGDs have been associated with advantages including superior colonization, adaptations, and increased effective population size. However, the lack of a comprehensive mapping of WGDs within a major plant clade has led to questions regarding the potential association of WGDs and higher diversification rates.

- Using seven chloroplast and nuclear ribosomal genes, we constructed a phylogeny of 5,036 species of Caryophyllales, representing nearly half of the extant species. We phylogenetically mapped putative WGDs as identified from analyses on transcriptomic and genomic data and analyzed these in conjunction with shifts in climatic niche and lineage diversification rate.

- Thirteen putative WGDs and twenty-seven diversification shifts could be mapped onto the phylogeny. Of these, four WGDs were concurrent with diversification shifts, with other diversification shifts occurring at shallower nodes than WGDs. Five WGDs were associated with shifts to colder climatic niches.

- While we find that many diversification shifts occurred after WGDs it is difficult to directly associate these and consider diversification and duplication to be tightly correlated. Our findings suggest that duplications may often along with shifts in either diversification rate, climatic niche, or rate of evolution.

Keywords: Caryophyllales, duplications, climatic niche, diversification rates, phylogenomics

Introduction

Understanding the causes and correlates of diversification within flowering plants has been a central goal of evolutionary biologists. Genomic and transcriptomic data have reinvigorated hypotheses associating whole genome duplication (WGD) with lineage diversification rate increases (e.g., Levin, 1983; Levin 2002; Barker et al. 2009; Soltis *et al.*, 2014; Edger et al. 2015; Puttick et al. 2015; Tank *et al.*, 2015; Barker et al. 2016; Huang et al. 2016; McKain et al. 2016). It is not self-evident why WGDs would be associated with increases in lineage diversification. One hypothesis suggests that the additional genetic material provides a basis to generate new adaptations (Edger *et al.*, 2015), although this itself assumes a co-occurrence of adaptation and lineage proliferation (Levin, 1983). The apparent lack of precise co-occurrence of

adaptation and lineage proliferation has been explained by the potential of a lag model (Tank et

al. 2015) where diversification may follow WGD events. In the absence of overwhelming

correlative signal, we are often unable to discern true ancient WGD events from aneuploidy

without advanced genomic information such as synteny mapping (Dohm *et al.*, 2012). Because it

is often difficult to distinguish the two, for simplicity we will define WGD broadly to include

putative ancient WGD events (paleopolyploidy) and ancient aneuploidy events. WGD events are

thought to be a common occurrence and have been associated with an estimated 15% of

angiosperm speciation events (Wood *et al.*, 2009). However, whether speciation by WGD is

correlated with higher diversification rates remains highly debated (Mayrose *et al.*, 2011; Soltis

et al., 2014; Tank *et al.*, 2015). Analyses based on recent WGD events have concluded that

immediate extinction rates are higher for polyploid plants (Mayrose *et al.*, 2011; Arrigo and

Barker, 2012). This may result from small initial population sizes and an increased dependence

on selfing. Alternatively, despite the disadvantages of WGD, others have suggested that

polyploids may be superior colonizers (Soltis and Soltis, 2000).

Indeed, extreme environments are associated with high levels of WGD, with up to 87%

of species restricted to areas that were glaciated during the last ice age consisting of polyploids

(Brochmann, 2004). However, in the example from Arctic plants, the high level of WGD has

occurred post-glaciation representing a micro-evolutionary period whereas previous studies often

focus at much deeper macro-evolutionary time scales (Mayrose *et al.*, 2011; Tank *et al.*, 2015;

Soltis *et al.*, 2014). From the perspective of a short timescale, polyploidy has the disadvantages

of higher error rates in mitosis (Storchová et al., 2006) and masking of deleterious mutations

allowing them to accumulate to higher frequencies in a population (Otto and Whitton, 2000). A

suite of advantages however may also arise, including gain of asexuality (Miller *et al.*. 2000) and

varying effects of heterosis (Comai, 2005). The net role these advantages and disadvantages play

on the macroevolutionary scale is difficult to determine from either the purely short-term or

purely long-term time scales previously used.

The long-term consequence of WGD is a central question in macroevolution and

comparative genomics. However, with a suite of advantages and disadvantages, much debate

surrounds the importance and patterns of correlation of WGD (Comai 2005). While

polyploidization events can cause instant speciation, there is no reason to assume that these

singular speciation events in themselves would influence large-scale diversification rate shifts

when considering lineage survivorship. Instead, there may be other factors, such as the increase

in genetic material, perhaps increasing genetic diversity or enabling adaptation, that cause long

term shifts in rates of diversification. Adaptations need not be associated with shifts in the tempo

of diversification and those adaptations and shifts in diversification may not co-occur on the

same branch (i.e., there may be a lag time; Donoghue 2005; Smith et al. 2011, Donoghue and

Sanderson 2015; Tank et al. 2015). In the broader context of plant evolution, there are several

possible outcomes of WGD in relation to the evolution and diversification of clades: 1) no

relationship between WGD and speciation rate or habitat shift/adaptation, 2) WGD coincides

with an increase of speciation rate, with or without a lag time, 3) WGD promotes dispersal and

habitat shifts, which has mixed relationship with speciation rate, and 4) a mixture (some

association, some not), similar to the previous hypothesis but without explicitly promoting

dispersal or habitat shift or speciation (e.g., adaptation could be more prominent than dispersal

and habitat shift). Here, we contribute to this discussion on diversification and WGDs with an in-

depth examination of the intersection of diversification and WGDs happening at a range of scales

within the hyperdiverse Caryophyllales.

The Caryophyllales contains ~12,500 species in 39 families (Thulin *et al.*, 2016; APG IV:

Chase *et al.*, 2016), representing approximately 6% of extant angiosperm species diversity. The

estimated crown age of Caryophyllales is approximately 67–121 millions of years ago (mega-

annum, Ma) (Bell *et al.*, 2010; Moore *et al.*, 2010) and species of the Caryophyllales exhibit

extreme life-history diversity, ranging from tropical trees to temperate annual herbs, and from

desert succulents (e.g., Cactaceae) to a diverse array of carnivorous plants (e.g., the sundews

Drosera and pitcher plants *Nepenthes*). Such extraordinary diversity makes Caryophyllales a

particularly useful system for investigating the relationship between WGD vs. diversification and

niche evolution. Our previous analyses using 62 transcriptomes representing 60 species across

the Caryophyllales identified 13 well-supported ancient WGD events (Yang *et al.*, 2015). We

have since nearly tripled the taxon sampling and assembled a data set comprising high-coverage

transcriptomes and genomes from 169 species across the Caryophyllales (Yang *et al.*,

submitted), providing even greater power for resolving the number and phylogenetic locations of

WGD events. Moreover, the growth in the number of plant taxa on GenBank that are represented

by traditional targeted sequences (e.g., *rbc*L, *mat*K, ITS, etc.) and the growth of publicly

available collections data (e.g., GBIF, iDigBio) provide excellent opportunities to apply

megaphylogeny and niche diversification approaches at fine scales in Caryophyllales.

By examining WGDs and diversification within the Caryophyllales, we present an

important example. Not only does the dataset examined have a high density of transcriptomic

sampling, the diversification of the bulk of Caryophyllales occurred during a time frame

intermediate to that of most published studies that have probed a link between WGD and

macroevolution. This time frame, between 10 and 100 Ma, is important for angiosperms as much

of the diversification that has led to the modern flora occurred during this period and most

modern angiosperm families appeared by this time. Discussion of speciation rate, niche shift, and

WGD would be flawed without accurate mappings of WGD events within this time scale. We

compiled a data set with extensive and precise mapping of WGD combined with a species-level

phylogeny. The megaphylogeny approach has been used extensively in the past to combine data

from many gene regions and across broad taxonomic groups to address evolutionary questions

(Smith *et al.*, 2009). Here, we use this approach to help inform analyses from phylogenomic

studies, and provide a broad context in which to examine these genomic phenomena. With half

of the species sampled, this represents to date the largest and most exhaustive study of WGD vs.

rate and adaptive shift.

Materials and Methods

Sanger sequencing and assembly. —A total of 248 new *matK* sequences were included in this

study (Table 1). To generate these sequences, leaf samples were collected in silica in the field or

from cultivated material, or were collected from herbarium sheets. DNA was isolated using

either the Nucleon Phytopure kit (GE Healthcare Life Sciences, Pittsburgh, PA, USA), using the

0.1 g protocol and following manufacturer's instructions, or using the Doyle and Doyle (1987)

protocol, with the addition of 1% PVP-40. An approximately 950 bp region in the middle of the

matK gene was amplified and sequenced using custom-designed primers (Table 2). PCRs were

performed in 12.5 μL volumes with 0.5 μL of 5 mM primer for both primers, 5-20 ng of DNA

template, 0.1 μL of GoTaq (Promega, Madison, WI, USA), 6.25 μL of Failsafe Premix B

(Epicentre, Madison, WI, USA), and 4.7 μL of sterile, deionized water. Reactions were run on a

Bio-Rad PTC 200 thermocycler (Bio-Rad, Hercules, CA, USA) at Oberlin College. Individual

PCRs were cleaned in 16.5 μL reactions containing 10 U of Exonuclease I (Affymetrix,

ThermoFisher Scientific, Waltham, MA, USA), 2 U of shrimp alkaline phosphatase

(Affymetrix), 8 μL of PCR product, and 8.5 μL of sterile, deionized water. Sanger sequencing of

the resulting cleaned PCRs was conducted by Neogenomics (formerly SeqWright; Houston, TX,

USA) using an ABI 3730xl automated sequencer (Applied Biosystems, ThermoFisher

Scientific). The resulting forward and reverse sequences for each reaction were trimmed and *de*

novo assembled using default parameters of the Geneious assembler in Geneious versions 5-7

(Biomatters, Auckland, New Zealand).

Molecular Data for Phylogenetic Reconstruction. —Nucleotide data from the nuclear

ribosomal internal transcribed spacers (ITS) and *phyC* gene, and the plastid loci *matK, ndhF,*

rbcL, trnH-psbA spacer, and *trnL-trnF* spacer were used to reconstruct the phylogeny. These data

were gathered first using PHLAWD (Smith and Donoghue, 2008; Smith *et al.*, 2009) and then

curated and combined with newly sequenced *matK* data for 124 additional species. This yielded

the following sampling: ITS 2,969 species, *matK* 2,270 species, *ndhF* 417 species, *phyC* 172

species, *rbcL* 947 species, *trnH-psbA* 240 species, and *trnL-trnF* 1,996 species. We used *mat*K,

*rbc*L, and *ndh*F sequences from *Aextoxicon, Apium, Berberidopsis, Campanula, Clethra, Coffea,*

Echinops, Helwingia, Ilex, Ipomoea, Lamium, Lonicera, Nyssa, Polysoma, Primula, Santalum,

Valeriana, and *Viburnum* to represent outgroups.

Phylogenetic Reconstruction. —We conducted phylogenetic analyses with RAxML v7.2.8

(Stamatakis, 2014) using the full analysis command, -f a, which conducts a rapid bootstrap and

then a full maximum likelihood search. The combined bootstrap and maximum likelihood search

allows for a more thorough for maximum likelihood analysis where the initial rapid bootstrap

results prime the maximum likelihood analysis. However, we did not use the rapid bootstrap

trees from this analysis and instead, we conducted a full bootstrap, generating the bootstrap

dataset using phyx (Brown *et al.*, 2017) and then conducting individual maximum likelihood

runs on each constructed bootstrap dataset. We conducted bootstraps within gene regions and we

retained the individual bootstrap alignments to conduct additional analyses (i.e., bootstrapped

alignments contained the same number of gene-specific sites as the empirical alignment). On

each of the resulting trees of the bootstrap and the maximum likelihood tree, we conducted SH-

like approximate likelihood ratio tests (SH-aLRT; Guindon *et al.*, 2010) as implemented in

RAxML. These analyses calculate support for each edge while also finding the NNI-optimal

topology. RAxML completed the likelihood search for each of these bootstrap replicates,

however the SH-aLRT analyses often resulted in an improved maximum likelihood topology.

The trees that resulted from the SH-aLRT, ML and bootstrap samples, were used for further

analyses. Because several deep relationships within Caryophyllales are hard to resolve without

large amounts of molecular data that are unavailable for most of the taxa included in this analysis

(Yang *et al.*, 2015), for all phylogenetic analyses we applied the following topological constraint:

(Droseraceae, (*Microtea*, (Stegnospermataceae, Limeaceae, (Lophiocarpaceae, (Barbeuiaceae,

Aizoaceae))))) as per previous analysis (Brockington et al. 2009; Yang *et al.*, 2015).

Divergence Time Estimation. — Few tractable options for divergence time estimation exist for

datasets of the size presented here. We use the penalized likelihood approach (Sanderson, 2003)

as implemented in the program treePL (Smith and O'Meara, 2012), which can handle large-scale

phylogenies. The early fossil record of the Caryophyllales is sparse with only a few known

records (Friis *et al.*, 2011; Arakaki *et al.*, 2011): (1) fossil pollen has been ascribed to

Amaranthaceae (*Chenopodipollis*) from the Paleocene of Texas (Nichols and Traverse, 1971);

(2) a putative fossil infructescence from within the Phytolaccaceae in the Campanian has also

been reported (Cevallos-Ferriz *et al.*, 2008), but this phylogenetic position has been disputed

(pers. comm. S. Manchester) and hence we excluded it; (3) Jordan and Macphail (2003) describe

a middle to late Eocene inflorescence from the species *Caryophylloflora paleogenica,* ascribed to

Caryophyllaceae; (4) pollen from Argentina within the Nyctaginaceae has been reported from the

middle Eocene (Zetter *et al.*, 1999); and (5) (Degreef, 1997). The penalized likelihood method

performs better when a calibration is used at the root. For this calibration, and because there is no

fossil record for the earliest Caryophyllales, we use a secondary calibration from the

comprehensive angiosperm divergence time analyses of Bell *et al.* (2010). Several other

secondary calibrations were attached to major clades where fossils are not available (Ocampo

and Columbus 2010; Arakaki et al. 2011; Schuster et al. 2013; Valente et al. 2013; see Supp.

Table S1 for detail on placement and calibrations). We conducted a priming analysis to

determine the best optimization parameter values. We then performed a cross validation analysis

using the random cross validation setting to determine the optimal smoothing parameter value.

215 ***Climate analyses.*** —We downloaded 6,592,700 georeferenced occurrences for the

216 Caryophyllales from GBIF (accessed on 6/1/2015; http://gbif.org). After removing samples

217 present in living collections, and therefore not necessarily representative of native climates, and

218 removing samples whose localities were over water, there were 6,009,552 samples. We extracted

219 bioclimatic values for each coordinate using the 2.5 arc-minute resolution data from WorldClim

220 (http://worldclim.org). We only included taxa that had at least three samples in these analyses to

221 reduce potential errors and to have the minimum number of samples required to calculate mean

222 and variance. The resulting overlap of the taxa represented in both the geographic and genetic

223 data was 2,843 taxa. We conducted principal component analyses (PCA) on these extracted

224 values. With both the bioclimatic values and the first two axes of the PCA, we conducted

225 ancestral state reconstruction analyses.

226  We also conducted contrast analyses and calculated Brownian motion rates of evolution

227 between sister clades (comparing duplicated lineages with their sisters) for mean annual

228 precipitation, mean annual temperature, and principal component axis 1. Contrasts were

229 calculated using phylogenetic independent contrasts. Brownian motion rates were calculated on

230 sister lineages independently using the analytical solution for rate: $\sigma^2 = \frac{1}{n}\sum_{i=1}^{n}\frac{u_i^2}{v_i^2}$.

232 ***Diversification analyses.*** —To map diversification rate shifts, we conducted MEDUSA (Alfaro

233 *et al.*, 2009; Pennell *et al.*, 2014) analyses on the maximum likelihood tree and the bootstrap

234 trees. MEDUSA is far more computationally tractable than some other diversification estimation

235 methods (e.g., BAMM). Furthermore, we required the ability to feasibly integrate over the

236 phylogenetic uncertainty within the phylogenetic dataset because of both the nature of the larger

237 phylogenetic dataset and the inherent biological uncertainty within the Caryophyllales.

238 MEDUSA fits a birth-death model of diversification (with parameters *r*: net diversification (birth

239 - death), and ε: relative extinction (death / birth)) before using stepwise AIC (Burnham and

240 Anderson, 2002) to identify shifts in rates of diversification. These analyses allow

241 complementary analyses targeted at accommodating topological and branch length uncertainty.

242 We performed these diversification analyses using a birth-death model on 97 chronograms

243 generated from nonparametric bootstrapping of the original matrix, inferring ML trees in

244 RAxML, and estimating divergence times in treePL using the temporal constraints described

above. We discarded three trees based on poor fossil placement resulting from phylogenetic

uncertainty causing fossil placements to conflict.

Whole Genome Duplication Identification. — To identify WGDs (procedure described below),

we generated a tree based on transcriptomic data. For this tree, we used 178 ingroup data sets

(175 transcriptomes, 3 genomes) representing 169 species in 27 families and 40 outgroup

genomes (Table S1-S2 in Yang et al. submitted). We mapped putative WGD events using

multiple strategies: gene tree topology, plotting synonymous distance, and chromosome counts

(Yang et al. 2015; Yang *et al.*, submitted). For gene tree topology analyses, we performed two

alternative strategies for mapping duplication events from gene trees to the species tree: mapping

to the most recent common ancestor (MRCA), or mapping to species tree only when gene tree

and species tree topologies are compatible.

To conduct synonymous distance analyses, we performed the following procedure. For

all ingroup Caryophyllales transcriptome data sets, we calculated the distribution of paralog

synonymous distance following the same procedure as (Yang *et al.*, 2015). We reduced highly

similar peptide sequences with CD-HIT (-c 0.99 -n 5). We also carried out an all-by-all BLASTP

within each taxon using an E value cutoff of 10 and -max_target_seq set to 20. Resulting hits

with pident < 20% or niden < 50 amino acids were removed. We removed sequences with ten or

more hits to avoid overrepresentation of gene families that experienced multiple recent

duplications. We used the remaining paralog pairs and their corresponding CDS to calculate Ks

values using the pipeline https://github.com/tan ghaibao/bio-

pipeline/tree/master/synonymous_calculation (accessed November 29, 2014). The pipeline first

carries out pairwise protein alignment using default parameters in ClustalW (Larkin *et al.*, 2007),

back-translates the alignment to a codon alignment using PAL2NAL (Suyama *et al.*, 2006), and

calculates the synonymous substitution rate (Ks) using yn00 as part of the PAML package

(Yang, 2007), with Nei–Gojobori correction for multiple substitutions (Nei and Gojobori, 1986).

We obtained chromosome counts from the Chromosome Counts Database (CCDB;

http://ccdb.tau.ac.il accessed Oct 5, 2015). When multiple counts were reported from different

authors or different plants, we erred on the conservative estimate and recorded the lowest

number. For species that were not available in the database, we found counts from the literature

(e.g., Jepson eFlora http://ucjeps.berkeley.edu/eflora/ and Flora of North America

http://floranorthamerica.org) or by a consensus from species of the same genera.

Results and Discussion

Phylogenetic results. —Phylogenetic analyses showed strong support based on bootstrap and

SH-aLRT values for the monophyly of most Caryophyllales families (see Fig. S1). We find

strong support for the carnivorous clade including Droseraceae, Ancistrocladaceae,

Nepenthaceae, Drosophyllaceae, and Dioncophyllaceae. There is also strong support for this

clade as sister to a clade including Frankeniaceae, Tamaricaceae, Plumbaginaceae, and

Polygonaceae. However, relationships among the families show more varied support. There is

weak support for the placement of other families relative to other early diverging Caryophyllales

(see Fig. S1). There is strong support for Caryophyllaceae sister to Amaranthaceae. There is very

weak support for Aizoaceae sister to Phytolaccaceae+Nyctaginaceae. As with previously

published analyses, there was no support for the monophyly of Phytolaccaceae in the traditional

sense (i.e., including Phytolaccaceae s.s., Petiveriaceae, and *Agdestis*; APG IV) and very weak

support for the placement of Sarcobataceae. There is also weak support for the relationships

among Limeaceae, Molluginaceae, and the Portulacineae. Many of these relationships have been

found to be strongly supported but conflicting in different analyses (Brockington *et al.*, 2009;

Soltis *et al.*, 2011; Yang *et al.*, 2015; Smith *et al.*, 2015; Walker *et al.*, 2017). Here, we focus less

on the systematic resolution within the Caryophyllales and instead examine the potential

relationship of diversification and climate shifts to WGDs. Therefore, we place more emphasis

on including more taxa over that of more gene regions (i.e., transcriptomes) at the cost of more

missing data. Confident resolution of many of the systematic relationships will require genomic

and transcriptomic sampling, and more thorough taxon sampling (Yang *et al.*, submitted).

Climate tolerance reconstruction results. —We performed climate ancestral reconstruction

analyses on the phylogeny of 2,843 taxa that included taxa with at least three sampled

geographic coordinates (Figs. 1-3). These analyses were conducted for visualization and for

comparison with diversification and WGD results (see below). Results for individual bioclimatic

variables and principal components can be found in Figs. S2-S4. Bioclimatic variable 1 (mean

annual temperature, Fig. 1) shows that there are several strong phylogenetic patterns of clades

with preferences for colder or warmer regions. For example, Polygonaceae, Caryophyllaceae,

and Montiaceae each are dominated by taxa with preferences for cold environments, although

each also contains early-diverging taxa with preferences to warm environments. In contrast, taxa

inhabiting warm environments predominate in Cactaceae, Amaranthaceae, Aizoaceae, the

carnivorous clade (Droseraceae, Drosophyllaceae, Nepenthaceae, Ancistrocladaceae,

Dioncophyllaceae), and the phytolaccoid clade (Nyctaginaceae, Phytolaccaceae, Petiveriaceae,

Sarcobataceae, and *Agdestis*). Bioclimatic variable 12 (mean annual precipitation) shows a

relatively consistent pattern of relatively dry to intermediately wet clades throughout the group.

Indeed, only a few clades inhabiting wet ecosystems (in this case, the wet tropics) exist in the

Caryophyllales, specifically small groups within the carnivorous clade, the phytolaccoids, early-

diverging Polygonaceae, and other small groups throughout the Caryophyllales. The principal

component loadings are presented in Fig. 2 and Fig. S5. Principal component 1, PCA1, shows

significant differentiation throughout the Caryophyllales, as for example, early-diverging

Polygonaceae vs the rest of Polygonaceae, early diverging Caryophyllaceae vs the rest of

Caryophyllaceae, phytolaccoids vs Aizoaceae, and Portulacineae + relatives vs Cactaceae, to

mention a few. These results generally reflect the extensive ecological diversification throughout

the group. They also reflect significant diversification in the temperate regions of the world

especially within the Caryophyllaceae and Polygonaceae contrasted with extensive

diversification in the succulent lineages (especially Aizoaceae and Cactaceae) found in relatively

dry and warm environments.

Diversification. —Significant shifts in diversification were detected in most major clades (Table

4, Fig. 1). The results from diversification analyses on the maximum likelihood tree and

bootstrap tree set are generally congruent with each other. However, there are discrepancies (Fig.

1). The bootstrap set recovered many shifts in Polygonaceae, the carnivorous clade,

Caryophyllaceae, some shifts within Cactaceae, phytolaccoids, and Amaranthaceae.

Disagreements on the existence and placement of shifts are primarily within Portulacineae,

Aizoaceae, and Amaranthaceae. Overall, MEDUSA detected 27 increases in diversification rate

using the ML tree and 16 increases using the bootstrap trees. Given the relative lack of support of

some of the branches in the phylogeny, we find the MEDUSA results on the set of bootstrapped

trees to be the most conservative while the ML results are suggestive but not definitive of

diversification shifts.

Duplications, diversification, and climate. —WGD analyses show thirteen putative WGDs that

can be mapped to clades (i.e., involve more than 1 taxon in the dataset; Table 3 and Figs. 1-3).

Many of these are found in early diverging lineages as opposed to nested deep within families,

though there are WGDs identified in *Amaranthus* and *Claytonia*. We also find evidence of nested

WGDs as within the phytolaccoids and Portulacineae. In addition to these deeper WGDs, there

are larger numbers of more recent WGDs that are present in Ks plots but cannot be mapped to a

clade (Yang *et al.*, submitted). By sampling more extensively, Yang *et al.* (submitted) and

Walker *et al.* (2017) found additional WGD events within the Caryophyllales. While it is

possible, this is unlikely to be phenomenon specific to the Caryophylalles and we will likely find

additional WGDs events in other lineages as more effort is placed on denser taxon sampling

using genomes and transcriptomes. We do not explore WGDs that can only be mapped to one tip

any further and more discussion of specific results related to the WGDs themselves can be found

in Yang et al. (submitted) and Walker et al. (2017).

To better examine whether WGDs coincide with diversification rate shifts, increases and

decreases, or notable changes in climate tolerance, we mapped WGDs onto the large phylogenies

and summarized the number of species and climate information for each clade (Tables 3-4, Figs.

1-3). Some WGD events are associated with synchronous diversification events. For example,

within Nyctaginaceae, a WGD event occurs on the same branch (leading to Tribe Nyctagineae;

Douglas and Spellenberg, 2010) as an increase in diversification rate in both the ML tree and the

bootstrapped dataset (Fig. 1, dup:1 div:n). These events are further associated with a shift in life

history and niche from an ancestral woody habit in the tropics to the largely herbaceous, arid-

adapted temperate Nyctagineae. This is also the case for *Amaranthus* (Fig. 1, dup:5 div:x). Other

coincident diversification and WGD events in the Droseraceae and Nepenthaceae are only

supported by the ML tree. Although these correlated events may, in fact, be accurate, we will

reserve more comments for when these are more confidently resolved. Other than these

simultaneous shifts, and excepting one diversification shift at the base of the MRCA of

Nyctaginaceae+Cactaceae, all other shifts in diversification occur more recently than WGD

events. Tank *et al.* (2015) suggested that this lagging pattern may be common at the broader

angiosperm scale, though the expected distance of the diversification shift from the WGD event
was not specified (this is discussed more below). In the results presented here, some
diversification events occur shortly after the WGD event, such as within the Amaranthaceae
(dup: 6) and Portulacineae (dup: 4). For others, it is difficult to determine whether the
diversification events that occur after the WGD events are significantly close to the WGD to
warrant suggestion of an association (e.g., dup: 7, dup: 10, dup: 8). More description of a model
that would generate a null expectation would be necessary to determine what is "close enough"
(see discussion below).
Many of the other inferred lineage diversification rate shifts are associated with very
recent, rapid radiations within genera such as those documented within *Commicarpus*
(Nyctaginaceae), *Dianthus* (Caryophyllaceae), *Cerastium* (Caryophyllaceae), *Arenaria*
(Caryophyllaceae), and *Salicornia* (Amaranthaceae), to name a few (Table 4). Although
polyploids were reported in these clades, we are unable to pinpoint the phylogenetic location of
any WGD with our current taxon sampling (e.g., *Dianthus*; Carolin, 1954; Weiss *et al.* 2002).
Increased sampling of transcriptomes and genomes will shed more light in these areas. While we
only find a few WGDs that coincide well with diversification rate shifts, it is important to note
that the uncertainty in the phylogenies makes it difficult to map anything but the strongest
diversification signals. This discrepancy can be seen in the difference between the number of
events supported by the ML analyses and those supported by the bootstrap analyses. It is possible
that additional sequence data will improve phylogenetic resolution and confidence, and that
consequently additional diversification events will emerge.
Equally interesting to the few WGD events associated directly with diversification are the
WGD events associated with general shifts in climate tolerance. WGDs in the Polygonaceae,
Caryophyllaceae, Montiaceae, and the Tribe Nyctagineae appear to be associated with movement
into colder environments (Figs. 1-2 and Figs. S2-S3). Species arising after the WGD within the
Amaranthaceae occupy wetter environments than the sister clade. The WGDs within the
carnivorous plants are also associated with shifts in environment as *Nepenthes* are found in very
wet environments and the Droseraceae are found in somewhat drier environments, at least
comparatively. However, in these cases, perhaps the development of the wide array of
morphologies associated with carnivory, apart from *Drosophyllum*, is more obviously associated
with the WGD (Walker *et al.*, 2017).

While these qualitative assessments suggest potential correlations of shift in the climate

occupied and WGDs, more specific and direct comparisons are necessary to quantify the extent

of the shifts. For many of the clades experiencing WGD, a direct comparison with a sister clade

is difficult because the sister may consist of a single species, another clade with WGD, or

another complication. For example, there are WGDs at the base of both Polygonaceae and

Plumbaginaceae as well as Nepenthaceae and Droseraceae. However, we made direct

comparison of five duplicated lineages (see Fig. 3) in both means (i.e., character contrasts

between sister clades) and variances (rate of Brownian motion) of climatic variables. In each

case, the duplicated lineage occupies a colder mean annual temperature. This is also the case

with the nested WGDs of Portulacineae and the Tribe Nyctagineae. Of course, we do not suggest

that all WGDs are associated with a shift to a colder climate. While such a pattern may exist in

some groups such as Caryophyllaceae, we emphasize the observation that there is a shift in the

climate occupied rather than the direction of the shift. Mean annual precipitation is not as clear

with some clades occupying a higher precipitation and some occupying lower precipitation.

Perhaps the best summary of climatic niche is the principal components of all the climatic

variables. Here, while the shift in units is less easily interpreted, duplicated clades occupy

different niches than sister lineages. This generally supports the hypothesis that WGD events are

associated with adaptations, in this case, that are associated with shifts in climatic niches. This

necessitates further examination in other angiosperm clades.

The rates of niche evolution show more complicated patterns. While some clades, such as

the Portulacineae, show significant increase in a rate of niche evolution as compared to the sister

clade (e.g., MAT), no clear pattern emerges across all comparisons. There are other shifts in rate

such as with MAT and MAP in the Nyctaginaceae and Montiaceae, but these are not as strong as

the pattern of climate itself discussed above.

With each of these patterns presented here, it is important to consider them in the context

of uncertainty, both inherent in the biological processes that generate the phylogeny and in the

analyses associated with large scale datasets. These large phylogenies and datasets allow for

more thorough examination of the clades, but uncertainty makes precise mapping of weaker

signals difficult. As mentioned above, this is demonstrated by both the mapping of

diversification events and duplications. Furthermore, the comparisons of the sister clades for

climatic niche analyses assumes accurate identification of sister lineages. Increasing taxon

sampling may help, but additional sequence data and specimen data for phylogenetic analyses,

WGD mapping analyses, and climate niche characterization will surely improve our precision in

these investigations.

What emerges from these analyses of WGD, diversification, and climate? It would appear

as though, perhaps not unexpectedly, the patterns are complex and mixed. Some WGD are

associated directly with diversification events, some WGD are associated with shifts in climate

tolerance, some WGD are coincident with shifts in rates of niche evolution, and still other WGD

are associated with known adaptations (carnivory, habit shifts associated with montane habitats,

etc.). Some diversification shifts follow WGD events. However, it is unclear whether these

events are linked or correlated and, if so, if they are correlated more with diversification than an

additional adaptation or other evolutionary pattern or process. As data increase in these groups

and as confidence increases in the phylogenetic relationships as well as the placement of both

diversification and WGD events, we will be able to better address these questions. However, at

least for the Caryophyllales, it does not appear as though diversification is tightly linked with

WGD. Instead, for the clades that can be tested, we find shifts in climate correspond well to

WGD.

Suggestions for moving forward. —WGD are almost certainly one of the dominant processes

that contribute to major evolutionary events within plant lineages. This may be in the form of

increased diversification, development of novel traits, adaptation to new environments, and many

other events (e.g., Schubert and Vu, 2016; Clavijo et al. 2017). However, for several reasons,

these events (i.e., WGD and other evolutionary events) may not occur simultaneously. In fact,

there may be little to no expectation for the events to occur simultaneously (e.g., Donoghue,

2005; Donoghue and Sanderson, 2015; Tank *et al.*, 2015). In any case, however, more precise

expectations and null models need to be developed to allow for reasonable tests of the

correlations among these events. For example, there may be shifts in diversification that follow a

WGD, but is it close enough, or frequent enough to infer that the two events are related? Is

correlation possible or identifiable if, as is expected, intervening lineages have gone extinct?

Furthermore, more precise connections should be made to the biology of speciation and genome

WGDs to better determine why, specifically, WGDs would be expected to correspond with any

diversification pattern instead of adaptations, which may or may not correspond with increases or

decreases in speciation. While still challenging, investigating the fate of and patterns of selection

within individual genes (e.g., subfunctionalization and neofunctionalization) may shed light into

the genomic basis of post-WGD and possibly allow for more concrete expectations for

diversification. With the availability of genomes and transcriptomes, this is now beginning to

become a possibility (e.g., Brockington et al., 2015, Walker *et al.*, 2017). Only when these

suggestions are linked to more specific biological hypotheses will we be able to better

understand the ultimate impact of WGD in plant evolution.

Acknowledgements

We thank Gregory Stull, Oscar Vargas, Ning Wang, Sonia Ahluwalia, Jordan Shore, Lijun Zhao,

Alex Taylorm, and Drew Larson for helpful discussion on the manuscript. The authors thank

Hilda Flores, Helga Ochoterena, Tom Wendt and the staff at the Plant Resources Center at the

University of Texas at Austin, the Lyon Arboretum, David Anderson, John Brittnacher, Anna

Brunner, Joseph Charboneau, Arianna Goodman, Heather-Rose Kates, Patricia Herrnández

Ledesma, Lucas Majure, Nidia Mendoza, Michael Powell, Rick Ree, Carl Rothfels, Flora Samis,

Jeffrey Sanders, Elizabeth Saunders, Rich Spellenberg, Greg Stull, Mats Thulin, Erin Tripp, and

Sophia Weinmann for help with obtaining material. This work was supported by NSF DEB

awards 1352907 and 1354048.

Author contributions

S.A.S., J.F.W., Y.Y., M.J.M., C.P.-F., and S.F.B. designed research. S.A.S., J.W.B., C.P.F., and

Y.Y. analyzed the data. S.A.S. led the writing. All authors read and contributed to the

manuscript.

References

Alfaro ME, Santini F, Brock C, Alamillo H, Dornburg A, Rabosky DL, Carnevale G,

Harmon LJ. 2009. Nine exceptional radiations plus high turnover explain species

diversity in jawed vertebrates. *Proceedings of the National Academy of Sciences, USA*

106: 13410–13414.

Angiosperm Phylogeny Group. 2016. An update of the Angiosperm Phylogeny Group
classification for the orders and families of flowering plants: APG IV. Botanical Journal
of the Linnean Society 181: 1-20.

Arakaki M, Christin PA, Nyffeler R, Lendel A, Eggli U, Ogburn RM, Spriggs E, Moore
MJ, Edwards EJ. 2011. Contemporaneous and recent radiations of the world's major
succulent plant lineages. *Proceedings of the National Academy of Sciences, USA* **108:**
8379–8384.

Arrigo N, Barker MS. 2012. Rarely successful polyploids and their legacy in plants genomes.
Current Opinion in Plant Biology **15:** 140–146.

Barker MS, Vogel H, Schranz ME. 2009. Paleopolyploidy in the Brassicales: Analyses of
the *Cleome* transcriptome elucidate the history of genome duplications in *Arabidopsis*
and other Brassicales. *Genome Biology and Evolution* **1**: 391–399.

Barker MS, Husband BC, Pires JC. 2016. Spreading Winge and flying high: The evolutionary
importance of polyploidy after a century of study. *Am. J. Bot.* **103**: 1139-1145.

Bell CD, Soltis DE, Soltis PS. 2010. The age and diversification of the angiosperms re-revisited.
American Journal of Botany **97:** 1296–1303.

Brochmann C, Brysting AK, Alsos IG, Borgen L, Grundt HH, Scheen AC, Elven R. 2004.
Polyploidy in arctic plants. *Biological Journal of the Linnean Society* **82:** 521–536.

Brockington SF, Alexandre R, Ramdial J, Moore MJ, Crawley S, Dhingra A, Hilu K, Soltis
DE, Soltis PS. 2009. Phylogeny of the Caryophyllales *sensu lato*: revisiting hypotheses
on pollination biology and perianth differentiation in the core Caryophyllales.
International Journal of Plant Sciences **170:** 627–643.

Brockington SF, Yang Y, Gandia-Herrero F, Covshoff S, Hibberd JM, Sage RF, Wong GK,
Moore MJ, Smith SA. 2015. Lineage-specific gene radiations underlie the evolution of
novel betalain pigmentation in Caryophyllales. *New Phytologist* **207:** 1170–1180.

Brown JW, Walker JF, Smith SA. 2017. phyx: Phylogenetic tools for Unix. *Bioinformatics*
btx063.

Burnham KP, Anderson DR. 2003. Model Selection and Multimodel Inference: A Practical
Information-Theoretic Approach. Springer, New York.

Carolin RC. 1954. Stomatal size, density and morphology in the genus *Dianthus*. *Kew Bulletin*
9: 251-258.

Cevallos-Ferriz SRS, Estrada-Ruiz E, Perez-Hernandez BR. 2008. Phytolaccaceae

infructescence from Cerro del Pueblo formation, upper Cretaceous (late Campanian),

Coahuila, Mexico. *American Journal of Botany* **95:** 77–83.

Chase MW, Christenhusz MJM, Fay MF, Byng JW, Judd WS, Soltis DE, Mabberley DJ,

Sennikov AN, Soltis PS, Stevens PF. 2016. An update of the Angiosperm Phylogeny

Group classification for the orders and families of flowering plants: APG IV. *Botanical*

Journal Linnean Society **181:** 1-20.

Clavijo BJ, Venturini L, Schudoma C, Accinelli GG, Kaithakottil G, Wright J, Borrill P,

Kettleborough G, Heavens D, Chapman H, Lipscombe J, Barker T, Lu FH,

McKenzie N, Raats D, Ramirez-Gonzalez RH, Coince A, Peel N, Percival-Alwyn L,

Duncan O, Trösch J, Yu G, Bolser DM, Namaati G, Kerhornou A, Spannagl M,

Gundlach H, Haberer G, Davey RP, Fosker C, Palma FD, Phillips A, Millar AH,

Kersey PJ, Uauy C, Krasileva KV, Swarbreck D, Bevan MW, Clark MD. 2017. An

improved assembly and annotation of the allohexaploid wheat genome identifies

complete families of agronomic genes and provides genomic evidence for chromosomal

translocations. *Genome Research* doi: 10.1101/gr.217117.116

Comai L. 2005. The advantages and disadvantages of being polyploid. *Nature Review Genetics*

6: 836–846.

Degreef JD. 1997. Fossil *Aldrovanda*. *Carnivorous Plant Newsletter* **26:** 93–97. Available at:

http://www.carnivorousplants.org/cpn/articles/CPNv26n3p93_97.pdf%5Cnpapers3://public

ation/uuid/A61A0327-61CF-4683-BF3F-EE5AFFD69F49.

Dodsworth S, Chase M, Leitch A. 2016. Is post-polyploidization diploidization the key to the

evolutionary success of angiosperms. *Botanical Journal of the Linnean Society* **180:**

1095–8339.

Dohm JC, Lange C, Holtgräwe D, Sörensen TR, Borchardt D, Schulz B, Lehrach H,

Weisshaar B, Himmelbauer H. 2012. Palaeohexaploid ancestry for Caryophyllales

inferred from extensive gene‑based physical and genetic mapping of the sugar beet

genome (Beta vulgaris). *The Plant Journal* **70:** 528–40.

Donoghue MJ. 2005. Key innovations, convergence, and success: macroevolutionary lessons

from plant phylogeny. *Paleobiology* **31:**77-93.

Donoghue MJ, Sanderson MJ. 2015. Confluence, synnovation, and depauperons in plant

diversification. *New Phyt.* **207:**260-274.

Douglas N, Spellenberg R. 2010. A new tribal classification of Nyctaginaceae. *Taxon* **59:** 905–

910.

Doyle JJ, Doyle JL. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf

tissue. *Phytochemical Bulletin* **19**: 11-15.

Edger PP, Heidel-Fischer HM, Bekaert M, Rota J, Glöckner G, Platts AE, Heckel DG, Der

JP, Wafula EK, Tang M, Hofberger JA. 2015. The butterfly plant arms-race escalated

by gene and genome duplications. *Proceedings of the National Academy of Sciences,*

USA **112:** 8362–8366.

Friis EM, Crane P, Pedersen KR. 2011. Early Flowers and Angiosperm Evolution. Cambridge.

Guindon S, Dufayard J, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New

algorithms and methods to estimate maximum-likelihood phylogenies: assessing the

performance of PhyML 3.0. *Systematic Biology* **59:** 307–321.

Huang, CH, Zhang C, Liu M, Hu Y, Gao T, Qi J, Hong Ma. 2016. Multiple Polyploidization

Events across Asteraceae with Two Nested Events in the Early History Revealed by

Nuclear Phylogenomics. *Mol Biol Evol.* **33:** 2820-2835.

Jordan GJ, Macphail MK. 2003. A middle-late Eocene inflorescence of Caryophyllaceae from

Tasmania, Australia. *American Journal of Botany* **90:** 761–768.

Levin DA. 1983. Polyploidy and novelty in flowering plants. *American Naturalist* **122:** 1–25.

Levin DA. 2002. The Role of Chromosomal Change in Plant Evolution. Oxford University

Press: New York.

McKain MR, Tang H, McNeal JR, Ayyampalayam S, Davis JI, dePamphilis CW, Givnish

TJ, Pires JC, Stevenson DW, Leebens-Mack JH. 2016. A Phylogenomic Assessment

of Ancient Polyploidy and Genome Evolution across the Poales. *Genome Biol. Evol.* **8**:

1150-1164.

Mayrose I, Zhan SH, Rothfels CJ, Magnuson-Ford K, Barker MS, Rieseberg LH, Otto SP.

2011. Recently formed polyploid plants diversify at lower rates. *Science* **333:** 1257.

Miller JS, Venable DL. 2000. Polyploidy and the evolution of gender dimorphism in plants.

Science **289:** 2335–2338.

Moore MJ, Soltis PS, Bell CD, Burleigh JG, Soltis DE. 2010. Phylogenetic analysis of 83
plastid genes further resolves the early diversification of eudicots. *Proceedings of the*
National Academy of Sciences, USA **107:** 4623–4628.

Nei M, Gojobori T. 1986. Simple methods for estimating the numbers of synonymous and
nonsynonymous nucleotide substitutions. *Molecular Biology and Evolution* **3:** 418–426.

Nichols DJ, Traverse A. 1971. Palynology, petrology, and depositional environments of some
early Tertiary lignites in Texas. *Geoscience and Man* **3:** 37–48.

Ocampo G, Columbus T. 2010. Molecular phylogenetics of suborder Cactineae
(Caryophyllales), including insights into photosynthetic diversification and historical
biogeography. *Am. J. Bot.* **97**: 1827-1847.

Otto SP, Whitton J. 2000. Polyploid incidence and evolution. *Annual Review of Genetics* **34:**
401–437.

Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, Fitzjohn RG, Alfaro ME,
Harmon LJ. 2014. geiger v2.0: an expanded suite of methods for fitting
macroevolutionary models to phylogenetic trees. *Bioinformatics* **30:** 2216–2218.

Puttick, MN, Clark J., Donoghue P. 2015. Size is not everything: rates of genome size
evolution, not C-value, correlate with speciation in angiosperms. *Proc. Roy. Soc. B.* **282**:
20152289.

Sanderson MJ. 2003. r8s: inferring absolute rates of molecular evolution and divergence times
in the absence of a molecular clock. *Bioinformatics* 19: 301–302.

Schubert I, Vu GTH. 2016. Genome Stability and Evolution: Attempting a Holistic View.
Trends in Plant Sciences 21: 749-757.

Schuster TM, Setaro SD, Kron KA. 2013. Age Estimates for the Buckwheat Family
Polygonaceae Based on Sequence Data Calibrated by Fossils and with a Focus on the
Amphi-Pacific *Muehlenbeckia*. *Plos ONE* **8**(4): e61261.

Smith SA, Beaulieu JM, Donoghue MJ. 2009. Mega-phylogeny approach for comparative
biology: an alternative to supertree and supermatrix approaches. *BMC Evolutionary*
Biology **9:** 37.

Smith SA, Beaulieu JM, Stamatakis A, Donoghue MJ. Understanding angiosperm
diversification using small and large phylogenetic trees. *American Journal of Botany* **98:**
404–414.

Smith SA, Donoghue MJ. 2008. Rates of molecular evolution are linked to life history in
flowering plants. *Science* 322: 86–89.

Smith SA, O'Meara BC. 2012. treePL: divergence time estimation using penalized likelihood
for large phylogenies. *Bioinformatics* **28:** 2689–2690.

Smith SA, Moore MJ, Brown JW, Yang Y. 2015. Analysis of phylogenomic datasets reveals
conflict, concordance, and gene duplications with examples from animals and plants.
BMC Evolutionary Biology **15:** 150.

Soltis PS, Soltis DS. 2000. The role of genetic and genomic attributes in the success of
polyploids. *Proceedings of the National Academy of Sciences, USA* **97:** 7051–7057.

Soltis DE, Smith SA, Cellinese N, Wurdack KJ, Tank DC, Brockington SF, Refulio-
Rodriguez NF, Walker JB, Moore MJ, Carlsward BS, Bell CD. 2011. Angiosperm
phylogeny: 17 genes, 640 taxa. *American Journal of Botany* **98:** 704–730.

Soltis PS, Liu X, Marchant DB, Visger CJ, Soltis DE. 2014. Polyploidy and novelty:
Gottlieb's legacy. *Philosophical Transactions of the Royal Society of London B:*
Biological Sciences **369:** 20130351.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* 30: 1312–1313.

Storchová Z, Breneman A, Cande J, Dunn J, Burbank K, O'toole E, Pellman D. 2006.
Genome-wide genetic analysis of polyploidy in yeast. *Nature* **443:** 541.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. Nucleic Acids Research **34 (suppl**
2): W609–W612.

Tank DC, Eastman JM, Pennell MW, Soltis PS, Soltis DE, Hinchliff CE, Brown JW, Sessa
EB, Harmon LJ. 2015. Nested radiations and the pulse of angiosperm diversification:
increased diversification rates often follow whole genome duplications. *New Phytologist*
207: 454–467.

Thulin M, Moore AJ, El-Seedi H, Larsson A, Christin P-A, Edwards EJ. 2016. Phylogeny
and generic delimitation in Molluginaceae, new pigment data in Caryophyllales, and the
new family Corbichoniaceae. *Taxon* **65:** 775–793.

Valente LM, Britton AW, Powell MP, Papadopulos AST, Burgoyne PM, Savolainen V.

2014. Correlates of hyperdiversity in southern African ice plants (Aizoaceae). *Botanical*

Journal of the Linnean Society Linnean Society of London. **174**:110-129.

Walker JF, Yang Y, Moore MJ, Mikenas J, Timoneda A, Brockington SF, Smith SA. 2017.

Widespread paleopolyploidy, gene tree conflict, and recalcitrant relationships among the

carnivorous Caryophyllales. *bioRxiv*: 115741.

Weiss H, Dobes C, Schneeweiss GM, Greimler J. 2002. Occurrence of tetraploid and

hexaploid cytotypes between and within populations in *Dianthus* sect. *Plumaria*

(Caryophyllaceae). *New Phyto.* **156:** 85-94.

Wood TE, Takebayashi N, Barker MS, Mayrose I, Greenspoon PB, Rieseberg LH. 2009.

The frequency of polyploid speciation in vascular plants. *Proceedings of the National*

Academy of Sciences, USA **106:** 13875–13879.

Yang Y, Moore MJ, Brockington SF, Soltis DE, Wong GK-S, Carpenter EJ, Zhang Y,

Chen L, Yan Z, Xie Y, Sage RF. 2015. Dissecting molecular evolution in the highly

diverse plant clade Caryophyllales using transcriptome sequencing. *Molecular Biology*

and Evolution **32:** 2001–2014.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and*

Evolution **24:** 1586–1591.

Zetter R, Hofmann CC, Draxler I, Durango de Cabrera J, Del MVergel M, Vervoorst F.

1999. A rich middle Eocene microflora at Arroyo de los Mineros, near Cañadón Beta, NE

Tierra del Fuego province, Argentina. *Abhandlungen der Geologischen Bundesanstalt*

56: 439–460.

Tables

Table 1 Voucher information and GenBank accession numbers for newly reported plastid *matK*

sequences. Families follow APG IV (Angiosperm Phylogeny Group 2016).

Family	Taxon	Voucher specimen (Herbarium acronym)	Collection locality	NCBI accession number
Achatocarpaceae	*Achatocarpus gracilis* H.Walter	Silvia H. Salas Morales et al. 5608 (TEX)	Mexico: Oaxaca	KY952292
Achatocarpaceae	*Phaulothamnus spinescens* A.Gray	Michael J. Moore et al. 976 (OC)	United States: Texas	KY952477
Achatocarpaceae	*Phaulothamnus spinescens* A.Gray	William R. Carr 27176 (TEX)	United States: Texas	KY952478
Amaranthaceae	*Allenrolfea occidentalis* (S.Watson) Kuntze	Michael J. Moore 474 (OC)	United States: Texas	KY952314
Amaranthaceae	*Alternanthera caracasana* Kunth	Michael J. Moore 1808 (OC)	United States: Texas	KY952319
Amaranthaceae	*Amaranthus cruentus* L.	Michael J. Moore 356 (OC)	United States: Ohio (cultivated)	KY952320
Amaranthaceae	*Amaranthus* sp.	Michael J. Moore 1801 (OC)	United States: Texas	KY952321
Amaranthaceae	*Amaranthus* sp.	Michael J. Moore 2186 (OC)	United States: Ohio	KY952322
Amaranthaceae	*Amaranthus* sp.	Michael J. Moore 2187 (OC)	United States: Illinois	KY952323
Amaranthaceae	*Atriplex prosopidum* I.M.Johnst.	Hilda Flores Olvera et al. 1658 (MEXU)	Mexico: Coahuila	KY952340
Amaranthaceae	*Atriplex* sp.	Michael J. Moore 1689 (OC)	United States: Texas	KY952338
Amaranthaceae	*Atriplex* sp.	Michael J. Moore 1699 (OC)	United States: Texas	KY952339
Amaranthaceae	*Celosia argentea* L. var. *plumosa*	Michael J. Moore 359 (OC)	United States: Ohio (cultivated)	KY952359
Amaranthaceae	*Charpentiera ovata* Gaudich. var. *ovata*	Flora K. Samis 7 (Lyon Arboretum living collection, accession 2011.0034)	United States: Hawaii	KY952360
Amaranthaceae	*Charpentiera tomentosa* Sohmer var. *maakuaensis* Sohmer	Flora K. Samis 6 (Lyon Arboretum living collection, accession 88.0141)	United States: Hawaii	KY952361
Amaranthaceae	*Chenopodium album* L.	Michael J. Moore 344 (OC)	United States: Ohio	KY952362
Amaranthaceae	*Gossypianthus lanuginosus* (Poir.) Moq.	Michael J. Moore 1807 (OC)	United States: Texas	KY952408
Amaranthaceae	*Guilleminea densa*	Michael J. Moore et	Mexico:	KY952412

	(Humb. & Bonpl. ex Schult.) Moq.	al. 2445 (OC)	Chihuahua	
Amaranthaceae	*Kali tragus* (L.) Scop.	Michael J. Moore 453 (OC)	United States: Texas	KY952506
Amaranthaceae	*Nototrichium divaricatum* D.H.Lorence	Flora K. Samis 3 (Lyon Arboretum living collection, accession 96.0036 #3)	United States: Hawaii	KY952468
Amaranthaceae	*Nototrichium humile* Hillebr.	Flora K. Samis 2 (Lyon Arboretum living collection, accession 2001-0254)	United States: Hawaii	KY952469
Amaranthaceae	*Suaeda jacoensis* I.M.Johnst.	Hilda Flores Olvera et al. 1662 (MEXU)	Mexico: Coahuila	KY952514
Amaranthaceae	*Suaeda jacoensis* I.M.Johnst.	Michael J. Moore et al. 2617 (OC)	Mexico: Nuevo Leon	KY952515
Amaranthaceae	*Suaeda mexicana* (Standl.) Standl.	Hilda Flores Olvera et al. 1654 (MEXU)	Mexico: Coahuila	KY952516
Amaranthaceae	*Tidestromia lanuginosa* (Nutt.) Standl.	Michael J. Moore 1128 (OC)	United States: Texas	KY952521
Amaranthaceae	*Zuckia brandegeei* (A.Gray) S.L.Welsh & Stutz var. *plummeri* (Stutz & S.C.Sand.) Dorn	Joseph L. M. Charboneau 9672 (RM)	United States: Colorado	KY952528
Cactaceae	*Leuenbergeria quisqueyana* (Alain) Lodé	Flora K. Samis 11 (Lyon Arboretum living collection, accession 2000.0281)	United States: Hawaii	KY952473
Caryophyllaceae	*Moehringia macrophylla* (Hook.) Fenzl	Arianna Goodman 1 (OC)	United States: Oregon	KY952464
Caryophyllaceae	*Paronychia lundellorum* Torr. & A.Gray	William R. Carr 17607 (MEXU)	United States: Texas	KY952472
Caryophyllaceae	*Saponaria officinalis* L.	Michael J. Moore et al. 1819 (OC)	United States: Indiana	KY952507
Caryophyllaceae	*Schiedea kaalae* Wawra	Flora K. Samis 5 (Lyon Arboretum living collection, accession 92.0513)	United States: Hawaii	KY952509
Caryophyllaceae	*Spergularia salina* J.Presl & C.Presl	Michael J. Moore 1693 (OC)	United States: Texas	KY952512
Didiereaceae	*Alluaudia ascendens* (Drake) Drake	Michael J. Moore 1645	United States (cultivated)	KY952318
Dioncophyllaceae	*Triphyophyllum peltatum* (Hutch. & Dalziel) Airy Shaw	Carel C. H. Jongkind et al. 7136 (WAG)	Liberia	KY952524
Droseraceae	*Drosera burmannii* Vahl cv. Pilliga Red	Michael J. Moore 1814 (OC)	United States (cultivated)	KY952400
Droseraceae	*Drosera peltata* Thunb.	Michael J. Moore	Australia:	KY952401

		1817 (OC)	Tasmania (cultivated)	
Droseraceae	*Drosera regia* Stephens	Michael J. Moore 1812 (OC)	United States (cultivated)	KY952402
Drosophyllaceae	*Drosophyllum lusitanicum* (L.) Link	Michael J. Moore 1816 (OC)	United States (cultivated)	KY952403
Frankeniaceae	*Frankenia gypsophila* I.M.Johnst.	Michael J. Moore et al. 1880 (OC)	Mexico: Nuevo Leon	KY952406
Microteaceae	*Microtea debilis* Sw.	Manuel Rimachi 11128 (TEX)	Peru: Loreto	KY952415
Montiaceae	*Claytonia sibirica* L.	Arianna Goodman 2 (OC)	United States: Oregon	KY952363
Montiaceae	*Phemeranthus parviflorus* (Nutt.) Kiger	Michael J. Moore et al. 2214 (OC)	United States: New Mexico	KY952479
Nyctaginaceae	*Abronia angustifolia* Greene	Michael J. Moore et al. 2063 (OC)	Mexico: Coahuila	KY952281
Nyctaginaceae	*Abronia angustifolia* Greene	Michael J. Moore et al. 896 (OC)	United States: New Mexico	KY952282
Nyctaginaceae	*Abronia bigelovii* Heimerl	Michael J. Moore et al. 704 (OC)	United States: New Mexico	KY952283
Nyctaginaceae	*Abronia elliptica* A.Nelson	Norman A. Douglas 2039 (DUKE)	United States: Arizona	KY952284
Nyctaginaceae	*Abronia fragrans* Nutt. ex Hook.	Billie L. Turner 20-22 (SRSC)	United States: Texas	KY952285
Nyctaginaceae	*Abronia fragrans* Nutt. ex Hook.	Glenn Kroh et al. 3021 (TEX)	United States: Texas	KY952286
Nyctaginaceae	*Abronia macrocarpa* L.A.Galloway	Steve L. Orzell et al. 6492 (TEX)	United States: Texas	KY952287
Nyctaginaceae	*Abronia mellifera* Douglas ex Hook.	N. Elizabeth Saunders BP 19 (SIU)	United States: Wyoming	KY952288
Nyctaginaceae	*Abronia mellifera* Douglas ex Hook.	N. Elizabeth Saunders BP 20 (SIU)	United States: Wyoming	KY952289
Nyctaginaceae	*Abronia nana* S.Watson var. *nana*	Robert C. Sivinski et al. 3108 (NMC)	United States: Arizona	KY952290
Nyctaginaceae	*Abronia umbellata* Lam.	N. Elizabeth Saunders LU 45 (SIU)	United States: California	KY952291
Nyctaginaceae	*Acleisanthes acutifolia* Standl.	James Henrickson et al. 22916 (TEX)	Mexico: Coahuila	KY952293
Nyctaginaceae	*Acleisanthes angustifolia* (Torr.) R.A.Levin	Michael J. Moore 460 (OC)	United States: Texas	KY952294
Nyctaginaceae	*Acleisanthes* cf. *purpusiana* (Heimerl) R.A.Levin	James Henrickson 23026 (TEX)	Mexico: Coahuila	KY952309
Nyctaginaceae	*Acleisanthes chenopodioides* (A.Gray) R.A.Levin	Michael J. Moore et al. 733 (OC)	United States: Texas	KY952295

Nyctaginaceae	*Acleisanthes crassifolia* A.Gray	Michael J. Moore et al. 569 (OC)	United States: Texas	KY952296
Nyctaginaceae	*Acleisanthes diffusa* (A.Gray) R.A.Levin var. *diffusa*	Michael J. Moore et al. 624 (OC)	United States: Texas	KY952297
Nyctaginaceae	*Acleisanthes lanceolata* (Wooton) R.A.Levin var. *lanceolata*	Michael J. Moore et al. 870 (OC)	United States: New Mexico	KY952298
Nyctaginaceae	*Acleisanthes lanceolata* (Wooton) R.A.Levin var. *lanceolata*	Michael J. Moore et al. 903 (OC)	United States: Texas	KY952299
Nyctaginaceae	*Acleisanthes lanceolata* (Wooton) R.A.Levin var. *megaphylla* (B.A.Fowler & B.L.Turner) Spellenb. & J.Poole	Alfred T. Richardson 1666 (TEX)	Mexico: Chihuahua	KY952300
Nyctaginaceae	*Acleisanthes longiflora* A.Gray	Michael J. Moore 435 (OC)	United States: Texas	KY952301
Nyctaginaceae	*Acleisanthes longiflora* A.Gray	Michael J. Moore et al. 571 (OC)	United States: Texas	KY952302
Nyctaginaceae	*Acleisanthes nana* I.M.Johnst.	Jackie Smith et al. 798 (TEX)	Mexico: San Luis Potosi	KY952303
Nyctaginaceae	*Acleisanthes obtusa* (Choisy) Standl.	Michael J. Moore et al. 984 (OC)	United States: Texas	KY952304
Nyctaginaceae	*Acleisanthes palmeri* (Hemsley) R.A.Levin	George S. Hinton 28620 (TEX)	MexicoNuevo Leon	KY952305
Nyctaginaceae	*Acleisanthes parvifolia* (Torr.) R.A.Levin	Michael J. Moore 452 (OC)	United States: Texas	KY952306
Nyctaginaceae	*Acleisanthes purpusiana* (Heimerl) R.A.Levin	James Henrickson 22709 (TEX)	Mexico: Coahuila	KY952307
Nyctaginaceae	*Acleisanthes purpusiana* (Heimerl) R.A.Levin	Billie L. Turner 6205 (TEX)	Mexico: Coahuila	KY952308
Nyctaginaceae	*Acleisanthes undulata* (B.A.Fowler & B.L.Turner) R.A.Levin	James Henrickson 23195 (TEX)	Mexico: Coahuila	KY952310
Nyctaginaceae	*Acleisanthes wrightii* (A.Gray) Benth. & Hook.	Michael J. Moore et al. 620 (OC)	United States: Texas	KY952311
Nyctaginaceae	*Allionia choisyi* Standl.	Norman A. Douglas 2187 (DUKE)	Mexico: Coahuila	KY952315
Nyctaginaceae	*Allionia incarnata* L.	Michael J. Moore et al. 1352 (OC)	Mexico: Nuevo Leon	KY952316
Nyctaginaceae	*Allionia* sp.	Michael J. Moore 424 (OC)	United States: Texas	KY952317
Nyctaginaceae	*Andradea floribunda* Allemão	André M. Amorim 2294 (NY)	Brazil	KY952324
Nyctaginaceae	*Andradea floribunda* Allemão	Jacquelyn Ann Kallunki 701 (NY)	Brazil	KY952325
Nyctaginaceae	*Anulocaulis annulatus*	Richard W.	United States:	KY952326

	(Coville) Standl.	Spellenberg 3162 (NMC)	California	
Nyctaginaceae	*Anulocaulis eriosolenus* (A.Gray) Standl.	James Henrickson et al. 23103 (TEX)	Mexico: Coahuila	KY952327
Nyctaginaceae	*Anulocaulis eriosolenus* (A.Gray) Standl.	Michael J. Moore et al. 611 (OC)	United States: Texas	KY952328
Nyctaginaceae	*Anulocaulis hintoniorum* B.L.Turner	Patricia Hernández Ledesma 52 (MEXU)	Mexico: Coahuila	KY952329
Nyctaginaceae	*Anulocaulis leiosolenus* (Torr.) Standl. var. *gypsogenus* (Waterf.) Spellenb. & T.Wootten	Michael J. Moore 402 (OC)	United States: New Mexico	KY952330
Nyctaginaceae	*Anulocaulis leiosolenus* (Torr.) Standl. var. *howardii* Spellenb. & T.Wootten	Thomas Wootten et al. s.n. (NMC)	United States: New Mexico	KY952331
Nyctaginaceae	*Anulocaulis leiosolenus* (Torr.) Standl. var. *lasianthus* I.M.Johnston	Michael J. Moore et al. 610 (OC)	United States: Texas	KY952332
Nyctaginaceae	*Anulocaulis leiosolenus* (Torr.) Standl. var. *leiosolenus*	Michael J. Moore 493 (OC)	United States: Texas	KY952333
Nyctaginaceae	*Anulocaulis leiosolenus* (Torr.) Standl. var. *leiosolenus*	Michael J. Moore et al. 825 (OC)	United States: Nevada	KY952334
Nyctaginaceae	*Anulocaulis leiosolenus* (Torr.) Standl. var. *leiosolenus*	Michael J. Moore et al. 853 (OC)	United States: Arizona	KY952335
Nyctaginaceae	*Anulocaulis reflexus* I.M.Johnst.	Michael J. Moore et al. 242 (TEX)	Mexico: Chihuahua	KY952336
Nyctaginaceae	*Anulocaulis reflexus* I.M.Johnst.	Michael J. Moore 483 (OC)	United States: Texas	KY952337
Nyctaginaceae	*Boerhavia anisophylla* Torr.	Norman A. Douglas 2194 (DUKE)	Mexico: Durango	KY952341
Nyctaginaceae	*Boerhavia ciliata* Brandegee	Norman A. Douglas 2145 (DUKE)	United States: Texas	KY952342
Nyctaginaceae	*Boerhavia coccinea* Mill.	Michael J. Moore 366 (OC)	United States: New Mexico	KY952343
Nyctaginaceae	*Boerhavia coulteri* (Hook.f.) S.Watson var. *palmeri* (S.Watson) Spellenb.	Richard W. Spellenberg 13273 (NMC)	United States: Arizona	KY952344
Nyctaginaceae	*Boerhavia dominii* Meikle & Hewson	H. Smyth 42 (NY)	Australia: South Australia	KY952345
Nyctaginaceae	*Boerhavia gracillima* Heimerl	Richard W. Spellenberg 12447 (NMC)	United States: Texas	KY952347
Nyctaginaceae	*Boerhavia intermedia* M.E.Jones	Richard W. Spellenberg 13279	United States: Arizona	KY952348

		(NMC)		
Nyctaginaceae	*Boerhavia lateriflora* Standl.	Norman A. Douglas 2161 (DUKE)	Mexico: Sonora	KY952349
Nyctaginaceae	*Boerhavia linearifolia* A.Gray	Michael J. Moore et al. 581 (OC)	United States: Texas	KY952350
Nyctaginaceae	*Boerhavia purpurascens* A.Gray	Richard W. Spellenberg 13261 (NMC)	United States: Arizona	KY952351
Nyctaginaceae	*Boerhavia repens* L.	J. S. Rose 2	United States: Hawaii	KY952352
Nyctaginaceae	*Boerhavia repens* L.	Richard W. Spellenberg 7183 (NMC)	Yemen: Sana	KY952353
Nyctaginaceae	*Boerhavia* sp.	Erin Tripp et al. 4090 (OC)	Namibia	KY952346
Nyctaginaceae	*Boerhavia torreyana* (S.Watson) Standl.	Michael J. Moore et al. 633 (OC)	United States: Texas	KY952354
Nyctaginaceae	*Bougainvillea campanulata* Heimerl	Michael Nee 51257 (TEX)	Bolivia: Santa Cruz	KY952355
Nyctaginaceae	*Bougainvillea glabra* Choisy	Michael J. Moore 538 (OC)	United States: Ohio (cultivated)	KY952356
Nyctaginaceae	*Bougainvillea spinosa* (Cav.) Heimerl	J. Saunders et al. 3371 (TEX)	Argentina: San Juan	KY952357
Nyctaginaceae	*Bougainvillea stipitata* Griseb.	Michael Nee 50723 (TEX)	Bolivia: Santa Cruz	KY952358
Nyctaginaceae	*Colignonia glomerata* Griseb.	Michael Nee 52523 (NY)	Bolivia	KY952364
Nyctaginaceae	*Colignonia scandens* Benth.	Martin Grantham 63 (SFBG living collection, accession 1996-0202)	Ecuador	KY952365
Nyctaginaceae	*Commicarpus ambiguus* Meikle	Mats Thulin 11015 (UPS)	Somalia: Sanaag	KY952366
Nyctaginaceae	*Commicarpus arabicus* Meikle	Mats Thulin et al. 9294 (UPS)	Yemen: Taizz	KY952367
Nyctaginaceae	*Commicarpus arabicus* Meikle	Richard W. Spellenberg 7217 (NMC)	Yemen: Ibb	KY952368
Nyctaginaceae	*Commicarpus arabicus* Meikle	Richard W. Spellenberg 7297 (NMC)	Yemen: Ibb	KY952369
Nyctaginaceae	*Commicarpus australis* (Meikle) Govaerts	Richard W. Spellenberg et al. 9469 (NMC)	Australia: Western Australia	KY952370
Nyctaginaceae	*Commicarpus boissieri* (Heimerl) Cufod.	Mats Thulin 11423 (UPS)	Oman: Dhofar	KY952371
Nyctaginaceae	*Commicarpus boissieri* (Heimerl) Cufod.	Carl J. Rothfels et al. 4331	Oman: Ash Sharqiyah	KY952373
Nyctaginaceae	*Commicarpus*	Patricia Hernández	Mexico: Baja	KY952372

	brandegeei Standl.	Ledesma 55 (MEXU)	California Sur	
Nyctaginaceae	*Commicarpus coctoris* N.A.Harriman	Richard W. Spellenberg et al. 12883 (NMC)	Mexico: Oaxaca	KY952374
Nyctaginaceae	*Commicarpus commersonii* (Baill.) Cavaco	Mats Thulin et al. 11836 (UPS)	Madagascar: Toliara	KY952380
Nyctaginaceae	*Commicarpus decipiens* Meikle	Erin Tripp et al. 4127 (NMC)	Namibia	KY952375
Nyctaginaceae	*Commicarpus grandiflorus* (A.Rich.) Standl.	Mats Thulin et al. 9311 (UPS)	Yemen: Taizz	KY952376
Nyctaginaceae	*Commicarpus greenwayi* Meikle	Mats Thulin 606 (UPS)	Tanzania: Iringa	KY952377
Nyctaginaceae	*Commicarpus helenae* (Roem. & Schult.) Meikle	Richard W. Spellenberg et al. 7504 (NMC)	Yemen: Dhamar	KY952378
Nyctaginaceae	*Commicarpus hiranensis* Thulin	Mats Thulin et al. 11225 (UPS)	Ethiopia: Harerge	KY952379
Nyctaginaceae	*Commicarpus mistus* Thulin	Mats Thulin et al. 9786 (UPS)	Yemen: Mahrah	KY952381
Nyctaginaceae	*Commicarpus parviflorus* Thulin	Mats Thulin 6318 (UPS)	Somalia: Banaadir	KY952382
Nyctaginaceae	*Commicarpus pedunculosus* (A.Rich.) Cufod.	Mats Thulin 1301 (UPS)	Ethiopia: Arussi	KY952383
Nyctaginaceae	*Commicarpus plumbagineus* (Cav.) Standl.	Mats Thulin 10747 (UPS)	Somalia: Togdheer	KY952384
Nyctaginaceae	*Commicarpus plumbagineus* (Cav.) Standl.	Mats Thulin et al. 11330 (UPS)	Ethiopia: Harerge	KY952385
Nyctaginaceae	*Commicarpus plumbagineus* (Cav.) Standl.	Richard W. Spellenberg et al. 7374 (NMC)	Yemen: Ta'izz	KY952386
Nyctaginaceae	*Commicarpus praetermissus* N.A.Harriman	Richard W. Spellenberg et al. 12905 (NMC)	Mexico: Michoacán	KY952387
Nyctaginaceae	*Commicarpus reniformis* (Chiov.) Cufod.	Mats Thulin 4200 (UPS)	Somalia: Sool	KY952388
Nyctaginaceae	*Commicarpus reniformis* (Chiov.) Cufod.	Mats Thulin et al. 8337 (UPS)	Yemen: Hadramaut	KY952389
Nyctaginaceae	*Commicarpus scandens* (L.) Standl.	Michael J. Moore 1127 (OC)	United States: Texas	KY952390
Nyctaginaceae	*Commicarpus scandens* (L.) Standl.	Richard W. Spellenberg et al. 12887 (NMC)	Mexico: Puebla	KY952391
Nyctaginaceae	*Commicarpus sinuatus* Meikle	Mats Thulin 10737 (UPS)	Somalia: Woqooyi Galbeed	KY952392

Nyctaginaceae	*Commicarpus sinuatus* Meikle	Richard W. Spellenberg 7144 (NMC)	Yemen: Sana'a	KY952393
Nyctaginaceae	*Commicarpus sinuatus* Meikle	Richard W. Spellenberg 7506 (NMC)	Yemen: Dhamar	KY952394
Nyctaginaceae	*Commicarpus squarrosus* (Heimerl) Standl. var. *squarrosus*	Erin Tripp et al. 4049 (NMC)	Namibia	KY952395
Nyctaginaceae	*Commicarpus stenocarpus* (Chiov.) Cufod.	Mats Thulin et al. 8062 (UPS)	Yemen: Hadramaut	KY952396
Nyctaginaceae	*Cuscatlania vulcanicola* Standl.	José L. Linares 12938 (MEXU)	El Salvador: Sonsonate	KY952397
Nyctaginaceae	*Cuscatlania vulcanicola* Standl.	José L. Linares 13440 (MEXU)	El Salvador: Sonsonate	KY952398
Nyctaginaceae	*Cyphomeris gypsophiloides* (M.Martens & Galeotti) Standl.	Michael J. Moore et al. 582 (OC)	United States: Texas	KY952399
Nyctaginaceae	*Grajalesia fasciculata* (Standl.) Mirand*a*	José L. Linares 13416 (MEXU)	El Salvador: Sonsonate	KY952409
Nyctaginaceae	*Guapira discolor* (Spreng.) Little	Richard W. Spellenberg 13294 (NMC)	United States: Florida	KY952410
Nyctaginaceae	*Guapira eggersiana* (Heimerl) Lundell	Scott A. Mori 25542/40 (NY)	French Guiana	KY952411
Nyctaginaceae	*Mirabilis albida* (Walter) Heimerl	Norman A. Douglas 2035 (DUKE)	United States: Arizona	KY952416
Nyctaginaceae	*Mirabilis albida* (Walter) Heimerl	William R. Carr 11075 (TEX)	United States: Texas	KY952417
Nyctaginaceae	*Mirabilis alipes* (S.Watson) Pilz	Arnold Tiehm 13461 (TEX)	United States: Nevada	KY952418
Nyctaginaceae	*Mirabilis bigelovii* A.Gray var. *retrorsa* (A. Heller) Munz	James D. Morefield et al. 3780 (TEX)	United States: California	KY952419
Nyctaginaceae	*Mirabilis* cf. *glabrifolia* (Ortega) I.M.Johnst.	Michael J. Moore et al. 1244 (OC)	Mexico: San Luis Potosi	KY952428
Nyctaginaceae	*Mirabilis* cf. *nesomii* B.L.Turner	George S. Hinton 25567 (TEX)	Mexico: Nuevo Leon	KY952449
Nyctaginaceae	*Mirabilis coccinea* (Torr.) Benth. & Hook.f.	Norman A. Douglas 2133 (DUKE)	United States: Arizona	KY952420
Nyctaginaceae	*Mirabilis coccinea* (Torr.) Benth. & Hook.f.	Steven P. McLaughlin et al. 9354 (ARIZ)	United States: Arizona	KY952421
Nyctaginaceae	*Mirabilis comata* (Small) Standl.	Norman A. Douglas 2084 (DUKE)	United States: Arizona	KY952422
Nyctaginaceae	*Mirabilis decumbens* (Nutt.) Daniels	Richard W. Spellenberg et al. 4073 (TEX)	Mexico: Zacatecas	KY952423

Nyctaginaceae	*Mirabilis donahooiana* Le Duc	Alice Le Duc et al. 247 (TEX)	Mexico: Michoacán	KY952424
Nyctaginaceae	*Mirabilis exserta* Brandegee	Pedro Tenorio 10586 (MEXU)	Mexico	KY952425
Nyctaginaceae	*Mirabilis gigantea* (Standl.) Shinners	J. Quayle et al. 752 (TEX)	United States: Texas	KY952426
Nyctaginaceae	*Mirabilis glabra* (S.Watson) Standl.	Michael J. Moore et al. 674 (OC)	United States: New Mexico	KY952446
Nyctaginaceae	*Mirabilis glabrifolia* (Ortega) I.M.Johnst.	Guy Nesom et al. 7654 (TEX)	Mexico: Coahuila	KY952427
Nyctaginaceae	*Mirabilis glabrifolia* (Ortega) I.M.Johnst.	Michael J. Moore et al. 1325 (OC)	Mexico: Nuevo Leon	KY952429
Nyctaginaceae	*Mirabilis gracilis* (Standl.) LeDuc	Alice Le Duc et al. 71 (TEX)	Mexico: Jalisco	KY952430
Nyctaginaceae	*Mirabilis grandiflora* (Standl.) Standl.	EDL 1863 (MEXU)	Mexico	KY952431
Nyctaginaceae	*Mirabilis greenei* S.Watson	George E. Pilz 998 (TEX)	United States: California	KY952432
Nyctaginaceae	*Mirabilis himalaica* (Edgew.) Heimerl var. *chinensis* Heimerl	D. E. Boufford et al. 32449 (F)	China: Xizang (Tibet)	KY952433
Nyctaginaceae	*Mirabilis himalaica* (Edgew.) Heimerl var. *chinensis* Heimerl	D. E. Boufford et al. 41198 (F)	China: Xizang (Tibet)	KY952434
Nyctaginaceae	*Mirabilis himalaica* (Edgew.) Heimerl var. *chinensis* Heimerl	D. E. Boufford et al. 41435 (F)	China: Xizang (Tibet)	KY952435
Nyctaginaceae	*Mirabilis hintoniorum* Le Duc	Patricia Hernández Ledesma 118 (MEXU)	Mexico: Michoacán	KY952436
Nyctaginaceae	*Mirabilis jalapa* L.	Michael J. Moore s.n.	United States (cultivated)	KY952437
Nyctaginaceae	*Mirabilis laevis* (Benth.) Curran	Andrew C. Sanders et al. 29410 (TEX)	United States: California	KY952438
Nyctaginaceae	*Mirabilis latifolia* (A.Gray) Diggs, Lipscomb & O'Kennon	Victor L. Cory 24549 (GH)	United States: Texas	KY952439
Nyctaginaceae	*Mirabilis linearis* (Pursh) Heimerl	Billie L. Turner 21-854 (TEX)	United States: Texas	KY952440
Nyctaginaceae	*Mirabilis linearis* (Pursh) Heimerl var. *decipiens* (Standl.) S.L.Welsh	Michael J. Moore et al. 1984 (OC)	Mexico: Coahuila	KY952441
Nyctaginaceae	*Mirabilis longiflora* L.	Michael J. Moore et al. 1230 (OC)	Mexico: San Luis Potosi	KY952442
Nyctaginaceae	*Mirabilis longiflora* L. var. *wrightiana* (A.Gray ex Britton & Kearney) Kearney & Peebles	Alice Le Duc 185 (TEX)	United States: New Mexico	KY952443
Nyctaginaceae	*Mirabilis melanotricha*	Michael J. Moore et	Mexico: San	KY952444

	(Standl.) Spellenb.	al. 1191 (OC)	Luis Potosi	
Nyctaginaceae	*Mirabilis melanotricha* (Standl.) Spellenb.	Norman A. Douglas 2067 (DUKE)	United States: New Mexico	KY952445
Nyctaginaceae	*Mirabilis multiflora* (Torr.) A.Gray	Michael J. Moore 1110 (OC)	United States: Texas	KY952447
Nyctaginaceae	*Mirabilis multiflora* (Torr.) A.Gray	Norman A. Douglas 2037 (DUKE)	United States: Arizona	KY952448
Nyctaginaceae	*Mirabilis nesomii* B.L.Turner	Michael J. Moore et al. 2179 (NMC)	Mexico: Nuevo Leon	KY952450
Nyctaginaceae	*Mirabilis nesomii* B.L.Turner	Michael J. Moore et al. 2643 (NMC)	Mexico: Nuevo Leon	KY952451
Nyctaginaceae	*Mirabilis nyctaginea* (Michx.) MacMill.	William R. Carr 14590 (TEX)	United States: Texas	KY952452
Nyctaginaceae	*Mirabilis oligantha* (Standl.) Standl.	José L. Panero 2816 (MEXU)	Mexico: Baja California	KY952453
Nyctaginaceae	*Mirabilis oxybaphoides* (A.Gray) A.Gray	George S. Hinton 25572 (TEX)	Mexico: Nuevo Leon	KY952454
Nyctaginaceae	*Mirabilis polonii* Le Duc	Alice Le Duc 259 (MEXU)	Mexico: Nuevo Leon	KY952455
Nyctaginaceae	*Mirabilis pringlei* Weath.	Alice Le Duc et al. 63 (TEX)	Mexico: Jalisco	KY952456
Nyctaginaceae	*Mirabilis pudica* Barneby	Arnold Tiehm 10971 (TEX)	United States: Nevada	KY952457
Nyctaginaceae	*Mirabilis texensis* (J.M.Coult.) B.L.Turner	Billie L. Turner 22-417 (TEX)	United States: Texas	KY952458
Nyctaginaceae	*Mirabilis triflora* Benth.	Ramón Cuevas G. et al. 3415 (MEXU)	Mexico: Jalisco	KY952459
Nyctaginaceae	*Mirabilis urbani* Heimerl	Mark Fishbein et al. 5107 (MEXU)	Mexico: Michoacan	KY952460
Nyctaginaceae	*Mirabilis violacea* (L.) Heimerl	Patricia Hernández Ledesma 63 (MEXU)	Mexico: Distrito Federal	KY952461
Nyctaginaceae	*Mirabilis viscosa* Cav.	Michael J. Moore et al. 1824 (NMC)	Mexico: San Luis Potosi	KY952462
Nyctaginaceae	*Mirabilis viscosa* Cav.	Patricia Hernández Ledesma 13 (MEXU)	Mexico	KY952463
Nyctaginaceae	*Neea belizensis* Donn.Sm.	Cyrus L. Lundell 17692 (TEX)	Guatemala: Petén	KY952465
Nyctaginaceae	*Neea cauliflora* Poepp. & Endl.	Schanke S15106 (NY)	Peru	KY952466
Nyctaginaceae	*Neea psychotrioides* Donn.Sm.	Robert L. Wilbur 63654	Costa Rica: Heredia	KY952467
Nyctaginaceae	*Nyctaginia capitata* Choisy	Michael J. Moore et al. 617 (OC)	United States: Texas	KY952470
Nyctaginaceae	*Okenia hypogaea* Schltdl. & Cham.	Thomas R. Van Devender et al. 92-1069 (NMC)	Mexico: Sonora	KY952471
Nyctaginaceae	*Pisonia aculeata* L.	C. Martínez 1209 (TEX)	Mexico: Oaxaca	KY952483
Nyctaginaceae	*Pisonia brunoniana*	J. S. Rose 3	United States:	KY952484

	Endl.		Hawaii	
Nyctaginaceae	*Pisonia capitata* (S.Watson) Standl.	Ana L. Reina Guerrero et al. 2000-193 (NMC)	Mexico: Sonora	KY952485
Nyctaginaceae	*Pisonia capitata* (S.Watson) Standl.	Thomas R. Van Devender et al. 2003-17 (TEX)	United States: Arizona	KY952486
Nyctaginaceae	*Pisonia macranthocarpa* (Donn.Sm.) Donn.Sm.	Dennis E. Breedlove et al. 30361 (TEX)	Mexico: Chiapas	KY952487
Nyctaginaceae	*Pisonia sandwicensis* Hillebr.	Flora K. Samis 1 (Lyon Arboretum living collection)	United States: Hawaii	KY952488
Nyctaginaceae	*Pisonia sylvatica* Standl.	José L. Linares 13403 (MEXU)	El Salvador: Sonsonate	KY952489
Nyctaginaceae	*Pisonia umbellifera* (J.R.Forst. & G.Forst.) Seem.	Flora K. Samis 12 (Lyon Arboretum living collection, accession 68.0453)	United States: Hawaii	KY952490
Nyctaginaceae	*Pisonia zapallo* Griseb.	Israel G. Vargas et al. 2001 (TEX)	Bolivia: Santa Cruz	KY952491
Nyctaginaceae	*Pisoniella arborescens* (Lag. & Rodr.) Standl.	Alice Le Duc et al. 231 (NMC)	Mexico: Oaxaca	KY952492
Nyctaginaceae	*Pisoniella arborescens* (Lag. & Rodr.) Standl.	William R. Anderson 13522 (NY)	Mexico: Oaxaca	KY952493
Nyctaginaceae	*Ramisia brasiliensis* Oliv.	Jomar G. Jardim 1507 (NY)	Brazil	KY952495
Nyctaginaceae	*Reichenbachia hirsuta* Spreng.	Michael Nee 47813 (NY)	Bolivia	KY952496
Nyctaginaceae	*Reichenbachia paraguayensis* (D.Parodi) Dugand & Daniel	Maria Maguidaura Hatschbach 49218 (NY)	Brazil	KY952497
Nyctaginaceae	*Salpianthus arenarius* Bonpl.	Richard W. Spellenberg 12903 (NMC)	Mexico: Michoacán	KY952503
Nyctaginaceae	*Salpianthus macrodontus* Standl.	Thomas R. Van Devender et al. 91-894 (NMC)	Mexico: Sonora	KY952504
Nyctaginaceae	*Salpianthus purpurascens* (Cav. ex Lag.) Hook. & Arn.	Richard W. Spellenberg et al. 12885 (NMC)	Mexico: Oaxaca	KY952505
Nyctaginaceae	*Tripterocalyx carneus* (Greene) L.A.Galloway	Norman A. Douglas 2060 (DUKE)	United States: New Mexico	KY952525
Nyctaginaceae	*Tripterocalyx crux-maltae* (Kellogg) Standl.	Arnold Tiehm et al. 12213 (TEX)	United States: Nevada	KY952526
Nyctaginaceae	*Tripterocalyx micranthus* (Torr.) Hook.	B. MacLeod et al. 751 (TEX)	United States: Colorado	KY952527
Phytolaccaceae	*Agdestis clematidea* Moc. & Sessé ex DC.	George S. Hinton 25023 (TEX)	Mexico: Tamaulipas	KY952313

Phytolaccaceae	*Gallesia integrifolia* (Spreng.) Harms	Michael Nee et al. 50072 (TEX)	Bolivia: Santa Cruz	KY952407
Phytolaccaceae	*Hilleria latifolia* (Lam.) H.Walter	Michael Nee 33807 (TEX)	Bolivia: Santa Cruz	KY952413
Phytolaccaceae	*Petiveria alliacea* L.	Lucas C. Majure 4132 (FLAS)	United States: Florida	KY952476
Phytolaccaceae	*Phytolacca americana* L.	Michael J. Moore 342 (OC)	United States: Ohio	KY952480
Phytolaccaceae	*Phytolacca icosandra* L.	Mark H. Mayfield et al. 1001 (TEX)	Mexico: Guerrero	KY952481
Phytolaccaceae	*Phytolacca octandra* L.	Juan A. Encina et al. 1545 (TEX)	Mexico: Nuevo Leon	KY952482
Phytolaccaceae	*Rivina humilis* L.	Michael J. Moore 1129 (OC)	United States: Texas	KY952499
Phytolaccaceae	*Seguieria aculeata* Jacq.	Elsa Zardini et al. 22101 (TEX)	Paraguay	KY952510
Phytolaccaceae	*Seguieria paraguariensis* Morong	Michael Nee 48735 (TEX)	Bolivia: Santa Cruz	KY952511
Phytolaccaceae	*Trichostigma octandrum* (L.) H.Walter	Michael Nee 47094 (TEX)	Bolivia: Santa Cruz	KY952522
Phytolaccaceae	*Trichostigma peruvianum* (Moq.) H.Walter	Flora K. Samis 10 (Lyon Arboretum living collection, accession 94.0377)	United States: Hawaii	KY952523
Plumbaginaceae	*Aegialitis annulata* R.Br.	Christopher T. Martine 4043 (OC)	Australia: Western Australia	KY952312
Plumbaginaceae	*Limonium limbatum* Small	Michael J. Moore et al. 694 (OC)	United States: New Mexico	KY952414
Plumbaginaceae	*Plumbago scandens* L.	Michael J. Moore et al. 1828 (OC)	Mexico: San Luis Potosi	KY952494
Polygonaceae	*Eriogonum longifolium* Nutt. var. *longifolium*	Michael J. Moore 1796 (OC)	United States: Texas	KY952404
Polygonaceae	*Eriogonum rotundifolium* Benth.	Michael J. Moore 1769 (OC)	United States: New Mexico	KY952405
Polygonaceae	*Persicaria odorata* LaLlave	Flora K. Samis 9 (Lyon Arboretum living collection, accession 88.0439)	United States: Hawaii	KY952475
Polygonaceae	*Persicaria* sp.	Michael J. Moore 1177	United States: Ohio	KY952474
Polygonaceae	*Reynoutria japonica* (Houtt.) Ronse Decr.	Michael J. Moore 2188 (OC)	United States: Ohio	KY952498
Polygonaceae	*Rumex albescens* Hillebr.	Flora K. Samis 4 (Lyon Arboretum living collection, accession 2008-0119)	United States: Hawaii	KY952500
Polygonaceae	*Rumex* sp.	Michael J. Moore 1800 (OC)	United States: Texas	KY952501
Polygonaceae	*Rumex* sp.	Michael J. Moore	United States:	KY952502

		1805 (OC)	Texas	
Sarcobataceae	*Sarcobatus vermiculatus* (Hook.) Torr.	Michael J. Moore et al. 813 (OC)	United States: Utah	KY952508
Stegnospermataceae	*Stegnosperma cubense* A.Rich.	Silvia H. Salas Morales 2649 (NY)	Mexico: Oaxaca	KY952513
Talinaceae	*Talinum* cf. *aurantiacum* Engelm.	Michael J. Moore et al. 1985 (OC)	Mexico: Coahuila	KY952517
Talinaceae	*Talinum fruticosum* (L.) Juss.	Flora K. Samis 8 (Lyon Arboretum living collection, accession 2012.0008)	United States: Hawaii	KY952518
Talinaceae	*Talinum paniculatum* (Jacq.) Gaertn.	Michael J. Moore 1789 (OC)	United States (cultivated)	KY952520
Talinaceae	*Talinum* sp.	Michael J. Moore et al. 1974 (MEXU)	Mexico: Coahuila	KY952519

667 **Table 2** List of primers used to amplify the *matK* sequences newly reported here. Within each

668 primer name, the number indicates the approximate position of the primer in nucleotides

669 downstream from the start of *matK*.

Primer name	Sequence (5'→3')	Notes
matK.300F.Car	TTG CAG TCA TTG TGG AAA TTC C	works broadly across most of Caryophyllales, but generally fails in Caryophyllaceae and Frankeniaceae
matK.1350R.Car	GCC AAA GTT CTA GCA CAA GAA AG	works broadly across most of Caryophyllales
matK.210F.Car	TTC GGC TAA TGA TTC TCA CCA A	designed specifically for Caryophyllaceae
matK.1345R.Car	GAG CCA AAG TTC TAG CAC AAG AA	designed specifically for Caryophyllaceae
matK.1355R.Car	TGT GTT TAC GAG CTA AAG TTC TAG	designed specifically for Caryophyllaceae
matK.300F.Fra	TCG CTG TCT TTG CTG AAA TTC C	designed specifically for Frankeniaceae

672 **Table 3** Summary of WGD events at identified clades with distance to diversification shift and

673 climate information. Numbers correspond to those in Figs. 1 and 2.

#	Putative WGD	Distance to diversification shift in nodes ML(BS)	Subtending species (sister)	Mean annual temp °C (sister)	Mean annual precip mm (sister)
1	Tribe Nyctagineae within the Nyctaginaceae	0 (0)	123 (40)	17.49 (20.08)	482.9 (997.08)
2	Phytolaccoid clade	6 (6)	182 (407)	19.64 (18.36)	1007.58 (452.47)

3	*Claytonia*	NA	38 (15)	5.28 (7.25)	790.5 (970.36)
4	Portulacineae	1 (1)	1600 (38)	16.19 (19.35)	699.87 (736.42)
5	*Amaranthus*	0 (0)	28 (1)	16.27 (27.09)	797.74 (117.63)
6	Tribe Gomphrenoideae within Amaranthaceae	7 (7)	172 (41)	17.91 (16.65)	871.95 (1289.5)
7	in Caryophyllaceae (Alsinoideae + Caryophylloideae sensu Greenberg and Donoghue 2011)	9 (9)	793 (13)	11.44 (12.06)	761.43 (720.00)
8	Polygonaceae	13 (13)	670 (70)	16.3. (16.89)	1084.17 (794.28)
9	Plumbaginaceae	NA	70 (670)	16.89 (16.3)	794.28 (1084.17)
10	Droseraceae	8 (NA)	67 (108)	16.3 (19.08)	1280.57 (1491.72)
11	Nepenthaceae	4 (NA)	89 (19)	22.52 (20.05)	2170.5 (1611.63)
12	Ancistrocladaceae	0 (NA)	15 (3)	24.17 (25.6)	1899.13 (2882.4)
13	Tamaricaceae	NA	19 (3)	14.09 (16.21)	568.32 (469.61)

676   **Table 4** Summary of diversification shifts. Letters correspond to those in Figs. 1 and 2.

#	Family	Diversification shift	Mean shift (ML)	Mean shift (BS)
a	Cactaceae	*Echinops*	1.7957	2.2008
b	Cactaceae	within *Gymnocalycium*	6.9152	
c	Cactaceae	*Gymnocalycium*	-0.001	0.0555
d	Cactaceae	*Hylocereus+Selenicereus*	0.1175	
e	Cactaceae	*Rhipsalis+Schlumbergera+Echinocereus*+relatives	0.0514	
f	Cactaceae	*Stenocactus*	-0.057	-0.019
g	Anacampserotaceae	*Anacampseros*	0.2624	
h	Portulacaceae	*Portulaca*	0.0427	0.0447
i	Montiaceae	*Montiopsis*	0.9418	
j	Montiaceae	Montiaceae	0.0325	
k	Aizoaceae	*Drosanthemum+Delosperma+Hereroa*+relatives	0.1469	
l	Nyctaginaceae	*Boerhavia*		0.0747
m	Nyctaginaceae	*Commicarpus*	0.9642	
n	Nyctaginaceae	Tribe Nyctagineae	0.0484	0.0485
o	Nyctaginaceae	*Abronia*		-0.084
p	Nyctag.+Aizo+Cact.+relatives	Nyctag.+Aizo+Cact.+relatives	0.0168	0.019
r	Amaranthaceae	*Salicornia*	0.2732	0.1649
s	Amaranthaceae	*Suaeda* clade 1	0.1027	
t	Amaranthaceae	*Suaeda* clade 2	-0.036	-0.028
u	Amaranthaceae	*Atriplex*	0.0384	
v	Amaranthaceae	*Corispermum*	0.1186	
w	Amaranthaceae	*Froelichia+Gomphrena*+relatives	0.0217	0.0132
x	Amaranthaceae	*Amaranthus*	0.335	0.2049
y	Caryophyllaceae	*Dianthus*	0.0662	0.0409
z	Caryophyllaceae	*Cerastium*	0.7137	
aa	Caryophyllaceae	*Arenaria*	0.4606	0.425
bb	Caryophyllaceae	*Moehringia*	1.0971	0.995
cc	Caryophyllaceae	*Schiedea*	0.2339	0.2767
dd	Polygonaceae	*Fagopyrum*	-0.04	-0.034
ee	Polygonaceae	*Eriogonum*+relatives	0.0432	0.0364
ff	Nepenthaceae	within *Nepenthes*	0.042	
gg	Ancistrocladaceae	*Ancistrocladus*	0.1426	
hh	Droseraceae	within *Drosera 1*	0.2237	0.2076
ii	Droseraceae	within *Drosera 2*		0.1622

Figures

Fig. 1 Chronogram of the Caryophyllales with putative WGD mapped along with identified

diversification shifts. Diversification analyses were performed on the maximum likelihood tree

(ML) as well as the bootstrap tree set (set) and those shifts that were identified in both groups are

shown. The branches are colored based on Bioclim variable 1 (Mean Annual Temperature).

Fig. 2 The chronograms and mapping of diversification and WGD are as in Fig. 1 (see caption

for details). A) The branches are colored based on Bioclim variable 12 (Mean Annual

Precipitation), and B) based on the principal component analyses (PCA) axis 1.

Fig. 3 Summary of WGD events, with numbers corresponding to those in Table 3, climatic

variables, and diversification shifts. Numbers along branches denote WGD, with the numbers

corresponding to those in Fig. 1 and Table 3. Numbers inside clades denote the number of

diversification rate shifts. Estimated species numbers are listed beside clade names. Box plots

show the values estimated (ancestral values are listed in the top rows, rates in the bottom rows)

for both the left and right clades across bootstrap samples. Clades shaded grey denote a WGD.

b), c), and d) have nested WGD.

Supporting Information

Fig. S1 The cladogram with support mapped for the bootstrap replicates described in the

methods.

Fig. S2 The chronograms and mapping of temperature variables (bioclimatic variables 13-19)

that are not presented in Fig. 1.

Fig. S3 The chronograms and mapping of precipitation variables (bioclimatic variables 13-19)

that are not presented in Fig. 2.

Fig. S4 The chronograms and mapping of PCA axis 2 on the broader Caryophyllales.

Fig. S5 Principal component loadings for bioclimatic variables.