Population genomics of *Cryptococcus neoformans* var. *grubii* reveals new biogeographic relationships and finely maps hybridization

Johanna Rhodes[1*], Christopher A. Desjardins[2*], Sean M. Sykes[2], Mathew A. Beale[1,3,4], Mathieu Vanhove[1], Sharadha Sakthikumar[2], Yuan Chen[5], Sharvari Gujja[2], Sakina Saif[2], Anuradha Chowdhary[6], Daniel John Lawson[7], Vinicius Ponzio[8], Arnaldo L. Colombo[8], Wieland Meyer[9,10], David M. Engelthaler[11], Ferry Hagen[12], Maria Teresa Illnait-Zaragozi[13], Alexandre Alanio[14], Jo-Marie Vreulink[15], Joseph Heitman[16], John R. Perfect[5], Anastasia Litvintseva[16], Tihana Bicanic[3], Thomas S. Harrison[3], Matthew C. Fisher[1#], Christina A. Cuomo[2#]

1 - Department of Infectious Disease Epidemiology, Imperial College London, London, W2 1PG United Kingdom
2 - The Broad Institute of MIT and Harvard, Cambridge, Massachusetts 02142, USA
3 - Institute of Infection and Immunity, St. George's University London, London WC1E 6BT, United Kingdom
4- Wellcome Trust Sanger Institute, Wellcome Genome Campus, Cambridge, United Kingdom
- Division of Infectious Diseases, Department of Medicine, Duke University Medical Center, Durham, North Carolina 27710, USA
- Department of Medical Mycology, Vallabhbhai Patel Chest Institute, University of Delhi, Delhi 110007, India
- Integrative Epidemiology Unit, School of Social and Community Medicine, University of Bristol, Bristol BS8 1TH, United Kingdom
– Division of Infectious Diseases of the Federal University of São Paulo-UNIFESP, São Paulo 04039-032, Brazil
- Molecular Mycology Research Laboratory, Centre for Infectious Diseases and Microbiology, Sydney Medical School-Westmead Hospital, Marie Bashir Institute for Infectious Diseases and Biosecurity, The University of Sydney, Westmead Institute for Medical Research, Sydney, Australia
- Mycology Laboratory, Evandro Chagas National Institute of Infectious Diseases, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil
- TGen North, Translational Genomics Research Institute, Flagstaff, Arizona, USA
- Department of Medical Microbiology and Infectious Diseases, Canisius-Wilhelmina Hospital, Nijmegen, The Netherlands
- Departamento Bacteriología-Micologia, Centro de Investigación, Diagnóstico y Referencia, Instituto de Medicina Tropical Pedro Kourí La Habana, Cuba
- Laboratoire de Parasitologie-Mycologie, AP-HP, Groupe Hospitalier Saint-Louis-Lariboisière-Fernand-WidalParis, France; Université Paris Diderot, Sorbonne Paris CitéParis, France; Unité de Mycologie Moléculaire, Institut Pasteur, Centre National de la Recherche Scientifique, Centre National de Référence Mycoses Invasives et Antifongiques, URA3012 Paris, France.
- Department of Microbiology, Stellenbosch University, Stellenbosch, South Africa
- Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina, USA

*These authors contributed equally to this work

Corresponding authors

Abstract

Cryptococcus neoformans var. *grubii* is the causative agent of cryptococcal meningitis, a significant source of mortality in immunocompromised individuals, typically HIV/AIDS patients from developing countries. Despite the worldwide emergence of this ubiquitous infection, little is known about the global molecular epidemiology of this fungal pathogen. Here we sequence the genomes of 188 diverse isolates and characterized the major subdivisions, their relative diversity and the level of genetic exchange between them. While most isolates of *C. neoformans* var. *grubii* belong to one of three major lineages (VNI, VNII, and VNB), some haploid isolates show hybrid ancestry including some that appear to have recently interbred, based on the detection of large blocks of each ancestry across each chromosome. Many isolates display evidence of aneuploidy, which was detected for all chromosomes. In diploid isolates of *C. neoformans* var. *grubii (*serotype A/A) and of hybrids with *C. neoformans* var. *neoformans* (serotype A/D) such aneuploidies have resulted in loss of heterozygosity, where a chromosomal region is represented by the genotype of only one parental isolate. Phylogenetic and population genomic analyses of isolates from Brazil revealed that the previously 'African' VNB lineage occurs naturally in the South American environment. This suggests migration of the VNB lineage between Africa and South America prior to its diversification, supported by finding ancestral recombination events between isolates from different lineages and regions. The results provide evidence of substantial population structure, with all lineages showing multi-continental distributions demonstrating the highly dispersive nature of this pathogen.

Author Summary

Cryptococcus neoformans var. *grubii* is a human fungal pathogen of immunocompromised individuals that has global clinical impact, causing half a million deaths per year. Substantial genetic substructure exists for this pathogen, with two lineages found globally (VNI, VNII) whereas a third has appeared confined to sub-Saharan Africa (VNB). Here, we utilized genome sequencing of a large set of global isolates to examine the genetic diversity, hybridization, and biogeography of these lineages. We found that while the three major lineages are well separated, recombination between the lineages has occurred, notably resulting in hybrid isolates with segmented ancestry across the genome. In addition, we showed that isolates from South America are placed within the VNB lineage, formerly thought to be confined to Africa, and that there is phylogenetic separation between these geographies that substantially expands the diversity of these lineages. Our findings provide a new framework for further studies of the dynamics of natural populations of *C. neoformans* var. *grubii.*

Introduction

The environmental basidiomycetous yeast *Cryptococcus neoformans* is capable of causing invasive fungal infections primarily in immunocompromised individuals. Meningitis is the most serious manifestation of cryptococcosis. The HIV/AIDS pandemic increased the population of these susceptible individuals and led to an increase in *C. neoformans* infection rates (Day, 2004). *C. neoformans* is the leading cause of mortality in HIV/AIDS patients worldwide, particularly in sub-Saharan Africa, where approximately half a million deaths occur annually (Park et al., 2009). While cryptococcal infection rates in HIV positive individuals have declined due to highly active antiretroviral therapy (HAART), new estimates continue to suggest there are more than 100,000 deaths/year (Rajasingham et al., 2017); recent data also suggests that the incidence of cryptococcosis has plateaued at a high number despite HAART availability. Furthermore, the increasing number of people living with other immunodeficiencies, including transplant and cancer patients, represents a growing population at risk for cryptococcosis (Maziarz and Perfect, 2016).

There are three major serotypes of *C. neoformans* distinguished by different capsular antigens, which include two separate varieties (*Cryptococcus neoformans* var. *grubii* and *Cryptococcus neoformans* var. *neoformans*, serotypes A and D respectively) and a hybrid between the two (serotype AD). While *C. neoformans* isolates are primarily haploid, diploid AD hybrid isolates consisting of both serotype A (*Cryptococcus neoformans* var. *grubii*) and serotype D (*Cryptococcus neoformans* var. *neoformans*) have been isolated from both clinical and environmental sources mostly in Europe (Cogliati, 2013; Desnos-Ollivier et al., 2015; Franzot et al., 1999). Serotype A isolates are the most common cause of infection, accounting for 95% of all *C. neoformans* infections globally (Casadevall and Perfect, 1998; Heitman et al., 2011). Genomes of serotype A and D isolates differ by 10-15% at the nucleotide level (Janbon et al., 2014; Kavanaugh et al., 2006; Loftus et al., 2005), and laboratory crosses of A and D isolates are possible but show reduced viability of meiotic spores (Lengeler et al., 2001; Vogan and Xu, 2014).

Cryptococcus neoformans var. *grubii* can be divided into three molecular types, or lineages: VNI, VNII and VNB (Litvintseva et al., 2006; Meyer et al., 1999, 2009). The VNI and VNII lineages are isolated globally, while the VNB lineage is predominantly located in sub-Saharan Africa (Litvintseva et al., 2006), although there is some evidence for VNB occurring in South America (Bovers et al., 2008; Ngamskulrungroj et al., 2009) and in the USA, Italy, and China in AD hybrid isolates (Litvintseva et al., 2007). Apart from clinical isolation, the VNI lineage is primarily associated with avian excreta (Lugarini et al., 2008; Nielsen et al., 2007) while the VNB lineage is found mostly in association with specific tree species

predominantly mopane trees (Litvintseva and Mitchell, 2012; Litvintseva et al., 2011). These

and recent studies have shown that VNI infections are associated with urbanized

populations where an avian-associated reservoir, pigeon guano, is also found, while the

VNB lineage is widely recovered in the African arboreal environment (Litvintseva et al.,

2011; Vanhove et al., 2017).

Mating in *C. neoformans* occurs between cells of opposite mating types (*MAT***a** and *MAT*α)

(Kwon-Chung, 1975, 1976), although unisexual mating can also occur (Lin et al., 2005).

*MAT*α isolates are capable of unisexual mating both within and between the two serotypes

(Lin et al., 2005, 2007), and recombination was shown to occur at similar levels in bisexual

and unisexual mating in serotype D isolates (Desnos-Ollivier et al., 2015; Sun et al., 2014).

Due to the rarity of *MAT***a** isolates of both serotypes in the environment (Lengeler et al.,

2000a; Litvintseva et al., 2003; Viviani et al., 2001), unisexual mating may have evolved to

enable meiotic recombination and genetic exchange between isolates. Several studies have

found evidence of recombination within VNI, VNII, and VNB populations although not

between these lineages (Bui et al., 2008; Litvintseva et al., 2003, 2005).

An additional level of genome diversity detected in *Cryptococcus neoformans* var. *grubii*

includes the presence of cryptic diploid isolates and variation in the copy number of

individual chromosomes or regions. Close to 8% of *Cryptococcus neoformans* var. *grubii*

global isolates appear diploid; these isolates contain the *MAT*α locus and many appear

autodiploid, thought to result either from endoreduplication or self-mating (Lin et al., 2009).

While the vast majority of serotype A or D isolates appear haploid, individual chromosomes

can be present at diploid or triploid levels (Hu et al., 2011). For chromosome 1, a specific

advantage of aneuploidy is copy number amplification of the azole drug targets or efflux

transporters, associated with drug resistance (Sionov et al., 2010). While the specific

selective advantage of other chromosomal aneuploidies is unknown, same-sex mating of

*MAT*α isolates generates aneuploid progeny at high frequency, some of which also exhibit

azole resistance (Ni et al., 2013). Titan cells, polyploid yeast cells produced in the lung of

infected animals, also generate aneuploid progeny under stress conditions (Gerstein et al.,

2015).

Previous studies examining the global population structure of *Cryptococcus neoformans* var.

grubii have used typing methods for a few genetic loci or focused on particular geographical

regions or countries (Hiremath et al., 2008; Khayhan et al., 2013; Litvintseva et al., 2006;

Oliveira et al., 2004). Recent approaches have applied whole genome sequencing (WGS) to

trace the microevolution of *Cryptococcus*, identifying variation that occurs during the course

of infection (Chen et al., 2017; Ormerod et al., 2013; Rhodes et al., 2017) or in the

environment (Vanhove et al., 2017). Here, we use WGS of 188 isolates to provide a

comprehensive view of the population variation between the three major lineages; the

sequenced isolates were selected to represent the diversity of *Cryptococcus neoformans* var.

grubii including each of the three major lineages and global geographical sampling. We

identify contributions to genomic diversity generated through inter-lineage meiotic exchange

to create haploid hybrids, generation of AD diploid hybrids, and regional copy number

amplification. Furthermore, we finely analyze the phylogenetic relationships and trace the

evolution of *Cng* at the global population level.

Results

Population subdivisions and detection of genetic hybrids

To examine the evolution of *C. neoformans* var. *grubii*, we sampled the population by

sequencing the genomes of 188 isolates (**Table 1**, **Table S1**) representing each of the three

major genetic subpopulations (VNI, VNII, and VNB) previously defined using multi-locus

sequence typing (MLST) (Litvintseva et al., 2006; Meyer et al., 2009). These isolates are

geographically diverse, originating from North America, South America, the Caribbean, Asia,

Europe, and Africa (**Table S1**). The VNI global lineage is the most geographically diverse,

whereas VNII is represented by a smaller number of locations and VNB appears most highly

prevalent in southern Africa. For VNI and VNB, both clinical and environmental isolates were

included, with 25 VNI isolates originating from avian guano or trees and 7 VNB isolates from

trees or other environmental sources (**Table S1**). For each isolate we identified SNPs using

GATK by aligning Illumina reads to the H99 reference genome assembly (**Methods**, (Janbon

et al., 2014)). Whereas 164 isolates appeared haploid, 24 isolates were determined to be

heterozygous diploids (**Methods, Table 1**) and analyzed separately. An initial phylogeny of

the 164 haploid isolates separated the three lineages but intermediate placement of five

isolates suggested the presence of hybrid haploid genotypes (**Figure S1**). As the

phylogenetic placement of such hybrid isolates is complicated by recombination, we

removed these isolates from the phylogenetic analysis and analyzed them using alternative

approaches (see below).

A phylogeny inferred from the SNPs for all non-hybrid isolates strongly supports the

three major lineages of *C. neoformans* var. *grubii*: VNI, VNII, and VNB (**Figure 1**). Of these

159 isolates, only 6 (4%) contain the rare *MATa* allele, including four VNB isolates (Bt63,

Bt85, Bt206, and CCTP15) and two VNI isolates (125.91 and Bt130). Based on these whole

genome SNP comparisons, none of these *MATa* isolates appeared highly related to each

other or to any *MATα* isolate. The two VNI *MATa* isolates are well separated within this

group, with Bt130 found in a subgroup of African isolates and 125.91 most closely related to

a pair of isolates from Africa and North America (**Figure 1**). Phylogenetic analysis showed

that VNB has the highest diversity between isolates, showing the longest tip branches

compared to VNI or VNII. In addition, VNB consisted of two diverged subgroups, VNBI and

VNBII, as suggested previously by MLST (Chen et al., 2015; Litvintseva et al., 2006, 2011)

and genomic analysis (Desjardins et al., 2017; Vanhove et al., 2017).

To better understand the population structure of the three lineages and identify potential

inter-lineage recombination, we compared results of two independent approaches. First, we

used principle components analysis (PCA) to identify the major groups in the population

using the SNP data. By comparing the SNP variants across isolates using PCA, we found

there are three major clusters corresponding to the VNI, VNII, and VNB lineages (**Figure 2**).

The five isolates that showed intermediate positions in phylogenetic analysis (**Figure S1**)

also appeared at intermediate positions by PCA, placed between VNI and VNB. In addition,

two isolates were separated from the VNII cluster and shifted towards the VNB cluster. All of

these seven isolates were collected from southern Africa, and all had a clinical origin except

isolate Ftc260-1, which was isolated from the environment (**Table S1**). Of the seven, two

sets of isolates share nearly identical ancestry ratios and are closely linked on the

phylogenetic tree. Isolates Bt131, Bt162, and Bt163 differed by an average of only 39 SNP

positions; similarly CCTP51 and MW_RSA852 differed by 200 SNP positions, suggesting

these five isolates are descended from two hybridization events. Therefore, four unique

hybridization events were detected in total, three for VNI-VNB and one for VNII-VNB. While

the basal branching VNB isolates from Brazil could suggest a hybrid ancestry, all appear to

be uniformly VNB (>99% of sites).

Next, we identified the ancestry contribution of each isolate using STRUCTURE with three

population subdivisions. This confirmed that most isolates have a single dominant ancestry

assigned to the VNI, VNB, and VNII lineages. In addition, the isolates with intermediate

positions indicated by PCA were found to have mixed ancestry contributions by

STRUCTURE. SNP sites for the VNI-VNB hybrids contain an average of 40.8% VNI

ancestry and 59.2% VNB ancestry whereas the VNII-VNB hybrids have 85.8% VNII and

14.2% VNB ancestry (**Table S2**). The similar fraction of ancestry in the VNI-VNB hybrids

suggests they could be recent mixtures of the two lineages, whereas the VNII-VNB hybrids

may be more ancient mixtures with additional crosses to VNII isolates biasing the final ratio

of parental SNPs.

Evidence of recent meiotic exchange generating haploid hybrids

To examine the degree of intermixing of ancestry for these hybrid genotypes across the

genome, we identified the most likely ancestry for each SNP site using the site-by-site mode

in STRUCTURE. Selecting positions where the ancestry assignment was most confident

(0.9 or greater, **Methods**), we examined the distribution of these sites by ancestry across

the fourteen chromosomes (**Figure 3**). Each of the three VNI-VNB hybrids displayed

different patterns of large regions corresponding to a single ancestry. For example,

chromosome 1 has three large blocks of different ancestry in Bt125, four in Bt131, and two

in Ftc260-1 (**Figure 3A-C**). While all chromosomes contained regions of both VNI and VNB

ancestry groups in Bt125 and Ftc260-1, two chromosomes of Bt131, chromosome 6 and 9,

have only large regions of VNB ancestry. By contrast, CCTP51, which contains a lower

fraction of the second ancestry (VNB), appears more highly intermixed with smaller ancestry

blocks (**Figure 3D**). Notably, three of the four unique genotypes (Bt131, CCTP51, and

Ftc260-1) contain the rare *MAT***a** locus; in all *MAT***a** isolates, the mating type locus region is

of VNB ancestry, whereas the mating locus region in the *MAT*α isolate (Bt125) is of VNI

ancestry (**Methods**). Overall these patterns suggest a recent hybridization of VNI and VNB

isolates, with recombination during meiosis generating chromosome-wide intermixing

resulting in distinct parental haplotype blocks. In Bt125, a 205 kb region of scaffold 6 is

present at nearly twice (1.92 fold) the average depth. Otherwise this isolate, and the other

six hybrid isolates, was found to contain even levels of ploidy across the 14 chromosomes

based on read depth.

For the three VNI-VNB hybrids showing large ancestry blocks, we also utilized the site

ancestry predictions to finely map the genotypes within each population. Given the roughly

equal contribution of the two ancestry sites and the large block size for each in these

genomes, we hypothesized that these hybrids could have resulted from recent mating of one

genotype of each lineage, which we could reconstruct using separate phylogenies of each

site class. For each genotype, sites mapped to either the VNI or VNB ancestry were

selected and a separate phylogeny constructed for each of these two sets of sites. For VNI

ancestry sites, these isolates had very different genotypes, with Ftc260-1 most closely

related to a diverse set of African isolates in VNI, whereas both Bt125 and Bt131 are more

closely related to highly clonal clades of VNI isolates (**Figure S2A,C,E**). Similarly for a

separate phylogenetic analysis of VNB ancestry sites, Bt125 and Bt131 were placed within

one of the two major subclades of VNB while Ftc260-1 was placed in the other (**Figure**

S2B,D,F). This supports that these three hybrids originated from very different genotypes of

VNI and VNB parental isolates.

Diploid isolates and genome plasticity

As noted above, a total of 24 sequenced isolates displayed heterozygous SNP positions

across the genome. Four of these isolates had higher rates of polymorphism overall and

appear to be hybrids within or between VN lineages (Bt66, Cng9, PMHc.1045.ENR.STOR,

and 102-14) (**Figure S3**). Each of these isolates contain two copies of the *MAT*α mating type

locus, suggesting they arose from same sex mating of two *MAT*α parental isolates. In

addition, 11 serotype A diploids showed very low rates of heterogyzosity (**Figure S3**),

consistent with AFLP and MLST-based evidence that they arose from endoreduplication or

self-mating (Lin et al., 2009). The remaining isolates include eight serotypeA/serotypeD

diploids, of which seven contain both *MAT***a** and *MAT*α mating types and one is homozygous

for the *MAT***a** locus, and one serotype A/*Cryptococcus gattii* hybrid containing two copies of

*MAT*α.

All types of diploid isolates in our set, including A/A diploids, exhibit regions of loss of

heterozygosity (LOH) in the genome, where alleles of only one parental isolate are present.

Three of the A/A diploids (Bt66, Cng9, and 102-14) are heterozygous throughout nearly all of

the genome; Cng9 exhibited only a small LOH region at the start of chromosome 2, which

also has haploid levels of genome coverage. Isolate PMHc1045 by contrast has large LOH

regions on six scaffolds, including a 1.1 Mb region of chromosome 6 (**Figure S3**). Some of

these regions of LOH in PMHc1045 are linked to aneuploid chromosome segments,

including a region of chromosome 12 reduced to haploid levels and or triploid levels of the

region adjacent to a LOH on chromosome 6. All LOH regions are telomere-linked,

reminiscent of what has previously been reported across diverse isolates of *Candida*

albicans (Hirakawa et al., 2015).

We next inferred the ancestry of the two parental isolates contributing to the A/A hybrids by

examining the frequency of SNP alleles that are highly predictive for VNI, VNII, or VNB

(**Methods**). Three of the isolates (Cng9, PMHc1045, and 102-14) have similar frequencies

of such VNII and VNB alleles, whereas Bt66 is comprised of VNI and VNB predictive alleles

(**Table S3**). Comparing Cng9 and PMHc1045 directly, 89.2% of variant sites are identical;

this fraction increases to 97.3% when LOH regions are excluded and a similar fraction of

sites are shared with 102-14. By comparison, both isolates share 23% of variant positions

with Bt66 when LOH regions are excluded from both. Notably, LOH has resulted in a mixing

of genotypes; examining predictive alleles for each of the eight LOH regions of PMHc1045

revealed regions corresponding to each lineage. Two regions encompassing 1.4% of the

genome share the highest fraction of private alleles with other VNB isolates whereas the

remaining six regions encompassing 10.2% of the genome share most private alleles with

other VNII isolates. Thus, LOH has led to large differences between otherwise highly similar

Cng9 and PMHc1045 isolates and resulted in blended ancestry by converting regions to

each of the two parents in PMHc1045.

The eight AD hybrids also showed evidence of even more extreme aneuploidy and LOH
related to loss of one of the two parental chromosomes. All isolates displayed evidence of
aneuploidy, by examining read coverage across both the H99 serotype A and JEC21
serotype D reference genomes (**Figure S4**). While some isolates have retained
chromosomes of both A and D origin, others have lost a chromosome from one parent and
duplicated the corresponding chromosome of the other (**Figure 4, Figure S4**). For example
in RCT14, two copies of chromosome 1 are present but both have serotype A origin;
similarly in IFNR21, both copies of chromosome 10 have serotype D origin. Both of these
isolates display additional aneuploidies, with 3 copies of some chromosomes. Notably,
CCTP50 appears mostly triploid, with either 2:1 or 1:2 ratios of the A:D ratio for each
chromosome (**Figure 4**); this pattern is also observed in IFN26 (**Figure S4**). In IFN-R26, loss
of chromosome 4 in JEC21, balanced by gain of chromosome 5 in H99 (**Figure S4**), has
resulted in a *MAT**a***/*MAT**a*** genotype. While the mating type of the original JEC21 parent can
not be determined, this suggests that generation of *MAT**a***/*MAT**a*** diploids can occur via
chromosome loss and duplication. All other isolates are *MAT**a***/*MAT*α, suggesting that they
originated from opposite sex mating. While diploid AD hybrids have been isolated from both
environmental or clinical sources (Litvintseva et al., 2006), all eight AD hybrids in our set are
of clinical origin.

To examine the diversity of these AD hybrids, SNPs were identified by comparison to a
combined A (H99) and D (JEC21) genome reference. Phylogenetic analysis of A and D
genome SNPs revealed that both the A and D copies of each hybrid are closely related for
these isolates (**Figure S5**). On average, the A genomes differ by 6,108 SNP positions and
the D genomes by 3,935 SNP positions. The A genomes are from the VNB lineage, most
closely related to Bt206 in our analysis (**Figure S5**). The low diversity of both the A and D
genomes between isolates suggests that this set of 8 AD hybrids may have originated from
a single hybrid isolate or from a set of closely related A and D parental isolates.

Chromosomal copy number variation
On a smaller scale than whole-genome hybridization, chromosomal copy number variants
appear to be common in *C. neoformans* and may be an adaptive mechanism for virulence
(Rhodes et al., 2017). In the set of 164 primarily haploid isolates, 25 exhibited whole or
partial chromosomal aneuploidies (**Figure S6**). In 13 of the 25 isolates, an entire
chromosome or region thereof showed a doubling of sequencing coverage, consistent with a
diploid chromosome in an otherwise haploid isolate. The remaining 12 isolates show a 50%
gain in coverage better explained by a diploid isolate with a triploid chromosome or region.

These likely diploid isolates do not display heterozygous base calls, suggesting a recent

endoreduplication of the genome and associated aneuploidy of additional chromosomes.

Aneuploidies of particular chromosomes may provide a specific biological advantage or

alternatively be better tolerated. In general, the smallest chromosomes (12 and 13) are the

most frequently observed to exhibit aneuploidy (**Figure S6**). Several isolates have an

increased copy number of chromosome 1; amplification of the lanosterol-14-α-demethylase

ERG11 and the major efflux transporter *AFR1* located on chromosome 1 can confer

resistance to azole drugs (Sionov et al., 2010). Of the four isolates that contain chromosome

1 aneuploidies, either *ERG11* (CCTP34) or *AFR1* (IFN-R11 and RCT6) or both genes

(CCTP9) are present at elevated copy number. The elevated copy number of *AFR1* appears

correlated with increased drug resistance; both CCTP9 and RCT6 displayed fluconazole

MIC values of 256 ug/ml, whereas CCTP34 appeared more susceptible at an MIC of 8 ug/ml

(**Methods**). Notably, all of the isolates with chromosome 1 aneuploidies are of clinical origin,

as are 24 of all 25 isolates with detected aneuploidies (**Figure S6, Table S1**). Of the seven

isolates with hybrid ancestry, only Bt125 included a small region of chromosome 6 at higher

copy number; otherwise this and the other hybrid isolates appeared to be haploid. Across

the diploid and haploid isolates, we detected aneuploidies affecting all chromosomes

(**Figures S3, S4, and S6**).

Conservation of gene content and structure across lineages

To examine the extent of gene content variation across the three major lineages of *C.*

neoformans var. *grubii*, we assembled and annotated genomes of 39 representative isolates

(**Methods**). Previously a high quality reference genome was produced for the H99 VNI

isolate (Janbon et al., 2014); our data set includes new annotated assemblies for 9 diverse

VNB isolates, 27 VNI isolates, and three VNII isolates (**Table S4**). The gene sets across all

40 assemblies (including H99) were compared to each other and to those of four *C. gattii*

(representing VGI, VGII, VGIII, and VGIV) and one *C. neoformans* var. *neoformans*

(serotype D) reference genomes (**Methods**) in order to evaluate gene conservation. Based

on orthologs identified across these genomes (**Methods**), an average of 4,970 genes are

conserved across all 45 compared *Cryptococcus* gene sets; within serotype A, an average

of 5,950 genes are conserved in all 40 genomes (**Figure S7**). A phylogeny inferred from

4,616 single copy genes supports VNII in an ancestral position relative to the more recently

diverging VNI and VNB (**Figure S7**).

Gene content is highly conserved across *C. neoformans* var. *grubii* with few examples of

genes specific to the separate lineages (**Supplementary Note**). Based on ortholog profiling,

a total of 11 genes are specific to VNI, three specific to VNB, and 25 specific to VNII (**Table**
S5). These include two clusters of genes specific to VNI or VNII located within otherwise
syntenic regions of the genome (**Figure 5**). The cluster of five genes unique to VNI genomes
include a predicted haloacid dehydrogenase, an amidohydrolase and an allantoate
permease, which could be involved in uptake of uric acid products. The cluster of six genes
unique to the VNII genomes includes a predicted transcription factor, amino acid transporter,
hydrolase, dihydropyrimidinase, and oxygenase superfamily protein. While both clusters are
also missing from the JEC21 *C. neoformans* var *neoformans* genome, the more distantly
related *C. gattii* genomes contain syntenic orthologs of all of the VNII-specific cluster genes
and between 1 and 3 non-syntenic orthologs of the VNI-specific cluster. These patterns
suggest gene loss and perhaps lateral transfer in some species and lineages account for
these differences. There was little other evidence of lineage-specific gene loss; orthologs
missing in only one lineage included only hypothetical proteins. In addition, we further
searched for genes with loss-of-function mutations in all members of each lineage using
SNP data, to find genes that may be disrupted but still predicted in the assemblies. However,
we found no convincing evidence of disrupted genes with known functions in any of the
three lineages (**Supplementary Note**).

Given the high level of gene conservation between lineages, we sought to identify rapidly
evolving genes that might be involved in phenotypic differences between *C. neoformans*
lineages. For each gene, we built a consensus sequence for each lineage and then
calculated pairwise d_N and d_S of these fixed sites. As d_S was uniformly low throughout the
dataset due to limited genetic diversity, we identified differences in d_N, which measures both
the mutation rate and selection. The top 10 annotated genes with the largest d_N for each
pairwise comparison are shown in Table 2, and the three comparisons in total include 18
unique genes. The set is dominated by transcription factors (*GLN3*, *PDR802*, *SXI1α*,
YOX101, and *ZNF2*) and transferases (*ATG2602*, *CDC43*, *GPI18*, *HOC1* and *RAM1*), many
of which have already been implicated in virulence (Esher et al., 2016; Jung et al., 2015; Lee
et al., 2015; Selvig et al., 2013; Wang et al., 2012) or resistance to oxidative stress (Jung et
al., 2015). In particular, *CDC43* and *RAM1* are both rapidly evolving; these genes represent
the two major independent methods of prenylation, key in proper subcellular localization of
many proteins, often to the membrane (Esher et al., 2016; Selvig et al., 2013). Other rapidly
evolving genes include β-glucan synthase *KRE63,* superoxide dismutase *SOD1*, and mating
regulator *SXI1α*, the latter of which is highly divergent between VNII and both VNI and VNB,
and could play a role in reproductive isolation of the VNII lineage.

Population measures and biogeography

Strikingly, recently identified VNB genotypes from South America are placed in the
phylogeny as basally branching clades for each VNB subgroup, which otherwise consist of
genotypes from Africa (**Figure 1**). All of the six South American VNB isolates contain the
*MAT*α genotype. By contrast, both VNI and VNII consist of more closely related though more
geographically diverse sets of isolates; one large clonal group is found in VNII, whereas
several are observed for VNI, which is oversampled owing to its higher prevalence in
patients and environments worldwide. Overall, VNB showed the highest average pairwise
diversity (pi=0.00736), nearly four times the level in VNI (pi=0.00200), with the lowest value
for VNII (pi=0.00105) (**Table 3**). Genetic diversity within the VNB lineage was similar
between the South America and African isolates (pi=0.00727 and 0.00736, respectively).
However, genetic diversity of VNI isolates in India was lower than VNI isolates in Africa
(pi=0.00146 and 0.00337). VNB also contained the largest fraction of private alleles
compared to VNI and VNII, reflecting the higher variation within VNB (**Table 4**). By contrast,
VNI and VNII had the highest number of fixed differences, reflecting the long branches
leading to these clades. The average divergence (dXY) between lineages ranges is 0.012
comparing isolates from VNI and VNB and 0.015 for comparison of either to VNII (**Table 4**),
highlighting the low nucleotide divergence between the lineages.   VNI and VNII were the
most differentiated of the three lineages as shown by pairwise whole genome fixation
indexes (F_{st}) (Weir and Cockerham 1984). The highest average chromosome F_{st} value is
0.874 between VNI and VNII isolates, while the average chromosome F_{st} values of VNI-VNB
and VNB-VNII are 0.595 and 0.707, respectively (**Table 4**).
419
To further examine the evolutionary history of the novel South American VNB isolates, we
subdivided VNB into four subclades (VNBI-South America, VNBI-Africa, VNBII-South
America, and VNBII-Africa) and calculated alleles unique to each subclade and shared
across VNB groups or geography (Methods). These subclades represent all combinations of
the two previously identified VNB groups (VNBI and VNBII) and the two geographies (South
America and Africa). One South American VNB isolate (V53), nested deeply within African
isolates on the phylogeny, was excluded from the analysis. All four of the subclades
contained more unique alleles than were shared across either VNB group or geography
(**Figure 7**), suggesting both a high level of genetic diversity within each subclade and some
degree of reproductive isolation between each subclade. Furthermore, there was greater
number of unique alleles shared within the VNB groups from different geographic regions
than were shared across VNB groups within the same geographic region (**Figure 7**). This
geographically and phylogenetically segregated diversity suggests that one or more ancient
migration events occurred between South America and Africa during the diversification of
VNB, followed by geographic isolation. In contrast, the VNI and VNII lineages showed a

pattern consistent with more rapid current migration, where isolates from different

geographic regions in many cases differed by fewer than 200 SNPs.

Recombination between and within lineages

The basal branching of Brazilian VNB isolates revealed in the phylogenetic analysis

suggested that South America could be a global center of *C. neoformans* var. *grubii* diversity.

To further investigate this hypothesis, and to explore recombination in the context of

population structure, we implemented the chromosome painting approach of fineStructure

(Lawson et al., 2012), which identifies shared genomic regions between individuals and

thereby ancestral relationships among individuals and populations. Our linked co-ancestry

model found the highest level of sharing among VNB isolates; in addition, there is evidence

of strong haplotype donation from South American VNB isolates (V2, V31, and V87) to all

other lineages and continents, suggestive of ancestral recombination (**Figure 6**).

Independent confirmation of ancestry using STRUCTURE confirmed that V87 includes

primarily VNB ancestry with ~1% VNI alleles (**Table S7**). Interrogating the chunk counts,

which are lengths of DNA shared by a donor to other individuals, and lengths produced by

fineStructure revealed that the haplotype chunks donated by these 'ancestral' isolates were

substantially higher than seen for other isolates, with other African VNB isolates receiving

significant chunks and lengths (Bt102, Bt63, Bt85, Tu229-1, Tu360-1, Tu369-1, and Tu401-

1) from the South American VNB isolates. Isolate V53 donated less strongly than these

three isolates to all lineages. Other South American VNB isolates (WM 1408 and V17)

donated strongly to specific lineages: WM 1408 to VNII and VNB, whilst V17 donated to VNI

and VNB. However, these findings for WM 1408 and V17 were not corroborated using

STRUCTURE. Despite their allocation to separate VNB subpopulations, V2 and V17 (VNB-I

and VNB-II respectively) donate the most genetic material (when interrogating the chunk

counts) to VNI isolates in Africa, India, and Thailand.

Within the VNI lineage, fineStructure analysis identified a subset of isolates with a high

frequency of haplotype sharing (**Figure 8**). Notably, a group of African (Tu259-1, 125.91,

RCT52, Bt100, Bt207 and Bt30) and Indian (INCr213 and INE071) isolates show strong

haplotype donation with many other VNI isolates, suggestive of ancestral recombination

events. These isolates are dispersed over four subpopulations within the VNI lineage.

Though the geographical distance between these populations should preclude frequent

intermixing, these isolates from Africa and India may include a higher fraction of ancestral

alleles, leading to a lack of phylogeographic structure among these highly geographically

distinct populations.

Finding that ancestral recombination in the VNB lineage contributed to VNI lineage diversity
suggested that there will be a signature of admixture linkage disequilibrium (LD) in these two
populations. Linkage disequilibrium differs between lineages (**Figure S8**), with VNII LD
decaying slowly with physical distance, and manifesting an LD50 (where linkage
disequilibrium has decayed to half its maximum value) at >150 kb. However, this value may
reflect the highly clonal nature and relatively small number of sequenced VNII isolates. LD
decay is relatively slow for VNI with an LD50 of 4,500 bp, whereas LD decays more rapidly
in the VNB lineage, with an LD50 of 1,500 bp. When separated into geographical origin of
isolation (**Figure S8** (b)), LD50 for South American VNB appears greater (> 150 kbp) than
that seen in African VNB (2,000 bp). The slower decay of LD in VNI and VNII relative to VNB
may reflect a predominately asexual mode of reproduction owing to the rarity of the *MATa*
idiomorph.
483

Discussion
This population genomic analysis of *Cryptococcus neoformans* var. *grubii* has revealed new
biogeographic relationships and highlighted a complex history of hybridization events
between groups. Analysis of genome-wide variation of 188 geographically diverse isolates
greatly increases the resolution of the VNI, VNII, and VNB phylogenetic groups and
precisely measures the level of genetic differentiation between isolates within each group
and across geographic scales. This data supports a much higher diversity of isolates in the
VNB group compared to VNI and VNII isolates. Notably, we show that hybridization between
these groups can result in genome mixing suggestive of recent and ongoing meiotic
exchange. A recent taxonomic proposal to divide the *C. neoformans* and *C. gattii* species
complexes into seven monophyletic species did not subdivide *C. neoformans* var *grubii* into
separate species; although VNI, VNII, and VNB were strongly supported clades in a
multilocus phylogeny, coalescent based approaches did not clearly support these three
lineages as separate species (Hagen et al., 2015). Our genome-wide analysis has
uncovered new biogeographic structure and ongoing hybridization between lineages of *C.*
neoformans var. grubii, suggesting that further subdivision is not straightforward. In addition,
such hybridization events may be a biological feature that extends across other lineages
within the *C. neoformans* and *C. gattii* species complexes (Farrer et al., 2015; Hagen et al.,
2015), prompting a need for wider investigation of the population genomic structure of this
whole complex is needed to support formal changes in taxonomy (see perspective by
Kwon-Chung'17).
505

The placement of isolates from Brazil at basal branching positions of the two VNB subclades

phylogenetically separates the South American and African isolates within both the VNBI

and VNBII groups. This finding, along with the presence of a large number of unique alleles

in each of these four subclades and strong haplotype sharing seen with fineStructure

analysis (**Figure 7**), suggests that there was ancient migration of the VNB group between

Africa and South America following the initial divergence of VNBI and VNBII but prior to each

group's diversification. This finding appears consistent with a prior report of diverse isolates

from Brazil in a new VNI genotype 1B (Oliveira et al., 2004). While the lack of a reliable

molecular clock combined with substantial rates of recombination prevents dating the time of

divergence between VNB from South America and Africa with confidence, this division

clearly occurred after these continents split over 110 million years ago, and also after VNB

itself subdivided into two lineages – VNBI and VNBII. As is the case with VNI, cross-Atlantic

migration events may have also vectored VNB between these two continents. However,

despite evidence for these migration events, the majority of VNI and VNII migrations were

likely much more recent than is seen with VNB, with nearly clonal isolates of VNI and VNII

found in disparate geographic regions. The presence of one South American VNB isolate

(V53) nested within African isolates on the phylogeny does, however, suggest a limited

number of more recent migration events may be occurring between the two regions even

within VNB despite the large degree of reproductive isolation that we observed. Identification

of additional South American VNB isolates is necessary to determine their diversity and

relationship to isolates from African continental regions. Although the sequenced isolates all

contain the *MAT*α genotype, our sample size was small and likely under-represents the true

diversity of this lineage in South America and the ecological reservoirs that it occupies.

Given the propensity of *Cryptococcus neoformans* var. *grubii* VNI and VNII for having an

environmental reservoir in bird excreta (unlike VNB which is principally associated with

arboreal reservoirs (Litvintseva et al., 2011; Vanhove et al., 2017)), it has been proposed

that pigeons globally dispersed *Cryptococcus neoformans* var. *grubii* from a genetically

diverse population in southern Africa (Litvintseva et al., 2011), resulting in an expansion of

the *Cng* VNI out of Africa. Litvintseva et al. (2011) hypothesized that this "out-of-Africa"

model for the evolution of VNI explains the origin of the global VNI population. Other studies

showing lower genetic diversity of VNI populations in Southeast Asia (Simwami et al., 2011)

and in South America (Ferreira-Paim et al., 2017), further supporting an African origin of

Cryptococcus neoformans var. *grubii*. An alternative explanation for the higher diversity of

African VNI could be that this lineage originated elsewhere and became more diverse in this

continent by mating with the 'native' VNB population or due to other factors. Our analysis did

not find a large subset of VNB alleles within the African VNI isolates based on ancestry

analysis. In addition, we found one VNI subclade composed mostly of African isolates that

appears to be recombining at higher frequency than other VNI groups. The phylogenetic

intermixing of isolates from India and Africa strongly support the hypothesis that there is

long-range dispersal and ancient recombination in environmental populations in India and

Africa, indicative of multiple migratory events across time and into the present. Did VNI

therefore evolve 'Out of Africa'? Further sampling of environmental isolates from across

South America as well as more diverse regions of Africa, as well as estimation of the

mutation rate in *Cryptococcus neoformans* var. *grubii* to allow calibration of a molecular

clock, is needed to further test this hypothesis.

While gene content is very similar across the *Cryptococcus neoformans* var. *grubii* lineages,

we found examples of lineage specific genes including clusters unique to VNI or VNII. While

this suggests that the *Cryptococcus neoformans* var. *grubii* gene inventory based on H99

(Janbon et al., 2014) is largely representative of all lineages, additional genes specific to

VNII and VNB are important to consider in studies focusing on isolates of these lineages.

Differences in gene expression may also differentiate the lineages, and it is important to note

that these will include lineage-specific genes that may contribute to variation in clinical

profiles and virulence that occur among lineages of *Cryptococcus neoformans* var. *grubii*

(Beale et al., 2015). In addition, we found the most rapidly evolving genes between each of

the lineages include transcription factors and transferases, suggesting phenotypic diversity

may be generated through transcriptional reprogramming and protein modification rather

than changes in gene content. The *SXI1* gene detected in comparisons of VNII with both

VNI and VNB appears to be highly substituted in the VNII lineage; this sequence divergence

of *SXI1* in VNII could contribute to differences in mating with this group. Truncated alleles of

SXI1 are frequently observed in the serotype D *MATα* chromosome of AD hybrids and

suggested to contribute to increased mating efficiency (Lin et al., 2007).

Our analysis revealed that hybrid isolates originate from multiple lineages, and resolved the

parental genotypes. Prior analysis with MLST loci suggested that some isolates contain a

mix of multiple genotypes (Chen et al., 2015; Litvintseva et al., 2003). However the

sensitivity and precision of these methods is limited by the small number of loci examined,

the use of genes involved in virulence that may be under different selective pressure, as well

as incomplete lineage sorting in some groups. Analysis of genome-wide variation revealed

that some isolates appear to be a recent mix of different ancestries, based on the detection

of large blocks of sites with each ancestry; this could result from a small number of crossing

over events for each chromosome during meiosis. Other isolates contain more highly

intermixed ancestry across the genome and predominantly of a single ancestry; these may

have occurred by more historical hybridization followed by subsequent mating within a single

lineage group. The demonstration of genome mixing in hybrid isolates raises interesting
questions about whether such fundamentally new assortments of the three lineages could
generate genotypes with new phenotypes, which perhaps have a fitness or selective
advantage.

Analysis of hybrids between serotypes A and D revealed a remarkable degree of genome
reassortment. All of the 8 sequenced AD isolates show evidence of aneuploidy, affecting the
copy number of 12 of 14 serotype A derived chromosomes and all 14 serotype D derived
chromosomes. This is consistent with the high rate of AD isolate aneuploidy previously
reported using flow analysis of DNA content (Lengeler et al., 2001) or comparative genome
hybridization (Hu et al., 2008). For some chromosomes, only one parental genotype was
detected in a subset of five isolates; this includes a loss of the serotype D copy of
chromosome 1, as previously observed in analysis of three AD hybrid isolates (Hu et al.,
2008). However, we further find that loss of heterozygosity (LOH) in some cases is due to
partial copies of several chromosomes, suggesting that genomic instability in AD hybrids
may result in chromosomal breakage. LOH was also observed for smaller regions in diploid
AA hybrids. Similar LOH events are frequently observed in diploid fungi including *Candida*
albicans (Hirakawa et al., 2015) and may contribute to the generation of genetic diversity in
both species.

Aneuploidy was also commonly observed in the haploid *Cryptococcus neoformans* var.
grubii isolates. Additional copies of regions of Chromosome 1 that include *AFR1* or *ERG11*
are associated with drug resistance, though aneuploidies of additional chromosomes are
also observed (Sionov et al., 2010). Although functional significance of aneuploidy of other
chromosomes is less well understood, most of the isolates exhibiting aneuploidy are of
clinical origin, suggesting increased copy of other genes may provide an advantage or that
there is higher genome instability during infection. An isochromosome of the left arm of
chromosome 12 that arose during infection has been reported (Ormerod et al., 2013) and
chromosome 12 aneuploidy is widely seen in African patients with relapsed infections (Chen
et al., 2017; Rhodes et al., 2017) although the specific role of this duplication is unclear. Our
data suggests that there could be additional isochromosomes based on the detection of
partial chromosomes using sequencing read depth; alternatively these regions could be
represented in the genome as fusions with other chromosomes.

Previous studies of *Cryptococcus gattii* have pointed towards South America as a source of
the diversity for the *C. gattii* VGII lineage (Engelthaler et al., 2014; Hagen et al., 2013).
Given the shared evolutionary history of *C. gattii* and *Cryptococcus neoformans* var. *grubii*
(Xu et al., 2000), South America could also represent a major diversity center of

Cryptococcus neoformans var. *grubii*. Our data suggests that *Cryptococcus neoformans* var.

grubii VNB isolates in both subgroups from South America have undergone ancestral

recombination events, donating genetic material to all lineages across multiple geographical

locations. Our study also provides clear evidence for recombination within the VNI and VNII

lineages, where nearly all the isolates contain the *MAT*α mating type. This suggests that

mating likely occurs between *MAT*α isolates, as is found in *C. neoformans* var. *neoformans*

(Sun et al., 2014). Previous studies have hypothesized that *Cryptococcus neoformans* var.

grubii can complete its sexual reproductive life cycle in environmental niches, such as plants

(Xue et al., 2007) and pigeon guano (Nielsen et al., 2007; Vanhove et al., 2017). Our

observations that all lineages of *Cryptococcus neoformans* var. *grubii* show the ability to

widely disperse, to recombine, and to hybridize, illustrates that this pathogen has a high

degree of evolutionary plasticity that is likely related to its success in infecting the

immunosuppressed 'human environment', thereby causing a high burden of mortality

worldwide (Armstrong-James et al., 2014).

Methods

Isolate selection

A total of 188 *C. neoformans* var. *grubii* isolates were selected from previous studies, which

include 146 clinical isolates, 36 environmental isolates, 4 animal isolates and 2 isolates of

unknown isolation source; these isolates were collected from 14 different countries:

Argentina, Australia, Botswana, Brazil, China, Cuba, France, India, Japan, South Africa,

Tanzania, Thailand, Uganda and USA (**Table S1**). Most of the clinical isolates were isolated

from the cerebrospinal fluid of patients. Eight of the 36 environmental isolates were isolated

from pigeon guano, and most of the remaining isolates were collected from Mopane and

other tree species.

Details of clinical trials and ethical review

French isolates were collected during the Crypto A/D study (Dromer et al., 2007). The study

was approved by the local ethical committee and reported to the French Ministry of Health

(registration # DGS970089). For clinical trials undertaken in South Africa (Bicanic et al.,

2007, 2008; Jarvis et al., 2012; Loyse et al., 2012) and Thailand (Brouwer et al., 2004),

ethical approval was obtained from the Wandsworth Research Ethics Committee covering St.

George's University of London. Local ethical approval was obtained from the University of

Cape Town Research Ethics Committee in South Africa the ethical and scientific review

subcommittee of the Thai Ministry of Public Health. Clinical isolates from India were

collected during routine diagnostic service; local ethical approval was obtained from the
Institutional Ethical Committee of Vallabhbhai Patel Chest Institute, University of Delhi, India.

<u>Fluconazole sensitivity testing</u>

Fluconazole MICs were determined for two isolates by the NHLS laboratory in Green Point,
Cape Town using the E-test method (Biomerieux) (Bicanic et al., 2006).

<u>DNA isolation and sequencing</u>

Each yeast isolate was recovered from a freezer stock and purely cultured on an YPD or SD
agar plate for 48-60 h. Next, a single colony was inoculated to another YPD plate and
cultured for 24 h. Approximately 100 µl of yeast cells were used for DNA isolation using the
MasterPure yeast DNA purification kit (Epicenter, Madison, WI) according to the
manufacturer's instructions. Alternatively, a single colony was inoculated into 6ml YPD broth
supplemented with 0.5M NaCl and cultured for 40 hours at 37°C, prior to extraction using the
MasterPure Yeast DNA purification kit (Epicentre) as previously described (Rhodes et al.,
2017).

DNA was sequenced using Illumina technology; for each isolate, a small insert library was
constructed and used to generate between 14 and 150 Million paired-end reads with 101bp
per isolate, which results in 56 to 603 fold average coverage of reads aligned to the H99
genome. In addition, large insert libraries were constructed for 15 isolates (**Table S4**) and
also used to generate 101bp paired-end reads. Isolates were sequenced at Imperial College
London and the Broad Institute (**Table S1**).

<u>Read alignment, variant detection, and ploidy analysis</u>

Illumina reads were aligned to the *Cryptococcus neoformans* var. *grubii* reference genome
H99 (Janbon et al., 2014) using the Burrows-Wheeler Aligner (BWA) 0.7.12 mem algorithm
(Li, 2013) with default parameters. BAM files were sorted and indexed using Samtools (Li et
al., 2009) version 1.2. Picard version 1.72 was used to identify duplicate reads and assign
correct read groups to BAM files. BAM files were locally realigned around INDELs using
GATK (McKenna et al., 2010) version 3.4-46 'RealignerTargetCreator' and 'IndelRealigner'.

SNPs and INDELs were called from all alignments using GATK version 3.4-46
'HaplotypeCaller' in GVCF mode with ploidy = 1, and genotypeGVCFs was used to predict
variants in each isolate. All VCFs were then combined and sites were filtered using

variantFiltration with QD < 2.0, FS > 60.0, and MQ < 40.0. Individual genotypes were then

filtered if the minimum genotype quality < 50, percent alternate allele < 0.8, or depth < 10.

In examining isolates with a high proportion of sites that were removed by these filters,

inspection of the allele balance supported that these isolates were diploid. For heterozygous

diploid isolates, haplotypeCaller was run in diploid mode. VariantFiltration was the same,

with the added filter of ReadPosRankSum < -8.0. Then for individual genotype filtration there

was no allele depth filter but otherwise was the same. The filters were kept as similar as

possible to maximize combinability. For AD hybrids, a combined reference of H99 (Janbon

et al., 2014) and JEC21 (Loftus et al., 2005) was used for alignment and SNP identification.

To examine variations in ploidy across the genome, the depth of bwa alignments at all

positions was computed using Samtools mpileup, and then the average depth computed for

5kb windows across the genome.

MAT locus determination

To evaluate the mating type alleles present in each isolate, Illumina reads were aligned

using bwa mem to a multifasta of both versions of the mating type locus (AF542529.2 and

AF542528.2 (Lengeler et al., 2000b)). Depth at all positions was computed using Samtools

mpileup, and then the average depth computed for the SXI1 and *STE20* genes for both

idiomorphs. Nearly all isolates showed unique mapping to either the *MAT***a** or *MAT*α alleles

of both genes; one isolate, Ftc158, showed significant mapping to both *MAT***a** and *MAT*α,

though 2-fold more to *MAT*α. For the hybrid haploid isolates, the ancestry of the *MAT* locus

was determined from the Structure site by site output.

Genome assembly and annotation

Illumina sequence for each isolate was assembled using Allpaths for 36 isolates (see Table

S4 for release numbers for each assembly) or SPAdes 3.6.0 (with parameter –careful) for

the remaining 3 isolates. Assemblies with both fragment and jump libraries were more

contiguous than those with fragment only data (average of 84 or 561 scaffolds, respectively,

Table S4). However there was little difference in the total contig length between assemblies

with or without jump data (average 18.4Mb and 18.5Mb, respectively, **Table S4**).

The predicted protein coding gene set for each assembly was generated by combining three

primary lines of evidence. Genes were transferred to each new assembly from the well

annotated H99 assembly (Janbon et al., 2014) based on whole genome nucleotide

alignments from nucmer. Genemark-ES (Ter-Hovhannisyan et al., 2008) was run on each

assembly to generate a de novo set of calls. These two sets were combined and improved

using PASA (Haas et al., 2008) with RNA-Seq data of three in vitro conditions (YPD, Limited

media, and Pigeon guano) generated for H99 (Janbon et al., 2014) and for the VNB isolate

Bt85 also input. Repetitive elements were removed from the gene set based on

TransposonPSI (http://transposonpsi.sourceforge.net/) alignments or PFAM domains found

only in transposable elements. The filtered set was assigned sequential locus identifiers

across each scaffold. The average number of 6,944 predicted genes across all assemblies

(**Table S4**) is close to the 6,962 predicted on the H99 reference.

Ortholog identification and comparison

To identify orthologs across the set of 45 *Cryptococcus* genomes (**Table S4**), proteins

clustered based on BLASTP pairwise matches with expect< 1e-5 using ORTHOMCL v1.4 (Li

et al., 2003). To identify orthologs specific to each of the serotype A lineages, we required

that genes were present in 90% of the assembled genomes for VNI (36 or more) or VNB (8

or more) or all VNII (3 genomes). To confirm that orthologs were missing in the other two

lineages, synteny was examined around each gene; in some cases this identified candidate

orthologs missed by OrthoMCL, which were confirmed by BLASTP similarity and removed.

Phylogenetic analysis

A phylogeny for the sets of 159 or 164 isolates was inferred from SNP data using RAxML

version 8.2.4 (Stamatakis, 2014) with model GTRCAT and 1,000 bootstrap replicates. A

separate analysis of the phylogenetic relationship based on gene content included 40

Cryptococcus neoformans var. *grubii* serotype A genomes (28 VNI, 3 VNII, and 9 VNB), 1

Cryptococcus neoformans var. *neoformans* serotype D genome (JEC21), and 4 *C. gattii*

genomes (WM276, R265, CA1873, and IND107) (**Table S4**). The total of 4616 single copy

orthologs identified in all genomes were aligned individually with MUSCLE (Edgar, 2004) at

the protein level, converted to the corresponding nucleotide sequence to maintain reading

frame alignment, poorly aligning regions removed trimal (Capella-Gutiérrez et al., 2009), and

invariant sites removed. A phylogeny was inferred using RAxML version 7.7.8 in rapid

bootstrapping mode with model GTRCAT and 1,000 bootstrap replicates.

Population structure

To examine major population subdivisions, we examined how isolates clustered in a

principal components analysis (PCA). SNP calls for all the isolates were compared using

SMARTPCA (Patterson et al., 2006). To identify the major ancestry subdivisions and their

contributions to the isolates appearing at intermediate positions in the PCA, a total of

338562 randomly subsampled positions containing variants in at least two isolates and less

than 5% missing data were clustered using the Bayesian model-based program

STRUCTURE v2.3 (Pritchard et al., 2000) in the site-by-site mode. Ancestry was plotted

across the genome for each isolate using the maplotlib plotting package in Python.

For analysis of *Cryptococcus neoformans* var. *grubii* diploid isolates (**Table S3**), diagnostic

SNPs for VNB and VNII were present exclusively in the respective group, and called for all

VNB, VNII, and >=100 VNI isolates. Diagnostic SNPs for VNI were present exclusively in

VNII and VNB, and called for all VNB, VNII, and >=100 VNI isolates.

Population genetic measures including Pi, Fst, and Tajima's D were calculated using

popGenome (Pfeifer et al., 2014). d_N and d_S measures were calculated from fixed SNPs in

each lineage using codeml version 4.9c (Yang, 2007). To examine the distribution of the

alleles within VNB, we first identified 445,193 alleles private to VNB (present in at least 1

VNB isolate and no VNI or VNII isolates). We subdivided VNB into four clades (VNBI-South

America, VNBI-Africa, VNBII-South America, and VNBII-Africa) and calculated the number

of those private alleles unique to each clade (present in that one clade and no others) and

shared across VNB groups or geography (present in the two compared clades but no others).

Linkage disequilibrium

Linkage disequilibrium was calculated in 500 bp windows of all chromosomes except for the

~100kb mating type locus on chromosome 5 with vcftools version 1.14 (Danecek et al.,

2011), using the --hap-r2 option with a minimum minor allele frequency of 0.1.

Population inference by fineStructure

Model-based clustering by fineStructure (Lawson et al., 2012) assigns individuals to

populations based on a coancestry matrix created from SNP data, using either Markov chain

Monte Carlo or stochastic optimisation. The algorithm uses chromosome painting, which is

an efficient way of identifying important haplotype information from dense data, such as SNP

data, and efficiently describes shared ancestry within a recombining population. Each

individual is painted using all the other individuals as donors. For example, if an isolate *x* is

clonal and a donor, the clonally related recipients will receive almost all of their genetic

material from isolate *x*, and its closest relatives. This approach has been applied to analyze

recombination in fungal (Engelthaler et al., 2014) and bacterial studies (Yahara et al., 2013).

fineStructure analysis (Lawson et al. 2012) was performed using an all lineage SNP matrix,

with one representative of each clonal VNI population in order to infer recombination,

population structure, and ancestral relationships of all lineages. A separate analysis of all

VNI lineage isolates was also performed. This approach was based on the presence or

absence of shared genomic haplotypes. ChromoPainter reduced the SNP matrix to a

pairwise similarity matrix under the linked model, which utilises information on linkage

disequilibrium, thus reducing the within-population variance of the coancestry matrix relative

to the between-population variance. Since the MAT idiotypes introduce large bias into SNP

analysis, they were removed to enable characterisation of more defined populations. There

was no significant loss of sharing of genetic material when compared to retaining the MAT

locus.

Acknowledgments

We thank the Broad Institute Genomics Platform for generating DNA sequence for this study

and Jose Munoz for helpful comments on the manuscript.

Funding Statement

This project has been funded in whole or in part with Federal funds from the National

Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of

Health and Human Services, under grant number U19AI110818 and by the National Human

Genome Research Institute grant number U54HG003067 to the Broad Institute. Support to

J.R.P. came from Public Health Service Grants AI73896, AI93257. JR and MAB were

supported by a UK Medical Research Council Grant awarded to MCF, TB and TH (MRC

MR/K000373/1). MV was supported by a UK Natural Environment Research Council PhD

studentship. JH was supported by NIH grants AI39115-19 and AI50113-13. The funders had

no role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript.

Data access

All sequence data from this study have been submitted to GenBank under BioProject ID

PRJNA 384983 (http://www.ncbi.nlm.nih.gov/bioproject); individual accession numbers are

listed in Supplemental Tables S1 and S4.

Author contributions

Investigation: JR, CAD, SMS, SS, CAC
Validation: JR, CAD, SMS, CAC
Visualization: JR, CAD, SMS, SS, CAC
Writing – Original Draft Preparation: JR, CAD, CAC
Writing – Review & Editing: JR, CAD, MCF, CAC, AA, MAB, DME, WM, FH, JMV, JH, AL,
JP
Resources: MCF, CAC, MV, YC, JP, TB, TH, VP, ALC, AC, FH, MTI-Z, WM, DME, AA, JMV,

JH
Supervision: CAC, MCF
Funding Acquisition : CAC, MCF
Conceptualization : CAC, MCF, AL

Tables

Table 1. Properties of Sequenced isolates.

Haploid isolates

Population	Isolates (#)	$MAT\alpha$	$MAT\mathbf{a}$
VNI	111	109	2
VNII	23	23	0
VNB	25	21	4
VNI/VNB	5	1	4
VNII/VNB	2	2	0

Diploid isolates

Population	Isolates (#)	$MAT\alpha/MAT\alpha$	$MAT\mathbf{a}/MAT\mathbf{a}$	$MAT\mathbf{a}/MAT\alpha$
VNI/VNB	1	1	0	0
VNII/VNB	2	2	0	0
VNB/Cnn	8	0	1	7
VNB/Cng	1	1	0	0

823 **Table 2.** Rapidly evolving genes in the three lineages of *C. neoformans* var. *grubii*.
824 Consensus sequences were built for each lineage, and d_N and d_S were calculated for each
825 lineage pair. As d_S was uniformly low throughout the dataset due to limited genetic diversity,
826 for each pair of lineages we identified the 10 genes with assigned names (Inglis et al., 2014)
827 with the highest d_N, which measures both the mutation rate and selection.

Comparison	dN	Locus	Gene	Annotation
VNI vs VNB	0.0181	CNAG_01841	*GLN3*	transcription factor, deletion sensitive to organic peroxides (Jung et al., 2015)
	0.0155	CNAG_03894	*PDR802*	transcription factor, deletion with reduced virulence (Jung et al., 2015)
	0.0095	CNAG_03213	*UVE1*	UV damage endonuclease
	0.0092	CNAG_02756	*CDC43*	geranylgeranyltransferase-I, essential for virulence (Selvig et al., 2013)
	0.0090	CNAG_06655	*GPI18*	GPI-anchor transamidase
	0.0089	CNAG_01908	*HEM4*	uroporphyrinogen-III synthase
	0.0085	CNAG_03133	*ATG2602*	UDP-glucose sterol transferase
	0.0084	CNAG_03617	*CLP1*	clampless protein 1
	0.0076	CNAG_05740	*RAM1*	farnesyltransferase β-subunit, essential for virulence (Esher et al., 2016)
	0.0068	CNAG_03637	*YKU80*	Double strand break repair factor and silencing regulator, deletion has reduced virulence (Liu et al., 2008)
VNI vs VNII	0.0610	CNAG_05836	*HOC1*	α 1,6-mannosyltransferase (Lee et al., 2015)
	0.0408	CNAG_05838	*RGD1*	Rho GTPase activating protein, deletion has increased virulence (Liu et al., 2008)
	0.0214	CNAG_06031	*KRE63*	β-glucan synthase, involved in capsule and cell wall formation, deletion has decreased virulence (Gilbert et al., 2010)
	0.0149	CNAG_06814	*SXI1α*	α cell type transcription factor, required for mating (Hull et al., 2002)
	0.0142	CNAG_01841	*GLN3*	see above
	0.0135	CNAG_03229	*YOX101*	transcription factor, deletion sensitive to organic peroxides (Jung et al., 2015)
	0.0127	CNAG_03398	*ZIP2*	zinc ion transporter
	0.0113	CNAG_03133	*ATG2602*	see above
	0.0110	CNAG_03366	*ZNF2*	transcription factor, overexpression results in reduced virulence (Wang et al., 2012)
	0.0104	CNAG_01019	*SOD1*	superoxide dismutase
VNB vs VNII	0.0617	CNAG_05836	*HOC1*	see above
	0.0402	CNAG_05838	*RGD1*	see above
	0.0171	CNAG_06031	*KRE63*	see above
	0.0128	CNAG_03366	*ZNF2*	see above
	0.0122	CNAG_06814	*SXI1α*	see above
	0.0114	CNAG_03213	*UVE1*	see above
	0.0104	CNAG_01019	*SOD1*	see above
	0.0104	CNAG_03398	*ZIP2*	see above
	0.0102	CNAG_01841	*GLN3*	see above
	0.0102	CNAG_02756	*CDC43*	see above

Table 3. Comparison of VN group statistics

Populations	Isolates (#)	Segregating sites	Pi	Tajima's D
VNI	111	190,716	0.00200	-0.107179
VNII	23	337,990	0.00105	-1.005950
VNB	25	613,991	0.00736	-0.232596

Table 4. Pairwise comparison of VN group statistics

Comparisons	Fixed	Shared	Private_A	Private_B	dXY	Fst
VNB vs VNI	54,719	52,536	446,566	102,817	IvB: 0.0119	IvB: 0.595
VNB vs VNII	118,329	68,211	405,406	78,444	BvII: 0.0154	BvII: 0.707
VNI vs VNII	188,590	38,501	116,845	83,802	IvII: 0.0152	IvII: 0.874

 Figure Legends

Figure 1. Phylogenetic analysis supports three major lineages of *C. neoformans* var. *grubii*.
Using a set of 876,121 SNPs across the 159 non-hybrid isolates, the phylogenetic
relationship was inferred using RAxML. The percentage of 1,000 bootstrap isolates that
support each node is shown for major nodes with at least 90% support. For each isolate, the
geographical site of isolation is noted by colored boxes.

Figure 2. Ancestry characterization of three major groups highlights hybrid isolates. A. The
fraction of ancestry (k=3) estimated by STRUCTURE is shown within a column for each
isolate. B. Principal components analysis separates the 3 major lineages, with the hybrid
isolates showing a mix of VNB ancestry with either VNI or VNII.

Figure 3. Large blocks of ancestry suggest recent recombination between lineages. For
each of the four isolates depicted (A-D), the STRUCTURE assigned ancestry for each site
along each chromosome is depicted as a colored bar corresponding to VNI, VNII, and VNB
ancestry. Locations of centromeres are marked with black bars.

Figure 4. Chromosome ancestry and ploidy variation of AD hybrids. For three AD hybrid
isolates (RCT14, IFNR21, and CCTP50), the contribution and copy number of A (green) and
D (blue) ancestry chromosomal regions was measured by aligning all sequence reads to a
combined AD reference (H99, left and JEC21, right). The copy number of each chromosome
is depicted, with either full or partial chromosomal regions shown; see Figure S4 for detailed
coverage plots for all AD hybrid isolates.

Figure 5. Lineage-specific gene clusters. Two large-lineage specific clusters were detected
in the VNI genomes or VNII genomes; these are depicted using a representative genome
from each lineage. A. Insertion of CNAG_06649 to CNAG_06653 in H99 (blue, VNI);
syntenic genes in Bt85 (pink, VNB) and MW_RSA852 (green, VNII) are connected with grey
bars. B. Insertion of C358_04097 to C358_04102 in MW_RSA852.

Figure 6. Genome-sharing analysis of *Cryptococcus neoformans* var. *grubii* using
fineStructure was performed on a SNP matrix using a representative of each clonal
population within the VNI lineage. These genomes were reduced to a pairwise similarity
matrix, which facilitates the identification of population structure based on haplotype sharing
within regions of the genome. The *x*-axis represents the "donor" of genomic regions, while
the *y*-axis represents the recipient of shared genomic regions. The scale bar represents the
amount of genomic sharing, with black representing the largest amount of sharing of genetic
material, and white representing the least amount of shared genetic material (no sharing).
The geographical site of isolation is illustrated with coloured boxes as in Figure 1, and
lineages are also shown.

Figure 7.VNB alleles in population subdivisions and across geography. A. Phylogeny of
VNB lineage showing major subdivisions (VNB-I and VNB-II) and inferred ancestral
geography (South America or Africa, depicted as continent shapes). B. Classification of all
445,193 private VNB alleles (present in at least 1 VNB isolate and no VNI or VNII isolates)
by subdivisions and geography. Most VNB alleles are specific for the each VNB subdivision
and for the geographic subdivisions within each group. More alleles are shared between
geographic locations in the same subdivision (VNBI or VNBII) than are shared within
geographic locations across subdivisions.

Figure 8. Genome-sharing analysis of the *Cng* VNI lineage using fineStructure on a SNP
matrix of 111 genomes. The *x*-axis represents the "donor" of genomic regions, whilst the *y*-
axis represents the recipient of shared genomic regions. The scale bar represents the
amount of genomic sharing, with blue representing the largest amount of sharing of genetic
material, and yellow representing the least amount of shared genetic material (no sharing).

Supplementary Figure Legends

Figure S1. Phylogenetic analysis of all sequenced isolates. Using a set of 876,121 SNPs
across 165 isolates, the phylogenetic relationship was inferred with RAxML. The percentage
of 1,000 bootstrap replicates that support each node is shown for major nodes with at least
90% support. Isolates with hybrid ancestry based on Structure are colored red.

Figure S2. Phylogenetic analysis of ancestry typed SNPs in hybrid isolates. SNPs with VNI
and VNB ancestry were separated for each hybrid isolate, and combined with the wider set
of SNPs for sequenced VNI or VNB isolates. Phylogenies were inferred using RAxML and
the percentage of 1,000 bootstrap replicates that support each node is shown. The hybrid
isolate in each phylogeny is highlighted with red text.

Figure S3. Loss of heterozygosity in AA diploid isolates. For each of the three diploid
heterozygous isolates of AA ancestry, the frequency of heterozygous SNPs/kb within each
isolate and the sequence depth is depicted for each of 14 chromosomes of H99. A. Bt66
has no apparent loss of heterozygosity. B. Cng9 shows a small LOH region at the start of
scaffold 2, which also appears haploid in sequence depth. C. PMHc1045 shows more
extensive LOH on multiple chromosomes, some of which are also associated with
aneuploidy regions.

Figure S4. AD hybrids show high chromosomal aneuploidy. For each AD hybrid isolate the
normalized depth of reads aligned to a combined AD reference is depicted across the
chromosomes of the H99 and JEC21 genomes. In some isolates, the loss of a chromosome
of one ancestry appears compensated by the gain of an extra copy in the other ancestry,
noted by brackets, resulting in homozygosity of some regions of the hybrid genome.
906
907 Figure S5. Phylogenetic analysis of AD hybrid isolates. For each isolate, SNPs called
908 against the H99 or JEC21 reference genome were separated, and those for H99 were
909 combined with those from other VNB isolates. Phylogenies were inferred using RAxML.
910
911 Figure S6. Chromosomal aneuploidy highlighted by normalized read depth. The average
912 depth of Illumina reads aligned to the 14 scaffolds of the H99 genome was plotted across
913 the genome; whole and partial chromosomal aneuploidies were detected for 25 isolates.
914
Figure S7. Phylogenetic relationship and gene conservation for selected *Cryptococcus*.
Using de novo assemblies and associated gene calls for the genome each isolate, orthologs
were identified using OrthoMCL, and the aligned protein sequences of 4,616 single-copy
genes were used to infer a phylogeny using RAxML. Bar graphs represent the numbers of
core protein clusters shared across all genomes (green), conserved protein clusters found in
a subset (grey), species specific protein clusters blue or orange) and clusters specific to a
single genome (yellow).
922

Figure S8. Linkage disequilibrium as a function of distance expressed as the correlation
coefficient r^2 for a) the three lineages, VNB, VNII and VNI and b) South American and
African VNB.

References

Armstrong-James, D., Meintjes, G., and Brown, G.D. (2014). A neglected epidemic:
fungal infections in HIV/AIDS. Trends Microbiol. *22*, 120–127.

Beale, M.A., Sabiiti, W., Robertson, E.J., Fuentes-Cabrejo, K.M., O'Hanlon, S.J.,
Jarvis, J.N., Loyse, A., Meintjes, G., Harrison, T.S., May, R.C., et al. (2015).
Genotypic Diversity Is Associated with Clinical Outcome and Phenotype in
Cryptococcal Meningitis across Southern Africa. PLoS Negl. Trop. Dis. *9*, e0003847.

Bicanic, T., Harrison, T., Niepieklo, A., Dyakopu, N., and Meintjes, G. (2006).
Symptomatic Relapse of HIV-Associated Cryptococcal Meningitis after Initial
Fluconazole Monotherapy: The Role of Fluconazole Resistance and Immune
Reconstitution. Clin. Infect. Dis. *43*, 1069–1070.

Bicanic, T., Meintjes, G., Wood, R., Hayes, M., Rebe, K., Bekker, L.-G., and Harrison,
T. (2007). Fungal burden, early fungicidal activity, and outcome in cryptococcal
meningitis in antiretroviral-naive or antiretroviral-experienced patients treated with
amphotericin B or fluconazole. Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am. *45*,
76–80.

Bicanic, T., Wood, R., Meintjes, G., Rebe, K., Brouwer, A., Loyse, A., Bekker, L.-G.,
Jaffar, S., and Harrison, T. (2008). High-dose amphotericin B with flucytosine for the
treatment of cryptococcal meningitis in HIV-infected patients: a randomized trial. Clin.
Infect. Dis. Off. Publ. Infect. Dis. Soc. Am. *47*, 123–130.

Bovers, M., Hagen, F., Kuramae, E.E., and Boekhout, T. (2008). Six monophyletic
lineages identified within *Cryptococcus neoformans* and *Cryptococcus gattii* by multi-
locus sequence typing. Fungal Genet. Biol. FG B *45*, 400–421.

Brouwer, A.E., Rajanuwong, A., Chierakul, W., Griffin, G.E., Larsen, R.A., White,
N.J., and Harrison, T.S. (2004). Combination antifungal therapies for HIV-associated
cryptococcal meningitis: a randomised trial. Lancet Lond. Engl. *363*, 1764–1767.

Bui, T., Lin, X., Malik, R., Heitman, J., and Carter, D. (2008). Isolates of
Cryptococcus neoformans from infected animals reveal genetic exchange in
unisexual, alpha mating type populations. Eukaryot. Cell *7*, 1771–1780.

Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. (2009). trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses. Bioinforma. Oxf.
Engl. *25*, 1972–1973.

Casadevall, A., and Perfect, J.R. (1998). *Cryptococcus neoformans* (Washington,
D.C.: ASM Press).

Chen, Y., Litvintseva, A.P., Frazzitta, A.E., Haverkamp, M.R., Wang, L., Fang, C.,
Muthoga, C., Mitchell, T.G., and Perfect, J.R. (2015). Comparative analyses of
clinical and environmental populations of *Cryptococcus neoformans* in Botswana.
Mol. Ecol. *24*, 3559–3571.

Chen, Y., Farrer, R.A., Giamberardino, C., Sakthikumar, S., Jones, A., Yang, T.,
Tenor, J.L., Wagih, O., Wyk, M.V., Govender, N.P., et al. (2017). Microevolution of

Serial Clinical Isolates of *Cryptococcus neoformans* var. *grubii* and *C. gattii*. mBio *8*,
e00166-17.

Cogliati, M. (2013). Global Molecular Epidemiology of *Cryptococcus neoformans* and
Cryptococcus gattii: An Atlas of the Molecular Types. Scientifica *2013*, 675213.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A.,
Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., et al. (2011). The variant call
format and VCFtools. Bioinformatics *27*, 2156–2158.

Day, J.N. (2004). Cryptococcal Meningitis. Pract. Neurol. *4*, 274–285.

Desjardins, C., Giamberardino, C., Sykes, S., Yu, C.-H., Tenor, J., Chen, Y., Yang,
T., Jones, A., Sun, S., Haverkamp, M., et al. (2017). Population Genomics And The
Evolution Of Virulence In The Fungal Pathogen *Cryptococcus neoformans*. bioRxiv
118323.

Desnos-Ollivier, M., Patel, S., Raoux-Barbot, D., Heitman, J., Dromer, F., and
French Cryptococcosis Study Group (2015). Cryptococcosis Serotypes Impact
Outcome and Provide Evidence of *Cryptococcus neoformans* Speciation. mBio *6*,
e00311.

Dromer, F., Mathoulin-Pélissier, S., Launay, O., Lortholary, O., and French
Cryptococcosis Study Group (2007). Determinants of disease presentation and
outcome during cryptococcosis: the CryptoA/D study. PLoS Med. *4*, e21.

Edgar, R.C. (2004). MUSCLE: multiple sequence alignment with high accuracy and
high throughput. Nucleic Acids Res *32*, 1792–1797.

Engelthaler, D.M., Hicks, N.D., Gillece, J.D., Roe, C.C., Schupp, J.M., Driebe, E.M.,
Gilgado, F., Carriconde, F., Trilles, L., Firacative, C., et al. (2014). *Cryptococcus
gattii* in North American Pacific Northwest: Whole-Population Genome Analysis
Provides Insights into Species Evolution and Dispersal. mBio *5*, e01464-14.

Esher, S.K., Ost, K.S., Kozubowski, L., Yang, D.-H., Kim, M.S., Bahn, Y.-S.,
Alspaugh, J.A., and Nichols, C.B. (2016). Relative Contributions of Prenylation and
Postprenylation Processing in *Cryptococcus neoformans* Pathogenesis. mSphere *1*.

Farrer, R.A., Desjardins, C.A., Sakthikumar, S., Gujja, S., Saif, S., Zeng, Q., Chen,
Y., Voelz, K., Heitman, J., May, R.C., et al. (2015). Genome Evolution and
Innovation across the Four Major Lineages of *Cryptococcus gattii*. mBio *6*, e00868-
815.

Ferreira-Paim, K., Andrade-Silva, L., Fonseca, F.M., Ferreira, T.B., Mora, D.J.,
Andrade-Silva, J., Khan, A., Dao, A., Reis, E.C., Almeida, M.T.G., et al. (2017).
MLST-Based Population Genetic Analysis in a Global Context Reveals Clonality
amongst *Cryptococcus neoformans* var. *grubii* VNI Isolates from HIV Patients in
Southeastern Brazil. PLoS Negl. Trop. Dis. *11*, e0005223.

Franzot, S.P., Salkin, I.F., and Casadevall, A. (1999). *Cryptococcus neoformans* var.
grubii: separate varietal status for *Cryptococcus neoformans* serotype A isolates. J.
Clin. Microbiol. *37*, 838–840.

Gerstein, A.C., Fu, M.S., Mukaremera, L., Li, Z., Ormerod, K.L., Fraser, J.A., Berman,
J., and Nielsen, K. (2015). Polyploid titan cells produce haploid and aneuploid
progeny to promote stress adaptation. mBio *6*, e01340-1315.

Gilbert, N.M., Donlin, M.J., Gerik, K.J., Specht, C.A., Djordjevic, J.T., Wilson, C.F.,
Sorrell, T.C., and Lodge, J.K. (2010). KRE genes are required for β-1,6-glucan
synthesis, maintenance of capsule architecture and cell wall protein anchoring in
Cryptococcus neoformans. Mol. Microbiol. *76*, 517–534.

Haas, B.J., Salzberg, S.L., Zhu, W., Pertea, M., Allen, J.E., Orvis, J., White, O., Buell,
C.R., and Wortman, J.R. (2008). Automated eukaryotic gene structure annotation
using EVidenceModeler and the Program to Assemble Spliced Alignments. Genome
Biol. *9*, R7.

Hagen, F., Ceresini, P.C., Polacheck, I., Ma, H., van Nieuwerburgh, F., Gabaldón, T.,
Kagan, S., Pursall, E.R., Hoogveld, H.L., van Iersel, L.J.J., et al. (2013). Ancient
dispersal of the human fungal pathogen *Cryptococcus gattii* from the Amazon
rainforest. PloS One *8*, e71148.

Hagen, F., Khayhan, K., Theelen, B., Kolecka, A., Polacheck, I., Sionov, E., Falk, R.,
Parnmen, S., Lumbsch, H.T., and Boekhout, T. (2015). Recognition of seven species
in the *Cryptococcus gattii*/*Cryptococcus neoformans* species complex. Fungal Genet.
Biol. FG B.

Heitman, J., Kozel, T.R., Kwon-Chung, K.J., Perfect, J.R., and Casadevall, A. (2011).
Cryptococcus: From Human Pathogen to Model Yeast (American Society of
Microbiology).

Hirakawa, M.P., Martinez, D.A., Sakthikumar, S., Anderson, M.Z., Berlin, A., Gujja,
S., Zeng, Q., Zisson, E., Wang, J.M., Greenberg, J.M., et al. (2015). Genetic and
phenotypic intra-species variation in *Candida albicans.* Genome Res. *25*, 413–425.

Hiremath, S.S., Chowdhary, A., Kowshik, T., Randhawa, H.S., Sun, S., and Xu, J.
(2008). Long-distance dispersal and recombination in environmental populations of
Cryptococcus neoformans var. *grubii* from India. Microbiol. Read. Engl. *154*, 1513–
1524.

Hu, G., Liu, I., Sham, A., Stajich, J.E., Dietrich, F.S., and Kronstad, J.W. (2008).
Comparative hybridization reveals extensive genome variation in the AIDS-
associated pathogen *Cryptococcus neoformans.* Genome Biol. *9*, R41.

Hu, G., Wang, J., Choi, J., Jung, W.H., Liu, I., Litvintseva, A.P., Bicanic, T., Aurora,
R., Mitchell, T.G., Perfect, J.R., et al. (2011). Variation in chromosome copy number
influences the virulence of *Cryptococcus neoformans* and occurs in isolates from
AIDS patients. BMC Genomics *12*, 526.

Hull, C.M., Davidson, R.C., and Heitman, J. (2002). Cell identity and sexual
development in *Cryptococcus neoformans* are controlled by the mating-type-specific
homeodomain protein Sxi1alpha. Genes Dev. *16*, 3046–3060.

Inglis, D.O., Skrzypek, M.S., Liaw, E., Moktali, V., Sherlock, G., and Stajich, J.E.
(2014). Literature-Based Gene Curation and Proposed Genetic Nomenclature for
Cryptococcus. Eukaryot. Cell *13*, 878–883.

Janbon, G., Ormerod, K.L., Paulet, D., Byrnes, E.J., Yadav, V., Chatterjee, G.,
Mullapudi, N., Hon, C.-C., Billmyre, R.B., Brunel, F., et al. (2014). Analysis of the
genome and transcriptome of *Cryptococcus neoformans* var. *grubii* reveals complex
RNA expression and microevolution leading to virulence attenuation. PLoS Genet.
10, e1004261.

Jarvis, J.N., Meintjes, G., Rebe, K., Williams, G.N., Bicanic, T., Williams, A., Schutz,
C., Bekker, L.-G., Wood, R., and Harrison, T.S. (2012). Adjunctive interferon-γ
immunotherapy for the treatment of HIV-associated cryptococcal meningitis: a
randomized controlled trial. AIDS Lond. Engl. *26*, 1105–1113.

Jung, K.-W., Yang, D.-H., Maeng, S., Lee, K.-T., So, Y.-S., Hong, J., Choi, J., Byun,
H.-J., Kim, H., Bang, S., et al. (2015). Systematic functional profiling of transcription
factor networks in *Cryptococcus neoformans*. Nat. Commun. *6*.

Kavanaugh, L.A., Fraser, J.A., and Dietrich, F.S. (2006). Recent Evolution of the
Human Pathogen *Cryptococcus neoformans* by Intervarietal Transfer of a 14-Gene
Fragment. Mol. Biol. Evol. *23*, 1879–1890.

Khayhan, K., Hagen, F., Pan, W., Simwami, S., Fisher, M.C., Wahyuningsih, R.,
Chakrabarti, A., Chowdhary, A., Ikeda, R., Taj-Aldeen, S.J., et al. (2013).
Geographically structured populations of *Cryptococcus neoformans* Variety *grubii* in
Asia correlate with HIV status and show a clonal population structure. PloS One *8*,
e72222.

Kwon-Chung, K.J. (1975). A new genus, *filobasidiella*, the perfect state of
Cryptococcus neoformans. Mycologia *67*, 1197–1200.

Kwon-Chung, K.J. (1976). A new species of *Filobasidiella*, the sexual state of
Cryptococcus neoformans B and C serotypes. Mycologia *68*, 943–946.

Lawson, D.J., Hellenthal, G., Myers, S., and Falush, D. (2012). Inference of
Population Structure using Dense Haplotype Data. PLoS Genet *8*, e1002453.

Lee, D.-J., Bahn, Y.-S., Kim, H.-J., Chung, S.-Y., and Kang, H.A. (2015). Unraveling
the novel structure and biosynthetic pathway of O-linked glycans in the Golgi
apparatus of the human pathogenic yeast *Cryptococcus neoformans*. J. Biol. Chem.
290, 1861–1873.

Lengeler, K.B., Davidson, R.C., D'Souza, C., Harashima, T., Shen, W.C., Wang, P.,
Pan, X., Waugh, M., and Heitman, J. (2000a). Signal transduction cascades
regulating fungal development and virulence. Microbiol Mol Biol Rev *64*, 746–785.

Lengeler, K.B., Wang, P., Cox, G.M., Perfect, J.R., and Heitman, J. (2000b).
Identification of the MATa mating-type locus of *Cryptococcus neoformans* reveals a
serotype A MATa strain thought to have been extinct. Proc. Natl. Acad. Sci. U. S. A.
97, 14455–14460.

Lengeler, K.B., Cox, G.M., and Heitman, J. (2001). Serotype AD strains of
Cryptococcus neoformans are diploid or aneuploid and are heterozygous at the
mating-type locus. Infect. Immun. *69*, 115–122.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. ArXiv13033997 Q-Bio.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and
SAMtools. Bioinformatics *25*, 2078–2079.

Li, L., Stoeckert, C.J., and Roos, D.S. (2003). OrthoMCL: identification of ortholog
groups for eukaryotic genomes. Genome Res *13*, 2178–2189.

Lin, X., Hull, C.M., and Heitman, J. (2005). Sexual reproduction between partners of
the same mating type in *Cryptococcus neoformans*. Nature *434*, 1017–1021.

Lin, X., Litvintseva, A.P., Nielsen, K., Patel, S., Floyd, A., Mitchell, T.G., and Heitman,
J. (2007). αADα Hybrids of *Cryptococcus neoformans*: Evidence of Same-Sex
Mating in Nature and Hybrid Fitness. PLOS Genet. *3*, e186.

Lin, X., Patel, S., Litvintseva, A.P., Floyd, A., Mitchell, T.G., and Heitman, J. (2009).
Diploids in the *Cryptococcus neoformans* Serotype A Population Homozygous for
the α Mating Type Originate via Unisexual Mating. PLoS Pathog *5*, e1000283.

Litvintseva, A.P., and Mitchell, T.G. (2012). Population Genetic Analyses Reveal the
African Origin and Strain Variation of *Cryptococcus neoformans* var. *grubii*. PLoS
Pathog *8*, e1002495.

Litvintseva, A.P., Marra, R.E., Nielsen, K., Heitman, J., Vilgalys, R., and Mitchell, T.G.
(2003). Evidence of sexual recombination among *Cryptococcus neoformans*
serotype A isolates in sub-Saharan Africa. Eukaryot. Cell *2*, 1162–1168.

Litvintseva, A.P., Kestenbaum, L., Vilgalys, R., and Mitchell, T.G. (2005).
Comparative Analysis of Environmental and Clinical Populations of *Cryptococcus
neoformans*. J. Clin. Microbiol. *43*, 556–564.

Litvintseva, A.P., Thakur, R., Vilgalys, R., and Mitchell, T.G. (2006). Multilocus
sequence typing reveals three genetic subpopulations of *Cryptococcus neoformans*
var. *grubii* (serotype A), including a unique population in Botswana. Genetics *172*,
2223–2238.

Litvintseva, A.P., Lin, X., Templeton, I., Heitman, J., and Mitchell, T.G. (2007). Many
Globally Isolated AD Hybrid Strains of *Cryptococcus neoformans* Originated in Africa.
PLOS Pathog. *3*, e114.

Litvintseva, A.P., Carbone, I., Rossouw, J., Thakur, R., Govender, N.P., and Mitchell,
T.G. (2011). Evidence that the human pathogenic fungus *Cryptococcus neoformans*
var. *grubii* may have evolved in Africa. PloS One *6*, e19688.

Liu, O.W., Chun, C.D., Chow, E.D., Chen, C., Madhani, H.D., and Noble, S.M.
(2008). Systematic genetic analysis of virulence in the human fungal pathogen
Cryptococcus neoformans. Cell *135*, 174–188.

Loftus, B.J., Fung, E., Roncaglia, P., Rowley, D., Amedeo, P., Bruno, D.,
Vamathevan, J., Miranda, M., Anderson, I.J., Fraser, J.A., et al. (2005). The genome
of the basidiomycetous yeast and human pathogen *Cryptococcus neoformans*.
Science *307*, 1321–1324.

Loyse, A., Wilson, D., Meintjes, G., Jarvis, J.N., Bicanic, T., Bishop, L., Rebe, K.,
Williams, A., Jaffar, S., Bekker, L.-G., et al. (2012). Comparison of the early
fungicidal activity of high-dose fluconazole, voriconazole, and flucytosine as second-
line drugs given in combination with amphotericin B for the treatment of HIV-
associated cryptococcal meningitis. Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am.
54, 121–128.

Lugarini, C., Goebel, C.S., Condas, L.A.Z., Muro, M.D., de Farias, M.R., Ferreira,
F.M., and Vainstein, M.H. (2008). *Cryptococcus neoformans* Isolated from Passerine
and Psittacine bird excreta in the state of Paraná, Brazil. Mycopathologia *166*, 61–69.

Maziarz, E.K., and Perfect, J.R. (2016). Cryptococcosis. Infect. Dis. Clin. North Am.
30, 179–206.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,
Garimella, K., Altshuler, D., Gabriel, S., Daly, M., et al. (2010). The Genome Analysis
Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing
data. Genome Res *20*, 1297–1303.

Meyer, W., Marszewska, K., Amirmostofian, M., Igreja, R.P., Hardtke, C., Methling,
K., Viviani, M.A., Chindamporn, A., Sukroongreung, S., John, M.A., et al. (1999).
Molecular typing of global isolates of *Cryptococcus neoformans* var. *neoformans* by
polymerase chain reaction fingerprinting and randomly amplified polymorphic DNA-a
pilot study to standardize techniques on which to base a detailed epidemiological
survey. Electrophoresis *20*, 1790–1799.

Meyer, W., Aanensen, D.M., Boekhout, T., Cogliati, M., Diaz, M.R., Esposto, M.C.,
Fisher, M., Gilgado, F., Hagen, F., Kaocharoen, S., et al. (2009). Consensus multi-
locus sequence typing scheme for *Cryptococcus neoformans* and *Cryptococcus
gattii*. Med. Mycol. *47*, 561–570.

Ngamskulrungroj, P., Gilgado, F., Faganello, J., Litvintseva, A.P., Leal, A.L., Tsui,
K.M., Mitchell, T.G., Vainstein, M.H., and Meyer, W. (2009). Genetic Diversity of the
Cryptococcus Species Complex Suggests that *Cryptococcus gattii* Deserves to Have
Varieties. PLoS ONE *4*, e5862.

Ni, M., Feretzaki, M., Li, W., Floyd-Averette, A., Mieczkowski, P., Dietrich, F.S., and
Heitman, J. (2013). Unisexual and Heterosexual Meiotic Reproduction Generate
Aneuploidy and Phenotypic Diversity De Novo in the Yeast *Cryptococcus
neoformans*. PLOS Biol. *11*, e1001653.

Nielsen, K., Obaldia, A.L.D., and Heitman, J. (2007). *Cryptococcus neoformans*
Mates on Pigeon Guano: Implications for the Realized Ecological Niche and
Globalization. Eukaryot. Cell *6*, 949–959.

Oliveira, M.T.B. de, Boekhout, T., Theelen, B., Hagen, F., Baroni, F.A., Lazera, M.S.,
Lengeler, K.B., Heitman, J., Rivera, I.N.G., and Paula, C.R. (2004). *Cryptococcus*
neoformans Shows a Remarkable Genotypic Diversity in Brazil. J. Clin. Microbiol. *42*,
1356–1359.

Ormerod, K.L., Morrow, C.A., Chow, E.W.L., Lee, I.R., Arras, S.D.M., Schirra, H.J.,
Cox, G.M., Fries, B.C., and Fraser, J.A. (2013). Comparative Genomics of Serial
Isolates of *Cryptococcus neoformans* Reveals Gene Associated With Carbon
Utilization and Virulence. G3 Bethesda Md *3*, 675–686.

Park, B.J., Wannemuehler, K.A., Marston, B.J., Govender, N., Pappas, P.G., and
Chiller, T.M. (2009). Estimation of the current global burden of cryptococcal
meningitis among persons living with HIV/AIDS. AIDS Lond. Engl. *23*, 525–530.

Patterson, N., Price, A.L., and Reich, D. (2006). Population structure and
eigenanalysis. PLoS Genet. *2*, e190.

Pfeifer, B., Wittelsbürger, U., Onsins, S.E.R., and Lercher, M.J. (2014). PopGenome:
An efficient swiss army knife for population genomic analyses in R. Mol. Biol. Evol.
msu136.

Pritchard, J.K., Stephens, M., and Donnelly, P. (2000). Inference of population
structure using multilocus genotype data. Genetics *155*, 945–959.

Rajasingham, R., Smith, S., Park, B., Jarvis, J., Govender, N., and Chiller, T. (2017).
Global burden of disease if HIV-associated Cryptococcal meningitis: An updated
analysis. Lancet Infect. Dis. *17*, in press.

Rhodes, J., Beale, M.A., Vanhove, M., Jarvis, J.N., Kannambath, S., Simpson, J.A.,
Ryan, A., Meintjes, G., Harrison, T.S., Fisher, M.C., et al. (2017). A Population
Genomics Approach to Assessing the Genetic Basis of Within-Host Microevolution
Underlying Recurrent Cryptococcal Meningitis Infection. G3 Genes Genomes Genet.
7, 1165–1176.

Selvig, K., Ballou, E.R., Nichols, C.B., and Alspaugh, J.A. (2013). Restricted
substrate specificity for the geranylgeranyltransferase-I enzyme in *Cryptococcus*
neoformans: implications for virulence. Eukaryot. Cell *12*, 1462–1471.

Simwami, S.P., Khayhan, K., Henk, D.A., Aanensen, D.M., Boekhout, T., Hagen, F.,
Brouwer, A.E., Harrison, T.S., Donnelly, C.A., and Fisher, M.C. (2011). Low diversity
Cryptococcus neoformans variety *grubii* multilocus sequence types from Thailand
are consistent with an ancestral African origin. PLoS Pathog. *7*, e1001343.

Sionov, E., Lee, H., Chang, Y.C., and Kwon-Chung, K.J. (2010). *Cryptococcus*
neoformans Overcomes Stress of Azole Drugs by Formation of Disomy in Specific
Multiple Chromosomes. PLoS Pathog *6*, e1000848.

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. Bioinformatics *30*, 1312–1313.

Sun, S., Billmyre, R.B., Mieczkowski, P.A., and Heitman, J. (2014). Unisexual
reproduction drives meiotic recombination and phenotypic and karyotypic plasticity in
Cryptococcus neoformans. PLoS Genet. *10*, e1004849.

Ter-Hovhannisyan, V., Lomsadze, A., Chernoff, Y.O., and Borodovsky, M. (2008).
Gene prediction in novel fungal genomes using an ab initio algorithm with
unsupervised training. Genome Res. *18*, 1979–1990.

Vanhove, M., Beale, M.A., Rhodes, J., Chanda, D., Lakhi, S., Kwenda, G., Molloy, S.,
Karunaharan, N., Stone, N., Harrison, T.S., et al. (2017). Genomic epidemiology of
Cryptococcus yeasts identifies adaptation to environmental niches underpinning
infection across an African HIV/AIDS cohort. Mol. Ecol. *26*, 1991–2005.

Viviani, M.A., Esposto, M.C., Cogliati, M., Montagna, M.T., and Wickes, B.L. (2001).
Isolation of a *Cryptococcus neoformans* serotype A MATa strain from the Italian
environment. Med. Mycol. *39*, 383–386.

Vogan, A.A., and Xu, J. (2014). Evidence for genetic incompatibilities associated
with post-zygotic reproductive isolation in the human fungal pathogen Cryptococcus
neoformans. Genome *57*, 335–344.

Wang, L., Zhai, B., and Lin, X. (2012). The link between morphotype transition and
virulence in *Cryptococcus neoformans*. PLoS Pathog. *8*, e1002765.

Xu, J., Vilgalys, R., and Mitchell, T.G. (2000). Multiple gene genealogies reveal
recent dispersion and hybridization in the human pathogenic fungus *Cryptococcus*
neoformans. Mol. Ecol. *9*, 1471–1481.

Xue, C., Tada, Y., Dong, X., and Heitman, J. (2007). The human fungal pathogen
Cryptococcus can complete its sexual cycle during a pathogenic association with
plants. Cell Host Microbe *1*, 263–273.

Yahara, K., Furuta, Y., Oshima, K., Yoshida, M., Azuma, T., Hattori, M., Uchiyama, I.,
and Kobayashi, I. (2013). Chromosome painting in silico in a bacterial species
reveals fine population structure. Mol. Biol. Evol. *30*, 1454–1464.

Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol.
Evol. *24*, 1586–1591.

1235

A

VNI   VNB   VNII

B