Mitochondria are physiologically maintained at close to 50 °C

Dominique Chrétien[1,2], Paule Bénit[1,2], Hyung-Ho Ha[3], Susanne Keipert[4], Riyad El-Khoury[5],

Young-Tae Chang[6], Martin Jastroch[4], Howard T Jacobs[7,8], Pierre Rustin[1,2], Malgorzata Rak[1,2]

[1]INSERM UMR1141, Hôpital Robert Debré, 48, Boulevard Sérurier, 75019, Paris, France

[2]Université Paris 7, Faculté de Médecine Denis Diderot, Paris, France

[3]College of Pharmacy, Sunchon National University, Sunchon 540-742, Republic of Korea

[4]Institute for Diabetes and Obesity, Helmholtz Centre Munich, German Research Center for

Environmental Health, 85764 Neuherberg, Germany

[5] Neuromuscular Diagnostic Laboratory, Department of Pathology & Laboratory Medicine,

American University of Beirut Medical Center, Beirut, Lebanon

[6]Department of Chemistry and MedChem Program of Life Sciences Institute, National

University of Singapore, 117543, Singapore

[7]BioMediTech and Tampere University Hospital, FI-33014 University of Tampere, Finland

[8]Institute of Biotechnology, FI-00014 University of Helsinki, Finland

Correspondence to: Pierre Rustin, Inserm U1141, Hôpital Robert Debré, Bâtiment Bingen, 48,

Boulevard Sérurier, 75019, Paris, France

email : pierre.rustin@inserm.fr   Tel : 00 (33) 1 40 03 19 89

Abstract

In warm-blooded species, heat released as a by-product of metabolism ensures stable internal temperature throughout the organism, despite varying environmental conditions. Mitochondria are major actors in this thermogenic process. The energy released by the oxidation of respiratory substrates drives ATP synthesis and metabolite transport, with a variable proportion released as heat. Using a temperature-sensitive fluorescent probe targeted to mitochondria, we measured mitochondrial temperature in situ under different physiological conditions. At a constant external temperature of 38 °C, mitochondria were more than 10 °C warmer when the respiratory chain was fully functional, both in HEK cells and primary skin fibroblasts. This differential was abolished by respiratory inhibitors or in cells lacking mitochondrial DNA, but enhanced by expressing thermogenic enzymes such as the alternative oxidase or the uncoupling protein UCP1. The activity of various RC enzymes was maximal at, or slightly above, 50 °C. Our study prompts a re-examination of the literature on mitochondria, taking account of the inferred high temperature.

Main Text

Mitochondrial targeting of the recently developed, temperature-sensitive fluorescent probe (Fig. S1A), Mito-Thermo-Yellow (MTY)(*1*) was confirmed in both HEK293 cells and primary skin fibroblasts, based on colocalization with the well-characterized MitoTracker green (MTG; Fig. 1A). Initial mitochondrial capture was dependent on the maintenance of a minimal membrane potential(*1*). Fluorescence remained stable over 2 hours in HEK293 cells, but mitochondrial MTY retention varied between cell lines (Fig. S2A). The exact sub-mitochondrial location of the probe is yet to be established, although it has been postulated to reside within the inner membrane. No toxicity of MTY (100 nM in culture medium) could be detected after 3 days.

The fluorescence of MTY (measured in solution at 562 nm) was progressively and reversibly

decreased by as the medium temperature increased (about 50% from 25 to 45°C; Fig. 1B, a,

b). Using a thermostated, magnetically stirred, closable 750 µl quartz-cuvette fitted with an

oxygen sensitive optode device(2) we simultaneously studied oxygen consumption or tension

and changes in MTY fluorescence (Fig. 1C). Adherent cells were harvested and pre-loaded

for 20 min with 100 nM MTY, harvested and washed, then kept as a concentrated pellet at

38°C for 10 min, reaching anaerobiosis in < 1 min. When cells were added to the oxygen-rich

medium they immediately started to consume oxygen (red trace), accompanied by a

progressive decrease of MTY fluorescence (blue trace; phase I; Fig. 1C). In the absence of

any inhibitor, the fluorescence gradually reached a stable minimum (phase II). Once all the

oxygen in the cuvette was exhausted (red trace) the shift of MTY fluorescence reversed

(phase III), returning gradually almost to the starting value (phase IV). To calibrate the

fluorescence signal, the temperature of the extra-cellular medium was increased stepwise

(green trace). MTY fluorescence returned to the prior value when the medium was cooled to

38 °C (Fig. 1C, phase V). Based on this calibration, we estimate the rise in mitochondrial

temperature due to the activation of respiration as ~10 °C (n=10, range 7-12 °C). At the

lowest (phase II) and highest fluorescence value (38°C, imposed by the water bath; phase IV),

the signal was proportional to the amount of added cells (Fig. 1D, a). Cell number did not

affect the maximal rate of fluorescence decrease (phase I), but, once anaerobic conditions had

been reached (phase III, initial), it was inversely related to the initial rate of fluorescence

increase (Fig. 1D, b). To confirm that the observed fluorescence changes were due to

mitochondrial respiration and not some other cellular process, we depleted HEK cells of their

mtDNA with ethidium bromide (EtBr) to a point where cytochrome c oxidase activity was

less than 2% of that in control cells (Fig. 1E, a; Fig. S1C). In EtBr-treated cells, no MTY

fluorescence changes were observed under aerobiosis and cyanide treatment had no effect

(Fig. 1E, b). Because MTY is derived from the membrane potential-sensitive dye rhodamine,

whose fluorescence is essentially unaffected by temperature (Figure 1B, b), we investigated

whether the observed changes MTY fluorescence could be influenced by altered membrane

potential, or by an associated parameter, *e.g.* pH. We therefore compared the response of

MTY fluorescence to cyanide or oligomycin (Fig. 2A, B; Fig. S1D) exerting opposite effects

on membrane potential and pH gradient (Fig. 2C). To avoid the possibly confounding effect

of anaerobiosis, the quartz cuvette was kept uncapped in this experiment, with the oxygen

tension rather than the rates of oxygen uptake being recorded (red traces). Once MTY

fluorescence was stabilized (maximal mitochondrial heating), and the medium re-oxygenated,

cyanide was added, causing a progressive increase in MTY fluorescence to the starting value

(Fig. 2A). Note that, when cyanide was present from the start of the reaction, fluorescence

changes and oxygen uptake were both abolished (Fig. 2A, dotted lines). Adding oligomycin in

lieu of cyanide decreased oxygen consumption and, as observed with cyanide, brought about

an increase in MTY fluorescence (Fig. 2B). Added first, oligomycin strongly decreased

oxygen uptake, abolishing MTY fluorescence decrease (Fig. 2B, dotted lines). Taken

together, these experiments imply that electron flow through the respiratory chain (RC) rather

than membrane potential controls mitochondrial temperature. This conclusion is supported by

the similar effects observed with two other respiratory inhibitors (Fig. S1D), affecting either

RC complex I (CI; rotenone, Fig. 2D) or III (CIII; antimycin, Fig. 2E). Despite their different

effects on the redox state of the various RC electron carriers, these inhibitors blocked oxygen

uptake and again triggered an increase in MTY fluorescence. Taking advantage of cyanide

removal from cytochrome oxidase to form cyanohydrin in the presence of α-ketoacids under

aerobiosis[3], we confirmed that the blockade of the respiratory chain did not result in MTY

leakage from the mitochondria since pyruvate addition resulted in oxygen uptake resuming

and MTY fluorescence decrease, both being inhibited by a further addition of antimycin (Fig.

2F). Noticeably the fluorescence of an endoplasmic reticulum (ER)-targeted version of the

probe in HEK cells and skin fibroblasts was essentially unaffected by the activity of the

mitochondria when modulated by cyanide, pyruvate or antimycin (Fig. S3).

We next studied MTY probe behavior in HEK cells made partially deficient for CI by varying

amounts of added rotenone. The rate of MTY fluorescence decrease was proportional to the

residual respiratory electron flux (Fig. 3A) while the maximal temperature as judged from

MTY fluorescence (phase II) was essentially unchanged (Fig. 3A; inset). We next tested the

effect of expressing the cyanide-insensitive non-proton motive alternative oxidase from *Ciona*

intestinalis (AOX; Fig. 3B, S1E), whose activity is unmasked in the presence of cyanide(*4*).

Before cyanide addition, the decrease in MTY fluorescence in AOX-expressing cells was

similar to control cells, consistent with previous inferences that the enzyme does not

significantly participate in uninhibited cell respiration. However upon cyanide addition,

AOX-endowed cells maintained low MTY fluorescence (Fig. 3B, blue trace), despite oxygen

consumption decreasing by more than 50% (red trace). The increased ratio of heat generated

to respiration is consistent with the predicted thermogenic properties of AOX. Subsequent

addition of 0.1 mM propylgallate, which inhibits AOX, almost completely abolished the

residual respiration and brought MTY fluorescence back to its starting value (corresponding

to 38°C).

So as to circumvent the fact that we were not able to use chemical uncouplers with this

probe(*1*), we used HEK cells engineered to express the uncoupling protein 1 (UCP1; Fig. 3C,

S1F). As expected, UCP1 conferred an increased rate of respiration, which was only partially

inhibited by oligomycin (red trace), and accompanied by an even greater drop in MTY

fluorescence, equivalent to a temperature at least 15 °C above the cellular environment.

The surprisingly high inferred mitochondrial temperatures prompted us to check the

dependence on assay medium temperature of RC enzyme activities measured under V_{max}

conditions (Fig. 4A, B) in crude extracts, where mitochondrial membrane integrity is

maintained. Antimycin-sensitive CIII, malonate-sensitive succinate:cytochrome c reductase

(CII+CIII) and cyanide-sensitive cytochrome c oxidase (CIV) activities all showed

temperature optima at or slightly above 50°C, whilst these activities tended gradually to

decrease again as the temperatures were raised further (Fig. 4A). This was not so for those

enzymes whose activities can be measured in vitro only after osmotic disruption of both outer

and inner mitochondrial membranes (Fig. 4B). Oligomycin-sensitive ATPase (CV) activity

was optimal around 46 °C, whereas rotenone-sensitive NADH quinone-reductase (CI) activity

declined sharply at temperatures above 38 °C. Under the even more disruptive conditions of

clear-native electrophoresis, which requires detergent-dispersal of the membrane-bound

complexes, in-gel activity even of CII or CIV was impaired at temperatures above 42 °C and

46 °C, respectively (Fig. 4C).This strongly suggests a vital role for the lipid components of

the inner mitochondrial membrane in the stabilization of the RC complexes at high

temperature. We next analyzed the temperature profile of RC activity of primary skin

fibroblasts. For CII+CIII, CIII and CIV (Fig. 4D), as well as CV (Fig. 4E) similar temperature

optima were observed as in HEK cells, whilst MTY fluorescence (Fig. 4F) also indicated

mitochondria being maintained at least 6-10 °C above environmental temperature.

Our findings raise numerous questions concerning the biochemistry, physiology and

pathology of mitochondria. The physical, chemical and electrical properties of the inner

mitochondrial membrane and of mitochondria in general, will need to be re-evaluated, given

that almost all previous literature reflects experiments conducted far from physiological

temperature. Traditional views of the lipid component of the respiratory membrane as a lake

in which the RC complexes are floating resulting in a random-diffusion model of electron

transfer or, more recently, as a sealant occupying the space between tightly packed

proteins(*5*), need to be revised in favour of one that considers it more as a glue that maintains

the integrity of the respiratory complexes. The effects of respiratory dysfunction need to be

reconsidered, to include those attributable to temperature changes, such as effects on

membrane fluidity, electrical conductance and transport. RC organization into

supercomplexes(*6, 7*) should be re-examined at more realistic temperatures, using methods

other than CNE, which does not maintain functional integrity even of the RC complexes

individually. Finally, whilst the subcellular distribution of mitochondria (e.g. perinuclear, or

synaptic) has previously been considered to reflect ATP demand, mitochondria should also be

considered as a source of heat, potentially relevant in specific cellular or physiological

contexts, not just in specifically thermogenic tissues like brown fat.

Legends to Figures

Figure 1 Determination of mitochondrial temperature in intact human cells.

A: The temperature-sensitive probe MitoThermoYellow (MTY: a, b) co-localizes with

MitoTrackerGreen (MTG: c, d, merge: e, f) in HEK293 cells and in primary skin fibroblasts

(e, f). B: a, Fluorescence excitation (green) and emission (red) spectra of MTY (1 mM) in

solution in 2 ml PBS at 25°C and 45°C; b, Response of MTY (blue and red) and rhodamine

(green) fluorescence (1 mM) in solution in 2 ml PBS to temperature (34 to 64°C). Note that

the linear decrease of MTY fluorescence to increasing temperature (blue) is essentially

reversed upon cooling (red) of the medium to the initial temperature. c, Linear increase of

fluorescence of HEK cells (preloaded 10 min before trypsinization with 100 μM MTY)

according to cell number (using cell protein concentration as surrogate parameter). C, The

definition of the various phases of MTY-preloaded HEK cell fluorescence adopted in this

study. phase I: cell respiration (red trace) after cells are exposed to aerobic conditions in PBS,

resulting in decreased MTY fluorescence (blue trace) as mitochondria heat up; phase II: cell

respiration under aerobic conditions, which has reached a steady-state of MTY fluorescence

(maximal warming of mitochondria). Cells were initially maintained for 10 min at 38°C under

anaerobic conditions, before being added to the cuvette; phase III: cell respiration that has

arrested due to oxygen exhaustion - MTY fluorescence progressively increases to the starting

value, as mitochondria cool down; phase IV: respiration remains stalled due to anaerobiosis;

after reaching steady-state, MTY fluorescence is dictated only by the water bath temperature;

phase V, respiration remains stalled due to anaerobiosis; temperature of the cell suspension

medium (green trace) shifted by stepwise adjustments to water-bath temperature, followed by

return to 38°C. Measurements were carried out in a quartz chamber closed except for a 0.6

mm addition hole in the hand-made cap. D, Maximal rate of decrease of MTY fluorescence

(%, blue circles; corresponding with mitochondrial warming) is not significantly affected by

cell number, whereas initial fluorescence increase in presence of cyanide (%, green circles,

corresponding with initial rate of mitochondrial cooling) is modulated by cell number. D: a,

HEK cells were made severely deficient for cytochrome oxidase by culture (10 days) in the

presence of ethidium bromide (EtBr; 1 µg/ml). Cytochrome c oxidase activity (blue circles)

declined to a few percent of the activity measured at t=0, whilst citrate synthase activity

(green circles) was little changed; b, The fluorescence of EtBr-treated HEK cells (10 days of

EtBr treatment) pre-loaded with MTY (blue continuous line) does not decrease following

suspension in oxygenated medium, whilst that of control HEK cells (blue dotted lines)

follows the profile documented in Fig. 1C; in contrast to control cells (red dotted line), EtBr-

treated HEK cells do not consume oxygen (red continuous line).

Figure 2. The rate of respiratory electron flow determines the temperature of mitochondria in

intact HEK cells.

A: The effect of 1 mM cyanide on MTY fluorescence (blue lines) and cell respiration (red

lines), when added under aerobic conditions (continuous lines), or when present from the start

of the experiment (dotted lines). Changes in the temperature of the cell suspension medium

(green line), imposed by water bath adjustment, were used to calibrate the MTY fluorescence

changes. B: The effect of 12.5 µM oligomycin on MTY fluorescence (blue lines) and oxygen

tension (affected by cell respiration balanced by medium stirring) in the uncapped quartz-

cuvette (red lines), when added to free respiring cells (continuous lines), or when present from

the start of the experiment (dotted lines). C: a, effects of different inhibitors on rhodamine

fluorescence, in digitonin (0.001%)-permeabilized HEK293 cells supplied with 10 mM

succinate and 0.1 mM ADP as indicated. Under state 3 conditions 12.5 µM oligomycin and

0.8 mM KCN have qualitatively opposite effects on rhodamine fluorescence, used as an

indicator of membrane potential ($\Delta\Psi$). The effects of 0.6 µM antimycin (D) and 2 µM

rotenone (E) on MTY fluorescence and oxygen tension, plotted as for oligomycin in (B, b). F:

The effect of adding pyruvate on MTY fluorescence (blue line) and oxygen uptake (red line)

by KCN-inhibited HEK cells. Temperature calibration (green line) of MTY fluorescence as in

A.

Figure 3. Effects on mitochondrial temperature of respiratory inhibitors, uncouplers and

expression of heterologous mitochondrial proteins.

A: Effect of variable rotenone addition to control HEK cells on the rates of oxygen uptake and

fluorescence decrease of MTY. Rotenone was added at t=4 min, rates calculated from 4 to 7

min expressed as a percent of initial rate. Inset: Maximal warming up of HEK cell

mitochondria in the absence or presence of 20 µM rotenone. B, C: Changes in MTY

fluorescence (blue lines), cell respiration (B, C) (red lines) and temperature of cell suspension

medium (green line), with additions of inhibitors as shown (KCN, n-propyl gallate and/or

oligomycin), alongside Western blots confirming expression or knockdown of the indicated

genes *Ciona intestinalis* alternate oxidase, AOX (B), UCP1 (C), alongside loading controls as

indicated. Traces for control cells are shown by dotted lines.

Figure 4. Effects of assay medium temperature on RC activities. Temperature profile of (A,

D) cytochrome c oxidase (CIV), malonate-sensitive succinate-cytochrome c reductase

(CII+CIII) and antimycin-sensitive decylubiquinol:cytochrome c reductase (CIII) activity in

(A) HEK293 cells, and (D) primary skin fibroblasts, after two freeze-thaw cycles, and (B, E)

oligomycin-sensitive ATPase (CV) and rotenone-sensitive NADH:decylubiquinone (CI)

activity, after disruption of inner mitochondrial membrane in frozen (B) HEK293 cells and

(E) primary skin fibroblasts, by osmolysis with water(*8*). Colours denote optimal (green),

minimal (red) and degrees of intermediate (pale greens, yellow) activity. Grey bars indicate

optimal temperature range. C: CNE in-gel activities of CI, CIV and CII extracted from

mitochondria and incubated for 10 min at (a) 4 °C , (b) 37°C, (c) 42 °C, (d) 46 °C and (e) 55

°C, also plotted graphically (lower panel). F: Changes in MTY fluorescence (blue), cell

respiration (red) and temperature of cell suspension medium (green), for primary skin

fibroblasts, as denoted in Fig. 2A, with addition of KCN as shown.

Material and Methods

Cell culture

Human cells derived from Embryonic Kidney (HEK293), Hepatoma Tissue Culture (HTC-

116) and from large cell lung cancer (NCI-H460) cells (American Type Culture Collection,

Manassas, VA 20108 USA) were cultured in DMEM medium containing 5 g/l glucose

supplemented by 2 mM glutamine (as Glutamax™), 10% foetal calf serum, 1 mM pyruvate,

100 µg/ml penicillin/streptomycin each. AOX-(*9*) or UCP-endowed(*10, 11*) HEK293 cells

were obtained as previously described.

Primary skin fibroblasts were derived from healthy individuals and grown under standard

condition in DMEM glucose (4.5 g/l), 6 mM glutamine, 10% FCS, 200 µM uridine,

penicillin/streptomycin (100 U/ml) plus 10 mM pyruvate.

Immunoblot analyses and *in gel* enzyme activity assays

For the Western blot analysis, mitochondrial proteins (50 µg) were separated by SDS–PAGE

on a 12% polyacrylamide gel, transferred to a nitrocellulose membrane and probed overnight

at 4°C with antibodies against the protein of interest, AOX 1:10,000[10] , UCP1 1:10,000[11].

Membranes were then washed in TBST and incubated with mouse or rabbit peroxidase-

conjugated secondary antibodies for 2 h at room temperature. The antibody complexes were

visualized with the Western Lightning Ultra Chemiluminescent substrate kit (Perkin Elmer).

For the analysis of respiratory chain complexes, mitochondrial proteins (100 μg) were

extracted with 6% digitonin and separated by hrCN-PAGE, on a 3.5–12% polyacrylamide gel.

Gels were stained by in gel activity assay (IGA) detecting CI, CII and CIV activity as

described[12].

Staining procedures and life cell imaging

Cells (HEK293, MTC, NIC, primary skin fibroblasts) were seeded on glass coverslips and

grown inside wells of a12 well-plate dish for 48 h in standard growth media at 37 °C, 5%

CO2. The culture medium was replaced with pre-warmed medium containing fluorescent

dyes, 100 nmol MitoTracker Green (Invitrogen M7514) and 100 nmol MitoThermo

Yellow(*1*) or 100 nmol ER Thermo Yellow(*12*). After 10 min the staining medium was

replaced with fresh pre-warmed medium or PBS buffer and cells were observed immediately

by Leica TCS SP8 confocal laser microscopy.

Assay of mitochondrial respiratory chain activity

The measurement of RC activities was carried out using a Cary 50 spectrophotometer (Varian

Australia, Victoria, Australia), as described(*13*). Protein was estimated using the Bradford

assay.

Simultaneous spectrofluorometric, temperature, oxygen uptake assay

Detached sub-confluent HEK (NCI, HTS) cells (25 cm^2 flask) or trypsinized sub-confluent

skin fibroblasts (75 cm^2 flask) were treated for 20 min with 100 μM MTY in 10 ml DMEM,

and spun down (1,500 *g* x 5 min). The pellet is resuspended in 1 ml PBS, cells being next

spun down (1,500 *g* x 5 min) and kept as a concentrated pellet. After anaerobiosis (checked

by inserting an optic fiber equipped with an oxygen-sensitive fluorescent terminal sensor

(Optode device; FireSting O_2, Bionef, Paris, France) was established (10 min incubation of

the pellet at 38°C;), cells (1 mg prot) were added to 750 µl of 38°C-thermostated medium.

The fluorescence (excitation 542 nm, emission 562 nm for MTY; excitation 559 nm, emission

581 nm for ERTY), the temperature of the medium in the cuvette and the respiration of the

intact cell suspension were simultaneously measured in a magnetically-stirred, 38°C-

thermostated 1 ml-quartz cell in 750 µl of PBS using the Xenius XC spectrofluorometer

(SAFAS, Monaco). Oxygen uptake was measured with an optode device fitted to a handmade

cap, ensuring either closing of the quartz-cell yet allowing micro-injections (hole with 0.6 mm

diameter) or leaving the quartz-cell open to allow for constant oxygen replenishment.

Alternatively, untreated HEK293 cells (250 µg protein) were added to 750 µl of medium

consisting of 0.25 M sucrose, 15 mM KCl, 30 mM KH_2PO_4 (pH 7.4), 5 mM $MgCl_2$, EGTA 1

mM, followed by addition of rhodamine to 100 nM and digitonin to 0.01 % w/v. The

permeabilized cells were successively given a mitochondrial substrate (10 mM succinate) and

ADP (0.1 mM) to ensure state 3 (phosphorylating) conditions, under which either 6.5 µM

oligomycin or 1 mM cyanide was added.

Statistics

Data are presented as mean ± SD statistical significance was calculated by standard unpaired

one-way ANOVA with Bonferroni post-test correction; a $p< 0.05$ was considered statistically

significant (GraphPad Prism).

Data availability statement.

The authors declare that all data supporting the findings of this study are

available within the paper and its supplementary information files.

References

1.   S. Arai *et al.*, *Chem Commun (Camb)* **51**, 8044 (May 11, 2015).
2.   R. El-Khoury *et al.*, *PLoS Genet* **9**, e1003182 (2013).
3.   D. E. Green, S. Williamson, *Biochem J* **31**, 617 (Apr, 1937).
4.   J. T. Bahr, W. D. Bonner, Jr., *J Biol Chem* **248**, 3446 (May 25, 1973).
5.   G. Lenaz *et al.*, *Biochim Biophys Acta* **1797**, 633 (Jun-Jul, 2010).
6.   H. Schagger, *Biochim Biophys Acta* **1555**, 154 (Sep 10, 2002).
7.   M. Wu, J. Gu, R. Guo, Y. Huang, M. Yang, *Cell* **167**, 1598 (Dec 01, 2016).
8.   D. Chretien *et al.*, *Biochem Biophys Res Commun* **301**, 222 (Jan 31, 2003).
9.   R. El-Khoury *et al.*, *Free Radic Biol Med* **96**, 57 (Jul, 2016).
10.   R. Oelkrug *et al.*, *Nat Commun* **4**, 2140 (2013).
11.   S. Keipert, M. Jastroch, *Biochim Biophys Acta* **1837**, 1075 (Jul, 2014).
12.   S. Arai, S. C. Lee, D. Zhai, M. Suzuki, Y. T. Chang, *Sci Rep* **4**, 6701 (Oct 21, 2014).
13.   P. Rustin *et al.*, *Clin Chim Acta* **228**, 35 (Jul, 1994).

330
331

End notes

Abbreviations

AOX, Alternative oxidase; CNE, Clearnative Electrophoresis; ERTY, Endoplasmic

Reticulum Thermo Yellow; EtBr, Ethidium bromide; MTG, MitoTracker Green; MTY,

MitoThermo Yellow; RC, Respiratory chain; UCP1 Uncouplin protein 1

Acknowledgements

This work was supported by French (ANR MITOXDRUGS to DC, PB, MR and PR) and

European (E-rare Genomit to DC, PB, MR and PR) institutions and patient's associations to

PB and PR Association d'Aide aux Jeunes Infirmes (AAJI), Association contre les Maladies

Mitochondriales (AMMi), Association Française contre l'Ataxie de Friedreich (AFAF), and

Ouvrir Les Yeux (OLY).

Authors contribution

M.R., HT.J. and P.R. conceived the project. HT.J., M.J., YT. C. and P.R. wrote the

manuscript. D.C., P.B., R. EK., HH.H., S.K., M.R. and P.R. conducted research. All authors

contributed to data analysis and manuscript preparation.

Author information

Reprints and permissions information is available at www.nature.com/reprints.

We confirm that there are no known conflicts or competing financial interests of interest

associated with this publication and there has been no significant financial support for this

work that could have influenced its outcome.

Correspondence and requests for materials should be addressed to Pierre Rustin,

pierre.rustin@inserm.fr, or Malgorzata Rak, malgorzata.rak@inserm.fr.

A

B

C

    Supplemental Material   *3 Figures and their legends*

Figure S1. The structure of the MitoThermo Yellow probe and a series of schematized views

of the various conditions used for testing warming up of mitochondria *in situ* in cells.

A, The structure of the rhodamine-derived MTY probe. B, The concurrent synthesis of ATP

and heat generation by the respiratory chain. C, The defective respiratory chain of EtBr

treated cells. D, The sites of action of the several inhibitors used in this study. The respiratory

chain of AOX- (E) or UCP1- (F) expressing HEK cells. I, II, III, IV, V, the various complexes

of the respiratory chain; Aa, antimycin A; AOX, Alternative oxidase; Fum, fumarate; Oligo,

oligomycin; Q, ubiquinone 10 (coenzyme Q); Rot, rotenone; Succ, succinate; UCP1,

uncoupling protein 1.

Figure S2. MTY probe well preserved *in situ* in HEK mitochondria is slowly (NCI cells) or

rapidly (HTC cells) expulsed from mitochondria.

A, Initial MTY fluorescence is mostly localized to mitochondria in HEK (a), NCI (b), and

HTC (c) cell lines as shown by the overlapping staining of MTY and MitoTracker green

(yellow color). After 2 hours, a significant amount of the probe is excluded from

mitochondria in NCI cells (e), resulting in a number of cells being green or red colored.

Noticeably red (MTY) fluorescence is observed in cytosolic small granules (h). A similar but

much more pronounced phenomenon is observed in HTC cells where large granules can be

observed (f, i, j). B, Quantification of MTY-stained (red) granules (a) and clustered-granules

(b) in HEK, NCI and HTC cells. C, MTY-fluorescence changes (as in Fig. 2A) in HEK, NCI

and HTC cells upon shift from anaerobic to aerobic conditions and the effect of a subsequent

addition of cyanide. Noticeably while cyanide restores the initial fluorescence value in HEK

cells it does not in NCI and even less in HTC cells. All together these experiments indicated

that depending on cell types MTY can be either preserved for long (2 hours) in mitochondria

(HEK cells) or more or less rapidly excluded as cytosolic granules (NCI, HTC cells) causing

an irreversible loss of MTY fluorescence as measured in the spectrofluorometer quartz

cuvette (C).

A

B

C

Figure S3. Modulating respiratory chain activity does not change the fluorescence of an

endoplasmic reticulum-targeted (ER thermos yellow; ERTY) version of MTY.

A, ERTY- (red) and MitoTracker green (green) fluorescence did not overlap (bottom) in HEK

cells. B, Tested as MTY in the spectrofluorometer (Fig. 1), the fluorescence of ERTY within

HEK cells (brown line) was unaffected by the activity of the mitochondria modulated by

cyanide, pyruvate or antimycin, chemicals that fully controlled oxygen uptake (red line).

Noticeably, fluorescence decrease of ERTY was similar in the initial absence or presence of

cyanide (not shown). C, ERTY- (red) and MitoTracker green (green) fluorescence did not

overlap either in skin fibroblasts (C) at variance with MTY and MitoTracker green which

staining perfectly overlaps in the same cells (Fig. 1Af).