Concentration Dependent Chromatin States Induced by the Bicoid Morphogen

Gradient

Colleen E. Hannon[1], Shelby A. Blythe[1] and Eric F. Wieschaus[1*]

[1]Department of Molecular Biology/Howard Hughes Medical Institute, Princeton University, Princeton, NJ

08544, USA

*Correspondence: efw@princeton.edu

ABSTRACT

In *Drosophila*, graded expression of the maternal transcription factor Bicoid (Bcd)

provides positional information to activate target genes at different positions along the

anterior-posterior axis. We have measured the genome-wide binding profile of Bcd

using ChIP-seq in embryos expressing single, uniform levels of Bcd protein, and

grouped Bcd-bound targets into four classes based on occupancy at different

concentrations. By measuring the biochemical affinity of target enhancers in these

classes *in vitro* and genome-wide chromatin accessibility by ATAC-seq, we found that

the occupancy of target sequences by Bcd is not primarily determined by Bcd binding

sites, but by chromatin context. Bcd drives an open chromatin state at a subset its

targets. Our data support a model where Bcd influences chromatin structure to gain

access to concentration-sensitive targets at high concentrations, while concentration-

insensitive targets are found in more accessible chromatin and are bound at low

concentrations.

INTRODUCTION

During embryonic development, multicellular organisms must generate the

patterned tissues of an adult organism from a single undifferentiated cell. This process

requires highly regulated control of gene expression, both in developmental time and at

reproducible positions in an embryo. These complex gene regulatory networks are

controlled by systems of transcription factors, which bind to DNA and control the

expression of genes required for development (Levine and Davidson, 2005). In early

Drosophila melanogaster embryos, the Bicoid transcription factor forms an anterior-to-

posterior protein gradient the embryo (Driever and Nüsslein-Volhard, 1988b). Bcd

functions as transcriptional activator to pattern the embryo, binding to target gene

enhancers and activating gene expression at distinct positions along the AP axis,

corresponding to different concentrations of Bcd protein (Driever and Nüsslein-Volhard,

1988a; Struhl et al., 1989).

Recent studies of Bcd function suggest that its interaction with its targets may be

more complex than the simple concentration-dependent activation originally proposed

for morphogen gradients (Wolpert, 1969). In the absence of a strong Bcd gradient,

embryos still exhibit patterned expression of Bcd target genes, and these genes can be

activated at lower concentrations of Bcd than these nuclei would be exposed to in a

wild-type embryo (Chen et al., 2012; Liu et al., 2013; Ochoa-Espinosa et al., 2009).

While changing Bcd dosage shifts cell fates, the shifts deviate quantitatively from those

expected of strict concentration dependence, especially as expression patterns are

assayed progressively later during development (Liu et al., 2013). These studies have

consequently raised doubts about the extent to which the local concentration of Bcd

along its gradient determines the spatial patterns of target gene expression in the

embryo.

While these studies argue against a strict application of the morphogen

hypothesis for Bcd, the patterned expression of target genes is also influenced by other

maternal patterning systems and interactions among the Bcd targets themselves (Chen

et al., 2012; Jaeger, 2010; Löhr et al., 2009). Chen, *et al*. have shown that the posterior

boundaries of Bcd target genes are positioned by a system of repressors including

Runt, Krüppel, and Capicua. This work suggests that the Bcd gradient does not directly

establish expression domains of its target genes but rather is just one player in a

network of patterning genes that influence cell fates in the embryo. However, using

target gene expression as a metric for Bcd function does not address how information

from the Bcd gradient initially establishes distinct cell fates.

Part of the difficulty in evaluating direct roles of the Bcd gradient arises from the

unknown nature of the molecular mechanism by which Bcd establishes concentration

thresholds different positions along the gradient. A simple model of the positioning of

Bcd target genes predicts that *cis*-regulatory elements of different genes respond to

different concentrations of Bcd. Genes in the anterior would have low affinity Bcd

binding sites and could therefore only be activated by high Bcd concentrations, whereas

genes expressed in more posterior positions would have higher affinity binding sites

(Driever et al., 1989b). Direct measurements of Bcd binding affinity have been

conducted *in vitro* using DNA probes (Burz et al., 1998; Gao and Finkelstein, 1998; Ma,

1996; Ma et al., 1996) and have demonstrated that Bcd is able to bind cooperatively to

achieve sharp concentration thresholds. While these measurements lend some support

to a simple affinity model, little correlation has been shown between predicted binding

site affinity and AP position of gene expression (Ochoa-Espinosa et al., 2005; Segal et

al., 2008). However, neither *in vitro* measurements of Bcd binding nor computational

predictions of binding sites can capture interactions between Bcd and its target

enhancers in the context of local chromatin structure.

Using high throughput sequencing approaches, we measured *in vivo* genome-

wide Bcd-DNA binding and chromatin accessibility in transgenic embryos expressing

different concentrations of uniform Bcd protein. These data reveal distinct classes of

enhancers that differ in their sensitivity to Bcd concentration. We find that these classes

differ both in the DNA binding motifs that they contain and in their local chromatin

accessibility. We also find that Bcd influences the accessibility of a subset of its target

enhancers, primarily at highly concentration-sensitive enhancers that drive gene

expression in the anterior of the embryo. This leads us to a model in which target

enhancers throughout the genome have a broad range of sensitivities for Bcd protein,

and can therefore respond to a range of Bcd concentrations along the gradient.

However, rather than arising from differences in Bcd binding site composition, these *in*

vivo interactions are chromatin context-dependent, and Bcd influences the chromatin

structure of its target enhancers.

RESULTS

Bicoid target gene expression boundaries are influenced by other maternal

factors, but its physical interaction with enhancers is not

To investigate the mechanism whereby Bcd functions to pattern the AP axis, we

performed chromatin immunoprecipitation followed by high throughput sequencing

(ChIP-seq) to determine the genome-wide binding profile of Bcd to its targets. We

performed the ChIP-seq experiments on embryos expressing GFP-tagged Bcd in a *bcd*

null mutant background that were staged precisely in nuclear cycle 14 (NC14), and

established a list of robust and reproducible list of 1,027 peak Bcd binding regions (see

Supplemental File 1 and Experimental Procedures). These peaks successfully identify

63 out of 66 of the previously identified Bcd target enhancers (Chen et al., 2012) and

overall associate with enhancers whose expression patterns span broadly across the

AP axis.

As a transcriptional regulator, Bcd activates the expression of a subset of its

targets whose expression domains are predominantly located in the anterior half of the

embryo. In *bcd* mutant embryos, such targets are not expressed. For example, the gap

genes *buttonhead* (*btd*) and *knirps* (*kni*) have anterior expression domains that are not

present in *bcd* mutant embryos (Figure 1A). The posterior *kni* expression domain,

however, is expressed in *bcd* embryos, albeit with shifted positional boundaries. These

distinct domains of *kni* expression are controlled by separate enhancer elements

(Pankratz et al., 1992; Schroeder et al., 2004), both of which are bound by Bcd *in vivo*

(Supplemental File 1). In the absence of all maternal AP patterning inputs (*bicoid nanos*

hunchback torsolike quadruple mutants), the *kni* posterior enhancer is not expressed.

However, in embryos where Bcd is the sole source of maternal patterning information

(*nanos hunchback torsolike* triple mutants), the *kni* posterior domain is expressed with a

near wild-type anterior expression boundary (Figure 1A and Figure 1 Figure

Supplement 1A). The *kni* posterior domain therefore represents a second class of Bcd

target gene, which depends on Bcd to determine the position of its expression but does

not demonstrate an absolute requirement of Bcd for transcriptional activation.

Both classes of Bcd target genes receive positional cues both from Bcd and from

other patterning systems. We considered the possibility that, given their influence on the

expression domains of Bcd target genes, the posterior and terminal patterning systems

may impact Bcd binding to its target enhancers in different nuclei along the AP axis. We

therefore tested whether loss of the posterior and terminal systems (*nanos* and

torsolike) would alter the Bcd ChIP-seq profile. We used the statistical package EdgeR

(Robinson et al., 2010) to test for differential Bcd binding between wild-type and *nanos*

torsolike embryos and found that we could not detect any significant change in binding

at any of these 1,027 regions (Figure 1 Figure Supplement 1B). Therefore, although the

expression domains of Bcd target genes are ultimately influenced by inputs from other

AP patterning systems, the physical interaction of Bcd with the DNA in the enhancers of

these genes occurs independently of other maternal AP patterning inputs.

Embryos expressing Bcd uniformly show developmental fates reflecting the

concentration of Bcd

We set out test whether incremental changes in Bcd concentration along the

gradient can be read out directly at the level of binding to target enhancers. Due to the

graded distribution of Bcd, each nucleus along the AP axis is exposed to a different

concentration of the protein. To measure the Bcd binding state at individual

concentrations, we performed ChIP-seq on embryos expressing Bcd at single, uniform

concentrations in every nucleus along the AP axis. Several previous studies have

included genetic manipulations in which the Bcd gradient has been flattened to assess

its activity independently of its distribution (Chen et al., 2012; Driever and Nüsslein-

Volhard, 1988a; Löhr et al., 2009; Ochoa-Espinosa et al., 2009). However, genetically

disrupting the gradient does not result in a total flattening, and transgenic approaches to

date have not allowed for precise and reproducible control over the level of expression

of the flattened Bcd. We therefore generated transgenic lines expressing GFP tagged

Bcd in which the endogenous 3'UTR responsible for graded localization is flanked by

FRT sites that allow it to be replaced with the unlocalized *spaghetti squash* 3'UTR. To

generate different expression levels of uniform Bcd, we coupled transgenes to different

maternally active promoters that yield embryos in which individual uniform Bcd

concentration approximates single positions along the gradient (see Figure 1 Figure

Supplement 1C).

To determine the expression levels of the uniform lines, we imaged GFP

fluorescence in live embryos expressing either uniform or graded GFP-Bcd (Gregor et

al., 2007a) (Figure 1B). The endogenous *bcd* promoter drives a level of uniformly

expressed Bcd equivalent to that measured at approximately 65% egg length of the

wild-type gradient. The *matrimony (mtrm)* and *αTubulin67C (αTub67C)* promoters drive

expression levels corresponding to approximately 45% and 25% egg length,

respectively. For simplicity, we refer to the uniform lines as low (*bcd* promoter), medium

(*mtrm* promoter), and high (*αTub67C* promoter). (See also Figure 1 Figure Supplement

1D)

Uniform expression of Bcd confers gene expression profiles and developmental

programs representative of distinct positions along the AP axis. The head gap gene

buttonhead (btd) is expressed in an anterior stripe in wild-type embryos (Figure 1A), but

expands to fill the entire middle of the embryo at the highest level of uniform Bcd (Figure

1C). At the medium level, the Btd anterior stripe is duplicated at the posterior, and at the

lowest Bcd level, it is not expressed. The gap gene *knirps*, which is expressed in an

anterior domain and a posterior stripe, shows a duplication of its anterior domain in the

posterior in high uniform Bcd embryos. There is also a weaker duplication at the

medium Bcd level. There is no apparent anterior expression at the lowest level, but an

expanded posterior stripe is present. The gene expression patterning we observe in the

presence of uniform Bcd likely result from the activity of additional maternal patterning

cues (*nanos* and *torso*) as well as interactions between the Bcd target genes

themselves. The concentration-dependent activity of uniform Bcd is also apparent in

cuticle preparations of embryos expressing the transgenic constructs. The transgenic

constructs specify increasingly anterior structures along larval body plan as the

concentration of Bcd increases (Figure 1D). These effects on the body plan indicate that

the uniform Bcd transgenes are capable of specifying cell fates that reflect their relative

expression levels.

Bcd binding to genomic targets is concentration dependent

We next determined genome-wide Bcd binding profiles at each individual

concentration by ChIP-seq and used these measurements to assign each of the 1,027

peak regions to classes distinguished by their degree of concentration-dependent Bcd

binding (Figure 2A). Using EdgeR (Robinson et al., 2010), we selected peak regions

that exhibited statistically significant (FDR ≤ 0.05) differences in binding by performing

pairwise exact tests between the three uniform Bcd concentrations. This yielded four

different classes of peaks, one concentration-insensitive class, and three classes with

increasing sensitivity to Bcd concentration.

The Concentration-Insensitive peak class (n = 143) shows no significant

differences between any of the concentrations of uniform Bcd we tested. Concentration-

Sensitive III peaks (n = 593) are significantly reduced in binding between the highest

and lowest Bcd concentrations, but reductions are not significant between high and

medium, or medium and low. Concentration-Sensitive II peaks (n = 138) are significantly

reduced in binding at the lowest Bcd level compared to either the medium or the high

levels. Finally, Concentration-Sensitive I peaks (n = 152) are significantly reduced in

binding at both the medium and the low Bcd levels compared to the highest level

(Figure 2A). These different groups suggest that Bcd binds differentially to target

enhancers at specific concentrations, and furthermore that certain subsets of enhancers

are bound only in anterior nuclei whereas others are bound broadly across the entire AP

axis.

Although 63 out of 66 previously characterized Bcd-dependent enhancers are

identified in our ChIP peaks, the majority of the 1,027 peaks identified have not been

extensively examined. Within the set of known Bcd targets, there is strong correlation

between position of expression and the associated Bcd sensitivity class (Figure 2A). To

extend this observation to previously uncharacterized Bcd target enhancers, we queried

the Fly Enhancer resource generated from the Vienna Tile GAL4 reporter library (Kvon

et al., 2014). The Fly Enhancer collection is a library of candidate enhancer DNA

fragments driving expression of GAL4 that covers 13.5% of the non-coding genome.

Each fragment's expression pattern has been measured and scored by developmental

stage. A total of 293 enhancer candidates overlap with at least one peak in our data set.

Of these, 163 drive gene expression in stage 4-6 (which includes NC14), and these

active enhancers overlap with a total of 151 (14.7%) of the Bcd-peaks. The remaining

overlapping fragments either are active later in development (75), or are not functional

(55). Given the large fraction of the queried enhancers that are active during early

development (163 out of 293 overlapping enhancers), it remains possible that a similar

fraction of the of the 876 peaks (77.2%) that do not overlap with the Fly Enhancer

candidates may correspond to enhancers active in the early embryo.

The Bcd sensitivity classes are predictive of the expression domains of

associated enhancer fragments. Enhancers overlapping with both the Concentration-

Sensitive I and II classes drive expression in anterior regions of the embryo, with the

Concentration-Sensitive III and Concentration-Insensitive classes driving broad and

posterior expression, respectively (Figure 2B). This indicates that our classifications of

the Bcd-bound peaks reflect unique groups of Bcd targets with differing abilities to bind

Bcd protein and consequently activate gene expression in different positions along the

AP axis. The boundary positions of anteriorly expressed Bcd targets may be refined at

the transcriptional level by interactions with opposing gradients of repressors like Runt

(Chen et al., 2012), that are Bicoid targets themselves. We addressed whether such

repression could account for the restricted expression of Concentration-Sensitive I and

II targets to anterior regions of the embryo by examining whether they were enriched for

binding of such repressors. By comparison with genome-wide binding profiles of

transcription factors in the BDTNP ChIP database (Li et al., 2008; MacArthur et al.,

2009), we instead find that the factors associated with each Bcd peak class are

generally those whose expression patterns overlap with the average expression

domains of each class (Figure 2 Figure Supplement 1A). Peaks in the Concentration-

Sensitive I class, for example, are enriched for binding of the terminal gap gene

Huckebein whereas those in the Concentration-Sensitive II class are enriched for

Krüppel and Giant binding. However, we find no evidence that the Bcd sensitivity

classes are predominantly defined by repressive interactions.

Sequence composition of ChIP sensitivity classes does not account for *in vivo*

sensitivity to Bcd concentration

We next wanted to determine whether the Bcd-bound regions in each sensitivity

class differ at the level of DNA sequence. *In vitro,* Bcd binds with high affinity to the

consensus 5'-TCTAATCCC-3', and that variations on this consensus sequence

constitute weak binding sites (Burz et al., 1998; Driever and Nüsslein-Volhard, 1989;

Driever et al., 1989b). If the affinity of a given enhancer for Bcd were encoded primarily

at the level of its DNA sequence, we would expect to see a higher representation of

strong Bcd binding sites in the less sensitive classes, and weaker sites in the more

sensitive classes. To test this, we performed *de novo* motif discovery using the RSAT

peak-motifs algorithm (Thomas-Chollier et al., 2012; 2008). We identified the top motifs

in the entire Bcd ChIP peak list, ranked by their e-value, and found that the top three

most highly ranked motifs were the consensus binding site for the proposed pioneer

factor Zelda (Zld) (Bosch et al., 2006; De Renzis et al., 2007; Harrison et al., 2011; Nien

et al., 2011), and a strong (TAATCC) and weak (TAAGCC) Bcd binding site (Figure 2C).

We next calculated the frequency with which these motifs appear in each peak, and

tested for enrichment between sensitivity classes by permutation test (Figure 2D). We

found that despite their failure to bind Bcd at low concentrations, the Concentration-

Sensitive I and II classes are enriched for both the strong and weak Bcd sites relative to

the peak set as a whole. Given our result that these classes drive expression primarily

in the anterior of the embryo (Figure 2B), the higher density of Bcd binding sites in these

enhancers contrasts with previous studies that have found little correlation between

number of binding sites and position of gene expression (Ochoa-Espinosa et al., 2005).

This difference likely reflects the larger sample size used in our study, as well as our

method for classifying Bcd bound peaks. The Concentration-Sensitive III class did not

contain an enrichment of any site over the total peak set. The Concentration-Insensitive

class, however, showed a higher prevalence of the Zld binding site relative to the total

peak set than any other class.

These results indicate that, in contrast to a binding site affinity model for Bcd

function, Bcd target enhancers that behave as concentration-sensitive and -insensitive

in vivo are not distinguished by their representation of strong versus weak Bcd binding

sites, confirming previous studies (Ochoa-Espinosa et al., 2005). In further support of

this concept, we found little correlation between *in vitro* binding affinity by

electrophoretic mobility shift assay and the *in vivo* binding properties we observe by

ChIP for a selected subset of peaks (Figure 2 Figure Supplement 2). At the level of

sequence composition, they instead appear to differ in their balance of Bcd and Zld

binding sites. Although both strong and weak Bcd sites and Zld sites are enriched in the

Bcd ChIP peaks as a whole, there is a bias toward both Bcd sites in peaks that show

concentration-sensitive binding properties by ChIP-seq and a bias toward Zld sites in

the concentration-insensitive peaks. Zld, a ubiquitously expressed early embryonic

transcription factor, has been implicated in chromatin remodeling prior to zygotic

genome activation (Harrison et al., 2011; Nien et al., 2011; Sun et al., 2015). The

predominance of Zld motifs over Bcd motifs in the Concentration-Insensitive class

suggests that *in vivo* chromatin structure also plays a role in the sensitivity of a given

target to transcription factor concentration in the context of the developing embryo.

Taken together, these findings suggest that the chromatin context of an enhancer may

play a greater role in its overall affinity for a transcription factor *in vivo* than the

sequences of the binding sites that it contains. We therefore set out to test the

hypothesis that sensitivity classes are distinguished at the level of chromatin structure.

Bcd is required for chromatin accessibility at a subset of concentration-sensitive

target sites

To measure genome-wide patterns of chromatin accessibility and nucleosome

positioning, we performed ATAC-seq (Buenrostro et al., 2015) on single wild-type

embryos precisely staged at 12 minutes after the onset of NC14, and identified 13,266

peaks of chromatin accessibility (see Experimental Procedures). Of the 1,027 Bcd-

bound regions identified by ChIP-seq, 855 (83.3%) of them overlap with ATAC-seq

peaks.

Given Zelda's role in influencing chromatin accessibility and the presence of its

binding sites at Bcd-bound regions of genome, we measured the effect of Zld on

accessibility at Bcd sites by ATAC seq (Figure 3A). Of the total 13,226 accessible

regions at NC14, 2,675 (20.2%) show a significant reduction in accessibility in *zld*

mutant embryos. This fraction is higher in Bcd-bound peaks; 402 (39.1%; or 379

[44.3%] of 855 the Bcd peaks that overlap with ATAC open regions, see Table S5)

show reduced accessibility in *zld* mutants, indicating that Bcd bound regions are more

likely to be dependent on Zld for their accessibility than the genome as a whole.

However, the Zld-dependent peaks are distributed across each sensitivity class

determined by ChIP, with no particular class being significantly more Zld-dependent.

This contrasts with the distribution of binding sites in the peak classes, which revealed

that the Concentration-Insensitive peaks were more likely to contain Zld binding sites.

These results suggest that while Zld contributes to the accessibility of a subset of Bcd

target gene enhancers, it is unlikely to determine the differential concentration sensitivity

of Bcd peaks as a whole.

Given the enrichment for both strong and weak Bcd binding sites in the

Concentration-Sensitive I and II classes, we next examined the impact of Bcd protein

itself on chromatin accessibility by ATAC seq (Figure 3A). In *bcd* mutants, 326 (2.4%) of

the 13,266 open regions in wild-type embryos show significantly reduced accessibility

accompanied by increased nucleosome occupancy in those same regions (Figure 3A

and B). These regions are therefore either directly or indirectly dependent on Bcd for

their accessibility. More strikingly, 132 (12.9%) of the 1,027 Bcd ChIP-seq peaks show

reduced accessibility in the absence of Bcd and likely represent regions where Bcd's

impact is direct. These regions dependent on Bcd for accessibility are significantly

enriched for peaks in the Concentration-Sensitive I and II classes (32.9% and 31.9% of

each class, with Fisher's exact test P-values of 4.29×10^{-12} and 1.37×10^{-10}, respectively).

In contrast, the Concentration-Sensitive II and Concentration-Insensitive classes are

both significantly underrepresented (6.07% and 0.7% and P-values = 7.88×10^{-13} and

2.29×10^{-8}) (Figure 3B). This suggests that Bcd binding influences chromatin

accessibility preferentially at a subset of highly concentration-sensitive enhancers.

Because the Concentration-Sensitive I and II classes are bound primarily at high

Bcd concentrations, Bcd's effects on chromatin accessibility at these targets likely

occurs only in anterior regions of the embryo. In support of this, we find that chromatin

accessibility at Bcd-dependent, concentration-sensitive targets is responsive to Bcd

concentration. Expressing uniform Bcd confers accessibility to peaks that are not

accessible in *bcd* mutant embryos (Figure 3B). The degree of chromatin accessibility

conferred by Bcd correlates positively with the concentration of uniform Bcd expressed

(Figure 3B). This observation, along with the overrepresentation of the Concentration-

Sensitive I and II classes in the Bcd-dependent peaks, suggests that Bcd influences the

chromatin state of these targets primarily at the high concentrations found in the anterior

of the embryo.

A second feature that distinguishes the chromatin structure of Bcd binding sites

is the presence of DNA sequences favorable for nucleosome occupancy (Segal et al.,

2006). Bcd bound regions in wild type embryos are generally depleted of nucleosomes

(Figure 4A). However, predicting nucleosome positioning sequences using the NuPoP

algorithm (Xi et al., 2010) suggests that the Concentration-Sensitive I and II Bcd

enhancer classes are more likely to bind nucleosomes than the Concentration-Sensitive

III and Concentration-Insensitive classes. (Figure 4B). The contrast between predicted

occupancy and observed depletion suggests that these regions are actively restructured

for Bcd and other transcription factors to bind. The increased nucleosome preference of

the more Concentration-Sensitive peaks, combined with the observation that these sites

become occupied by nucleosomes in *bcd* mutants suggests a model where Bcd, either

directly or in combination with cofactors is able to direct chromatin remodeling events,

which may play a significant role in distinguishing concentration-sensitive and -

insensitive targets. Additionally, we find that Bcd-bound regions that are dependent on

either Zld or Bcd for their accessibility are more likely to have a higher nucleosome

preference than regions that are independent of both factors (Figure 4C). This further

suggests that Bcd is able to overcome a high nucleosome barrier in a manner similar to

Zld (Sun et al., 2015) at a subset of its target enhancers.

These effects of chromatin accessibility impact the availability of sequence motifs

for binding Bcd. In wild type embryos, there is a gradual increase in average motif

accessibility from high to low sensitivity, and this difference becomes more pronounced

in *bcd* mutant embryos (Figure 4D), consistent with a role for Bcd in driving changes in

accessibility at more sensitive sites in a concentration dependent manner. This is also

evident at the level of nucleosome organization. Calculating the fraction of motifs that

overlap nucleosomes in either wild type or *bcd* mutant chromatin conformations, we find

that whereas on average across all sensitivity classes 55 ± 2% of Bcd motifs are in

nucleosome-free tracts in wild-type embryos, in *bcd* mutant embryos motifs have lower

overall accessibility and a graded association with nucleosomes that correlates with the

sensitivity classes (41%, 46%, 50%, and 53% of motifs are accessible from high to low

sensitivity). These results indicate that the mechanistic determinants of concentration-

dependent Bcd action likely involve a complex interaction between Bcd, DNA, and

chromatin structure.

A truncated Bcd protein shows reduced binding specifically at concentration-

sensitive target enhancers

A chromatin remodeling activity associated with Bcd has not been previously

described. We hypothesize that Bcd renders its target sites accessible either by

competing with nucleosomes to access its binding sites and bind to DNA at high

concentrations or by recruiting chromatin-remodeling enzymes to accessible motifs and

subsequently driving local nucleosome remodeling to render more sites accessible. We

reasoned that if Bcd can displace nucleosomes simply by competing with them for

access to its binding sites, it should be possible for the Bcd DNA-binding homeodomain

to compete. However, if Bcd instead drives remodeling via recruitment of cofactors, it is

likely that these interactions or activities are carried out through regions of the protein

outside of the DNA binding domain. To distinguish between these two possibilities, we

designed a transgenic GFP-Bcd construct that is truncated downstream of the

homeodomain. We modeled the truncated Bcd protein after the bcd^{085} allele, which was

originally classified as an "intermediate allele" of bcd (Frohnhöfer and Nüsslein-Volhard,

1986) and reported to have weak transcriptional activating activity (Struhl et al., 1989).

The truncation occurs 28 amino acids downstream of the homeodomain (Figure 5A),

and the GFP-tagged protein was therefore expected to bind DNA but lack functions

requiring its C-terminus. The truncated protein (known as GFP-Bcd^{085}) forms a gradient

from the anterior of the embryo, and is expressed at a similar level as a full-length GFP-

Bcd (Figure 5B).

By ChIP-seq, we found that compared to wild-type, GFP-Bcd^{085} binding in the

Concentration-Sensitive I and II enhancer classes was significantly reduced relative to

the Concentration-Sensitive II and Concentraiton-Insensitive classes (p-value < 0.0001

in permutation test with n = 10,000 trials) (Figure 5C). Our ATAC-seq experiments

revealed that these classes have reduced chromatin accessibility in *bcd* mutants

(Figures 3B and 4D). Taken together, these results suggest that Bcd's ability to access

its concentration-sensitive targets is dependent upon activities carried out by domains in

the C-terminus of the protein, likely via recruitment of a cofactor, and that its DNA

binding activity alone is insufficient to drive chromatin accessibility.

Bicoid binding sites confer anterior expression to a posterior target

393   Overall, highly concentration-dependent targets are expressed in the anterior and

are dependent on Bcd for accessibility, while less sensitive targets show more posterior

expression patterns and a greater enrichment for Zld binding sites. An enhancer for

caudal is a Concentration-Insensitive Bcd target and drives expression in the posterior

of the embryo (Figure 6). This enhancer depends on Zld for chromatin accessibility, and

consequently is not functional in *zld* mutants (Supplemental File 1 and Figure 6B). This

supports previous findings that Zld binding contributes to allowing Bcd activation at low

concentrations in posterior nuclei (Xu et al., 2014). Like the *kni* posterior enhancer

(Figure 1A), the *caudal* enhancer is Bcd independent for chromatin accessibility and its

expression boundary shifts anteriorly in *bcd* mutant embryos. We tested whether we

could convert the properties of the *caudal* enhancer from low to high sensitivity by

manipulating DNA motifs. We identified the Zld binding sites in the *caudal* enhancer

sequence and mutated them to Bcd binding sites (Figure 6A). These mutations result in

a shift of *caudal* reporter expression to the anterior of the embryo. Anterior expression

of the mutated reporter is Bcd dependent, as it is lost in *bcd* mutant embryos.

Importantly, the mutant enhancer is functional in *zld* mutant embryos, retaining a distinct

anterior expression domain. In the absence of Zld, the wild type enhancer does not

drive expression. (Figure 6B) By replacing Zld motifs with Bcd motifs, the enhancer

retains functionality, but the spatial domains of expression are now restricted to regions

of high Bcd concentration. These results are consistent with a model where Bcd

operates at high concentrations to confer chromatin accessibility at target sites, thereby

delineating distinct gene expression and chromatin states at specific positions along its

concentration gradient.

DISCUSSION

A model for chromatin accessibility thresholds at Bcd target genes

The results presented here demonstrate that the positional information in the Bcd

gradient is read out as differential binding between Bcd and the *cis*-regulatory regions of

its target genes. The overrepresentation of enhancers for anteriorly expressed target

genes in the more sensitive classes provides support for the classic French flag model,

as their enhancers are only capable of binding Bcd at high levels. However, motif

analysis and *in vitro* EMSA experiments reveal that the differences in binding affinities

that we observe *in vivo* cannot be explained entirely by the sequence of Bcd binding

sites in the target enhancers. Instead we find that a subset of the enhancers in the

concentration-sensitive classes require Bcd for chromatin accessibility. Taken together,

this leads us to model in which the Bcd morphogen establishes concentration

thresholds along the AP axis of the developing embryo by driving opening chromatin

states at high concentrations, thereby gaining access to its most sensitive target

enhancers. At lower concentrations in more posterior nuclei, Bcd is unable to access

these enhancers, and therefore does not bind and activate their transcription. (Figure 7)

In this way, expression of these concentration-sensitive target genes is restricted to

anterior regions of the embryo. The higher density of Bcd binding sites in highly

concentration-sensitive target enhancers (shown in Figure 2B) suggests that these

enhancers may require a larger number of Bcd molecules to be bound at a given time to

keep the enhancers free of nucleosomes and accessible to the additional regulatory

factors. We therefore provide a model for morphogen function in which the

concentration thresholds in the gradient are read out molecularly at the level of

chromatin accessibility, rather than through the strength of binding sites in the target

enhancers.

It is important to note that the discrete sensitivity classes described here were

generated by Bcd binding data, and this binding occurs prior to the activation of target

genes and refinement of their expression domains. In our model Bcd establishes these

initial patterns not by competing with its own target genes, but with default nucleosome

positions in the early embryo. We predict that this initial interaction with chromatin is an

essential event for establishing distinct, positionally defined patterns of gene

expression. The chromatin landscapes established early by Bcd are then elaborated

upon by additional patterning factors, including Bcd target genes themselves, as well as

the repressor gradients proposed by Chen, *et al.* (Chen et al., 2012) Thus, the pre-

transcriptional information presented by Bcd in the form of differential binding states is

refined at the level of gene expression domains both by Bcd and other transcription

factors active in the early embryo.

Relationship of Bicoid and Zelda at Bicoid-bound enhancers

The prominence of the Zld binding motif in the Bcd-bound ChIP peaks and

ATAC-seq in *zld* mutants reveals that Zld also contributes to the accessibility of Bcd

targets in the genome, in part at those targets that are not dependent on Bcd for their

accessibility (61/1,027 peaks are dependent on both Bcd and Zld for accessibility). Zld

is therefore likely to be one component that influences the accessibility and therefore

the apparent *in vivo* affinity of the enhancers that are bound by Bcd but insensitive to its

local concentration (Figure 7). Previous work has suggested that Zld contributes to

activation of target genes at low concentrations of Bcd protein (Xu et al., 2014). That

study, in combination with the work presented here, allows us to predict that

transforming a concentration-sensitive Bcd target enhancer into a Zld dependent

enhancer would increase the accessibility and therefore the sensitivity of that region *in*

vivo. Indeed, Xu, *et al.* have previously demonstrated that adding increasing numbers of

Zld sites to an inactive Bcd-bound enhancer can drive increasingly posterior gene

expression (Xu et al., 2014). The reporter construct used to demonstrate this effect

(HC_45) is identified as a Concentration-Sensitive I target in our study. We posit that

the increase in gene expression from this reporter observed in their work is the result of

increasing the accessibility of the enhancer region.

Alternately, when we replace Zld sites with Bcd sites in an enhancer that drives

posterior expression, the expression domain shifts to the anterior of the embryo. This

demonstrates that without Zld to keep the enhancer open in posterior nuclei, activation

of the reporter gene becomes entirely dependent on Bcd, effectively shifting this

reporter from a concentration-insensitive to concentration-sensitive enhancer. This

finding fits with both previously reported findings and the model proposed in our study.

Namely, that Zld contributes to the accessibility of Bcd target genes throughout the

embryo, while high levels of Bcd can drive accessibility independently and activate gene

expression at a subset of targets in the anterior of the embryo.

The reduced binding by a truncated Bcd protein at the most concentration-

sensitive targets indicates that Bcd does not displace nucleosomes by simply by

competing for binding to genomic targets, but rather that the C-terminus of the Bcd

protein is required for accessing its nucleosome-associated DNA targets. Previous work

has shown that various domains of the Bcd protein are required for interactions with

both co-activators and co-repressors. The N-terminus of Bcd is required for interactions

with components of the Sin3A/HDAC repressor complex, and these interactions are

proposed to play a role in reducing Bcd's transcriptional activation activity (Zhao et al.,

2003; Zhu et al., 2001). Multiple Bcd domains, including the C-terminus, are required for

interaction with CREB-binding protein (CBP), which has histone acetyltransferase

activity (Fu and Ma, 2005; Fu et al., 2004). It is possible that in our truncated Bcd

construct, this interaction with CBP is disrupted. As CBP is thought to play a role in

increasing chromatin accessibility for transcription factors (Chan and La Thangue,

2001), the loss of this interaction could lead to the reduced binding to sensitive targets

that we observe in embryos with truncated Bcd. The enhancers that we have classified

as concentration-insensitive do not depend on Bcd to establish an open chromatin

state. This suggests that these sites are opened by other chromatin remodeling factors,

or are inherently more likely to be nucleosome-free based on their underlying sequence.

It has previously been suggested that transcription factors can compete with

nucleosomes for access to their DNA binding sites (Mirny, 2010; Wang et al., 2011).

This could occur through cooperative binding to nucleosome-associated enhancers: if

one Bcd molecule could gain access to a binding site that was protected by a

nucleosome, it could recruit additional Bcd protein molecules to bind to nearby sites and

occlude nucleosome binding. This cooperativity would require a high concentration of

Bcd protein, fitting with our observation that Bcd influences accessibility more strongly

at high concentrations. However, our experiments with a truncated Bcd protein reveal

that Bcd cannot bind to its most sensitive targets without its C-terminal domains. As

many of the residues that have been implicated in cooperative binding reside in the Bcd

homeodomain (Burz and Hanes, 2001), we would expect this truncated Bcd to bind

cooperatively. This finding therefore supports a model in which Bcd is actively

remodeling chromatin, either directly or more likely by interacting with chromatin

remodeling factors through its C-terminus.

As a maternally supplied factor, Bcd provides one of the first cues to the break

the symmetry of the embryonic body plan. Our results suggest that this symmetry

breaking occurs first at the level of chromatin accessibility, as Bcd drives the opening of

its most concentration-sensitive target enhancers in anterior nuclei. Another maternal

factor, Zld, is proposed to act as a pioneer factor at early embryonic enhancers with a

high intrinsic nucleosome barrier. By binding to these enhancers, Zld depletes them of

nucleosomes and allows patterning transcription factors to bind and activate gene

expression (Sun et al., 2015). We have demonstrated here that Bcd influences the

accessibility primarily of its concentration-sensitive targets, which also exhibit a high

predicted nucleosome barrier (Figure 4B). This raises the possibility that Bcd may be

exhibiting pioneer-like activity at high concentrations, driving accessibility of these sites

prior to transcriptional activation. It is unlikely that Bcd is unique in its ability to influence

the local chromatin accessibility of its targets. Recent work in mouse embryos has

shown that another homeodomain transcription factor, Cdx2, influences the chromatin

accessibility of its targets during posterior axial elongation (Amin et al., 2016). This may

be a common property of developmental transcription factors that must gain early

access their target enhancers while the chromatin state of the genome is being

remodeled during large-scale transitions in the gene regulatory landscape.

EXPERIMENTAL PROCEDURES

Fly stocks and Genetics

bcd mutants refers to embryos derived from bcd^{E1} homozygous mothers. The bcd^{E1} and

$bcd^{E1}\ nos^{L7}\ tsl^4$ stocks were from the Wieschaus/Schüpbach stock collection maintained

at Princeton University. *zld* mutants are embryos derived from $zelda^{294}$ germline clones.

Zelda mutant embryos were generated from the zld^{294} allele (kind gift of Christine

Rushlow) as germline clones as described previously (Blythe and Wieschaus, 2015).

Uniform Bcd and Bcd^{085} transgenes were expressed in a bcd^{E1} mutant background.

Germline clones possessing only positional information from Bicoid were generated by

heat shocking $hsFLP;\ FRT82B\ hb^{FB}\ nos^{BN}\ tsl^4/\ FRT82B\ tsl^4\ Ovo^D$ larvae. Germline

clones lacking Bicoid positional information as well were generated by heat shocking

$hsFLP;\ FRT82B\ bcd^{E1}\ hb^{FB}\ nos^{BN}\ tsl^4/\ FRT82B\ tsl^4\ Ovo^D$ larvae. Embryos from

homozygous $eGFP\text{-}Bcd;\ bcd^{E1}\ nos^{L7}\ tsl^4$ mothers were used in ChIP-seq experiments to

determine the impact of removing other maternal factors on Bcd binding to its targets.

All ATAC-seq experiments were performed in a *His2Av-GFP* (Bloomington) background

to facilitate scoring of nuclear density.

The uBcd and Bcd[085] constructs were injected for site directed transgenesis into

embryos from a *y,w;attp40* stock by Genetic Services (bcd-uBcd and αTub67C-uBcd) or

BestGene (mtrm-uBcd and Bcd[085]) and stable transformant lines were established. The

mutant cad-GAL4 reporter was injected into a *M{vas-int.Dm}ZH-2A, P{CaryP}attP2*

stock by Rainbow Transgenic Flies, Inc.

The uBcd transformants expressed eGFP-tagged Bcd in a graded distribution in the

embryo and RFP in the eyes. Transgenic flies containing the uBcd constructs were

crossed into a *bcd*^{E1} background. To achieve uniform Bcd expression, the uBcd flies

were crossed to a stock expressing a heat shock-inducible *flippase* in a *bcd*^{E1}

background and the resulting larvae were heat shocked at 37°C. Recombination of the

FRT-flanked cassette containing the *bcd* 3'UTR and 3xP3-RFP was scored by a mosaic

loss of RFP expression in the eyes. Mosaic flies were further outcrossed to *bcd*^{E1} and

progeny lacking the bcd 3'UTR were sorted by loss of RFP expression. The resulting

flies produced embryos in which the bcd 3'UTR was replaced by the sqh 3'UTR causing

a uniform distribution of Bcd along the AP axis. (Figure 1 Figure Supplement 1C)

Transgenic Constructs

The uniform Bcd constructs were generated using a pBABR plasmid containing an N-

terminal GFP-tagged *bcd* cDNA in which the *bcd* 3'UTR was replaced by the *sqh* 3'UTR

(pBABR GFP-Bcd3'sqh) (Oliver Grimm, unpublished). This results in a loss of mRNA

localization at the anterior pole of the oocyte. A sequence containing the *bicoid* 3'UTR

and a 3xP3-RFP reporter flanked by FRT sites was synthesized by GenScript and

cloned by Gibson Assembly into the pBABR GFP-Bcd3'sqh plasmid. The FRT-flanked

cassette was inserted between the *bcd* coding sequence and the *sqh* 3'UTR. The *bcd*

promoter was removed by digesting with AgeI and KpnI and replaced with either the

mtrm or the *αTub67C* promoter to generate the *bcd-uBcd*, *mtrm-uBcd*, and *αTub67C-*

uBcd constructs.

The GFP-Bcd[085] truncation was generated from eGFP-Bcd (Gregor et al., 2007a) in

pBlueScript by amplifying with primers to create a stop codon after amino acid 179 as in

the *bcd[085]* hypomorphic EMS allele (Rivera-Pomar et al., 1996). The primers inserted an

AvrII restriction site 3' to the deletion site.

F Primer: 5'-TTGtagCCTAGGCCTGGATGAGAGGCGTGT-3'

R Primer: 5'-TCCAGGCCTAGGctaCAAGCTGGGGGGATC-3'

The plasmid was amplified by PCR and the linear product was digested and ligated to

create the Bcd[085] truncation. The GFP-Bcd[085] construct was digested from pBlueScript

with BamHI and EcoRI and ligated into pBabr.

The wild-type cad-GAL4 reporter (VT010589, coordinates chr2L: 20767347–20768825)

was ordered from the Vienna Drosophila Resource Center (VDRC ID 205848/construct

ID 210589). The mutated *cad* enhancer sequence was synthesized by GenScript,

amplified, and cloned into the pBPGUw vector (Pfeiffer et al., 2008).

Primers, sequences, and plasmids are available upon request.

Western Blots

Live embryos were dechorionated in bleach, rinsed in salt solution (NaCl with TritonX-

100), and embryos at NC14 were sorted under a light microscope and flash frozen on

dry ice. Western blots were performed using a using a rabbit anti-GFP antibody

(Millipore Cat # AB3080P) and mouse anti-tubulin antibody (Sigma Cat # T9026) as a

loading control. For quantification, the GFP band intensities were normalized to α-

tubulin band intensities in each lane. Two biological replicates of 50 embryos were

homogenized in 50μL buffer for each genotype, and 10μL (= 10 embryos) was loaded

per lane.

Western blots were used to generate an estimate of Bcd concentration in each of the

uniform Bcd lines. Drocco, *et al.* used western blots to measure Bcd protein

accumulation in the embryo during development, and calculated the total amount of Bcd

in the embryo at NC14 to be $1.5\pm0.2\times10^8$ molecules (Drocco et al., 2011). Given that the

volume of the nucleus is ~1/10 (or 1/1+9) the volume of the cytoplasm and Bcd

partitions between the nucleus and the cytoplasm at a ratio of ~4:1 (Gregor et al.,

2007a), we can generate a ratio of 4/4+9 or 0.31 for nuclear/cytoplasmic Bcd. Using this

value, we can convert $1.5\pm0.2\times10^8$ molecules/embryo into 4.6×10^7 molecules/nucleus at

NC14. At this stage, there are 6,000 nuclei at the cortex of the embryo, which would be

~7,750 Bcd molecules/nucleus if the Bcd were distributed uniformly. Additionally, optical

measurements estimate a nuclear concentration of Bcd as 8±1 nM and 690 Bcd

molecules at the *hunchback* expression boundary (~48% x/L) at NC14 (Gregor et al.,

2007b). We used these values to generate a conversion factor of 0.011594203

nM/molecule and calculate the approximate nuclear concentrations given below for

each uniform Bcd line. See also Figure 1 Figure Supplement 1D.

Genotype	Expression/WT	Number of Molecules	Nuclear Concentration
bcd>uBcd	0.14	1085	12.58 nM

mtrm>uBcd	1.1	8525	98.84 nM
αTub67C>uBcd	2.7	20925	242.61 nM

Table S1. Estimated nuclear concentrations of Bcd protein in each uniform line.

Immunostaining and Imaging

Embryos of indicated genotypes were collected from 0-4 hour laying cages, and fixed

and stained essentially as described in (Dubuis et al., 2013), with rabbit anti-Bcd,

guinea pig anti-Kni, and rat anti-Btd primary antibodies, followed by fluorophore-

conjugated secondary antibodies Alexa-488 (guinea pig), Alexa-568 (rat), and Alexa-

647 (rabbit) from Invitrogen. Stained embryos were imaged on a Leica SP5 laser-

scanning confocal microscope.

Live Imaging and Image Analysis

Dechorionated embryos of the indicated genotypes were mounted on coverslips

overlaid with halocarbon oil and imaged in the mid-sagittal plane on a Leica SP5 laser

scanning confocal microscope. Image analysis was performed in MATLAB

(http://www.mathworks.com). GFP intensity along the dorsal profile of each embryo was

extracted for each frame of the live movies in nuclear cycle 14. The frame with the

highest overall intensity in each movie was plotted.

Bicoid homeodomain expression and protein purification

A cDNA coding for amino acids 89-154 of the Bicoid protein (including the

homeodomain) as described in (Burz et al., 1998) with a C-terminal HA epitope tag was

cloned into the pET-15b plasmid, which contains an N-terminal 6xHis tag and T7

promoter, to make plasmid pET-15B-BcdHD. Expression was induced in BL21 (DE3)

pLysS *E. coli* cells using 2 mM IPTG. The protein was purified by affinity

chromatography using HisPur Cobalt Resin (Fisher Scientific Cat # 89965) followed by

ion exchange chromatography with SP Sepharose Fast Flow resin (GE Healthcare Cat

#17-0729-01).

EMSAs and K_d Calculations

EMSAs were performed using purified Bicoid homeodomain and biotin-labeled DNA

probes were designed to span ~200 bp in the maximal peak region of Bcd-bound peaks

identified by ChIP and corresponding to previously characterized enhancers. Effective

K_d values for each enhancer probe were calculated using the ratio of total shifted probe

to free probe.

Primers		Sequence (5'->3')
hbP2 probe F	**Forward primer**	/bio/GTCAAGGGATTAGATGGGCA
hbP2 Probe R	**Reverse primer**	/bio/GTCGACTCCTGACCAACGTA
kni post F	**Forward primer**	/bio/AGAAAAAATGAGAACAATGTGAC
kni post R	**Reverse primer**	/bio/AGCCAGCGATTTCGTTACCT
kni ant F	**Forward primer**	/bio/ACAACACCGACCCGTAATCC
kni ant R	**Reverse primer**	/bio/GTCATGTTGGCTAATCTGGC
kr ant F	**Forward primer**	/bio/CAGAAAAGAAAAAGTGTAACGCC

Kr ant R	**Reverse primer**	/bio/GCGAAAAAACGCGTCGCGCT
otd intron F	**Forward primer**	/bio/ATCGTTCCTTGCGGTTTAAT
otd intron R	**Reverse primer**	/bio/AGAACAGGACAAAGGGAATTTAATC
otd early F	**Forward primer**	/bio/CTCGCCTCGCGTGCGACATT
otd early R	**Reverse primer**	/bio/CCTGCGGCAGGACTTCACTT
btd F	**Forward primer**	/bio/ACGAAGTCAAAACTTTTCCA
btd R	**Reverse primer**	/bio/AGCTAAGAGATCTCAACCAAC
gt -3 F	**Forward primer**	/bio/TTACAACTGCCCATTCAGGG
gt -3 R	**Reverse primer**	/bio/GAAGGGCTCGGGTTCGG
gt -10 F	**Forward primer**	/bio/AGATCCAGGCGAGCACTTGA
gt -10 R	**Reverse primer**	/bio/TTAAATTAAAATGTCGCAGGAAGGCG

Table S2: Primer sequences for EMSA probes.

ChIP-seq and Data Analysis

Sample Collection

Drosophila embryos were collected from 0-4 hour laying cages, dechorionated in bleach

and crosslinked in with 180 mL 20% paraformaldehyde in 2 ml PBS + 0.5% Triton X-100

and 6 mL Heptane for 15 minutes. Crosslinking was quenched with 125 mM Glycine in

PBS + 0.5% Triton X-100. Fixed embryos were visually staged and sorted using a

dissecting microscope, and all experimental replicates consisted of 200 embryos in

nuclear cycle 14. Chromatin immunoprecipitation was performed with an anti-GFP

antibody (Millipore) in embryos expressing GFP-tagged Bcd either in a wild-type graded

distribution (*eGFP-Bcd;;bcd^{E1}* and *eGFP-Bcd;;bcd^{E1} nos^{L7} tsl^4*) or uniformly (*GFP-*

uBcd;;bcd^{E1}). Sequencing libraries were prepared using the NEBNext ChIP-seq Library

Prep master mix kit and sequenced as described in (Blythe and Wieschaus, 2015;

Drocco et al., 2011).

Defining a Peak List

Barcode split sequencing files were mapped to *Drosophila melanogaster* genome

assembly BDGP R5/dm3 using Bowtie2 (Gregor et al., 2007a; Langmead and Salzberg,

2012) using default parameters. To generate a conservative, high-confidence list of

Bcd-bound peaks, peaks were called on each replicate of wild-type and uniform Bcd

ChIP-seq data using MACS2 (Gregor et al., 2007b; Zhang et al., 2008) with settings -p

1e-3 --to-large --nomodel --shiftsize 130 for wild-type samples and -p 0.000001 --slocal

5000 --llocal 50000 --keep-dup all for uBcd samples. The most reproducible peaks from

each genotype were selected using an irreproducible discovery rate (IDR) of 1%

(Dubuis et al., 2013; Landt et al., 2012; Li et al., 2011). Given evidence that highly

transcribed (i.e., highly accessible) regions often give false positive results in ChIP

experiments (Burz et al., 1998; Teytelman et al., 2013), we used our ATAC-seq data to

filter our ChIP-seq peaks. We compared the number of CPM-normalized ATAC-seq

reads to ChIP-seq reads in each peak, and performed permutation tests (n = 1,000) to

determine the probability of selecting open regions of the genome at random that had

higher ATAC-seq counts (i.e., regions that were more accessible) than the ATAC-seq

counts in the Bcd ChIP-peaks. We determined that at a ratio of 5.4 ATAC-seq/ChIP-seq

counts, 95% of the ChIP peaks (permutation test p value = 0.05) were no more open

than a random selection of open regions. We filtered out the remaining ChIP peaks with

ATAC/ChIP ratios above 5.4, as these peaks are more likely to correspond to highly

transcribed open regions where most false positive signals can be found. We then

chose the peaks that were common to wild-type and uniform Bcd embryos, which

resulted in a list of 1,027 Bcd-bound peak regions. The number of peaks at each step of

this filtering is shown in Table S3.

Filter Applied	Number of Peaks			
	Wild-Type	tub>uBcd	mtrm>uBcd	bcd>uBcd
MACS2	29,090	15,429	11,812	38,392
IDR	9,815	4,245	1,464	1,329
Euchromatic only	2,319	4,123	1,429	1,257
Common peaks (2/3)		4,126		
ATAC-seq ratios	2,143	2,087		
Common Peaks	1,027			

Table S3. Number Bcd ChIP-seq peaks at each step of filtering.

Comparing Binding Between Uniform Bcd Levels

Mapped BAM files were imported into R as GenomicRanges objects (Lawrence et al.,

2013), filtering out reads with map quality scores below 30. Significant differences

between the uBcd levels were assessed on a pairwise basis using edgeR (Robinson et

al., 2010) in the 1,027 pre-defined peak plus 50,000 additional non-peak noise regions

selected from the dm3 genome.

Data Normalization and Display

Sequencing data was z-score normalized for display in heatmaps. Sequencing read

count coverage was calculated for 10 base pair windows across the genome, and the

mean counts per million reads were determined in each ChIP peak, as well as the

additional noise peaks. Z-scores were computed for each peak using

697
$$z = \frac{CPM - \mu}{\sigma}$$

698   where μ = mean CPM in noise peaks and σ = standard deviation of CPM in noise

699   peaks.

700   **Overlaps with Vienna Tile-GAL4 Enhancers**

	Overlaps		Total Vienna Tiles
	Bcd Peaks	**Vienna Tiles**	
Expressed (all stages)	193	238	3604
Expressed (stage 4-6)	151	163	666
Not Expressed	41	55	4189
Total	234	293	7793

701   Table S4. Number of overlapping Bcd ChIP peaks and Vienna Tile-GAL4 enhancer
702   reporters. The reporters expressed at stage 4-6 that overlapped with more than one Bcd
703   peak were excluded from the plot in Figure 2B.
704

ATAC-seq and Data Analysis

Sample Collection

Live embryos expressing a histone (H2Av)-GFP or RFP construct were individually

staged on an epifluorescence microscope in halocarbon oil. After the onset of nuclear

cycle 14, single embryos were dechorionated in bleach and macerated in cold lysis

buffer at t = 12 minutes into NC14. Samples were pelleted in lysis buffer at 4°C (3000

rpm for 10 minutes), buffer was removed, and the embryo pellet was flash frozen on dry

ice. Frozen pellets were resuspended in Nextera Tagment DNA Buffer + Enzyme and

incubated at 37°C for 30 minutes shaking at 800 rpm. Tagged DNA was purified using a

Qiagen Minelute column and eluted in 10 µL. Barcoded sequencing libraries were

generated by PCR amplifying the purified DNA using the Nextera DNA Sample Prep kit.

Paired-end sequencing was performed on six samples per genotype by the Lewis Sigler

Institute for Integrative Genomics Sequencing Core Facility on an Illumina HiSeq 2500.

Data Processing

Initial processing of the data was performed essentially as described in (Blythe and

Wieschaus 2016, *submitted*). Sequencing files were barcode split and adaptors were

trimmed using TrimGalore. Trimmed reads were mapped to the BDGP R5/dm3 genome

assembly using BWA (Li and Durbin, 2009) with default parameters. Optical and PCR

duplicates were marked using Picard Tools MarkDuplicates

(https://broadinstitute.github.io/picard/). Mapped reads were filtered using samtools (Li

et al., 2009) to remove reads with a map quality score ≤ 30, unmapped reads,

improperly paired reads, and duplicate reads. To distinguish reads corresponding to

open chromatin reads from nucleosome protected reads, the size of the ATAC-seq

fragments were fit to the sum of an exponential and Gaussian distribution as described

in (Buenrostro et al., 2013). We used a fragment size cutoff of ≤ 100 bp to identify

fragments originating from open chromatin. Filtered open reads were imported into R as

GenomicRanges objects.

Peak Calling

Regions of open chromatin at NC14+12 minutes were determined by calling peaks on

the merged open chromatin reads from wild-type replicates using Zinba (Rashid et al.,

2011) with the parameters:

input = 'none', winSize = 300, offset = 50, extension = 65, selectmodel = FALSE,

formula = exp_count ~ exp_cnvwin_log + align_perc, formulaE = exp_count ~

exp_cnvwin_log + align_perc, formulaZ = exp_count ~ align_perc, FDR = TRUE,

threshold = 0.05, winGap = 0, cnvWinSize = 2.5E+4, refinepeaks = TRUE.

Nucleosome Positioning

Nucleosome positioning was determined samples from all genotypes using

NucleoATAC (Schep et al., 2015), with default settings. Peak regions used for

NucleoATAC were open chromatin peaks from Zinba combined with the Bcd ChIP

peaks and widened to 2500 bp centered over the peak maxima. This combined peak list

was then reduced and used as NucleoATAC input. For each genotype, BAM files from

ATAC-seq were merged and used as input for NucleoATAC.

Differential Accessibility between Wild-Type, bcd^{E1}, zld$^-$ Embryos

EdgeR was used to determine significant differences in accessibility between different

genotypes, with an exact test FDR ≤ 0.05 used as the significance cutoff. For edgeR

comparisons, the ATAC-seq peaks called by Zinba from wild-type embryos at NC14+12

minutes were combined with Bcd ChIP-seq peaks resized to 300 bp centered around

the peak summit and 25,000 background regions. The background regions were

generated by extending each of the ATAC open peaks to 10 kb and subtracting them

from the dm3 genome assembly. The remaining non-peak regions were then sampled

randomly 25,000 times and widened to reflect the distribution of sizes in the ATAC-seq

peaks. Bcd- or Zld-dependent peaks were those peaks identified as having reduced

accessibility *bcd*^{E1} or *zld*$^-$ embyos. A summary of the differential accessibility in the

ATAC-seq vs. Bcd ChIP-seq peaks is shown in Table S5.

Peak List	Bcd Dependent	Zld Dependent	Bcd+Zld Dependent	Total
ATAC Open Peaks	326	2,675	206	13,226
Bcd ChIP Peaks	132	402	61	1,027
ATAC+ChIP Common Peaks	121	379	58	855

759 Table S5. Number of peaks dependent on Bcd or Zld for chromatin accessibility. ATAC
760 + ChIP common peaks are peaks that overlap between the Bcd ChIP-seq peaks and
761 the wild-type ATAC-seq open chromatin peaks.
762

To measure differential accessibility of Bcd motifs between wild type and bcdE1 mutant

embryos, the positions of Bcd motifs within ChIP-seq peaks were found, and ATAC-seq

accessibility scores were calculated for the 10 bp window containing the midpoint of

each Bcd motif. Scores for all motifs within a single peak were averaged prior to

plotting. Motifs were found using the "strong" Bcd position weight matrix via the R

function 'matchPWM' in the Biostrings package with an 80% match threshold.

To estimate the overlap between nucleosomes and Bcd motifs, predicted nucleosome

dyad centers from NucleoATAC were widened to 160 bp and motifs overlapping these

intervals were scored as 'nucleosome associated'. Motifs not overlapping widened

nucleosome intervals were scored as 'open'. The fraction of open Bcd motifs per peak

was calculated by dividing the number of open Bcd motifs by the total number of

encoded Bcd motifs over each peak.

Data Normalization and Display

For each genotype, BAM-formatted ATAC-seq reads for each replicate, filtered by

quality and duplicates removed as described above, were merged into a single file.

Coverage was calculated from these BAM files in 10 bp windows in the dm3 genome

assembly. This coverage was then normalized by counts per million reads. The genome

coverage in 2 kb regions flanking Bcd ChIP peaks was then z-score normalized as

described above. The z-score values are displayed as heatmaps of open chromatin

displayed in Figure 3 A and B. Heatmaps are plotted in order of decreasing accessibility

z score in wild-type embryos.

Predicted dyad centers from NucleoATAC were widened to 147 bp to model

nucleosome positions. Occupancy scores from NulcleoATAC in the 147 bp

nucleosomes were computed in 10 bp windows across the dm3 genome assembly.

Occupancy scores overlapping Bcd ChIP peaks are plotted in the heatmaps in Fig 4 A

and B. Nucleosome heatmaps are plotted in the same order as open chromatin

heatmaps, by decreasing accessibility z score in wild-type embryos.

Supplemental File 1 shows the genomic (BDGP Release 5/dm3) coordinates of Bcd-

bound peaks identified by ChIP-seq as described in Table S3. Additional columns

indicate the nearest gene to the peak that shows maternal or zygotic expression in the

embryo (Blythe and Wieschaus, 2015) and classifications of each peak as Bcd or Zld

dependent for accessibility (determined by ATAC-seq) and their sensitivity group

classifications (determined by ChIP-seq). The tileID column gives the name of each

Vienna Tile-GAL4 construct (if any) that overlaps with each peak, and the HC_ID

column indicates reporter constructs from (Chen, et al., 2012) that overlap with each

peak.

***In situ* hybridizations and Cad-GAL4 reporters**

The GAL4 coding sequence was amplified from a genomic DNA preparation generated

from a *Drosophila* stock carrying a GAL4 reporter, using primers 5'-

TGCGATATTTGCCGACTTA-3' and 5'-

<u>TGTAATACGACTCACTATAG</u>GGAACATCCCTGTAGTGATTCCA-3'. The amplified

sequence was used as a template in the MEGAscript T7 Transcription Kit

(ThermoFisher Cat. #AM1334) to with digoxygenin-labeled UTP to generate an anti-

GAL4 RNA probe. *In situ* hybridizations were performed according to standard

protocols.

Author Contributions

C.E.H and S.A.B. designed experiments. C.E.H. performed all experiments and C.E.H.

and S.A.B. performed data analysis. E.F.W. discussed results and performed difficult

meiotic recombination. C.E.H. wrote the first draft of the manuscript and S.A.B. and

E.F.W. contributed to the final version.

Acknowledgements

We thank T. Gregor, E. Gavis, O. Grimm, and members of the Wieschaus and

Schüpbach labs for helpful feedback and discussion, H. Chen, M. Levine and S. Little

for comments on the manuscript, and W. Wang and the staff of the Lewis-Sigler Institute

Sequencing Core Facility. This work was supported by Ruth L. Kirschstein NRSA pre-

doctoral fellowship F31HD082940 (C.E.H.) and postdoctoral fellowship F32HD072653

(S.A.B.).

References

Amin, S., Neijts, R., Simmini, S., van Rooijen, C., Tan, S.C., Kester, L., van
Oudenaarden, A., Creyghton, M.P., and Deschamps, J. (2016). Cdx and T Brachyury
Co-activate Growth Signaling in the Embryonic Axial Progenitor Niche. Cell Rep *17*,
3165–3177.

Berleth, T., Burri, M., Thoma, G., Bopp, D., Richstein, S., Frigerio, G., Noll, M., and
Nüsslein-Volhard, C. (1988). The role of localization of bicoid RNA in organizing the
anterior pattern of the Drosophila embryo. Embo J. *7*, 1749–1756.

Blythe, S.A., and Wieschaus, E.F. (2015). Zygotic Genome Activation Triggers the DNA
Replication Checkpoint at the Midblastula Transition. Cell *160*, 1–13.

Bosch, ten, J.R., Benavides, J.A., and Cline, T.W. (2006). The TAGteam DNA motif
controls the timing of Drosophila pre-blastoderm transcription. Development *133*, 1967–
1977.

Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y., and Greenleaf, W.J. (2013).
Transposition of native chromatin for fast and sensitive epigenomic profiling of open
chromatin, DNA-binding proteins and nucleosome position. Nat. Methods *10*, 1213–
1218.

Buenrostro, J.D., Wu, B., Chang, H.Y., and Greenleaf, W.J. (2015). ATAC-seq: A
Method for Assaying Chromatin Accessibility Genome-Wide. Curr Protoc Mol Biol *109*,
21.29.1–.29.9.

Burz, D.S., and Hanes, S.D. (2001). Isolation of mutations that disrupt cooperative DNA
binding by the Drosophila bicoid protein. J. Mol. Biol. *305*, 219–230.

Burz, D.S., Rivera-Pomar, R., Jäckle, H., and Hanes, S.D. (1998). Cooperative DNA-
binding by Bicoid provides a mechanism for threshold-dependent gene activation in the
Drosophila embryo. Embo J. *17*, 5998–6009.

Chan, H.M., and La Thangue, N.B. (2001). p300/CBP proteins: HATs for transcriptional
bridges and scaffolds. J. Cell. Sci. *114*, 2363–2373.

Chen, H., Xu, Z., Mei, C., Yu, D., and Small, S. (2012). A System of Repressor
Gradients Spatially Organizes the Boundaries of Bicoid-Dependent Target Genes. Cell
149, 618–629.

Crauk, O., and Dostatni, N. (2005). Bicoid Determines Sharp and Precise Target Gene
Expression in the Drosophila Embryo. Current Biology *15*, 1888–1898.

De Renzis, S., Elemento, O., Tavazoie, S., and Wieschaus, E.F. (2007). Unmasking
Activation of the Zygotic Genome Using Chromosomal Deletions in the Drosophila
Embryo. PLoS Biol. *5*, e117.

Driever, W., and Nüsslein-Volhard, C. (1988a). The bicoid protein determines position in
the Drosophila embryo in a concentration-dependent manner. Cell *54*, 95–104.

Driever, W., and Nüsslein-Volhard, C. (1989). The bicoid protein is a positive regulator
of hunchback transcription in the early Drosophila embryo. Nature *337*, 138–143.

Driever, W., Ma, J., Nüsslein-Volhard, C., and Ptashne, M. (1989a). Rescue of bicoid
mutant Drosophila embryos by bicoid fusion proteins containing heterologous activating
sequences. Nature *342*, 149–154.

Driever, W., and Nüsslein-Volhard, C. (1988b). A gradient of bicoid protein in Drosophila
embryos. Cell *54*, 83–93.

Driever, W., Thoma, G., and Nüsslein-Volhard, C. (1989b). Determination of spatial
domains of zygotic gene expression in the Drosophila embryo by the affinity of binding
sites for the bicoid morphogen. Nature *340*, 363–367.

Drocco, J.A., Grimm, O., Tank, D.W., and Wieschaus, E. (2011). Measurement and
Perturbation of Morphogen Lifetime: Effects on Gradient Shape. Biophysical Journal
101, 1807–1815.

Dubuis, J.O., Samanta, R., and Gregor, T. (2013). Accurate measurements of dynamics
and reproducibility in small genetic networks. Molecular Systems Biology *9*, 1–15.

Frohnhöfer, H.G., and Nüsslein-Volhard, C. (1986). Organization of anterior pattern in
the Drosophila embryo by the maternal gene bicoid. Nature *324*, 120–125.

Fu, D., and Ma, J. (2005). Interplay between positive and negative activities that
influence the role of Bicoid in transcription. Nucleic Acids Res. *33*, 3985–3993.

Fu, D., Wen, Y., and Ma, J. (2004). The co-activator CREB-binding protein participates
in enhancer-dependent activities of bicoid. J. Biol. Chem. *279*, 48725–48733.

Gao, Q., and Finkelstein, R. (1998). Targeting gene expression to the head: the
Drosophila orthodenticle gene is a direct target of the Bicoid morphogen. Development
125, 4185–4193.

Goto, T., Macdonald, P., and Maniatis, T. (1989). Early and late periodic patterns of
even skipped expression are controlled by distinct regulatory elements that respond to
different spatial cues. Cell *57*, 413–422.

Gregor, T., Wieschaus, E.F., McGregor, A.P., and Bialek, W. (2007a). Stability and
Nuclear Dynamics of the Bicoid Morphogen Gradient. Cell *130*, 141–152.

Gregor, T., Tank, D.W., Wieschaus, E.F., and Bialek, W. (2007b). Probing the limits to
positional information. Cell *130*, 153–164.

Harrison, M.M., Li, X.-Y., Kaplan, T., Botchan, M.R., and Eisen, M.B. (2011). Zelda

binding in the early Drosophila melanogaster embryo marks regions subsequently
activated at the maternal-to-zygotic transition. PLoS Genet. *7*, e1002266.

Häder, T., Wainwright, D., Shandala, T., Saint, R., Taubert, H., Brönner, G., and Jäckle,
H. (2000). Receptor tyrosine kinase signaling regulates different modes of Groucho-
dependent control of Dorsal. Curr. Biol. *10*, 51–54.

Hoch, M., Seifert, E., and Jäckle, H. (1991). Gene expression mediated by cis-acting
sequences of the Krüppel gene in response to the Drosophila morphogens bicoid and
hunchback. Embo J. *10*, 2267–2278.

Jaeger, J. (2010). The gap gene network. Cell. Mol. Life Sci. *68*, 243–274.

Janody, F., Sturny, R., Catala, F., Desplan, C., and Dostatni, N. (2000). Phosphorylation
of bicoid on MAP-kinase sites: contribution to its interaction with the torso pathway.
Development *127*, 279–289.

Janody, F., Sturny, R., Schaeffer, V., Azou, Y., and Dostatni, N. (2001). Two distinct
domains of Bicoid mediate its transcriptional downregulation by the Torso pathway.
Development *128*, 2281–2290.

Kvon, E.Z., Kazmar, T., Stampfel, G., Yáñez-Cuna, J.O., Pagani, M., Schernhuber, K.,
Dickson, B.J., and Stark, A. (2014). Genome-scale functional characterization of
Drosophila developmental enhancers in vivo. Nature.

Landt, S.G., Marinov, G.K., Kundaje, A., Kheradpour, P., Pauli, F., Batzoglou, S.,
Bernstein, B.E., Bickel, P., Brown, J.B., Cayting, P., et al. (2012). ChIP-seq guidelines
and practices of the ENCODE and modENCODE consortia. Genome Res. *22*, 1813–
1831.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2.
Nat. Methods *9*, 357–359.

Lawrence, M., Huber, W., Pagès, H., Aboyoun, P., Carlson, M., Gentleman, R., Morgan,
M.T., and Carey, V.J. (2013). Software for computing and annotating genomic ranges.
PLoS Comput. Biol. *9*, e1003118.

Levine, M., and Davidson, E.H. (2005). Gene regulatory networks for development.
Proceedings of the National Academy of Sciences *102*, 4936–4942.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R., 1000 Genome Project Data Processing Subgroup (2009). The
Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-
Wheeler transform. Bioinformatics *25*, 1754–1760.

Li, Q., Brown, J.B., Huang, H., and Bickel, P.J. (2011). Measuring Reproducibility of

High-Throughput Experiments. The Annals of Applied Statistics *5*, 1752–1779.

Li, X.-Y., MacArthur, S., Bourgon, R., Nix, D., Pollard, D.A., Iyer, V.N., Hechmer, A.,
Simirenko, L., Stapleton, M., Luengo Hendriks, C.L., et al. (2008). Transcription factors
bind thousands of active and inactive regions in the Drosophila blastoderm. PLoS Biol.
6, e27.

Liu, F., Morrison, A.H., and Gregor, T. (2013). Dynamic interpretation of maternal inputs
by the Drosophila segmentation gene network. Proc Natl Acad Sci U S A *110*, 6724–
6729.

Löhr, U., Chung, H.-R., Beller, M., and Jäckle, H. (2009). Antagonistic action of Bicoid
and the repressor Capicua determines the spatial limits of Drosophila head gene
expression domains. Proceedings of the National Academy of Sciences *106*, 21695–
21700.

Ma, X. (1996). Sequences Outside the Homeodomain of Bicoid Are Required for
Protein-Protein Interaction. Journal of Biological Chemistry *271*, 21660–21665.

Ma, X., Yuan, D., Diepold, K., Scarborough, T., and Ma, J. (1996). The Drosophila
morphogenetic protein Bicoid binds DNA cooperatively. Development *122*, 1195–1206.

MacArthur, S., Li, X.-Y., Li, J., Brown, J.B., Chu, H.C., Zeng, L., Grondona, B.P.,
Hechmer, A., Simirenko, L., Keränen, S.V., et al. (2009). Developmental roles of 21
Drosophila transcription factors are determined by quantitative differences in binding to
an overlapping set of thousands of genomic regions. Genome Biol. *10*, R80.

Mirny, L.A. (2010). Nucleosome-mediated cooperativity between transcription factors.
Proceedings of the National Academy of Sciences *107*, 22534–22539.

Nien, C.-Y., Liang, H.-L., Butcher, S., Sun, Y., Fu, S., Gocha, T., Kirov, N., Manak, J.R.,
and Rushlow, C. (2011). Temporal coordination of gene networks by Zelda in the early
Drosophila embryo. PLoS Genet. *7*, e1002339.

Ochoa-Espinosa, A., Yu, D., Tsirigos, A., Struffi, P., and Small, S. (2009). Anterior-
posterior positional information in the absence of a strong Bicoid gradient. Proceedings
of the National Academy of Sciences *106*, 3823–3828.

Ochoa-Espinosa, A., Yucel, G., Kaplan, L., Pare, A., Pura, N., Oberstein, A.,
Papatsenko, D., and Small, S. (2005). The role of binding site cluster strength in Bicoid-
dependent patterning in Drosophila. Proc Natl Acad Sci U S A *102*, 4960–4965.

Pankratz, M.J., Busch, M., Hoch, M., Seifert, E., and Jäckle, H. (1992). Spatial control of
the gap gene knirps in the Drosophila embryo by posterior morphogen system. Science
255, 986–989.

Perry, M.W., Boettiger, A.N., and Levine, M. (2011). Multiple enhancers ensure
precision of gap gene-expression patterns in the Drosophila embryo. Proc Natl Acad Sci

U S A *108*, 13570–13575.

Pfeiffer, B.D., Jenett, A., Hammonds, A.S., Ngo, T.-T.B., Misra, S., Murphy, C., Scully,
A., Carlson, J.W., Wan, K.H., Laverty, T.R., et al. (2008). Tools for neuroanatomy and
neurogenetics in Drosophila. Proceedings of the National Academy of Sciences *105*,
9715–9720.

Rashid, N.U., Giresi, P.G., Ibrahim, J.G., Sun, W., and Lieb, J.D. (2011). ZINBA
integrates local covariates with DNA-seq data to identify broad and narrow regions of
enrichment, even within amplified genomic regions. Genome Biol. *12*, R67.

Rechsteiner, M., and Rogers, S.W. (1996). PEST sequences and regulation by
proteolysis. Trends Biochem. Sci. *21*, 267–271.

Rivera-Pomar, R., Niessing, D., Schmidt-Ott, U., Gehring, W.J., and Jäckle, H. (1996).
RNA binding and translational suppression by bicoid. Nature *379*, 746–749.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics *26*, 139–140.

Sauer, F., Hansen, S.K., and Tjian, R. (1995). DNA Template and Activator-Coactivator
Requirements for Transcriptional Synergism by Drosophila Bicoid. Science *270*, 1825–
1828.

Schep, A.N., Buenrostro, J.D., Denny, S.K., Schwartz, K., Sherlock, G., and Greenleaf,
W.J. (2015). Structured nucleosome fingerprints enable high-resolution mapping of
chromatin architecture within regulatory regions. Genome Res. *25*, 1757–1770.

Schroeder, M.D., Pearce, M., Fak, J., Fan, H., Unnerstall, U., Emberly, E., Rajewsky,
N., Siggia, E.D., and Gaul, U. (2004). Transcriptional control in the segmentation gene
network of Drosophila. PLoS Biol. *2*, E271.

Segal, E., Fondufe-Mittendorf, Y., Chen, L., Thåström, A., Field, Y., Moore, I.K., Wang,
J.-P.Z., and Widom, J. (2006). A genomic code for nucleosome positioning. Nature *442*,
772–778.

Segal, E., Raveh-Sadka, T., Schroeder, M., Unnerstall, U., and Gaul, U. (2008).
Predicting expression patterns from regulatory sequence in Drosophila segmentation.
Nature *451*, 535–540.

Struhl, G., Struhl, K., and Macdonald, P.M. (1989). The gradient morphogen bicoid is a
concentration-dependent transcriptional activator. Cell *57*, 1259–1273.

Sun, Y., Nien, C.-Y., Chen, K., Liu, H.-Y., Johnston, J., Zeitlinger, J., and Rushlow, C.
(2015). Zelda overcomes the high intrinsic nucleosome barrier at enhancers during
Drosophila zygotic genome activation. Genome Res. *25*, 1703–1714.

Teytelman, L., Thurtle, D.M., Rine, J., and van Oudenaarden, A. (2013). Highly
expressed loci are vulnerable to misleading ChIP localization of multiple unrelated
proteins. Proceedings of the National Academy of Sciences *110*, 18602–18607.

Thomas-Chollier, M., Herrmann, C., Defrance, M., Sand, O., Thieffry, D., and van
Helden, J. (2012). RSAT peak-motifs: motif analysis in full-size ChIP-seq datasets.
Nucleic Acids Res. *40*, e31.

Thomas-Chollier, M., Sand, O., Turatsinze, J.-V., Janky, R., Defrance, M., Vervisch, E.,
Brohée, S., and van Helden, J. (2008). RSAT: regulatory sequence analysis tools.
Nucleic Acids Res. *36*, W119–W127.

Wang, X., Bai, L., Bryant, G.O., and Ptashne, M. (2011). Nucleosomes and the
accessibility problem. Trends Genet. *27*, 487–492.

Wimmer, E.A., Simpson-Brose, M., Cohen, S.M., Desplan, C., and Jäckle, H. (1995).
Trans- and cis-acting requirements for blastodermal expression of the head gap gene
buttonhead. Mech. Dev. *53*, 235–245.

Wolpert, L. (1969). Positional information and the spatial pattern of cellular
differentiation. Journal of Theoretical Biology *25*, 1–47.

Xi, L., Fondufe-Mittendorf, Y., Xia, L., Flatow, J., Widom, J., and Wang, J.-P. (2010).
Predicting nucleosome positioning using a duration Hidden Markov Model. BMC
Bioinformatics *11*, 346.

Xu, Z., Chen, H., Ling, J., Yu, D., Struffi, P., and Small, S. (2014). Impacts of the
ubiquitous factor Zelda on Bicoid-dependent DNA binding and transcription in
Drosophila. Genes Dev. *28*, 608–621.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,
Nusbaum, C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of
ChIP-Seq (MACS). Genome Biol. *9*, R137.

Zhao, C., Fu, D., Dave, V., and Ma, J. (2003). A composite motif of the Drosophila
morphogenetic protein bicoid critical to transcription control. J. Biol. Chem. *278*, 43901–
43909.

Zhu, W., Foehr, M., Jaynes, J.B., and Hanes, S.D. (2001). Drosophila SAP18, a
member of the Sin3/Rpd3 histone deacetylase complex, interacts with Bicoid and
inhibits its activity. Development Genes and Evolution *211*, 109–117.

1033

1034

Figure 1. Uniform Bcd expression specifies cell fates corresponding to levels of expression.

(A) Wild-type, *bcd* null mutant (*bcd^E1^*)*,* and maternal *hunchback, nanos, torso-like (hb nos tsl)* triple mutant and *bcd hb nos tsl* mutant embryos at NC14 immunostained with antibodies against Bcd, Btd, and Kni. Embryos are oriented with anterior to the left. The anterior Kni domain (filled arrow) is absent in *bcd* but restored in *hb nos tsl* embryos, while the posterior stripe (open arrow) shifts anteriorly in in *bcd* but expands posteriorly in *hb nos tsl.* Neither Btd nor Kni exhibit patterned expression in *bcd hb nos tsl*. Images are maximum z-projections and image contrast was adjusted uniformly across the entire image for display. See Figure S1A for quantification of Kni intensity between genotypes.

(B) Expression levels of uniform GFP-Bcd transgenic constructs relative to wild-type Bcd expression. Live embryos were imaged in during NC14, and dorsal profiles were plotted. Error bars are standard error of the mean. For wild-type, n = 23 embryos; bcd-uBcd n = 13; mtrm-uBcd n = 7; and αTub67C -uBcd n = 14. See also Figures S1 and S2 and Table S1.

(C) Immunostaining as (A), for each level of uniform Bcd. Anterior target gene expression is absent at the lowest level. At intermediate (*mtrm*) and high (*αTub67C*) levels of uBcd, anterior expression patterns are expanded and/or duplicated in the posterior, and posterior expression of Kni is absent.

(D) Larval cuticle preparations for the indicated genotypes. Embryos are oriented with anterior at the top. Head structures are indicated with open arrows and tail structures with filled arrows. *αTub67C*>uBcd embryos develop essentially no cuticle tissue, but form only what appear to be anteriorly-derived mouth structures. mtrm>uBcd results in a

duplication of the anterior-most abdominal denticles in the anterior and posterior of the embryo, with no clear terminal structures forming at either end. bcd>uBcd embryos have a normal posterior and all abdominal segments, but no thoracic or head structures. Images of individual embryos were rotated and cropped to exclude nearby embryos and air bubbles.

Figure 2. Bcd-bound regions are classified into groups of increasing sensitivity to Bcd concentration.

(A) ChIP-seq data in Bcd-bound peaks. Data is displayed as a heatmap of z-score normalized ChIp-seq reads, in a 2 kb region centered around each peak. Peaks in each class are arranged in order of decreasing z-scores in wild-type embryos. One peak (peak 549, see Table S6) was not classified, as it showed increasing binding at decreasing Bcd concetrations. Previously characterized enhancers overlapping with each class are indicated at right. Concentration-Insensitive: the posterior stripe enhancers for both *knirps* (Pankratz et al., 1992) and *giant* (Schroeder et al., 2004), and the *Kr* CD1 enhancer (Hoch et al., 1991).

Concentration-Sensitive III: *cap'n'collar* (Schroeder et al., 2004), and *huckebein* (Häder et al., 2000) enhancers.

Concentration-Sensitive II: the *hunchback* P2 proximal (Struhl et al., 1989) and shadow enhancers (Perry et al., 2011), the *even-skipped* stripe 2 enhancer (Goto et al., 1989), an early *paired* enhancer (Ochoa-Espinosa et al., 2005), and an anterior enhancer for *giant* (Schroeder et al., 2004).

Concentration-Sensitive I: *buttonhead* (Wimmer et al., 1995)*, orthodenticle* (Gao and Finkelstein, 1998)*,* and anterior enhancers for both *knirps* and *giant* (Schroeder et al., 2004)*.*

(B) Mean expression patterns of Vienna Tile-GAL4 enhancer reporters overlapping with Bcd peaks in each sensitivity class. Peaks and tiles with more than one overlap were excluded from the plot.

(C) Top DNA motifs discovered by RSAT peak-motifs. The e-value for is a p-value computed from a binomial distribution for a given motif in the dataset, corrected for multiple testing. See Figure S2B for *de novo* motif discovery in each sensitivity class.

(D) Heatmap displaying the enrichment of a given motif in each sensitivity class, relative to the peak list as a whole. P-values were generated from permutation tests (n = 10,000 tests).

A

	Wild-type	bcd^{E1}	bcd>uBcd	mtrm>uBcd	αTub67C>uBcd	zld$^-$
Concentration-Insensitive						
Concentration-Sensitive III						
Concentration-Sensitive II						
Concentration-Sensitive I						

Accessibility Score (max–min)

Concentration-Insensitive
Concentration-Sensitive III
Concentration-Sensitive II
Concentration-Sensitive I

Probability of Nucleosome Occupancy (max–min)

-1 kb peak +1 kb

B

Bcd Dependent Peaks

Wild-type | bcd^{E1} | bcd>uBcd | mtrm>uBcd | αTub67C>uBcd

-1 kb peak +1 kb

Enrichment in Bcd Dependent Peaks

	p-val
Sensitive I	p-val= 6.39e-13
Sensitive II	p-val= 1.01e-10
Sensitive III	p-val= 7.35e-14
Insensitive	p-val= 2.36e-08

log2(Odds Ratio)

Figure 3. Bcd drives chromatin accessibility primarily at concentration-sensitive targets.

(A) Heatmaps showing chromatin accessibility (top) and probability of nucleosome occupancy (bottom) around Bcd-bound peaks from ATAC-seq experiments. Peak regions are arranged by decreasing accessibility in wild-type embryos. bcd^{E1} mutant embryos show a loss of accessibility and increased nucleosome occupancy most strongly at the Concentration-Sensitive I and II peaks. zld^- embryos show reduced accessibility across all sensitivity classes.

(B) Subset of 132 Bcd-bound peaks selected from (A) that become inaccessible in the absence of Bcd. Accessibility at these peaks increases with increasing concentrations of uniform Bcd. Odds ratios and p-values calculated from Fisher's exact test show significant overrepresentation of the Concentration-Sensitive I and II classes in the Bcd-dependent peaks.

A

B

C

D

Figure 4. Bcd sensitivity classes differ in both predicted and observed nucleosome occupancy.

(A) Metaprofiles of nucleosome occupancy in each sensitivity class in wild-type embryos. Background represents random selection of regions outside of Bcd peaks shows a genome-wide average nucleosome probability of ~0.6. Bcd-bound peak regions show reduced nucleosome occupancy compared to unbound regions.

(B) Predicted nucleosome occupancy using NuPoP show higher modeled probability of nucleosome occupancy in Bcd-bound peaks relative to background regions, with higher probability of occupancy at the Concentration-Sensitive I and II classes.

(C) Predicted nucleosome occupancy in peaks dependent on Bcd vs. Zld (n_{Bcd}=132 peaks, n_{Zld}=402 peaks, with n=61 peaks dependent on both Bcd and Zld) for accessibility show higher predicted occupancy than peaks independent of both Bcd and Zld (n=554).

(D) Mean wild type (black) or *bcd⁻* (red) ATAC accessibility scores for Bcd motifs were calculated for each peak and plotted by sensitivity group. Boxplots depict the distribution of accessibility scores for each group in each genotype, and individual data points are shown as points. P-values were calculated by one-sided permutation test and indicate the likelihood in a randomly selected population of observing a difference between means greater than the observed values (p < 1e-6 for Concentration-Sensitive I and II groups, p = 0.001207 for Concentration-Sensitive III, and p = 0.988167 for Concentration-Insensitive).

A

Full-Length Bcd

Bcd085 Truncation

B

C

Figure 5. Bicoid requires C-terminal protein domains to bind to concentration-sensitive targets.

(A) GFP-Bcd[085] construct is truncated within the S/T domain downstream of the homedomain. Wild-type protein domains modified from (Janody et al., 2001) and (Crauk and Dostatni, 2005) . The N-terminus of the protein includes a PRD repeat, followed by the DNA-binding homeodomain (HD) (Berleth et al., 1988). The serine/threonine-rich (S/T) domain is the target of MAPK phosphorylation by the terminal patterning Torso pathway (Janody et al., 2000) and contains a PEST sequence implicated in targeting the protein for degradation (Rechsteiner and Rogers, 1996). The C-terminus contains three domains implicated in transcriptional activation. The glutamine-rich (Q)/OPA and alanine-rich (A) domains are required for interactions with TAFII110 and TAFII60, respectively (Sauer et al., 1995). The acidic (C) domain has been demonstrated to play a role in transcriptional activation in yeast, but is not required for Bicoid activity in the embryo (Driever et al., 1989a).

(B) GFP-Bcd[085] forms a protein gradient comparable to wild-type GFP-Bcd. GFP fluorescence intensity was extracted from dorsal profiles of live embryos. Error bars are standard error of the mean: GFP-Bcd embryos, n = 8; and GFP-Bcd[085] embryos, n = 8.

(C) Boxplots displaying log transformed CPM normalized ChIP-seq data from GFP-Bcd;;bcd[E1] (wild-type) and GFP-Bcd[085];bcd[E1] (Bcd[085]) embryos show significant reduction binding of Bcd[085] in Concentration-Sensitive I and II peaks. P-values were calculated from permutation tests (n = 10,000). See also Figure 2 Figure Supplement 2.

A

○ Zld binding site
○ Bcd binding site

Cad-GAL4

GAL4

Cad (bcd->zld)-GAL4

✗ Zld>Bcd binding site

GAL4

B

	Cad-GAL4	**Cad (bcd->zld)-GAL4**
Wild-type		
bcd		
zld		

Figure 6. Replacing Zld sites with Bcd sites shifts gene expression to the anterior.

(A) Schematic of the Vienna Tile enhancer reporter for *caudal*, containing 5 Zld and 6

Bcd binding sites. The mutated reporter contains 11 Bcd binding sites and no Zld sites.

(B) Expression of the wild-type and mutated reporter in wild-type, *bcd⁻* or *zld⁻* embryos.

Concentration
Insensitive

Concentration
Sensitive

Figure 7. Model for Bicoid function along the AP axis.

Bcd drives accessibility of concentration-sensitive, Bcd-dependent enhancers at high concentrations in anterior nuclei, and these sites are closed in posterior nuclei. concentration-insensitive targets remain accessible in both anterior and posterior nuclei, likely through inputs from other factors such as Zld and more open local chromatin structure with a lower nucleosome preference.

Figure 1 Figure Supplement 1.

A) Quanitification of Knirps intensity in wild-type, triple mutant (*hb nos tsl*) and quadruple mutant (*bcd hb nos tsl*). Bicoid activates patterned expression of Knirps. In embryos in which Bicoid is the only source of maternal patterning information (*hb nos tsl*), a broad domain of Kni is expressed in the posterior of the embryo. In quadruple mutant embryos, a low level of uniform Knirps is expressed ubiquitously, suggesting that Bcd is required for activating expression of *knirps* above a background level. Heat-fixed embryos from wild-type (Oregon-R) mothers, *hunchback nanos torso-like* germline clones and *bicoid hunchback nanos torso-like* germline clones were pooled and immunostained in a single tube with a rat anti-Knirps primary antibody and Alexa-647 rat antibody. Embryos were mounted on a single slide and imaged by confocal microscopy. Representative embryos for each genotype are shown. Fluorescence intensity of Knirps was extracted from dorsal profiles of midsagittal sections of embryos and plotted using MATLAB. Data are fluorescence intensity minus background, and error bars are standard error of the mean for n = 5 wild-type, n = 8 *hb nos tsl,* and n = 6 *bcd hb nos tsl* embryos.

B) Smear plot generated in EdgeR (Robinson et al., 2010) showing the log transformed fold-change in Bcd binding between mutant and wild-type embryos for each Bcd peak, vs. the average log transformed sequencing read counts per million (CPM). Bcd binding shows no significant changes between wild-type and *nos tsl* mutant embryos. Significance was determined using EdgeR to perform a pairwise exact test with a cutoff of FDR ≤ 0.05, comparing binding between *eGFP-Bcd;;bcd^{E1}* and *eGFP-Bcd;; bcd^{E1} hb^{FB} nos^{L7} tsl^{4}* in the 1,027 Bcd peaks.

C) Schematic of the uniform Bcd transgene. The uniform Bcd transgene contains an N-terminal GFP-tagged Bcd driven by the various maternal promoters discussed in the text. Downstream of the *bcd* coding sequence is a cassette containing the endogenous *bcd* 3'UTR and a 3xP3-hsp70 promoter driving promoter of RFP. This cassette is flanked by FRT sites. The *sqh* 3'UTR lies downstream of the FRT cassette. Flies expressing this version of the transgene can be identified by RFP expression in their eyes, and females produce embryos in which Bcd is distributed in a gradient. Males from this transgenic stock are crossed to females expressing a heat shock inducible flippase (*hsFLP)*, and heat shocking the F1 larvae results in recombination and excision of the cassette at the FRT sites, bringing the *sqh* 3'UTR directly downstream of the *bcd* coding sequence. This initially results in mosaic F1 flies with a mosaic graded/uniform Bcd germline. The F1 are further outcrossed to bcd^{E1} mutants and F2 individuals producing embryos with uniform Bcd distributions can be identified by the lack of RFP expression in the eyes.

D) Expression levels of uniform Bcd constructs measured by western blots. Western blots for GFP-Bcd were performed on embryos at NC14. Representative gels and quantifications are shown for the bcd promoter-driven transgene (A), mtrm promoter-driven transgene (B) and α-tub67C promoter-driven transgene (C). In the barplots, band intensities are reported relative to wild-type (GFP-Bcd). All lanes are normalized to an α-tubulin loading control. Error bars are standard deviation between two biological replicates for each sample. MW = molecular weight marker, * = skipped lane.

A

kni anterior enhancer

[Bcd HD]

Free DNA →

Bcd-DNA

kni anterior
Kd: 91.5 ± 4.64 nM
Hill coeff: 2.42 ± 0.261

kni posterior enhancer

[Bcd HD]

Free DNA →

Bcd-DNA

kni posterior
Kd: 82.6 ± 7.03 nM
Hill coeff: 1.49 ± 0.168

B

Legend:
- otd early
- otd intron
- btd
- kni anterior
- gt anterior
- hbP2
- kr
- gt posterior
- kni posterior

Effective Kd (nM)

- Sensitive I
- Sensitive II
- Insensitive

otd early, otd intron, btd, kni anterior, gt anterior, hbP2, Kr, gt posterior, kni posterior

Figure 2 Figure Supplement 1. *In vitro* **binding affinity of target enhancers for Bcd protein is insufficient to explain** *in vivo* **binding behavior.**

(A) Representative gels from EMSAs with *kni* anterior or posterior enhancer sequence used as DNA probe. Binding curves display the log transformed Bcd concentration is plotted vs. ratio of bound to shifted probe (p.bound).

(B) Binding curves for nine EMSA probes show largely overlapping profiles of *in vitro* affinity for Bcd.

(C) Effective K_d measurements for nine EMSA probes do not correspond to *in vivo* behavior of the same DNA sequences. *In vivo* sensitivity classifications determined by ChIP-seq are indicated by color of bars. Error bars are standard error from 2-3 technical replicates per DNA probe.

Figure 2 Figure Supplement 2. Enrichment for binding and motifs of transcription factors in Bcd sensitivity classes.

(A) Heatmap depicting enrichment of Berkeley Drosophila Transcription Network Project (BDTNP) ChIP-chip peaks for AP factors in Bcd-bound sensitivity classes. ChIP data (MacArthur et al., 2009) was downloaded from bdtnp.lbl.gov, and overlap between ChIP peaks for the indicated factors/antibodies and either the Bcd sensitivity classes or ATAC-seq accessibility dependence groups were calculated. One-sided Fisher's exact tests were performed to test for enrichment of a BDTNP ChIP peak set within given Bcd peak class. P-values are plotted as -log10 values, where white indicates non-significant values.

(B) *De novo* motif discovery performed with RSAT as in Figure 2B, for each of the Bcd sensitivity classes individually. The top five enriched motifs are displayed for each sensitivity class.