Psychophysiological Interactions in a Visual Checkerboard Task: Reproducibility, Reliability, and

the Effects of Deconvolution

Xin Di and Bharat B. Biswal[*]

Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ, 07102, USA.

*Corresponding author:

Bharat B. Biswal, PhD

607 Fenster Hall, University Height

Newark, NJ, 07102, USA

bbiswal@yahoo.com

Running head: Reproducibility and reliability of PPI

Abstract:

Psychophysiological interaction (PPI) is a regression based method to study task modulated brain

connectivity. Despite its popularity in functional MRI (fMRI) studies, its reliability and reproducibility

have not been evaluated. We investigated reproducibility and reliability of PPI effects during a simple

visual task, and examined the effect of deconvolution on the PPI results. A large open-access dataset was

analyzed (n = 138), where a visual task was scanned twice with repetition times (TRs) of 645 ms and

1400 ms, respectively. We first replicated our previous results by using the left and right middle occipital

gyrus as seeds. Then ROI-wise (regions of interest) analysis was performed among twenty visual-related

thalamic and cortical regions, and negative PPI effects were found between many ROIs with the posterior

fusiform gyrus as a hub region. Both the seed-based and ROI-wise results were similar between the two

runs and between the two PPI methods with and without deconvolution. The non-deconvolution method

and the short TR run in general had larger effect sizes and greater extents. However, the deconvolution

method performed worse in the 645 ms TR run than the 1400 ms TR run in the voxel-wise analysis.

Given the general similar results between the two methods and the uncertainty of deconvolution, we

suggest that deconvolution may be not necessary for PPI analysis on block-designed data. Lastly,

intraclass correlations between the two runs were much lower for the PPI effects than the activation main

effects, which raise cautions on performing inter-subject correlations and group comparisons on PPI

effects.

Keywords: Reproducibility, reliability, test-retest, psychophysiological interaction, deconvolution.

1. Introduction

Psychophysiological interaction (PPI) is a widely used method to study task related brain functional connectivity changes (Friston et al., 1997). It employed simple regression-based method to model task modulated connectivity effects, thus enabling whole brain exploratory analysis. Therefore, even though there are more sophisticated methods available, e.g. dynamic causal modeling (Friston et al., 2003), PPI is still a valuable method for fMRI data, given that our knowledge on large-scale task related connectivity is still quite limited. Several modifications of the PPI method have been made after it was proposed, including adding a deconvolution step to deal with the asynchrony between task design and fMRI hemodynamic response (Gitelman et al., 2003) and introducing a generalized framework to model more than two experimental conditions (McLaren et al., 2012).

A PPI effect is defined as an interaction between the time series of a brain region (physiological variable) and a (or more) task design variable (psychological variable). Noises of both the physiological and psychological variables go into the interaction term, so that the interaction effect is much noisier than the main effects of task free connectivity (physiological main effect) and task activation (psychological main effect). This makes PPI analysis having lower statistical power than simple connectivity and conventional activation analysis. Since PPI analysis has been increasingly used to study group differences and inter-subjects variability, it is important to evaluate the reproducibility and reliability of the PPI methods (Dubois and Adolphs, 2016; Vul et al., 2009). Voxel-based meta-analysis has been used to examine consistency of PPI results across studies (Di et al., 2017a). However, because the tasks used in different studies varied greatly, the motivation of a meta-analysis on PPI was rather to identify different connectivity that were modulated by different tasks, than to simply identify consistent connectivity cross studies with different tasks (Di et al., 2017a). Nevertheless, the reliability of PPI effect has not been directly examined.

One critical step for the PPI method is to properly deal with the asynchrony between task design and observed blood-oxygen-level dependent (BOLD) signals. An earlier solution is to convolve the

63 psychological variable with hemodynamic response function (HRF). Then the PPI term x^1_{PPI} could be

64 expressed as:

$$x^1_{PPI} = x_{Physio} \cdot (z_{Psych} * hrf) \qquad (1)$$

66 where x_{Physio} represents the physiological variable, z_{Psych} represents the psychological design variable, and

67 * represents convolution operator. However, this calculation is not appropriate if the interaction

68 happened faster than the slow hemodynamic response. Therefore, a deconvolution procedure is required

69 (Gitelman et al., 2003) to find a variable z_{Physio} that:

$$x_{Physio} = z_{Physioi} * hrf \qquad (2)$$

71 If this could be achieved, then the interaction could be calculated at the neuronal level and then convolve

72 with HRF:

$$x^2_{PPI} = (z_{Psych} \cdot z_{Physio}) * hrf \qquad (3)$$

74 We can also put equation (2) to equation (1), so that:

$$x^1_{PPI} = (z_{Psych} * hrf) \cdot (z_{Physio} * hrf) \qquad (4)$$

Mathematically, x^1_{PPI} and x^2_{PPI} are not equivalent. Therefore, deconvolution seems necessary.

Effective deconvolution depends on assumptions such as known HRF and noise characteristics in the

BOLD signals (O'Reilly et al., 2012; Roebroeck et al., 2011). Unfortunately, there are substantial amount

of variability in HRF both across brain regions and across subjects (Handwerker et al., 2004). On the

other hand, if a task design is slower than the hemodynamic response, e.g. a blocked design, the PPI terms

calculated from the above mentioned two methods could be very similar. We have demonstrated that the

PPI results of a block-designed visual task are spatially corresponding very well between the

deconvolution and non-deconvolution PPI methods (Di et al., 2017b). Whether to perform deconvolution

then needs to compromise between the deviation between the PPI terms calculated in different ways and

the uncertainty of deconvolution (Di et al., 2017b). Therefore, it might be better to not perform

deconvolution for an block-designed task, which is actually recommended by FSL (FMRIB Software

Library) (Jenkinson et al., 2012; O'Reilly et al., 2012). For event-related designed task, however,

deconvolution may be still necessary, because the PPI terms calculated from the deconvolution and non-

deconvolution methods may be dramatically different.

We recently demonstrated negative PPI effects (reduced connectivity) between the middle

occipital gyrus to the fusiform gyrus and supplementary motor areas in a simple block-designed

checkerboard task compared with a fixation baseline (Di et al., 2017b). Here, we further analyzed a

larger sample of checkerboard data (n = 138) of two separate runs with two repetition times (TR: 645 ms

and 1400 ms) (Nooner et al., 2012). The aims of the current study are to first evaluate reproducibility and

reliability of PPI effects in the checkerboard task. Additionally, we investigated the impact of PPI

calculation methods on the PPI results and their reproducibility and reliability. We operationally defined

reproducibility as whether previously reported clusters could be observed in the current analysis, and

whether the clusters reported in one run could be observed in the other run. Quantitatively, we utilized

Dice coefficient to quantify overlaps of voxels on thresholded maps (Rombouts et al., 1998; Taylor et al.,

2012). Next, we used intraclass correlation (ICC) to quantify test-retest reliability. Because the short TR

run has about twice the number of time points as the long TR one, we predict that statistical results would

be better for the short TR run compared with the long TR run. In addition, shorter sampling rate may

provide more accurate estimate of hemodynamic response, therefore deconvolution PPI method should

work better for the short TR than the long TR runs.

2. Methods

2.1. Simulations on the correlations between PPI terms

The hemodynamic response is a slow response compared with neuronal events, which can be understood

as a low-pass filter. Intuitively, if a task design is slow enough, e.g. a blocked design, the convolution

with the HRF may not affect PPI calculations much. To directly demonstrate this relationship between

design alternating length and the effect of convolution on PPI calculation, we firstly performed a

simulation. In this simulation, we defined a simple block-designed task with equal on and off periods

with different cycle lengths (from 8 s to 80 s), and a simple event-related design with fixed inter trial

interval of 12 s (Figure 1A). We used a typical sampling rate of 2 s, so that the event-related design could

be expressed as alterations of one time bin (2 s) of a trial and 5 time bins (10 s) of the baseline condition

(The first column in Figure 1A). The remaining columns in Figure 1A show block designs with different

frequencies of repetition. For example, 80 secs cycle means 40-s on and 40-s off of the task condition

related to the baseline. We then simulated the physiological variable of neuronal activities as a Gaussian

variable for 1,000 times. For each design and simulated "neuronal" physiological variable, we calculated

PPI terms using two ways: 1) each variable convolved with the canonical HRF and then the two

convolved variables were multiplied to form a PPI term (corresponding to x^1_{PPI} in equation 4); 2) the two

variables were multiplied and then convolved with the canonical HRF (corresponding to x^2_{PPI} in equation

3). We then calculated the correlations of the PPI terms calculated from the two methods. The code for

this simulation can be found at: https://github.com/dixy0/PPI_correlation_demo .

Figure 1 Simulations of the correlations between PPI terms calculated from deconvolution and non-

deconvolution methods. Panel A illustrates different task designs that were used for the simulation. Each

column represents a task design. E in the x axis represents the event-related design, with 1 time bin (2 s)

of trial condition and 5 time bins (10 s) of baseline condition. The remaining columns show block

designs with different frequencies of repetition. For example, 80 secs cycle means 40-s on and 40-s off of

the task condition related to the baseline. Physiological variables at the neuronal level were generated

using Gaussian random variables for 1,000 times. Penal B shows boxplots of correlations across the

1,000 simulations between PPI terms calculated from two methods: 1) the two simulated variables were

convolved with the HRF and then multiplied to form the PPI term; 2) the two simulated variables were

135   multiplied and then convolved with the HRF.

2.2. FMRI data and task design

We used the checkerboard fMRI data with TRs of 645 ms and 1400 ms from the release 1 of Enhanced

Nathan Kline Institute - Rockland Sample (http://fcon_1000.projects.nitrc.org/indi/enhanced/). 146

subjects' data with age equal or larger than 20 years old were included for analysis. Six subjects' data

were discarded due to large head motion during fMRI scanning in any of the two scans (maximum frame-

wise displacement (FD) (Di and Biswal, 2015) greater than 1.5 mm or 1.5^o). One subject's data were

deleted because of poor coverage of the lower occipital lobe, and another subject's data were deleted

because of failure of coregistration and normalization. The effective number of subjects was 138 (89

females, 45 males, 1 unidentified). The mean age of the sample was 47.8 years (20 to 83 years).

146   The checkerboard task consisted of 20 s fixation block and 20 s flickering checkerboard block

repeated three times. A blank screen was presented after the third checkerboard block until fMRI scan

was complete. The task was scanned for two separate runs with two TRs: 645 ms and 1400 ms,

respectively. For the 645 ms run, 239 or 240 fMRI images were scanned for each subject. The following

parameters were used: TR = 645 ms; TE = 30 ms; flip angle = 60 deg; voxel size = 3 x 3 x 3 mm^3

isotropic; number of slices = 40. For the 1400 ms run, 98 fMRI images were scanned for each subject.

The following parameters were used: TR = 1400 ms; TE = 30 ms; flip angle = 65 deg; voxel size = 2 x 2

x 2 mm^3 isotropic; number of slices = 64. Anatomical T1 images were scanned using MPRAGE

(magnetization-prepared rapid acquisition with gradient echo) sequence with the following parameters:

TR = 1900 ms; TE = 2.52 ms; flip angle = 9 $^{\circ}$; voxel size = 1 x 1 x 1 mm^3 isotropic. More information of

the data can be found in Nooner et al. (Nooner et al., 2012).

2.3. FMRI data analysis

2.3.1. FMRI data preprocessing

Functional MRI data preprocessing and analysis were performed using SPM12 software

(http://www.fil.ion.ucl.ac.uk/spm/) under MATLAB environment (http://www.mathworks.com/). For the

645 ms run, the first 14 images (9 s) were discarded from analysis, resulting in 225 images for each

subject. For the 1400 TR run, the first five images (7 s) were discarded from analysis, resulting in 93

images for each subject. The functional images were motion corrected, and coregistered to subject's

anatomical images. The anatomical images were segmented, and the deformation field images were used

to normalize the functional images into MNI space. The data from the two TR runs were both resliced

and resampled at a spatial resolution of 3 x 3 x 3 mm^3. Lastly, the functional images were smoothed

using a 6 mm full width at half maximum (FWHM) Gaussian kernel.

2.3.2. Activation analysis

We first defined functional ROIs of the visual thalamus and lower visual area by performing general

linear model (GLM) analysis on the checkerboard task. The checkerboard task was modeled as a box-car

function, with 1 representing the checkerboard condition and 0 representing the fixation or blank screen.

The box-car function was convolved with the canonical hemodynamic response function (HRF) to form a

predictor of BOLD responses. Two regressors of the first eigenvariate of BOLD signals in white matter

and cerebrospinal fluid (CSF), and 24 regressors of Friston's autoregressive head motion model (Friston

et al., 1996) were also added in the model as covariates. An implicit high-pass filter of 1/128 Hz was also

implemented in the model. The high-pass filtering is accomplished in SPM by using discrete cosine

transform functions. The effective high-pass filtering cutoffs were then 0.0069 Hz (1 / 145.125 s) for the

645 ms TR run and 0.0077 Hz (1 / 130.2 s) for the 1400 ms TR run. The GLM model was estimated for

each voxel in the brain to identify regions that showed similar patterns of activations as the task design.

The beta maps of task activation were used for group level analysis using a one sample t-test model.

Statistical significant clusters were identified by using cluster level statistics based on random field theory.

Clusters were first identified using a one-tailed t-test at $p < 0.001$, and cluster extent was determined

using false discovery rate (FDR) at $p < 0.05$.

2.3.3. Definition of regions of interest

We performed two types of PPI analyses, voxel-wise analysis using seed regions that were activated by

the checkerboard task and ROI-based analysis among visual thalamus and cortical visual areas

independently defined from other toolbox. In the activation analysis of the current data, the posterior

visual cortex and the posterior portion of the thalamus were robustly activated by the visual checkerboard

stimulation in both TR runs. We therefore defined the left and right middle occipital gyrus (LMOG and

RMOG) and the thalamus as regions of interest (ROIs) based on the activations. To define the ROIs with

proper size, we increase the threshold to $t > 16$ to define the LMOG and RMOG, and made an intersection

between the two runs. The size of LMOG was 222 voxels, and the size of RMOG was 259 voxels.

Thalamus was defined using a threshold of $p < 0.001$, with an intersection between the two runs. Because

the visual thalamus is small, left and right ROIs were combined to form a single thalamus ROI (171

voxels). Different thresholds were chosen to ensure that these ROIs are similar in size. The eigenvariate

of a ROI was extracted with adjustment of effects of no interests (head motion, WM/CSF variables, and

low frequency drifts).

We defined the visual thalamus as the regions that show functional associations with the lateral

visual network in resting-state (Yuan et al., 2016). Cortical visual areas were defined by using

probabilistic cytoarchitectonic maps. These areas include the OC1/OC2 (occipital cortex) (Amunts et al.,

2000), ventral and dorsal OC3 and OC4 (Kujovic et al., 2013; Rottschy et al., 2007), OC5 (Malikovic et

al., 2006), and FG1/FG2 (fusiform gyrus) (Caspers et al., 2013). For the probabilistic maps of these

regions, we first performed a winner-takes-all algorithm to define unique regions of each area, and then

split them into left and right regions. As a result, there are 20 ROIs (left and right thalamus, OC1, OC3,

OC3d, OC3v, OC4d, OC4v, OC5, FG1, and FG2). The eigenvariate of a ROI was extracted with

adjustment of effects of no interests (head motion, WM/CSF variables, and low frequency drifts).

2.3.4. Psychophysiological interaction analysis

PPI analysis was performed using SPM12 with updates 6685. PPI terms were calculated by using both

deconvolution method and non-deconvolution method. For the deconvolution method, the time series of

a seed region was deconvolved with the canonical HRF, multiplied with the centered psychological box-

car function, and convolved back with the HRF to form a predicted PPI time series at hemodynamic

response level. For the non-deconvolution method, the box-car function of psychological design was

convolved with the HRF to form a psychological variable, and it was centered and multiplied with the raw

seed time series. Figure 2 shows examples of PPI terms calculated from the two methods in the two TR

runs.

Figure 2 Examples of PPI terms calculated by the deconvolution and non-deconvolution methods for the

two TR runs.

For voxel-wise PPI analysis, separate GLMs were built for the LMOG, RMOG, and thalamus

seeds, and for the two TR runs. The models included one regressor representing task activation, one

regressor representing the seed time series, the PPI term, and covariates the same as the activation GLMs

described above. Group-level one sample t-test was used on the corresponding PPI effects, to test where

in the brain showed consistent PPI effects with a seed region. For both positive and negative contrasts, a

one-tailed t-test of p < 0.001 was first used to define clusters, and then a FDR cluster threshold of p< 0.05

was used to identify statistical significant clusters. For ROI-wise analysis, PPI GLM models were built

for each of the 20 ROIs, and applied to all other ROIs as a dependent variable. The GLM model included

one psychological variable, one physiological variable, one PPI variable, and one constant term. The

covariates were not included because they have already been regressed out from all ROI time series. PPI

effects were calculated between each pair of ROIs, resulting in a 20 x 20 matrix of beta values for each

subject. The matrices were symmetrized by averaging corresponding upper and lower diagonal elements

(Di et al., 2017b), with a total of 190 (20 x 19 / 2) unique effects. Group-level one-sample t-test was

performed on each element of the matrix. For both positive and negative contrasts, a one-tailed t-test of p

< 0.001 was used to identify significant PPI effects. This threshold was chosen to match with voxel-wise

analysis. We also used FDR correction on the total of 190 effects. And the results are similar to what

using a p < 0.001 threshold. However, FDR depends on the distribution of all tested p values, making it

difficult to compare between two runs. Therefore, we adopted p < 0.001 to report ROI-based PPI results.

2.3.5. Reproducibility and reliability

We operationally define reproducibility as overlaps of supra-threshold clusters. Dice coefficient was used

to quantify reproducibility (Rombouts et al., 1998). Two strategies were used to threshold the maps or

matrix from the two TR runs. First, statistical t maps or t matrices from the two TR runs were

thresholded using a common t value, ranging from 1.7 (approximately corresponds to p < 0.05) to 7.

However, it is possible that the effect sizes in the two TR runs are systematically different, so that using a

same t value could generate very different numbers of supra-threshold voxels or elements in the two runs.

Therefore, we also thresholded t maps or t matrices based on the percentile of t values within a map or

matrix. This could ensure that the numbers of supra-threshold voxels or elements are the same between

the two TR runs.

We operationally define reliability as test-retest reliability between the two TR runs, as quantified

as ICC (Zuo et al., 2010a). Voxel-wise ICC maps or each ROI and ICC matrices across 20 ROIs were

calculated between two TR runs for each PPI method. At each voxel or matrix element, ICC was

calculated from a 138 (subject) by 2 (run) matrix by using a MATLAB function written by Zuo et al. (Zuo

et al., 2010a). Because only voxels that have significant effects might show meaningful reliability, we

displayed histograms of ICCs within significant voxels or elements with reference to those in the whole

brain. For task activations, the significant voxels were determined using intersection of the two TR runs

each thresholded at p < 0.001. For PPI effects of each ROI, the significant voxels were determined using

intersection of the two TR runs and two methods each thresholded at p < 0.01. This slightly liberal

threshold was chosen to ensure enough number of voxels survived in the conjunction of the four scenarios.

The whole brain mask was determined as all voxels in the brain, including WM and CSF.

2.3.6. Coefficient of variation

We calculated coefficient of variation to estimate measurement error of task activations and PPI effects.

Coefficient of variation was calculated in ROIs that showed significant activation effects. Specifically,

the LMOG, RMOG, and thalamus ROIs that were used as seed in the PPI analysis were used to represent

activation effects. For the PPI results, we performed a conjunction analysis of the voxel-wise negative

PPI effects across all the eight contrasts (2 PPI methods x 2 TR runs x 2 seeds) using a threshold of p <

0.01, and identified 27 ROIs that showed common negative PPI effects. Beta values of activations or PPI

effects of these ROIs were extracted. Coefficient of variation was calculated based on the method

assuming the variation is proportional to the mean (Bland and Altman, 1996). It measures within subject

variations (across the two TR runs in the current case) relative to the mean effects of the two runs.

Specifically, coefficient of variation was calculated based on a 138 (subject) x 2 (run) matrix. The beta

values were first logarithmic transformed. Variation was then calculated for each subject, and a square

root of mean variations across subjects was calculated. The resulting value was then transformed back

using an exponential function, and subtracted by 1. The script for calculating coefficient of variation is

available at: https://github.com/dixy0/PPI_correlation_demo. The resulting value represents the

percentage of variation of a measure relative to the mean. Coefficients of variation were calculated on the

LMOG, RMOG, and thalamus ROIs to reflect measurement errors of the task activations, and were

calculated on the 27 ROIs and two ROI seeds to reflect measurement errors of the PPI effects.

3. Results

3.1. Simulations on the correlations between PPI terms

The distributions of PPI correlations for each task design are shown in Figure 1B. For the block designs,

the PPI correlations are a function of block cycle length. With longer design cycle, e.g. greater than 40 s

(20-s on and 20-s off), the correlations of PPI terms could be higher than 0.9. Practically, most of the

block-designed fMRI experiments have longer block cycles than 20-s on and 20-s off. If the block

alterations become faster, the correlation between PPI terms decreased. And for the event-related design,

the mean PPI correlations were below 0.5 and with large variations. This simulation demonstrates that if

a neuronal activity time series is known, using convolved time series to calculate PPI term (i.e. x_{PPI}^1)

could be very similar to what calculated by first multiplying the two variables and then convolving (i.e.

x_{PPI}^2) for typical block designed experiments. In real fMRI data, the "neuronal" physiological variable is

not known, and has to be estimated by using deconvolution. Considering the similarities of the PPI terms

and the caveats of deconvolution, PPI calculations without deconvolution may be a better choice for block

designed experiments. On the other hand, the PPI correlations in the event-related design are much

smaller (r < 0.5, meaning less than 25% of shared variance). So that deconvolution is still a necessary

step for PPI analysis in event-related designed experiments.

3.2. Activations of the checkerboard task

Both TR runs showed highly significant activations in the visual cortex, as well as in the posterior portion

of the thalamus (Figure 3A). The overlaps (Dice coefficients) of thresholded t maps between the two TR

runs were as high as 0.7 (Figure 3B) at most of the shown t range percentile range. And Dice coefficients

went down when only extremely activated voxels were thresholded. The visual cortex regions also

showed high test-retest reliability (ICC greater than 0.7) (Figure 3C). However, the activations of the

thalamus only showed small test-retest reliability around 0.2. The histograms of ICCs in the significant

voxels and in the whole brain are shown on the right of Figure 3C.

Figure 3 A) Activations (*t* maps) of visual checkerboard presentation for the 645 ms TR run (upper) and

1400 ms TR run (lower). The threshold t value corresponds to one-tailed significance at p < 0.001. B)

Overlaps (Dice coefficients) between the two TR runs using t threshold (left) and percentile threshold

(right). C) Test-retest reliability map (intraclass correlations, ICC) of activations between the two runs is

shown on the left, which were thresholded at ICC > 0.2. The histograms of ICC of activations between

the two TR runs in significant voxels and whole brain are shown on the right. The significant voxels

were determined using intersection of the two runs each thresholded at p < 0.001.

3.3. Psychophysiological interactions

The voxel-wise PPI analysis of the LMOG and RMOG seeds conveyed very similar patterns. The PPI

effects of the LMOG seed for the two TRs and two methods are shown in Figure 4. We first observed

that even though spatial extents of PPI effects varied across the two TR runs and two PPI methods, the

negative PPI effects in previously reported regions, i.e. supplementary motor area and higher visual

cortex, could be observed from all four scenarios. The deconvolution method in 645 ms TR run had the

smallest spatial extent and statistical significance, while the non-deconvolution method in 645 ms TR run

had the largest spatial extent and strongest statistical significance. Both methods in TR of 1400 ms

showed similar spatial extent and significance levels. The last row in Figure 4 demonstrates the overlaps

of negative effects in the four scenarios. Similar results were found in the analysis of the RMOG seed

(Figure 5).

Figure 4 Psychophysiological interaction (PPI) results for the left middle occipital gyrus (LMOG) seed

during checkerboard presentation in the two TR (repetition time) runs of 645 ms and TR 1400 ms. The

resulting clusters were thresholded at $p < 0.001$ (approximated $t = 3.15$), with DF (degree of freedom) of

137. The last row illustrates the number of overlapped negative PPI results in the four scenarios.

Numbers on the bottom represent z coordinates in MNI (Montreal Neurology Institute) space.

Figure 5 Psychophysiological interaction (PPI) results for the right middle occipital gyrus (RMOG) seed
during checkerboard presentation in the two TR (repetition time) runs of 645 ms and TR 1400 ms. The
resulting clusters were thresholded at p < 0.001 (approximated t = 3.15), with DF (degree of freedom) of
137. The last row illustrates the number of overlapped negative PPI results in the four scenarios.
Numbers on the bottom represent z coordinates in MNI (Montreal Neurology Institute) space.

The voxel-wise PPI analysis of the thalamus seed only showed significant effects in the 645 TR
run, but with different brain regions with opposite effects (Figure 6). With deconvolution method, the
thalamus seed showed significant positive PPI effects with middle cingulate gyrus, anterior portion of the

thalamus, bilateral anterior insula, basal ganglia, and right fursiform gyrus. Whereas with non-

deconvolution method, the thalamus seed showed significant negative PPI effects with the bilateral

occipital pole regions. There were no consistent results between two TR runs and two methods.

Therefore subsequent analysis was only performed on the LMOG and RMOG seeds.

Figure 6 Psychophysiological interaction (PPI) results for the thalamus seed during checkerboard

presentation in the TR (repetition time) run of 645 ms. There is no significant PPI effects of the thalamus

seed in TR run of 1400 ms. The resulting clusters were thresholded at $p < 0.001$ (approximated t = 3.15),

with DF (degree of freedom) of 137. Numbers on the bottom represent z coordinates in MNI (Montreal

Neurology Institute) space.

We next performed ROI-based PPI analysis among 20 regions of visual thalamus and cortical

visual areas (Figure 7). The 645 ms TR run showed more significant PPI effects than the 1400 ms TR run.

And non-deconvolution method showed more significant PPI effects than the deconvolutio method. A

prominent number of connectivity changes are between the bilateral FG1 regions and other lower level

visual areas ranging from OC1, OC2, to OC4. We performed a conjunction analysis of PPI results across

the four scenarios, and identified five connections with reduced connectivity in checkerboard than in

fixation. The regions and connections are highlighted in Figure 8.

Figure 7 Matrices of psychophysiological interaction (PPI) results among the 20 regions of interest of

visual thalamus and visual cortex for the two TR (repetition time) runs and two methods. The resulting

clusters were thresholded at p < 0.001.

Figure 8 Illustration of consistently reduced connectivity during checkerboard presentation compared

with fixation in the ROI-based (region of interest) psychophysiological interaction (PPI) analysis in the

365   two TR (repetition time) runs and two methods. Numbers on the bottom represent z coordinates in MNI

366   (Montreal Neurology Institute) space.

3.4. Reproducibility of PPI effects

Since we observed similarities of spatial clusters and connectivity between the two TR runs, we next

examined reproducibility of PPI effects by calculating Dice coefficients of thresholded statistical maps or

PPI matrices between the two TR runs (Figure 9). For voxel-wise analysis of both LMOG and RMOG

seeds, when varying t threshold, the non-deconvolution method showed higher level overlap compared

with the deconvolution method (Figure 9A). When thresholding statistical maps with matched number of

surviving voxels, a similar pattern could still be observed that the non-deconvolution method produced

larger overlaps than the deconvolution method (Figure 9B). For the ROI-wise analysis, however, Dice

coefficients were at similar level between two PPI methods at most t and percentile thresholds. But at

very high t threshold or percentile thresholds, the deconvolution method seemed to produce larger

overlaps (higher Dice coefficients) (Figure 9C and 9D).

Figure 9 Dice coefficients of thresholded negative PPI effects between the two TR runs as functions of t

threshold (A) and percentile threshold (B) for the two seeds and two PPI methods. The lowest t used for

calculating overlap is 1.7, which approximately corresponds to p < 0.05. The largest percentile is 80 to

99.9 percentile, which is approximately corresponds to the largest proportions of voxels at p < 0.05.

3.5. Reliability of PPI effects

Lastly, we calculated ICC between the two TR runs to reflect reliability of PPI effects. The voxel-wise

maps of ICC showed that there were typically low reliability in both methods and ROIs, even in the

regions that showed consistent negative PPI effects (Supplementary Figure S1). We then plotted the

histograms of ICCs in voxels from the whole brain (gray lines) and within regions that showed significant

PPI effects (red lines) (Figure 10A through 10D). It turns out that the distributions of ICCs within

significant regions are only slightly different from the distributions of correlations in the whole brain,

with means around 0.07. The distributions of ICCs were not different between deconvolution and non-

deconvolution methods. Similar distributions of ICCs were also found for the ROI-wise analysis (Figure

10E, 10F, and supplementary Figure S2). We found five PPI effects that were consistently significant in

both TR runs and methods. And the ICCs for the five effects were also small and close to zero.

Figure 10 Histograms (normalized) of intraclass correlations of PPI effects between the two TR runs

across the whole brain (gray lines) and in statistically significant voxels (red lines). The significant

voxels were determined using intersection of the two runs and two methods each thresholded at $p < 0.01$.

400   Left and right masks were calculated separately.

3.6. Measurement error

We calculated coefficients of variation (Bland and Altman, 1996) on task activations and PPI effects to

reflect measurement error (Figure 11). The variations of activation in the LMOG and RMOG were about

70% of the mean activation, while the variation of activation in the thalamus was about 270% of the mean

activation (Figure 11A). In contrast, the variations of PPI effects through the 27 ROIs were about 500%

of the mean effects for both the LMOG and RMOG seeds (Figure 11 B and 11C), which indicated much

larger variation of PPI effects compared with activations. The deconvolution and non-deconvolution

methods had similar level of coefficients of variations. But when directly comparing the two methods,

there was a trend that the non-deconvolution method had smaller coefficients of variation than the

deconvolution method in most of the ROIs (Figure 11D and 11E).

Figure 11 Measurement errors as revealed by coefficients of variations (CV) (Bland and Altman, 1996)

for the activation (A) and psychophysiological interaction (PPI) results (B and C). Please notice the

different scales in y axes. D) and E) demonstrate the differences of CV on PPI effects between the

deconvolution and non-deconvolution methods. LMOG, left middle occipital gyrus; RMOG, right middle

417   occipital gyrus; Tha, thalamus.

3.7. Miscellaneous analysis

To gain further insight to the cases of deconvolution failure, we calculated correlations of PPI terms

between deconvolution and non-deconvolution methods for the LMOG and RMOG seeds (Figure 12A).

In both TR runs, the distributions of correlations centered approximately on 0.7, and there were outliers

whose correlations were only 0.2 or 0.3. This is in contrast with the simulation results (Figure 1B, 40 sec

cycle), where the correlations were around 0.9.

Figure 12 A) Histograms of correlations between PPI terms with and without deconvolution across all

subjects from both the LMOG and RMOG ROIs for the two TR runs. B) For the worst case as spotted by

the black arrow in A), we show the raw time series and the time series with deconvolution and re-

convolution with hemodynamic response function.

We identified the worst case in Figure 12A (black arrow indicated), and deconvolved and

reconvovled it with HRF using SPM's method (Figure 12B). The raw and reconvovled signals look

dramatically different, with the reconvolved signal resembling a smoothed version of the original signal.

Smoothness is indeed the case for the SPM version of deconvolution (Gitelman et al., 2003), because it

utilizes regularization to suppress high frequency components of cosine basis functions those were used to

approximate the neuronal level physiological variable. To directly illustrate this point, we performed fast

Fourier transformation on the time series of the RMOG for all the subjects on the raw, deconvolved, and

reconvolved time series for the two TR runs (Figure 13). It could be seen that after deconvolution, high

frequency components have been suppressed in both TR runs. Particularly, there is a black line that

shows higher power between frequencies of 0.2 to 0.4 Hz in the raw data plot of 645 ms TR run, which

coincides to be the outlier observed in Figure 12. The high frequency component was suppressed, so that

the reconvolved signal looks smooth.

Figure 13 Power spectrums of time series from the right middle occipital gyrus seed for each of the 138

subjects for the 645 ms run (upper panels) and 1400 ms run (lower panels). Each line in a plot represents

one subject. Left, middle, and right panels show the power spectrum of the raw, deconvolved, and re-

convolved time series, respectively.

4. Discussion

By analyzing two separate runs of visual checkerboard task from a large sample (n = 138), the current

study first replicated previously reported negative PPI effects between visual cortex and widespread brain

regions, and then showed negative PPI effects among visual areas centered in the bilateral fusiform gyrus.

By comparing results from two separate runs, we showed that group averaged effects were largely

reproducible; however, the inter-subject reliabilities of the PPI effects were typically low. By comparing

the deconvolution and non-deconvolution PPI methods, we demonstrated that the results by the two

methods were in general very similar, but the non-deconvolution produced larger statistical effects and

spatial extents. The non-deconvolution method may reduce inter-subject variations and increase overlaps

of results between the two runs in some circumstances compared with the deconvolution method.

4.1. Functional connectivity during checkerboard stimulation

The voxel-wise analysis of the LMOG and RMOG seeds replicated our previous results which only

analyzed a sub-set of 26 subjects (Di et al., 2017b, 2015). In our previous work (Di et al., 2017b) we

could only identify significant PPI effects using the RMOG seed, while the current study demonstrated

similar PPI effects from both the LMOG and RMOG seeds. Furthermore, we illustrated that the spatial

extent of regions that showed reduced connectivity with the MOG seed could be much larger and

extended to other brain regions such as the insula and bilateral sensorimotor cortex. This further suggests

a higher extent of functional segregation between the visual cortex and other brain systems during such a

simple visual stimulation task compared with the fixation. The current study also extended previous

study by analyzing task modulated connectivity effects among cytoarchitectonically defined visual areas.

Reduced functional connectivity was observed among many visual areas, with the bilateral FG1 as hub

regions. FG1 is the most posterior portion of the fusiform gyrus, which just laid anterior to the occipital

cortex (Caspers et al., 2013). It is thought a transition zone between lower retinotopic visual areas and

higher category specific brain areas, and integrates information from different retinotopic visual areas to

higher category specific brain areas (Caspers et al., 2014). Therefore, it is reasonable to see that the FG1

showed reduced functional connectivity with many lower visual areas in the checkerboard condition,

because the simple stimuli cannot form a meaningful percept of a specific category.

The thalamus is a critical subcortical structure in the brain, which not only relay sensory

information to the cortex, but also thought to mediate corticocortical communications (Guillery and

Sherman, 2002; Saalmann and Kastner, 2011). The PPI analysis of the thalamus, however, did not show

consistent effects in different TR runs, and different methods. It may because that the visual thalamus is

small in size compared with cortical visual areas, and the signals in the thalamus are not reliable enough.

The current results do suggest some reduced connectivity between the visual thalamus to the primary

visual cortex, and increased connectivity between the visual thalamus to the anterior portion of the

thalamus, basal ganglia, and insula. However, the results are weak and unreliable, especially considering

that the current analysis had included 138 subjects.

4.2. Reproducibility and reliability of PPI effects

To our knowledge, the current study is the first one to evaluate reproducibility and reliability on PPI

effects. The current analysis did not only reproduce the results reported previously (Di et al., 2017b), but

also examined the reproducibility between two runs. Although the two runs were scanned using different

parameters, most importantly the temporal and spatial resolutions, the patterns of PPI effects turned out to

be quite similar between the two runs. The run with 645 ms TR seemed to generate larger spatial extent

in the voxel-wise analysis and more statistically significant results in the ROI-wise analysis. This is

consistent with our prediction, because there are more time points in the 645 ms TR run than in the 1400

ms TR run, which could yield higher statistical power. We do notice that in some scenarios, i.e. voxel-

wise analysis with deconvolution, the PPI results in 645 ms TR run had smaller effect size and spatial

extent, which might be due to failure of deconvolution.

On the other hand, the results indicated that inter-subject reliabilities are typically low (around

0.07) no matter which PPI method was used. The low reliability should be compared with those of

simple task activations, which showed reasonably high reliability regardless of the scan length. The

reliability of PPI effects in the current analysis are also much lower than previous reported test-retest

reliabilities on task activations (Plichta et al., 2012; Raemaekers et al., 2007) and resting-state functional

connectivity (Guo et al., 2012; Zuo et al., 2010b). Of course the short scan lengths could be one factor

that explains the low reliability of PPI effects. But it should be also emphasized that the reliability of

higher order interaction effects (i.e. the PPI) should be much lower than the main effects of task

activations and task-free functional connectivity. A scan length that is sufficient for obtaining reliable

task activations may not be necessarily enough to yield reliable task modulated connectivity estimates.

This factor should be taken into account when designing studies on task based connectivity.

4.3. Deconvolution and PPI

The PPI results using both the deconvolution and non-deconvolution methods are in general very similar.

This is consistent with the simulation showing that the PPI term calculated from the convolution then

multiplication method is very similar to the hypothetical PPI term with a known neural activity in a block-

designed task. When comparing the differences of PPI results with these two methods, the non-

deconvolution method seems to be able to generate larger statistical effects and greater spatial extents or

number of significant effects. The non-deconvolution method also increased the Dice coefficients of

thresholded PPI maps between the two TR runs. However, the Dice coefficients of thresholded PPI

matrices between the two TR runs are quite similar between the two PPI methods, and the deconvolution

method may be even benefiting at higher thresholds. These results highlighted the uncertainty of

deconvolution method in PPI analysis.

519   We have shown that the correlations of PPI terms between deconvolution and non-deconvolution

methods may have outliers whose correlations were only 0.2 or 0.3 (Figure 12), which is in contrast with

the simulation results (Figure 1B). The lower correlations of PPI terms from empirical data compared

with the simulations imply that there might be some uncountable variations introduced during the

deconvolution/convolution of real fMRI data. Indeed, deconvolution is rather a practical problem to

recover underlying signals from some recorded measures, than a simple mathematical problem as

depicted in equation 2. In the practical context, measurement noises need to be taken into account in the

deconvolution model. For fMRI, the goal of deconvolution is to recover neuronal activities from

observed BOLD signals, where there are plenty of noises during MRI recording. The deconvolution

should be expressed as follows with an additional error term:

$$x_{Physio} = z_{Physio} * hrf + \varepsilon \qquad (5)$$

In this circumstance, some noises would be removed during deconvolution so that a signal deconvolved

and convolved back with a HRF will no longer be the same as the original signal. The noise

characteristics and regularization methods for recovering z_{Physio} become critical to the success of

deconvolution.

As have been shown in Figure 13, SPM's deconvolution method explicitly suppresses high

frequency components with the intention that the hemodynamic response is slow therefore high frequency

components may represent noises. But this may overly smooth the data and remove useful information in

higher frequency bands, thus making PPI results with the deconvolution method less sensitive than those

with the direct PPI method. This problem may be more severe for short TR data, because there are more

high frequency components in the data. On the other hand, high frequency signals in BOLD have been

increasingly recognized as functionally meaningful (Chen and Glover, 2015; Gohel and Biswal, 2015;

Lewis et al., 2016), and high frequency components may be critical for connectivity dynamics. Given

that multiband imaging technique has made fMRI sampling rate much faster, proper treatment of high

frequency signals may be critical in deconvolution of fMRI signals and connectivity analysis in general.

Given the facts that the two PPI methods can generate similar results for the current block-

designed task and the non-deconvolution method may increase statistical power, we lean toward a

conclusion that the non-deconvolution PPI method may be a better choice for a block-designed task. This

is in line with the recommendation by FSL (O'Reilly et al., 2012). Of course, deconvolution is still

necessary for an event-related task design, because the PPI terms calculated from the convolution then

multiplication method are dramatically different from those calculated from the multiplication then

convolution method (Figure 1). It's also worth mentioning that it has been suggested that the beta series

method (Rissman et al., 2004) might be an alternative method for event-related designed data (Cisler et al.,

2014). Lastly, there are indeed many variety of deconvolution methods (Havlicek et al., 2011; Makni et

al., 2008; Wu et al., 2013), and some of the methods may be more suitable for fMRI signals and PPI

analysis. Systematic comparisons between these different methods are needed in the future.

The current analyses are mostly based on empirical fMRI data. One limitation of empirical

analysis is that there is no known ground truth to compare with. Simulation may be an alternative way to

approach the question. However, development of biological realistic models for task modulated

connectivity is still challenging, so that the deconvolution problem is difficult to study using simulations

at the current stage. In addition, the similarities and differences between PPI results of the deconvolution

and non-deconvolution methods depend on the variability of hemodynamic response in real fMRI data,

which cannot be simply derived from simulations. Therefore, we believe that the current empirical

analysis is suitable for the question of deconvolution.

4.4. Practical implications on PPI analysis

The current study analyzed data from a simple task design with one task condition and one baseline

condition. In real fMRI experiments, however, there are usually more than two conditions. To deal with

multiple conditions, it was recommended that each task condition is modeled separately with respect to all

other conditions (McLaren et al., 2012). In such "generalized PPI" framework, each experimental

condition is modeled as the same way as the checkerboard condition in the current study. It is reasonable

to conclude that the similarities of PPI results with and without deconvolution could be generalized to

experiments with more than two conditions.

Task related functional connectivity as measured by PPI analysis is typically much smaller, in

terms of effect size, reproducibility, and reliability, than simple task activations, and has much larger

measurement error. To ensure enough statistical power and reliability, a larger sample size than typical

activation studies and enough scan length for each subject are necessary. The design for an fMRI task

needs to consider scan length as a critical factor, if the goal of the study is to examine task related

connectivity. To date, it is still largely unknown how long a scan is needed for reliability capture task

related connectivity. We can only get some insights from resting-state connectivity research, where large

scale test-retest datasets are available (Biswal et al., 2010; Zuo et al., 2014). In resting-state literature, it

has been suggested that at least five minutes of scan is needed for reliability estimate functional

connectivity (Birn et al., 2013; Van Dijk et al., 2010). Then at least five minutes of scan length for a

single task condition is needed for task based fMRI. If the PPI effects are going to be compared between

two experimental conditions, which is usually the case for a well-designed cognitive neuroimaging study,

the required scan length would be much longer. Of course, direct examinations of the effect of scan

length on task related connectivity estimates are still needed in future research.

The PPI method takes advantages of the dynamic aspect of the BOLD signals. Therefore, it's

preferable to adopt faster sampling rate to capture temporal dynamics, which may in turn lead to sacrifice

of other aspects of the signals, e.g. spatial resolution. The current results support the idea that shorter TR

may be beneficial for PPI analysis. Of course, faster sampling rate could be accomplished by new

developments of MRI techniques such as multi-band acquisition (Feinberg and Yacoub, 2012). However,

the current results also suggested some pitfalls of using short TR data. The currently used HRF models

and deconvolution method may be not quite suitable for fast TR data, so that the PPI method with

deconvolution may fail in some cases in short TR data. More work is still needed to validate and

optimize models on high speed fMRI data. Of course, high spatial resolution has its own advantage on

mapping small brain structures such as the thalamus. So that the considerations of temporal and spatial

resolutions may also need to take into account the spatial scales of regions of interest.

5. Conclusion

We demonstrated that the deconvolution and non-deconvolution PPI methods generated similar results on

a simple block-designed task. The deconvolution method may be beneficial in terms of statistical power

and reproducibility. Taken together, deconvolution may be not necessary for PPI analysis for block-

designed fMRI data. When using a large sample, group mean PPI effects are reproducible; however,

inter-subject reliabilities of the PPI effects are quite limited. Systematic evaluations on scan length and

reliability may be necessary before studying inter-subject differences or group differences of PPI effects.

Acknowledgement

This research was supported by National Institute of Health grants R01 AG032088 and R01 DA038895.

Reference

Amunts, K., Malikovic, A., Mohlberg, H., Schormann, T., Zilles, K., 2000. Brodmann's Areas 17 and 18

Brought into Stereotaxic Space—Where and How Variable? Neuroimage 11, 66–84.

doi:10.1006/nimg.1999.0516

Birn, R.M., Molloy, E.K., Patriat, R., Parker, T., Meier, T.B., Kirk, G.R., Nair, V.A., Meyerand, M.E.,

Prabhakaran, V., 2013. The effect of scan length on the reliability of resting-state fMRI connectivity

estimates. Neuroimage 83, 550–558. doi:10.1016/j.neuroimage.2013.05.099

Biswal, B.B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S.M., Beckmann, C.F., Adelstein, J.S.,

Buckner, R.L., Colcombe, S., Dogonowski, A.-M., Ernst, M., Fair, D., Hampson, M., Hoptman,

M.J., Hyde, J.S., Kiviniemi, V.J., Kötter, R., Li, S.-J., Lin, C.-P., Lowe, M.J., Mackay, C., Madden,

D.J., Madsen, K.H., Margulies, D.S., Mayberg, H.S., McMahon, K., Monk, C.S., Mostofsky, S.H.,

Nagel, B.J., Pekar, J.J., Peltier, S.J., Petersen, S.E., Riedl, V., Rombouts, S.A.R.B., Rypma, B.,

Schlaggar, B.L., Schmidt, S., Seidler, R.D., Siegle, G.J., Sorg, C., Teng, G.-J., Veijola, J., Villringer,

A., Walter, M., Wang, L., Weng, X.-C., Whitfield-Gabrieli, S., Williamson, P., Windischberger, C.,

Zang, Y.-F., Zhang, H.-Y., Castellanos, F.X., Milham, M.P., 2010. Toward discovery science of

human brain function. Proc. Natl. Acad. Sci. U. S. A. 107, 4734–9. doi:10.1073/pnas.0911855107

Bland, J.M., Altman, D.G., 1996. Statistics Notes: Measurement error proportional to the mean. BMJ 313,

106–106. doi:10.1136/bmj.313.7049.106

Caspers, J., Zilles, K., Amunts, K., Laird, A.R., Fox, P.T., Eickhoff, S.B., 2014. Functional

characterization and differential coactivation patterns of two cytoarchitectonic visual areas on the

human posterior fusiform gyrus. Hum. Brain Mapp. 35, 2754–2767. doi:10.1002/hbm.22364

Caspers, J., Zilles, K., Eickhoff, S.B., Schleicher, A., Mohlberg, H., Amunts, K., 2013.

632   Cytoarchitectonical analysis and probabilistic mapping of two extrastriate areas of the human

633   posterior fusiform gyrus. Brain Struct. Funct. 218, 511–526. doi:10.1007/s00429-012-0411-8

634 Chen, J.E., Glover, G.H., 2015. BOLD fractional contribution to resting-state functional connectivity

635   above 0.1Hz. Neuroimage 107, 207–218. doi:10.1016/j.neuroimage.2014.12.012

636 Cisler, J.M., Bush, K., Steele, J.S., 2014. A comparison of statistical methods for detecting context-

637   modulated functional connectivity in fMRI. Neuroimage 84, 1042–1052.

638   doi:10.1016/j.neuroimage.2013.09.018

639 Di, X., Biswal, B.B., 2015. Characterizations of resting-state modulatory interactions in the human brain.

640   J. Neurophysiol. 114, 2785–96. doi:10.1152/jn.00893.2014

641 Di, X., Fu, Z., Chan, S.C., Hung, Y.S., Biswal, B.B., Zhang, Z., 2015. Task-related functional

642   connectivity dynamics in a block-designed visual experiment. Front. Hum. Neurosci. 9, 1–11.

643   doi:10.3389/fnhum.2015.00543

644 Di, X., Huang, J., Biswal, B.B., 2017a. Task modulated brain connectivity of the amygdala: a meta-

645   analysis of psychophysiological interactions. Brain Struct. Funct. 222, 619–634.

646   doi:10.1007/s00429-016-1239-4

647 Di, X., Reynolds, R.C., Biswal, B.B., 2017b. Imperfect (de)convolution may introduce spurious

648   psychophysiological interactions and how to avoid it. Hum. Brain Mapp. 38, 1723–1740.

649   doi:10.1002/hbm.23413

650 Dubois, J., Adolphs, R., 2016. Building a Science of Individual Differences from fMRI. Trends Cogn. Sci.

651   20, 425–443. doi:10.1016/j.tics.2016.03.014

652 Feinberg, D.A., Yacoub, E., 2012. The rapid development of high speed, resolution and precision in fMRI.

653   Neuroimage 62, 720–725. doi:10.1016/j.neuroimage.2012.01.049

654 Friston, K.J., Buechel, C., Fink, G.R., Morris, J., Rolls, E., Dolan, R.J., 1997. Psychophysiological and

655   modulatory interactions in neuroimaging. Neuroimage 6, 218–29.

656 Friston, K.J., Harrison, L., Penny, W., 2003. Dynamic causal modelling. Neuroimage 19, 1273–1302.

doi:10.1016/S1053-8119(03)00202-7

Friston, K.J., Williams, S., Howard, R., Frackowiak, R.S., Turner, R., 1996. Movement-related effects in

fMRI time-series. Magn. Reson. Med. 35, 346–55. doi:DOI 10.1002/mrm.1910350312

Gitelman, D.R., Penny, W.D., Ashburner, J., Friston, K.J., 2003. Modeling regional and

psychophysiologic interactions in fMRI: the importance of hemodynamic deconvolution.

Neuroimage 19, 200–7.

Gohel, S.R., Biswal, B.B., 2015. Functional Integration Between Brain Regions at Rest Occurs in

Multiple-Frequency Bands. Brain Connect. 5, 23–34. doi:10.1089/brain.2013.0210

Guillery, R.W., Sherman, S.M., 2002. Thalamic Relay Functions and Their Role in Corticocortical

Communication. Neuron 33, 163–175.

Guo, C.C., Kurth, F., Zhou, J., Mayer, E.A., Eickhoff, S.B., Kramer, J.H., Seeley, W.W., 2012. One-year

test–retest reliability of intrinsic connectivity network fMRI in older adults. Neuroimage 61, 1471–

1483. doi:10.1016/j.neuroimage.2012.03.027

Handwerker, D.A., Ollinger, J.M., D'Esposito, M., 2004. Variation of BOLD hemodynamic responses

across subjects and brain regions and their effects on statistical analyses. Neuroimage 21, 1639–

1651. doi:10.1016/j.neuroimage.2003.11.029

Havlicek, M., Friston, K.J., Jan, J., Brazdil, M., Calhoun, V.D., 2011. Dynamic modeling of neuronal

responses in fMRI using cubature Kalman filtering. Neuroimage 56, 2109–2128.

doi:10.1016/j.neuroimage.2011.03.005

Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. FSL. Neuroimage.

doi:10.1016/j.neuroimage.2011.09.015

Kujovic, M., Zilles, K., Malikovic, A., Schleicher, A., Mohlberg, H., Rottschy, C., Eickhoff, S.B.,

Amunts, K., 2013. Cytoarchitectonic mapping of the human dorsal extrastriate cortex. Brain Struct.

Funct. 218, 157–172. doi:10.1007/s00429-012-0390-9

Lewis, L.D., Setsompop, K., Rosen, B.R., Polimeni, J.R., 2016. Fast fMRI can detect oscillatory neural

activity in humans. Proc. Natl. Acad. Sci. 201608117. doi:10.1073/pnas.1608117113

Makni, S., Beckmann, C., Smith, S., Woolrich, M., 2008. Bayesian deconvolution fMRI data using

bilinear dynamical systems. Neuroimage 42, 1381–1396. doi:10.1016/j.neuroimage.2008.05.052

Malikovic, A., Amunts, K., Schleicher, A., Mohlberg, H., Eickhoff, S.B., Wilms, M., Palomero-Gallagher,

N., Armstrong, E., Zilles, K., 2006. Cytoarchitectonic Analysis of the Human Extrastriate Cortex in

the Region of V5/MT+: A Probabilistic, Stereotaxic Map of Area hOc5. Cereb. Cortex 17, 562–574.

doi:10.1093/cercor/bhj181

McLaren, D.G., Ries, M.L., Xu, G., Johnson, S.C., 2012. A generalized form of context-dependent

psychophysiological interactions (gPPI): A comparison to standard approaches. Neuroimage 61,

1277–1286. doi:10.1016/j.neuroimage.2012.03.068

Nooner, K.B., Colcombe, S.J., Tobe, R.H., Mennes, M., Benedict, M.M., Moreno, A.L., Panek, L.J.,

Brown, S., Zavitz, S.T., Li, Q., Sikka, S., Gutman, D., Bangaru, S., Schlachter, R.T., Kamiel, S.M.,

Anwar, A.R., Hinz, C.M., Kaplan, M.S., Rachlin, A.B., Adelsberg, S., Cheung, B., Khanuja, R., Yan,

C., Craddock, C.C., Calhoun, V., Courtney, W., King, M., Wood, D., Cox, C.L., Kelly, A.M.C., Di

Martino, A., Petkova, E., Reiss, P.T., Duan, N., Thomsen, D., Biswal, B., Coffey, B., Hoptman, M.J.,

Javitt, D.C., Pomara, N., Sidtis, J.J., Koplewicz, H.S., Castellanos, F.X., Leventhal, B.L., Milham,

M.P., 2012. The NKI-Rockland Sample: A Model for Accelerating the Pace of Discovery Science in

Psychiatry. Front. Neurosci. 6, 152. doi:10.3389/fnins.2012.00152

O'Reilly, J.X., Woolrich, M.W., Behrens, T.E.J., Smith, S.M., Johansen-Berg, H., 2012. Tools of the

Trade: Psychophysiological Interactions and Functional Connectivity. Soc. Cogn. Affect. Neurosci.

nss055-. doi:10.1093/scan/nss055

Plichta, M.M., Schwarz, A.J., Grimm, O., Morgen, K., Mier, D., Haddad, L., Gerdes, A.B.M., Sauer, C.,

Tost, H., Esslinger, C., Colman, P., Wilson, F., Kirsch, P., Meyer-Lindenberg, A., 2012. Test–retest

reliability of evoked BOLD signals from a cognitive–emotive fMRI test battery. Neuroimage 60,

1746–1758. doi:10.1016/j.neuroimage.2012.01.129

Raemaekers, M., Vink, M., Zandbelt, B., van Wezel, R.J.A., Kahn, R.S., Ramsey, N.F., 2007. Test–retest

reliability of fMRI activation during prosaccades and antisaccades. Neuroimage 36, 532–542.

doi:10.1016/j.neuroimage.2007.03.061

Rissman, J., Gazzaley, A., D'Esposito, M., 2004. Measuring functional connectivity during distinct stages

of a cognitive task. Neuroimage 23, 752–63. doi:10.1016/j.neuroimage.2004.06.035

Roebroeck, A., Formisano, E., Goebel, R., 2011. The identification of interacting networks in the brain

using fMRI: Model selection, causality and deconvolution. Neuroimage 58, 296–302.

doi:10.1016/j.neuroimage.2009.09.036

Rombouts, S. a, Barkhof, F., Hoogenraad, F.G., Sprenger, M., Scheltens, P., 1998. Within-subject

reproducibility of visual activation patterns with functional magnetic resonance imaging using

multislice echo planar imaging. Magn. Reson. Imaging 16, 105–113.

Rottschy, C., Eickhoff, S.B., Schleicher, A., Mohlberg, H., Kujovic, M., Zilles, K., Amunts, K., 2007.

Ventral visual cortex in humans: Cytoarchitectonic mapping of two extrastriate areas. Hum. Brain

Mapp. 28, 1045–1059. doi:10.1002/hbm.20348

Saalmann, Y.B., Kastner, S., 2011. Cognitive and Perceptual Functions of the Visual Thalamus. Neuron

71, 209–223. doi:10.1016/j.neuron.2011.06.027

Taylor, P.A., Gohel, S., Di, X., Walter, M., Biswal, B.B., 2012. Functional covariance networks:

obtaining resting-state networks from intersubject variability. Brain Connect. 2, 203–17.

doi:10.1089/brain.2012.0095

Van Dijk, K.R.A., Hedden, T., Venkataraman, A., Evans, K.C., Lazar, S.W., Buckner, R.L., 2010.

Intrinsic functional connectivity as a tool for human connectomics: theory, properties, and

optimization. J. Neurophysiol. 103, 297–321. doi:10.1152/jn.00783.2009

Vul, E., Harris, C., Winkielman, P., Pashler, H., 2009. Puzzlingly High Correlations in fMRI Studies of

Emotion, Personality, and Social Cognition. Perspect. Psychol. Sci. 4, 274–290. doi:10.1111/j.1745-

6924.2009.01125.x

Wu, G.-R., Liao, W., Stramaglia, S., Ding, J.-R., Chen, H., Marinazzo, D., 2013. A blind deconvolution

approach to recover effective connectivity brain networks from resting state fMRI data. Med. Image

Anal. 17, 365–74. doi:10.1016/j.media.2013.01.003

Yuan, R., Di, X., Taylor, P.A., Gohel, S., Tsai, Y.-H., Biswal, B.B., 2016. Functional topography of the

thalamocortical system in human. Brain Struct. Funct. 221, 1971–1984. doi:10.1007/s00429-015-

1018-7

Zuo, X.-N., Anderson, J.S., Bellec, P., Birn, R.M., Biswal, B.B., Blautzik, J., Breitner, J.C.., Buckner,

R.L., Calhoun, V.D., Castellanos, F.X., Chen, A., Chen, B., Chen, J., Chen, X., Colcombe, S.J.,

Courtney, W., Craddock, R.C., Di Martino, A., Dong, H.-M., Fu, X., Gong, Q., Gorgolewski, K.J.,

Han, Y., He, Y., He, Y., Ho, E., Holmes, A., Hou, X.-H., Huckins, J., Jiang, T., Jiang, Y., Kelley,

W., Kelly, C., King, M., LaConte, S.M., Lainhart, J.E., Lei, X., Li, H.-J., Li, K., Li, K., Lin, Q., Liu,

D., Liu, J., Liu, X., Liu, Y., Lu, G., Lu, J., Luna, B., Luo, J., Lurie, D., Mao, Y., Margulies, D.S.,

Mayer, A.R., Meindl, T., Meyerand, M.E., Nan, W., Nielsen, J.A., O'Connor, D., Paulsen, D.,

Prabhakaran, V., Qi, Z., Qiu, J., Shao, C., Shehzad, Z., Tang, W., Villringer, A., Wang, H., Wang,

K., Wei, D., Wei, G.-X., Weng, X.-C., Wu, X., Xu, T., Yang, N., Yang, Z., Zang, Y.-F., Zhang, L.,

Zhang, Q., Zhang, Z., Zhang, Z., Zhao, K., Zhen, Z., Zhou, Y., Zhu, X.-T., Milham, M.P., 2014. An

open science resource for establishing reliability and reproducibility in functional connectomics. Sci.

Data 1, 140049. doi:10.1038/sdata.2014.49

Zuo, X.-N., Di Martino, A., Kelly, C., Shehzad, Z.E., Gee, D.G., Klein, D.F., Castellanos, F.X., Biswal,

B.B., Milham, M.P., 2010a. The oscillating brain: Complex and reliable. Neuroimage 49, 1432–

1445. doi:10.1016/j.neuroimage.2009.09.037

Zuo, X.-N., Kelly, C., Adelstein, J.S., Klein, D.F., Castellanos, F.X., Milham, M.P., 2010b. Reliable

intrinsic connectivity networks: Test–retest evaluation using ICA and dual regression approach.

Neuroimage 49, 2163–2177. doi:10.1016/j.neuroimage.2009.10.080

Figure legend

Figure 1 Simulations of the correlations between PPI terms calculated from deconvolution and non-deconvolution methods. Panel A illustrates different task designs that were used for the simulation. Each column represents a task design. E in the x axis represents the event-related design, with 1 time bin (2 s) of trial condition and 5 time bins (10 s) of baseline condition. The remaining columns show block designs with different frequencies of repetition. For example, 80 secs cycle means 40-s on and 40-s off of the task condition related to the baseline. Physiological variables at the neuronal level were generated using Gaussian random variables for 1,000 times. Penal B shows boxplots of correlations across the 1,000 simulations between PPI terms calculated from two methods: 1) the two simulated variables were convolved with the HRF and then multiplied to form the PPI term; 2) the two simulated variables were multiplied and then convolved with the HRF.

Figure 2 Examples of PPI terms calculated by the deconvolution and non-deconvolution methods for the two TR runs.

Figure 3 A) Activations (*t* maps) of visual checkerboard presentation for the 645 ms TR run (upper) and 1400 ms TR run (lower). The threshold t value corresponds to one-tailed significance at $p < 0.001$. B) Overlaps (Dice coefficients) between the two TR runs using t threshold (left) and percentile threshold (right). C) Test-retest reliability map (intraclass correlations, ICC) of activations between the two runs is shown on the left, which were thresholded at $ICC > 0.2$. The histograms of ICC of activations between the two TR runs in significant voxels and whole brain are shown on the right. The significant voxels were determined using intersection of the two runs each thresholded at $p < 0.001$.

Figure 4 Psychophysiological interaction (PPI) results for the left middle occipital gyrus (LMOG) seed during checkerboard presentation in the two TR (repetition time) runs of 645 ms and TR 1400 ms. The

resulting clusters were thresholded at p < 0.001 (approximated t = 3.15), with DF (degree of freedom) of

137. The last row illustrates the number of overlapped negative PPI results in the four scenarios.

Numbers on the bottom represent z coordinates in MNI (Montreal Neurology Institute) space.

Figure 5 Psychophysiological interaction (PPI) results for the right middle occipital gyrus (RMOG) seed

during checkerboard presentation in the two TR (repetition time) runs of 645 ms and TR 1400 ms. The

resulting clusters were thresholded at p < 0.001 (approximated t = 3.15), with DF (degree of freedom) of

137. The last row illustrates the number of overlapped negative PPI results in the four scenarios.

Numbers on the bottom represent z coordinates in MNI (Montreal Neurology Institute) space.

Figure 6 Psychophysiological interaction (PPI) results for the thalamus seed during checkerboard

presentation in the TR (repetition time) run of 645 ms. There is no significant PPI effects of the thalamus

seed in TR run of 1400 ms. The resulting clusters were thresholded at p < 0.001 (approximated t = 3.15),

with DF (degree of freedom) of 137. Numbers on the bottom represent z coordinates in MNI (Montreal

Neurology Institute) space.

Figure 7 Matrices of psychophysiological interaction (PPI) results among the 20 regions of interest of

visual thalamus and visual cortex for the two TR (repetition time) runs and two methods. The resulting

clusters were thresholded at p < 0.001.

Figure 8 Illustration of consistently reduced connectivity during checkerboard presentation compared

with fixation in the ROI-based (region of interest) psychophysiological interaction (PPI) analysis in the

two TR (repetition time) runs and two methods. Numbers on the bottom represent z coordinates in MNI

(Montreal Neurology Institute) space.

Figure 9 Dice coefficients of thresholded negative PPI effects between the two TR runs as functions of t

threshold (A) and percentile threshold (B) for the two seeds and two PPI methods. The lowest t used for

calculating overlap is 1.7, which approximately corresponds to $p < 0.05$. The largest percentile is 80 to

99.9 percentile, which is approximately corresponds to the largest proportions of voxels at $p < 0.05$.

Figure 10 Histograms (normalized) of intraclass correlations of PPI effects between the two TR runs

across the whole brain (gray lines) and in statistically significant voxels (red lines). The significant

voxels were determined using intersection of the two runs and two methods each thresholded at $p < 0.01$.

Left and right masks were calculated separately.

Figure 11 Measurement errors as revealed by coefficients of variations (CV) (Bland and Altman, 1996)

for the activation (A) and psychophysiological interaction (PPI) results (B and C). Please notice the

different scales in y axes. D) and E) demonstrate the differences of CV on PPI effects between the

deconvolution and non-deconvolution methods. LMOG, left middle occipital gyrus; RMOG, right middle

occipital gyrus; Tha, thalamus.

Figure 12 A) Histograms of correlations between PPI terms with and without deconvolution across all

subjects from both the LMOG and RMOG ROIs for the two TR runs. B) For the worst case as spotted by

the black arrow in A), we show the raw time series and the time series with deconvolution and re-

convolution with hemodynamic response function.

Figure 13 Power spectrums of time series from the right middle occipital gyrus seed for each of the 138

subjects for the 645 ms run (upper panels) and 1400 ms run (lower panels). Each line in a plot represents

one subject. Left, middle, and right panels show the power spectrum of the raw, deconvolved, and re-

convolved time series, respectively.