Exon-based phylogenomics strengthens the phylogeny of Neotropical cichlids and

identifies remaining conflicting clades (Cichlomorphae: Cichlidae: Cichlinae)

Katriina L. Ilves[1,2*,] Dax Torti[3,4], and Hernán López-Fernández[1]

[1] Department of Natural History, Royal Ontario Museum, 100 Queen's Park, Toronto,

ON M5S 2C6 Canada

[2] Current address: Biology Department, Pace University, 1 Pace Plaza, New York, NY

10038

[3] Donnelly Sequencing Center, University of Toronto, 160 College Street, Toronto, ON

M5S 3E1 Canada

[4] Current address: Ontario Institute for Cancer Research, MaRS Center, 661 University

Avenue, Toronto, ON M5G 0A3 Canada

*Corresponding author at: Biology Department, Pace University, 1 Pace Plaza, New

York, NY 10038; email: katriina.ilves@gmail.com

Abstract

The phenotypic, geographic, and species diversity of cichlid fishes have made them a

group of great interest for studying evolutionary processes. Here we present a targeted-

exon next-generation sequencing approach for investigating the evolutionary

relationships of cichlid fishes (Cichlidae), with a particular focus on the Neotropical

subfamily Cichlinae using a set of 923 primarily single-copy exons designed through

mining of the Nile tilapia (*Oreochromis niloticus*) genome. Sequence capture and

assembly were robust, leading to a complete dataset of 415 exons for 139 species (147

terminals) that consisted of 128 Neotropical species, six African taxa, and five Indo-

Malagasy cichlids. Gene and species trees were calculated using alternative partitioning

schemes and reconstruction methods. In general, all methods yielded similar topologies

to previously hypothesized relationships within the Cichlinae and clarified several

relationships that were previously poorly supported or in conflict. Additional work will

be needed to fully resolve all aspects of Cichlinae phylogeny. Overall, this approach

yielded a well-resolved phylogeny of Neotropical cichlids that will be of utility for future

assessments of the evolutionary and ecological processes within this diverse group of

fishes. Furthermore, the general methodology employed here of exon targeting and

capture should be applicable to any group of organisms with the availability of a

reference genome.

1. Introduction

Neotropical cichlid fishes are rapidly becoming a model to understand the

evolutionary history and biogeography of the exceptionally diverse Neotropical

freshwater fish fauna (e.g. Hulsey and García de León 2005, 2006; López-Fernández et

al. 2013; Arbour and López-Fernández 2014, 2016; McMahan et al. 2013; Říčan et al.

2013; Astudillo-Clavijo et al. 2015; Tagliacollo et al. 2015; Burress 2016). Likewise, the

emergence of Neotropical cichlids as models of adaptive diversification in riverine

environments is starting to provide a meaningful complement to the long-established

studies of adaptive radiation of lacustrine cichlids in Africa (e.g. Fryer and Iles 1972;

Kocher 2004; Wagner et al. 2012; Brawand et al. 2014; Seehausen 2015). Nevertheless,

continuing the evolutionary study of Neotropical cichlids depends on the availability of a

robust phylogenetic framework that allows reliable reconstruction of divergence times,

supports comparative analysis of lineage and phenotype divergence, and clarifies our

understanding of biogeographic history.

Numerous studies have addressed the intergeneric and higher level relationships

of Neotropical cichlids or some of their clades, and a relatively clear phylogenetic

structure for the subfamily has emerged over the last two decades (e.g. Kullander 1998,

Farias et al. 1999, 2000; Hulsey et al. 2004; Concheiro-Pérez et al. 2007; Musilová et al.

2008, 2009; Smith et al. 2008; López-Fernández et al. 2010; Hulsey and Hollingsworth

2010; McMahan et al. 2013; Říčan et al, 2013, 2016). While these analyses have resulted

in an increasingly stable understanding of Cichlinae relationships, a well-established

taxonomy at the tribe level, and a relatively robust set of relationships among genera, a

fully resolved and unambiguously supported phylogeny of Neotropical cichlids has yet to

be achieved. This is particularly true of several basal relationships among genera or

groups of genera within the three main tribes, Geophagini, Cichlasomatini and Heroini

that remain poorly resolved or supported (e.g. López-Fernández et al. 2010; McMahan et

al. 2013; Říčan et al. 2016).

Most analyses of Neotropical cichlid phylogeny have been based on relatively

few loci (usually 10 or less) and often have been heavily informed by mitochondrial data

(e.g. Říčan et al. 2008, Smith et al, 2008, López-Fernández et al. 2010, McMahan et al.

2013; Friedman et al. 2013). These studies are therefore limited in their ability to provide

robust phylogenetic analyses, especially in the light of sequence saturation (e.g. Farias et

al. 2001; López-Fernández et al. 2005), conflicting signal between nuclear and

mitochondrial data (Dornburg et al. 2014; Říčan et al. 2016), and extensive basal short

branches that often receive poor statistical support (López-Fernández et al. 2005, 2010).

Beyond these well-known limitations, the small size of these datasets does not allow the

incorporation of species tree approaches to phylogenetic analyses, and thus, all these

studies are susceptible to producing misleading relationships due to conflict between

gene trees and species trees (e.g. Maddison 2007; Edwards 2009). A recent study by

Říčan et al. (2016) attempted to circumvent some of these potential limitations by using a

large dataset of concatenated Single Nucleotide Polymorphisms (SNPs) derived from

Restriction Enzyme Associated DNA (ddRAD). They obtained a largely well-supported

tree with the largest taxon sampling of Central American cichlids to date, but their

analysis was not able to unambiguously resolve some relationships and was limited to

only one clade within Cichlinae. Moreover, Říčan et al.'s (2016) dataset is not amenable

to be analyzed under coalescent-based methods and therefore is not suitable to identify

potential conflicts in phylogenetic relationships derived from the effects of deep

coalescence (e.g. Edwards 2009, Heled and Drummond 2010).

This latter point is relevant because the Neotropical cichlid phylogeny is plagued

by short basal branches that, in previous work, have been interpreted as evidence of rapid

diversification (López-Fernández et al. 2005, 2010). Concordant with this interpretation,

comparative studies are revealing patterns of rapid early lineage and phenotype

diversification through habitat and diet-related morphological diversification (e.g. López-

Fernández et al. 2013; Arbour and López-Fernández 2014.). These short branches,

however, also represent a problem in phylogenetic reconstruction because rapid

divergence can result in marked incongruence between gene divergence patterns and

species divergence patterns due to population-level phenomena such as deep coalescence

(e.g. Kubatko and Degnan 2007; Edwards 2009). Traditional phylogenetic methods such

as those generally used in Neotropical cichlid analyses assume that loci in the genome

diverge in concert with the species-level divergence of their corresponding species and

that the phylogenetic signal is additive across loci; however, it is widely understood that

genes have their own phylogenetic histories (gene trees) and that those histories do not

always coincide node-to-node with the phylogenetic history of evolving lineages (species

trees) (e.g. Maddison 2007). These incongruences between gene trees and species trees

can be caused by several mechanisms, including hybridization and introgression, but for

phylogenetic purposes the most widespread and potentially problematic is incomplete

lineage sorting (ILS) or deep coalescence (Edwards 2009). From a phylogenetic

reconstruction point of view, the challenge is how to identify the signal of individual

gene divergence that is congruent with species divergence (Heled & Drummond 2010). It

has been repeatedly shown that concatenation of sequences from genes with incongruent

gene trees can produce erroneous topologies with high statistical support (Edwards 2009,

Heled & Drummond 2010, Mendes 2017). In the case of several Neotropical cichlid

clades, short, poorly supported basal branches could result from poor sampling of that

portion of the tree or represent times of divergence during which ILS resulted from fast

species divergence. In the latter case, concatenated phylogenetic analyses would result in

misleading topological reconstructions. Clearly separating these two scenarios, lack of

data versus deep coalescence, may not be entirely possible, but the use of many

independent nuclear loci (to increase gene tree sampling) and comparing concatenation

methods and gene tree methods should help clarifying whether the pattern observed is

due to a real evolutionary process or is the result of lack of data, improper analysis or

both. Until recently, both the availability of methods to analyze species trees and the

technical difficulties associated with identifying and sequencing many nuclear loci made

this type of analysis very difficult or plainly inaccessible. With the advent of massively

parallel sequencing and the development of methods that allow reconstructing

phylogenies using models based on the coalescent, it is now possible to re-evaluate the

phylogeny of Neotropical cichlids using phylogenomic datasets of hundreds of loci along

with methods that correct for the effects of ILS. Despite these promising developments,

however, computational limitations still restrict the options available to analyze truly

phylogenomic datasets (e.g. hundreds or thousands of loci) with relatively large numbers

of terminals and explicit coalescent analyses of sequence alignments.

In this study we used massively parallel sequencing and a custom-designed exon

target-capture toolkit (Ilves and López-Fernández 2014) to generate a phylogenomic

analysis of Neotropical cichlids under both concatenation and summary coalescent

approaches. The main goals of the study were to update currently available hypotheses of

Cichlinae relationships with a much larger dataset than previously available, and to

leverage the phylogenomic dataset to perform coalescent-based analyses that should

account for the possible effects of incomplete lineage sorting on our ability to accurately

reconstruct Cichlinae relationships.

2. Materials and Methods

2.1. Taxon selection

Selection of taxa aimed at including as many lineages of Neotropical cichlids as

possible. Representatives of all recognized genera were included for all tribes except the

Heroini. A recent proliferation of new generic names for taxa in the clade has

taxonomically separated several monophyletic lineages that were previously considered

congenerics (McMahan et al. 2015, Říčan et al. 2016). More importantly, many

previously "orphan" clades variously referred to as '*Cichlasoma' sensu lato* or '*Heros'*

have been formally assigned names. Given that these changes occurred after the dataset

for this study was generated, representatives for some of these new genera are absent, but

our dataset includes members of every clade within Heroini. Moreover, the study revises

the identification of a few lineages of questionable identity in a previous analysis led by

the senior author of this paper (López-Fernández et al. 2010, and see Říčan et al. 2013).

Although those changes did not greatly affect the overall results of those analyses, they

did incorporate additional uncertainty to the resolution of relationships within a clade of

Heroini and, therefore, it is pertinent to correct them.

The dataset used herein closely resembles that of López-Fernández et al. (2010),

and focuses on improving sampling of portions of the tree that were most problematic in

that study in hopes of improving resolution and support for uncertain groupings such as

those among the Cichlasomatini *Acaronia* and *Laetacara* ,as well as some basal

relationships among Geophagini and the heroin clade "amphilophines" sensu López-

Fernández et al. (2010). Taxon sampling also was designed to test the depth of

phylogenetic divergence that could be resolved using the targeted capture probes used in

this study (and see Ilves and López-Fernández 2014). At the most recent variation end of

the spectrum, we included two individuals of several species to test whether the dataset

had enough phylogenetic signal to identify species as monophyletic. We also tested the

ability of the dataset to resolve species-level relationships within genera by including a

comparatively large number of species within the large *Crenicichla-Teleocichla* clade of

Geophagini.

Altogether, the dataset included 147 terminals representing 139 species. Of these,

128 were Neotropical species and six were African taxa, which represent the sister group

of Cichlinae. These taxa include *Heterochromis multidens*, the sister to all African

cichlids, and both the reference sequence and a newly generated set of sequences for the

Nile Tilapia, *Oreochromis niloticus*, the species from which the probes for target capture

were developed (Ilves and López-Fernández 2014). Finally, we included five Indo-

Malagasy cichlids which are sister to the African-Neotropical clade. The Sri Lankan

cichlid *Pseudetroplus suratensis* was used as the outgroup in all analyses. *Pseudetroplus*

and its sister genus, *Etroplus*, are part of the Etroplinae subfamily which forms the sister

group to the rest of the family (e.g. Stiassny 1991, Sparks and Smith 2004, Sparks 2008,

López-Fernández et al. 2010). The complete list of species, museum catalog numbers,

accession numbers (when available) and general locality data are given in Appendix A.

2.2. Library preparation for exon capture and sequencing

DNA extraction procedures followed those from Ilves and López-Fernández

(2014). Exon capture of 923 targets and sequencing was performed at the Donnelly

Sequencing Centre (DSC) at the University of Toronto

(http://dsc.utoronto.ca/dsc/index.html) led by D. Torti, using the probes and general

protocol described in Ilves and López-Fernández (2014). A double-hybridization

procedure of probes to templates was performed, as this was found to significantly

increase yield in previous work (Ilves and López-Fernández 2014). Details about the

probe clean-up, library preparation, and exon capture procedure can be found in

Appendix B. Paired-end sequencing was conducted on an Illumina HiSeq platform.

2.3. Automation of data processing and analysis

189   Multiple custom scripts were used to automate data processing and analysis from

190   the initial step of read quality control to the creation of maximum likelihood phylogenetic

191   gene trees. An overview file that describes each step can be found in Appendix C and

192   subsequent appendices contain the specific script files used to perform each particular

193   task or set of tasks.

195   *2.4. Sequence quality control, read assembly and consensus sequence generation*

196   The general procedures and parameters for the quality control of the sequencing

reads, assembly of the reads into contigs, and generation of consensus sequences for each

exon and species follow those from Ilves and López-Fernández (2014). A custom script

was run that automated read quality control, contig assembly, and consensus sequence

generation (Appendix D). Briefly, the standalone version of PRINSEQ (Schmieder and

Edwards 2011) was used to retain only high-quality reads based on read length and base

quality, bowtie2 version 2.1.0 (Langmead and Salzberg 2012) was used to map the

contigs to a set of reference sequences from the Nile tilapia (*Oreochromis niloticus*)

genome (Appendix E), and SAMtools version 0.1.19-44428cd (Li et al. 2009) was used

to generate consensus sequences of the assembled contigs. Custom scripts were used to

convert the FASTQ files to FASTA files for alignment (Appendix F) and convert all low

quality base calls to "N" and trim terminal "N"s (Appendix G). FASTA sequences were

imported into Geneious version 7.1.8 (http://www.geneious.com, Kearse et al. 2012),

from which only sequences with a minimum of 100 bp were exported for subsequent use

in alignment and phylogenetic analyses.

2.5. Sequence alignment and gene tree and species tree phylogenetic analyses

A custom script was used to combine all sequences for each exon into a single file

(Appendix H). Sequence alignment for each exon was performed with muscle version

3.8.31 (Edgar 2004) using default parameters and was automated using a custom script

(Appendix I - MuscleAlignedScript_144taxa_X600_q20_d10_100bp.txt). Each

alignment was then manually inspected for quality and completeness in Geneious version

7.1.8 (http://www.geneious.com, Kearse et al. 2012). Only alignments that included a

sequence of at least 100 bp for each taxon were retained for phylogenetic analyses. The

32 opsin exon alignments in the target kit were excluded from all analyses because they

represent a family of genes for which duplication and pseudogenization events have been

documented, complicating their use in phylogenetics due to paralogy (e.g. Bowmaker

2008, Weadick et al. 2012).

Coding and non-coding regions of each alignment were inferred from the

annotated reference genome. Although the probes, as originally designed were intended

to target exon-coding regions, analyses of early alignments during the design of the

protocol revealed non-open reading frames for some sequences (see Ilves and López-

Fernández 2014). Additionally, subsequent iterations of annotation in the Nile Tilapia

genome revealed that some regions originally annotated as coding exonic regions actually

corresponded to non-coding fragments. Coding and non-coding regions were identified

and correspondingly separated for analyses in this paper. Bootstrapped gene trees (1000

replicates) for each exon alignment were generated using RAxML version 8.0.10

(Stamatakis 2008) with a GTRGAMMA model and a corresponding partition file with

the codon positions and non-coding regions of the exon. This procedure was automated

with a custom script (Appendix J).

236   ASTRAL-II version 4.8.0 (Mirarab and Warnow 2015) and STAR (Liu et al.

2009) species tree methods with multi-locus bootstrapping (Seo 2008) were run on the

resulting bootstrapped RAxML datasets. ASTRAL-II analyses with 500 bootstrap

replicates were run locally on a desktop Apple® iMac whereas STAR analyses were run

on the STRAW server (Shaw et al. 2013). Because ASTRAL analyses require a

maximum likelihood (ML) tree for each gene in addition to the set of bootstrapped tree

files, RAxML was used to conduct 40 ML searches on each exon alignment (custom

script Appendix K). To assess possible conflicts between species-tree and total-evidence

concatenated analyses, we also performed a Maximum Likelihood analysis of

concatenated 415 loci in RAxML with one thousand bootstrap replicates.

2.6. Computing resources used:

248   Read quality control, mapping, contig assembly, consensus sequence generation,

and sequence alignment and some individual gene alignment RAxML analyses were

performed locally on a desktop Apple iMac (3.5 GHz Intel Quad Core i7 with 32GB

RAM and 3TB hard drive). Some individual gene alignment RAxML analyses were

performed on the GPC supercomputer at the SciNet HPC Consortium (Loken et al.

2010). SciNet is funded by the Canada Foundation for Innovation under the auspices of

Compute Canada, the Government of Ontario, Ontario Research Fund - Research

Excellence, and the University of Toronto.

2.7. Data availability

Alignments for all exons with complete taxon representation of 100 bp or greater

as well as all bootstrapped phylogenetic trees, will be available on Dryad and all other

data files, including raw fastq files, are available from corresponding author K.L. Ilves,

following publication.

3. Results

3.1. Target capture

923 exons are included in the probe set first developed by Ilves and López-

Fernández (2014). 32 exons correspond to opsins, which were excluded from this study *a*

priori due to their histories of gene duplication, which left a total of 891 exons as

potential targets. Enforcing the restrictions of a complete dataset where every species

must have a sequence of at least 100 bp, resulted in a dataset of 428 captured exons. 13 of

these alignments were deemed to be 'poor' after visual inspection due to large blocks of

ambiguous base pair calls (N) present in multiple species, often comprising over 50% of

the total sequence length, thereby resulting in a dataset of 415 exons (~47% of target set)

for phylogenetic analysis. Although the minimum sequence length was set at 100 bp, the

average minimum sequence length relative to the average reference sequence length of

1136 bp was ~62% (SD 23%), indicating that most species had sequence for most of the

length of each target. The final dataset comprised 147 taxa representing 139 species for

415 exons with a total length of 471,448 bp.

3.2. Depth of phylogenetic resolution in the exon-capture dataset

Analyses of multiple individuals within a species consistently resulted in

monophyly at the species level with unambiguous support, as revealed by the

concatenated analyses (Appendix L). These results were consistent at various levels of

divergence, ranging from resequencing of the African reference *Oreochromis niloticus* to

newly sequenced Neotropical species in all major tribes (e.g. Geophagini: *Crenicichla*

sveni, Cichlasomatini: *Krobia petitella*, Heroini: *Symphysodon aequifasciatus*). Both

concatenated and consensus coalescence analyses of species-level divergence within

genera also revealed a large phylogenetic signal that resulted in generally well-resolved

and supported relationships among species within genera (e.g. *Crenicichla*). These results

generally support the notion that the target capture probes used in this study are adequate

to resolve phylogenetic relationships within Cichlidae spanning family to species levels

of divergence.

3.3. Phylogenetic analyses

All analyses, regardless of methods, recovered the expected monophyletic

Neotropical subfamily Cichlinae as sister to a monophyletic African Pseudocrenilabrinae,

both of which in turn are sister to a paraphyletic arrangement of Indian and Malagasy

lineages in the subfamilies Etroplinae and Ptychochrominae (Fig. 1). These relationships

have long been well established through numerous studies of molecular and

morphological datasets and will not be discussed further (Stiassny 1991, Sparks 2004,

2008, Smith et al. 2008, López-Fernández et al. 2010, McMahan et al. 2013, Friedman et

al. 2013, Matschiner et al. 2016). Relationships among the seven recognized tribes of

Neotropical cichlids (Cichlini, Retroculini, Astronotini, Chaetobranchini, Geophagini,

Cichlasomatini and Heroini), were likewise recovered unambiguously across analyses.

Results herein coincide with previous work in placing the Retroculini and Cichlini clade

as sister to the remainder of Cichlinae. The tribes Cichlasomatini and Heroini form a

monophyletic clade sister to a clade formed by Astronotini, Chaetobranchini and

Geophagini. Interestingly, all analyses recovered Astronotini (genus *Astronotus*) as sister

to the Chaetobranchini plus Geophagini clade. This relationship was also found by

Matschiner et al. (2016), but has not been universally recovered in previous analyses:

Smith et al. (2008) found *Astronotus* as sister to all Cichlinae except Cichlini and

Retroculini, López-Fernández et al. (2010) found it as sister to the Cichlasomatini and

Heroini clade, and McMahan et al. (2013) recovered the genus as sister to a clade of

Chaetobranchini and Geophagini. All these studies were based on a limited set of genes

and a large amount of mitochondrial data, suggesting that the much larger dataset used

herein resolves the previously unstable placement of the Astronotini.

In general, intergeneric relationships within each of the main Neotropical tribes, --

Geophagini, Cichlasomatini and Heroini--, were recovered with unambiguous support.

Nevertheless, considerable ambiguity was observed across methods in the placement of

some lineages within these tribes (see Fig. 1). Given the widespread similarity among

topologies, for the remainder of the paper we use the ASTRAL-II coalescent species-tree

(Fig. 1) as a reference because of the superior performance of Astral-II as a tool for

generating coalescent-based consensus species trees from gene trees (e.g. Arcila et al.

2016). Nevertheless, disagreements between this topology and those derived from the

STAR species-tree method and the concatenated super-matrix topology are highlighted

when pertinent (Fig. 2). Later we discuss the potential impact of topological uncertainty

on macroevolutionary analyses and historical biogeographic studies of Neotropical

cichlids.

All analyses coincide in placing the South American lineages as a paraphyletic

arrangement at the base of Heroini, with the genus *Pterophyllum* as the earliest diverging

lineage, followed by a clade of *Hoplarchus* and *Hypselecara* and then by a clade in which

Mesonauta is sister to *Heros*, which is in turn sister to *Symphysodon* and *Uaru*. A similar

arrangement has been found by other studies (e.g. McMahan et al. 2013), but the position

of *Pterophyllum* can be flipped with that of *Hypselecara* and *Hoplarchus* clade (e.g.

López-Fernández et al. 2010). Likewise, all analyses resulted in less than complete

support for the position of *Heros*, but the placement does not vary across trees in this

study. Support for the placement of *Heros* in previous studies (e.g. López-Fernández et

al. 2010) was relatively weak, and despite orders of magnitude increase in the number of

loci used herein, the phylogenomic analyses still result in less than 100% bootstrap

support for the position of the genus. Contrastingly, the current analyses removed all

ambiguity from the previously uncertain placement of the genus *Uaru* (López-Fernández

et al. 2010).

As in previous studies, the remainder of Heroini is comprised by a geographically

non-monophyletic arrangement of South and Central American lineages. Among these, a

clade including the genera *Rocio*, *Tomocichla*, *Herotilapia* and *Astatheros* is sister to the

rest of Heroini. This arrangement is equivalent to and in the same position of the

astatheroines clade sensu Říčan et al. (2016), but it does not include their newly

described genus *Cribroheros*, which was not represented in our dataset. Říčan et al.

(2016) recently split *Astatheros* sensu López-Fernández et al.'s (2010) into *Astatheros*,

Rocio and *Cribroheros*, but phylogenetically there is no incongruence among the groups

in both studies. The astatheroine clade is sequentially followed by the genus

Australoheros as sister to all other heroins, an arrangement identical with that found by

Říčan et al's (2016) analysis based on concatenated single nucleotide polymorphisms

from a restriction site associated DNA (RAD) dataset. This result differs from that of

López-Fernández et al. 2010 because the only species of *Australoheros* in that study was

inadvertently switched with that of the unrelated *Cryptoheros nanoluteus* (see Appendix

A for details, and Říčan et al. 2013). McMahan et al. (2013) also find a different

placement for *Australoheros*.

Our analyses unambiguously find a monophyletic clade of purely Central

American taxa that corresponds with the amphilophines of Říčan et al. (2016), even

though our analysis did not include their genera *Cryptoheros*, *Talamancaheros* and

Isthmoheros. Despite this correspondence, however, our analyses produced different

internal relationships among the included genera. For example, while Říčan et al. found

Parachromis in a subclade with *Amatitlania*, we find it in a different subclade that

includes *Amphilophus* (Fig. 1). Likewise, our grouping corresponds roughly with the

amphilophines of López-Fernández et al. 2010, but excludes *Trichromis ('Cichlasoma'*

salvini in their Fig. 1). Even though the results of our phylogenomic analyses have

stronger statistical support that those of López-Fernández et al. (2010) and of Říčan et al.

(2016, see their Fig. 5), the amphilophine clade found herein contains the largest number

of weakly supported nodes in our analyses. It is interesting to note that, among

amphilophines, the sequential position of *Petenia* and *Chortiheros* as sister to a clade of

Amatitlania, *Hypsophrys* and *Neetroplus* was consistently recovered in all analyses, but

support for this arrangement was ambiguous (Fig. 1, Appendices L & M). Říčan et al.

(2016) recovered these two genera as sister to each other, but none of our analyses

supported that relationship.

The largest disagreement among topologies obtained in this study, as well as with

those from previously published analyses, involves the genera in the informal clade

herichthyines, the Caribbean genus *Nandopsis*, and the caquetaines clade containing the

genera *Caquetaia*, *Heroina* and the newly named genus *Kronoheros* (Fig. 2A). The

relationship between these lineages and the amphilophines also varies among our three

analyses, and often differs from relationships found in other studies. It is interesting that

our concatenated and STAR coalescent analyses are more similar with each other than

either is to the ASTRAL-II topology. In Říčan et al.'s (2016) analysis, the caquetaines

were sister to a clade of amphilophines and herichthyins with *Nandopsis* sister to the

latter. This is a similar arrangement to that found in our STAR and concatenated

analyses, but in our ASTRAL-II topology (Fig. 2A) *Nandopsis* was sister to

amphilophines, and in turn, the two were sister to the caquetaines. In all analyses, at least

some of these relationships are inconclusively supported, although the concatenated

analysis obtained the highest bootstrap values of the three. In Říčan et al's (2016)

analysis, *Chiapaheros* remained in a polytomy. In our analysis, it is part of a well-

supported clade along with two other well-supported clades: *Thorichthys* and *Trichromis*

and a clade comprising *Herichthys*, *Vieja*, *Wajpamheros*, *Chuco* and *Theraps*, but the

relationships among these three lineages remains unclear (Fig. 2A).

Relationships among Geophagini in all analyses were generally compatible with

those previously described by López-Fernández et al. (2012) and comprised by two major

clades compatible with those described by López-Fernández et al. (2010, 2012). In the

first of these, a clade of *Guianacara* and *Mazarunia* is sister to a clade of

"apistogrammines" and "crenicichlines" sensu López-Fernández et al. (2010). Except for

a few nodes within the *Crenicichla-Teleocichla* group and another one within *Mazarunia*,

relationships within this large clade are largely congruent with previous results and well

supported. Additionally, the sister relationship between *Acarichthys* and *Biotoecus* was

unambiguously recovered, but support of its sister relationship to *Crenicichla* was always

below 100%.

A second clade of Geophagini comprised the genus *Biotodoma*, the

"geophagines", "mikrogeophagines" and "crenicaratines" of López-Fernández et al.

(2010). Monophyly of the "geophagines" (genera *Geophagus*, *Gymnogeophagus* and the

'Geophagus' steindachneri group) was unambiguously supported by all analyses, but

relationships within the group were different in the three topologies (Fig. 2B). Moreover,

the relative position of the genus *Biotodoma* and the "mikrogeophagines" (genera

Mikrogeophagus and *'Geophagus' brasiliensis*) was different in the coalescent-based

analyses compared to the concatenated topology. As discussed below, ambiguity in the

placement of the "geophagines" genera could have implications for estimating the age of

cichlids.

413 Relationships within Cichlasomatini were generally identical among analyses. In

414 al cases, a clade including *Nannacara*, *Ivanacara* and *Cleithracara* was recovered as the

415 unambiguously supported sister group to the rest of Cichlasomatini. This clade is

416 equivalent to the "nannacarines" sensu López-Fernández et al. (2010), but its position

417 with respect to the rest of the tribe is novel with respect to previous studies. Musilová et

418 al. (2009) found the "nannacarines" as sister to *Laetacara* but with low support, and

419 López-Fernández et al. (2010) found it as sister to a clade of *Laetacara*, *Acaronia* and

420 their "andinoacarines", but also with low support. The genera *Cichlasoma* and *Aequidens*

421 were found as sister to *Krobia* in all studies and as expected from previous work

422 (Musilová et al. 2009, López-Fernández et al. 2010). Our analyses also unambiguously

423 recovered a well-supported monophyly of *Andinoacara* and *Bujurquina* which in turn are

424 sister to *Tahuantinsuyoa*. *Acaronia* and *Laetacara* were recovered as sister to the

425 "andinoacarines" but the relative placement of the genera with respect to each other was

426 not unambiguously supported in any of the analyses. Placement of these two genera has

427 varied across studies: Musilová et al. (2009) found *Acaronia* to be sister to all

428 Cichlasomatini, with *Laetacara* either sister to the remainder of the tribe or to the

429 *Cichlasoma*, *Aequidens* and *Krobia* clade; López-Fernández et al. (2010) recovered a

430 weakly supported sister clade that in turn was sister to "andinoacarines", but support was

431 low.

433 **4. Discussion**

434 Even though higher-level relationships among clades and genera of Neotropical

435 cichlids have become increasingly resolved and supported by recent work, the position of

several groups remains uncertain or weakly supported. In this study, we used a recently

developed set of exon-targeting probes and massive parallel next generation sequencing

to generate a large dataset aimed at resolving Cichlinae relationships that remain poorly

supported. Our phylogenomic analyses confirm many relationships previously found

among Neotropical cichlids, and provide unprecedented resolution and support for many

relationships that were previously weakly supported, especially near the base of the tree.

This is particularly true among some genera of Central American lineages in the

amphilophines clade, the unambiguous clarification of the relationship among

Geophagini clades crenicichlines, apistogrammines and guianacarines (sensu López-

Fernández et al. 2010), and the strongly supported position of the Cichlasomatini clade

nannacarines as the sister group to the rest of the tribe (Fig. 1 and see Appendices L &

M). Nevertheless, despite an increase of two orders of magnitude in data when compared

with other sequencing studies (e.g. López-Fernández et al. 2010; Říčan et al. 2013), some

relationships remain unclearly established or poorly supported. The three main regions of

the tree that continue to resist clear resolution include the Central American

herichthyines, the position of the Cichlasomatini genera *Laetacara* and *Acaronia*, and the

order of divergence among genera in the Geophagini clade including *Geophagus*,

Gymnogeophagus and the *'Geophagus' steindachneri* clade (Fig. 2).

Previous studies have repeatedly found that Neotropical cichlids diversified over a

relatively short time period, as evidenced by short branches and frequently weakly

supported relationships near the base of the tree (e.g. Farias et al. 1999, López-Fernández

et al. 2005, 2010). Most of the previously unresolved or weakly supported relationships

in the López-Fernández et al. (2010) study, which has the most comparable taxon

sampling to this study, are resolved with unambiguous support by the phylogenomic

analyses presented here regardless of the method used (compare Fig. 1 of both studies).

Thus, in combination, the generalized stability of higher-level phylogenetic hypotheses

obtained through both summary coalescent and concatenation methods in our analyses,

suggests that conflict between gene trees and the species tree is relatively rare among

Neotropical cichlids. It appears that incomplete lineage sorting and other sources of

misleading phylogenetic signal, such as introgression or hybridization, do not frequently

disrupt our ability to reconstruct relationships. We make this assertion with caution,

however, because detailed phylogeographic studies of Neotropical cichlids have shown

that at least some taxa may be affected by these problems, particularly at more recent

levels of divergence (e.g. Willis et al. 2013, 2017). Even more germane to our study, the

persistence of unresolved or poorly supported "deep" relationships within a few clades of

Cichlinae may indicate a role for deep coalescence or other confounding effects in some

early events of divergence among Neotropical cichlids (Fig. 2).

473   This point is particularly relevant because the tribes containing the remaining

unresolved or conflictive clades underwent relatively quick adaptive diversification

giving origin to a variety of lineages (López-Fernández et al. 2013, Arbour and López-

Fernández 2014). It is suggestive that the largest remaining conflicts within the tree

involve the Central American herichthyines, including some South American genera with

South American distribution and the Caribbean genus *Nandopsis* (Fig. 2A). Recent work

has shown that invasion of Central America by the tribe Heroini provided renewed

ecological opportunity that allowed this clade to rapidly diversify into a broad variety of

ecologically specialized forms (Arbour and López-Fernández 2016). It is conceivable that

such rapid adaptive divergence produced incomplete lineage sorting among some heroine

lineages, resulting in reduced phylogenetic resolution (e.g. Kubatko and Degnan 2007,

Edwards 2009). In fact, the early radiation of the Neotropical cichlid tribes in South

America has been similarly shown to have occurred quickly, potentially leaving a

similarly conflictive gene tree divergence patterns in other regions of the tree, particularly

within Geophagini, which dominates the lineage and functional diversity of South

American Neotropical cichlids (López-Fernández et al. 2013, Arbour and López-

Fernández 2014, Astudillo-Clavijo et al. 2015). Even if early adaptive radiation in

Neotropical cichlids resulted in incomplete lineage sorting in some clades, our results

suggest that its effects may not be extensive because, with the exceptions pointed out

above, both our concatenated and coalescent analyses recover largely congruent

phylogenies. Moreover, most of the relationships recovered herein are congruent with

those founds in most previous studies based on much smaller, concatenated datasets (e.g.

Říčan et al. 2008, Musilová et al. 2009, López-Fernández et al. 2010, McMahan et al.

2013) and with the ddRAD-based analysis of Central American heroines of Říčan et al.

(2016).

The ability to generate a reliable phylogeny has important consequences beyond

the mere systematic applications of the study. Uncertainty about the order of divergence

and relationships among genera and higher clades can affect our ability to reconstruct the

history of evolutionary divergence in cichlids. Three types of studies could be affected by

diminishing but still present uncertainty in Neotropical cichlid relationships. Robust

topologies are critical for using the fossil record to calibrate molecular phylogenies and

reconstruct the timeline of diversification of a clade. The fossil record of cichlids is

relatively scarce, and considerable debate has ensued regarding the identity and

placement of fossils, particularly the recently described Eocene fossils from the Lumbrera

formation in Argentina (e.g. Malabarba et al. 2010, 2014, Friedman et al. 2013, López-

Fernández et al. 2013). On the one hand, the strong support received by most nodes in

our analyses should provide a solid scaffold for time calibration. Unfortunately, one of

the most unstable relationships, that involving the geophagine genera *Gymnogeophagus*,

Geophagus and *'Geophagus' steindachneri*, affects the certainty with which the Eocene

fossil †*Gymnogeophagus eocenicus* can be placed on the tree. As can be gleaned from

Fig. 2B, the position of *Gymnogeophagus* and its putative fossil taxon in the phylogeny is

unstable. As a consequence, calibration of *Gymnogeophagus* remains uncertain because it

is unclear whether *Gymnogeophagus* is sister to the broadly distributed *Geophagus*

sensus stricto or to the northern Andes clade including *'Geophagus' steindachneri* (Fig.

2B). Therefore, the inconclusively established position of a fossil-bearing clade can have

important consequences in both reconstruction of the timeline of divergence and the

historical biogeography of South American cichlids.

From a biogeographic point of view, the uncertainty observed among Central

American cichlids is likely to have even more dramatic consequences. The incongruent

placement of the Caribbean genus *Nandopsis* and of the South American caquetaines

limits our ability to accurately recreate the history of heroine invasion of Central America

and for understanding the events driving the potential recolonization of South America by

some of the Mesoamerican lineages such as *Mesoheros* and the caquetaines. Recent

studies that have addressed the historical biogeography of Central American cichlids and

their relationships to South America have relied on single reconstructions of the

phylogeny relying on either a small number of concatenated loci (e.g. Říčan et al. 2013,

Tagliacollo et al. 2015) or on phylogenomic approaches based on concatenation of single

nucleotide polymorphisms (Říčan et al. 2016). These studies focus on interpretations of

the particular trees found in each of their analyses, but our analyses suggest that the

historical biogeography of Central America may require either a more exhaustive

analysis of phylogenetic relationships or the reconstruction of competing, alternative

scenarios that reflect the current uncertainty in the phylogeny. Based purely on

methodological arguments of performance and accuracy (e.g. Mirarab and Warnow 2015,

Arcila et al. 2017), it is possible that our ASTRAL-II topology provides a more stable

framework for analysis, but the weak support received by nodes in that arrangement

suggest that any interpretations of historical biogeography should be done cautiously.

Phylogenies are also becoming increasingly important as the framework to

perform macroevolutionary analyses of lineage and phenotypic divergence. With a

proliferation of comparative methods and the increased availability of well supported and

increasingly better dated trees, our ability to infer patterns and process of divergence

continues to improve. With several studies recently addressing the evolution of

Neotropical cichlids (e.g. Hulsey 2006, López-Fernández et al. 2013, Arbour and López-

Fernández 2013, 2014, 2016, Astudillo-Cavijo et al. 2015, Burress 2016), it is pertinent

to ask whether changes in the topology may require modification of our emerging

understanding of Cichlinae macroevolution. The phylogeny obtained herein is

remarkably similar to the López-Fernández et al. (2010) tree used in most of the

macroevolutionary analyses listed above. When analyses are performed on a sample of

the posterior distribution of the dated phylogeny and not just on the Maximum Clade

Credibility (MCC) tree, topological uncertainty should be reflected in the results such

that observed macroevolutionary patterns should be robust to moderate topological

changes. This should be particularly true for analyses of phenotypic divergence based on

the construction of morphospaces, such as disparity through time and adaptive landscape

inferences (e.g. Ingram and Mahler 2013, Slater and Pennel 2014). Likely more sensitive

to changes in the tree are lineage through time and rate analyses because they depend on

branch lengths in ultrametric trees, which in turn depend on the accuracy of both the

topology and of age estimates. Because coalescent summary analyses such as ASTRAL-

II and STAR do not provide branch lengths, the species trees estimated here cannot be

used directly in comparative analyses that require ultrametric topologies derived from

estimates of absolute time.

Finally, generating an ultrametric topology by dating phylogenomic datasets

remains a challenge. In principle, there is no reason that phylogenomic species-tree

hypotheses cannot be dated, but in practice it is not clear if the incorporation of large

phylogenomic datasets and fossil data into actual analyses is computationally feasible

(e.g. Bouckaert et al. 2014, Matschiner et al. 2016). Moreover, among other problems,

traditional node dating methods require a priori placement of fossils on topologies,

further complicating the use of fossils in unresolved clades, such as *†Gymnogeophagus*

eocenicus. Alternatively, emerging total evidence dating methods that simultaneously

generate phylogenies and age estimates combining molecular, morphological and fossil

data may provide more flexibility and accuracy *vis a vis* uncertainty in the molecular

phylogenies (e.g. Heath et al. 2014). Whether these methods can be employed in truly

phylogenomic contexts with hundreds of loci and taxa is not yet clear (e.g. Ronquist et al.

2012, Gavryushkina et al. 2016).

5. Conclusion

Phylogenomic analyses of Neotropical cichlids (subfamily Cichlinae) using 415

exons for 139 species in both concatenated and summary statistic coalescent frameworks

resulted in generally well-resolved, strongly-supported and broadly-congruent topologies.

The topologies obtained are similar to previous hypotheses of relationships among

Cichlinae but, in general, provide stronger support for many relationships that previously

had weak or conflicting support. The results of our analyses also suggest that the targeted

genomic regions contain phylogenetic signal capable of resolving relationships at all

levels of divergence within the clade. Nevertheless, we identified several regions of the

tree in which relatively early divergence events cannot be reconstructed with certainty

because different methods provide conflicting results, none of them with conclusive

support. We suggest this incongruence may result from incompletely lineage sorting

associated with the early adaptive divergence of the clades in which incongruence is

observed. We argue that, despite some disagreements in the placement of these lineages

both in this study and in previous analyses, most studies provide a broadly common

signal of evolutionary relationships among Neotropical cichlids. A generally better

supported phylogeny derived from our phylogenomic analyses should continue to provide

a solid framework for the evolutionary analysis of lineage and phenotypic divergence in

Neotropical cichlids. Future work will focus in further clarifying relationships within the

few recalcitrant clades identified herein, and in leveraging the current dataset as a

uniquely strong molecular framework for clarifying the timeline of evolutionary

divergence among Neotropical cichlids.

Acknowledgements

The authors are grateful to M. Burridge, M. Zur and E. Holm (Royal Ontario Musem

[ROM]), M, Stiassny and B. Brown (American Museum of Natural History, New York),

M. Sabaj (Academy of Natural Sciences of Drexel University, Philadelphia), R. Rodíles-

Hernández (ECOSUR), L.R. Malabarba (Universidade Federal do Rio Grande do Sul,

Porro Alegre, Brazil), J. Armbruster (Auburn University Museum, Auburn, Alabama)

and Yves Fermon (Muséum National d'Histoire Naturelle, Paris) for curatorial assistance

and the loan of tissue samples not available at the ROM collection. We are grateful to the

people who assisted during fieldwork related to this work: J.H. Arbour, D. Bloom, R.R.

Hernández, F.E. Hauser, N.K. Lujan, N. Meliciano, E.A. Liverpool, C.G. Montaña, G.

Ortí, S. Refvik, M. A.P. Röepke, Soria, D.C. Taphorn, M. Tobler, S. C. Willis, K.O.

Winemiller, Funding for this work, including fieldwork and laboratory analyses was

provided by a Discovery Grant from the Natural Sciences and Engineering Research

Council of Canada, ROM Governors and the National Geographic Society grants to HLF.

Additional fieldwork was supported by a Coypu Foundation Grant to N.K. Lujan and

ROM Trust Funds through HLF. KLI was supported by a Rebanks Postdoctoral

Fellowship from the ROM Governors granted to HLF to develop a phylogenomic toolkit

for the phylogenetic analysis of cichlids.

References

Arbour JH, López-Fernández H. 2014. Adaptive landscape and functional diversity of

Neotropical cichlids: implications for the ecology and evolution of Cichlinae

(Cichlidae; Cichliformes). J Evol Biol 27:2431–42. doi: 10.1111/jeb.12486

Arbour JH, López-Fernández H. 2016. Continental cichlid radiations: functional diversity

reveals the role of changing ecological opportunity in the Neotropics. Proc R Soc

B Biol Sci 283:20160556. doi: 10.1098/rspb.2016.0556

Arcila D, Ortí G, Vari R, Armbruster JW, Stiassny MLJ, Ko KD, Sabaj MH, Lundberg J,

Revell LJ, Betancur-R. R, Much. 2017. Genome-wide interrogation advances

resolution of recalcitrant groups in the Tree of Life. Nat Ecol Evol. doi:

10.1038/s41559-016-0020

Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu CH, Xie D, Suchard M a., Rambaut

A, Drummond AJ. 2014. BEAST 2: A Software Platform for Bayesian

Evolutionary Analysis. PLoS Comput Biol 10:1–6. doi:

10.1371/journal.pcbi.1003537

Bowmaker JK. 2008. Evolution of vertebrate visual pigments. Vision Res 48:2022–41.

doi: 10.1016/j.visres.2008.03.025

Brawand D, Wagner CE, Li YI, Malinsky M, Keller I, Fan S, Simakov O, Ng AY, Lim

ZW, Bezault E, Turner-Maier J, Johnson J, Alcazar R, Noh HJ, Russell P, Aken

B, Alföldi J, Amemiya C, Azzouzi N, Baroiller J-F, Barloy-Hubler F, Berlin A,

Bloomquist R, Carleton KL, Conte M a., D'Cotta H, Eshel O, Gaffney L, Galibert

F, Gante HF, Gnerre S, Greuter L, Guyon R, Haddad NS, Haerty W, Harris RM,

Hofmann H a., Hourlier T, Hulata G, Jaffe DB, Lara M, Lee AP, MacCallum I,

Mwaiko S, Nikaido M, Nishihara H, Ozouf-Costaz C, Penman DJ, Przybylski D,

Rakotomanga M, Renn SCP, Ribeiro FJ, Ron M, Salzburger W, Sanchez-Pulido

L, Santos ME, Searle S, Sharpe T, Swofford R, Tan FJ, Williams L, Young S, Yin

S, Okada N, Kocher TD, Miska E a., Lander ES, Venkatesh B, Fernald RD,

Meyer A, Ponting CP, Streelman JT, Lindblad-Toh K, Seehausen O, Di Palma F.

2014. The genomic substrate for adaptive radiation in African cichlid fish. Nature

513:375–381. doi: 10.1038/nature13726

Dornburg A, Townsend JP, Friedman M, Near TJ. 2014. Phylogenetic informativeness

reconciles ray-finned fish molecular divergence times. BMC Evol Biol 14:169.

doi: 10.1186/s12862-014-0169-0

Edwards SV. 2009. Is a new and general theory of molecular systematics emerging?

Evolution 63:1–19. doi: 10.1111/j.1558-5646.2008.00549.x

Farias, I.P., Ortí, G., Sampaio, I., Schneider, H., Meyer, A., 1999. Mitochondrial DNA

phylogeny of the family Cichlidae: monophyly and fast molecular evolution of

the Neotropical assemblage. J. Mol. Evol. 48, 703–711.

Farias IP, Ortí G, Meyer A. 2000. Total evidence: molecules, morphology, and the

phylogenetics of cichlid fishes. J Exp Zool 288:76–92.

Friedman M, Keck BP, Dornburg A, Eytan RI, Martin CH, Darrin C, Wainwright PC,

Near TJ, Hulsey CD. 2013. Molecular and fossil evidence place the origin of

cichlid fishes long after Gondwanan rifting. Proc R Soc B Biol Sci 280:20131733.

doi: http://dx.doi.org/10.1098/rspb.2013.1733

Gavryushkina A, Heath TA, Ksepka DT, Stadler T, Welch D, Drummond AJ,

Avryushkina ALG, Eath TRAH, Sepka DATK, Tadler TAS, Elch DAW. 2016.

664   Bayesian Total-Evidence Dating Reveals the Recent Crown Radiation of

665   Penguins. Syst Biol 0:syw060. doi: 10.1093/sysbio/syw060

666 Heath TA, Huelsenbeck JP, Stadler T. 2014. The fossilized birth-death process for

667   coherent calibration of divergence-time estimates. Proc Natl Acad Sci. doi:

668   10.1073/pnas.1319091111

669 Heled J, Drummond AJ. 2010. Bayesian inference of species trees from multilocus data.

670   Mol Biol Evol 27:570–80. doi: 10.1093/molbev/msp274

671 Hulsey CD, García de León FJ, Rodiles-Hernández R. 2006. Micro- and

672   macroevolutionary decoupling of cichlid jaws: a test of Liem's key innovation

673   hypothesis. Evolution 60:2096–109.

674 Hulsey CD, García De León FJ. 2005. Cichlid jaw mechanics: Linking morphology to

675   feeding specialization. Funct Ecol 19:487–494. doi: 10.1111/j.1365-

676   2435.2005.00987.x

677 Ilves, K.L., López-Fernández, H. 2014. A targeted next-generation sequencing toolkit for

678   exon-based cichlid phylogenomics. Mol. Ecol. Res. 14, 802–811.

679 Ingram T, Mahler DL. 2013. SURFACE: Detecting convergent evolution from

680   comparative data by fitting Ornstein-Uhlenbeck models with stepwise Akaike

681   Information Criterion. Methods Ecol Evol 4:416–425. doi: 10.1111/2041-

682   210X.12034

683 Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton,

684 S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Mentjies, P., &

685   Drummond, A. 2012. Geneious Basic: an integrated and extendable desktop

software platform for the organization and analysis of sequence data.

Bioinformatics 28, 1647-1649.

Kocher TD. 2004. Adaptive evolution and explosive speciation: the cichlid fish model.

Nat Rev Genet 5:288–298. doi: 10.1038/nrg1316

Kubatko LS, Degnan JH. 2007. Inconsistency of phylogenetic estimates from

concatenated data under coalescence. Syst Biol 56:17–24. doi:

10.1080/10635150601146041

Kullander, S.O., 1998. A Phylogeny and Classification of the Neotropical Cichlidae

(Teleostei: Perciformes). In: Malabarba, L.R., Reis, R.E., Vari, R.P., Lucena,

Z.M., Lucena, C.A.S. (Eds.), Phylogeny and classification of Neotropical fishes.

EDIPUCRS, Porto Alegre, pp. 461–498.

Liu, L., Yu, L., Pearl, D.Kk, Edwards, S.V., 2009. Estimating species phylogenies using

coalescence times among sequences. Syst. Biol. 58, 468–477.

Loken, C., Gruner, D., Groer, L., Peltier, R., Bunn, N., Craig, M., Henriques, T.,

Dempsey, J., Yu. C.-H., Chen, Dursi, J.L., Chong, J., Northrup, S., Pinto, J., Knecht, N.,

Van Zon, R., 2010. SciNet: lessons learned from building a power-efficient top-

20 system and data centre. J. Phys.: Conf. Ser. 256, 012026. doi: 10.1088/1742-

6596/256/1/012026.

López-Fernández H, Winemiller KO, Honeycutt RL. 2010. Multilocus phylogeny and

rapid radiations in Neotropical cichlid fishes (Perciformes: Cichlidae: Cichlinae).

Mol Phylogenet Evol 55:1070–1086. doi: 10.1016/j.ympev.2010.02.020

López-Fernández H, Taphorn DC, Liverpool EA. 2012. Phylogenetic diagnosis and

expanded description of the genus Mazarunia Kullander,1990 (Teleostei :

709   Cichlidae) from the upper Mazaruni River , Guyana , with description of two new

710   species. 10:465–486.

711 López-Fernández H, Arbour JH, Winemiller KO, Honeycutt RL. 2013. Testing for

712   ancient adaptive radiations in neotropical cichlid fishes. Evolution 67:1321–37.

713   doi: 10.1111/evo.12038

714 Maddison W. 1997. Gene trees in species trees. Syst Biol 46:523–536.

715 Mendes F, Hahn M. 2017. Why Concatenation Fails in the Anomaly Zone. BioRxiv.

716   https://doi.org/10.1101/116509

717 Matschiner M, Musilová Z, Barth JMI, Starostová Z, Salzburger W, Steel M, Bouckaert

718   R (2016) Bayesian Phylogenetic Estimation of Clade Ages Supports Trans-

719   Atlantic Dispersal of Cichlid Fishes. Syst Biol syw076. doi:

720   10.1093/sysbio/syw076

721 McMahan CD, Chakrabarty P, Sparks JS, Smith WL, Davis MP. 2013. Temporal Patterns

722   of Diversification across Global Cichlid Biodiversity (Acanthomorpha:

723   Cichlidae). PLoS One 8:e71162. doi: 10.1371/journal.pone.0071162

724 McMahan CD, Matamoros WA, Piller KR, Chakrabarty P. 2015. Taxonomy and

725   systematics of the herichthyins (Cichlidae: Tribe Heroini), with the description of

726   eight new Middle American Genera. Zootaxa 3999:211–234. doi:

727   10.11646/zootaxa.3999.2.3

728 Mirarab, S., Warnow, T. 2015. ASTRAL-II: coalescent-based species tree estimation

729   with many hundreds of taxa and thousands of genes. Bioinformatics 31, i44–i52.

730   doi: 10.1093/bioinformatics/btv234

Musilová Z, Říčan O, Novák J. 2009. Phylogeny of the Neotropical cichlid fish tribe

Cichlasomatini (Teleostei: Cichlidae) based on morphological and molecular data,

with the description of a new genus. J Zool Syst Evol Res 47:234–247. doi:

10.1111/j.1439-0469.2009.00528.x

Říčan O, Piálek L, Dragová K, Novák J. 2016. Diversity and evolution of the Middle

American cichlid fishes (Teleostei: Cichlidae) with revised classificatio. Vertebr

Zool 66:1–102.

Ronquist F, Klopfstein S, Vilhelmsen L, Schulmeister S, Murray DL, Rasnitsyn AP.

2012. A total-evidence approach to dating with fossils, applied to the early

radiation of the hymenoptera. Syst Biol 61:973–999. doi: 10.1093/sysbio/sys058

Seehausen O. 2015. Process and pattern in cichlid radiations - inferences for

understanding unusually high rates of evolutionary diversification. New Phytol

207:304–312. doi: 10.1111/nph.13450

Seo, T.K. 2008. Calculating bootstrap probabilities of phylogeny using multilocus

sequence data. Mol. Biol. Evol. 25, 960–971.

Shaw, T.I, Ruan, Z., Glenn, T.C., Liu, L. 2013. STRAW: Species TRee Analysis Web

server. Nucleic Acids Res. 41, W238–W241. doi: 10.1093/nar/gkt377

Slater GJ, Pennell MW. 2014. Robust regression and posterior predictive simulation

increase power to detect early bursts of trait evolution. Syst Biol 63:293–308. doi:

10.1093/sysbio/syt066

Smith L, Chakrabarty P, Sparks J. 2008. Phylogeny, taxonomy, and evolution of

Neotropical cichlids (Teleostei: Cichlidae: Cichlinae). Cladistics 24:625–641.

Sparks JS. 2008. Phylogeny of the Cichlid Subfamily Etroplinae and Taxonomic

754   Revision of the Malagasy Cichlid Genus Paretroplus (Teleostei: Cichlidae). Bull

755   Am Museum Nat Hist 314:1. doi: 10.1206/314.1

756   Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-

757   analysis of large phylogenies. Bioinformatics 30, 1312–1313.

758   Thompson AW, Betancur-R R, López-Fernández H, Ortí G. 2014. A Time-Calibrated,

759   Multi-locus Phylogeny of Piranhas, Pacus, and Allies (Characiformes:

760   Serrasalmidae) and a Comparison of Species Tree Methods. Mol Phylogenet Evol

761   In review:242–257. doi: 10.1016/j.ympev.2014.06.018

762   Tonini J, Moore A, Stern D, Shcheglovitova M, Ortí G. 2015. Concatenation and Species

763   Tree Methods Exhibit Statistically Indistinguishable Accuracy under a Range of

764   Simulated Conditions. 1–15. doi:

765   10.1371/currents.tol.34260cc27551a527b124ec5f6334b6be.Abstract

766   Wagner CE, Harmon LJ, Seehausen O. 2012. Ecological opportunity and sexual selection

767   together predict adaptive radiation. Nature 487:366–369. doi:

768   10.1038/nature11144

769   Weadick CJ, Loew ER, Rodd FH, Chang BSW. 2012. Visual pigment molecular

770   evolution in the Trinidadian pike cichlid (*Crenicichla frenata*): a less colorful

771   world for neotropical cichlids? Mol Biol Evol 29:3045–60. doi:

772   10.1093/molbev/mss115

773   Willis SC, Farias IP, Ortí G. 2013. Testing mitochondrial capture and deep coalescence

774   in Amazonian cichlid fishes (Cichlidae: Cichla). Evolution 68:256–68. doi:

775   10.1111/evo.12230

Willis SC. 2017. One species or four? Yes!...and, no. Or, arbitrary assignment of lineages

to species obscures the diversification processes of Neotropical fishes. PLoS One

12:e0172349. doi: 10.1371/journal.pone.0172349

Figure 1. Species tree generated with ASTRAL-II for 415 loci comprising 471,448 bp.

Colored clades depict formally recognized Neotropical cichlid tribes Heroini (Green),

Cichlasomatini (Orange), Geophagini (Blue), Chaetobranchini (Magenta), Astronotini

(Red), Retroculini (Brown), and Cichlini (Purple). See text for further discussion of

relationships among and within tribes. Node bootstrap support is indicated when

pertinent; nodes without labels received 100% support in this analysis. Nodes labeled in

red received <75% bootstrap support. Nodes marked with an asterisk (*) represent

weakly supported intergeneric relationships with incongruent resolution among two

different species-tree and one concatenated phylogenetic analyses. See Figure 2 and

Discussion for further analyses of these results. Complete topologies not shown here are

provided in Appendices L and M along with node support.

A. Centra American Heroini

B. Geopgagini: *Geophagus-Gymnogeophagus* clade

C. Cichlasomatini: *Laetacara-Acaronia*

Figure 2. Conflicting results among analyses. Each panel depicts the alternative

topological arrangements found in the clades highlighted with asterisks in Fig. 1. Colors

are meant to represent lineages within each panel and not to be used as comparison

among clades in different panels. Numbers by nodes represent bootstrap support in each

case; nodes without numbers received 100% bootstrap support in their respective

analyses. See Fig. 1 for the ASTRAL-II species tree and Appendices L and M to see the

concatenated and STAR topologies.

Appendix L – Maximum Likelihood concatenated tree produced from analysis of 415

loci with no partitioning. Numbers in each node denote bootstrap support based on 1000

replicates in RAxML. See text and Figures 1 and 2 for details.

Appendix M. Summary coalescent tree generated with the program STAR. Numbers in

each node denote bootstrap support. See text and Figures 1 and 2 for details.