On Model Averaging Partial Regression Coefficients

Jeffrey A. Walker[*][a]

[a]Department of Biological Sciences, University of Southern Maine,
Portland, ME 04103, USA

November 19, 2017

1   Running Head: Model averaging partial regression coefficients

[*]walker@maine.edu

Abstract

Model averaging partial regression coefficients has been criticized because the coefficients from different submodels estimate different parameters, so the coefficients have different meanings from model to model. A partial regression coefficient estimates two very different parameters, a regression (conditional effect) parameter and a (causal) effect parameter. The regression parameter is a descriptor, and its meaning is conditional on the covariates in the model. By contrast, an effect parameter derives its meaning from a specified causal hypothesis and not from the set of covariates. A multiple regression model implicitly specifies a causal model with direct, causal paths from each predictor to the response. Consequently, the partial regression coefficient for any predictor has the same meaning across all sub-models if the goal is estimation of the causal effects that generated the response. In a recent article, Cade (2015) suggested that, in the presence of *any* multicollinearity, partial regression coefficients not only estimate different parameters but their averaging is invalid because they have no defined units. I argue that Cade's interpretation of the math is incorrect. While partial regression coefficients may be meaningfully averaged, model averaging may not be especially useful. To clarify this, I compare effect estimates using a small Monte-Carlo simulation. The simulation results show that model-averaged (and ridge) estimates have increasingly better performance, relative to full model estimates, as multicollinearity increases, despite the full regression model correctly specifying the causal effect structure (that is, even when we know the truth, a method that averages over incorrectly specified models outperforms the correctly specified model).

Keywords multiple regression, causal model, conditional effect, monte carlo simulation, ridge regression, model selection, multicollinearity.

Introduction

Model averaging is an alternative to model selection for either effect estimation or prediction. Model averaging is attractive because it recognizes that for data typical in biology, *all* measured predictors will have some non-zero association with the response variable independent of that shared with other predictors. Consequently, model averaging encourages the worthy goal of emphasizing effect estimation and not simply the identification of some "best" subset of predictors. Model-selection often follows an all-subsets regression, a practice that is criticized for mindless model building. Nevertheless, averaging across all or a best subset of models shrinks regression coefficients toward zero, which has the effect of contracting error variance. Consequently, model-averaging can outperform model selection or the full model under some conditions, where performance is measured by a summary of the long run frequency of error (Raftery et al., 1997; Hjort and Claeskens, 2003).

Despite these features of model-averaging, model averaged partial regression coefficients have been criticized in the recent ecology literature because their computation requires averaging over a set of coefficients that are conditional on a specific set of covariates (Cade, 2015; Banner and Higgs, 2017). That is, coefficients from different models have different interpretations and cannot be meaningfully averaged. Cade (2015) took this criticism further, and specifically argued that in the presence of any multicollinearity, averaging coefficients is meaningless because an averaged coefficient has "no defined units." Cade's criticism is not the typical caution against the estimation of partial regression coefficients in the presence of high multicollinearity because of a high variance inflation factor but an argument that model-averaged coefficients in the presence of *any* correlation among the predictors are, quite literally, meaningless. Cade's critique is receiving much attention, as evidenced by the 108 Google Scholar citations

in about two years.

These critiques are noteworthy given that model averaging has developed a rich lit-

erature in applied statistics over the last 25 years (Hoeting et al., 1999; Burnham and

Anderson, 2002; Hjort and Claeskens, 2003; Hansen, 2007; Liang et al., 2011; Zhang

et al., 2014) with only limited attention to the meaning of the parameter estimated by

a model averaged coefficient (Draper, 1999; Candolo et al., 2003; Raftery and Zheng,

2003). Berger et al. (2001) noted the issue not in the context of a meaningless aver-

age but in the context of modeling the prior distribution. For similar purposes, Con-

sonni and Veronese (2008) showed that the parameters of a single submodel have four

different interpretations. In one of these, the parameter for X_j in a submodel has the

same meaning as that in the full model if we model the submodel as a variant of the

full model with the parameters for the excluded predictors set to zero (Equation 1)

(this is a more formal version of the reply given by Hoeting et al., 1999, in response

to Draper).

$$Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \varepsilon_i \quad \text{full model: } \beta_2 \neq 0 \tag{1}$$

$$Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \varepsilon_i \quad \text{sub model: } \beta_2 = 0 \tag{2}$$

Here, I offer a defense of model averaging partial regression coefficients that is re-

lated to this zero effect interpretation of the parameters: the coefficients $b_{j.m}$ from dif-

ferent submodels can be meaningfully averaged if they estimate a common effect pa-

rameter (β_j), which is the direct causal or generating effect of X_j on Y. In short, if a

researcher wishes to *describe* a statistical relationship conditional on a specific set of

covariates, then model averaging would indeed be averaging different things and an av-

eraged value would have an awkward (or not especially useful) interpretation. Often,

however, researchers use multiple regression to explicitly (or more commonly implic-

itly) estimate the causal effects that *generated the data*, that is, the effect parameters.

Importantly, an effect parameter derives its meaning from a pre-specified causal hy-

pothesis and this meaning is *independent of the set of variables in the full model* (Pearl,

2009). Consequently, averaging estimates of these parameters is perfectly meaningful.

Here, I show this independence using the math of causal modeling with causal (path or

structural) diagrams (Wright, 1934; Shipley, 2002; Pearl, 2009).

A motivating example: the effect of parental sex on off-spring calling in owls

A recent article on best practices involving regression-like models (Zuur and Ieno, 2016)

used as an example the data of Roulin and Bersier (2007), who showed that – and

entitled their paper – "nestling barn owls beg more intensely in the presence of their

mother than in the presence of their father." This title might simply be a description of

the major result, that is, a difference in conditional means (on the set of covariates in

the study, including time of arrival, time spent in nestbox, a food manipulation treat-

ment, and all interactions with parental sex). In the discussion, however, Roulin and

Bersier (2007) state that "differential begging to mother and father implies that off-

spring should be able to recognize the identity of each parent." That is, the chick be-

havior is in direct response to the sex of the parent, or, put differently, the sex of the

parent bringing food causally modifies the intensity of chick calling.

This example serves to introduce a major argument of the present note: the param-

eters estimated by the coefficients of a linear model used for *causal* inference are funda-

mentally different from the parameters estimated by the coefficients of a linear model

used for *describing* differences in conditional means. A partial regression coefficient $b_{j.m}$

is a difference in conditional means – it is the difference in the mean response between two groups that vary in x_j by one unit but have the same values for all other covariates (X_m). The partial regression coefficient $b_{j.m}$ is an estimator of two parameters

$$\theta_{j.m} = E(Y|X_j = x_j + 1, X_m = x_m) - E(Y|X_j = x_j, X_m = x_m) \tag{3}$$

$$\beta_j = E(Y|do(X_j = x_j + 1)) - E(Y|do(X_j = x_j)) \tag{4}$$

$\theta_{j.m}$ is the regression (or conditional effect) parameter, a descriptive parameter describing the population difference in conditional means. β_j is an effect parameter (Pearl, 2009), the direct, generating effect of X_j on Y. The concept of effect coefficients goes back to at least the seminal work of Sewell Wright (1918, 1934) in his method of path analysis. The *do* operator represents what would happen in an experimental intervention that modifies X_j but leaves all other variables unchanged (Pearl, 2009). The *do* operator is what we *do* when we conduct a randomized experiment and when we simulate a data generating process. Importantly, β_j is not a conditional effect, and its definition does not refer to other variables in the model. This is because β_j derives its meaning from a pre-specified causal hypothesis and this meaning is independent of other variables that might causally influence Y (Pearl, 2009). A reasonable concern is, how does an effect parameter, which represents the effect of an intervention, apply to observational data, where a change in the value of X_j is always associated with changes in other predictor variables? The answer is, the formal definition clarifies the assumptions needed to use a partial regression coefficient as an estimate of an effect parameter. More specifically, a partial regression coefficient $b_{j.m}$ is a consistent estimator of the effect parameter β_j if the regression model correctly specifies the generating process. This specification includes the inclusion of all confounders of the effect of X_j on Y, which are variables that both causally effect Y by a path independent of that through

X_j and are correlated with X_j. A partial regression coefficient is a biased estimate of

β_j if the regression model excludes confounders for X_j – a bias known as omitted vari-

able bias. Omitted variable bias is an introductory textbook concept in disciplines that

commonly use regression for causal modeling, including econometrics and epidemiol-

ogy, but not disciplines where explicit causal modeling is uncommon, including biology

generally, and ecology and evolution, specifically.

Model-averaged coefficients are meaningful estimates

of effect parameters β_j

Figure 1:

To highlight the difference between the parameters, $\theta_{j.m}$ and β_j, I use the Roulin

and Bersier (2007) example and show three simplified, generative models of $\log(NCalls)$

(Figure 1). The effect coefficients β_1 and β_2 are the direct, causal effect of $ParentSex$

and $ArrivalTime$ on Y. In Model A, $ArrivalTime$ is measured and included in the re-

gression model. In Models B, $ArrivalTime$ is unmeasured and designated $U_{ArrivalTime}$.

Both models also include U, which represents all un-modeled factors, other than $ArrivalTime$,

that contribute to the variance in Y. The generating models also include a correlation

r_{12} between $ParentSex$ and $ArrivalTime$. An important assumption of these models

is that the unmeasured factor U is not correlated with either of the X_j as indicated by

a lack of a double-headed arrow (otherwise it would be a confounding variable – see

below).

In Model A, the partial regression coefficients $b_{1.2}$ and $b_{2.1}$ are unbiased estimates

of the parameters $\theta_{1.2}$ and $\theta_{2.1}$. Because the model is correctly specified, the regres-

sion coefficients are also unbiased estimates of the generating parameters β_1 and β_2.

In Model A, then, the parameters $\theta_{j.m}$ and β_j coincide, in the sense that both have the

same value. This coincidence occurs only under very limited conditions.

Model B is the same generating model as Model A, but the regression model omits

ArrivalTime. The effect parameter β_1 has precisely the same meaning in Model B as

it did in Model A, but the regression coefficient b_1 is a biased estimate of β_1 because

of the omitted variable. This bias is $r_{12}\beta_2$. By contrast, the regression parameter θ_1 in

Model B has a different meaning than $\theta_{1.2}$ in Model A, and differs from the latter by

$r_{12}\beta_2$, but the regression coefficient b_1 is an unbiased estimate of θ_1. In Model B, then,

the parameters θ_1 and β_1 do not coincide.

Note that the missing confounder *ArrivalTime* in Model B does not bias the esti-

mate of θ_1 but does bias the estimate of β_1 (again, by $r_{12}\beta_2$). Consequently, if the goal

is mere description, omitted confounders are not a concern. But if the goal is causal

modeling, missing confounders are (or should be) a major concern. Omitted confounders

result in standard errors of effect coefficients that are (often far) too small, which re-

sults in inflated confidence in effect magnitudes and even signs (Walker, 2014).

Importantly, while the meaning and/or value of the regression parameter $\theta_{j.m}$ dif-

fers among the models, the meaning and value of the effect parameter β_j is constant

among the models. Because the meaning of the β_j is invariable across generating mod-

els, estimates of β_j have the same meaning – they are estimates of β_j – across regression

models, regardless of the set of covariates specified in the model. And because partial

regression coefficients *as estimates of* β_j have the same meaning across regression mod-

els, partial regression coefficients from different models can be meaningfully averaged.

Model-averaged coefficients outperform full-model coefficients when power is low

Even though partial regression coefficients as estimates of effect coefficients can be meaningfully averaged, the averaged coefficients may not be very useful, compared to the coefficients computed from the full model. Here, I show that, despite averaging over incorrectly specified models, model-averaged coefficients can outperform the full-model coefficients for estimating effect parameters even when the full regression model correctly specifies the generating model. I use a Monte Carlo simulation experiment to measure the total (root mean square) error in the estimates relative to the known, generating parameters. Freckleton (2010) used a similar simulation to show how full model and model-averaged estimates perform with increased multicollinearity and showed that the error variance of the full model estimates increases more rapidly than that of the model-averaged estimates but that the model-averaged estimates were increasingly biased with increased multicollinearity. Unfortunately, Freckleton (2010) did not combine the two sources of error into a single measure (such as mean square error) for comparison and, as a consequence, I cannot determine from his results which method performs better *overall*. Because I am specifically comparing the relative performance of the estimators at different levels of multicollinearity, I also compare ridge regression estimates. The simulation is not meant to be a comprehensive comparison of model averaging estimators but simply a pedagogical case study of why model average estimators should be considered as reasonable alternatives to the full model.

183   Data simulating the owl call data (Roulin and Bersier, 2007) were generated with

$$NCalls \sim \text{Poisson}(\mu_i) \tag{5}$$

$$\log(\mu_i) = \beta_0 + \beta_1 ParentSex_i + \beta_2 ArrivalTime_i \tag{6}$$

$$ParentSex_i = Z_i \tag{7}$$

$$ArrivalTime_i \sim N(\beta_z Z_i, 1) \tag{8}$$

$$Z_i = 0 \text{ or } 1 \tag{9}$$

$NCalls$ are sampled for $n = 27$ nests, once for each parent. There is no nest effect in

the generation of the data (nor is one modeled in the regression). $\text{Exp}(\beta_0)$ is the ex-

pected number of calls (175) during the mother's visit. $\text{Exp}(\beta_1)$ was set to one of four

values 0.99, 0.98, 0.97, and 0.96, which is equivalent to standardized effects (Cohen's d)

of -0.13, -0.27, -0.40, and -0.53. The expected reductions in calls during the father's vis-

its holding $ArrivalTime$ constant are 1.75, 3.50, 5.25, and 7.00. $\text{Exp}(\beta_2)$ was set to 0.9.

Z is the common cause of $ParentSex$ and $ArrivalTime$, which creates a correlation

(collinearity) between the two effects. Z and $ParentalSex$ are equal numerically (fe-

male=0, male= 1) but are not conceptually equivalent. Z is the sex determining factor

while $ParentalSex$ is the phenotypic feature that allows a chick to identify the parent

as dad or mom. The expected correlation between $ParentSex$ and $ArrivalTime$ is set

(using the parameter β_z) to one of four values: 0.2, 0.4, 0.6, and 0.8. Each iteration,

the empirical correlation is checked and only used if it is within 0.02 of the expected

correlation. 5000 iterations were run for each combination of β_1 (controlling effect size

and power) and β_z (controlling collinearity). The power to reject a null direct effect

($\beta_1 = 0$) at a type I error rate of 5% was computed using the 5000 runs for each com-

bination of the causal parameters β_1 and β_z. The performance (the ability to estimate

β_1) of model averaged, full model, and ridge estimates were quantified using the long-run error $RMSE = \sqrt{\frac{\sum(b_1 - \beta_1)^2}{5000}}$, which accounts for both error variance and bias.

The entire simulation was implemented in R (R Core Team, 2015) and the script is available in Supplement 1. In each run, the generating coefficients were estimated using the full model, model averaging, and ridge regression. Model-averaged coefficients were computed using AICc weights and over all models using the dredge and model.avg function in the MuMIn package. Coefficients of predictors excluded from a model were assigned a value of zero (using the row of the coefficient table with the label "full"). For the ridge regression, I used the cv.glmnet function (setting alpha=0) in the glmnet package (Friedman et al., 2010) and used the default 10-fold cross-validation to compute the optimal tuning parameter.

The performance of the three methods are shown in (Figure 2, where the X-axis is the effect parameter of $ParentSex$ standardized by the average expected variance to generalize the results. More specifically, $\beta_1' = \beta_1/\sigma$, where σ is the square root of $(\text{Exp}(\beta_0)/2 + \text{Exp}(\beta_0 + \beta_1)/2)$. Again, these standardized effects are -0.13, -0.27, -0.40, and -0.53. While -0.13 is a "small" standardized effect, this is a common value in ecology, where an estimated average standardized effect size is 0.18-0.19 (Møller and Jennions, 2002). The label for each panel shows the mean correlation (over the 5000 runs) between $ParentSex$ and $ArrivalTime$ (again, all of the correlations within a batch of 5000 runs were within 0.02 of the expected correlation).

The simulation shows increased RMSE for all estimators as collinearity increases and the qualitative pattern of relative performance among models remains about the same as collinearity increases (Figure 2). Quantitatively, the full model RMSE increases 71% (averaged over the four levels of β_1 as the correlation increases from 0.2 to 0.8. By contrast, the model averaged and ridge RMSE increase 40% and 37%. Model averag-

Figure 2: **Error as a function of increased collinearity**. Root mean square error in the estimate of β_1 (the effect parameter of $ParentSex$) over the 5000 runs for each combination of β_1 and the correlation r between $ParentSex$ and $ArrivalTime$. The effect parameter is standardized and presented as Cohen's d. Key to methods: MA = model average, OLS = full model, Ridge = ridge regression.

ing outperforms the full model when power is relatively low despite the full model correctly specifying the generating model. When collinearity is high, power is relatively low at all tested effect sizes of β_1' and, consequently, model averaging outperforms the full model at all levels of β_1'. Ridge regression outperforms model averaging over much of the space except at the lowest power.

Partial regression coefficients from different models have the same units

If Cade (2015) were simply cautioning against model averaging when data have high multicollinearity, the simulation above would suggest otherwise, as it shows that model averaging is less affected by multicollinearity than the full model. But a warning is not Cade's point; instead, Cade argues that model averaging is meaningless if there is *any* correlation among predictors because model-averaged coefficients "have no defined units

Table 1: Two interpretation of statements in Cade (2015)

Statement	Interpretation
1. "Their AIC model averaging of regression coefficients acts as if they are just numbers without any units attached to them"	the units of $b_{j.m}$ differ among models
2. "It is impossible to interpret the model-averaged regression coefficients (Tables 2–5) in terms of a $\Delta y/\Delta X_i$ because we do not know what units should apply to the denominator because it no longer refers to any specific covariance structure among the predictor variables"	
3. "Multicollinearity implies that the scaling of units in the denominators of the regression coefficients may change across models such that neither the parameters nor their estimates have common scales"	a unit difference in X_j differs among models
4. "the model averaging of regression coefficients ends up being done across estimates ($\beta_i = \Delta y/\Delta X_i$) without common denominators and is nonsensical because a unit change in the predictor variable (ΔX_i) is not the same across all models."	

in the presence of multicollinearity." As a consequence, Cade (2015) simply rejects the only experiments where we know the truth and where we can empirically compare the efficacy of different estimators.

It is therefore imperative that we explore what Cade means by "No defined units." This might mean that 1) the units of $b_{j.m}$ differ among submodels, or 2) a unit difference in X_j differs among submodels (Table 1). The first interpretation is simply false; a partial regression coefficient $b_{j.m}$ has the units of the simple regression coefficient of Y on X_j regardless of the other predictors in the model (Supplement 1). Cade clarifies the second interpretation using the Frisch-Waugh decomposition of $\mathbf{b} = (\mathbf{X}^\top \mathbf{X})^{-1}(\mathbf{X}^\top \mathbf{y})$

$$b_{j.m} = \frac{\mathrm{COV}(\tilde{X}_{j.m}, Y)}{\mathrm{VAR}(\tilde{X}_{j.m})} \qquad (10)$$

where $\tilde{X}_{j.m}$ is the component of the variation of X_j not shared with the other X, which is simply the vector of residuals of the regression of X_j on the set of covariates X_m. I

interpret Cade as stating that, because the variance of the residuals $\tilde{X}_{j.m}$ have different values among the models (because of different patterns of shared variance among the X), the scaling of the numerator differs among the models (I agree) and, therefore, a unit difference in X_j differs among models, or, equivalently, the units of $\tilde{X}_{j.m}$ differ among models. This conclusion (everything after "therefore") is a misunderstanding of the math; the magnitude of a unit or unit difference is defined by the actual units and not the variance of $\tilde{X}_{j.m}$. The units of the residuals of weight on height are full kilograms, not partial kilograms. In the owl example, if $ArrivalTime$ is measured in hours, a one unit difference is one hour regardless if we are referring to the raw measures or the residuals of $ArrivalTime$ on $ParentSex$. One hour (or one kilogram or one degree Celsius) does not shrink or swell among models due to differences in the magnitude of $\mathrm{VAR}(\tilde{X}_{j.m})$. What does shrink or swell with differences in $\mathrm{VAR}(\tilde{X}_{j.m})$ is the magnitude of the difference in $E(Y)$ given a one unit difference in X_j, but this is just re-stating the definition of a partial regression coefficient.

Conclusion

In their abstract, Banner and Higgs (2017) state that the "use of model averaging implicitly assumes the same parameter exists across models so that averaging is sensible. While this assumption may initially seem tenable, regression coefficients associated with particular explanatory variables may not hold equivalent interpretations across all of the models in which they appear, making explanatory inference about covariates challenging." This statement fails to recognize that an effect parameter β_j is distinct from a regression (difference in conditional means) parameter $\theta_{j.m}$ and the former but not the latter has the same meaning across models, because a causal hypothesis takes its meaning from a specified path and not the combination of variables in the model

(Pearl, 2009) (Figure 1). As emphasized by Pearl (2009), the two parameters only coin-

cide if the regression model correctly specifies the generating model.

Cade (2015) explicitly rejects simulation experiments similar to that above, arguing

that "the statistical performance suggested by distributions of their simulated model-

averaged estimates is of questionable merit" because model averaging is "nonsensical"

because the averaged coefficients have no defined units. This is somewhat surprising; if

a method has pretty good empirical results relative to other estimators, I'd be inclined

to use it, even if it's not entirely intellectually satisfying (Breiman, 2001). But averag-

ing partial regression coefficients is intellectually satisfying if causal modeling because

both a unit difference in X_j and the parameter estimated by $b_{j.m}$ is the same among

models.

Both Cade (2015) and Banner and Higgs (2017) consider model averaging the pre-

dicted outcome to be sensible, because the estimated parameter is the same among

models. This computation is also uncontroversial in the applied statistics literature.

But if Cade's interpretation of partial regression coefficients is correct (that these con-

tain partial units of X_j), how do we generate a prediction with sensible units, as this

computation requires postmultiplication of the model matrix, which includes full units

of X, by the vector of coefficients, which have partial units of X? Cade's interpreta-

tion implies that all model-averaging is meaningless and the rich literature developed

over the last twenty five years should simply be rejected. Indeed, if we accept Cade's

interpretation, then we must throw out additional tools in our kit. For example, meta-

analysis requires averaging effects over multiple studies, many of which have been con-

ditioned on different sets of covariates.

Given the long and rich history of model averaging within several applied fields, in-

cluding economics and epidemiology, its application to a diverse array of problems, and

its relationship to other well known methods, model averaging as a method probably does not need defending. Here, I am not advocating model averaging as the best choice among the many for estimating effect parameters, but simply defending it as a meaningful choice. Ultimately, this may prove to be an empirical issue, which can often only be obtained via simulation, where the true effects are known.

Acknowledgments

I thank three colleagues and multiple reviewers for improving the clarity of the manuscript.

References

Banner, K. M. and M. D. Higgs, 2017. Considerations for assessing model averaging of regression coefficients. *Ecological Applications* **27**:78–93.

Berger, J. O., L. R. Pericchi, J. K. Ghosh, T. Samanta, F. De Santis, J. O. Berger, and L. R. Pericchi, 2001. Objective Bayesian methods for model selection: introduction and comparison. *Lecture Notes-Monograph Series* pages 135–207.

Breiman, L., 2001. Statistical Modeling: The Two Cultures. *Statistical Science* **16**:199–231.

Burnham, K. P. and D. R. Anderson, 2002. Model selection and multimodel inference, 2nd Edition. Springer-Verlag, New York, NY.

Cade, B. S., 2015. Model averaging and muddled multimodel inferences. *Ecology* **96**:2370–2382.

Candolo, C., A. C. Davison, and C. G. B. Demétrio, 2003. A note on model uncer-

tainty in linear regression. *Journal of the Royal Statistical Society: Series D (The Statistician)* **52**:165–177.

Consonni, G. and P. Veronese, 2008. Compatibility of Prior Specifications Across Linear Models. *Statistical Science* **23**:332–353.

Draper, D., 1999. Comment—Bayesian model averaging: a tutorial. *Statistical Science* **14**:405–409.

Freckleton, R. P., 2010. Dealing with collinearity in behavioural and ecological data: model averaging and the problems of measurement error. *Behavioral Ecology and Sociobiology* **65**:91–101.

Friedman, J., T. Hastie, and R. Tibshirani, 2010. Regularization paths for generalized linear models via coordinate descent. *Journal of Statistical Software* **33**:1–22.

Hansen, B. E., 2007. Least squares model averaging. *Econometrica* **75**:1175–1189.

Hjort, N. L. and G. Claeskens, 2003. Frequentist model average estimators. *Journal of the American Statistical Association* **98**:879–899.

Hoeting, J. A., D. Madigan, A. E. Raftery, and C. T. Volinsky, 1999. Bayesian model averaging: a tutorial. *Statistical Science* **14**:382–417.

Liang, H., G. Zou, A. T. K. Wan, and X. Zhang, 2011. Optimal weight choice for frequentist model average estimators. *Journal of the American Statistical Association* **106**:1053–1066.

Møller, A. P. and M. D. Jennions, 2002. How much variance can be explained by ecologists and evolutionary biologists? *Oecologia* **132**:492–500.

Pearl, J., 2009. Causal inference in statistics: An overview. *Statistics Surveys* **3**:96–146.

R Core Team, 2015. R: A Language and Environment for Statistical Computing. R
  Foundation for Statistical Computing, Vienna, Austria.

Raftery, A. E., D. Madigan, and J. A. Hoeting, 1997. Bayesian model averaging for
  linear regression models. *Journal of the American Statistical Association* **92**:179–191.

Raftery, A. E. and Y. Zheng, 2003. Discussion: performance of bayesian model averag-
  ing. *Journal of the American Statistical Association* **98**:931–938.

Roulin, A. and L.-F. Bersier, 2007. Nestling barn owls beg more intensely in the pres-
  ence of their mother than in the presence of their father. *Animal Behaviour* **74**:1099–
  1106.

Shipley, B., 2002. Cause and correlation in biology: a user's guide to path analysis,
  structural equations and causal inference. Cambridge: Cambridge University Press.

Walker, J. A., 2014. The effect of unmeasured confounders on the ability to estimate
  a true performance or selection gradient (and other partial regression coefficients).
  Evolution **68**:2128–2136.

Wright, S., 1918. On the nature of size factors. *Genetics* **3**:367–374.

Wright, S., 1934. The method of path coefficients. *The Annals of Mathematical Statis-
  tics* **5**:161–215.

Zhang, X., G. Zou, and H. Liang, 2014. Model averaging and weight choice in linear
  mixed-effects models. *Biometrika* **101**:205–218.

Zuur, A. F. and E. N. Ieno, 2016. A protocol for conducting and presenting results of
  regression-type analyses. *Methods in Ecology and Evolution* **7**:636–645.