**Active trachoma cases in the Solomon Islands have varied polymicrobial community structures but do not associate with individual non-chlamydial pathogens of the eye.**

Robert M R Butcher[1]*, Oliver Sokana[2], Kelvin Jack[2], Eric Kalae[4], Leslie Sui[4], Charles Russell[5], Joanna Houghton[1], Christine Palmer[1], Martin J Holland[1], Richard T Le Mesurier[6], Anthony W Solomon[1,3], David C W Mabey[1,3], Chrissy h. Roberts[1].

Author contact details:

[1] Clinical Research Department, London School of Hygiene & Tropical Medicine, Keppel Street, London WC1E 7HT, UK

[2] Eye Department, Solomon Island Ministry of Health and Medical Services, Honiara, Solomon Islands

[3] Hospital for Tropical Diseases, University College London Hospitals, Mortimer Market, London

[4] Primary Care Department, Lata Hospital, Lata, Santa Cruz, Solomon Islands

[5] Bellona Rural Health Centre, Bellona, Solomon Islands

[6] The Fred Hollows Foundation, Carlton, Victoria, Australia

**Abstract**

Background

Several non-chlamydial microbial pathogens are associated with clinical signs of active trachoma in trachoma endemic communities with a low prevalence of ocular *Chlamydia trachomatis* (*Ct*) infection. We observed a low prevalence of *Ct* in the Solomon Islands, therefore hypothesised that the high prevalence of active trachoma could be explained by a common non-chlamydial infection or by a dominant polymicrobial community dysbiosis.

Methods

We studied DNA from conjunctival swabs collected from 257 Solomon Islanders with active trachoma and 257 matched controls. Droplet digital PCR was used to test for *Adenoviridae, Haemophilus influenzae, Moraxella catarrhalis, Staphylococcus aureus,* coagulase-negative *Staphylococcus* and *Streptococcus pneumoniae*. Polymicrobial community diversity and composition were studied by sequencing of hypervariable regions of the 16S ribosomal ribonucleic acid gene in a subset of 54 cases and 53 controls.

Results

Although *Ct* was associated with active trachoma, the number of infections was low (cases: 3.9%, controls: 0.4%). Estimated prevalence (cases, controls) of each nonchlamydial infection was as follows: *S. aureus* (1.9%, 1.9%), *Adenoviridae* (1.2%, 1.2%), coagulase-negative *Staphylococcus (5.8%, 4.3%), H. influenzae* (7.4%, 11.7%)*, M. catarrhalis* (2.3%, 4.7%) and *S. pneumoniae* (7.0%, 6.2%). There was no statistically significant association between clinical signs of trachoma and presence or load of any of the non-*Ct* infections that were assayed. Inter-individual variations in the conjunctival microbiome were characterised by differences in the levels of *Corynebacterium*, *Proprionibacterium*, *Helicobacter* and *Paracoccus*, but diversity and relative abundance of these specific genera did not differ between cases and controls.

53 Discussion

54 It is unlikely that the prevalent trachoma-like follicular conjunctivitis in the Solomon Islands has a

55 dominant bacterial aetiology. Before implementing community-wide azithromycin distribution for

56 trachoma, policy makers should consider that clinical signs of trachoma can occur without the

57 involvement of any azithromycin-susceptible organism.

**Introduction**

Trachoma, caused by *Chlamydia trachomatis* (*Ct*), is the leading infectious cause of preventable blindness (1), and is targeted for elimination as a public health problem by 2020 through the SAFE strategy (**S**urgery, **A**ntibiotics, **F**acial cleanliness and **E**nvironmental improvement). The decision to implement community-wide trachoma control interventions, which include mass drug administration (MDA) with azithromycin, is based on population prevalence estimates of one clinical sign of active trachoma (trachomatous inflammation–follicular [TF]) in the 1–9-year-old age group (2). In the Solomon Islands, the prevalence of active trachoma is sufficient to warrant MDA but the prevalence of TT suggests trachoma may not pose a significant public health problem. A recent study of trachoma in a treatment-naïve population of the Solomon Islands estimated the TF prevalence at 26% but, surprisingly, conjunctival *Ct* infection was detected in only 1.3% of 1–9 year olds (3). Whilst *Ct* infection is not always detectable in TF cases (4, 5), infection prevalence in the Solomon Islands is far lower than is seen in other countries with similar TF prevalence (3).

*Ct* is not the only cause of follicular conjunctivitis. *Moraxella lacunata*, Parinaud's oculo-glandular syndrome, atopic conjunctivitis (6), adenovirus (7), and other *Chlamydia* serotypes (8) and species (9), among others (10), have been suggested as possible causes of TF in trachoma-endemic populations. A greater diversity of pathogens can be cultured from the conjunctivae of people with TF (10) and, among bacterial species isolated during a study in Tanzanian children, *Streptococcus pneumoniae*, *Haemophilus influenzae* B and *Haemophilus influenzae* non-type B associated more strongly than *Ct* with clinical signs of active trachoma (10). In communities which had received MDA in The Gambia, *S. pneumoniae* and *H. influenzae* type b infection correlated closely with signs of active trachoma, whereas *Moraxella catarrhalis* and *Staphylococcus aureus* did not (11). Common non-chlamydial pathogens are also associated with trachomatous scarring (TS) (12) and recurrence of trachomatous trichiasis (TT) after surgery (13).

Although traditionally thought to be 'sterile', several studies have now described polymicrobial communities colonising the conjunctival epithelium (14, 15). *Staphylococcus*, *Streptococcus*, *Haemophilus* and *Moraxella* genera can be readily cultured from swabs taken from the inferior fornix (10–12). Attempts to detect bacteria from the conjunctiva have shown community diversity and composition to vary significantly between individuals. It is unclear whether a 'core' microbiota persists at the conjunctiva, but profiles closely related to that of the skin have been reported (16). *Corynebacterium*, *Propionibacterium*, *Staphylococcus* and *Streptococcus* have been consistently dominant, whereas other genera such as *Acinetobacter, Brevundimonas, Pseudomonas, Bradyrhizobium, Sphingomonas, Bacillus, Simonsiella* and *Elizabethkingia* have been identified more sporadically (17–19). Conjunctival polymicrobial community composition is known to vary with age and season (20) and appears to be responsive to external stimuli such as regular contact lens wear (21), however, significant associations with disease have yet to be described (20, 22).

We hypothesised that clinical signs of active trachoma in the Solomon Islands where ocular *Ct* is uncommon could be explained by a common non-chlamydial infection or by a dominant polymicrobial community dysbiosis.

**Methods**

Study ethics

Ethical approval was granted by the London School of Hygiene & Tropical Medicine (6319/6360) and the Solomon Islands National Health Research (HRC13/18) Ethics Committees. A parent or guardian provided written consent for each child participant.

*110* **Table 1.** Oligonucleotides used in this study, with assay concentrations derived from *in vitro* optimisation.

| Organism / target | Oligo | Sequence (5'-3') | Concentration in final assay (nM) | Amplicon size | Ref |
|---|---|---|---|---|---|
| *C. trachomatis / Plasmid open reading frame 2 (pORF2)* | F | CAG CTT GTA GTC CTG CTT GAG AGA | 900 | 109 | (3, 23, 24) |
| | R | CAA GAG TAC ATC GGT CAA CGA AGA | 900 | | |
| | Probe | [FAM] CCC CAC CAT TTT TCC GGA GCG A [BHQ1] | 200 | | |
| *S. aureus* / Staphylococcal protein A (*SpA*) | F | CAG CAA ACC ATG CAG ATG CTA | 900 | 101 | (25, 26) |
| | R | CGC TAA TGA TAA TCC ACC AAA TAC A | 900 | | |
| | Probe | [VIC] TCA AGC ATT ACC AGA AAC [MGBNFQ] | 250 | | |
| Coagulase-negative *Staphylococcus* / translation elongation factor (*tuf*) | F | TAT CCA CGA AAC TTC TAA AAC AAC TGT TAC T | 450 | 204 | (25) |
| | R | TCT TTA GAT AAT ACG TAT ACT TCA GCT TTG AAT TT | 450 | | |
| | Probe | [FAM] TAT TAG ACT ACG CTG AAG CTG GTG ACA ACA T [BHQ1] | 125 | | |
| *S. pneumoniae /* N-acetylmuramoyl-L-alanine amidase (*lytA*) | F | ACG CAA TCT AGC AGA TGA AGC A | 500 | 75 | (27) |
| | R | TCG TGC GTT TTA ATT CCA GCT | 500 | | |
| | Probe | [FAM] GCC GAA AAC GCT TGA TAC AGG GAG [BHQ1] | 150 | | |
| *H. influenzae /* L-fuculokinase (*fucK*)† | F | ATG GCG GGA ACA TCA ATG A | 900 | 102 | (28) |
| | R | ACG CAT AGG AGG GAA ATG GTT | 900 | | |
| | Probe* | [FAM] CGg TAa TTg GGa TCc AT [BHQ1] | 125 | | |
| *Adenoviridae /* capsid assembly protein (hexon) | F | GCC ACG GTG GGG TTT CTA AAC TT | 500 | 132 | (29) |
| | R | GCC CCA GTG GTC TTA CAT GCA CAT C | 500 | | |
| | Probe | [HEX] TGC ACC AGA CCC GGG CTC AGG TAC TCC GA [BHQ1] | 125 | | |
| *M. catarrhalis /* outer membrane protein (*copB*) | F | GTG AGT GCC GCT TTT ACA ACC | 900 | 72 | (30, 31) |
| | R | TGT ATC GCC TGC CAA GAC AA | 900 | | |
| | Probe | [HEX] TGC TTT TGC AGC TGT TAG CCA GCC TAA [BHQ1] | 250 | | |

*111* * Lower case symbols denote locked nucleic acid bases

112 † These primers do not differentiate between encapsulated and noncapsulated subtype

Study population

The samples tested during this study were a sub-set of specimens from a population-based trachoma prevalence survey of 3674 people (1135 children aged 1–9 years) in 32 clusters from Temotu and Rennell & Bellona Provinces, Solomon Islands; data from that survey are presented elsewhere (3). Sterile swabs were passed three times over the right conjunctiva of every child before storage in 300 μL RNALater® (Life Technologies). Swabs were kept cool in the field for up to 24hrs then frozen (3). Inclusion criteria for cases (n = 257) were age 1–9 years, detectable human DNA in the conjunctival swab and a field grade of TF or trachomatous inflammation—intense (TI) in the right eye. An equal number (n = 257) of age, gender and island matched controls without TF or TI were selected using the 'e1071' R package. A random subset of cases (n = 54) and controls (n = 53) were further characterized using V1-V3 16S rRNA gene sequencing. Field controls (clean swabs passed within 20 cm of seated participants, without touching them, and then treated identically to subjects' specimens), extraction controls (extraction carried out in the absence of a specimen and PCR carried out on the eluate) and PCR controls (PCR carried out in the absence of any added material) were used to determine background levels of microbial contamination in the 16S rRNA gene sequencing protocol.

Droplet digital PCR

All individuals in this sample have previously been tested for *Ct* and human RPP30 using DNA extracted from conjunctival swabs with the Qiagen AllPrep DNA/RNA kit. We chose not to use mechanical lysis on these low biomass specimens due to the reported lower yields compared to chemical lysis (32).

Duplex ddPCR assays were developed for (1) *Adenoviridae* (29) and *S. pneumoniae* (27)*, (2) *H. influenzae* (28) and *M. catarrhalis* (30, 31), and (3) *S. aureus* and coagulase-negative *Staphylococcus* (25, 26) based on published assays. Assay performance was assessed by repeat

testing of synthetic standard dilution series' and cultured pathogen material before rolling out to

clinical samples (Supplementary Table 1). Each assay contained 10 μL 2× ddPCR Supermix for

probes (Bio-rad, Hemel Hempstead, UK), primers and probes at custom concentrations (Table 1)

and 8 μL of template DNA. Thermal cycling was 10'00'' at 95°C then 40x (0'15'' at 95°C | 1'00'' at

60°C) then 12'00'' at 98°C.(23) A positive result for a clinical specimen was defined as >95%

confidence of non-zero target load, as described previously.(23)

<u>16S rRNA gene sequencing</u>

An approximately 530-bp region of the 16S ribosomal RNA (rRNA) gene (variable regions 1-3) was

amplified using forward (modified 27F; 5'-[adaptor]-AGAGTTTGGATCCTGGCTCAG-3') and custom

barcoded reverse primers (534R; 5'-[adaptor]-[barcode]-

AGTCAGTCAGCCATTACCGCGGCTGCTGG-3'). Each 15 μL reaction contained 7.5 μL 2x

Phusion High Fidelity Master Mix (New England Biosciences, MA, USA), 0.45 μL DMSO, 0.1 μM

primers and 5.55 μL DNA. The thermal cycling conditions were 0'30'' at 98°C then 31x (0'10'' at

98°C | 0'30'' at 62°C | 0'15'' at 72°C) then 7'00'' at 72°C. Amplicons were cleaned using 0.6 v/v

AMPure XP beads (Beckman Coulter, CA, USA), quantified using Qubit (Thermo Fisher Scientific,

MA, USA) then pooled at equimolar concentrations. The 4 nM sequencing library was mixed 0.75

v/v with a 4 nM Phi-X control library (Illumina, CA, USA). 3 μL of 100 μM custom read primers were

mixed in to wells 12 (Read 1; CTACACTATGGTAATTGTAGAGTTTGGATCCTGGCTCAG), 13

(Index; CCAGCAGCCGCGGTAATGGCTGACTGACT) and 14 (Read 2;

AGTCAGTCAGCCATTACCGCGGCTGCTGG) of the MiSeq reagents cartridge before 2 x 300 bp

paired-end sequencing with the 600-cycle MiSeq v3 sequencing reagent kit on the MiSeq platform

using a standard protocol (Illumina, CA, USA). 96 uniquely barcoded specimens were run in

multiplex on each MiSeq run.

<u>Data analysis</u>

ddPCR data were analysed using R version 3.2.2 (33). Binomial univariate regression was used to test the relationship between each individual infection and active trachoma. The most accurate final multivariate model, determined by lowest Akaike Information Criterion (AIC) value, was determined by step-wise removal of variables from a binomial multivariate regression model incorporating all tested pathogens. The chance of the *Ct* association occurring due to chance was assessed by counting the number of significant results obtained from randomly re-ordering the *Ct* data 1000 times.

Raw 16S amplicon sequences were directly assigned to genera using Illumina BaseSpace '16S Metagenomics' app version 1.0.1.0, which uses algorithms from Wang and colleagues (34). Two clinical specimens yielded fewer than 1000 reads in total and were removed from the analysis. Amplicons were sequenced from six no-template control (NTC) specimens (two 'field' controls, two 'extraction' controls, two 'PCR' controls, defined above) to identify any contaminants endogenous to the collection, PCR or sequencing process. Any genera represented by more than 600 reads in the six combined NTCs (average 100 reads per NTC) were eliminated from the clinical specimen analysis. This resulted in the removal of 21 genera, which accounted for 95.1% of the reads from NTCs and 88.3% of the reads from clinical specimens. The genera removed included common contaminants of reagent kits (35) such as *Pelomonas*, as well as some previously described conjunctival microbiota constituents such as *Brevundimonas*, *Staphylococcus* and *Streptococcus* (17, 20).

Between-group differences in the Shannon Diversity Index (a measure of diversity which increases with increasing species abundance and evenness) and Inverse Simpson Index (another measure of diversity where 1 is infinite diversity and 0 is no diversity) were compared using a t-test. Discriminant Analysis of Principal Components (DAPC), a multivariate clustering method to analyse complex datasets, was performed using the 'adegenet' package in R (36) and cross-validation was

used to determine the optimal number of principal components to include in a discriminant function

aimed at separating cases from controls on the basis of their polymicrobial community structures.

**Results**

<u>Specimen set demographics</u>

A total of 257 cases and 257 controls (n = 514) were tested. All 257 cases had TF, and one also

had TI. Case status was defined by clinical signs in the right eye but respectively 8/257 (3.1%) and

236/257 (91.8%) of the controls and cases had TF in the left eye. Both case and control groups

were 38% female. Mean age was 5.6 years (cases) and 5.5 years (controls, student's t test p =

0.76). There was no significant difference between cases and controls in terms of the clusters

represented (Kolmogorov-Smirnov test p = 0.97) and all 32 clusters were represented. The case

specimens had higher loads of human DNA than the controls (18030 versus 9354 copies/swab; p =

0.00003). There were no significant differences in age, gender or location within the subset selected

for 16S-amplicon sequencing.

<u>Quantitative PCR tests for ocular pathogens</u>

In this study, 19.8% of children had evidence of infection with at least one of the targeted organisms

(table 2). We considered the prevalence of *Ct* in this sample set to be too low (96.1% of TF cases

were infection-negative) to account for the level of active disease. The prevalence of *Adenoviridae*

(1.2%) and *S. aureus* (1.9%) were both very low. There was no association between active

trachoma and infection with *H. influenzae, S. pneumoniae, S. aureus,* coagulase-negative

*Staphylococcus* or *M. catarrhalis. Ct* infection was associated with active trachoma (logistic

regression p = 0.026; odds ratio: 10.4). The association between *Ct* and active trachoma was still

significant in a multivariate analysis (p = 0.025). The permuted p-value for the association between

*Ct* and active trachoma was highly significant (p = 0.001). Active disease was neither associated

with infection with at least one pathogen (any pathogen, p = 0.185) nor the number of concurrent

infections in the same eye (number of concurrent pathogens, p = 0.207). Infectious loads of those

who tested positive are shown figure 2. Despite some numerical differences between groups, none

of the differences were statistically significant.

**Figure 2.** Target copies per swab of each pathogen identified in conjunctival swabs collected from children with and

without active trachoma in the Solomon Islands. Numbers show p-values for logistic regression comparison between

active trachoma case and control groups for each pathogen. None of the differences observed were statistically

significant.

**Table 2.** Cases of infection in specimens from children with and without active disease. The relationship of those

infections to trachoma has been tested with univariate logistic regression, and then stepwise removal of variables from

multivariate regression model was used to determine the final multivariate regression model that provided the best fit for

the data.

| Pathogen | ddPCR result | No TF/TI (%; n=257) | TF/TI (%; n=257) | Total (%; n=514) | Univariate odds ratio (95% CI) | Univariate p-value | Multivariate odds ratio (95% CI) | Multivariate p-value |
|---|---|---|---|---|---|---|---|---|
| *Adenoviridae* | Positive | 3 (1.2) | 3 (1.2) | 6 (1.2) | 1.00 (0.67 – 1.50) | 1.000 | - | - |
| | Negative | 254 (98.8) | 254 (98.8) | 508 (98.8) | | | | |
| *Chlamydia trachomatis** | Positive | 1 (0.4) | 10 (3.9) | 11 (2.1) | 10.36 (1.96 – 190.92) | **0.026** | 10.64 (2.15 – 196.00) | **0.025** |
| | Negative | 256 (99.6) | 247 (96.1) | 503 (97.9) | | | | |
| Coagulase-negative *Staphylococcus* | Positive | 15 (5.8) | 11 (4.3) | 26 (5.1) | 0.72 (0.32 – 1.59) | 0.423 | - | - |
| | Negative | 242 (94.2) | 246 (95.7) | 488 (94.9) | | | | |
| *Haemophilus influenzae* | Positive | 19 (7.4) | 30 (11.7) | 49 (9.5) | 1.66 (0.91 – 3.07) | 0.101 | - | - |
| | Negative | 238 (92.6) | 227 (88.3) | 465 (90.4) | | | | |
| *Moraxella catarrhalis* | Positive | 6 (2.3) | 12 (4.7) | 18 (3.5) | 2.05 (0.78 – 5.96) | 0.158 | 2.13 (0.81 – 6.20) | 0.137 |
| | Negative | 251 (97.7) | 245 (95.3) | 496 (96.5) | | | | |
| *Staphylococcus aureus* | Positive | 5 (1.9) | 5 (1.9) | 10 (1.9) | 1.00 (0.28 – 3.64) | 1.000 | - | - |
| | Negative | 252 (98.1) | 252 (98.1) | 504 (98.1) | | | | |
| *Streptococcus pneumoniae* | Positive | 18 (7.0) | 16 (6.2) | 34 (6.6) | 0.88 (0.43 – 1.77) | 0.723 | - | - |
| | Negative | 239 (93.0) | 241 (93.8) | 480 (93.4) | | | | |
| Any pathogen | Positive | 45 (17.5) | 57 (22.2) | 102 (19.8) | 0.75 (0.48 – 1.15) | 0.185 | Not included | |
| | Negative | 212 (82.4) | 200 (77.8) | 412 (80.2) | | | | |
| Different pathogen species in same eye | 0 | 212 (82.4) | 200 (77.8) | 412 (80.2) | 1.21 (0.90 – 1.63) | 0.207 | Not included | |
| | 1 | 28 (10.9) | 35 (13.6) | 63 (12.3) | | | | |
| | >1 | 17 (6.6) | 22 (8.6) | 39 (7.6) | | | | |

*Data taken from Butcher *et al.*(3)

16S rRNA gene sequencing

The total number of genera identified across all clinical specimens and NTCs was 659, however,

125/659 (19%) had a cumulative total of ten reads or fewer. The median number of reads per

clinical specimen was 55,104 (IQR: 40,705 – 89,541) and the median percentage of reads mapped

to genus level was 51.9%. The median number of genera identified was 151 (IQR: 117 – 212) per

clinical specimen. Following removal of presumed contaminants, approximately 40% of cleaned

reads were assigned to genera that each constituted less than 1% of the total polymicrobial

community. 40 genera were represented by at least 1% of remaining reads in clinical samples.

The diversity of bacterial communities was in general low; the mean Inverse Simpsons Diversity

index over all specimens was 0.061 (IQR: 0.046 – 0.092). There was no significant difference in

alpha-diversity between cases and controls by either measure of diversity. The median Inverse

Simpson's Diversity index in cases was 0.061 [IQR: 0.048 – 0.095] versus controls 0.060 [IQR:

0.044 – 0.078] (Student's t-test p = 0.56). The median Shannon Diversity Index was 3.38 in cases

and 3.37 in controls (Student's t-test p = 0.96). In cases of active trachoma, the most dominant

bacterial genera were *Corynebacterium* (12.0% of total reads), *Proprionibacterium* (6.2%) and

*Helicobacter* (4.8%). In controls, the most dominant genera were *Corynebacterium* (13.9%),

*Paracoccus* (5.2%), *Proprionibacterium* (4.7%), and *Neisseria* (4.1%) (supplementary figure 2). The

genus level membership of the bacterial community varied between cases and controls. Of 21

genera found in specimens from those with active trachoma, 10 were not found in controls,

including *Helicobacter*, *Mesoplasma, Brachybacterium* and *Haemophilus*. Of 23 genera found in

controls, 12 were not found in cases, including *Neisseria, Prevotella, Rhodococcus* and

*Porphyromonas*.

Principal Components (PC) Analysis (figure 2) revealed that 47% of the total variation in the

microbiomes of children in the Solomon Islands is dominated by *Corynebacterium* (PC1),

*Paracoccus* (PC2), *Propionibacterium* (PC2,PC3) and *Helicobacter* (PC3). While individual cases or

controls have distinctive profiles dominated by these genera, the majority of cases and controls are

indistinguishable (figure 2) using the first three PCs (explaining 45% of total variation). We

condensed PCs 1-20 into a single discriminant function, which explained 92% of variation in the

data. In this analysis, the genus-level polymicrobial community structure is significantly different

between cases and controls (logistic regression p = 0.0000013) (figure 3). That discriminant

function was dominated by a number of genera such as *Curvibacterium* and *Mesoplasma*

(Supplementary Figure 4). Cross validation to predict group membership using discriminant

functions of between 10 and 90 PCs had a low success rate (median 53.3% success, range: 50.1 –

62.3%). This suggests that while differences do exist between the polymicrobial communities in

cases and controls, they are too subtle and varied to be predictive of phenotype, at least when

working with this number of specimens.

**A**

**B**

**Figure 2. (A)** First and second and **(B)** second and third principal components describing variation between the 16S

sequences identified in Solomon Island children with and without active trachoma. Dark blue spots indicate controls. Light

blue spots indicate cases. Red arrows show principal component loadings.

*286*

*287*   **Figure 3:** Discriminant analysis of the association of 20 combined principal components with active trachoma, showing a

*288*   significant discrimination of phenotype groups (p = 0.0000013).

*289*

*290*

**Discussion**

Follicular conjunctivitis meeting WHO criteria for the trachoma phenotype "TF" is highly prevalent in the Solomon Islands, but the prevalence of *Ct* infection is curiously low. Although *Ct* was the only pathogen to associate with TF in this study, the prevalence was much lower than may be expected of a population with this level of TF (3). By ddPCR testing for a number of bacteria and viruses that have previously been linked to TF, we can discount the possibility that any of them can account either singly or *en masse* for the high TF prevalence. Both *S. pneumoniae* and *H. influenzae* were detected at moderate prevalence in our population, whilst we found very few cases of *Adenoviridae* and *M. catarrhalis* infection. Comparator data from other populations are scarce, but it is clear that staphylococcal species were detected in the conjunctivae in the Solomon Islands at a substantially lower prevalence (7%) than in children in either Tanzania (14.8% *S. epidermidis*) (10), Sierra Leone (20% *S. aureus,* 29% coagulase-negative *Staphylococcus*) (37) or The Gambia (post MDA, 14.7% *S. aureus*) (11). Through 16S amplicon sequencing and community profiling, we have shown that there is apparently no dominant bacterial genus associated with this disease. The dominant features of variation in conjunctival bacterial communities in Solomon Islands children (*Corynebacterium, Propionibacterium)* have consistently been identified in other studies. Other genera (e.g., *Paracoccus*, *Helicobacter, Haemophilus)* have not previously been identified in 16S studies, whilst some reported in other studies (e.g., *Simonsella, Pseudomonas*) were not found in these specimens.(17–20) Many studies have sequenced the V3-4 region of the 16S gene whereas we targeted V1-3. V region choice has been shown to have an impact on the genera identified at other mucosal sites (38) and, while no data have yet emerged on how this affects profiles from the eye, it is possible this could account for some of the differences. Consistent with previous studies (35), we found a background of genera that amplified in NTCs that also appeared in clinical specimens. The microbiological biomass at the conjunctiva is known to be very low, even compared with other nearby sites such as skin or oral mucosa (18). When true resident bacteria are scarce, 16S rRNA gene PCR readily amplifies reagent contaminants. We took stringent measures to exclude reagent contaminants, but this resulted in the removal of some important genera such as

*Staphylococcus* and *Streptococcus.* Among the biggest contributors to between-conjunctiva

variation were *Corynebacterium* and *Propionibacterium*, but these genera did not differentiate TF

cases from controls (figure 2). Previous investigations of the role of the conjunctival microbiome in

ocular disease have also not shown significant differences. People with ocular manifestations of

rosacea, Sjögren's syndrome and healthy controls did not have significant differences in diversity or

relative abundance of key phyla identified across all three groups (22). In The Gambia, there was an

increased abundance of *Haemophilus* in cases of TF, as compared to controls. The abundance of

*Corynebacterium* and *Streptococcus* varied between scarred and non-scarred individuals. However,

neither trachoma-related difference was significant (20).

The microbiota of these children were heterogeneous and had subtle variations that appeared to

reach significant association with TF when the full community structure was considered in

comparative statistical models. It remains possible (though perhaps unlikely) that a multitude of

factors, operating at the level of single bacterial species, polymicrobial communities or viral species,

all contribute to the presentation of the phenotype. It is perhaps more likely that as yet unidentified

viral or allergic causes could explain the prevalence of TF in the Solomon Islands.

There are some limitations to our study. Although the ddPCR assays are based on validated assays

and appear to be accurate (supplementary table 1), they have not been formally evaluated in this

format. By not using mechanical lysis in our extraction process, the absolute prevalence of some

difficult-to-lyse Gram positive genera may have been under-estimated, although this would not

impair the comparison of cases to controls, between which protocols were consistent. 16S amplicon

sequencing of low biomass samples is known to result in amplification of reagent and environmental

contaminants (35). This study focused on gross differences in community structure between those

with and without disease and these should be independent of contaminating reads. We also

employed stringent quality control measures to ensure data were not confused by artefacts of the

sequencing process.

Public health scenarios such as the one in the Solomon Islands will become increasingly common

as trachoma elimination programmes reduce the global prevalence of *Ct,* and the positive predictive

value of TF for ocular Ct infection declines as a result (39). MDA might be inappropriately delivered

when clinical signs of trachoma are the only indicator used for programmatic decision making. While

MDA is effective for the treatment of trachoma (40), mass antibiotic exposure may increase

macrolide resistance in nonchlamydial bacteria (41) and theoretically could make the population

more susceptible to later infections by preventing accumulation of acquired immunity to *Ct* (42) and

decisions to undertake such a program should therefore be carefully considered. The case for using

tests for *Ct* infection during trachoma surveys is strengthened by our data. The value of nucleic acid

amplification tests for detecting non-chlamydial infection remains questionable, as multiple

infectious agents might be important, and these are likely to differ between populations. Key

questions now include whether non-chlamydial follicular conjunctivitis such as this responds to MDA

and whether it is linked to incident or progressive scarring which may lead to blindness.

Acknowledgements

We thank the residents of the communities who took part in this study, Victoria Miari and Emma

Cobb (Pathogen Molecular Biology Department, London School of Hygiene & Tropical Medicine,

UK) for providing bacterial culture isolates for PCR assay validation, and David Nelson (Indiana

University, Indianapolis, US) for the provision of 16S sequencing primers and protocols.

Financial statement

Fieldwork was jointly funded by the United Kingdom's Department for International Development

Global Trachoma Mapping Project grant (ARIES: 203145) to Sightsavers, and by the Fred Hollows

Foundation, Australia (1041). Laboratory costs were funded by the Fred Hollows Foundation,

Australia (1041).

RMRB and AWS were funded by a Wellcome Trust Intermediate Fellowship to AWS (098521).

OS, KJ, EK, LS and CR were employed by the Solomon Islands Ministry of Health and Medical

Services for the duration of the survey.

MJH is supported by the Wellcome Trust (093368/Z/10/Z).

ChR is supported by the Wellcome Trust Institutional Strategic Support Fund (105609/Z/14/Z).

Author Contributions

Conceived and designed the study: RMRB, OS, RTLM, AWS, DCWM, ChR.

Performed the fieldwork: RMRB, OS, EK, KJ, LS, CR.

Performed the experiments: RMRB, MJH, JH, CP, ChR.

Analysed the data: RMRB, ChR.

Wrote the manuscript: RMRB, ChR.

Revised and approved the manuscript: RMRB, OS, EK, KJ, LS, CR, JH, CP, MJH, RTLM, AWS,

DCWM, ChR.

References

1.    Bourne RA, Stevens GA, White RA, Smith JL, Flaxman SR, Price H, Jonas JB, Keeffe J,

Leasher J, Naidoo K, Pesudovs K, Resnikoff S, Taylor HR, Vision Loss Expert G. 2013.

Causes of vision loss worldwide, 1990-2010: a systematic analysis. Lancet Glob Heal

1:e339-49.

2.    World Health Organization. 2003. Report of the 2nd Global Scientific Meeting on Trachoma.

25-27 August. Geneva, Switzerland.

3.   Butcher RMR, Sokana O, Jack K, Macleod CK, Marks ME, Kalae E, Sui L, Russell C, Tutill

HJ, Williams RJ, Breuer J, Willis R, Le Mesurier RT, Mabey DCW, Solomon AW, Roberts CH.

2016. Low Prevalence of Conjunctival Infection with Chlamydia trachomatis in a Treatment-

Naïve Trachoma-Endemic Region of the Solomon Islands. PLoS Negl Trop Dis 10:e0004863.

4.   Thein J, Zhao P, Liu H, Xu J, Jha H, Miao Y, Pizzarello L, Tapert L, Schachter J, Mabon M,

Osaki-Holm S, Lietman T, Paxton A. 2002. Does clinical diagnosis indicate ocular chlamydial

infection in areas with a low prevalence of trachoma? Ophthalmic Epidemiol 9:263–9.

5.   Grassly NC, Ward ME, Ferris S, Mabey DC, Bailey RL. 2008. The natural history of trachoma

infection and disease in a Gambian cohort with frequent follow-up. PLoS Negl Trop Dis

2:e341.

6.   Dawson CR, Jones BR, Tarizzo ML, World Health Organization. 1981. Guide to trachoma

control in programmes for the prevention of blindness. Geneva, Switzerland.

7.   Jhanji V, Chan TCYY, Li EYMM, Agarwal K, Vajpayee RB. 2015. Adenoviral

keratoconjunctivitis. Surv Ophthalmol 60:435–443.

8.   Andersson P, Harris SR, Smith HMBS, Hadfield J, O'Neill C, Cutcliffe LT, Douglas FP, Asche

LV, Mathews JD, Hutton SI, Sarovich DS, Tong SYC, Clarke IN, Thomson NR, Giffard PM.

2016. Chlamydia trachomatis from Australian Aboriginal people with trachoma are

polyphyletic composed of multiple distinctive lineages. Nat Commun 7:10688.

9.   Lietman T, Brooks D, Moncada J, Schachter J, Dawson C, Dean D. 1998. Chronic follicular

conjunctivitis associated with Chlamydia psittaci or Chlamydia pneumoniae. Clin Infect Dis

26:1335–40.

10.  Burton MJ, Hu VH, Massae P, Burr SE, Chevallier C, Afwamba IA, Courtright P, Weiss HA,

Mabey DCW, Bailey RL. 2011. What is causing active trachoma? The role of nonchlamydial

bacterial pathogens in a low prevalence setting. Invest Ophthalmol Vis Sci 52:6012–6017.

11.  Burr SE, Hart JD, Edwards T, Baldeh I, Bojang E, Harding-Esch EM, Holland MJ, Lietman

TM, West SK, Mabey DCW, Sillah A, Bailey RL. 2013. Association between ocular bacterial

carriage and follicular trachoma following mass azithromycin distribution in The Gambia.

PLoS Negl Trop Dis 7:e2347.

12. Hu VH, Massae P, Weiss HA, Chevallier C, Onyango JJ, Afwamba IA, Mabey DCW, Bailey RL, Burton MJ. 2011. Bacterial infection in scarring trachoma. Invest Ophthalmol Vis Sci 52:2181–6.

13. Burton MJ, Adegbola RA, Kinteh F, Ikumapayi UN, Foster A, Mabey DCW, Bailey RL. 2007. Bacterial infection and trachoma in the gambia: a case control study. Invest Ophthalmol Vis Sci 48:4440–4.

14. Kugadas A, Gadjeva M. 2016. Impact of Microbiome on Ocular Health. Ocul Surf 14:342–9.

15. Lu LJ, Liu J. 2016. Focus: Microbiome: Human Microbiota and Ophthalmic Disease. Yale J Biol Med 89:325.

16. Zhou Y, Gao H, Mihindukulasuriya KA, La Rosa PS, Wylie KM, Vishnivetskaya T, Podar M, Warner B, Tarr PI, Nelson DE, Fortenberry JD, Holland MJ, Burr SE, Shannon WD, Sodergren E, Weinstock GM. 2013. Biogeography of the ecosystems of the healthy human body. Genome Biol 14:R1.

17. Dong Q, Brulc JM, Iovieno A, Bates B, Garoutte A, Miller D, Revanna K V, Gao X, Antonopoulos DA, Slepak VZ, Shestopalov VI. 2011. Diversity of bacteria at healthy human conjunctiva. Invest Ophthalmol Vis Sci 52:5408–13.

18. Doan T, Akileswaran L, Andersen D, Johnson B, Ko N, Shrestha A, Shestopalov V, Lee CS, Lee AY, Van Gelder RN. 2016. Paucibacterial Microbiome and Resident DNA Virome of the Healthy Conjunctiva. Investig Opthalmology Vis Sci 57:5116.

19. Huang Y, Yang B, Li W. 2016. Defining the normal core microbiome of conjunctival microbial communities. Clin Microbiol Infect 22:643.e7-643.e12.

20. Zhou Y, Holland MJ, Makalo P, Joof H, Roberts CH, Mabey D, Bailey RL, Burton MJ, Weinstock GM, Burr SE. 2014. The conjunctival microbiome in health and trachomatous disease: a case control study. Genome Med 6:99.

21. Shin H, Price K, Albert L, Dodick J, Park L, Dominguez-Belloa MG. 2016. Changes in the eye microbiota associated with contact lens wearing. MBio 7:1–6.

22. de Paiva CS, Jones DB, Stern ME, Bian F, Moore QL, Corbiere S, Streckfus CF, Hutchinson DS, Ajami NJ, Petrosino JF, Pflugfelder SC. 2016. Altered Mucosal Microbiome Diversity and

Disease Severity in Sjögren Syndrome. Sci Rep 6:23561.

23.  Roberts CH, Last A, Molina-Gonzalez S, Cassama E, Butcher R, Nabicassa M, McCarthy E,

Burr SE, Mabey DC, Bailey RL, Holland MJ. 2013. Development and Evaluation of a Next-

Generation Digital PCR Diagnostic Assay for Ocular Chlamydia trachomatis Infections. J Clin

Microbiol 51:2195–203.

24.  Last AR, Roberts CH, Cassama E, Nabicassa M, Molina-Gonzalez S, Burr SE, Mabey DCW,

Bailey RL, Holland MJ. 2013. Plasmid copy number and disease severity in naturally

occurring ocular Chlamydia trachomatis infection. J Clin Microbiol 52:324.

25.  Okolie CE, Wooldridge KG, Turner DP, Cockayne A, James R. 2015. Development of a new

pentaplex real-time PCR assay for the identification of poly-microbial specimens containing

Staphylococcus aureus and other staphylococci, with simultaneous detection of

staphylococcal virulence and methicillin resistance markers. Mol Cell Probes 29:144–50.

26.  Nakagawa S, Taneike I, Mimura D, Iwakura N, Nakayama T, Emura T, Kitatsuji M, Fujimoto

A, Yamamoto T. 2005. Gene sequences and specific detection for Panton-Valentine

leukocidin. Biochem Biophys Res Commun 328:995–1002.

27.  Carvalho M da GS, Tondella ML, McCaustland K, Weidlich L, McGee L, Mayer LW,

Steigerwalt A, Whaley M, Facklam RR, Fields B, Carlone G, Ades EW, Dagan R, Sampson

JS. 2007. Evaluation and improvement of real-time PCR assays targeting lytA, ply, and psaA

genes for detection of pneumococcal DNA. J Clin Microbiol 45:2460–6.

28.  Meyler KL, Meehan M, Bennett D, Cunney R, Cafferkey M. 2012. Development of a

diagnostic real-time polymerase chain reaction assay for the detection of invasive

Haemophilus influenzae in clinical samples. Diagn Microbiol Infect Dis 74:356–62.

29.  Heim A, Ebnet C, Harste G, Pring-Akerblom P. 2003. Rapid and quantitative detection of

human adenovirus DNA by real-time PCR. J Med Virol 70:228–39.

30.  Kais M, Spindler C, Kalin M, Ortqvist A, Giske CG. 2006. Quantitative detection of

Streptococcus pneumoniae, Haemophilus influenzae, and Moraxella catarrhalis in lower

respiratory tract samples by real-time PCR. Diagn Microbiol Infect Dis 55:169–78.

31.  Greiner O, Day PJR, Altwegg M, Nadal D. 2003. Quantitative detection of Moraxella

catarrhalis in nasopharyngeal secretions by real-time PCR. J Clin Microbiol 41:1386–90.

32.  Abusleme L, Hong B-Y, Dupuy AK, Strausbaugh LD, Diaz PI. 2014. Influence of DNA

extraction on oral microbial profiles obtained via 16S rRNA gene sequencing. J Oral Microbiol

6.

33.  R Core Team. 2014. R: A Language and Environment for Statistical Computing. R Found

Stat Comput.

34.  Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naive Bayesian Classifier for Rapid

Assignment of rRNA Sequences into the New Bacterial Taxonomy. Appl Environ Microbiol

73:5261–5267.

35.  Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, Turner P, Parkhill J,

Loman NJ, Walker AW, Kunin V, Engelbrektson A, Ochman H, Hugenholtz P, Fv W, Göbel U,

Stackebrandt E, Kulakov L, McAlister M, Ogden K, Larkin M, O'Hanlon J, McAlister M,

Kulakov L, O'Hanlon J, Larkin M, Ogden K, Kéki Z, Grébner K, Bohus V, Márialigeti K, Tóth

E, Bohus V, Kéki Z, Márialigeti K, Baranyi K, Patek G, Schunk J, Tóth E, McFeters G,

Broadaway S, Pyle B, Egozy Y, Nogami T, Ohto T, Kawaguchi O, Zaitsu Y, Sasaki S, Shen

H, Rogelj S, Kieft T, Rand K, Houck H, Maiwald M, Ditton H, Sonntag H, Doeberitz M von K,

Tanner M, Goebel B, Dojka M, Pace N, Corless C, Guiver M, Borrow R, Edwards-Jones V,

Kaczmarski E, Fox A, Grahn N, Olofsson M, Ellnebo-Svedlund K, Monstein H, Jonasson J,

Newsome T, Li B, Zou N, Lo S, Mohammadi T, Reesink H, Vandenbroucke-Grauls C,

Savelkoul P, Barton H, Taylor N, Lubbers B, Pemberton A, Laurence M, Hatzis C, Brash D,

Oberauner L, Zachow C, Lackner S, Högenauer C, Smolle K, Berg G, Duc M La, Kern R,

Venkateswaran K, Ling Z, Liu X, Luo Y, Yuan L, Nelson K, Wang Y, Xiang C, Li L, Benítez-

Páez A, Álvarez M, Belda-Ferre P, Rubido S, Mira A, Tom I, Amar J, Serino M, Lange C,

Chabo C, Iacovoni J, Mondot S, Lepage P, Klopp C, Mariette J, Bouchez O, Perez L,

Courtney M, Marre M, Klopp P, Lantieri O, Doré J, Charles M, Balkau B, Burcelin R, Branton

W, Ellestad K, Maingat F, Wheatley B, Rud E, Warren R, Holt R, Surette M, Power C,

Borewicz K, Pragman A, Kim H, Hertz M, Wendt C, Isaacson R, Dong Q, Brulc J, Iovieno A,

Bates B, Garoutte A, Miller D, Revanna K, Gao X, Antonopoulos D, Slepak V, Shestopalov V,

Xuan C, Shamonki J, Chung A, DiNome M, Chung M, Sieling P, Lee D, Kuehn J, Gorden P,

Munro D, Rong R, Dong Q, Plummer P, Wang C, Phillips G, Srinivas G, Möller S, Wang J,

Künzel S, Zillikens D, Baines J, Ibrahim S, Boissière A, Tchioffo M, Bachar D, Abate L, Marie

A, Nsango S, Shahbazkia H, Awono-Ambene P, Levashina E, Christen R, Morlais I,

McKenzie V, Bowers R, Fierer N, Knight R, Lauber C, Carlos C, Torres T, Ottoboni L, Cheng

X, Tian X, Wang Y, Lin R, Mao Z, Chen N, Xie B, Davidson S, Powell R, James S, Knowlton

C, Veerapaneni R, D'Elia T, Rogers S, Shtarkman Y, Koçer Z, Edgar R, Veerapaneni R,

D'Elia T, Morris P, Rogers S, DeLeon-Rodriguez N, Lathem T, Rodriguez-R L, Barazesh J,

Anderson B, Beyersdorf A, Ziemba L, Bergin M, Nenes A, Konstantinidis K, Turner P, Turner

C, Jankhot A, Helen N, Lee S, Day N, White N, Nosten F, Goldblatt D, Willerslev E, Hansen

A, Poinar H, Kearney M, Spindler J, Wiegand A, Shao W, Anderson E, Maldarelli F, Ruscetti

F, Mellors J, Hughes S, Grice S Le, Coffin J, Cooper A, Poinar H, Roberts C, Ingham S,

Segal L, Alekseyenko A, Clemente J, Kulkarni R, Wu B, Chen H, Berger K, Goldring R, Rom

W, Blaser M, Weiden M, Bhatt A, Freeman S, Herrera A, Pedamallu C, Gevers D, Duke F,

Jung J, Michaud M, Walker B, Young S, Earl A, Kostic A, Ojesina A, Hasserjian R, Ballen K,

Chen Y, Hobbs G, Antin J, Soiffer R, Baden L, Garrett W, Hornick J, Marty F, Meyerson M,

Deragon J, Sinnett D, Mitchell G, Potier M, Labuda D, Sarkar G, Sommer S, Klaschik S,

Lehmann L, Raadts A, Hoeft A, Stuber F, Tamariz J, Voynarovska K, Prinz M, Caragine T,

Hilali F, Saulnier P, Chachaty E, Andremont A, Heininger A, Binder M, Ellinger A, Botzenhart

K, Unertl K, Döring G, Silkie S, Tolcher M, Nelson K, Carroll N, Adamson P, Okhravi N,

Mohammadi T, Reesink H, Vandenbroucke-Grauls C, Savelkoul P, Hughes M, Beck L, Skuce

R, Vaishampayan P, Probst A, Duc M La, Bargoma E, Benardini J, Andersen G,

Venkateswaran K, Rueckert A, Morgan H, Patel P, Garson J, Tettmar K, Ancliff S, McDonald

C, Pitt T, Coelho J, Tedder R, Champlot S, Berthelot C, Pruvost M, Bennett E, Grange T,

Geigl E, Chang S, Hsu H, Cheng J, Tseng C, Bonfert T, Csaba G, Zimmer R, Friedel C,

Knights D, Kuczynski J, Charlson E, Zaneveld J, Mozer M, Collman R, Bushman F, Knight R,

Kelley S, Eren A, Maignen L, Sul W, Murphy L, Grim S, Morrison H, Sogin M, Morris A, Beck

J, Schloss P, Campbell T, Crothers K, Curtis J, Flores S, Fontenot A, Ghedin E, Huang L,

Jablonski K, Kleerup E, Lynch S, Sodergren E, Twigg H, Young V, Bassis C, Venkataraman

A, Schmidt T, Weinstock G, Lane D, Klindworth A, Pruesse E, Schweer T, Peplies J, Quast

C, Horn M, Gl°Ckner F, Cooper P, Walker A, Reyes J, Chico M, Salter S, Vaca M, Parkhill J,

Schloss P, Westcott S, Ryabin T, Hall J, Hartmann M, Hollister E, Lesniewski R, Oakley B,

Parks D, Robinson C, Sahl J, Stres B, Thallinger G, Horn D Van, Weber C, Kozich J,

Westcott S, Baxter N, Highlander S, Schloss P, Quince C, Lanzen A, Davenport R,

Turnbaugh P, Bolger A, Lohse M, Usadel B, Kiełbasa S, Wan R, Sato K, Horton P, Frith M,

Huson D, Mitra S, Ruscheweyh H, Weber N, Schuster S, Schloss P, Gevers D, Westcott S,

White J, Nagarajan N, Pop M. 2014. Reagent and laboratory contamination can critically

impact sequence-based microbiome analyses. BMC Biol 12:87.

36.   Jombart T, Devillard S, Balloux F, Falush D, Stephens M, Pritchard J, Pritchard J, Stephens

M, Donnelly P, Corander J, Waldmann P, Sillanpaa M, Tang J, Hanage W, Fraser C,

Corander J, Lee C, Abdool A, Huang C, Jombart T, Jombart T, Devillard S, Dufour A, Pontier

D, Jombart T, Pontier D, Dufour A, McVean G, Novembre J, Stephens M, Patterson N, Price

A, Reich D, Price A, Patterson N, Plenge R, Weinblatt M, Shadick N, Reich D, Hotelling H,

Hotelling H, Pearson K, Liu N, Zhao H, Fisher R, Lachenbruch P, Goldstein M, Aitchison J,

Reyment R, Beharav A, Nevo E, Fraley C, Raftery A, Cann H, Toma C de, Cazes L, Legrand

M, Morel V, Piouffre L, Bodmer J, Bodmer W, Bonne-Tamir B, Cambon-Thomsen A,

Ramachandran S, Deshpande O, Roseman C, Rosenberg N, Feldman M, Cavalli-Sforza L,

Rosenberg N, Pritchard J, Weber J, Cann H, Kidd K, Zhivotovsky L, Feldman M, Wang S,

Lewis C, Jakobsson M, Ramachandran S, Ray N, Bedoya G, Rojas W, Parra M, Molina J,

Gallo C, Balloux F, Rosenberg N, Mahajan S, Ramachandran S, Zhao C, Pritchard J,

Feldman M, Rambaut A, Pybus O, Nelson M, Viboud C, Taubenberger J, Holmes E, Russell

C, Jones T, Barr I, Cox N, Garten R, Gregory V, Gust I, Hampson A, Hay A, Hurt A, Smith D,

Lapedes A, Jong J de, Bestebroer T, Rimmelzwaan G, Osterhaus A, Fouchier R, Holmes E,

Ghedin E, Miller N, Taylor J, Bao Y, George KS, Grenfell B, Salzberg S, Fraser C, Lipman D,

Young J, Palese P, Benson D, Karsch-Mizrachi A, Lipman D, Ostell J, Wheeler D, Larkin M,

Blackshields G, Brown N, Chenna R, McGettigan P, McWilliam H, Valentin F, Wallace I, Wilm

A, Lopez R, Waterhouse A, Procter J, Martin D, Clamp M, Barton G, Paradis E, Claude J,

Strimmer K, Handley L, Manica A, Goudet J, Balloux F, Serre D, Paabo S, Corander J,

Marttinen P, Siren J, Tang J, Francois O, Ancelet S, Guillot G, Hunley K, Healy M, Long J,

Kittles R, Weiss K, Manica A, Prugnolle F, Balloux F, Prugnolle F, Manica A, Balloux F,

Romero I, Manica A, Handley L, Balloux F, Amos W, Hoffman J, Fraley C, Raftery A, Peres-

Neto P, Jackson D, Somers K, Saporta G, Paradis E, Dray S, Dufour A, Schwarz G, Evanno

G, Regnaut S, Goudet J, Jakobsson M, Rosenberg N, Chessel D, Dufour A, Thioulouse J,

Dray S, Dufour A, Chessel D, Venables W, Ripley B, Nei M. 2010. Discriminant analysis of

principal components: a new method for the analysis of genetically structured populations.

BMC Genet 11:94.

37.    Capriotti JA, Pelletier JS, Shah M, Caivano DM, Ritterband DC. 2009. Normal ocular flora in

healthy eyes from a rural population in Sierra Leone. Int Ophthalmol 29:81–4.

38.    Walker AW, Martin JC, Scott P, Parkhill J, Flint HJ, Scott KP. 2015. 16S rRNA gene-based

profiling of the human infant gut microbiota is strongly influenced by sample processing and

PCR primer choice. Microbiome 3:26.

39.    Ramadhani AM, Derrick T, Macleod D, Holland MJ, Burton MJ. 2016. The Relationship

between Active Trachoma and Ocular Chlamydia trachomatis Infection before and after Mass

Antibiotic Treatment. PLoS Negl Trop Dis 10:e0005080.

40.    Solomon AW, Holland MJ, Alexander NDE, Massae PA, Aguirre A, Natividad-Sancho A,

Molina S, Safari S, Shao JF, Courtright P, Peeling RW, West SK, Bailey RL, Foster A, Mabey

DCW. 2004. Mass treatment with single-dose azithromycin for trachoma. N Engl J Med

351:1962–71.

41.    Seidman JC, Coles CL, Silbergeld EK, Levens J, Mkocha H, Johnson LB, Muñoz B, West

SK. 2014. Increased carriage of macrolide-resistant fecal E. coli following mass distribution of

azithromycin for trachoma control. Int J Epidemiol 43:1105–13.

42.    Brunham RC, Rekart ML. 2008. The Arrested Immunity Hypothesis and the Epidemiology of

Chlamydia Control. Sex Transm Dis 35:53–54.

*593*

594 ## Supplementary information

| Assay | Standard curve * | | |
|---|---|---|---|
| | LoD | $R^2$ | CoV (%) |
| *S. aureus* | 4.0 | 0.991 | 14.4 |
| Coagulase-negative *Staphylococcus* | 1.2 | 0.994 | 14.7 |
| *S. pneumoniae* | 9.9 | 0.998 | 11.9 |
| *H. influenza* | 2.3 | 0.997 | 14.7 |
| *Adenoviridae* | 3.0 | 0.997 | 9.1 |
| *M. catarrhalis* | 1.0 | 0.993 | 7.8 |

CoV: Coefficient of variation; LoD: Limit of detection: $R^2$: Coefficient of determination

* Values calculated from 10-fold dilution series of PCR product between approximately $10^5$ and $10^1$ copies per target per test with 5

technical replicates at each dilution point.

**Supplementary table 1:** Each PCR product for dilution into standards was prepared using TaqMan Universal II PCR mix

(Life Technologies, Paisley, UK), cleaned with Qiagen MinElute PCR product kit (Qiagen, Manchester, UK) and serially

diluted from $1:10^6$ to $1:10^{12}$ in ten-fold dilution steps. Each series was tested in five technical replicates with duplex ddPCR

assays to determine the reproducibility of the assay. Target concentrations between one and 10 copies/µL were

reproducibly detected by all six assays. The coefficient of determination for all assays was in excess of 0.99 when fitted to

a linear regression model. All six assays were highly reproducible, and the mean coefficient of variance (CoV) was 12.1%

(range: 7.8 – 14.7%).

*609*

*610*

*611* **Supplementary figure 2.** Relative taxa abundance in age-, sex- and location-matched children with TF/TI (n = 54) and

*612* without (n = 53). Relative taxa abundance is expressed as percentage of total reads per specimen group. Genera with

*613* reads representing less than 1% of the total number of reads were combined into a group entitled 'Other'.

*614*

615 **A**

616 **B**

618 **Supplementary figure 3.** First and second principal components coloured by **(A)** age group and **(B)** gender. Red spots

619 indicate individuals. Red arrows indicate loadings.

621    **Supplementary figure 4.** Relative contributions of genera driving difference between active trachoma cases and controls