A specific non-bisphosphonate inhibitor of the bifunctional farnesyl/geranylgeranyl diphosphate synthase in malaria parasites

Jolyn E. Gisselberg[1], Zachary Herrera[1], Lindsey Orchard[4], Manuel Llinás[4,5,6], and Ellen Yeh[1,2,3*]

[1]Department of Biochemistry, [2]Pathology, and [3]Microbiology and Immunology, Stanford Medical School, Stanford University, Stanford, CA 94305, USA

[4]Department of Biochemistry & Molecular Biology, [5]Department of Chemistry and [6]Huck Center for Malaria Research, Pennsylvania State University, University Park, PA 16802

*Corresponding author and lead contact: ellenyeh@stanford.edu

Summary

Isoprenoid biosynthesis is essential for *Plasmodium falciparum* (malaria) parasites and contains multiple validated antimalarial drug targets, including a bifunctional farnesyl and geranylgeranyl diphosphate synthase (FPPS/GGPPS). We identified MMV019313 as an inhibitor of *Pf*FPPS/GGPPS. Though *Pf*FPPS/GGPPS is also inhibited by a class of bisphosphonate drugs, MMV019313 has significant advantages for antimalarial drug development. MMV019313 has superior physicochemical properties compared to charged bisphosphonates that have poor bioavailability and strong bone affinity. We also show that it is highly selective for *Pf*FPPS/GGPPS and showed no activity against human FPPS or GGPPS. Inhibition of *Pf*FPPS/GGPPS by MMV019313, but not bisphosphonates, was disrupted in an S228T variant, demonstrating that MMV019313 and bisphosphonates have distinct modes-of-inhibition against *Pf*FPPS/GGPPS. Altogether MMV019313 is the first specific, non-bisphosphonate inhibitor of *Pf*FPPS/GGPPS. Our findings uncover a new small molecule binding site in this important antimalarial drug target and provide a promising starting point for development of *Plasmodium*-specific FPPS/GGPPS inhibitors.

Keywords: malaria; drug discovery; and high-throughput screening; non-bisphosphonate inhibitor

Introduction

There is an urgent need for antimalarials with novel mechanisms-of-action to circumvent resistance to frontline drugs. The biosynthesis of cellular isoprenoids is an essential process in *Plasmodium* parasites that cause malaria. A number of antimalarial compounds target enzymes

in isoprenoid biosynthetic pathways leading to parasite growth inhibition. First, *Plasmodium*

parasites depend on the 7-enzyme prokaryotic 2-*C*-methyl-D-erythritol 4-phosphate (MEP)

pathway in its plastid organelle, the apicoplast, to produce isopentenyl pyrophosphate (IPP) and

its isomer dimethyallyl pyrophosphate (DMAPP) (Jomaa et al., 1999). IPP and DMAPP are the

C5 building blocks for all isoprenoids. The antibiotic fosmidomycin inhibits the MEP enzyme,

Dxr/IspC, in both bacteria and *Plasmodium* parasites (Jomaa et al., 1999).

Second, at least three isoprenoid synthases (PF3D7_1128400.1, PF3D7_0202700,

PF3D7_0826400) catalyze the condensation of IPP and DMAPP into longer prenyl chains (Artz

et al., 2011; Jordão et al., 2013; Tonhosolo et al., 2005). In particular, farnesyl pyrophosphate

synthase (FPPS) and geranylgeranyl pyrophosphate synthase (GGPPS) are key branch point

enzymes that synthesize C15 and C20 prenyl chains, respectively, for multiple downstream

enzymes. In *Plasmodium* parasites, these reactions are catalyzed by a single bifunctional

enzyme, the farnesyl/geranylgeranyl diphosphate synthase (Artz et al., 2011; Jordão et al., 2013).

Nitrogen-containing bisphosphonates, blockbuster drugs which inhibit human FPPS, also inhibit

the bifunctional *Plasmodium* FPPS/GGPPS (Ghosh et al., 2004; Jordão et al., 2011; Michael B

Martin et al., 2001; No et al., 2012; Singh et al., 2010).

Finally, prenyl chains are cyclized and/or conjugated to small molecule and protein

scaffolds by a variety of prenyltransferases to biosynthesize final isoprenoid products required

for parasite growth and replication. Tetrahydroquinolines (THQ) have been shown to potently

inhibit the *Plasmodium* protein farnesyltransferase (Eastman et al., 2005; 2007; Nallan et al.,

2005). Other inhibitors may interfere with isoprenoid biosynthesis indirectly by disrupting

transporters that supply starting substrates or export products or blocking pathways that provide

cofactors for isoprenoid biosynthetic enzymes.

Importantly fosmidomycin, bisphosphonates, and tetrahydroquinolines have all shown

efficacy in mouse models of malaria infection, validating the key importance of isoprenoid

biosynthesis as an antimalarial drug target (Jomaa et al., 1999; Nallan et al., 2005; No et al.,

2012; Singh et al., 2010). Fosmidomycin is currently being tested in human clinical trials, while

a THQ lead candidate was investigated in preclinical studies (Fernandes et al., 2015; Nallan et

al., 2005). However, novel chemical scaffolds that disrupt isoprenoid biosynthetic pathways in

Plasmodium remain desirable to overcome unfavorable drug properties of these known

inhibitors. For example, bisphosphonates avidly bind bone mineral, and both fosmidomycin and

THQs have short half-lives *in vivo* (Sinigaglia et al., 2007; Tsuchiya et al., 1982; Van Voorhis et

al., 2007).

In 2011 the Medicine for Malaria Venture (MMV) distributed the Open-Access Malaria

Box to accelerate antimalarial drug discovery (Spangenberg et al., 2013). The Malaria Box

consists of 400 structurally diverse compounds, curated from >20,000 hits generated from large-

scale screens, that inhibit the growth of blood-stage *Plasmodium falciparum* parasites (Gamo et

al., 2010; Guiguemde et al., 2010; Rottmann et al., 2010). A major goal of sharing these

compounds was to facilitate elucidation of their antimalarial mechanism-of-action and open new

classes of validated chemical scaffolds and drug targets. Compounds that disrupt isoprenoid

metabolism can be detected by "rescue" of their growth inhibition upon supplementation of

isoprenoids in the growth media (Yeh and Derisi, 2011). Previously, we and two other groups

screened the Malaria Box for compounds whose growth inhibition were rescued by addition of

IPP and identified MMV008138 (Bowman et al., 2014; Wu et al., 2015). We and our

collaborators demonstrated that MMV008138 inhibits IspD, an enzyme in the MEP pathway that

produces IPP (Wu et al., 2015).

Using a quantitative high-throughput screen (qHTS), we report a second compound in the

Malaria Box, MMV019313, that shows an IPP rescue phenotype but was not identified in

screens performed by other groups (Bowman et al., 2014; DeRisi, 2014; Van Voorhis et al.,

2016). We demonstrate that the target of MMV019313 is the *P. falciparum* FPPS/GGPPS, a

cytosolic isoprenoid synthase that utilizes IPP and the key branch point enzyme in isoprenoid

biosynthesis in parasites. MMV019313 represents the first new class of non-bisphosphonate

inhibitors of *Pf*FPPS/GGPPS.

Results

A quantitative high-throughput screen (qHTS) for growth and IPP rescue identifies

MMV019313 as an inhibitor of isoprenoid biosynthesis

Previous IPP rescue screens of the Malaria Box tested compounds at a single, high

concentration >5 μM (Bowman et al., 2014; DeRisi, 2014; Van Voorhis et al., 2016). While

testing compounds at a single concentration is useful for identifying phenotypes that occur at a

threshold concentration (e.g. growth inhibition), growth rescue is expected to occur within a

specific range of concentrations. For example, doxycycline inhibits *P. falciparum* growth with an

EC_{50}= 0.3 μM that increases to 3.2 μM upon addition of IPP (Yeh and Derisi, 2011). At

concentrations <0.3 μM, doxycycline does not cause growth inhibition. However, at

concentrations >3.2 μM, it is no longer specific for its target and causes growth inhibition

through additional targets that cannot be IPP rescued. Therefore the concentration range in which

IPP rescue can be observed is greater than the EC_{50} of the compound for its specific, IPP-

rescuable target but less than that for any nonspecific targets.

To increase the sensitivity for detecting IPP chemical rescue, we screened the Malaria

Box for growth inhibition of blood-stage *P. falciparum* in the presence and absence of IPP over

8-12 drug concentrations from 0.01-27 μM (Table S1). Of 397 compounds tested (3 compounds

were not available), 383 showed growth inhibition at ≤27 μM. Initial hits showing IPP rescue of

growth inhibition at one or more drug concentrations were commercially sourced and retested.

Along with the previously reported compound, MMV008138, we confirmed a second compound

in the Malaria Box showing an IPP rescue phenotype, MMV019313 (Figure 1A). MMV019313

inhibited *P. falciparum* growth measured in a single replication cycle with EC_{50}=268 nM (250-

289 nM) in the absence of IPP; in the presence of IPP (200 μM), the EC_{50} was over 13-fold more

at 3.6 μM (3.2-4.0 μM) (Figure 1B). Notably, at concentrations >3.6 μM, MMV019313 inhibits a

nonspecific target that can no longer be IPP rescued, which explains why it was not identified in

previous screens.

An unusual feature of blood-stage *Plasmodium* is that IPP can rescue complete loss of the

apicoplast, the plastid organelle which houses the MEP pathway, since production of IPP is the

only essential function of the apicoplast. Compounds like doxycycline that disrupt apicoplast

biogenesis cause growth inhibition rescued by IPP and result in parasites lacking an apicoplast

(Yeh and Derisi, 2011). In contrast, inhibitors of isoprenoid biosynthesis, like fosmidomycin and

MMV008138, cause growth inhibition rescued by IPP with an intact apicoplast (Amberg-

Johnson et al., 2017). We determined whether MMV019313 disrupted the biogenesis of the

apicoplast. Both the replication of the apicoplast genome and import of an apicoplast-targeted

GFP were intact in MMV019313-treated and IPP-rescued parasites (Figure S1) (Yeh and Derisi,

2011). Altogether MMV019313 causes growth inhibition rescued by IPP with no defect in

apicoplast biogenesis, suggesting that, like fosmidomycin and MMV008138, MMV019313

blocks isoprenoid biosynthesis (Wu et al., 2015).

MMV019313-resistant parasites contain a mutation in the farnesyl/geranylgeranyl

diphosphate synthase

Though metabolomic profiling of MMV019313 has been performed, isoprenoid

intermediates were not examined so a specific disruption of these pathways could not be inferred

(Allman et al., 2016; Creek et al., 2016). Instead to gain further insight into the mechanism-of-

action of MMV019313, we identified mutations that confer resistance to MMV019313. Our

initial attempt to select drug-resistant parasites from a bulk culture of 10^{10} parasites was not

successful, though this same protocol was effective in selecting drug resistance against

MMV008138 when performed in parallel (Wu et al., 2015). Therefore, we turned to chemical

mutagenesis to increase the likelihood of selecting MMV019313-resistant parasites. Chemical

mutagenesis followed by whole genome sequencing to identify mutations has been successfully

employed to perform phenotypic screens in *T. gondii* but has not commonly been used in

Plasmodium parasites (Desai et al., 2005; Farrell et al., 2014; Inselburg, 1984; 1985). We treated

10^8 *P. falciparum* W2 parasites with sub-lethal doses of ethyl methanesulfonate (EMS), an

alkylating agent. Because EMS is unstable, we tested a fresh batch of mutagen immediately prior

to treatment in a standard parasite growth inhibition assay to determine its EC_{50}. We then

selected the EC_{50} (2.025 mM) as the highest EMS concentration used and tested several lower

concentrations by serially diluting 3-fold. These relatively low concentrations were selected to

maximize the frequency of drug-resistant mutations while minimizing lethal mutations that

would result in a smaller pool of starting parasites or the presence of nonspecific passenger

mutations that would confound identification of causative mutations during whole genome

analysis. Following mutagenesis, parasites were continuously selected with a dose of

MMV019313 equal to its EC_{75} in the presence of IPP for 32 days before removal of the drug.

This concentration was chosen to maximize selection pressure for developing resistance in the

IPP-rescuable target, while minimizing that for developing resistance in nonspecific targets.

Resistant parasites emerged in all mutagenized cultures one week after drug removal. In

these resistant populations, MMV019313 showed EC_{50} values that ranged from 3-9-fold greater

than that observed in the initial susceptible population. Growth inhibition of two MMV019313-

resistant populations which showed the highest levels of resistance at the lowest EMS

concentrations used (8.3 µM and 25 µM, designated 019313R1 and 019313R2 respectively) are

shown in Figure 2A. Significantly the EC_{50} of MMV019313 in 019313R1 and 019313R2 was

similar to that observed in susceptible strains in the presence of IPP, and growth inhibition could

no longer be rescued by addition of IPP (Figure S2). These results suggest that, as expected,

019313R1 and 019313R2 populations were completely resistant to inhibition of its specific IPP-

rescuable target but had not developed resistance to inhibition of additional nonspecific targets.

This is the first example of using chemical mutagenesis to aid selection of drug-resistant

parasites in *P. falciparum.*

019313R1, 019313R2, and their respective mutagenized parent strains used to initiate

drug selection, WT1 and WT2, were subjected to whole-genome sequencing. Comparison of the

resistant and parent genome sequences identified a single nucleotide variant (SNV) common to

both resistance populations, which was present in 100% of reads in 019313R1 and 63% of reads

in 019313R2 but not present in either susceptible parent genomes (Table 1). No other SNV were

detected at >40% prevalence in either resistant populations relative to the corresponding parent

populations (Table S2). The identified SNV was a T-to-A mutation in the gene PF3D7_1128400

characterized as a bifunctional FPPS/GGPPS and resulted in a Ser228-to-Thr change in the

protein (Figure 2B) (Artz et al., 2011; Jordão et al., 2013). IPP is a known substrate of the

FPPS/GGPPS, and in the primary sequence Ser228 is adjacent to conserved KT residues and an

Asp-rich region required for catalysis in all homologous FPPS and GGPPS enzymes (Figure 2C).

*Pf*FPPS/GGPPS was a strong candidate for further validation as the molecular target of

MMV019313.

Overexpression and an S228T variant of *Pf*FPPS/GGPPS confers resistance to

MMV019313

We determined whether overexpression of wildtype *Pf*FPPS/GGPPS was sufficient to

confer resistance to MMV019313. A transgene encoding FPPS/GGPPS-GFP under the control of

either the ribosomal L2 protein (*RL2*; PF3D7_1132700) or calmodulin (*CaM*; PF3D7_1434200)

promoter was integrated into an engineered *att*B locus in Dd2[attb] parasites (Nkrumah et al.,

2006). Expression of the CaM promoter is 10 to 50-fold greater across the life cycle than that of

the RL2 promoter based on RNA-seq data, resulting in appreciably greater FPPS/GGPPS-GFP

protein (Figure S3A) (Otto et al., 2010). Therefore, we compared the effect of no, moderate

(RL2), and high (CaM) levels of FPPS/GGPPS-GFP overexpression on susceptibility to

MMV019313. As shown in Figure 3A and Table S3, overexpression of FPPS/GGPPS-GFP

results in a dose-dependent increase in the EC_{50} of MMV019313 with intermediate-level

resistance to MMV019313 observed in the RL2-FPPS/GGPPS-GFP strain and high-level

resistance observed in the CaM-FPPS/GGPPS-GFP strain. Notably growth inhibition by

MMV019313 in the CaM-FPPS/GGPPS-GFP strain could no longer be rescued by IPP,

indicating that complete resistance to inhibition of the specific IPP-rescuable target had been

achieved (Figure S3B).

We also determined whether the S228T variant of GGPPS confers MMV019313

resistance. Initially we attempted to introduce the S228T mutation into the *fpps/ggpps* gene in the

P. falciparum genome using CRISPR-Cas9 mutagenesis but were unable to recover mutant

parasites. As an alternative, we overexpressed the FPPS/GGPPS(S228T)-GFP variant using the

"moderate" RL2 promoter and compared its effect on MMV019313 susceptibility with that of

the wild-type RL2-FPPS/GGPPS-GFP. Overexpression of the FPPS/GGPPS(S228T)-GFP

variant, even at moderate levels, caused an 18-fold increase in the EC_{50} which did not increase

further with IPP rescue, indicating that complete resistance to inhibition of the specific, IPP-

rescuable target had been achieved (Figure 3B and S3C; Table S3). Because moderate

overexpression of the S228T variant caused greater MMV019313 resistance than moderate

overexpression of wildtype *Pf*FPPS/GGPPS, we conclude that the S228T variant is sufficient to

confer resistance independent of overexpression. Altogether these results clearly demonstrated

that the mechanism-of-action of MMV019313 is dependent on *Pf*FPPS/GGPPS.

MMV019313 specifically inhibits the enzymatic activity of *Pf*FPPS/GGPPS

To confirm that *Pf*FPPS/GGPPS is the molecular target of MMV019313, we directly

measured MMV019313 inhibition in enzymatic assays. Consistent with a previous report,

purified *Pf*FPPS/GGPPS catalyzed the production of both farnesyl(C15)-PP and

geranylgeranyl(C20)-PP (Jordão et al., 2013). To measure the inhibitory effect of MMV019313,

we determined its IC_{50} for FPP and GGPP production in two conditions. In the first "non-

saturating" condition, substrate concentrations equaled K_M values (Figure S4), the concentration

at which the rate of reaction is half-maximal. In the second "k_{cat}" condition, substrate

concentrations were saturating and the rate of reaction was maximal. Measuring inhibitor effects

in both "non-saturating" and "k_{cat}" conditions increases the sensitivity for detecting different

types of inhibitors (competitive, non-competitive, uncompetitive). We found the IC_{50} value was

330 nM for FPP production under "non-saturating" conditions, comparable to its EC_{50} value in

cellular growth inhibition assays (Figure 1B and S5). MMV019313 was identified by its IPP

rescue phenotype in cell growth assays wherein addition of exogenous substrate increases

enzymatic rates, indicating that physiological conditions are likely sub-saturating. Therefore its

IC_{50} value measured in sub-saturating condition is more relevant for comparison to its EC_{50} value

for growth inhibition. Under "k_{cat}" conditions, IC_{50} values were 2.0 µM for FPP production and

9.8 µM for GGPP production (Figure 4A-B).

Since inhibition was detectable in both conditions, assays of mutant and human enzymes

were performed at saturating conditions which gives higher signal-to-noise. Significantly

MMV019313 at concentrations up to 200 µM did not inhibit human FPPS or GGPPS activity,

indicating that it was at least 100-fold selective for *Pf*FPPS/GGPPS over human homologs

(Figure 4A-B; Table S3). In contrast, the most selective bisphosphonate identified by a previous

study, BPH-703, showed only a 2.6-fold selectivity for *Pf*FPPS/GGPPS versus human FPPS and

3.6-fold selectivity versus human GGPPS in enzymatic assays (Figure 4C-D; Table S3) (No et

al., 2012).

MMV019313 binds a novel site on *Pf*FPPS/GGPPS, distinct from that of bisphosphonates

The lack of inhibition of human FPPS and GGPPS by MMV019313 was intriguing since

*Pf*FPPS/GGPPS is structurally related to human FPPS and GGPPS and all are inhibited by

bisphosphonates (No et al., 2012). Co-crystal structures have shown that bisphosphonates,

zoledronate and its lipophilic analog BPH-703, bind in the allylic substrate (DMAPP, GPP, or

FPP) site in the *Plasmodium vivax* FPPS/GGPPS homolog (No et al., 2012). This binding is

similar to what is observed in bisphosphonate-mammalian FPPS complexes (Kavanagh et al.,

2006; Rondeau et al., 2006; Yokoyama et al., 2015; Zhang et al., 2013). To determine whether

MMV019313 and bisphosphonates share a common binding site, we compared the effect of the

S228T variant on inhibition by MMV019313 and the bisphosphonate BPH-703. MMV019313

and BPH-703 both inhibited wild-type *Pf*FPPS/GGPPS activity. However only MMV019313

inhibition was decreased by over 10-fold in the S228T variant (Figure 4A-D). This difference in

their *in vitro* enzyme inhibition was consistent with the effect on parasite growth inhibition, in

which moderate overexpression of the S228T variant conferred resistance to MMV019313 but

not BPH-703 (Figure S5). In addition, residue S228 is adjacent to the IPP binding site in the

bisphosphonate-*Pv*FPPS/GGPPS structures and distal from the bound bisphosphonate. These

results demonstrate that MMV019313 has a novel binding mode, distinct from the known allylic

substrate site occupied by bisphosphonates, one which may confer its specificity for *Plasmodium*

FPPS/GGPPS over human homologs.

Because our biochemical results indicated that MMV019313 binds a new site which is

unique to *Pf*FPPS/GGPPS, we performed molecular docking calculations to evaluate the

likelihood of MMV019313 binding at known small molecule binding sites in *Pf*FPPS/GGPPS.

Three small molecule binding sites have been identified in FPPS and GPPS enzymes: 1) the

allylic substrate binding site, which is also occupied by bisphosphonates, 2) the IPP binding site,

and 3) an allosteric site identified in *Hs*FPPS (Jahnke et al., 2010). Because both the allylic

substrate and IPP sites have been structurally characterized in *Pv*FPPS/GGPPS, we used the rigid

body docking program Glide to generate poses with MMV019313 bound to these sites and

estimate the binding free energy of each pose (Friesner et al., 2004; 2006; Halgren et al., 2004).

The binding affinity estimated for the MMV019313 poses were $>10^8$ weaker than

bisphosphonate at the allylic site and $>10^5$ weaker than IPP at its site (Table S4). Therefore, the

modeling suggests that MMV019313 is unlikely to bind either the allylic substrate or IPP site.

We also attempted to dock MMV019313 into other potential binding pockets in the

PvFPPS/GGPPS apo structure but with inconclusive results.

Discussion

Farnesyl and geranylgeranyl diphosphate synthase (FPPS and GGPPS) are key branch

point enzymes in isoprenoid biosynthesis. Human cells contain separate FPPS and GGPPS

enzymes. An important class of clinical drugs, nitrogen-containing bisphosphonates, inhibits

human FPPS in osteoclasts and block their function and proliferation (Kavanagh et al., 2006).

Because osteoclasts are responsible for bone resorption, bisphosphonates are highly effective for

treatment of osteoporosis and other bone remodeling diseases. Bisphosphonates are chemically

stable analogs of inorganic pyrophosphate containing a P-C-P bond in place of the

phosphodiester, which accounts for both its inhibition of FPPS (acting as an analog of the allylic

diphosphate substrate) and its high selectivity for osteoclasts (depositing in bone mineral which

is composed of calcium and phosphate). Unfortunately the charge state of bisphosphonates is a

major liability in other therapeutic applications, as they are poorly bioavailable, rapidly cleared

by the kidney, and do not achieve therapeutic levels in serum for treatment of non-bone diseases

(Cremers et al., 2005; Jordão et al., 2011; Singh et al., 2010; Van Voorhis et al., 2016).

*Pf*FPPS/GGPPS, the molecular target of MMV019313 as demonstrated in this study,

closely resembles mammalian FPPS enzymes in sequence, structure, and inhibition by

bisphosphonates (Jordão et al., 2013; No et al., 2012). Like human FPPS, it is a central node in

cellular isoprenoid biosynthesis vulnerable to drug inhibition (Artz et al., 2011; Kavanagh et al.,

2006; Luckman et al., 1998). In *Plasmodium*, FPP and GGPP are required for the biosynthesis of

prenylated proteins, the prenyl modification of ubiquinone, and other isoprenoid products, such

that inhibition of *Pf*FPPS/GGPPS disrupts multiple cellular pathways (Chakrabarti et al., 2002;

de Macedo et al., 2002; Gabriel et al., 2015; Tonhosolo et al., 2005; 2009). Previously lipophilic

bisphosphonates modified with an alkyl chain to increase their cell permeability were shown to

inhibit *Pv*FPPS/GGPPS homolog and clear both blood- and liver-stage *Plasmodium* parasites in

mice infection models (No et al., 2012; Singh et al., 2010). Importantly, these results validated

Plasmodium FPPS/GGPPS as an antiparasitic drug target for both acute malaria treatment and

malaria chemoprophylaxis.

Our identification of MMV019313 further addresses two key hurdles in the development

of *Pf*FPPS/GGPPS inhibitors as antimalarial drugs. First, MMV019313 represents the first non-

bisphosphonate class of *Plasmodium* FPPS/GGPPS inhibitors with superior physicochemical

properties. Many efforts have been made to develop modified bisphosphonates or non-

bisphosphonate compounds as FPPS and/or GGPPS inhibitors for treatment of soft-tissue

cancers and infectious diseases (Chen et al., 2013; Jahnke et al., 2010; Liu et al., 2014;

Marzinzik et al., 2015; Zhang et al., 2009). Unlike bisphosphonates, MMV019313 has drug-like

physicochemical properties satisfying the Rule of 5 and does not need to mimic a charged

diphosphate substrate to achieve FPPS/GGPPS inhibition (Van Voorhis et al., 2016). As a

compound in the Malaria Box library, it has already been tested in a panel of bioactivity and

pharmacokinetic assays with encouraging results (Van Voorhis et al., 2016). Furthermore the

results of >300 assays characterizing Malaria Box compounds as part of an innovative "open

source" drug discovery effort by many groups will be a rich source of information (Allman et al.,

2016; Paul et al., 2016; Ullah et al., 2017; Van Voorhis et al., 2016).

Second, MMV019313 has high selectivity for *Pf*FPPS/GGPPS over human FPPS and

GGPPS, minimizing the potential for mechanism-based (e.g. "on-target") toxicity. In fact, we

could not detect any inhibition of human FPPS or GGPPS indicating selectivity at the enzymatic

level of at least 100-fold. Consistent with this lack of enzymatic inhibition, MMV019313

showed no cytotoxicity against a panel of 60 human cancer cell lines at 10 μM (Van Voorhis et al.,

2016). In contrast, the most selective bisphosphonate BPH-703 identified by No *et al* showed

2.6-3.6-fold selectivity in our enzymatic assays (corresponding to a reported therapeutic index of

193 in growth inhibition assays) (No et al., 2012). Our results suggest that MMV019313 binds a

novel site in *Pf*FPPS/GGPPS that is either absent from or substantially different in the human

homologs, which may explain its high selectivity. Altogether MMV019313 offers a distinct and

promising starting point for development of antimalarial FPPS/GGPPS inhibitors, which

circumvents the inherent liabilities of the bisphosphonate pharmacophore and greatly improves

on their selectivity.

Our immediate priority is to obtain structures of the inhibitor-enzyme complex to aid in

structure-based design of MMV019313 derivatives with higher potency, as well as discovery of

additional chemical scaffolds that occupy this new binding site. A potential challenge is that the

FPPS/GGPPS structure is dynamic during catalysis. For example, the human FPPS is known to

undergo at least two critical conformational changes (Kavanagh et al., 2006): The first is from an

"open" apo form to a "closed" conformation upon binding of the allylic substrate, which is

associated with drastically increased affinity for the allylic substrate (and bisphosphonates that

mimic this substrate) as well as orders the IPP binding pocket. The second structural change

occurs upon IPP binding and sequesters the active site from bulk solvent. Several attempts to

obtain co-crystal structures of MMV019313 with *Pv*FPPS/GGPPS under conditions in which

bisphosphonate-*Pv*FPPS/GGPPS crystals were obtained (presumed to correspond to the "closed"

conformation) have been unsuccessful (personal communication, Dr. Raymond Hui). Additional

crystallization conditions may require testing in the presence and absence of different allylic

substrates and IPP, as their binding could affect the binding of MMV019313. Moreover the

conformational dynamics of FPPS and GGPPS enzymes preclude straightforward interpretation

of the mode-of-inhibition from conventional kinetic assays. Bisphosphonates that bind at the

allylic substrate site show slow, tight binding with characteristics of irreversible inhibitors rather

than classic competitive inhibitors (Dunford et al., 2008; Taylor, 2004). Indeed our preliminary

assays indicate that MMV019313 also shows slow binding likely dependent on conformational

changes.

Since the S228T variant distinguishes between MMV019313 and bisphosphonate binding

to *Pf*FPPS/GGPPS, obtaining its structure to compare with wildtype FPPS/GGPPS will also offer

clues to this new mode-of-inhibition. The resistance caused by the S228T variant could be

explained by a direct contact between Ser228 and MMV019313 in a new small molecule binding

pocket. But because the change from Ser to Thr is quite conservative, the addition of a methyl

group, it seems more likely that Ser228 is involved in conformational dynamics important for

catalysis. Structural analysis of this variant enzyme may reveal altered conformational states

underlying the resistance to MMV019313.

Optimization of the bioactivity and pharmacokinetic properties of MMV019313

derivatives will be priorities for their development as antimalarials. The luciferase-based

enzymatic assays performed in our study are robust for detection of effective inhibitors and can

be adapted to high-throughput screening of small molecule libraries (in contrast we found a

commercially-available colorimetric assay was not sufficiently sensitive) (Crowther et al., 2011).

Inhibitor-enzyme structures will greatly accelerate optimization for potency against the

Plasmodium FPPS/GGPPS, while simultaneously minimizing binding to the human homologs.

P. falciparum strains overexpressing wildtype or the S228T variant, generated during this study,

can also be re-tooled as secondary cellular assays for on-target specificity. Increased potency

against blood-stage parasites will likely improve its activity against liver stage which currently is

not detectable at 5 μM (Van Voorhis et al., 2016). Inhibition of isoprenoid biosynthesis by

fosmidomycin, bisphosphonates, or MMV019313 has not been shown to block gametocyte

development (Lell et al., 2003; Van Voorhis et al., 2016). The drug properties of MMV019313

derivatives will be optimized in standard absorption-distribution-metabolism-excretion-toxicity

(ADME-T) assays. For example, we found that while MMV019313 is stable to human liver

microsomal enzymes ($t_{1/2} > 159$ min), its $t_{1/2}$ in mouse liver microsomes was 4 min. This metabolic

instability may account for a <1μM peak serum concentration following oral administration in

mice and will need to be addressed before derivatives can be tested in mouse models of

Plasmodium infection (Van Voorhis et al., 2016).

Finally, our study demonstrates two methodological improvements. First, we took

advantage of a quantitative screen for IPP chemical rescue using a range of inhibitor

concentrations, rather than a single high dose. The increased sensitivity allowed us to identify

MMV019313, though it was missed in prior screens. It also permits the screen to be performed

with geranylgeraniol, which is significantly less costly than IPP, since so far all identified

inhibitors that rescue with IPP also show at least a partial rescue of parasite growth inhibition

with geranylgeraniol (Yeh and Derisi, 2011; Zhang et al., 2011). Second, we successfully

identified a drug-resistant mutation in chemically mutagenized *Plasmodium* parasites to facilitate

mechanism-of-action elucidation. Though the rate and type of mutations caused by EMS was not

characterized in our experiments, in the related apicomplexan parasite, *Toxoplasma gondii* EMS

preferentially mutated A/T residues with similar rates of transitions and transversions (Farrell et

al., 2014). Notably our study identified a T-to-A transversion in *Pf*FPPS/GGPPS in

MMV019313-resistant parasites which were only selected after chemical mutagenesis.

Characterization of the rate and type of mutations in *Plasmodium* parasites will facilitate

selection and dosing of specific chemical mutagens to optimize phenotype selection and

mutation identification by sequencing. Though limited, our experience suggests that the use of

chemical mutagens will increase successful selection for drug resistance in *Plasmodium*

parasites, which is the most common method to identify drug targets from phenotypic drug

screens (Gamo et al., 2010; Guiguemde et al., 2010; Rottmann et al., 2010). The low frequency

of *in vitro* MMV019313 resistance also indicates that clinical resistance will also be less frequent

to MMV019313 and its derivatives.

Significance

There is an urgent need for antimalarials with novel mechanisms-of-action to circumvent

resistance to frontline drugs. Isoprenoid biosynthetic pathways are essential for *Plasmodium*

(malaria) parasites and contain multiple validated drug targets. Using a chemical genetic

approach from a phenotypic growth rescue screen to target identification, we identified a new

antimalarial compound that inhibits a key branchpoint enzyme in isoprenoid biosynthesis. Our

compound has significant advantages over bisphosphonate inhibitors that inhibit the same drug

target with improved drug properties and specificity for the *Plasmodium* enzyme over human

homologs. Our findings uncover a new "druggable" site in this important antimalarial drug target

and provide a chemical starting point for development of *Plasmodium*-specific FPPS/GGPPS

inhibitors.

Materials and Methods

***P. falciparum in vitro* cultures.** *Plasmodium falciparum* W2 (MRA-157) and Dd2^{attB}

(MRA-843) were obtained from MR4. Parasites were grown in human erythrocytes (2%

hematocrit, obtained from the Stanford Blood Center) in RPMI 1640 media supplemented with

0.25% Albumax II (GIBCO Life Technologies), 2 g/L sodium bicarbonate, 0.1 mM

hypoxanthine, 25 mM HEPES (pH 7.4), and 50 µg/L gentamycin, at 37°C, 5% O_2, and 5% CO_2.

For passage of drug-treated, IPP-rescued parasites, the media was supplemented with 5 µM drug

and 200 µM IPP (Isoprenoids LC or NuChem). For comparison of growth between different

treatment conditions, cultures were carried simultaneously and handled identically with respect

to media changes and addition of blood cells

Chemical handling. Malaria Box compounds were received as 10 mM DMSO stocks in

96-well plates and diluted three-fold manually in DMSO. Fosmidomycin was included in control

wells. Two-fold serial dilutions of the 96-well plates were performed on Velocity11. Compound

stocks stored in DMSO were diluted for growth assays. MMV019313 was purchased from

ChemDiv Biotech.

qHTS screen for IPP chemical rescue. Growth assays were performed in 384-well clear

bottom assay plates (E and K scientific, Santa Clara, CA). Drug (100-400 nL) was added directly

to each well using PinTool (V&P Scientific) on a Sciclone ALH3000 (Caliper Sciences). Using

the Titertek Multidrop 384, first 40 µL of growth media with and without 375 µM IPP was

dispensed, followed by 10 µL ring-stage *P. falciparum* D10 parasites (parasitemia 0.8% in 10%

hematocrit) into 384-well plates using the Titertek Multidrop 384. The final assay consisted of

50 µL ring-stage cultures at 0.8% parasitemia/ 2% hematocrit and drug concentrations from

0.01-26.7 µM ± 300 µM IPP. The plate was incubated at 37°C for 72 h. Parasites were lysed

with 10 µL 5mM EDTA, 1.6% Triton-X, 20mM Tris-HCl and 0.16% Saponin containing 0.1%

Sybr Green I (Invitrogen). The plates were then incubated at -80 °C for 20 min and thawed at

room temperature overnight in the dark. Fluorescence was detected using Flexstation II- 384.

Compounds that showed IPP rescue of growth inhibition at 1 or more drug concentrations in the

initial 384-well high-throughput screen were re-tested in a 96-well growth assay.

Growth inhibition assays to determine EC$_{50}$ values. *P. falciparum* cultures (125 µL)

were grown in 96-well plates containing serial dilution of drugs in triplicate. Media was

supplemented with 200 µM IPP as indicated. Growth was initiated with ring-stage parasites at

1% parasitemia and 0.5% hematocrit. Plates were incubated for 72h. Growth was terminated by

fixation with 1% formaldehyde and parasitized cells were stained with 50 nM YOYO-1

(Invitrogen). Parasitemia was determined by flow cytometry. Data were analyzed by BD C6

Accuri C-Sampler software and fitted to a sigmoidal dose-inhibition function

(Inhibition=Bottom + (Top-Bottom)/(1+10^((LogIC50-[drug])*HillSlope) by GraphPad Prism.

***P. falciparum* mutagenesis and resistance selection.** Late-stage parasites were purified

using a SuperMACS II separator (Miltenyi Biotec) and incubated in complete medium with 8.3 –

2025 μM ethyl methanesulfonate (EMS, 6 concentrations total) for 2 hours. The concentrations

of EMS used were selected by determining the EC_{50} in 72h parasite growth inhibition assays.

The highest concentration used for mutagenesis was equal to the EC_{50} in order to maximize the

selection pressure. The mutagen was then serially diluted 1:3 to test lower concentrations that

might give lower mutation rates and therefore a lower frequency of passenger mutations.

Mutagenized parasites were washed and separated into wells of 10 mL total volume

(approximately 10% parasitemia, 2 % HCT). MMV019313 drug selection was applied to one

well for each mutagenesis condition at 600 nM (approximately EC_{75}), while the other well was

left untreated in order to serve as a control for whole genome sequencing. Parasites were fed

daily for the first week and every 3 days thereafter. Each culture was split in half every 6 days in

order to introduce fresh RBC. Drug pressure was maintained for 32 days, with no observable

parasite growth. After 32 days of selection, half of the culture from each EMS condition was

removed from drug pressure. In these cultures, parasites which showed resistance to

MMV019313 by a standard drug assay were observable after 7 days at all EMS conditions used.

The parasites treated with the two lowest concentrations of EMS were selected for whole

genome sequencing.

Whole Genome Sequencing. *Plasmodium falciparum* strains were sequenced by

Illumina sequencing as described previously (Straimer et al., 2012). Briefly, NEBNext DNA

library reagents (NEB) and NEXTflex DNA barcode adapters (Bioo Scientific) were used to

prepare PCR-free libraries (Kozarewa et al., 2009). Eight whole genome gDNA libraries were

multiplexed and spiked with 8% PhiX control. Single-end sequencing was performed across two

lanes on an Illumina HiSeq 2500 system. Data was analyzed using tools available in the Galaxy

platform (Blankenberg et al., 2010; Giardine et al., 2005; Goecks et al., 2010). Sequencing reads

were mapped against the *P. falciparum* 3D7v.10.0 reference genome using the Burrows-Wheeler

Alignment tool (Li and Durbin, 2009). Sequencing data was visualized using Integrative

Genomics Viewer (IGV) (Robinson et al., 2011; Thorvaldsdóttir et al., 2013). Variants were

called using Freebayes (Garrison and Marth) and filtered for Quality >100 and Read Depth >30

using GATK tools (Van der Auwera et al., 2013). SnpEff was used to annotate the list of variants

based on the *P. falciparum* 3D7v9.1 reference genome (Cingolani et al., 2012). Sequencing data

have been deposited to the NCBI SRA (www.ncbi.nlm.nih.gov/sra; SRP106479).

***P. falciparum* Transfections.** An *E. coli* codon optimized version of *Pf*FPPS/GGPPS

(PF3D7_1128400) was designed and synthesized by GeneWiz. Quick change mutagenesis was

used to mutate GGPPS serine 228 to a threonine in the pUC vector provided by GeneWiz. These

constructs were then moved into the pLN transfection plasmid designed for Bxb1

mycobacteriophage integrase system (Adjalley et al., 2010). The In-fusion cloning kit (Clontech)

was used for all cloning. Restriction enzymes *Av*rII and *Bs*iWI were used to linearize the pLN

vector. GGPPS was designed to have a *C*-terminal GFP tag. All transgenes were driven with

either the ribosomal L2 protein (RL2) promoter (PF3D7_1132700) or the calmodulin (CaM)

promoter (PF3D7_1434200) as denoted.

Transfections were carried out as previously described (Spalding et al., 2010). Briefly,

400 µL fresh red blood cells were preloaded with 100 µg of both pINT, which carries the

bacteriophage integrase, and pRL2, which carries the gene of interest and the blasticidin

resistance cassette, using a BioRad Gene-Pulser Xcell Electroporator. Electroporation conditions

were infinite resistance, 310 V, and 950 μF using a 2 mm cuvette. Preloaded red blood cells were

combined with 2.5 mL ~20% schizont Dd2^{attB} parasites and allowed to recover for 2 days before

selection pressure was applied. Transfected parasites were selected with 2.5 μg/mL blasticidin

are were detectable by thin smear within 15 days. Integration was confirmed by PCR and identity

of the transgene was confirmed by sanger sequencing.

Immunoblots. Parasites expressing either ACP$_L$-GFP or one of the FPPS/GGPPS

constructs generated in this study were isolated by saponin lysis and resuspended in 1xNuPage

LDS sample buffer (Invitrogen). Whole cell lysate was separated by SDS-PAGE using 4-12%

Bis-Tris gels (Initrogen) and transferred to nitrocellulose using a Trans Turbo-blot (BioRad).

Membranes were blocked with 3% BSA, probed with 1:5,000 monoclonal anti-GFP JL-8

(BioRad) overnight, washed, then probed with 1:10,000 IRDye 680LT goat-anti-mouse. The

Western was imaged using a Oddysey Imager (LiCor Biosciences).

Live microscopy. Infected red blood cells were treated with Hoescht to stain the nucleus.

Single z-stack images were collected on an epifluorescence Nikon eclipse.

Quantative PCR. Quantitative PCR was performed as previously published (Yeh and

Derisi, 2011). Briefly, parasites from 200 μL of culture were isolated by saponin lysis followed

by PBS wash to remove extracellular DNA. DNA was purified using DNeasy Blood and Tissue

kit (Qiagen). Primers were designed to target genes found on the apicoplast or nuclear

genome: *tufA* (apicoplast) 5′-GATATTGATTCAGCTCCAGAAGAAA-3′ / 5′-

ATATCCATTTGTGTGGCTCCTATAA-3′ and *CHT1* (nuclear) 5′-

TGTTTCCTTCAACCCCTTTT-3′ / 5′-TGTTTCCTTCAACCCCTTTT-3′. Reactions contained

template DNA, 0.15 μM of each primer, and 0.75× LightCycler 480 SYBR Green I Master mix

(Roche). PCR reactions were performed at 56°C primer annealing and 65°C template extension

for 35 cycles on a Lightcycler 6500 (Roche). For each time point, the apicoplast：nuclear genome

ratio of the fosmidomycin-treated positive control, chloramphenicol-treated negative control, or

MMV019313-treated experiment were calculated relative to that of an untreated control

collected at the same time.

Recombinant protein purification. Full length constructs of *Pf*FPPS/GGPPS, *Hs*FPPS,

and *Hs*GGPPS were cloned into pET28a with an n-terminal hexahistidine tag. The *P. falciparum*

FPPS/GGPPS was codon optimized for expression in *E. coli* (GeneWiz). *Pf*FPPS/GGPPS was

mutagenized (S228T) using quick change mutagenesis. When expressed in *E. coli,*

*Pf*FPPS/GGPPS (wt and S228T) and *Hs*FPPS were toxic. Cultures were supplemented with 0.4%

glucose and grown to OD_{600} of 0.8-1 and induced with 0.5 mM IPTG. *Hs*GGPPS was grown

without supplementation to an OD_{600} of 0.8-1. All cultures were induced for 4 hours at 37 °C,

then harvested. Cells were lysed in 20 mM HEPES pH 8.0, 150 mM NaCl, 2 mM $MgCl_2$, and

protease inhibitor cocktail using sonication. Cleared lysates were either mixed with Talon metal

affinity resin (Clonetech) or purified over 5 ml HisTrap columns (GE Healthcare). His tagged

protein was purified with a single step purification eluting with buffer with 300 mM imidazole.

Proteins were dialyzed to remove imidazole and flash frozen.

In vitro **FPPS/GGPPS assays.** FPPS/GGPPS activity was measured by monitoring

pyrophosphate release using the Lonza PPiLight kit under kinetic conditions. Drug and either 20

µg/ml *Pf*FPPS/GGPPS, 40 µg/ml *Hs*GGPPS, or 100 µg/ml *Hs*FPPS protein were incubated for

30 min room temperature. The reaction was initiated by the addition of Lonza PPiLight kit

reaction mix and substrates. Saturating substrate conditions were 100 µM GPP or FPP, and 200

µM IPP. Sub-saturation conditions were K_M conditions as shown in Figure S4. Luciferase

activity was monitored over time using a BioTek plate reader. Reaction rates were calculated

from the linear portion of the raw luminescence over time curves (R^2 >0.9). Data were fitted to a

sigmoidal dose-inhibition function (Inhibition=Bottom + (Top-Bottom)/(1+10^((LogIC50-

[drug])*HillSlope) by GraphPad Prism software.

MMV019313 ligand docking. Using a solved *Pv*FPPS/GGPPS structure (PDB: 3EZ3,

zoledronate and IPP bound) as a receptor model protein preparation wizard was used to add back

in missing hydrogens and side chains. All crystallographic waters were removed. Hydrogen

bonds were calculated using Epik at pH 8 (±1). The protein structure was minimized using OPL3

force field. MMV019313 was docked using glide to receptor grids generated from the relevant

crystallized small molecules.

Author Contributions

JEG, ZH, LO, and EY conducted experiments. JEG, ZH, LO, ML, and EY were

responsible for data analysis. JEG and EY designed experiments and wrote the paper. ML and

EY supervised the project.

Acknowledgements

We are grateful to Medicines for Malaria Ventures (MMV) for providing the Malaria

Box compounds and making this valuable library freely available, as well as GlaxoSmithKline

for their screening efforts that first identified MMV019313 (TCMDC-123889). We would like to

thank Dr. Susmitha Suresh for performing drug screens and Dr. Felice Kelly for advice on how

to chemically mutagenize our parasites for resistance selection. We are grateful to Dr. James

Dunford (University of Oxford) for advice in developing the *in vitro* enzyme activity assays. We

also would like to acknowledge Dr. Wei Zhu and Professor Eric Oldfield (University of Illinois,

Urbana-Champaign) for *in vitro P. vivax* GGPPS activity assays and providing BPH-703.

Funding support for this project was generously provided by the Stanford Consortium for

Innovation, Design, Evaluation and Action (C-IDEA), NIH 1K08AI097239 (EY), NIH

1DP5OD012119 (EY), the Burroughs Wellcome Fund Career Award for Medical Scientists

(EY), the Burroughs Wellcome Fund Investigators in Pathogenesis of Infectious Disease (PATH)

Award (ML), and the Stanford School of Medicine Dean's Postdoctoral Fellowship (JEG). The

authors declare no conflict of interest.

References

586
Adjalley, S.H., Lee, M.C.S., and Fidock, D.A. (2010). A method for rapid genetic integration
into *Plasmodium falciparum* utilizing mycobacteriophage Bxb1 integrase. Methods Mol. Biol.
634, 87–100.

Allman, E.L., Painter, H.J., Samra, J., Carrasquilla, M., and Llinás, M. (2016). Metabolomic
Profiling of the Malaria Box Reveals Antimalarial Target Pathways. Antimicrob. Agents
Chemother. *60*, 6635–6649.

Amberg-Johnson, K., Ganesan, S.M., Lorenzi, H.A., Niles, J.C., and Yeh, E. (2017). A first-in-

class inhibitor of parasite FtsH disrupts plastid biogenesis in human pathogens. bioRxiv 108910.

Artz, J.D., Wernimont, A.K., Dunford, J.E., Schapira, M., Dong, A., Zhao, Y., Lew, J., Russell,
R.G.G., Ebetino, F.H., Oppermann, U., et al. (2011). Molecular characterization of a novel
geranylgeranyl pyrophosphate synthase from *Plasmodium* parasites. J. Biol. Chem. *286*, 3315–
3322.

Blankenberg, D., Kuster, Von, G., Coraor, N., Ananda, G., Lazarus, R., Mangan, M.,
Nekrutenko, A., and Taylor, J. (2010). Galaxy: a web-based genome analysis tool for
experimentalists. Curr Protoc Mol Biol *Chapter 19*, Unit19.10.1–Unit19.10.21.

Bowman, J.D., Merino, E.F., Brooks, C.F., Striepen, B., Carlier, P.R., and Cassera, M.B. (2014).
Antiapicoplast and gametocytocidal screening to identify the mechanisms of action of
compounds within the malaria box. Antimicrob. Agents Chemother. *58*, 811–819.

Chakrabarti, D., Da Silva, T., Barger, J., Paquette, S., Patel, H., Patterson, S., and Allen, C.M.
(2002). Protein farnesyltransferase and protein prenylation in *Plasmodium falciparum*. J. Biol.
Chem.

Chen, S.-H., Lin, S.-W., Lin, S.-R., Liang, P.-H., and Yang, J.-M. (2013). Moiety-linkage map
reveals selective nonbisphosphonate inhibitors of human geranylgeranyl diphosphate synthase. J
Chem Inf Model *53*, 2299–2311.

Cingolani, P., Patel, V.M., Coon, M., Nguyen, T., Land, S.J., Ruden, D.M., and Lu, X. (2012).
Using *Drosophila melanogaster* as a Model for Genotoxic Chemical Mutational Studies with a
New Program, SnpSift. Front Genet *3*, 35.

Creek, D.J., Chua, H.H., Cobbold, S.A., Nijagal, B., MacRae, J.I., Dickerman, B.K., Gilson,
P.R., Ralph, S.A., and McConville, M.J. (2016). Metabolomics-Based Screening of the Malaria
Box Reveals both Novel and Established Mechanisms of Action. Antimicrob. Agents
Chemother. *60*, 6650–6663.

Cremers, S.C.L.M., Pillai, G., and Papapoulos, S.E. (2005).
Pharmacokinetics/pharmacodynamics of bisphosphonates: use for optimisation of intermittent
therapy for osteoporosis. Clin Pharmacokinet *44*, 551–570.

Crowther, G.J., Napuli, A.J., Gilligan, J.H., Gagaring, K., Borboa, R., Francek, C., Chen, Z.,
Dagostino, E.F., Stockmyer, J.B., Wang, Y., et al. (2011). Identification of inhibitors for putative
malaria drug targets among novel antimalarial compounds. Mol. Biochem. Parasitol. *175*, 21–29.

de Macedo, C.S., Uhrig, M.L., Kimura, E.A., and Katzin, A.M. (2002). Characterization of the
isoprenoid chain of coenzyme Q in *Plasmodium falciparum*. FEMS Microbiol. Lett. *207*, 13–20.

DeRisi, J.L. (2014). UCSF DeRisi Lab MMV Box Apicoplast Screening (EMBL-EBI).

Desai, S.A., Alkhalil, A., Kang, M., Ashfaq, U., and Nguyen, M.-L. (2005). Plasmodial surface
anion channel-independent phloridzin resistance in *Plasmodium falciparum*. J. Biol. Chem. *280*,
16861–16867.

Dunford, J.E., Kwaasi, A.A., Rogers, M.J., Barnett, B.L., Ebetino, F.H., Russell, R.G.G.,
Oppermann, U., and Kavanagh, K.L. (2008). Structure-activity relationships among the nitrogen
containing bisphosphonates in clinical use and other analogues: time-dependent inhibition of
human farnesyl pyrophosphate synthase. J. Med. Chem. *51*, 2187–2195.

Eastman, R.T., White, J., Hucke, O., Bauer, K., Yokoyama, K., Nallan, L., Chakrabarti, D.,
Verlinde, C.L.M.J., Gelb, M.H., Rathod, P.K., et al. (2005). Resistance to a protein
farnesyltransferase inhibitor in *Plasmodium falciparum*. J. Biol. Chem. *280*, 13554–13559.

Eastman, R.T., White, J., Hucke, O., Yokoyama, K., Verlinde, C.L.M.J., Hast, M.A., Beese,
L.S., Gelb, M.H., Rathod, P.K., and Van Voorhis, W.C. (2007). Resistance mutations at the lipid
substrate binding site of *Plasmodium falciparum* protein farnesyltransferase. Mol. Biochem.
Parasitol. *152*, 66–71.

Farrell, A., Coleman, B.I., Benenati, B., Brown, K.M., Blader, I.J., Marth, G.T., and Gubbels,
M.-J. (2014). Whole genome profiling of spontaneous and chemically induced mutations in
Toxoplasma gondii. BMC Genomics *15*, 354.

Fernandes, J.F., Lell, B., Agnandji, S.T., Obiang, R.M., Bassat, Q., Kremsner, P.G., Mordmüller,
B., and Grobusch, M.P. (2015). Fosmidomycin as an antimalarial drug: a meta-analysis of
clinical trials. Future Microbiol *10*, 1375–1390.

Friesner, R.A., Banks, J.L., Murphy, R.B., Halgren, T.A., Klicic, J.J., Mainz, D.T., Repasky,
M.P., Knoll, E.H., Shelley, M., Perry, J.K., et al. (2004). Glide: a new approach for rapid,
accurate docking and scoring. 1. Method and assessment of docking accuracy. J. Med. Chem. *47*,
1739–1749.

Friesner, R.A., Murphy, R.B., Repasky, M.P., Frye, L.L., Greenwood, J.R., Halgren, T.A.,
Sanschagrin, P.C., and Mainz, D.T. (2006). Extra precision glide: docking and scoring
incorporating a model of hydrophobic enclosure for protein-ligand complexes. J. Med. Chem.
49, 6177–6196.

Gabriel, H.B., Silva, M.F., Kimura, E.A., Wunderlich, G., Katzin, A.M., and Azevedo, M.F.
(2015). Squalestatin is an inhibitor of carotenoid biosynthesis in *Plasmodium falciparum*.
Antimicrob. Agents Chemother. *59*, 3180–3188.

Gamo, F.J., Sanz, L.M., Vidal, J., de Cozar, C., Alvarez, E., Lavandera, J.-L., Vanderwall, D.E.,
Green, D.V.S., Kumar, V., Hasan, S., et al. (2010). Thousands of chemical starting points for
antimalarial lead identification. Nature *465*, 305–310.

Garrison, E., and Marth, G. Haplotype-based variant detection from short-read sequencing.
Httpsarxiv.orgabs.V *2012*.

Ghosh, S., Chan, J.M.W., Lea, C.R., Meints, G.A., Lewis, J.C., Tovian, Z.S., Flessner, R.M.,
Loftus, T.C., Bruchhaus, I., Kendrick, H., et al. (2004). Effects of bisphosphonates on the growth
of *Entamoeba histolytica* and *Plasmodium* species in vitro and in vivo. J. Med. Chem. *47*, 175–
187.

Giardine, B., Riemer, C., Hardison, R.C., Burhans, R., Elnitski, L., Shah, P., Zhang, Y.,
Blankenberg, D., Albert, I., Taylor, J., et al. (2005). Galaxy: a platform for interactive large-scale
genome analysis. Genome Res. *15*, 1451–1455.

Goecks, J., Nekrutenko, A., Taylor, J., Galaxy Team (2010). Galaxy: a comprehensive approach
for supporting accessible, reproducible, and transparent computational research in the life
sciences. Genome Biol. *11*, R86.

Guiguemde, W.A., Shelat, A.A., Bouck, D., Duffy, S., Crowther, G.J., Davis, P.H., Smithson,
D.C., Connelly, M., Clark, J., Zhu, F., et al. (2010). Chemical genetics of *Plasmodium*
falciparum. Nature *465*, 311–315.

Halgren, T.A., Murphy, R.B., Friesner, R.A., Beard, H.S., Frye, L.L., Pollard, W.T., and Banks,
J.L. (2004). Glide: a new approach for rapid, accurate docking and scoring. 2. Enrichment factors
in database screening. J. Med. Chem. *47*, 1750–1759.

Inselburg, J. (1984). Induction and selection of drug resistant mutants of *Plasmodium*
falciparum. Mol. Biochem. Parasitol. *10*, 89–98.

Inselburg, J. (1985). Induction and isolation of artemisinine-resistant mutants of *Plasmodium*
falciparum. Am. J. Trop. Med. Hyg. *34*, 417–418.

Jahnke, W., Rondeau, J.-M., Cotesta, S., Marzinzik, A., Pellé, X., Geiser, M., Strauss, A., Götte,
M., Bitsch, F., Hemmig, R., et al. (2010). Allosteric non-bisphosphonate FPPS inhibitors
identified by fragment-based discovery. Nat. Chem. Biol. *6*, 660–666.

Jomaa, H., Wiesner, J., Sanderbrand, S., Altincicek, B., Weidemeyer, C., Hintz, M., Türbachova,
I., Eberl, M., Zeidler, J., Lichtenthaler, H.K., et al. (1999). Inhibitors of the nonmevalonate
pathway of isoprenoid biosynthesis as antimalarial drugs. Science *285*, 1573–1576.

Jordão, F.M., Gabriel, H.B., Alves, J.M.P., Angeli, C.B., Bifano, T.D., Breda, A., de Azevedo,
M.F., Basso, L.A., Wunderlich, G., Kimura, E.A., et al. (2013). Cloning and characterization of
bifunctional enzyme farnesyl diphosphate/geranylgeranyl diphosphate synthase from
Plasmodium falciparum. Malar. J. *12*, 184.

Jordão, F.M., Saito, A.Y., Miguel, D.C., de Jesus Peres, V., Kimura, E.A., and Katzin, A.M.
(2011). *In vitro* and *in vivo* antiplasmodial activities of risedronate and its interference with
protein prenylation in *Plasmodium falciparum*. Antimicrob. Agents Chemother. *55*, 2026–2031.

Kavanagh, K.L., Guo, K., Dunford, J.E., Wu, X., Knapp, S., Ebetino, F.H., Rogers, M.J.,
Russell, R.G.G., and Oppermann, U. (2006). The molecular mechanism of nitrogen-containing
bisphosphonates as antiosteoporosis drugs. Proc. Natl. Acad. Sci. U.S.a. *103*, 7829–7834.

Kozarewa, I., Ning, Z., Quail, M.A., Sanders, M.J., Berriman, M., and Turner, D.J. (2009).
Amplification-free Illumina sequencing-library preparation facilitates improved mapping and
assembly of (G+C)-biased genomes. Nat. Methods *6*, 291–295.

Lell, B., Ruangweerayut, R., Wiesner, J., Missinou, M.A., Schindler, A., Baranek, T., Hintz, M.,

Hutchinson, D., Jomaa, H., and Kremsner, P.G. (2003). Fosmidomycin, a novel
chemotherapeutic agent for malaria. Antimicrob. Agents Chemother. *47*, 735–738.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics *25*, 1754–1760.

Liu, J., Liu, W., Ge, H., Gao, J., He, Q., Su, L., Xu, J., Gu, L.-Q., Huang, Z.-S., and Li, D.
(2014). Syntheses and characterization of non-bisphosphonate quinoline derivatives as new
FPPS inhibitors. Biochim. Biophys. Acta *1840*, 1051–1062.

Luckman, S.P., Hughes, D.E., Coxon, F.P., Graham, R., Russell, G., and Rogers, M.J. (1998).
Nitrogen-containing bisphosphonates inhibit the mevalonate pathway and prevent post-
translational prenylation of GTP-binding proteins, including Ras. J. Bone Miner. Res. *13*, 581–
589.

Marzinzik, A.L., Amstutz, R., Bold, G., Bourgier, E., Cotesta, S., Glickman, J.F., Götte, M.,
Henry, C., Lehmann, S., Hartwieg, J.C.D., et al. (2015). Discovery of Novel Allosteric Non-
Bisphosphonate Inhibitors of Farnesyl Pyrophosphate Synthase by Integrated Lead Finding.
ChemMedChem *10*, 1884–1891.

Michael B Martin, Joshua S Grimley, Jared C Lewis, Huel T Heath, III, Brian N Bailey, Howard
Kendrick, Vanessa Yardley, Aura Caldera, Renee Lira, Julio A Urbina, et al. (2001).
Bisphosphonates Inhibit the Growth of *Trypanosoma brucei, Trypanosoma cruzi, Leishmania
donovani, Toxoplasma gondii,* and *Plasmodium falciparum*: A Potential Route to
Chemotherapy. J. Med. Chem. *44*, 909–916.

Nallan, L., Bauer, K.D., Bendale, P., Rivas, K., Yokoyama, K., Hornéy, C.P., Pendyala, P.R.,
Floyd, D., Lombardo, L.J., Williams, D.K., et al. (2005). Protein farnesyltransferase inhibitors
exhibit potent antimalarial activity. J. Med. Chem. *48*, 3704–3713.

Nkrumah, L.J., Muhle, R.A., Moura, P.A., Ghosh, P., Hatfull, G.F., Jacobs, W.R., and Fidock,
D.A. (2006). Efficient site-specific integration in *Plasmodium falciparum* chromosomes
mediated by mycobacteriophage Bxb1 integrase. Nat. Methods *3*, 615–621.

No, J.H., de Macedo Dossin, F., Zhang, Y., Liu, Y.-L., Zhu, W., Feng, X., Yoo, J.A., Lee, E.,
Wang, K., Hui, R., et al. (2012). Lipophilic analogs of zoledronate and risedronate inhibit
Plasmodium geranylgeranyl diphosphate synthase (GGPPS) and exhibit potent antimalarial
activity. Proc. Natl. Acad. Sci. U.S.a. *109*, 4058–4063.

Otto, T.D., Wilinski, D., Assefa, S., Keane, T.M., Sarry, L.R., Böhme, U., Lemieux, J., Barrell,
B., Pain, A., Berriman, M., et al. (2010). New insights into the blood-stage transcriptome of
Plasmodium falciparum using RNA-Seq. Mol. Microbiol. *76*, 12–24.

Paul, A.S., Moreira, C.K., Elsworth, B., Allred, D.R., and Duraisingh, M.T. (2016). Extensive
Shared Chemosensitivity between Malaria and Babesiosis Blood-Stage Parasites. Antimicrob.
Agents Chemother. *60*, 5059–5063.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and

Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.

Rondeau, J.-M., Bitsch, F., Bourgier, E., Geiser, M., Hemmig, R., Kroemer, M., Lehmann, S., Ramage, P., Rieffel, S., Strauss, A., et al. (2006). Structural Basis for the Exceptional in vivo Efficacy of Bisphosphonate Drugs. ChemMedChem *1*, 267–273.

Rottmann, M., McNamara, C., Yeung, B.K.S., Lee, M.C.S., Zou, B., Russell, B., Seitz, P., Plouffe, D.M., Dharia, N.V., Tan, J., et al. (2010). Spiroindolones, a potent compound class for the treatment of malaria. Science *329*, 1175–1180.

Singh, A.P., Zhang, Y., No, J.H., Docampo, R., Nussenzweig, V., and Oldfield, E. (2010). Lipophilic bisphosphonates are potent inhibitors of *Plasmodium* liver-stage growth. Antimicrob. Agents Chemother. *54*, 2987–2993.

Sinigaglia, L., Varenna, M., and Casari, S. (2007). Pharmacokinetic profile of bisphosphonates in the treatment of metabolic bone disorders. Clin Cases Miner Bone Metab *4*, 30–36.

Spalding, M.D., Allary, M., Gallagher, J.R., and Prigge, S.T. (2010). Validation of a modified method for Bxb1 mycobacteriophage integrase-mediated recombination in *Plasmodium falciparum* by localization of the H-protein of the glycine cleavage complex to the mitochondrion. Mol. Biochem. Parasitol. *172*, 156–160.

Spangenberg, T., Burrows, J.N., Kowalczyk, P., McDonald, S., Wells, T.N.C., and Willis, P. (2013). The open access malaria box: a drug discovery catalyst for neglected diseases. PLoS ONE *8*, e62906.

Straimer, J., Lee, M.C.S., Lee, A.H., Zeitler, B., Williams, A.E., Pearl, J.R., Zhang, L., Rebar, E.J., Gregory, P.D., Llinás, M., et al. (2012). Site-specific genome editing in *Plasmodium falciparum* using engineered zinc-finger nucleases. Nat. Methods *9*, 993–998.

Taylor, K.B. (2004). Slow and Tight Inhibition. In Enzyme Kinetics and Mechanisms, (Dordrecht: Kluwer Academic Publishers), pp. 122–146.

Thorvaldsdóttir, H., Robinson, J.T., and Mesirov, J.P. (2013). Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Brief. Bioinformatics *14*, 178–192.

Tonhosolo, R., D'Alexandri, F.L., de Rosso, V.V., Gazarini, M.L., Matsumura, M.Y., Peres, V.J., Merino, E.F., Carlton, J.M., Wunderlich, G., Mercadante, A.Z., et al. (2009). Carotenoid biosynthesis in intraerythrocytic stages of *Plasmodium falciparum*. J. Biol. Chem. *284*, 9974–9985.

Tonhosolo, R., D'Alexandri, F.L., Genta, F.A., Wunderlich, G., Gozzo, F.C., Eberlin, M.N., Peres, V.J., Kimura, E.A., and Katzin, A.M. (2005). Identification, molecular cloning and functional characterization of an octaprenyl pyrophosphate synthase in intra-erythrocytic stages of *Plasmodium falciparum*. Biochem. J. *392*, 117–126.

Tsuchiya, T., Ishibashi, K., Terakawa, M., Nishiyama, M., Itoh, N., and Noguchi, H. (1982).

Pharmacokinetics and metabolism of fosmidomycin, a new phosphonic acid, in rats and dogs.
Eur J Drug Metab Pharmacokinet *7*, 59–64.

Ullah, I., Sharma, R., Biagini, G.A., and Horrocks, P. (2017). A validated bioluminescence-
based assay for the rapid determination of the initial rate of kill for discovery antimalarials. J.
Antimicrob. Chemother. *72*, 717–726.

Van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., Del Angel, G., Levy-Moonshine,
A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., et al. (2013). From FastQ data to high
confidence variant calls: the Genome Analysis Toolkit best practices pipeline. Curr Protoc
Bioinformatics *43*, 11.10.1–.10.33.

Van Voorhis, W.C., Adams, J.H., Adelfio, R., Ahyong, V., Akabas, M.H., Alano, P., Alday, A.,
Alemán Resto, Y., Alsibaee, A., Alzualde, A., et al. (2016). Open Source Drug Discovery with
the Malaria Box Compound Collection for Neglected Diseases and Beyond. PLoS Pathog. *12*,
e1005763.

Van Voorhis, W.C., Rivas, K.L., Bendale, P., Nallan, L., Hornéy, C., Barrett, L.K., Bauer, K.D.,
Smart, B.P., Ankala, S., Hucke, O., et al. (2007). Efficacy, pharmacokinetics, and metabolism of
tetrahydroquinoline inhibitors of *Plasmodium falciparum* protein farnesyltransferase.
Antimicrob. Agents Chemother. *51*, 3659–3671.

Wu, W., Herrera, Z., Ebert, D., Baska, K., Cho, S.H., Derisi, J.L., and Yeh, E. (2015). A
chemical rescue screen identifies a Plasmodium falciparum apicoplast inhibitor targeting MEP
isoprenoid precursor biosynthesis. Antimicrob. Agents Chemother. *59*, 356–364.

Yeh, E., and Derisi, J.L. (2011). Chemical rescue of malaria parasites lacking an apicoplast
defines organelle function in blood-stage *Plasmodium falciparum*. PLoS Biol. *9*, e1001138.

Yokoyama, T., Mizuguchi, M., Ostermann, A., Kusaka, K., Niimura, N., Schrader, T.E., and
Tanaka, I. (2015). Protonation State and Hydration of Bisphosphonate Bound to Farnesyl
Pyrophosphate Synthase. J. Med. Chem. *58*, 7549–7556.

Zhang, B., Watts, K.M., Hodge, D., Kemp, L.M., Hunstad, D.A., Hicks, L.M., and Odom, A.R.
(2011). A second target of the antimalarial and antibacterial agent fosmidomycin revealed by
cellular metabolic profiling. Biochemistry *50*, 3570–3577.

Zhang, Y., Cao, R., Yin, F., Hudock, M.P., Guo, R.-T., Krysiak, K., Mukherjee, S., Gao, Y.-G.,
Robinson, H., Song, Y., et al. (2009). Lipophilic bisphosphonates as dual
farnesyl/geranylgeranyl diphosphate synthase inhibitors: an X-ray and NMR investigation. J.
Am. Chem. Soc. *131*, 5153–5162.

Zhang, Y., Zhu, W., Liu, Y.-L., Wang, H., Wang, K., Li, K., No, J.H., Ayong, L., Gulati, A.,
Pang, R., et al. (2013). Chemo-Immunotherapeutic Anti-Malarials Targeting Isoprenoid
Biosynthesis. ACS Med Chem Lett *4*, 423–427.

Tables

Table 1. Summary of mutations identified by whole genome sequencing

Gene ID[a]	Description[b]	Base Call[c]			AA change	Population	Read Number[d]		%[e]	
		Position	WT	EMS			WT	R	WT	R
PF3D7_1128400	Geranylgeranyl pyrophosphate synthase (GGPPS)	682	T	A	S228T	019313R1	219	163	100	100
						019313R2	186	181	100	62.4

815   [a] PlasmoDB gene Identification number
816   [b] Basic gene description based on PlasmoDB functional assignments
817   [c] WT calls match 3D7 reference genome
818   [d] Read depth corresponding to WT call (IGV) and Mut call (SnpEff) respectively
819   [e] Percent of reads corresponding to WT call (IGV) and Mut call (SnpEff) respectively
820
821
822   **Figures and Figures legends**
823

824
825
826   **Figure 1. IPP rescues growth inhibition by MMV019313. A.** The structure of MMV019313.
827   **B.** EC_{50} curves in the absence (solid line) and presence (dotted line) of IPP. Parasitemia is
828   normalized to that of an untreated control. Results shown for experiments performed in
829   biological triplicate with technical duplicate. Shown plotted as mean ±SD.
830

831
832
Figure 2. MMV019313-resistant parasites contain a mutation in the bifunctional farnesyl
and geranylgeranyl diphosphate synthase. **A.** EC_{50} curves of parental W2 parasites alone
(black, solid line) and with IPP supplementation (black, dotted line) and the two resistant
populations (019313R1, red line, and 019313R2, green line) without IPP added. Fold change in
EC_{50} compared to W2 parasites is shown in the right. Results shown for experiments performed
in biological triplicate with technical duplicate. Shown plotted as mean ±SD. **B**. Mutation
determined by whole genome sequencing of resistant populations 019313R1 and 019313R2,
highlighted in red **C.** Schematic of *Pf*FPPS/GGPPS protein. The S228T residue is highlighted in
bold black while conserved KT and SARM residues are highlighted in bold grey and underlined.
842

843
844
Figure 3. Overexpression of WT and an S228T variant of *Pf*FPPS/GGPPS confer resistance
to MMV019313. A. EC_{50} curves of MMV019313 against the parental Dd2^{attB} parasites (black)
and parasites over-expressing FPPS/GGPPS-GFP under the RL2 (red) or CaM (green) promoter.
B. EC_{50} curves of MMV019313 against the parental Dd2^{attB} parasites (black) and parasites over-
expressing wild type (red) or mutant (S228T, blue) FPPS/GGPPS-GFP under the RL2 promoter.
Results shown for experiments performed in biological duplicate with technical duplicate.
Shown plotted as mean ±SD.
852

853
854
Figure 4. **MMV019313 has a specific and distinct mode-of-inhibition against purified**
***Pf*FPPS/GGPPS.** Dose-dependent inhibition of wild-type PfFPPS/GGPPS (solid),
PfFPPS/GGPPS S228T variant (dotted), and either human FPPS or GGPPS (dashed) of **A**. FPP
(C15) production by MMV019313, **B**. GGPP (C20) production by MMV010313, **C**. FPP (C15)
production by BPH-703, and **D**. GGPP (C20) production by BPH-703. Partial inhibition of
*Hs*GGPPS by BPH-703 was observed under k_{cat} conditions, while complete inhibition was
observed under non-saturating conditions (Figure S5). Shown plotted as mean ±SD.

862
863