Multifactorial Chromosomal Variants Regulate Polymyxin Resistance in

Extensively Drug-Resistant *Klebsiella pneumoniae*

Miranda E Pitt[1], Alysha G Elliott[1], Minh Duc Cao[1], Devika Ganesamoorthy[1], Ilias

Karaiskos[2], Helen Giamarellou[2], Cely S Abboud[3], Mark AT Blaskovich[1], Matthew A

Cooper[1]* and Lachlan Coin[1]*

[1]Institute for Molecular Bioscience, University of Queensland, Brisbane, Australia

[2]6th Department of Internal Medicine, Hygeia General Hospital, Athens, Greece

[3]Instituto Dante Pazzanese de Cardiologia, São Paulo, Brazil

Running Title: Polymyxin Resistance in XDR *K. pneumoniae*

*Address correspondence to authors: Lachlan Coin (l.coin@imb.uq.edu.au) & Matthew A Cooper

(m.cooper@uq.edu.au)

ABSTRACT

Extensively drug-resistant *Klebsiella pneumoniae* (XDR-KP) infections cause high mortality and are disseminating globally. Identifying the genetic basis underpinning resistance allows for rapid diagnosis and treatment. XDR isolates sourced from Greece and Brazil, including nineteen polymyxin-resistant and five polymyxin-susceptible strains, underwent whole genome sequencing. Approximately 90% of polymyxin resistance was enabled by alterations upstream or within *mgrB*. The most common mutation identified was an insertion at nucleotide position 75 in *mgrB* via an IS*Kpn26*-like element in the ST258 lineage and IS*Kpn13* in one ST11 isolate. Three strains acquired an IS*1* element upstream of *mgrB* and another strain had an IS*Kpn25* insertion at 133 bp. Other isolates had truncations (C28STOP, Q30STOP) or a missense mutation (D31E) affecting *mgrB*. Complementation assays revealed all *mgrB* perturbations contributed to resistance. Missense mutations in *phoQ* (T281M, G385C) were also found to facilitate resistance. Several variants in *phoPQ* co-segregating with the IS*Kpn26*-like insertion were identified as potential partial suppressor mutations. Three ST258 samples were found to contain subpopulations with different resistance conferring mutations, including the IS*Kpn26*-like insertion colonising with a novel mutation in *pmrB* (P158R), both confirmed via complementation assays. We also characterized a new multi-drug resistant *Klebsiella quasipneumoniae* strain ST2401 which was susceptible to polymyxins. These findings highlight the broad spectrum of chromosomal modifications which can facilitate and regulate resistance against polymyxins in *K. pneumoniae*.

DATA SUMMARY

1. Whole genome sequencing of the 24 clinical isolates has been deposited under BioProject PRJNA307517 (www.ncbi.nlm.nih.gov/bioproject/PRJNA307517).

IMPACT STATEMENT

Klebsiella pneumoniae contributes to a high abundance of nosocomial infections and the rapid emergence of antimicrobial resistance hinders treatment. Polymyxins are predominantly utilized to treat multidrug-resistant infections, however, resistance to the polymyxins is arising. This increasing prevalence in polymyxin resistance is evident especially in Greece and Brazil. Identifying the genomic variations conferring resistance in clinical isolates from these regions assists with potentially detecting novel alterations and tracing the spread of particular strains. This study commonly found mutations in the gene *mgrB*, the negative regulator of PhoPQ, known to cause resistance in KP. In the remaining isolates, missense mutations in *phoQ* were accountable for resistance. Multiple novel mutations were detected to be segregating with *mgrB* perturbations. This was either due to a mixed heterogeneous sample of two polymyxin-resistant strains, or because of multiple mutations within the same strain. Of interest was the validation of novel mutations in *phoPQ* segregating with a previously known IS*Kpn26*-like element in disrupted *mgrB* isolates. Complementation of these *phoPQ* mutations revealed a reduction in minimum inhibitory concentrations and suggests the first evidence of partial suppressor mutations in KP. This research builds upon our current understanding of heteroresistance, lineage specific mutations and regulatory variations relating to polymyxin resistance.

INTRODUCTION

Klebsiella pneumoniae (KP) strains classified as extensively drug-resistant (XDR) are rapidly emerging due to the dissemination of plasmid-encoded resistance towards aminoglycosides, β-lactams, fluoroquinolones and carbapenems [1]. Notably, carbapenem-resistant KP have been linked to high morbidity and an overall mortality of 48% in infected patients [2]. Polymyxin B and colistin (polymyxin E) are now one of the last viable therapeutic options [3]. Unfortunately, resistance to this last-line antibiotic class is an increasing global burden, with countries particularly impacted including Asia (Korea [4, 5], India [6, 7]), Europe (Greece [8-10]), Italy [10, 11]) and Latin America (Brazil [12, 13]). There is considerable debate regarding the mortality associated with polymyxin-resistant infections. Combining several clinical cohorts has provided an overall mortality estimate ranging from 20 to 100% which was dependent on early detection of the outbreak [14].

Polymyxins infiltrate Gram-negative bacteria via initial binding to the basal component of lipopolysaccharide, lipid A. This causes the displacement of Mg^{2+} and Ca^{2+}, disrupting bacterial outer membrane integrity, allowing the polymyxins to traverse the inner membrane and act on intracellular targets. An extended exposure in KP triggers the activation of the two-component regulatory systems, PmrAB and PhoPQ [15-17]. These systems regulate a pathway that modulates *pmrC* and the *pmrHFIJKLM* operon facilitating the addition of phosphoethanolamine (pEtN) and/ or 4-amino-4-deoxy-L-arabinose to lipid A phosphate groups, resulting in impaired polymyxin binding interactions [18-20]. Disruption of *mgrB*, the negative regulator of PhoPQ, has been commonly observed in isolates of clinical origin [8, 21]. The constitutive up-regulation of *pmrC*

and the *pmrHFIJKLM* operon incurs a minimal fitness cost and appears to be stable, with minimal

reports of reversions [22, 23]. Heteroresistant populations, where only a subset of bacteria are

resistant, have been reported in KP which complicates diagnosis [24]. The emergence of pandrug-

resistant KP is of grave concern [25] and this acquisition of resistance is further exacerbated by

the recently reported plasmid-encoded colistin resistance gene *mcr-1*, which encodes a pEtN

transferase enzyme, albeit currently rare in KP [26].

This study aimed to investigate XDR-KP clinical isolates arising in Greece and Brazil during 2012

to 2014 to identify and validate genetic variants contributing to resistance. These alterations were

compared to prior clinical isolates to ascertain if these mutations have been previously detected

globally.

METHODS

Bacterial isolates

KP clinical isolates were acquired from the Hygeia General Hospital, Athens, Greece and Instituto Dante Pazzanese de Cardiologia, Brazil from patients in 2012 to 2014. Cultures were supplied as stabs/slants or on agar, and were subsequently cultured in Nutrient Broth. Cultures were made to 20% (v/v) glycerol and stored at -80 °C. When required for assay or extraction, glycerol stocks were struck out to obtain single colonies on either Nutrient Agar or Tryptic Soy Agar with 5% defibrinated sheep blood. Reference strains included *Escherichia coli* (ATCC 25922) and *Klebsiella spp*. (ATCC 13883, ATCC 700603, ATCC BAA-2146), which were obtained from the American Type Culture Collection (ATCC; Manassas, VA, USA).

Antimicrobial susceptibility assays

Species identification and susceptibility profiles of clinical isolates from Greece and Brazil were evaluated in the clinic using VITEK®2 (bioMérieux). Strains were further validated at the Institute for Molecular Bioscience (IMB) (The University of Queensland, Australia) using the standard Clinical & Laboratory Standards Institute (CLSI) approved broth microdilution (BMD) methods with cation-adjusted Mueller-Hinton Broth (caMHB). Resistance was determined as per CLSI guidelines [27] except for tigecycline and fosfomycin where The European Committee on Antimicrobial Susceptibility Testing (EUCAST) (Version 7.1, 2017) (see http://www.eucast.org) guidelines were implemented. Categorisation of drug resistance level was determined through guidelines previously outlined [28].

DNA extraction

DNA was extracted from overnight cultures using the DNeasy Blood and Tissue Kit (Qiagen) with the additional enzymatic lysis buffer pre-treatment as per manufacturer's instructions. DNA was quantified with Qubit®3.0 (ThermoFisher Scientific).

DNA library preparation and sequencing

Library preparation was performed using the Nextera XT kit (Illumina) with 1 ng input of DNA as per manufacturer's instructions. Quality of libraries were checked using a 2100 Bioanalyzer (Agilent Technologies). Libraries were sequenced on an Illumina MiSeq with 300 bp paired-end sequencing reads and >100X coverage per sample.

Sequencing analysis

Paired-end reads were trimmed with Trimmomatic [29] and assembled using SPAdes [30]. The Rapid Annotation using Subsystem Technology (RAST) was utilized to annotate assembled genomes [31]. Assemblies were also uploaded to the Centre for Genomic Epidemiology (CGE) to identify sequence types (STs) (MultiLocus Sequence Typing Server 1.8 [32]) and acquired antibiotic resistance genes (ResFinder 2.1 [33]). A neighbor-joining tree was constructed using the 2358 *Klebsiella pneumoniae/quasipneumoniae/variicola* genes known to form the core genome MLST (cgMLST) using Ridom SeqSphere+ v4.0.1 software [34]. The cgMLST was compared against complete assemblies of ST11 (HS11286), ST147 (MS6671), ST258 (NJST258_1, NJST258_2) and a reference for *K. quasipneumoniae*, ATCC 700603 [25, 35-37].

Variant detection

Alterations both in and flanking the genes *pmrAB*, *phoPQ* and *mgrB* were examined and sequence reads of all strains were aligned to the assembly of 20_GR_12, a polymyxin-susceptible ST258 strain with the least number of contigs, using BWA-MEM [38]. The alignment was analyzed through FreeBayes [39] to identify single nucleotide and small indel variation, using a diploid analysis in order to identify potential heterogeneity. Sites with more than 20% of reads mapping to the minor allele were considered potentially heterogeneous. The effects of variations were determined by snpEff [40]. The impact on protein sequence was further confirmed by the Protein Variation Effect Analyzer (PROVEAN) [41]. For the analysis of large chromosome changes, the gene sequences including 300 bp flanking were extracted from the assemblies. A multiple alignment of each gene was constructed from the pair-wise alignment to the longest gene sequence.

Insertion sequence element validation

ISFinder [42] was used for the identification of insertion sequence (IS) elements. To confirm disruptive IS elements, *mgrB* was amplified with primers displayed in Table S1 via 2X Phusion HF master mix (Invitrogen) under the following cycling conditions: 98 °C 10 seconds, 50 °C 30 seconds and 72 °C 60 seconds (35X). Amplicon identity was validated via Sanger sequencing.

Complementation assays

The contribution of variants to resistance was validated through complementation assays as previously described [43]. Briefly, genes (Table S1) were amplified from a polymyxin-susceptible isolate, 20_GR_12, and cloned into the pCR-BluntII-TOPO vector via the Zero Blunt TOPO PCR cloning kit (Invitrogen). Chemically competent *E. coli* TOP10 cells were transformed and selected by the addition of 50 mg/L kanamycin in MHA. Isolation of plasmids were via the PureLink™

Quick Plasmid Miniprep Kit (Invitrogen) and transformed into KP strains via electroporation (25

μF, 200 Ω, 1.25 kV/cm) with a Gene Pulser (Bio-Rad Laboratories). Selection was accomplished

through supplementation of ≥500 mg/L zeocin in MHA plates. Transformed colonies (n=≥2) were

acquired and placed in MHB containing 1500 mg/L zeocin and 1 mM isopropyl *β-D*-1-

thiogalactopyranoside (Sigma Aldrich). If polymyxin susceptibility was not restored upon

complementation, genes harboring mutations were further amplified and introduced into

20_GR_12. To discern the impact of additional mutations in *phoPQ* and *pmrB* segregating with

disrupted *mgrB*, mutant genes were introduced into a polymyxin-resistant isolate only harboring

an IS element *mgrB* disruption, 7_GR_13. Controls included transformation of WT genes into

20_GR_12, sequencing of amplicon prior to introduction in vector and KP transformed strains

undergoing a plasmid extraction and further PCR of the multiple cloning site. Antimicrobial testing

against polymyxin B were conducted as described above.

RESULTS

Characterization of clinical isolates

KP isolates were all characterized in the hospital microbiology facility using VITEK®2. Several discrepancies were detected between VITEK®2 and broth microdilution (BMD) results (Table 1, Table S2), predominantly the level of resistance towards aminoglycosides, tetracyclines, fosfomycin and tigecycline. A major dissimilarity was polymyxin susceptibility in 6_GR_12 (sensitive in BMD, resistant in VITEK®2) and resistant in 23_GR_13 (resistant in BMD, sensitive in VITEK®2). Polymyxin resistance was identified in 19 of the isolates. An abundance of acquired resistance genes (Table 2) were detected and this presence corresponded to the antimicrobial testing phenotype. This analysis did not identify *mcr-1* in these strains. Only 18_GR_14 and 19_GR_14 were not identified as extended-spectrum beta-lactamase producers amongst the polymyxin-resistant strains. Consequently, all polymyxin-resistant strains that harbored non-susceptibility to at least one antibiotic in 15 or more of the 17 antimicrobial categories hence were defined as XDR.

Sequence type determination

Two thirds of the Greece clinical strains were found to belong to ST258 and the remaining were ST11, ST147 or ST383 (Table 1). While 5_GR_13 and 6_GR_12 were both ST383, only 5_GR_13 was resistant to polymyxin. Among the two strains from Brazil, 11_BR_13 was ST437 and 12_BR_13 was ST11. 21_GR_13 had a profile previously undefined and has been newly designated ST2401. Further cgMLST studies were conducted on the isolates using complete assemblies of reference genomes for ST11 (HS11286), ST147 (MS6671) ST258 (NJST258_1,

NJST258_2) and KQ (ATCC 700603) (Fig. 1). For the ST258 isolates, these were more similar to

NJST258_2 rather than NJST258_1. Within this cluster, 7_GR_13, 9_GR_12 and 24_GR_13 were

closely related (≤15 allelic changes). Similarly, grouped together were 2_GR_12 and 23_GR_12;

3_GR_13 and 22_GR_12; 13_GR_14 and 14_GR_14; and 18_GR_14 and 19_GR_14. In ST11,

16_GR_13 and 17_GR_14 harbored only 3 allele differences and the Brazilian isolate, 12_BR_13,

had 206 variants apparent. ST383 isolates 5_GR_13 and 6_GR_12 only exhibited 1 allele change.

ST147 1_GR_13 was not clonal to the previous pandrug resistant KP, MS6671. Clustering analysis

revealed 21_GR_13 as *Klebsiella quasipneumoniae* (KQ) and diverged with reference genome

ATCC 700603 (ST489).

MgrB disruption

Seventeen of the nineteen polymyxin-resistant strains exhibited either missense mutations,

nonsense mutations or IS elements in *mgrB* (Table 3). Both 5_GR_13 and 19_GR_14 harbored a

truncation while an amino acid change, D31E, was apparent in 3_GR_13. IS element disruption

was prevalent in 53% of strains and commonly an IS*5*-like element was integrated at nucleotide

position 75 (Fig. S1). Sanger sequencing revealed this element was closely related to IS*Kpn26*,

herein known as IS*Kpn26*-like, except for 12_BR_13 which matched IS*Kpn13.* IS*1R* was detected

upstream of *mgrB* in 11_BR_13 and an IS*1R*-like (A>C, 393 bp; C>T, 396 bp) element in

16_GR_13 and 17_GR_14. Strain 15_GR_13 had a deletion of the *mgrB* locus from nucleotide

position 133 onwards. The 127 bp flanking region mapped to IS*Kpn25* with the transposase in the

same orientation as *mgrB*. All 3 of IS*1* element insertions, but only one of the 8 IS*Kpn26*-like

element insertions had the transposase in the same orientation as *mgrB*.

Single, multiple and heterogeneous mutations

Aberrations in genes commonly identified to confer polymyxin resistance in KP include *mgrB*, *phoPQ* and *pmrAB* (Table 2). Several non-synonymous mutations were identified across the isolates however, not all were predicted to be deleterious (Table S3). ST383 contained several mutations in *pmrAB* although only Q30STOP in polymyxin-resistant 5_GR_13 was predicted to have an impact. Similarly, neutral changes in all four of these genes were detected in polymyxin-susceptible KQ strains ATCC 700603 and 21_GR_13. 8_GR_13 and 9_GR_12 harbored a single detrimental missense mutation in *phoQ*. Alterations in *mgrB* were accompanied by one or more missense mutations in *phoPQ* and/ or *pmrB*. Predicted deleterious variants segregating with disrupted *mgrB* included *pmrB* (T140P, P158R), *phoP* (P74L, A95S) and *phoQ* (N253T, V446G), which were commonly in the ST258 lineage. V446G (*phoQ*) and P158R (*pmrB*) were heterogeneous in 13_GR_14 (65% (V446G), 66% (P158R) mutation allele frequency) and 14_GR_14 (52% (V446G) and 57% (P158R) mutation allele frequency). Assembly revealed 23_GR_12 harbored an IS*Kpn26*-like disrupted *mgrB* alongside the intact version with alterations in *phoP* and *phoQ* in 57% of the sample.

Role of *mgrB* disruptions and presence of heteroresistance via complementation assays

Complementation of the WT gene elucidated the role of these mutations in resistance (Fig. 2). Introduction of pTOPO-*mgrB* restored susceptibility in all resistant isolates with *mgrB* coding mutations or upstream disruptions, with the exception of two strains heterogeneous for the *mgrB* disruption and a *pmrB* coding mutation (13_GR_14 and 14_GR_14) (Fig. 2a). For these two strains, pTOPO-*mgrB* restored susceptibility in zero of three 13_GR_14 colonies and one of three 14_GR_14 colonies. Transformation of 1 out of 3 colonies for both 13_GR_14 and 14_GR_14 strains with pTOPO-*pmrB* restored susceptibility (Fig. 2d) and *mgrB* amplification of these

colonies revealed an intact *mgrB* locus (data not shown). Colonies which were reverted on

complementation were further passaged 3 times with no antibiotic pressure in order to remove the

plasmid and discern if these mutations were contributing to resistance. After passaging, pTOPO-

mgrB isolates harbored an MIC of ≥64 mg/L whilst pTOPO-*pmrB* colonies were 16 mg/L to

confirm two resistant populations in these samples. 23_GR_12 was also observed to have a

heterogeneous *mgrB* disruption but did not carry a corresponding *pmrB* mutation however,

harbored similar mutations to 2_GR_12 in *phoPQ*. Amplification of *mgrB* identified two of three

23_GR_12 transformed colonies contained the IS element disruption and were reverted to

susceptible upon complementation with pTOPO-*mgrB*.

Validation of resistance conferring mutations in *phoQ*

Strains 8_GR_13 and 9_GR_12 harbored a single mutation in *phoQ* potentially conferring

resistance (Table 2). When these isolates were transformed with pTOPO-*phoQ*, results remained

variable where a lack of growth was present in a susceptible range (MIC: ≤2 mg/L) however,

several wells containing high polymyxin B concentrations exhibited growth (Fig. 2c). To resolve

this, the mutated gene was introduced into a polymyxin-susceptible isolate, 20_GR_12, and

resistance was apparent (Fig. 2e).

Potential suppressor mutations in *phoPQ*

Several mutations co-segregating with disrupted *mgrB* were detected including *phoP* (P74L,

A95S), *phoQ* (N253T, V446G) and *pmrB* (T140P). Complementation of WT genes in these

isolates commonly facilitated a ≥2-fold increase in MIC with the exception of 10_GR_13, which

had an additional predicted neutral mutation in *phoQ* (A225T) (Table S3, Fig. 2b-d). To evaluate

the potential influence of these mutations on polymyxin resistance, mutated genes were placed

into a strain only containing the *mgrB* IS element disruption, 7_GR_13 (Fig. 2f). Complementation

of the mutant *phoQ* (N253T) decreased the MIC by 2-fold, potentially indicating a partial

suppressor mutation. Initially, the *phoQ* (V446G) mutation was anticipated to segregate with the

mgrB disrupted population in 13_GR_14 and 14_GR_14 however, when *phoQ* was amplified from

a colony reverted to susceptible via pTOPO-*mgrB* complementation, the WT *phoQ* was observed

(Fig. S3). The *phoQ* (V446G) mutation was successfully amplified from a 14_GR_14 colony

containing the *pmrB* (T158R) mutation and upon complementation in 7_GR_13, resulted in a 2-

fold reduction in MIC. Although this mutation did not segregate with disrupted *mgrB*, it may act

as a partial suppressor mutation when a resistance conferring mutation is present in *pmrB*.

Mutations in *phoP* (P74L, A95S) reduced the MIC in 7_GR_13 by ≥4-fold which identifies these

as partial suppressor mutations. Complementation of mutant *pmrB* (T140P) into 7_GR_13 did not

lead to an observable corresponding reduction in MIC however, once transformed into 20_GR_12,

a 2-fold increase in MIC was apparent (Fig. 2e).

DISCUSSION

Polymyxin resistance in XDR-KP is of grave concern given that this is a last-line antibiotic, and is increasingly prevalent in countries such as Greece and Brazil [10, 12-14, 44]. We evaluated the genetic basis of polymyxin resistance in a series of Greek and Brazilian clinical isolates from patients in 2012 to 2014 and found alterations in genes *mgrB*, *phoPQ* and *pmrAB*.

Inactivation of *mgrB* was highly prevalent in these strains with an IS*Kpn26*-like element being the predominant cause of resistance, as confirmed by complementation restoring susceptibility in all isolates. Several other studies have observed an IS*5*-like element integration in the same position, including reports from Greece, Italy, France, Turkey and Colombia [8, 9, 45, 46]. The IS*Kpn26*-like element resembled the same sequence from Greece isolates previously described [46]. We identified that this mutation still persisted in 2014, after being first detected in 2012 [9]. Disruptions in *mgrB* including the IS*Kpn26*-like forward insertion at nucleotide 75 in ST147, IS*Kpn13* integration at nucleotide 75 in ST11 and IS*Kpn25* in the ST258 lineage have yet to be reported. We identified IS*1R* or IS*1R*-like elements positioned upstream of *mgrB* in 3 isolates (11_BR_13, 16_GR_13, 17_GR_14) which were reverted upon complementation indicating an impact on the promoter region.

Truncations identified at position 28 and 30 of *mgrB* have been previously detected, although these were identified in differing STs indicating mutations potentially have arisen independently in Greece [21, 47]. Complementation restored susceptibility to polymyxins for these mutations and this study further revealed the amino acid change D31E in 3_GR_13 to be a resistance conferring alteration. These findings support the notion that intact MgrB is required to confer negative

feedback on PhoPQ [8]. The inactivation of *mgrB* is prevalent in polymyxin-resistant KP and may

arise owing to its capacity to promote virulence and further attenuate the early host defence

response, with little or no fitness cost [48].

Single predicted detrimental mutations were observed in the *phoQ* histidine kinase region, critical

for phosphorylation and interaction with *phoP*, in 8_GR_13 (G385C) and 9_GR_12 (T281M). The

G385C mutation had previously been reported, [21] however in a differing ST. Complementation

revealed an inconsistent MIC for these strains, although when a polymyxin-susceptible isolate was

transformed with the mutated gene, full resistance was restored. Dominance of mutated *phoQ* has

recently been highlighted and these results may imply the inability of pTOPO-*phoQ* to override

the resistance caused by these mutations [49].

Several non-synonymous changes were identified to be not deleterious according to PROVEAN

analysis. Notably, these were abundant in KQ strains ATCC 700603 [37] and 21_GR_13. This

was further identified in KP ST383 lineages and PROVEAN detected these neutral changes. These

mutations represent lineage specific alterations, however, this does not negate the possibility of

previously resistance conferring alterations being acquired in these loci with subsequent reversion

mutations to give rise to a susceptible phenotype.

Heterogeneity was apparent in several isolates. In near equal ratios, 13_GR_14 and 14_GR_14

possessed the IS*Kpn26*-like *mgrB* disruption and a new alteration conferring resistance in *pmrB*,

P158R as determined by complementation. 23_GR_12 consisted of approximately half the reads

mapping to the undisrupted genes and the other to the IS*Kpn26*-like strain with several additional

predicted deleterious mutations. This heterogeneity may explain the initial clinical detection for

this isolate to be polymyxin-susceptible.

Several isolates harboring IS*Kpn26*-like element disrupted *mgrB* were accompanied by mutations in *phoPQ* and/ or *pmrB*. These changes were present in ≥98% of reads to render the involvement of heterogeneity unlikely. Once complemented, an increase in resistance was commonly recorded. This potentially reflects partial suppressor mutations as strains which solely possessed this IS element disruption commonly exhibited a heightened MIC of ≥64 mg/L. One variant segregating with this disruption included *pmrB* T140P. This had formerly been identified in an ST258 lineage but even when the resistant gene was complemented, the MIC increased by 2-fold but was not defined as clinically resistant [21, 50].

When mutated *phoP* or *phoQ* were introduced into the *mgrB* disrupted isolate, a reduction in MIC was apparent. The involvement of additional mutations in PhoPQ to influence the level of polymyxin resistance has yet to be reported in KP. Previous research by Miller *et al* [51] determined additional mutations in PhoPQ altered polymyxin resistance in *Pseudomonas aeruginosa*. This prior study describes *phoP* mutations with the capacity to partially or fully suppress resistance-causing mutations in *phoQ*. These mutations in *phoP* were near or within the DNA binding site which differs to our results, where the alterations are impacting the response regulatory region that interacts with PhoQ. Conversely, all mutations partially suppressing the MIC were identified to be targeting the HAMP and histidine kinase component of PhoQ. These were in regions similar to revertant *P. aeruginosa* strains identified by Lee and Ko [52]. We postulate these mutations are perturbing the critical transfer of phosphoryl groups from the histidine kinase of PhoQ to PhoP and subsequent *pmrD* expression. Whether these mutations constitute a fitness advantage due to the reduction of metabolism required for the production of LPS modifications is yet to be discerned. Furthermore, due to variability in some of the

complementation data, a knockout *phoPQ* background and introduction of genes that are potential

suppressor mutations is required.

Rapid and accurate detection of mutations attributed to polymyxin resistance remains a

longstanding burden. Our research has contributed to the current understanding of the

dissemination and evolution of this resistance in KP. Although the sample size is limited, this study

highlights several issues arising from solely interrogating genomes for resistance detection

including ST specific non-synonymous changes, and heterogeneity. The study provides the first

potential report of suppressor mutations for polymyxin resistance. Through complementation

assays, we have discerned the role of these modifications and have identified resistance-causing

alterations that can be monitored during future genome-based diagnostics.

AUTHOR STATEMENTS

Funding information

LC is an ARC Future Fellow (FT110100972). MAC is an NHMRC Principal Research Fellow (APP1059354) and currently holds a fractional Professorial Research Fellow appointment at the University of Queensland with his remaining time as CEO of Inflazome Ltd. a company headquartered in Dublin, Ireland that is developing drugs to address clinical unmet needs in inflammatory disease by targeting the inflammasome. MEP is an Australian Postgraduate Award scholar. AGE and MATB are supported in part by a Wellcome Trust Strategic Award 104797/Z/14/Z. Research was supported by NHMRC grants (APP1005350, APP1045326), NIH grant (R21AI098731/R33AI098731) and AID sequencing Grant (2013) as well as funding from the Institute for Molecular Bioscience Centre for Superbug Solutions (610246).

Acknowledgements

We thank Dr Aurélie Jayol and Professor Patrice Nordmann for providing their complementation assay methodology. The authors thank Maite Amado and Angela M. Kavanagh for technical support with susceptibility assays and Rhia Stone for the quality control of antibiotics. We acknowledge the sequencing services provided by the Australian Genome Research Facility. We thank the team of the curators of the Institut Pasteur MLST system (Paris, France) for importing novel alleles, profiles and/or isolates at http://bigsdb.web.pasteur.fr.

Conflicts of interest

None.

ABBREVIATIONS

BMD, Broth microdilution; caMHB, cation-adjusted Mueller-Hinton broth; CLSI, Clinical & Laboratory Standards Institute; EUCAST, The European Committee on Antimicrobial Susceptibility Testing; IS, Insertion sequence; KP, *Klebsiella pneumoniae*; KQ, *Klebsiella quasipneumoniae*; MHA, Mueller-Hinton agar; MHB, Mueller-Hinton broth; MIC, Minimum Inhibitory Concentration; MLST, Multi-locus sequence type; pEtN, phosphoethanolamine; PROVEAN, Protein Variation Effect Analyzer; ST, Sequence type; XDR, Extensively drug resistant; XDR-KP, Extensively drug-resistant *Klebsiella pneumoniae.*

REFERENCES

1. **Ramirez MS, Traglia GM, Lin DL, Tran T, Tolmasky ME.** Plasmid-Mediated Antibiotic Resistance and Virulence in Gram-Negatives: the *Klebsiella pneumoniae* Paradigm. *Microbiol Spectr* 2014;2:1-15.

2. **Patel G, Huprikar S, Factor SH, Jenkins SG, Calfee DP.** Outcomes of carbapenem-resistant *Klebsiella pneumoniae* infection and the impact of antimicrobial and adjunctive therapies. *Infect Control Hosp Epidemiol* 2008;29:1099-1106.

3. **Biswas S, Brunel JM, Dubus JC, Reynaud-Gaubert M, Rolain JM.** Colistin: an update on the antibiotic of the 21st century. *Expert Rev Anti Infect Ther* 2012;10:917-934.

4. **Suh JY, Son JS, Chung DR, Peck KR, Ko KS *et al*.** Nonclonal emergence of colistin-resistant *Klebsiella pneumoniae* isolates from blood samples in South Korea. *Antimicrob Agents Chemother* 2010;54:560-562.

5. **Kim SY, Shin J, Shin SY, Ko KS.** Characteristics of carbapenem-resistant *Enterobacteriaceae* isolates from Korea. *Diagn Microbiol Infect Dis* 2013;76:486-490.

6. **Mohamudha PR, Harish BN, Parija SC**. Emerging carbapenem resistance among nosocomial isolates of *Klebsiella pneumoniae* in South India. *Int J Pharma Bio Sci* 2010;1:1-11.

7. **Goel G, Hmar L, Sarkar De M, Bhattacharya S, Chandy M.** Colistin-resistant *Klebsiella pneumoniae*: report of a cluster of 24 cases from a new oncology center in eastern India. *Infect Control Hosp Epidemiol* 2014;35:1076-1077.

8.   **Cannatelli A, Giani T, D'Andrea MM, Di Pilato V, Arena F *et al*.** MgrB inactivation is

a common mechanism of colistin resistance in KPC-producing *Klebsiella pneumoniae* of

clinical origin. *Antimicrob Agents Chemother* 2014;58:5696-5703.

9.   **Mavroidi A, Katsiari M, Likousi S, Palla E, Roussou Z *et al*.** Characterization of ST258

Colistin-Resistant, blaKPC-Producing *Klebsiella pneumoniae* in a Greek Hospital. *Microb*

Drug Resist 2016;22:392-398.

10.   **Giamarellou H.** Epidemiology of infections caused by polymyxin-resistant pathogens. *Int*

J Antimicrob Agents 2016;48:614-621.

11.   **Monaco M, Giani T, Raffone M, Arena F, Garcia-Fernandez A *et al*.** Colistin resistance

superimposed to endemic carbapenem-resistant *Klebsiella pneumoniae*: a rapidly evolving

problem in Italy, November 2013 to April 2014. *Euro Surveill* 2014;19:pii=20939.

12.   **Pereira GH, Garcia DO, Mostardeiro M, Fanti KS, Levin AS.** Outbreak of carbapenem-

resistant *Klebsiella pneumoniae*: two-year epidemiologic follow-up in a tertiary hospital.

Mem Inst Oswaldo Cruz 2013;108:113-115.

13.   **Gaspar GG, Bellissimo-Rodrigues F, Andrade LN, Darini AL, Martinez R.** Induction

and nosocomial dissemination of carbapenem and polymyxin-resistant *Klebsiella*

pneumoniae. Rev Soc Bras Med Trop 2015;48:483-487.

14.   **Ah YM, Kim AJ, Lee JY.** Colistin resistance in *Klebsiella pneumoniae. Int J Antimicrob*

Agents 2014;44:8-15.

15.   **Groisman EA, Kayser J, Soncini FC.** Regulation of polymyxin resistance and adaptation

to low-Mg^{2+} environments. *J Bacteriol* 1997;179:7040-7045.

16.   **Velkov T, Thompson PE, Nation RL, Li J.** Structure-activity relationships of polymyxin

antibiotics. *J Med Chem* 2010;53:1898-1916.

17.   **Cheng HY, Chen YF, Peng HL.** Molecular characterization of the PhoPQ-PmrD-PmrAB

mediated pathway regulating polymyxin B resistance in *Klebsiella pneumoniae* CG43. *J*

Biomed Sci 2010;17:60.

18.   **Helander IM, Kato Y, Kilpelainen I, Kostiainen R, Lindner B *et al.*** Characterization

of lipopolysaccharides of polymyxin-resistant and polymyxin-sensitive *Klebsiella*

pneumoniae O3. *Eur J Biochem* 1996;237:272-278.

19.   **Velkov T, Deris ZZ, Huang JX, Azad MA, Butler M *et al.*** Surface changes and

polymyxin interactions with a resistant strain of *Klebsiella pneumoniae*. *Innate Immun*

2014;20:350-363.

20.   **Wright MS, Suzuki Y, Jones MB, Marshall SH, Rudin SD *et al.*** Genomic and

transcriptomic analyses of colistin-resistant clinical isolates of *Klebsiella pneumoniae*

reveal multiple pathways of resistance. *Antimicrob Agents Chemother* 2015;59:536-543.

21.   **Olaitan AO, Diene SM, Kempf M, Berrazeg M, Bakour S *et al.*** Worldwide emergence

of colistin resistance in *Klebsiella pneumoniae* from healthy humans and patients in Lao

PDR, Thailand, Israel, Nigeria and France owing to inactivation of the PhoP/PhoQ

regulator *mgrB*: an epidemiological and molecular study. *Int J Antimicrob Agents*

2014;44:500-507.

22.   **Arena F, De Angelis LH, Cannatelli A, Di Pilato V, Amorese M *et al.*** Colistin resistance

caused by inactivation of the MgrB regulator is not associated with decreased virulence of

ST258 KPC carbapenemase-producing *Klebsiella pneumoniae*. *Antimicrob Agents*

Chemother 2016;60:2509-2512.

23.   **Lee JY, Choi MJ, Choi HJ, Ko KS.** Preservation of Acquired Colistin Resistance in

Gram-Negative Bacteria. *Antimicrob Agents Chemother* 2015;60:609-612.

24.   **Meletis G, Tzampaz E, Sianou E, Tzavaras I, Sofianou D.** Colistin heteroresistance in

carbapenemase-producing *Klebsiella pneumoniae*. *J Antimicrob Chemother* 2011;**66**:946-

947.

25.   **Zowawi HM, Forde BM, Alfaresi M, Alzarouni A, Farahat Y *et al*.** Stepwise evolution

of pandrug-resistance in *Klebsiella pneumoniae*. *Sci Rep* 2015;5:15082.

26.   **Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R *et al*.** Emergence of plasmid-mediated

colistin resistance mechanism MCR-1 in animals and human beings in China: a

microbiological and molecular biological study. *Lancet Infect Dis* 2016;16:161-168.

27.   Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial*

Susceptibility Testing– Twenty-sixth Edition: Approved standard M100S. CLSI, Wayne,

PA, 2016.

28.   **Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME *et al*.** Multidrug-

resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert

proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect*

2012;18:268-281.

29.   **Bolger AM, Lohse M, Usadel B.** Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics* 2014;30:2114-2120.

30.   **Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M *et al*.** SPAdes: a new

genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*

2012;19:455-477.

31.   **Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ *et al*.** The SEED and the Rapid

Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids*

Res 2014;42:D206-214.

32.   **Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H *et al*.** Multilocus sequence

typing of total-genome-sequenced bacteria. *J Clin Microbiol* 2012;50:1355-1361.

33.   **Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S *et al*.** Identification

of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 2012;67:2640-2644.

34.   **Jünemann S, Sedlazeck FJ, Prior K, Albersmeier A, John U *et al*.** Updating benchtop

sequencing performance comparison. *Nat Biotechnol* 2013;31:294-296.

35.   **Liu P, Li P, Jiang X, Bi D, Xie Y, Tai C *et al*.** Complete genome sequence of *Klebsiella*

pneumoniae subsp. pneumoniae HS11286, a multidrug-resistant strain isolated from

human sputum *J Bacteriol* 2012; 194:1841-1842.

36.   **Deleo FR, Chen L, Porcella SF, Martens CA, Kobayashi SD, Porter AR *et al*.**

Molecular dissection of the evolution of carbapenemresistant multilocus sequence type 258

Klebsiella pneumoniae. *Proc Natl Acad Sci USA* 2014; 111:4988–4993.

37.   **Elliott AG, Ganesamoorthy D, Coin L, Cooper MA, Cao MD.** Complete Genome

Sequence of *Klebsiella quasipneumoniae subsp. similipneumoniae* Strain ATCC 700603.

Genome Announc 2016;4:e00438-16.

38.   **Li H, Durbin R.** Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics 2009;25:1754-1760.

39.   **Garrison E, Marth G.** Haplotype-based variant detection from short-read sequencing.

arXiv preprint arXiv 2012;12073907.

40.   **Cingolani P, Platts A, Wang le L, Coon M, Nguyen T *et al*.** A program for annotating

and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome

of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly (Austin)* 2012;6:80-92.

41.   **Choi Y, Chan AP.** PROVEAN web server: a tool to predict the functional effect of amino

498   acid substitutions and indels. *Bioinformatics* 2015;31:2745-2747.

499   42.   **Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M.** ISfinder: the reference

500   centre for bacterial insertion sequences. *Nucleic Acids Res* 2006;34:D32-36.

501   43.   **Jayol A, Poirel L, Brink A, Villegas MV, Yilmaz M A *et al*.** Resistance to colistin

502   associated with a single amino acid change in protein PmrB among *Klebsiella pneumoniae*

503   isolates of worldwide origin. *Antimicrob Agents Chemother* 2014;58:4762-4766.

504   44.   **Neonakis IK, Samonis G, Messaritakis H, Baritaki S, Georgiladakis A *et al*.** Resistance

505   status and evolution trends of *Klebsiella pneumoniae* isolates in a university hospital in

506   Greece: ineffectiveness of carbapenems and increasing resistance to colistin.

507   *Chemotherapy* 2010;56:448-452.

508   45.   **Poirel L, Jayol A, Bontron S, Villegas MV, Ozdamar M *et al*.** The *mgrB* gene as a key

509   target for acquired resistance to colistin in *Klebsiella pneumoniae*. *J Antimicrob Chemother*

510   2015;70:75-80.

511   46.   **Cannatelli A, D'Andrea MM, Giani T, Di Pilato V, Arena F *et al*.** *In vivo* emergence of

512   colistin resistance in *Klebsiella pneumoniae* producing KPC-type carbapenemases

513   mediated by insertional inactivation of the PhoQ/PhoP *mgrB* regulator. *Antimicrob Agents*

514   *Chemother* 2013;57:5521-5526.

515   47.   **Aires CAM, Pereira PS, Asensi MD, Carvalho-Assef AP.** *mgrB* mutations mediating

516   polymyxin B resistance in *Klebsiella pneumoniae* isolates from rectal surveillance swabs

517   in Brazil. *Antimicrob Agents Chemother* 2016;60:6969-6972.

48. **Kidd TJ, Mills G, Sá-Pessoa J, Dumigan A, Frank CG *et al.*** A *Klebsiella pneumoniae* antibiotic resistance mechanism that subdues host defences and promotes virulence. *EMBO Mol Med* 2017;9:430-447.

49. **Halaby T, Kucukkose E, Janssen AB, Rogers MR, Doorduijn DJ *et al*.** Genomic characterization of colistin heteroresistance in *Klebsiella pneumoniae* during a nosocomial outbreak. *Antimicrob Agents Chemother* 2016;60:6837-6843.

50. **Deleo FR, Chen L, Porcella SF, Martens CA, Kobayashi SD *et al*.** Molecular dissection of the evolution of carbapenem-resistant multilocus sequence type 258 *Klebsiella pneumoniae*. *Proc Natl Acad Sci U S A* 2014;111:4988-4993.

51. **Miller AK, Brannon MK, Stevens L, Johansen HK, Selgrade SE *et al*.** PhoQ mutations promote lipid A modification and polymyxin resistance of *Pseudomonas aeruginosa* found in colistin-treated cystic fibrosis patients. *Antimicrob. Agents Chemother* 2011;55:5761–5769.

52. **Lee JY, Ko KS.** Mutations and expression of PmrAB and PhoPQ related with colistin resistance in *Pseudomonas aeruginosa* clinical isolates. *Diagn. Microbiol. Infect. Dis*. 2014;78:271–276.

DATA BIBLIOGRAPHY

1. NCBI Bioproject PRJNA307517 (2016).

2. Liu P, Li P, Jiang X, Bi D, Xie Y, Tai C *et al*. Complete genome sequence of *Klebsiella pneumoniae subsp. pneumoniae* HS11286, a multidrug-resistant strain isolated from human sputum J Bacteriol 2012; 194:1841-1842. NCBI BioProject PRJNA78789.

3. Zowawi HM, Forde BM, Alfaresi M, Alzarouni A, Farahat Y *et al*. Stepwise evolution of pandrug-resistance in *Klebsiella pneumoniae*. *Sci Rep* 2015; 5:15082. European Nucleotide PRJEB7538.

4. Deleo FR, Chen L, Porcella SF, Martens CA, Kobayashi SD, Porter AR *et al*. Molecular dissection of the evolution of carbapenemresistant multilocus sequence type 258 *Klebsiella pneumoniae*. Proc Natl Acad Sci USA 2014; 111:4988–4993. European Nucleotide PRJNA237670.

FIGURES AND TABLES

Table 1. Broth microdilution and VITEK®2 antimicrobial testing for the 24 clinical isolates

Strain[*]	Source[†]	Resistance Profile[‡]																							
		1			2	3	4		5	6			7	8	9	10	11	12	13	14	15	16	17	
		AMK	GEN	TOB	CPT	TZP	IPM	MEM	CFZ	FEP	CTX	CAZ	FOX	CIP	SXT	TGC	ATM	AMP	SAM	CHL	FOF	CST	MIN	TET
1_GR_13	St	R	R	R	R^N	R	R	R	R^N	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
2_GR_12	U	R	R	R	R^N	R	R	R	R^N	R	R	R	R	R	R	I^R	R	R	R	R	R^S	R	R	R
3_GR_13	S	R	R	R	R^N	R	R	R	R^N	R^I	R	R	R	R	R	I	R	R	R	R	R	R^I	I^R	R
4_GR_12	B	R	R	R	R^N	R	R	R	R^N	R	R	R	R	R	R	I^R	R	R	R	R	R	R	I^R	R
5_GR_13	St	S	S	I	R^N	R	R	R	R^N	R	R	R	R	R	R	I^R	R	R	R	R	R	R	R	R
6_GR_12	St	S	S	I^R	R^N	R	R	R	R^N	R	R	R	R	R	R	R	R	R	R	R	R	S^R	I^R	R
7_GR_13	St	R	S	R	R^N	R	R	R	R^N	R	R	R	R	R	R	R	R	R	R	R	R	R	S^R	S^I
8_GR_13	St	R	R	R	R^N	R	R	R	R^N	R	R	R	R	R	R	R	R	R	R	R	R	R	R	R
9_GR_12	Br	I^R	S	R	R^N	R	R	R	R^N	R	R	R	R	R	R	I^R	R	R	R	R	R	R	I^R	S^I
10_GR_13	B	S^R	S	R	R^N	R	R	R	R^N	R	R	R	R	R	R	I	R	R	R	R	R	R	S^R	S^I
11_BR_13	U	S	S	R^N	R^N	R	R	R	R^N	R	R	R	R	R	R^N	I	R	R	R	S^N	R^N	R	S^N	S^N
12_BR_13	Br	S	R^S	I^N	R^N	R	R	R	R^N	R^I	R	R	R	R	R^N	I^R	R	R	R	R^N	R^N	R	I^N	S^N
13_GR_14	Br	I^R	S	R	R^N	R	R	R	R^N	R	R^N	R	R	R	R	S^R	R	R	R	R	R	R	S^I	S^I
14_GR_14	U	I^R	S^R	R	R^N	R	R	R	R^N	R	R^N	R	R	R	R	S^R	R	R	R	R	R	R	S^R	S^R
15_GR_13	St	I^R	S	R	R^N	R	R	R	R^N	R^I	R^N	R	R	R	R	I^R	R	R	R	R	R	R	S^I	S
16_GR_13	St	R	R	R	R^N	R	R	R	R^N	R	R^N	R	R	R	R	R^I	R	R	R	R	R	R	I^R	R
17_GR_14	St	R	R	R	R^N	R	R	R	R^N	R	R^N	R	R	R	R	I	R	R	R	R	R	R	S^R	R
18_GR_14	St	I^R	S	R	R^N	R	R	R	R^N	R	R	R	R	R	R	I	R	R	R	R	R	R	S^I	S
19_GR_14	St	I^R	S	R	R^N	R	R	R	R^N	R	R	R	R	R	R	R	R	R	R	R	R	R	I^R	I^R
20_GR_12	St	R	S	R	R^N	R	R	R	R^N	R^I	R	R	R	R	R	R	R	R	R	R	R	S	R	R
21_GR_13	U	S	S^R	I	R^N	R	R	I^R	R^N	R^I	R	R	R	S	R	S	S	R	R	S	S	S	S	S
22_GR_12	S	I^R	S	R	R^N	R	R	R	R^N	R^I	R	R	R	R	R	I	R	R	R	R	R^S	S	S^I	S
23_GR_12	St	R	R	R	R^N	R	R	R	R^N	R	R	R	R	R	R	R	R	R	R	R	R^S	R^S	R	R
24_GR_13	St	I^R	S	R	R^N	R	R	R	R^N	R^I	R	R	R	R	R	I^R	R	R	R	R	R	S	I^R	S

*Strain identification, numerical order catalogued at IMB_Country (GR:Greece, BR:Brazil)_last two digits of isolation year.

†Source represented as B, Blood; Br; Bronchial secretion; U, Urine; S, Sputum; St, Stool.

‡Antibiotic resistance as determined by broth microdilution according to CLSI guidelines (EUCAST for fosfomycin (disk diffusion) and

tigecycline) and in superscript, any discrepancies identified in VITEK®2 results. Antibiotic classes tested include **1**, Aminoglycosides (Amikacin,

AMK; Gentamicin, GEN; Tobramycin, TOB); **2**, Anti-MRSA cephalosporins (Ceftaroline, CPT); **3**, Antipseudomonal penicillins + β-lactamase

inhibitors (Piperacillin-tazobactam, TZP); **4**, Carbapenems (Imipenem, IPM; Meropenem, MEM); **5**, Non-extended spectrum cephalosporins (1st

and 2nd generation) (Cefazolin, CFZ); **6**, Extended-spectrum cephalosporins (3rd and 4th generation) (Cefepime, FEP; Cefotaxime, CTX,

Ceftazidime, CAZ); **7**, Cephamycins (Cefoxitin, FOX); **8**, Fluoroquinolones (Ciprofloxacin, CIP); **9**, Folate pathway inhibitors (Trimethoprim-

sulfamethoxazole, SXT); **10**, Glycylcyclines (Tigecycline, TGC); **11**, Monobactams (Aztreonam, ATM); **12**, Penicillins (Ampicillin, AMP); **13**,

Penicillins + β-lactamase inhibitors (Amipicillin-sulbactam, SAM); **14**, Phenicols (Chloramphenicol, CHL); **15**, Phosphonic acids (Fosfomycin,

FOF); **16**, Polymyxins (Colistin, CST); **17**, Tetracyclines (Minocycline, MIN; Tetracycline, TET).

R, Resistant; I, Intermediate; S, Susceptible; N, Not tested.

Table 2. Potential mutations contributing to polymyxin resistance and acquired resistance genes

*Multilocus sequence type as identified through MultiLocus Sequence Typing Server 1.8.

†Variations detected in *mgrB*, *phoPQ* and *pmrAB* potentially causing polymyxin resistance (PMX-R). Significant non-synonymous changes

determined by PROVEAN analysis. WT (Wild-type) alleles in comparison to 20_GR_12. Displayed as gene impacted, initial amino acid, position

and new amino acid. If - shown in front of position, alteration is encoded upstream and if - is only displayed, no significant non-synonymous

changes were detected in these loci. Insertion sequences classified as Δ, identity as per ISFinder and orientation in superscript. Orientation

determined as forward, [F], if transposase is in the same direction as *mgrB* and conversely, reverse, [R], if in the opposite direction to *mgrB*.

‡Acquired antibiotic resistance genes detected via ResFinder 2.1. Classes of antibiotics impacted displayed as A, Aminoglycoside; B, Beta-lactam;

F; Fosfomycin; M, Macrolide; P, Phenicol; Q, Quinolone; R, Rifampicin; S, Sulphonamide; T, Tetracycline; Tr, Trimethoprim. Shading indicates

detection of gene (≥90% homology, ≥60% sequence length).

Fig. 1. Neighbor-joining tree of core genome MLST of 24 *Klebsiella* clinical isolates. Clustering of

sequence type (ST) indicated at base of diverging branch. cgMLST compared to completed assemblies

including ATCC 700603 (*K. quasipneumoniae*), HS11286 (ST11), MS6671 (ST147) and NJST258_1

and NJST258_2 (ST258).

Fig. 2. Complementation assays and influence of gene on polymyxin resistance. Polymyxin B MIC measured before (○) and after (■) complementation of wild-type gene (a) pTOPO-*mgrB*, (b) pTOPO-*phoP*, (c) pTOPO-*phoQ*, or (d) pTOPO-*pmrB* in indicated resistant isolates. (e) Mutated genes complemented into 20_GR_12 (polymyxin-susceptible isolate) to determine if variant induces polymyxin resistance. (f) Complementation of 9_GR_13 (IS element disrupted *mgrB* control) to detect potential suppressor mutations. Strains shown on x axis for (a-d) and superscript indicates variants in genes including *mgrB* (a), *phoP* (b), *phoQ* (c) and *pmrB* (d) that differ from 20_GR_12. For (e, f), the x axis shows the gene complemented with amino acid variation in brackets. Dotted line at 2 mg/L represents the breakpoint for polymyxin B. Values indicate mean±standard deviation where no error bars display no fluctuation in MIC (n≥2 colonies).