Concomitant duplication of the *Cid* and *Cenp-C* genes in *Drosophila*

José R. Teixeira[1], Guilherme B. Dias[1], Marta Svartman[1], Alfredo Ruiz[2], Gustavo C. S. Kuhn[1]

[1]Departamento de Biologia Geral, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil, Postal Code: 31270-901

[2]Departament de Genètica i de Microbiologia, Universitat Autònoma de Barcelona, Bellaterra (Barcelona), Spain, Postal Code: 08193

Corresponding author:

Prof. Dr. Gustavo C. S. Kuhn

Laboratório de Citogenômica Evolutiva, Departamento de Biologia Geral, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Av. Antônio Carlos 6627 – Pampulha, Belo Horizonte, MG, Brazil. Postal Code: 31270-901.

Fone: +55 (31) 3409-3062

E-mail: gcskuhn@ufmg.br

Running title: *Cid* and *Cenp-C* duplicates in *Drosophila*

Abstract

Despite their essential role in the process of chromosome segregation in eukaryotes, kinetochore

proteins are highly diverse across species, being lost, duplicated, invented, or diversified on the

sequence level. Based on comparative genomics, the duplication of the inner kinetochore proteins

CenH3 and Cenp-C, which are interdependent in their roles of stablishing centromere identity and

function, can be said to be rare in animals. Surprisingly, the *Drosophila CenH3* homolog *Cid*

underwent four independent duplication events during evolution. Particularly interesting are the

highly diverged and subfunctionalized *Cid1* and *Cid5* paralogs of the *Drosophila* subgenus, which

reveal that over one thousand *Drosophila* species encode two *Cid* genes, making those with a single

copy a minority. Given that CenH3 and Cenp-C likely co-evolve as a functional unit, we

investigated the molecular evolution of *Cenp-C* in species of *Drosophila*. We report yet another *Cid*

duplication within the *Drosophila* subgenus and that not only *Cid*, but also *Cenp-C* is duplicated in

the entire *Drosophila* subgenus. The *Cenp-C* paralogs, which we named *Cenp-C1* and *Cenp-C2*, are

highly divergent. The retention of key motifs involved in centromere localization by both Cenp-C1

and Cenp-C2 makes neofunctionalization unlikely. In contrast, the alternate conservation of some

functional motifs between the proteins is highly indicative of subfunctionalization. Interestingly,

both *Cid5* and *Cenp-C2* are male germline-biased and evolved adaptively. Our findings point

towards a specific inner kinetochore composition to a specific context (i.e., spermatogenesis), which

could prove valuable for the understanding of how the extensive kinetochore diversity is related to

essential cellular functions.

Keywords: CenH3, Cenp-C, gene duplication, centromere, kinetochore, Drosophila

Introduction

During eukaryotic cell division, accurate chromosome segregation requires the interaction of chromosomes with the microtubules from the spindle apparatus. This interaction is mediated by the kinetochore, a multiprotein structure that is hierarchically assembled on the centromeres. Upstream in the assembly of the kinetochore are CenH3 and Cenp-C, two proteins interdependent in their roles of establishing centromere identity and function (Erhardt *et al*. 2008; Orr and Sunkel 2011). Both CenH3 and Cenp-C have similar phylogenetic profiles (i.e., they have similar patterns of presences and absences across eukaryote evolution) and likely co-evolve as a functional unit (van Hooff *et al*. 2017). However, despite the essentiality of centromeres, centromeric DNA and proteins are remarkably diverse (Henikoff *et al*. 2000; Talbert *et al*. 2004; Plohl *et al*. 2008). This rapid evolution despite the expectation of constraint is referred to as the "centromere paradox" (Henikoff *et al*. 2001). This paradox can be explained by the centromere drive hypothesis, which proposes that genetic conflicts during female meiosis drive centromere evolution (Henikoff *et al*. 2001; Dawe and Henikoff 2006).

Female meiosis is asymmetric in animals and plants, where only one of the four meiotic products is viable. If a pair of homologous chromosomes have kinetochores that interact unequally with the spindle fibers, then there is potential for a biased inheritance (Ross and Malik 2014). The expansion of centromeric DNA sequences leads to the recruitment of more kinetochore proteins and, subsequently, to the recruitment of more spindle fibers, which confers an advantage to the homolog with the expanded centromere (Iwata-Otsubo *et al*. 2017). However, the spread of expanding centromeres throughout a population might be accompanied by deleterious effects, such as increased male sterility or a skewed sex ratio (Fishman and Saunders 2008; Rutkowska and Badyaev 2008; Malik and Henikoff 2009). Changes in CenH3 and Cenp-C related to more 'flexible' DNA-binding preferences are expected to counteract the transmission advantage gained by the homolog with the expanded centromere, thus restoring meiotic parity for both homologs (Henikoff et al. 2001; Dawe and Henikoff 2006).

The kinetochore is highly diverse across species, with proteins being lost, duplicated, invented, or diversified on the sequence level (van Hooff *et al*. 2017). Given that data directly supporting a correlation between the evolution of centromeric DNA sequences and CenH3 and Cenp-C are still absent, it is not known if and how such structural divergence is related to centromere drive suppression. However, the subfunctionalization of *CenH3* duplicates in some lineages of *Drosophila* have been hypothesized to be linked to centromere drive suppression (Kursel and Malik 2017). The *Drosophila CenH3* homolog *Cid* underwent four independent duplication events during evolution, and some *Cid* duplicates are primarily expressed in the male germline and evolve under

positive selection (Kursel and Malik 2017). Such duplications could have allowed for the rapid

evolution of centromeric proteins without compromising their essential function by separating

functions with divergent fitness optima.

The existence of *CenH3* duplicates in genetically tractable organisms provides an opportunity

to study the functional consequences of genetic variations for kinetochore-related processes without

interfering with essential mitotic functions. Thus, given the interdependence between CenH3 and

Cenp-C, we decided to further analyze the molecular evolution of the *Cid* and *Cenp-C* genes in

Drosophila species. Here, we report yet another *Cid* duplication within the *Drosophila* subgenus

and that not only *Cid*, but also *Cenp-C* is duplicated in the entire *Drosophila* subgenus. The *Cid* and

Cenp-C paralogs likely subfunctionalized, as some motifs are alternatively conserved between the

paralogs. Interestingly, both the *Cid* and *Cenp-C* duplications generated copies that are male-biased

and evolved under positive selection. Our findings point towards a specific kinetochore composition

to a specific context (i.e., the male germline), which could prove valuable for the understanding of

how the extensive kinetochore diversity is related to essential cellular functions.

Results and Discussion

***Cid1* was replaced by a new paralog in a clade within the *Drosophila* subgenus**

Duplicate *Cid* genes exist in *D. eugracilis* (*Cid1*, *Cid2*) and in the *montium* subgroup (*Cid1*,

Cid3, *Cid4*), both within the *Sophophora* subgenus, and in the entire *Drosophila* subgenus (*Cid1*,

Cid5). In all analyzed species from the *Drosophila* subgenus, *Cid1* is flanked by the *cbc* and *bbc*

genes, and *Cid5* is flanked by the *Kr* and *CG6907* genes (Kursel and Malik 2017). As expected, we

found two *Cid* genes while looking for the orthologs of *Cid1* and *Cid5* in the assembled genomes of

two cactophilic species from the *Drosophila* subgenus, *D. buzzatii* and *D. seriema* (*repleta* group).

Surprisingly, while one of the genes is present in the expected locus of *Cid5*, the other one is in an

entirely different locus, flanked by the *CG14341* and *IntS14* genes. We named this new paralog

Cid6.

By investigating the *Cid1* locus of *D. buzzatii*, we found a myriad of transposable elements

(TEs) surrounding a 116-bp fragment of the original *Cid1* gene (fig. 1, upper panel). Due to

fragmentary genome assembly, the *Cid1* locus of *D. seriema* could not be identified. *Cid6* and *Cid5*

of either *D. buzzatii* or *D. seriema* share ~40% amino acid identity but, in contrast, *Cid6* of either

species and *Cid1* of the closely related *D. mojavensis* are much more similar, sharing ~80%

identity. Fluorescent *in situ* hybridizations on polytene chromosomes showed that *Cid6* is distal (in

relation to the chromocenter) in the Muller element B of *D. buzzatii* and *D. seriema*, and that *Cid1*

is proximal in the Muller element C of *D. mojavensis* and the outgroup *D. virilis* (fig. 1, lower
panel). Therefore, we inferred that *Cid1* was degenerated by several TE insertions after the origin of
Cid6 by an inter-chromosomal duplication of *Cid1* in the lineage that gave rise to *D. buzzatii* and *D.*
seriema. The time of divergence between *D. buzzatii* and *D. seriema* has been estimated at ~4.6
mya, and the divergence between them and the closely related *D. mojavensis* has been estimated at
~11.3 mya (Oliveira *et al*. 2012). Accordingly, the *Cid1* duplication happened between ~4.6 and
11.3 mya.

Why *Cid6* persevered while *Cid1* degenerated? The *Cid1* locus of *D. buzzatii* is located in the
most proximal region of the Muller element C (scaffold 115; Guillén *et al*. 2014), which is very
close to the pericentromeric heterochromatin, where TEs are highly abundant (Pimpinelli *et al*.
1995; Casals *et al*. 2005; Rius *et al*. 2016). Natural selection is known to be less effective in
pericentromeric and adjacent regions due to low rates of crossing-over (Zhang and Kishino 2004;
Clément *et al*. 2006; Comeron *et al*. 2012; Nambiar and Smith 2016). Thus, it is reasonable to
suggest that the presence of *Cid6* in Muller element B alleviated the selective pressures over *Cid1* in
Muller element C, whose proximity to the pericentromeric heterochromatin fostered its degradation
by several posterior TE insertions.

Cenp-C is duplicated in the entire *Drosophila* subgenus

Given that CenH3 and Cenp-C are interdependent and co-evolve as a function unit, we
investigated if *Cenp-C* was also duplicated in species of the lineages where *Cid* was found
duplicated. *D. eugracilis*, species from the *montium* subgroup, and all the other species of the
Sophophora subgenus have only one copy of *Cenp-C*, which is always flanked by the *5-HT2B* gene.
Interestingly, the species of the *Drosophila* subgenus have two copies of *Cenp-C* with ~55%
nucleotide identity, which we named *Cenp-C1* and *Cenp-C2*: the former is flanked by the *5-HT2B*
and *CG1427* genes, and the latter is flanked by the *CLS* and *RpL27* genes. No additional copies of
Cenp-C were found in *D. buzzatii* and *D. seriema*. A maximum likelihood tree shows that *Cenp-C*
was likely duplicated after the split between the *Sophophora* and *Drosophila* subgenera but before
the split between *D. busckii* and the other species of the *Drosophila* subgenus (fig. 2). Thus, we
conclude that *Cenp-C2* originated from a duplication of *Cenp-C1* in the lineage that gave rise to
species of the *Drosophila* subgenus, at least 50 mya (Russo *et al*. 2013).

Why *Cenp-C* is duplicated only in the *Drosophila* subgenus if *Cid* is also duplicated in *D.*
eugracilis and the *montium* subgroup? Interestingly, only one Cid encoding gene was found in *D.*
busckii, meaning either that *Cid* and *Cenp-C* were concomitantly duplicated and that *Cid5* was
subsequently lost in *D. busckii* (Kursel and Malik 2017), or that *Cid1* was duplicated after the

divergence of *D. busckii*. It is likely that the *Cid* and *Cenp-C* duplications in the *Drosophila*

subgenus were concomitant but not correlated. Nevertheless, it seems that the *Cenp-C* duplication

had an effect on the evolution of the *Cid1* and *Cid5* paralogs.

As a histone H3 variant, CenH3 has the C-terminal histone fold domain, which is reasonably

conserved among species, and the N-terminal tail, which is highly variable among species

(Henikoff *et al*. 2000). In *D. eugracilis*, the *Cid2* paralog functionally replaced the pseudogenized

ancestral *Cid1* paralog (Kursel and Malik 2017). In the *montium* subgroup, all paralogs are

functional but highly diverged: they share 45-80% amino acid identity at their histone fold domains

and ~25% at their N-terminal tails (Kursel and Malik 2017). The *Cid* paralogs of *Drosophila*

subgenus species are also both functional: they share ~80% amino acid identity at their histone fold

domains but only ~15% at their N-terminal tails (Kursel and Malik 2017).

The conserved amino acid identity of the N-terminal tail is represented by the fact that it

evolves in a modular manner, with four core motifs always conserved when there is only one Cid

protein encoded in the genome (Kursel and Malik 2017). In species of the *montium* subgroup, these

four motifs are shared between the paralogs. In contrast, in species of the *Drosophila* subgenus, all

four motifs are conserved in Cid1 but only 1-2 are conserved in Cid5 (Kursel and Malik 2017).

Therefore, we propose that if the N-terminal tail of Cid interacts with Cenp-C, a new *Cenp-C* copy

would allow a higher divergence of the *Cid* paralogs by alleviating the selective pressures over the

Cid/Cenp-C interaction. Further studies focusing on the specific interactions between Cid and

Cenp-C could be valuable for the comprehension of why these two proteins can be so evolutionary

flexible.

Some Cenp-C motifs are alternatively conserved between Cenp-C1 and Cenp-C2

Cenp-C was previously thought to be absent in *Drosophila* (Talbert *et al*. 2004), but a protein

that interacts with the regulatory subunits of separase turned out to be a highly diverged Cenp-C

homolog (Heeger *et al*. 2005). The *D. melanogaster* Cenp-C1, as was characterized by Heeger *et al*.

(2005), has seven independent functional motifs, from N- to C-terminal (fig. 3A): arginine-rich (R-

rich), drosophilids Cenp-C homology (DH), AT hook 1 (AT1), nuclear localization signal (NLS),

CenH3 binding (also known as the Cenp-C motif), AT hook 2 (AT2), and C-terminal dimerization

(Cupin). The R-rich and DH motifs, as well as both AT1 and AT2 motifs (which could mediate

binding to the minor grove of DNA), are functionally poorly characterized. However, all except

AT1 appear to provide essential functions, as Cenp-C1 versions lacking these regions were unable

to prevent phenotypic abnormalities in *Cenp-C1* mutant embryos (Heeger *et al*. 2005). Furthermore,

present in the CenH3 binding motif is the arginine 1101 (R1101), which is crucial for centromere

localization (Heeger *et al*. 2005). Given the functional relevance of these motifs, we searched for
them in both Cenp-C1 and Cenp-C2.

With the exception of *D. kikkawai* (from the *montium* subgroup), in which the AT2 motif is
absent, all seven motifs are conserved in Cenp-C1 of all other species of the *Sophophora* subgenus
(data not shown). In contrast, the motifs are alternatively conserved between Cenp-C1 and Cenp-C2
of species of the *Drosophila* subgenus (fig. 3B). Both Cenp-C1 and Cenp-C2 of all species have the
DH, NLS, and CenH3 binding motifs, the corresponding R1101 of *D. melanogaster*, and lack the
AT1 motif. However, only Cenp-C2 has the R-rich and AT2 motifs conserved. Furthermore, both
Cenp-C1 and Cenp-C2 of most species have the Cupin motif, the exceptions being Cenp-C1 of *D.*
busckii, which lacks the final half of it, and Cenp-C2 of *D. grimshawi*, which lacks it entirely.
Interestingly, with the exception of the CenH3 binding and Cupin motifs, almost all motifs of Cenp-
C2 are more similar to those of Cenp-C1 of the *Sophophora* subgenus than to those of Cenp-C1 of
the *Drosophila* subgenus (table 1).

Given that the motifs responsible for centromere localization are conserved in both Cenp-C1
and Cenp-C2 (i.e., the NLS and CenH3 binding motifs, as well as the corresponding R1101 of *D.*
melanogaster), it is unlikely that any of the paralogs underwent neofunctionalization. The
(parcial)loss of the Cupin motif in *D. busckii* and *D. grimshawi* points towards subfunctionalization.
Given that its function is unknown, it is difficult to comprehend the loss of the AT1 motif in both
Cenp-C1 and Cenp-C2 of the *Drosophila* subgenus. However, even though that their specific
functions are also unknown, the loss of the R-rich and AT2 motifs in Cenp-C1, and their retention
in Cenp-C2 are highly indicative of subfunctionalization.

The *Cenp-C* paralogs are differentially expressed

Cid and Cenp-C are fundamentally interdependent because the centromeric localization of one
depends on the centromeric localization of the other (Erhardt *et al*. 2008; Orr and Sunkel 2011).
Given that Cenp-C is incorporated into centromeres concomitantly with Cid (Schuh *et al*. 2007),
and that the excess of both proteins can cause centromere expansion and kinetochore failure
(Schittenhelm *et al*. 2010), the expression of both proteins needs to be tightly regulated.

Kursel and Malik (2017) showed that *Cid5* expression is male germline-biased and proposed
that *Cid1* and *Cid5* subfuncionalized and now performed nonredundant centromeric roles. In order
to investigate if *Cenp-C1* and *Cenp-C2* are differentially expressed and correlated in some way with
the expression of the *Cid* paralogs, we analyzed the available transcriptomes from embryos, larvae,
pupae and adult females and males of *D. buzzatii* (Guillén *et al*. 2014), and from testes of *D. virilis*
and *D. americana* (BioProject Accession PRJNA376405).

While *Cid6* is transcribed in all stages of development of *D. buzzatii*, which confirms that *Cid6* functionally replaced *Cid1*, *Cid5* transcription is limited to pupae and adult males, being higher than *Cid6* transcription in the latter (fig. 4A). Also, *Cid5* transcription is elevated in testes of *D. virilis* and *D. americana*, whereas *Cid1* is virtually silent (fig. 4C). Our results further support the finding of Kursel and Malik (2017) that *Cid5* displays a male germline-biased expression. In this context, our finding that *Cid5* is also transcribed in pupae of *D. buzzatii* may be related to the ongoing development of the male gonads.

In contrast to the *Cid* paralogs, we found that both *Cenp-C1* and *Cenp-C2* are transcribed in almost all stages of development of *D buzzatii*, with the exception of larvae (fig. 4B). *Cenp-C1* transcription is higher than *Cenp-C2* transcription in embryos and adult females of *D. buzzatii*. On the other hand, *Cenp-C2* transcription is higher than *Cenp-C1* transcription in pupae and adult males of *D. buzzatii*. Also, *Cenp-C2* transcription is higher than *Cenp-C1* transcription in testes of *D. virilis*, but there is no significant difference between their expression in testes of *D. americana* (fig. 4D). Therefore, similar to what was found for the *Cid* paralogs, the differential expression between the *Cenp-C* paralogs support the subfunctionalization hypothesis. The male germline-biased expression of both *Cid5* and *Cenp-C2* points towards their interaction in spermatogenesis, but biochemical assays need to be performed to confirm some sort of interdependence.

Both *Cid5* and *Cenp-C2* evolve under positive selection in species of the *repleta* group

Centromeres are essential for the faithful segregation of chromosomes in cell divisions, yet the centromeric DNA sequences and both CenH3 and Cenp-C are highly variable across species (Henikoff *et al.* 2000; Talbert *et al.* 2004; Plohl *et al.* 2008). This paradox can be explained by the centromere drive hypothesis, which states that CenH3 and Cenp-C constantly evolve and acquire new binding preferences to rapidly evolving centromeric DNA sequences in an effort to suppress their selfishly spread throughout the population by female meiotic drive (Henikoff *et al.* 2001; Dawe and Henikoff 2006). However, the rapid evolution of CenH3 required for the "drive suppressor" function may be disadvantageous for canonical functions (e.g., mitosis; Finseth *et al.* 2015; Kursel and Malik 2017).

The possibility that the paralogs achieved fitness optima for divergent functions predicts that selection may act differently in each of the *Cid* and *Cenp-C* paralogs. To test this hypothesis, full-length alignments of the *Cid* and *Cenp-C* paralogs were analyzed for positive selection using maximum likelihood methods. We focused our analyses on five closely related cactophilic *Drosophila* species from the *repleta* group (*D. mojavensis*, *D. arizonae*, *D. navojoa*, *D. buzzatii* and *D. seriema*). Interestingly, both *Cid5* and *Cenp-C2* showed signs of positive selection (table 2).

Bayes Empirical Bayes analyses identified with a posterior probability > 95% four amino acids in
the N-terminal tail of Cid5 and six amino acids across Cenp-C2 as having evolved under positive
selection. Of the six Cenp-C2 amino acids, one is in the DH motif, one is in the Cupin motif, and
the remaining four are in inter-motif sequences.

Kursel and Malik (2017) found signs of positive selection in the *Cid3* paralog of the *montium*
subgroup and proposed that *Cid3* and *Cid5* are candidate suppressors of centromere drive given
their male germline-biased expression. Our results of positive selection on both *Cid5* and *Cenp-C2*
do support this hypothesis. However, male germline-biased genes are widely known to evolve
adaptively as the result of male-male or male-female competition (Ellegren and Parsch 2007;
Meisel 2011). Either way, molecular genetic data alone cannot reveal the underlying cause of
adaptive evolution. What our findings do suggest is that species of the *Drosophila* subgenus likely
have a specific inner kinetochore composition that seems to mainly function in spermatogenesis.
246

247 **Concluding remarks**
248

The extensive diversity of kinetochore compositions in eukaryotes poses numerous questions
regarding flexibility of essential cellular functions (van Hooff *et al*. 2017). Is the kinetochore less
conserved than other core eukaryotic cellular systems? And if so, why is it allowed to be less
conserved? Are many of the alterations adaptive to the species? To answer such questions, how a
specific kinetochore composition affect specific cellular features and lifestyles should be
investigated. Here, we show that Cid5 and Cenp-C2 offer such possibility, as both are inner
kinetochore protein variants likely specialized to function mainly in spermatogenesis. Thus, finding
out if and how Cid5 and Cenp-C2 play a role either in centromere drive suppression or reproductive
competition can shine new light into our understanding of centromere evolution.
258

259 **Materials and Methods**
260

261 **Identification of *Cid* and *Cenp-C* orthologs and paralogs in sequenced genomes**
262

Drosophila *Cid* and *Cenp-C* genes were identified by tBLASTx in sequenced genomes using
the *D. melanogaster Cid1* and *Cenp-C1* as queries (FlyBase IDs FBgn0040477 and FBgn0266916,
respectively). For the *D. seriema* genome assembly see Supplementary File S1. Since Cid is
encoded by a single exon in *Drosophila*, we selected the entire open reading frame for each *Cid*
gene hit, and since *Cenp-C* has multiple introns, we used the Augustus gene prediction algorithm
(Stanke and Morgenstern 2005) to identify the coding DNA sequences. For annotated genomes, we

recorded the 5' and 3' flanking genes for the *Cid* and *Cenp-C* genes of each species. For genomes
that are not annotated, we used the 5' and 3' nucleotide sequences flanking the *Cid* and *Cenp-C*
genes as queries to the *D. melanogaster* genome using BLASTn and verified the synteny in
accordance to the hits. All *Cid* and *Cenp-C* coding sequences and their database IDs can be found in
Supplementary Files S2 and S3, respectively.

Fluorescent *in situ* hybridizations (FISH) on polytene chromosomes

Probes for *Cid1*/*Cid6* were obtained by PCR from genomic DNA of *D. buzzatii* (strain st-1), *D.*
seriema (strain D73C3B), *D. mojavensis* (strain 14021-0248.25) and *D. virilis* (strain 15010-
1551.51). We cloned the PCR products into the pGEM-T vector (Promega) and sequenced them to
confirm identity. Recombinant plasmids were labeled with digoxigenin 11-dUTP by nick
translation (Roche Applied Science). FISH on polytene chromosomes was performed as described
in Dias *et al*. (2015). The slides were analyzed under an Axio Imager A2 epifluorescence
microscope equipped with the AxioCam MRm camera (Zeiss). Images were captured with the
AxioVision software (Zeiss) and edited in Adobe Photoshop. Chromosome arms were identified by
their morphology (Kuhn *et al*. 1996; González *et al*. 2005; Schaeffer *et al*. 2008).

Phylogenetic analyses

Cid and *Cenp-C* sequences were aligned at the codon level using MUSCLE (Edgar 2004) and
refined manually. Using the conserved motifs of Cenp-C (Heeger *et al*. 2005), we generated
maximum likelihood phylogenetic trees in MEGA6 (Tamura *et al*. 2013) with the GTR substitution
model and 1,000 bootstrap replicates for statistical support.

Expression analyses

RNA-seq data from *D. buzzatii* (Guillén *et al*. 2014), and from *D. virilis* and *D. americana*
(BioProject Accession PRJNA376405) were analyzed for the *Cid* and *Cenp-C* expression patterns
with Bowtie2 (Langmead and Salzberg 2012), as implemented to the Galaxy server (Afgan *et al*.
2016). Mapped reads were normalized by the transcripts per million (TPM) method (Wagner *et al*.
2012), and all normalizes values < 1 were set to 1 so that $\log_2$ TPM ≥ 0.

Positive selection analyses

Cid and *Cenp-C* alignments and gene trees were used as input into the CodeML NSsites models

of PAMLX version 1.3.1 (Xu and Yang 2013). To determine whether each paralog evolved under

positive selection, we compared three models that do not allow dN/dS to exceed 1 (M1a, M7 and

M8a) to two models that allow dN/dS > 1 (M2a and M8). Positively selected sites were classified as

those with a M8 Bayes Empirical Bayes posterior probability > 95%.

Acknowledgments

We are grateful to Dr. Maura Helena Manfrin (Univesity of São Paulo) for providing us with

the *D. seriema* strain used in the present work. This work was supported by a grant from "Fundação

de Amparo à Pesquisa do Estado de Minas Gerais" (FAPEMIG) to G.K. (grant number APQ-

01563-14).

References

Afgan E, Baker D, van den Beek M, Blankenberg D, Bouvier D, Čech M, Chilton J, Clements D, Coraor N, Eberhard C, et al. 2016. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2016 update. Nucleic Acids Res. 44:W3-W10.

Casals F, Cáceres M, Manfrin M, González J, Ruiz A. 2005. Molecular Characterization and Chromosomal Distribution of Galileo, Kepler and Newton, Three Foldback Transposable Elements of the Drosophila buzzatii Species Complex. Genetics 169:2047-2059.

Clément Y, Tavares R, Marais G. 2006. Does lack of recombination enhance asymmetric evolution among duplicate genes? Insights from the Drosophila melanogaster genome. Gene 385:89-95.

Comeron J, Ratnappan R, Bailin S. 2012. The Many Landscapes of Recombination in Drosophila melanogaster. PLoS Genet. 8:e1002905.

Dawe R, Henikoff S. 2006. Centromeres put epigenetics in the driver's seat. Trends Biochem Sci. 31:662-669.

Dias G, Heringer P, Svartman M, Kuhn G. 2015. Helitrons shaping the genomic architecture of Drosophila: enrichment of DINE-TR1 in α- and β-heterochromatin, satellite DNA emergence, and piRNA expression. Chromosome Res. 23:597-613.

Edgar R. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 32:1792-1797.

Erhardt S, Mellone B, Betts C, Zhang W, Karpen G, Straight A. 2008. Genome-wide analysis reveals a cell cycle–dependent mechanism controlling centromere propagation. J Cell Biol. 183:805-818.

Finseth F, Dong Y, Saunders A, Fishman L. 2015. Duplication and Adaptive Evolution of a Key Centromeric Protein in Mimulus, a Genus with Female Meiotic Drive. Mol Biol Evol. 32:2694-2706.

Fishman L, Saunders A. 2008. Centromere-Associated Female Meiotic Drive Entails Male Fitness Costs in Monkeyflowers. Science 322:1559-1562.

Gonzalez J, Nefedov M, Bosdet I, Casals F, Calvete O, Delprat A, Shin H, Chiu R, Mathewson C, Wye N, et al. 2005. A BAC-based physical map of the Drosophila buzzatii genome. Genome Res. 15:885-889.

Guillén Y, Rius N, Delprat A, Williford A, Muyas F, Puig M, Casillas S, Ràmia M, Egea R, Negre B, et al. 2014. Genomics of Ecological Adaptation in Cactophilic Drosophila. Genome Biol Evol. 7:349-366.

Heeger S, Leismann O, Schittenhelm R, Schraidt O, Heidmann S, Lehner C. 2005. Genetic interactions of separase regulatory subunits reveal the diverged Drosophila Cenp-C homolog. Genes Dev. 19:2041-2053.

Henikoff S, Ahmad K, Malik H. 2001. The Centromere Paradox: Stable Inheritance with Rapidly Evolving DNA. Science 293:1098-1102.

Henikoff S, Ahmad K, Platero J, van Steensel B. 2000. Heterochromatic deposition of centromeric histone H3-like proteins. Proc Natl Acad Sci USA 97:716-721.

Iwata-Otsubo A, Dawicki-McKenna JM, Akera T, Falk SJ, Chmátal L, Yang K, Sullivan BA, Schultz RM, Lampson MA, Black BE. 2017. Expanded Satellite Repeats Amplify a Discrete CENP-A Nucleosome Assembly Site on Chromosomes that Drive in Female Meiosis. Curr Biology doi:10.1016/j.cub.2017.06.069

Kohany O, Gentles A, Hankus L, Jurka J. 2006. Annotation, submission and screening of repetitive elements in Repbase: RepbaseSubmitter and Censor. BMC Bioinformatics 7:474.

Kuhn G, Ruiz A, Alves M, Sene F. 1996. The metaphase and polytene chromosomes of Drosophila seriema (repleta group; mulleri subgroup). Brazilian Journal of Genetics 19:209–216.

Kursel L, Malik H. 2017. Recurrent Gene Duplication Leads to Diverse Repertoires of Centromeric Histones in Drosophila Species. Mol Biol Evol. doi: 10.1093/molbev/msx091.

Langmead B, Salzberg S. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9:357-359.

Malik H, Henikoff S. 2009. Major Evolutionary Transitions in Centromere Complexity. Cell 138:1067-1082.

Nambiar M, Smith G. 2016. Repression of harmful meiotic recombination in centromeric regions. Semin Cell Dev Biol. 54:188-197.

Oliveira D, Almeida F, O'Grady P, Armella M, DeSalle R, Etges W. 2012. Monophyly, divergence times, and evolution of host plant use inferred from a revised phylogeny of the Drosophila repleta species group. Mol Phylogenet Evol. 64:533-544.

Orr B, Sunkel C. 2011. *Drosophila* CENP-C is essential for centromere identity. Chromosoma 120:83-96.

Pimpinelli S, Berloco M, Fanti L, Dimitri P, Bonaccorsi s, Marchetti E, Caizzi R, Caggese C, Gatti M. 1995. Transposable elements are stable structural components of Drosophila melanogaster heterochromatin. Proc Natl Acad Sci USA 92:3804-3808.

Plohl M, Luchetti A, Meštrović N, Mantovani B. 2008. Satellite DNAs between selfishness and functionality: Structure, genomics and evolution of tandem repeats in centromeric (hetero)chromatin. Gene 409:72-82.

Rius N, Guillén Y, Delprat A, Kapusta A, Feschotte C, Ruiz A. 2016. Exploration of the Drosophila buzzatii transposable element content suggests underestimation of repeats in Drosophila genomes. BMC Genomics 10:17-344.

Ross B, Malik H. 2014. Genetic Conflicts: Stronger Centromeres Win Tug-of-War in Female Meiosis. Current Biology 24:R966-R968.

Russo C, Mello B, Frazão A, Voloch C. 2013. Phylogenetic analysis and a time tree for a large drosophilid data set (Diptera: Drosophilidae). Zoological Journal of the Linnean Society 169:765-775.

Rutkowska J, Badyaev A. 2008. Meiotic drive and sex determination: molecular and cytological mechanisms of sex ratio adjustment in birds. Philosophical Transactions of the Royal Society B: Biological Sciences 363:1675-1686.

Schaeffer S, Bhutkar A, McAllister B, Matsuda M, Matzkin L, O'Grady P, Rohde C, Valente V, Aguade M, Anderson W, et al. 2008. Polytene Chromosomal Maps of 11 Drosophila Species: The Order of Genomic Scaffolds Inferred From Genetic and Physical Maps. Genetics 179:1601-1655.

Schittenhelm RB, Althoff F, Heidmann S, Lehner C. 2010. Detrimental incorporation of excess Cenp-A/Cid and Cenp-C into *Drosophila* centromeres is prevented by limiting amounts of the bridging factor Cal1. Journal of Cell Science 123:3768-3779.

Schuh M, Lehner CF, Heidmann S. 2007. Incorporation of Drosophila CID/Cenp-A and CENP-C into centromeres during early embryonic anaphase. Curr Biol. 17:237-243.

Stanke M, Morgenstern B. 2005. AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. Nucleic Acids Res. 33:W465-W467.

Talbert P, Bryson T, Henikoff S. 2004. Adaptive evolution of centromere proteins in plants and animals. J Biol. 3:18.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. Mol Biol Evol. 30:2725-2729.

van Hooff J, Tromer E, van Wijk LM, Snel B, Kops G. 2017. Evolutionary dynamics of the kinetochore network in eukaryotes as revealed by comparative genomics. EMBO reports doi:10.15252/embr.201744102

Wagner G, Kin K, Lynch V. 2012. Measurement of mRNA abundance using RNA-seq data: RPKM measure is inconsistent among samples. Theory Biosci. 131:281-285.

Xu B, Yang Z. 2013. PAMLX: a graphical user interface for PAML. Mol Biol Evol. 30:2723-2724

Zhang Z, Kishino H. 2004. Genomic Background Predicts the Fate of Duplicated Genes: Evidence From the Yeast Genome. Genetics 166:1995-1999.

Table 1. Distance between the *Cenp-C* paralogs.

	Sophophora C1 vs. Drosophila C1	Sophophora C1 vs. Drosophila C2	Drosophila C1 vs. Drosophila C2	Overall
R-rich	0.47540 (0.57118)	0.40707 (0.38894)	0.50030 (0.62341)	0.49964 (0.56808)
DH	0.39635 (0.45272)	0.37059 (0.39857)	0.38366 (0.45928)	0.40709 (0.46422)
NLS	0.40471 (0.41853)	0.39180 (0.37797)	0.45128 (0.49346)	0.44810 (0.46395)
CenH3 binding	0.37557 (0.41411)	0.38110 (0.38213)	0.32599 (0.32348)	0.37573 (0.39051)
AT2	0.57958 (0.76277)	0.45967 (0.57424)	0.55814 (0.74257)	0.55202 (0.71809)
Cupin	0.42511 (0.52160)	0.42624 (0.51710)	0.40136 (0.49603)	0.44415 (0.54587)

Note – Values refer to distances between coding DNA sequences, and values between brackets refer to amino acids distances.

Table 2. Summary of tests for positive selection performed on each *Cid* and *Cenp-C* paralog.

	Alignment length (#nts)	M1a vs. M2a	M7 vs. M8	M8a vs. M8
Cid1/Cid6	609	$P = 1$	$P = 1$	$P = 0.982$
Cid5	600	$P = 0.099$	$P = 0.069$	$P = 0.025*$
Cenp-C1	3,492	$P = 0.496$	$P = 0.163$	$P = 0.210$
Cenp-C2	3,696	$P = 0.194$	$P = 0.005*$	$P = 0.068$

Figure legends

Figure 1. *Cid1* **degenerated after its inter chromosomal duplication that originated** *Cid6*. (Upper panel) Comparison between the *Cid1* and *Cid6* loci of *D. buzzatii* and the corresponding regions of *D. mojavensis*. The black asterisk indicates a fragment of *Cid1*, 'N' indicates unidentified nucleotides, red boxes indicate transposable elements, and arrows indicate primers used for the fluorescent *in situ* hybridizations (FISH) experiments. (Lower panel) FISH on polytene chromosomes of *D. buzzatii* (Dbuz) and *D. seriema* (Dser) using *Cid6* probes, and of the closely related *D. mojavensis* (Dmoj) and the outgroup *D. virilis* (Dvir) using *Cid1* probes. The chromosome arm in which the *Cid* probe hybridized (red signal) is indicated by a letter that represents the corresponding Muller element. The chromocenter, a region in which all centromeres bundle together, is indicated by a white asterisk. The chromocenter of *D. buzzatii* and *D. mojavensis* ruptured during the fixation step.

Figure 2. *Cenp-C1* **duplicated in the lineage that gave rise to species of the *Drosophila* subgenus.** Maximum likelihood tree of the *Cenp-C1* and *Cenp-C2* nucleotide motif sequences. Red and green branches are respectively from *Cenp-C1* and *Cenp-C2* of species of the *Drosophila* subgenus. *D. busckii Cenp-C2* grouped with the outgroup *Phortica variegata Cenp-C1* likely due to long branch attraction. Bootstrap values are shown in each node. Scale bar represents number of substitutions per site.

Figure 3. Some Cenp-C motifs are alternatively conserved between Cenp-C1 and Cenp-C2.
(A) Schematic representation of the motif structure of *D. melanogaster* Cenp-C1. (B) Logo representations for each motif of the *Drosophila* subgenus Cenp-C1 (C1) and Cenp-C2 (C2). Motifs are as follow: R-rich, arginine-rich; DH, drosophilid Cenp-C homology; AT1, AT hook 1; NLS, nuclear localization signal; CenH3 binding, also known as Cenp-C motif; AT2, AT hook 2; Cupin, a dimerization domain near the C-terminal region. The asterisk in the CenH3 binding motif indicates the corresponding R1101 of *D. melanogaster*, which is crucial for the centromere localization of Cenp-C1.

Figure 4. ***Cid5* and *Cenp-C2* are male germline-biased.** *Cid* and *Cenp-C* expression patterns in *D. buzzatii* (A and B**)** and *D. virilis* and *D. americana* (C and D). *Cid* and *Cenp-C* transcripts are expected to be found in abundance only in mitotic active tissues. Therefore, *Cid* and *Cenp-C* transcripts are abundant in embryos and adult females likely due to oogenesis and oocyte mRNA accumulation required to sustain the high mitotic rate in early embryonic development, in adult males likely due to spermatogenesis, and in pupae likely due to morphogenesis. A low abundance of *Cid* and *Cenp-C* transcripts is expected in larvae, as most of its mitotic activity is limited to neuroblasts. Data are presented as mean ± 95% confidence interval and analyzed by one-way ANOVA (A and B) and Student's t-test (C and D): n.s., not significant; *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$. TPM, transcripts per million.