Multiple poliovirus-induced organelles suggested by comparison of spatiotemporal dynamics of membranous structures and phosphoinositides

Hyung S. Oh[1], Sravani Banerjee[1], David Aponte-Diaz, Jason Aligo[2], Maria F. Lodeiro and Craig E. Cameron*

Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, PA 16802

[1]These authors contributed equally to this study.

[2]Current address: Biologics Toxicology, Janssen Research and Development, LLC, Spring House, PA 19477

*Correspondence: cec9@psu.edu; 814-863-8705 (Phone); 814-863-7024 (Fax); @CameronLabPSU (Twitter)

Running Title: Multiple PV-induced organelles

ABSTRACT

1
Poliovirus (PV) infection induces membranes with elevated levels of phosphatidylinositol-4-phosphate
(PI4P) and invaginated, tubular structures that appear as vesicular clusters in cross section. Here, we
characterize PV mutants, termed EG and GG, which exhibit aberrant proteolytic processing of the P3
precursor that can delay the onset of genome replication and/or impair virus assembly. For WT PV,
changes to the PI4P pool were observed as early as 30 min post-infection. PI4P remodeling occurred even
in the presence of guanidine hydrochloride, a replication inhibitor. Vesicular clusters were not apparent
until 3 h post-infection, a time too slow for these structures to be responsible for genome replication.
Delays in the onset of genome replication observed for EG and GG PVs were explained completely by the
reduced kinetics of virus-induced remodeling of PI4P pools, consistent with PI4P serving as a marker of
the genome-replication organelle. Infectious virus produced by GG PV is known to be reduced nearly 5-
logs. We show that GG PV was unable to make virus-induced vesicular clusters. Instead, GG PV-infected
cells accumulated elongated tubules. Our results are consistent with the existence of distinct organelles
for genome replication and virus assembly. We suggest that the pace of formation and spatiotemporal
dynamics of PV-induced organelles may be set by the rate of P3 precursor processing to form 3AB and/or
3CD proteins.

AUTHOR SUMMARY – All positive-strand RNA viruses replicate their genomes in association with host cell
membranes. PV does not just remodel existing membranes, but induces membranes with unique
structure and lipid composition. There has been some suggestion that the functions of the PV-induced
structures observed during infection may not be those that perform genome replication. This study uses
kinetic analysis of virus-induced membrane formation and PI4P induction by two PV mutants to provide
evidence for the existence of a virus-induced genome-replication organelle distinct from a second
organelle, the absence of which impairs virus assembly. In addition, our studies suggest that formation
of both organelles may require participation of viral proteins, 3AB and/or 3CD. Therefore, this study
provides a new perspective on the cell biology of PV infection and should inspire a fresh look at
picornavirus-induced organelles, their functions and the role of P3 proteins in their formation.

INTRODUCTION

Positive-strand RNA viruses pose a great threat to public health because of their simplicity and

evolvability [1]. Introduction of a modestly-sized, mRNA-sense genome is sufficient to commandeer the

cell, establish infection and produce thousands of progeny in a matter of only hours. Among the least

understood aspect of positive-strand RNA virus biology is the eclipse phase of the life cycle, the events

post-entry but prior to exponential amplification of the genome and assembly of infectious virus. We use

poliovirus (PV), the type virus in the Enterovirus genus of the *Picornaviridae* family of viruses, as our model

system. Establishment of infection by PV likely requires entry of multiple genomes into the cell [2, 3].

This circumstance may be required to permit translation of infecting virion RNA to produce viral protein

in sufficient quantity to overwhelm host defenses that might otherwise curtail infection and to hijack host

factors essential to initiate a productive infection. It is becoming increasingly clear that picornaviruses

have evolved mechanisms to increase multiplicity of infection, for example by leaving cells in vesicles and

perhaps entering that way [4] or by attaching to bacterial surface polysaccharides and entering in

association with these organisms [5].

Another important function of viral proteins produced during the eclipse phase is establishment

of sites of genome replication. All positive-strand RNA viruses replicate in association with membranes;

three general strategies exist [6]. The first strategy is to commandeer an organellar membrane by

directing viral proteins to the organelle. Binding of the viral protein will induce invaginations with negative

curvature from the perspective of the cytoplasm that have been termed spherules. In response to actions

of the same or different viral proteins, the lipid and protein composition of the spherules may be

remodeled to reflect the genome-replication needs of the virus. The second strategy uses a two-step

process. The virus commandeers vesicular and/or organellar membranes, again by directing viral proteins

to their membranes. Hijacked membranes are then forced into a new form by remodeling lipid and/or

protein composition to create a virus-induced organelle best suited to genome replication. The third

strategy is a blend of the first two. Localize first to an organelle and later induce a virus-specific

organelle(s), which may be the case for enteroviruses [7].

Since the earliest studies of PV-infected cells using transmission electron microscopy (TEM), it has

been well documented that PV infection wreaks havoc on intracellular membranes, for example causing

dissolution of the Golgi apparatus (referred to throughout as Golgi), to create membranes of unique

benefit to the virus [8-12]. PV-induced membranes often appear as clusters of vesicles by TEM. These

structures have been shown to contain viral RNA and proteins [13-16]. More recent studies by Ehrenfeld,

Belov and colleagues using electron tomography have suggested that PV-induced membranes arise from

tubules [9]. The thought is that viral or viral-hijacked proteins induce invaginations of positive curvature

and promote folding of tubules, which would appear as vesicular clusters in cross section. In the Ehrenfeld

and Belov study, a kinetic analysis was performed that provided the first suggestion that the most

dramatic PV-induced changes to membranes occurred after genome-replication was well underway, if not

completed.

Prior to the tomography study, Altan-Bonnet and colleagues showed induction of

phosphatidylinositol-4-phosphate (PI4P) by another member of the Enterovirus genus, Coxsackievirus B3

(CVB3) [17]. In uninfected cells, PI4P localizes to the Golgi. During infection, PI4P starts at the Golgi then

relocalizes to a site between the endoplasmic reticulum (ER) and Golgi, perhaps the ER-Golgi intermediate

compartment (ERGIC). At later times post-infection, PI4P fills the entire perinuclear region of the cell and

now localizes with ER markers. During the entire infection period, PI4P localizes with viral RNA and

proteins. One conclusion of the study was that PI4P is a marker of the PV-induced organelle. When

considered in the context of the tomography study, it is possible that the perinuclear PI4P observed for

CVB3 reflects the invaginated and folded tubules observed for PV that might not function in genome

replication. Therefore, the earliest stages of PI4P induction and membrane remodeling may actually

represent the replication organelle, for which TEM images are not available.

The large polyprotein encoded by the PV genome can be divided into three smaller precursor

proteins: P1, P2 and P3 (**Fig. 1A**) [18]. P1 proteins form the capsid. P2 proteins are thought to be the

most functionally diverse of the PV proteins as these proteins shut off host cell pathways, derange host

cell membranes, function in genome replication and genome encapsidation, and can even leave their

mark on the encapsidated genome by somehow controlling its uncoating [12, 14, 19-30]. P3 proteins are

most intimately associated with genome replication as these proteins bind to the genome and catalyze

RNA synthesis [31-37].

PV uses differential cleavage of the polyprotein as one mechanism to expand its proteome.

Cleavage of the P3 precursor protein is mediated by the 3C-encoded protease activity. This protease

cleaves at Gln-Gly junctions, but the requirements for cleavage extend beyond a single dipeptide [38-40].

Two pathways exist for cleavage of P3 [41, 42]. The major pathway produces only two proteins: 3AB and

3CD (**Fig. 1B**). The minor pathway is more complex. First, P3 is cleaved to produce 3A and 3BCD proteins.

3BCD is then cleaved to produce 3BC and 3D. Finally, 3BC is cleaved to produce 3B (also known as VPg,

virion protein linked to the genome) and 3C.

Previously, we reported two PV mutants in which the cleavage site between 3B and 3C was

changed to Gly-Gly (GG PV) [42] or Glu-Gly (EG PV) [42, 43]. For GG PV, the products of the major pathway

were 3ABC and 3D, thus ablating the production of both 3AB and 3CD (see GG in **Fig. 1B**). With the

exception of 3BC, which cannot be cleaved, products of the minor pathway were essentially unchanged

[42]. For EG PV, the products of both pathways were the same as observed for wild type (see EG in **Fig.**

1B). However, the rate of cleavage at the 3BC junction was reduced, causing a reduced rate of

accumulation of 3AB and 3CD. Both mutants exhibited what appeared to be a reduced rate of genome

replication. For EG PV, ectopic expression of 3CD alone corrected this phenotype, consistent with the

phenotype arising solely from the reduced rate of 3CD production. In addition to the impact on genome

yield, GG PV exhibited a near-complete loss of infectious virus. Neither of the GG PV phenotypes could

be suppressed by ectopic expression of 3CD.

The extended duration of the eclipse phase observed for EG PV and the defect to virus assembly

observed for GG PV provided a unique opportunity to expand our understanding of these steps of the

virus lifecycle at the molecular and ultrastructural levels. We find that there is a clear temporal ordering

of PI4P induction and its initial induction and limited redistribution exhibit the expected kinetics for an

organelle involved in genome replication. During the early stages of PI4P remodeling, Golgi fragments

and ERGIC expands. It is only after genome replication is near completion that the perinuclear region of

the cell becomes filled with vesicular clusters. At the same stage of genome replication that vesicular

clusters predominate for WT PV, tubules predominate for GG PV. The inability of GG PV to assemble

infectious virus is consistent with the possibility that vesicular clusters contribute substantively to steps

after genome replication on path to production of infectious virus. We propose that two organelles are

induced during PV infection that serve distinct functions during the PV lifecycle and that P3-encoded

proteins contribute to formation and function of both.

 RESULTS

Delayed release of 3AB and 3CD proteins from the P3 precursor protein delays onset of genome

replication.

Our previous studies of EG PV showed that its genome-replication defect could be complemented

in trans by expression of 3CD [43]. This observation was unexpected because numerous studies have

shown that genome-replication defects can only be complemented in trans by expression of P3 or P2-P3

polyproteins, if at all [44-46]. Our study used a subgenomic replicon in which P1-coding sequence had

been replaced with firefly luciferase-coding sequence, and replication was monitored indirectly by

measuring luciferase activity. It was possible that measuring luciferase activity instead of RNA confounded

our interpretation of the experiment.

To test this possibility, we transfected WT or EG PV subgenomic replicon RNA into HeLa cells. We

monitored RNA synthesis either by measuring luciferase activity (**Fig. 2A**) or by measuring positive-sense

RNA using Northern blotting (**Figs. 2B and 2C**). We observed three discernible phases of luciferase activity

for WT: 0-2 h, 2-5 h, and 5-10 h; and for EG: 0-3 h, 3-7 h and 7-10 h (**Fig. 2A**). In contrast, we observed

only one phase of RNA accumulation for both (**Fig. 2B**). WT RNA accumulation began between 1-2 h post-

transfection; EG RNA accumulation began between 3-4 h post-transfection (**Fig. 2B**). We conclude that a

delay of two hours in the onset of replicon RNA replication can be obscured by monitoring luciferase

activity. Luciferase activity concealed the delay. Translation of transfected replicon RNA produced

luciferase activity for at least two hours in the presence of guanidine hydrochloride (GuHCl), a replication

inhibitor, and therefore without the need for replicon RNA replication (see WT +GuHCl in **Fig. 2A**). These

results support the hypothesis that delayed kinetics of P3 precursor processing to form 3AB and 3CD in

cells causes a delay in a step preceding the onset of RNA synthesis.

Kinetics of formation of PV-induced vesicular clusters are inconsistent with these structures serving as

the sites of genome replication.

A hallmark of PV-infected cells as observed by transmission electron microscopy (TEM) is the

presence of vesicular clusters [9, 12]. These vesicular clusters are also referred to as replication complexes

or replication organelles. Given the accepted role of these vesicular clusters in genome replication, we

reasoned that formation of these structures should occur either prior to the onset of genome replication

or concomitant with genome replication. Formation and/or function of these structures could therefore

represent the pre-genome-replication step perturbed for EG PV.

We infected HeLa cells with EG PV and then at various times post-infection processed samples to

measure genomic RNA by Northern blotting (**Figs. 3A** and **3B**), to measure infectious virus by plaque assay

(**Fig. 3A**) and to visualize formation of the vesicular clusters by TEM (**Fig. 3C**). The exponential phase of

genome replication occurred between 2 h and 6 h post-infection (**Figs. 3A** and **3B**). The exponential phase

of infectious virus production occurred between 4 h and 8 h post-infection (**Fig. 3A**). Vesicular clusters

became visible at 4 h post-infection and continued to increase in abundance and/or change in organization

throughout the 7 h time course (**Fig. 3C**). Formation of vesicular clusters neither preceded genome

replication nor occurred concomitant with genome replication as expected for an organelle responsible

for genome replication. In contrast, formation of vesicular clusters preceded infectious virus production.

The delay in induction of the vesicular clusters by EG PV provided a unique opportunity to monitor

the early steps in formation of these structures. In uninfected cells, it was very easy to observe

mitochondria and Golgi. However, closer inspection was required to observe the tubules of the

endoplasmic reticulum (ER) (uninfected in **Fig. 3C**). By 2 h post-infection, the density of mitochondria had

diminished (2 h in **Fig. 3C**). Golgi stacks were fragmented and swollen, and the tubules of the ER and/or

ER-Golgi intermediate compartment (ERGIC) were swollen and easier to visualize (2 h in **Fig. 3C**). At 3 h

post-infection, much of the Golgi had disappeared (3 h in **Fig. 3C**). The area between the remaining Golgi

and the now very swollen ER was filled with very small vesicles that may be the ever-expanding ERGIC

caused by the loss of the Golgi (3 h in **Fig. 3C**) [17]. At 4 h post-infection, vesicular clusters were first

observed (4 h in **Fig. 3C**). These vesicular clusters appear to be encased by the swollen tubules of the ER

(4 h in **Fig. 3C**).

In order to make certain that our observations above were not unique to the mutant virus, we

repeated the experiment using WT PV (**Fig. 4**). The exponential phase of genome replication occurred

between 1 h and 5 h post-infection (**Figs. 4A** and **4B**). The exponential phase of infectious virus production

occurred between 3 h and 7 h post-infection (**Fig. 4A**). Vesicular clusters were now clearly visible at 3 h

post-infection and again continued to increase in abundance or change in organization throughout the 7

h time course (**Fig. 4C**). Many of the early steps described for EG PV could not be observed for wild type

because they occurred so soon post-infection that we could not process the cells fast enough to capture

images of these steps by TEM.

In addition to the temporal difference in the formation of the vesicular clusters observed for EG

PV and WT PV, differences also existed in the overall size, interaction and localization of these structures

(compare 5 h for EG PV in **Fig. 3C** to 4 h for WT PV in **Fig. 4C**). Relative to EG PV, vesicular clusters induced

by WT PV appeared larger, clustered more tightly and were far more constrained to the perinuclear region

of the cell.

Both experiments were consistent with the surprising finding that the kinetics of formation of the

PV-induced vesicular clusters are inconsistent with these structures serving as the genome-replication

organelle. These structures became easily identifiable two hours after genome replication began (4 h for

EG PV in **Fig. 3C**; 3 h for WT PV in **Fig. 4C**). Interestingly, and perhaps coincidentally, this was the time

after genome replication began that we were able to detect infectious virus by plaque assay (**Figs. 3B** and

4B).

Kinetics of PI4P induction are consistent with this phosphoinositide demarcating the sites of genome

replication.

Coxsackievirus B3 (CVB3) infection induces PI4P and this phosphoinositide is a component of a

virus-induced organelle [17]. PI4P co-localizes with viral non-structural proteins and RNA [17]. Whether

or not PI4P localizes to the virus-induced vesicular clusters is not known. Given our results, it is possible

that the virus-induced PI4P localizes to membranes other than those observed by TEM well after genome

replication has begun.

Although it is assumed that PV infection induces PI4P based on the data for CVB3, PI4P induction

by PV has never been reported. We infected HeLa cells with WT PV or EG PV and measured PI4P as a

function of time post-infection by immunofluorescence microscopy (IFM) (**Fig. 5**). We used an antibody

specific for PI4P to detect this phosphoinositide.

It is well known that most PI4P in the cell localizes to the Golgi, giving rise to very nice staining

with anti-PI4P antibody (mock in **Fig. 5A**) [17]. For WT PV, induction and redistribution of Golgi-localized

PI4P was evident as early as 30 min post-infection WT in **Fig. 5B**). Both redistribution and induction

continued to 3 h post-infection, the last time point evaluated (WT in **Fig. 5B**). Observations with EG PV

were essentially identical, except for a one-hour delay of the changes relative to wild type (EG in **Fig. 5B**).

The kinetics of induction and redistribution of PI4P were consistent with a structure responsible for

genome replication. Given the TEM experiments reported above (**Figs. 3** and **4**), it is now clear that the

PI4P-containing membranes observed by IFM prior to two hours post-infection (**Fig. 5B**) must be distinct

from the large vesicular clusters that appear after genome replication (**Figs. 3** and **4**).

PI4P induction and redistribution during WT PV infection occurred as fast as we could reliably

process our cells for imaging. This observation suggested to us that viral proteins made by translation of

infecting viral RNA might be sufficient to initiate PI4P remodeling. We infected cells with WT or EG PV in

the presence of the genome-replication inhibitor, GuHCl. Under these conditions, the only event that can

occur is translation of the infecting genomes. PI4P induction and redistribution occurred in the presence

of GuHCl for both WT (WT+GuHCl in **Fig. 5C**) EG (EG+GuHCl in **Fig. 5C**) PVs. GuHCl did not have an impact

on PI4P localization in the absence of infection (Mock+GuHCl in **Fig. 5C**).

PV-induced vesicular clusters as sites of virus assembly, an assembly organelle?

The studies reported above suggested that PI4P was a marker for the PV genome-replication

organelle. These studies also suggested that PI4P-containing structures observed early are distinct from

PV-induced vesicular clusters. The question now was: what is the function of the vesicular clusters? One

possibility was that the vesicular clusters are structures required for virus assembly, an assembly

organelle, as suggested previously by Belov and Ehrenfeld [9]. Another possibility was that the vesicular

clusters are structures that represent accumulated, unused and unwanted viral proteins and nucleic acids,

a viral trash compacter.

GG PV replicated to within one log of wild type but exhibited a near 5-log reduction in the amount

of infectious virus produced [42]. We reasoned that this mutant might help us identify the function of the

PV-induced vesicular clusters. If the vesicular clusters contribute to virus assembly, then these structures

might be perturbed in cells infected with the assembly-defective GG PV. Conversely, if the vesicular

clusters function as a means to consolidate viral waste, then these structures might appear unchanged in

cells infected with GG PV.

We transfected HeLa cells simultaneously with two subgenomic replicons, one expressing

luciferase and the other expressing EGFP. Transfected cells were then grown in suspension culture and

harvested hourly to monitor luciferase activity and EGFP fluorescence. Luciferase specific activity,

measured as relative light units per microgram of protein in the extract, was plotted as a function of time

for WT and GG replicons (**Fig. 6A**). We were particularly interested in 5 h post-transfection for WT as

vesicular clusters reached their maximum level at that time (**Fig. 4C**). At 5 h post-transfection of the WT

replicon combination, the luciferase specific activity was approximately 10^4 RLU/μg (**Fig. 6A**). We

observed the equivalent luciferase specific activity at 14 h post-transfection of the GG replicon

combination (**Fig. 6A**). Based on this result, we concluded that 5 h and 14 h post-transfection represented

equivalent extents of the infection process for WT and GG PVs, respectively.

We used fluorescence activated cell sorting (FACS) to isolate transfected cells based on EGFP

expression. Cells replicating WT or GG EGFP-replicon RNA before and after sorting are shown in **Fig. 6B**.

We used the EGFP-expressing, sorted cells for TEM (**Fig. 6C**). Vesicular clusters were readily apparent in

images of WT replicon-transfected cells, and these clusters were of the same overall appearance as

observed using infected cells that did not require sorting (**Fig. 4C**). In contrast, vesicular clusters were few

or absent in images of GG replicon-transfected cells (**Fig. 6C**). Instead, we observed a tubular/reticular network (see enlargement of panel GG in **Fig 6C**).

These results support the hypothesis that vesicular clusters function as an assembly organelle, and the absence of these structures in GG PV contribute to the defect in virus production. We propose that the tubular/reticular network is an intermediate on the pathway to formation of the vesicular clusters. These structures were apparent between 2 h and 4 h post-infection for EG PV (**Fig. 3C**) but occurred too early to be observed for WT PV (**Fig. 4C**). The tubules appeared to fold in on themselves, creating the vesicular clusters (see for example 4 hpi for EG PV in **Fig. 3C** or 3 hpi for WT PV in **Fig. 4C**). Formation of tubules and their transformation into a net-like array of vesicles has been observed before [9]. Additional studies will also be required to further clarify the relationship between the tubular/reticular network in GG PV and the vesicular clusters in EG PV and WT PV.

Finally, we assessed formation of the PI4P-marked structures by GG PV RNA. We transfected HeLa cells with GG PV EGFP-expressing, subgenomic replicon RNA and used IFM to monitor the spatiotemporal dynamics of PI4P. Redistribution and induction of PI4P occurred substantially slower than observed for EG PV or WT PV (compare **Fig. 6D** to **Fig. 5**). Nevertheless, temporal dynamics of PI4P explains the 8 h or so pre-genome-replication phase observed previously for GG PV [42, 43]. These data are consistent with PI4P demarcating the replication organelle and a requirement for induction of this structure prior to the onset of genome replication.

High multiplicity of infection exaggerates spatiotemporal dynamics of vesicular clusters.

The spatiotemporal dynamics of WT PV-induced vesicular clusters reported here were different than observed previously by others [9, 11, 16]. In particular, formation of the clusters was slower, and the density of the clusters was lower (**Fig. 4C**) than previously reported. In these earlier experiments, multiplicity of infection ranged from 30-50, but this study used 10. Because translation of the infecting genomes is sufficient to begin the remodeling process, it was possible that elevated levels of protein produced at the higher multiplicities of infection might impact the size of the clusters. To address this possibility, we infected HeLa cells with WT PV at a multiplicity of 10, 50 or 100 and processed the infected cells for TEM at 3 h post-infection. This time point was chosen because of the modest density of the vesicular clusters observed at a multiplicity of infection of 10 (**Fig. 4C**) and should make any observed differences more noticeable. As shown in **Fig. 7**, the size of the vesicular cluster at the level of each "vesicle" and in terms of the area of the perinuclear region occupied by the vesicular clusters scaled directly with the multiplicity of infection. Therefore, it is imperative to consider multiplicity of infection when comparing micrographs.

DISCUSSION

This study was inspired by our observation that ectopic expression of PV 3CD protein was

sufficient to complement a genome-replication defect of the EG PV mutant, which exhibited a delay in

processing of the P3 precursor protein [43]. The rule of thumb for PV and other picornaviruses is that

non-structural proteins used to replicate a genome are produced in cis—that is, by that genome or by one

in the same replication complex [44-46]. However, it became clear that the genome-replication defect

was actually a defect at steps preceding genome replication (**Fig. 2B**). The confusion was caused by the

multiphasic kinetics of reporter activity produced by a subgenomic replicon (**Fig. 2A**). The first phase of

activity requires only translation of the transfected replicon. The second phase requires replicon

replication. The third phase appears to be translation of replicated replicon RNA that would normally be

packaged if capsid proteins had been present.

Once we knew that EG PV impaired steps leading up to genome replication, we turned our focus

to formation of the "replication organelle." Since the first TEM images of PV-infected cells were published

[8], multiple reports have shown that a hallmark of PV infection is the induction of membranous structures

[9-12]. These structures have many names: replication complexes, vesicular clusters, tubules and

replication organelle. Most of the early studies searching for viral factors contributing to formation of PV-

induced membranes implicated P2 proteins in this process [14, 15]. The most convincing experiment

showed that expression of the 2BC precursor proteins was sufficient to induce membranes with the same

buoyant density and appearance by TEM as those observed during PV infection [20, 21]. As a result, it

was unclear how processing of the P3 precursor to form 3AB and 3CD contributed to formation of the

replication organelle. However, because formation of this structure should precede genome replication,

it was possible that this step could be complemented in trans. 3A is known to inhibit guanine nucleotide

exchange factor (GEF), GBF1, which prevents trafficking between ER and Golgi and thereby causes

dissolution of the Golgi [47-52]. 3AB might also have this activity. Inhibition of ER-to-Golgi trafficking

should enlarge the ER-Golgi intermediate compartment, and this compartment has been implicated in

genome replication [17, 51]. 3AB is also known to promote invagination of membranes, an activity that

could contribute to formation of vesicular clusters [53]. 3CD, on the other hand, has not been shown to

perturb membrane form or function in cells. However, in cell-free extracts, the presence of 3CD promotes

recruitment of ADP-ribosylation factors (ARFs) and GEFs to membranes, which could modulate vesicular

trafficking and/or fusion [54, 55].

Our examination of the kinetics of formation of the vesicular clusters by EG PV relative to the

kinetics of genome replication was eye-opening. The kinetics of these two processes did not have the

expected behavior---that is, the site of RNA synthesis forming prior to RNA synthesis (**Fig. 3**). This
unexpected kinetic behavior was not an oddity of the mutant as WT PV also exhibited this behavior (**Fig.**
4). Our conclusion was that the vesicular clusters must not be the replication organelle. Belov and
Ehrenfeld floated this possibility several years ago [9]. In reviewing the literature on formation of PV-
induced vesicles, it became clear that the TEM images of others showed faster formation of vesicular
clusters in spite of the fact that kinetics of genome replication and/or production of infectious virus were
identical [9, 11, 16]. The one difference between our study and those of others was the multiplicity of
infection used. We used an MOI of 10 pfu/cell; others used an MOI of 30-50 pfu/cell. When we increased
the MOI to these higher levels, we observed faster kinetics of formation of the vesicular clusters (**Fig. 7**).
Under these conditions, the vesicular clusters were also substantially larger (**Fig. 7**). These observations
suggest that the kinetics of vesicular cluster formation can be modulated by multiplicity of infection
without a concomitant increase in the kinetics of genome replication or virus assembly, thus creating
some ambiguity to the temporal ordering of formation of the vesicular clusters relative to replication of
the genome.

Altan-Bonnet and colleagues showed that phosphatidylinositol-4-phosphate (PI4P) is induced by
Coxsackievirus B3 infection and redistributes from the Golgi to sites thought to be involved in genome
replication [17]. Therefore, we monitored the spatiotemporal dynamics of PI4P during PV infection. We
found that the spatiotemporal dynamics of the PI4P pool was completely consistent with PI4P
demarcating the replication organelle as induction and redistribution occurred well before the onset of
genome replication (**Fig. 5A**). Moreover, spatiotemporal dynamics of PI4P was delayed for EG PV (**Fig.**
5B). Unexpectedly, PI4P induction and redistribution did not require any genome replication as these
events occurred in the presence of a replication inhibitor. To the best of our knowledge, this observation
is among the first to link translation of infecting viral genomes alone to events essential for genome
replication. Under these conditions, the amount of viral proteins produced cannot be detected readily by
Western blotting or immunofluorescence [42, 43]. Therefore, very little of one or more viral proteins is
sufficient to hijack cellular factors needed for PI4P induction and redistribution.

We propose that 3AB and/or 3CD contribute to PI4P induction and/or redistribution. If this is the
case, then delays in P3 processing would be expected to have an impact on PI4P dynamics. Our initial
characterization of EG PV showed that its genome-replication phenotype could be complemented fully by
ectopic expression of 3CD [43]. It is therefore possible that 3CD or 3CD-containing precursor alone
contribute to PI4P dynamics. Interestingly, synthesis of PI4P by the beta isoform of the class III
phosphatidylinositol-4-kinase (PI4KIIIβ), the isoform thought to be responsible for induction of PI4P by

enteroviruses [17], is known to require Arf1 and GBF1 [17]. Perhaps the ability of 3CD to induce

translocation of Arf1 and GBF1 from the cytoplasm to membranes may be related to its role in PI4P

induction suggested here [54, 55].

339   The delay in formation of the replication organelle by EG PV created a window into the early

340   events of this process. It was possible to visualize Golgi fragmentation, proliferation of vesicles in the area

341   between ER and Golgi, and formation of short, thick tubules (**Fig. 3**). These events were suggested by

342   Altan-Bonnet and colleagues based on Golgi, ERGIC and ER marker co-localization with PI4P and CVB3

343   non-structural proteins and/or RNA [17]. Together, these data make a compelling case for these localized

344   membranous structures including and between ER and Golgi representing the replication organelle.

345   Clues into the formation and function of PV-induced vesicular clusters were obtained from studies

of GG PV. This mutant took even longer to initiate genome replication than EG PV, and this delay was

attributable to an even longer delay in PI4P induction and redistribution (**Fig. 6**). GG PV does not produce

any 3AB or 3CD, which is likely the cause of this phenotype [42, 43]. This mutant clearly establishes a role

for P3 proteins in formation of the replication organelle. However, TEM of GG PV-infected cells revealed

a cell full of tubules at times in which genome replication of GG PV RNA was occurring exponentially (**Fig.

6**). Hints of tubulation were evident in micrographs of EG and WT PVs (**Figs. 3 and 4**), but tubules never

really accumulated. We propose that the tubules observed in GG PV-infected cells are an intermediate

on path to formation of the vesicular clusters. Positive invagination of the tubules and interactions

between invaginated membranes should produce vesicular clusters, as seen in cross section by TEM. As

to function of the vesicular clusters, GG PV was unable to produce significant titers of infectious virus [42,

43]. We propose that formation of vesicular clusters requires 3AB and/or 3CD and facilitates virus

assembly. It is known that 3CD promotes cell-free synthesis of infectious PV [56, 57]. It is possible that

these membranes serve as an assembly organelle.

359   There was a time when poliovirus non-structural proteins were considered in two categories:

those that had an impact on interactions with the host; and those that had an impact on genome

replication. Emphasis was often placed on shut off of host protein synthesis, permeabilization of cellular

membranes and formation of vesicles by P2-encoded proteins. P3-encoded proteins bind to the viral

genome, produce the peptide/precursor used to initiate genome replication and function to replicate the

genome. This study highlights the almost unlimited functional capacity of PV non-structural proteins by

now implicating P3 proteins in events both before and after genome replication. Understanding how

these proteins contribute to these events requires further study. It is likely that the findings reported

here for PV will apply to other picornaviruses as well.

MATERIALS AND METHODS

Materials. Cell culture media and supplements were from Invitrogen Life Technologies (Gibco); Difco-

NZCYM for bacterial growth was from BD Biosciences; 5X Cell Culture Lysis Reagent was purchased from

Promega; primary antibody against PI4P was purchased from Echelon biosciences; Alexafluor-594-IgM

(goat anti-mouse) was from ThermoFisher Scientific; 16% formaldehyde, methanol-free, ultrapure was

purchased from polysciences; Digitonin was from Sigma-Aldrich; Bovine Serum Albumin (BSA)-Fraction V,

heat shock treated was from Fisher Bioreagents; Vectashield Antifade Mounting Medium with DAPI was

from Vector Laboratories; nitrocellulose membrane was from GE Healthcare; TransIT-mRNA transfection

kit was purchased from Mirus Bio LLC.

Construction of 3B-3C Gly-Gly mutant EGFP subgenomic replicon*.* Enhanced green fluorescence protein

(EGFP) integrated subgenomic replicon plasmid, pREGFP-, was generated by replacing luciferase gene in

pRLuc- [42] with EGFP. PCR was used to amplify the EGFP region using the oligonucleotides NotI-EGFP-

for (5'-AAT TCG GAG CGG CCG CTG TGA GCA AGG GCG AGG AGC-3') and EGFP-XhoI-rev (5'-GTC AGA TCC

TCG AGC TTG TAC AGC TCG TCC ATG-3') and pEGFP as a template. The PCR product replaced luciferase

gene in pRLuc-GG using NotI and XhoI sites to obtain pREGFP-GG subgenomic replicon. Sequencing at the

Penn State Nucleic Acid facility was used to confirm all clones.

RNA transcription. Linearization and RNA transcription were performed as described previously [42].

Briefly, the pMo-, pRLuc- and pREGFP- plasmids were linearized with EcoRI or ApaI and purified with Qiaex

II bead suspension (Qiagen) following manufacture's protocol. RNA was then transcribed from the

linearized plasmid DNAs in a 20 μL reaction containing 350 mM HEPES, pH 7.5, 32 mM magnesium acetate,

40 mM dithiothreitol (DTT), 2 mM spermidine, 28 mM nucleoside triphosphates (NTPs), 0.025 μg/μL

linearized DNA, and 0.025 μg/μL T7 RNA polymerase. The reaction mixture was incubated for 3 h at 37°C

and magnesium pyrophosphate was removed by centrifugation for 2 min. The supernatant was

transferred to a new tube and subjected to RQ1 DNase treatment (Promega) for 30 min at 37°C. RNA

quality was then verified by 0.8% agarose gel electrophoresis. The RNA concentration was determined by

comparing pre-known purified RNA side-by-side on an agarose gel.

Subgenomic luciferase replicon assay. Subgenomic luciferase assays were performed as described

previously [42] with the following modifications. Subgenomic replicon RNA (5 μg of *in vitro* transcribed

RNA) was electroporated into HeLa cells. The cells were incubated in normal growth media (DMEM/F12

supplemented with 10% fetal bovine serum, 1% penicillin/streptomycin, 5 mL/1 X 10^6 cells) and 1 X 10^5

cells were harvested and lysed using 100 μL of 1X cell culture lysis reagent (CCLR) at an indicated time

post-electroporation. Luciferase activity was measured by adding an equal volume of firefly luciferase

assay substrate to cell lysate and the reaction mixture was applied to a Junior LB 9509 luminometer

(Berthold Technologies) to read relative light units (RLU) for 10 s. Relative light units (RLU) were then

normalized based on the total protein concentration determined by Biorad protein assay reagent (BioRad)

for each sample. GuHCl when used, the final concentration was 3 mM and the inhibitor was kept on cells

during transfection and after, for the entire incubation period.

Total RNA preparation. One day prior to infection, 4 X 10^6 HeLa cells were seeded in a 100 mm dish. On

the day of infection, the prepared HeLa monolayers were infected at an MOI 10 with WT or EG PV for 30

min at room temperature. Infected cells were washed once with PBS to remove unattached virus and

then incubated in normal growth media (DMEM/F12 supplemented with 10% fetal bovine serum and 1%

penicillin/streptomycin) at 37°C. At the indicated times post-infection, cells were lysed with 1 mL of Trizol

(Invitrogen Life Technologies) and total RNAs were purified by following manufacture's protocol. In order

to purify subgenomic replicon RNA containing total RNA, HeLa cells were transfected with subgenomic

RNA as described above and 1.2 X 10^6 HeLa cells were lysed with 1 mL Trizol. The RNA quantity was

determined using NanoDrop 1000 (ThermoFisher Scientific) and the quality was evaluated by using 0.8%

agarose gel.

Northern-Blot analysis of RNA synthesis. Northern-Blot analysis was performed as described previously

[43]. Briefly, the RNA was separated on a 0.6% denaturing agarose gel (0.8 M formaldehyde in 1X MOPS)

by running at 120V for 2 h. The gel was washed in water for 30 min twice followed by soaking in 20X SSC

(Saline Sodium Citrate) buffer for 30 min. RNA was transferred to nylon membrane (Hybond XL, GE

Healthcare) using capillary blotting with 20X SSC for 16 h at room temperature. RNA was crosslinked to

membrane using UV Crosslinker (Stratalinker 2400, Stratagene). The membrane was washed twice with

wash buffer (1X SSC and 0.1% SDS) at 65°C for 30 min each and prehybridized in 100 mL of modified church

buffer (0.5 M sodium phosphate, pH 7.2, 7% SDS, and 1 mM EDTA) for 4 h at 65°C. Hybridization was

performed in a modified church buffer for 16 h at 65°C. The membrane was washed with wash buffer (1X

SSC and 0.1% SDS) for 20 min at 65°C twice and once at room temperature. RNA was visualized by

exposing the membrane to a phosphor screen followed by scanning the screen on a Typhoon 8600 scanner

(Promega) in phosphor mode. Hybridization probe was made by PCR using oligonucleotides: 3Dseq-100-

for (5'-GTT TGA AGG GGT GAA GGA A-3') and 3Dseq-1085-rev (5'-CTC CCA TGT GAC TGT TTC AAA TG-3')

and pRLuc-RA as a template. In the reaction, [α-^{32}P]-dATP (1 mCi/mL, 3000 Ci/mmol, GE Healthcare) was

used with cold dNTPs (300 µM for each dCTP, dGTP, and dTTP and 10 µM for dATP) in total 100 µl reaction.

The quality of PCR product was confirmed by agarose gel electrophoresis and cpm was determined by

using scintillation counter (LKB Wallac 1217 Rackbeta liquid scintillation counter).

One step growth curve assay. One day prior to infection, HeLa cells were seeded into 6-well plate at a

density of 5 X 10^5 cells/well. On the day of infection, the prepared HeLa monolayers were washed and

infected with WT or EG virus at an MOI of 10 for 30 min at room temperature. Infected cells were washed

once with PBS to remove unattached virus and then incubated at 37°C. At varying times post-infection,

virus was harvested from cells and media by three freeze-thaw cycles with vortexing in between. The

isolated virus was quantified by plaque assay.

Virus quantification by plaque assay. For plaque assay, 0.5 X 10^6 HeLa cells were plated one day prior to

infection in 6-well plates. On the day of infection, the media was removed and the cells were washed

once in PBS. Next, the cells were infected with the 10-fold dilutions of viral supernatants previously

prepared and incubated for 30 min at 37°C. The inoculum was then removed and the cells were overlaid

with 1% low-melting-point agarose (EMD). The overlay was allowed to solidify for 20 min at room

temperature, and the plates were incubated at 37°C for 2 days to allow for plaque formation. 30 min

prior to harvesting, the overlays were incubated with 4% formaldehyde/PBS to fix virus and cells. The

agarose monolayers were then removed and plaques stained with crystal violet and counted. Viral titers

were calculated in PFU/ml.

Sorting subgenomic replicon expressing cells with flow cytometer. WT and GG-subgenomic replicon

RNAs (2 µg of *in vitro* transcribed pRLuc RNA and 4 µg of *in vitro* transcribed pREGFP RNA) were

electroporated into 1.2 X 10^7 HeLa cells. The cells were incubated in normal growth media at 34°C and

luciferase activity was monitored. When the specific activity of luciferase was about 10,000 RLU/µg of

total protein (5 h post-transfection for pRLuc-/pREGFP-WT and 14 h post-transfection for pRLuc-/pREGFP-

GG), cells were harvested, chilled on ice for 5 min and sorted to collect transfection positive cells. For the

sorting, EGFP positive cells were excited with a 488 Argon laser, detected with a 532/40 nm band pass

filter and sorted by Cytopeia Influx sorter (Beckton-Dickinson, San Jose, CA) with 100 micron tip at 15 psi.

The sorted HeLa cells were processed for transmission electron microscopy.

Transmission electron microscopy. Virus-Infected HeLa cells or EGFP-positive post-sorted HeLa cells were

harvested, fixed and embedded for TEM studies as described previously [43]. Briefly, the harvested cells

were fixed with 1% glutaraldehyde, washed with 0.1 M cacodylate (Sodium dimethyl arsenate, Electron

Microscopy Sciences) twice for 5 min each, incubated in 1% reduced osmium tetroxide containing 1%

potassium ferricyanide in 0.1 M cacodylate for 60 min in the dark with one exchange and washed two

times with 0.1 M cacodylate again. En bloc staining was performed with 3% uranyl acetate in 50% ethanol

for 60 min in the dark. Dehydration was carried out with different concentrations of ethanol (50, 70, 95

and 100% for 5-10 min) and 100% acetonitrile. Embedding was performed overnight with 100% Epon at

65ºC. The embedded sample was sectioned with a diamond knife (DiATOME) to slice 60-90 nm thickness

by using ultra microtome (Reichart-Jung). The sectioned sample was place on copper grid (Electron

Microscopy Science), stained with 2% uranyl acetate in 50% ethanol followed by lead citrate staining for

12 min. The grid was washed with H_2O and the grid was dried completely. The image was obtained by

using Jeol JEM 1200 EXII and FEI Tecnai G2 Spirit BioTwin located in Electron Microscopy Facility in

Pennsylvania State University.

Immunofluorescence. 2.5 X 10^5 cells were seeded on coverslips in 6-well plates and next day infected

with WT or EG virus at an MOI of 10. GuHCl whenever used, the final concentration was 3 mM and the

inhibitor was kept on cells with the virus for 30 min and for the entire incubation period thereafter. For

subgenomic replicon RNA transfection, same number of cells were transfected with 2 µg pREGFP-GG RNA

using the RNA transfection kit (Mirus). Infected or transfected cells were fixed with 4% formaldehyde at

indicated times post-infection or transfection. Post-fixation, cells were permeabilized with 20 µM

digitonin for 10 min and washed with PBS. Cells were then blocked with 3% BSA in PBS for 1 h followed

by incubation with anti-PI4P (1:200) antibody for 1 h in blocking buffer. This was followed by washing with

PBS and incubating with Alexafluor594; IgM (1:1000) for 1 h. Finally, the cells were washed with PBS and

mounted on slides using mounting buffer with DAPI. Samples were imaged with Zeiss Axiovert 200 M

epifluorescence microscope.

ACKNOWLEDGMENTS

We thank Dr. Greg Ning and staff of the electron microscopy core facility for their guidance and assistance.

This study was funded by grant AI053531 from NIAID, NIH to CEC.

.

AUTHOR CONTRIBUTIONS

Conceptualization: CEC

Funding acquisition: CEC

Investigation: HSO, SB, DAD, JA, MFL

Project administration: HSO, SB, DAD, CEC

Supervision: CEC

Validation: HSO, SB, DAD

Visualization: HSO, SB, DAD

Writing- original draft: HSO, SB, CEC

Writing- review & editing: HSO, SB, JA, CEC

REFERENCES

1. Holmes EC, Drummond AJ. The evolutionary genetics of viral emergence. Curr Top Microbiol Immunol.

2007; 315:51-66. Epub 2007/09/13. PMID: 17848060.

2. Korboukh VK, Lee CA, Acevedo A, Vignuzzi M, Xiao Y, Arnold JJ, et al. RNA virus population diversity, an

optimum for maximal fitness and virulence. J Biol Chem. 2014; 289(43):29531-44. Epub 2014/09/13.

https://doi.org/10.1074/jbc.M114.592303 PMID: 25213864.

3. Aguilera ER, Erickson AK, Jesudhasan PR, Robinson CM, Pfeiffer JK. Plaques Formed by Mutagenized

Viral Populations Have Elevated Coinfection Frequencies. MBio. 2017; 8(2). Epub 2017/03/16.

https://doi.org/10.1128/mBio.02020-16 PMID: 28292984.

4. Chen YH, Du W, Hagemeijer MC, Takvorian PM, Pau C, Cali A, et al. Phosphatidylserine vesicles enable

efficient en bloc transmission of enteroviruses. Cell. 2015; 160(4):619-30. Epub 2015/02/14.

https://doi.org/10.1016/j.cell.2015.01.032 PMID: 25679758.

5. Robinson CM, Jesudhasan PR, Pfeiffer JK. Bacterial lipopolysaccharide binding enhances virion stability

and promotes environmental fitness of an enteric virus. Cell Host Microbe. 2014; 15(1):36-46. Epub

2014/01/21. https://doi.org/10.1016/j.chom.2013.12.004 PMID: 24439896.

6. den Boon JA, Ahlquist P. Organelle-like membrane compartmentalization of positive-strand RNA virus

replication   factories.   Annu   Rev   Microbiol.   2010;   64:241-56.   Epub   2010/09/10.

https://doi.org/10.1146/annurev.micro.112408.134012 PMID: 20825348.

7. Belov GA, Sztul E. Rewiring of cellular membrane homeostasis by picornaviruses. J Virol. 2014;

88(17):9478-89. Epub 2014/06/13. https://doi.org/10.1128/jvi.00922-14 PMID: 24920802.

8. Kallman F, Williams RC, Dulbecco R, Vogt M. Fine structure of changes produced in cultured cells

sampled at specified intervals during a single growth cycle of polio virus. J Biophys Biochem Cytol. 1958;

4(3):301-8. Epub 1958/05/25. PMID: 13549502.

9. Belov GA, Nair V, Hansen BT, Hoyt FH, Fischer ER, Ehrenfeld E. Complex dynamic development of

poliovirus membranous replication complexes. J Virol. 2012; 86(1):302-12. Epub 2011/11/11.

https://doi.org/10.1128/jvi.05937-11 PMID: 22072780.

10. Bienz K, Egger D, Pfister T, Troxler M. Structural and functional characterization of the poliovirus

replication complex. J Virol. 1992; 66(5):2740-7. Epub 1992/05/01. PMID: 1313898.

11. Bienz K, Egger D, Rasser Y, Bossart W. Kinetics and location of poliovirus macromolecular synthesis in

correlation to virus-induced cytopathology. Virology. 1980; 100(2):390-9. Epub 1980/01/30. PMID:

6243433.

12. Schlegel A, Giddings TH, Jr., Ladinsky MS, Kirkegaard K. Cellular origin and ultrastructure of
membranes induced during poliovirus infection. J Virol. 1996; 70(10):6576-88. Epub 1996/10/01. PMID:
8794292.

13. Troxler M, Egger D, Pfister T, Bienz K. Intracellular localization of poliovirus RNA by in situ hybridization
at the ultrastructural level using single-stranded riboprobes. Virology. 1992; 191(2):687-97. Epub
1992/12/01. PMID: 1333118.

14. Bienz K, Egger D, Troxler M, Pasamontes L. Structural organization of poliovirus RNA replication is
mediated by viral proteins of the P2 genomic region. J Virol. 1990; 64(3):1156-63. Epub 1990/03/01.
PMID: 2154600.

15. Bienz K, Egger D, Pasamontes L. Association of polioviral proteins of the P2 genomic region with the
viral replication complex and virus-induced membrane synthesis as visualized by electron microscopic
immunocytochemistry and autoradiography. Virology. 1987; 160(1):220-6. Epub 1987/09/01. PMID:
2820130.

16. Bienz K, Egger D, Rasser Y, Bossart W. Intracellular distribution of poliovirus proteins and the induction
of virus-specific cytoplasmic structures. Virology. 1983; 131(1):39-48. Epub 1983/11/01. PMID: 6316654.

17. Hsu NY, Ilnytska O, Belov G, Santiana M, Chen YH, Takvorian PM, et al. Viral reorganization of the
secretory pathway generates distinct organelles for RNA replication. Cell. 2010; 141(5):799-811. Epub
2010/06/01. https://doi.org/10.1016/j.cell.2010.03.050 PMID: 20510927.

18. Racaniello VR. Picornaviridae: the viruses and their replication. In: Knipe DM, Howley PM, editors.
Fields virology. 6th ed. Philadelphia, PA: Lippincott Williams & Wilkins. p. 453-89

19. Cho MW, Teterina N, Egger D, Bienz K, Ehrenfeld E. Membrane rearrangement and vesicle induction
by recombinant poliovirus 2C and 2BC in human cells. Virology. 1994; 202(1):129-45. Epub 1994/07/01.
https://doi.org/10.1006/viro.1994.1329 PMID: 8009827.

20. Teterina NL, Gorbalenya AE, Egger D, Bienz K, Ehrenfeld E. Poliovirus 2C protein determinants of
membrane binding and rearrangements in mammalian cells. J Virol. 1997; 71(12):8962-72. Epub
1997/11/26. PMID: 9371552.

21. Suhy DA, Giddings TH, Jr., Kirkegaard K. Remodeling the endoplasmic reticulum by poliovirus infection
and by individual viral proteins: an autophagy-like origin for virus-induced vesicles. J Virol. 2000;
74(19):8953-65. Epub 2000/09/12. PMID: 10982339.

22. Ventoso I, MacMillan SE, Hershey JW, Carrasco L. Poliovirus 2A proteinase cleaves directly the eIF-4G
subunit of eIF-4F complex. FEBS Lett. 1998; 435(1):79-83. Epub 1998/10/02. PMID: 9755863.

23. Haghighat A, Svitkin Y, Novoa I, Kuechler E, Skern T, Sonenberg N. The eIF4G-eIF4E complex is the

target for direct cleavage by the rhinovirus 2A proteinase. J Virol. 1996; 70(12):8444-50. Epub 1996/12/01.

PMID: 8970966.

24. Li JP, Baltimore D. An intragenic revertant of a poliovirus 2C mutant has an uncoating defect. J Virol.

1990; 64(3):1102-7. Epub 1990/03/01. PMID: 2154595.

25. Li JP, Baltimore D. Isolation of poliovirus 2C mutants defective in viral RNA synthesis. J Virol. 1988;

62(11):4016-21. Epub 1988/11/01. PMID: 2845120.

26. Vance LM, Moscufo N, Chow M, Heinz BA. Poliovirus 2C region functions during encapsidation of viral

RNA. J Virol. 1997; 71(11):8759-65. Epub 1997/10/29. PMID: 9343235.

27. Liu Y, Wang C, Mueller S, Paul AV, Wimmer E, Jiang P. Direct interaction between two viral proteins,

the nonstructural protein 2C and the capsid protein VP3, is required for enterovirus morphogenesis. PLoS

Pathog. 2010; 6(8):e1001066. Epub 2010/09/25. https://doi.org/10.1371/journal.ppat.1001066 PMID:

20865167.

28. Asare E, Mugavero J, Jiang P, Wimmer E, Paul AV. A Single Amino Acid Substitution in Poliovirus

Nonstructural Protein 2CATPase Causes Conditional Defects in Encapsidation and Uncoating. J Virol. 2016;

90(14):6174-86. Epub 2016/04/15. https://doi.org/10.1128/jvi.02877-15 PMID: 27076638.

29. Pincus SE, Diamond DC, Emini EA, Wimmer E. Guanidine-selected mutants of poliovirus: mapping of

point mutations to polypeptide 2C. J Virol. 1986; 57(2):638-46. Epub 1986/02/01. PMID: 3003395.

30. Baltera RF, Jr., Tershak DR. Guanidine-resistant mutants of poliovirus have distinct mutations in

peptide 2C. J Virol. 1989; 63(10):4441-4. Epub 1989/10/01. PMID: 2550675.

31. Hope DA, Diamond SE, Kirkegaard K. Genetic dissection of interaction between poliovirus 3D

polymerase and viral protein 3AB. J Virol. 1997; 71(12):9490-8. Epub 1997/11/26. PMID: 9371611.

32. Giachetti C, Hwang SS, Semler BL. cis-acting lesions targeted to the hydrophobic domain of a poliovirus

membrane protein involved in RNA replication. J Virol. 1992; 66(10):6045-57. Epub 1992/10/01. PMID:

1326655.

33. Andino R, Rieckhof GE, Achacoso PL, Baltimore D. Poliovirus RNA synthesis utilizes an RNP complex

formed around the 5'-end of viral RNA. Embo j. 1993; 12(9):3587-98. Epub 1993/09/01. PMID: 8253083.

34. Gamarnik AV, Andino R. Switch from translation to RNA replication in a positive-stranded RNA virus.

Genes Dev. 1998; 12(15):2293-304. Epub 1998/08/08. PMID: 9694795.

35. Parsley TB, Towner JS, Blyn LB, Ehrenfeld E, Semler BL. Poly (rC) binding protein 2 forms a ternary

complex with the 5'-terminal sequences of poliovirus RNA and the viral 3CD proteinase. Rna. 1997;

3(10):1124-34. Epub 1997/11/05. PMID: 9326487.

36. Paul AV, Rieder E, Kim DW, van Boom JH, Wimmer E. Identification of an RNA hairpin in poliovirus

RNA that serves as the primary template in the in vitro uridylylation of VPg. J Virol. 2000; 74(22):10359-

70. Epub 2000/10/24. PMID: 11044080.

37. Rieder E, Paul AV, Kim DW, van Boom JH, Wimmer E. Genetic and biochemical studies of poliovirus

cis-acting replication element cre in relation to VPg uridylylation. J Virol. 2000; 74(22):10371-80. Epub

2000/10/24. PMID: 11044081.

38. Toyoda H, Nicklin MJ, Murray MG, Anderson CW, Dunn JJ, Studier FW, et al. A second virus-encoded

proteinase involved in proteolytic processing of poliovirus polyprotein. Cell. 1986; 45(5):761-70. Epub

1986/06/06. PMID: 3011278.

39. Nicklin MJ, Harris KS, Pallai PV, Wimmer E. Poliovirus proteinase 3C: large-scale expression,

purification, and specific cleavage activity on natural and synthetic substrates in vitro. J Virol. 1988;

62(12):4586-93. Epub 1988/12/01. PMID: 2846872.

40. Ypma-Wong MF, Semler BL. In vitro molecular genetics as a tool for determining the differential

cleavage specificities of the poliovirus 3C proteinase. Nucleic Acids Res. 1987; 15(5):2069-88. Epub

1987/03/11. PMID: 3031587.

41. Lawson MA, Semler BL. Alternate poliovirus nonstructural protein processing cascades generated by

primary sites of 3C proteinase cleavage. Virology. 1992; 191(1):309-20. Epub 1992/11/01. PMID:

1329322.

42. Pathak HB, Oh HS, Goodfellow IG, Arnold JJ, Cameron CE. Picornavirus genome replication: roles of

precursor proteins and rate-limiting steps in oriI-dependent VPg uridylylation. J Biol Chem. 2008;

283(45):30677-88. Epub 2008/09/10. https://doi.org/10.1074/jbc.M806101200 PMID: 18779320.

43. Oh HS, Pathak HB, Goodfellow IG, Arnold JJ, Cameron CE. Insight into poliovirus genome replication

and encapsidation obtained from studies of 3B-3C cleavage site mutants. J Virol. 2009; 83(18):9370-87.

Epub 2009/07/10. https://doi.org/10.1128/jvi.02076-08 PMID: 19587035.

44. Towner JS, Mazanet MM, Semler BL. Rescue of defective poliovirus RNA replication by 3AB-containing

precursor polyproteins. J Virol. 1998; 72(9):7191-200. Epub 1998/08/08. PMID: 9696813.

45. Jurgens C, Flanegan JB. Initiation of poliovirus negative-strand RNA synthesis requires precursor forms

of p2 proteins. J Virol. 2003; 77(2):1075-83. Epub 2002/12/28. PMID: 12502823.

46. Novak JE, Kirkegaard K. Coupling between genome translation and replication in an RNA virus. Genes

Dev. 1994; 8(14):1726-37. Epub 1994/07/15. PMID: 7958852.

47. Wessels E, Duijsings D, Niu TK, Neumann S, Oorschot VM, de Lange F, et al. A viral protein that blocks

Arf1-mediated COP-I assembly by inhibiting the guanine nucleotide exchange factor GBF1. Dev Cell. 2006;

11(2):191-201. Epub 2006/08/08. https://doi.org/10.1016/j.devcel.2006.06.005 PMID: 16890159.

48. Wessels E, Duijsings D, Lanke KH, van Dooren SH, Jackson CL, Melchers WJ, et al. Effects of

picornavirus 3A Proteins on Protein Transport and GBF1-dependent COP-I recruitment. J Virol. 2006;

80(23):11852-60. Epub 2006/09/29. https://doi.org/10.1128/jvi.01225-06 PMID: 17005635.

49. Belov GA, Altan-Bonnet N, Kovtunovych G, Jackson CL, Lippincott-Schwartz J, Ehrenfeld E. Hijacking

components of the cellular secretory pathway for replication of poliovirus RNA. J Virol. 2007; 81(2):558-

67. Epub 2006/11/03. https://doi.org/10.1128/jvi.01820-06 PMID: 17079330.

50. Doedens JR, Kirkegaard K. Inhibition of cellular protein secretion by poliovirus proteins 2B and 3A.

Embo j. 1995; 14(5):894-907. Epub 1995/03/01. PMID: 7889939.

51. Doedens JR, Giddings TH, Jr., Kirkegaard K. Inhibition of endoplasmic reticulum-to-Golgi traffic by

poliovirus protein 3A: genetic and ultrastructural analysis. J Virol. 1997; 71(12):9054-64. Epub

1997/11/26. PMID: 9371562.

52. Choe SS, Dodd DA, Kirkegaard K. Inhibition of cellular protein secretion by picornaviral 3A proteins.

Virology. 2005; 337(1):18-29. Epub 2005/05/26. https://doi.org/10.1016/j.virol.2005.03.036 PMID:

15914217.

53. Wang J, Ptacek JB, Kirkegaard K, Bullitt E. Double-membraned liposomes sculpted by poliovirus 3AB

protein. J Biol Chem. 2013; 288(38):27287-98. Epub 2013/08/03.

https://doi.org/10.1074/jbc.M113.498899 PMID: 23908350.

54. Belov GA, Habbersett C, Franco D, Ehrenfeld E. Activation of cellular Arf GTPases by poliovirus protein

3CD correlates with virus replication. J Virol. 2007; 81(17):9259-67. Epub 2007/06/15.

https://doi.org/10.1128/jvi.00840-07 PMID: 17567696.

55. Belov GA, Fogg MH, Ehrenfeld E. Poliovirus proteins induce membrane association of GTPase ADP-

ribosylation factor. J Virol. 2005; 79(11):7207-16. Epub 2005/05/14.

https://doi.org/10.1128/jvi.79.11.7207-7216.2005 PMID: 15890959.

56. Franco D, Pathak HB, Cameron CE, Rombaut B, Wimmer E, Paul AV. Stimulation of poliovirus synthesis

in a HeLa cell-free in vitro translation-RNA replication system by viral protein 3CDpro. J Virol. 2005;

79(10):6358-67. Epub 2005/04/29. https://doi.org/10.1128/jvi.79.10.6358-6367.2005 PMID: 15858019.

57. Franco D, Pathak HB, Cameron CE, Rombaut B, Wimmer E, Paul AV. Stimulation of poliovirus RNA

synthesis and virus maturation in a HeLa cell-free in vitro translation-RNA replication system by viral

protein 3CDpro. Virol J. 2005; 2:86. Epub 2005/11/23. https://doi.org/10.1186/1743-422x-2-86 PMID:

16300678.

 FIGURE LEGENDS

Figure 1**. Poliovirus genome organization and P3-polyprotein processing. (A**) Schematic of the poliovirus

genome. The 7.5 kb long genome consists of a 5'-nontranslated region (NTR), an open reading frame, a

3'-NTR and a poly(rA) tail. The 5'-end of the genome is covalently linked to a peptide (VPg) encoded by

the 3B gene. The 5'-NTR contains a cis-acting replication element (CRE) termed oriL or cloverleaf followed

by a type II internal ribosome entry site (IRES). Two additional CREs exist: oriI and oriR, located within the

2C gene and 3'-NTR, respectively. IRES mediated translation yields a single polyprotein comprise of three

functional domains of structural (P1) and the non-structural (P2 and P3) proteins. (**B**) Processing of the P3

region by WT and mutant PVs. Two pathways, major and minor, exist for P3 processing. Major pathway

use for WT PV is shown and produces only 3AB and 3CD because of cleavage at Gln-Gly junction between

3B and 3C. EG PV changes the 3B-3C junction to Glu-Gly, producing the normal products at a reduced

rate. GG PV changes the 3B-3C junction to Gly-Gly, which is uncleavable, inducing aberrant processing

and producing 3ABC and 3D instead 3AB and 3CD.

Figure 2**. EG PV exhibits a delay in the onset of RNA synthesis. (A**) Kinetics of replication of subgenomic

replicon by WT (■) and EG (□) monitored indirectly by luciferase activity. HeLa cells were transfected with

in vitro transcribed replicon RNA, placed at 37 °C and luciferase activity (RLU/μg) measured at the

indicated times post-transfection. A control for translation of transfected replicon RNA was performed by

performing the experiment above in the presence of 3 Mm GuHCl (●). (**B**) Kinetics of replication of

subgenomic replicon by WT (●) and EG (○) monitored by Northern blotting. HeLa cells were transfected

with *in vitro* transcribed replicon RNA, placed at 37 °C, and at the indicated times post-transfection, cells

were harvested for total RNA isolation. Total RNA was separated on a 0.6% agarose gel containing 0.8 M

formaldehyde, transferred to nylon membrane and hybridized with a ^{32}P-labeled DNA probe. (**C**) Image of

a representative blot visualized by phosphorimaging. *In vitro* transcribed RNA (*****) is shown as reference.

Figure 3**. Kinetics of genome replication precedes the kinetics of vesicular cluster formation for EG PV.**

(**A**) Kinetics of RNA synthesis (○) and virus production (●) by EG PV. HeLa cells were infected with EG PV

at an MOI of 10, placed at 37 °C, and at the indicated times post-infection, total RNA was isolated and

subjected to either Northern blotting or assayed for virus production by standard plaque assay. (**B**) Image

of a representative blot visualized by phosphorimaging. **(C)** Kinetics of formation of virus-induced vesicular

clusters by transmission electron microscope (TEM). HeLa cells were infected with EG PV at MOI of 10,

placed at 37 °C, and at the indicated times post-infection, infected cells were fixed and visualized by TEM,

bar = 1 μm.

Figure 4. **Kinetics of genome replication precedes the kinetics of vesicular cluster formation for WT PV.**

(**A**) Kinetics of RNA synthesis (○) and virus production (●) by WT PV. HeLa cells were infected with WT PV

at an MOI of 10, placed at 37 °C, and at the indicated times post-infection, total RNA was isolated and

subjected to either Northern blotting or assayed for virus production by standard plaque assay. (**B**) Image

of a representative blot visualized by phosphorimaging. (**C**) Kinetics of formation of WT PV-induced

vesicular cluster formation was visualized by TEM. HeLa cells were infected with WT PV at an MOI of 10,

placed at 37 °C, and at the indicated times post-infection, the infected cells were fixed and visualized by

TEM, bar = 1 μm.

Figure 5. **EG PV exhibits delayed induction and redistribution of PI4P. (A)** Immunostaining of PI4P in

mock-infected HeLa cells. **(B)** Time-course of PI4P-staining in HeLa cells infected with WT or EG PV. HeLa

cells were infected with WT or EG virus at an MOI of 10, fixed at indicated times post-infection, and

immunostained for PI4P. **(C)** Impact of GuHCl on PI4P induction by WT and EG PVs. HeLa cells were either

incubated with PBS or infected with WT or EG virus (MOI 10) in presence of 3 mM GuHCl and

immunostained. In all cases, PI4P was stained using anti-PI4P antibody (red) and nuclei were stained with

DAPI (blue).

Figure 6**. GG PV induces and redistributes PI4P in spite of impaired formation of vesicular clusters.** (**A**)

Kinetics of RNA synthesis by WT (●) and GG (Δ) subgenomic replicon RNA. HeLa cells were co-transfected

with two different replicon RNAs, luciferase replicon (2 μg) and EGFP replicon (4 μg), placed at 34 °C and

at the indicated times post-transfection, luciferase activity was measured. (**B**) Cell sorting to isolate PV

replicon-positive cells. WT replicon RNA-transfected cells were 61% positive in pre-sort cells (top-left) and

98% positive in post-sort cells (bottom-left). GG replicon RNA transfected cells were 18% positive in pre-

sort cells (top-right) and 94% positive in post-sort cells (bottom-right). (**C**) WT and GG PV-induced

membranes visualized by TEM. HeLa cells were transfected with either WT or GG replicon RNA, placed at

34 °C for 5 h or 14 h, respectively, at which time cells were fixed and visualized by TEM. Bar =1 μm. **(D)**

Kinetics of PI4P induction and redistribution by the GG PV subgenomic replicon. HeLa cells were

transfected with replicon RNA expressing EGFP and samples were fixed at the indicated time post-

transfection and subjected to IFM using anti-PI4P antibody (red) and nuclei were stained with DAPI (blue).

Figure 7**. Multiplicity of infection exaggerates size and quantity of WT PV-induced vesicular clusters.**

Transmission electron micrographs (TEM) of HeLa cells infected with WT PV at an MOI of 10 **(A),** 50 **(B)** or

100 **(C)**. HeLa cells were infected with WT PV expressing EGFP at an MOI of 10, 50 or 100 by incubating

the monolayer with WT virus for 30 min at room temperature, followed by removal of the virus and

washing of the monolayer with PBS. Prewarmed media was then added to the monolayer and incubated

at 37 °C. Samples were fixed 3 h post-infection and visualized by TEM. Bar = 1 µm.

A

oriL

IRES

VPg (3B)

oriI

| VP0 | VP3 | VP1 | 2A | 2B | 2C | 3A | 3B | 3C | 3D |

oriR

poly A

5' NTR   P1   P2   P3   3' NTR

B

Gln-Gly

| 3A | 3B | 3C | 3D |

WT   EG   GG

| 3A | 3B | | 3C | 3D |

| 3A | 3B | 3C | | 3D |

Slower Kinetics

| 3A | 3B | | 3C | 3D |

*Reference *in vitro* transcribed RNA

A. RNA (volume) / Virus titer (PFU/ml) vs Time post-infection (h); ○ RNA synthesis, ● Virus production. B. Time course (0–8 h). C. Electron micrographs: UN, 1h, 2h, 4h, 5h, 7h.