The Genomic History of Southeastern Europe

Iain Mathieson[†] (1), Songül Alpaslan Roodenberg (1), Cosimo Posth (2,3), Anna Szécsényi-Nagy (4), Nadin Rohland (1), Swapan Mallick (1,5), Iñigo Olalde (1), Nasreen Broomandkhoshbacht (1,5), Francesca Candilio (6), Olivia Cheronet (6,7), Daniel Fernandes (6,8), Matthew Ferry (1,5), Beatriz Gamarra (6), Gloria González Fortes (9), Wolfgang Haak (2,10), Eadaoin Harney (1,5), Eppie Jones (11,12), Denise Keating (6), Ben Krause-Kyora (2), Isil Kucukkalipci (3), Megan Michel (1,5), Alissa Mittnik (2,3), Kathrin Nägele (2), Mario Novak (6,13), Jonas Oppenheimer (1,5), Nick Patterson (14), Saskia Pfrengle (3), Kendra Sirak (6,15), Kristin Stewardson (1,5), Stefania Vai (16), Stefan Alexandrov (17), Kurt W. Alt (18,19,20), Radian Andreescu (21), Dragana Antonović (22), Abigail Ash (6), Nadezhda Atanassova (23), Krum Bacvarov (17), Mende Balázs Gusztáv (4), Hervé Bocherens (24,25), Michael Bolus (26), Adina Boroneanţ (27), Yavor Boyadzhiev (17), Alicja Budnik (28), Josip Burmaz (29), Stefan Chohadzhiev (30), Nicholas J. Conard (31,25), Richard Cottiaux (32), Maja Čuka (33), Christophe Cupillard (34,35), Dorothée G. Drucker (25), Nedko Elenski (36), Michael Francken (37), Borislava Galabova (38), Georgi Ganetovski (39), Bernard Gély (40), Tamás Hajdu (41), Veneta Handzhyiska (42), Katerina Harvati (37,25), Thomas Higham (43), Stanislav Iliev (44), Ivor Janković (13,45), Ivor Karavanić (46,45), Douglas J. Kennett (47), Darko Komšo (33), Alexandra Kozak (48), Damian Labuda (49), Martina Lari (16), Catalin Lazar (50,51), Maleen Leppek (52), Krassimir Leshtakov (42), Domenico Lo Vetro (53,54), Dženi Los (29), Ivaylo Lozanov (42), Maria Malina (26), Fabio Martini (53,54), Kath McSweeney (55), Harald Meller (20), Marko Menđušić (56), Pavel Mirea (57), Vyacheslav Moiseyev (58), Vanya Petrova (42), T. Douglas Price (59), Angela Simalcsik (60), Luca Sineo (61), Mario Šlaus (62), Vladimir Slavchev (63), Petar Stanev (36), Andrej Starović (64), Tamás Szeniczey (41), Sahra Talamo (65), Maria Teschler-Nicola (66,7), Corinne Thevenet (67), Ivan Valchev (42), Frédérique Valentin (68), Sergey Vasilyev (69), Fanica Veljanovska (70), Svetlana Venelinova (71), Elizaveta Veselovskaya (69), Bence Viola (72,73), Cristian Virag (74), Joško Zaninović (75), Steve Zäuner (76), Philipp W. Stockhammer (52,2), Giulio Catalano (61), Raiko Krauß (77), David Caramelli (16), Gunita Zariņa (78), Bisserka Gaydarska (79), Malcolm Lillie (80), Alexey G. Nikitin (81), Inna Potekhina (48), Anastasia Papathanasiou (82), Dušan Borić (83), Clive Bonsall (55), Johannes Krause (2,3), Ron Pinhasi* (6,7), David Reich* (1,14,5)

* These authors contributed equally to the manuscript
[†] Present address; Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia PA 19104, USA
Correspondence to I.M. (mathi@upenn.edu) or D.R. (reich@genetics.med.harvard.edu) or R.P. (ron.pinhasi@ucd.ie)

(1) Department of Genetics, Harvard Medical School, Boston 02115 MA USA (2) Department of Archaeogenetics, Max Planck Institute for the Science of Human History, 07745 Jena, Germany (3) Institute for Archaeological Sciences, University of Tuebingen, Germany (4) Laboratory of Archaeogenetics, Institute of Archaeology, Research Centre for the Humanities, Hungarian Academy of Sciences, H-1097 Budapest, Hungary (5) Howard Hughes Medical Institute, Harvard Medical School, Boston 02115 MA USA (6) Earth Institute and School of Archaeology, University College Dublin, Belfield, Dublin 4, Republic of Ireland (7) Department of Anthropology, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria (8) CIAS, Department of Life Sciences, University of Coimbra, 3000-456 Coimbra, Portugal (9) Department of Life Sciences and Biotechnology, University of Ferrara, Via L. Borsari 46. Ferrara 44100 Italy (10) Australian Centre for Ancient DNA, School of Biological Sciences, The University of Adelaide, SA-5005 Adelaide, Australia (11) Smurfit Institute of Genetics, Trinity College Dublin, Dublin 2, Ireland (12) Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK (13) Institute for Anthropological Research, Ljudevita Gaja 32, 10000 Zagreb, Croatia (14) Broad Institute of Harvard and MIT, Cambridge MA (15) Department of Anthropology, Emory University, Atlanta, Georgia 30322, USA (16) Dipartimento di Biologia, Università di Firenze, 50122 Florence, Italy (17) National Institute of Archaeology and Museum, Bulgarian Academy of Sciences, 2 Saborna Str., BG-1000 Sofia, Bulgaria (18) Danube Private University, A-3500 Krems, Austria (19) Department of Biomedical Engineering and

Integrative Prehistory and Archaeological Science, CH-4123 Basel-Allschwil, Switzerland (20) State Office for Heritage Management and Archaeology Saxony-Anhalt and State Museum of Prehistory, D-06114 Halle, Germany (21) Romanian National History Museum, Bucharest, Romania (22) Institute of Archaeology, Belgrade, Serbia (23) Institute of Experimental Morphology, Pathology and Anthropology with Museum, Bulgarian Academy of Sciences, Sofia, Bulgaria (24) Department of Geosciences, Biogeology, Universität Tübingen, Hölderlinstr. 12, 72074 Tübingen, Germany (25) Senckenberg Centre for Human Evolution and Palaeoenvironment, University of Tuebingen, 72072 Tuebingen, Germany (26) Heidelberg Academy of Sciences and Humanities, Research Center ''The Role of Culture in Early Expansions of Humans'' at the University of Tuebingen, Rümelinstraße 23, 72070 Tuebingen, Germany (27) 'Vasile Pârvan' Institute of Archaeology, Romanian Academy (28) Human Biology Department, Cardinal Stefan Wyszyński University, Warsaw, Poland (29) KADUCEJ d.o.o Papandopulova 27, 21000 Split, Croatia (30) St. Cyril and Methodius University, Veliko Turnovo, Bulgaria (31) Department of Early Prehistory and Quaternary Ecology, University of Tuebingen, Schloss Hohentübingen, 72070 Tuebingen, Germany (32) INRAP/UMR 8215 Trajectoires, 21 Alleé de l'Université, 92023 Nanterre, France (33) Archaeological Museum of Istria, Carrarina 3, 52100 Pula, Croatia (34) Service Régional de l'Archéologie de Bourgogne-Franche-Comté, 7 rue Charles Nodier, 25043 Besançon Cedex, France (35) Laboratoire Chronoenvironnement, UMR 6249 du CNRS, UFR des Sciences et Techniques, 16 route de Gray, 25030 Besançon Cedex, France (36) Regional Museum of History Veliko Tarnovo, Veliko Tarnovo, Bulgaria (37) Institute for Archaeological Sciences, Paleoanthropology, University of Tuebingen, Rümelinstraße 23, 72070 Tuebingen, Germany (38) Laboratory for human bio-archaeology, Bulgaria, 1202 Sofia, 42, George Washington str (39) Regional Museum of History, Vratsa, Bulgaria (40) DRAC Auvergne - Rhône Alpes, Ministère de la Culture, Le Grenier d'abondance 6, quai Saint Vincent 69283 LYON cedex 01 (41) Eötvös Loránd University, Faculty of Science, Institute of Biology, Department of Biological Anthropology, H-1117 Pázmány Péter sétány 1/c. Budapest, Hungary (42) Department of Archaeology, Sofia University St. Kliment Ohridski, Bulgaria (43) Oxford Radiocarbon Accelerator Unit, Research Laboratory for Archaeology and the History of Art, University of Oxford, Dyson Perrins Building, South Parks Road, OX1 3QY Oxford, UK (44) Regional Museum of History, Haskovo, Bulgaria (45) Department of Anthropology, University of Wyoming, 1000 E. University Avenue, Laramie, WY 82071, USA (46) Department of Archaeology, Faculty of Humanities and Social Sciences, University of Zagreb, Ivana Lučića 3, 10000 Zagreb, Croatia (47) Department of Anthropology and Institutes for Energy and the Environment, Pennsylvania State University, University Park, PA 16802 (48) Department of Bioarchaeology, Institute of Archaeology, National Academy of Sciences of Ukraine (49) CHU Sainte-Justine Research Center, Pediatric Department, Université de Montréal, Montreal, PQ, Canada, H3T 1C5 (50) National History Museum of Romania, Calea Victoriei, no. 12, 030026, Bucharest, Romania (51) University of Bucharest, Mihail Kogalniceanu 36-46, 50107, Bucharest, Romania (52) Institute for Pre- and Protohistoric Archaeology and the Archaeology of the Roman Provinces, Ludwig-Maximilians-University, Schellingstr. 12, 80799 Munich, Germany (53) Dipartimento SAGAS - Sezione di Archeologia e Antico Oriente, Università degli Studi di Firenze, 50122 Florence, Italy (54) Museo e Istituto fiorentino di Preistoria, 50122 Florence, Italy (55) School of History, Classics and Archaeology, University of Edinburgh, Edinburgh EH8 9AG, United Kingdom (56) Conservation Department in Šibenik, Ministry of Culture of the Republic of Croatia, Jurja Čulinovića 1, 22000 Šibenik, Croatia (57) Teleorman County Museum, str. 1848, no. 1, 140033 Alexandria, Romania (58) Peter the Great Museum of Anthropology and Ethnography (Kunstkamera) RAS, 199034 St. Petersburg, Russia (59) University of Wisconsin, Madison WI, USA (60) Olga Necrasov Centre for Anthropological Research, Romanian Academy – Iaşi Branch, Theodor Codrescu St. 2, P.C. 700481, Iaşi, Romania (61) Dipartimento di Scienze e tecnologie biologiche, chimiche e farmaceutiche, Lab. of Anthropology, Università degli studi di Palermo, Italy (62) Anthropological Center, Croatian Academy of Sciences and Arts, 10000 Zagreb, Croatia (63) Regional Historical Museum Varna, Maria Luiza Blvd. 41, BG-9000 Varna, Bulgaria (64) National Museum in Belgrade, 1a Republic sq., Belgrade, Serbia (65) Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany (66) Department of Anthropology, Natural History Museum Vienna, 1010 Vienna, Austria (67) INRAP/UMR 8215 Trajectoires, 21 Allée de l'Université, 92023 Nanterre, France (68) CNRS/UMR 7041 ArScAn MAE, 21 Allée de l'Université, 92023 Nanterre, France (69) Institute of Ethnology and Anthropology, Russian Academy of Sciences, Leninsky Pr., 32a, Moscow, 119991, Russia (70) Archaeological Museum of Macedonia, Skopje (71) Regional museum of history, Shumen, Bulgaria (72) Department of Anthropology, University of Toronto, Toronto, Ontario, M5S 2S2, Canada (73) Institute of Archaeology & Ethnography, Siberian Branch, Russian Academy of Sciences, Lavrentiev Pr. 17, Novosibirsk 630090, Russia (74) Satu Mare County Museum Archaeology Department,V. Lucaciu, nr.21, Satu Mare, Romania (75) Municipal Museum Drniš, Domovinskog rata 54, 22320 Drniš, Croatia (76) anthropol - Anthropologieservice, Schadenweilerstraße 80, 72379 Hechingen, Germany (77) Institute for Prehistory, Early History and Medieval Archaeology, University of Tuebingen, Germany (78) Institute of Latvian History, University of Latvia, Kalpaka Bulvāris 4, Rīga 1050, Latvia (79) Department of Archaeology, Durham University, UK (80) School of Environmental Sciences: Geography, University of Hull, Hull HU6 7RX, UK (81) Department of Biology, Grand Valley State University, Allendale, Michigan, USA (82) Ephorate of Paleoanthropology and Speleology, Athens, Greece (83) The Italian Academy for Advanced Studies in America, Columbia University, 1161 Amsterdam Avenue, New York, NY 10027, USA.

Abstract

Farming was first introduced to southeastern Europe in the mid-7th millennium BCE – brought by migrants from Anatolia who settled in the region before spreading throughout Europe. To clarify the dynamics of the interaction between the first farmers and indigenous hunter-gatherers where they first met, we analyze genome-wide ancient DNA data from 223 individuals who lived in southeastern Europe and surrounding regions between 12,000 and 500 BCE. We document previously uncharacterized genetic structure, showing a West-East cline of ancestry in hunter-gatherers, and show that some Aegean farmers had ancestry from a different lineage than the northwestern Anatolian lineage that formed the overwhelming ancestry of other European farmers. We show that the first farmers of northern and western Europe passed through southeastern Europe with limited admixture with local hunter-gatherers, but that some groups mixed extensively, with relatively sex-balanced admixture compared to the male-biased hunter-gatherer admixture that prevailed later in the North and West. Southeastern Europe continued to be a nexus between East and West after farming arrived, with intermittent genetic contact from the Steppe up to 2,000 years before the migration that replaced much of northern Europe's population.

Introduction

The southeastern quadrant of Europe was the beachhead in the spread of agriculture from its source in the Fertile Crescent of southwestern Asia. After the first appearance of agriculture in the mid-7th millennium BCE,[1,2] farming spread westward via a Mediterranean and northwestward via a Danubian route, and was established in both Iberia and Central Europe by 5600 BCE.[3,4] Ancient DNA studies have shown that the spread of farming across Europe was accompanied by a massive movement of people[5-8] closely related to the farmers of northwestern Anatolia[9-11] but nearly all the ancient DNA from Europe's first farmers is from central and western Europe, with only three individuals reported from the southeast.[9] In the millennia following the establishment of agriculture in the Balkan Peninsula, a series of complex societies formed, culminating in sites such as the mid-5th millennium BCE necropolis at Varna, which has some of the earliest evidence of extreme inequality in wealth, with one individual (grave 43) from whom we extracted DNA buried with more gold than is known from any earlier site. By the end of the 6th millennium BCE, agriculture had reached eastern Europe, in the form of the Cucuteni-Trypillian complex in the area of present-day Moldova, Romania and Ukraine, including "mega-sites" that housed hundreds, perhaps thousands, of people.[12] After around 4000 BCE, these settlements were largely abandoned, and

archaeological evidence documents cultural contacts with peoples of the Eurasian steppe.[13]

However, the population movements that accompanied these events have been unknown due

to the lack of ancient DNA.

Results

We generated genome-wide data from 223 ancient humans (214 reported for the first time),

from the Balkan Peninsula, the Carpathian Basin, the North Pontic Steppe and neighboring

regions, dated to 12,000-500 BCE (Figure 1A, Supplementary Information Table 1,

Supplementary Information Note 1). We extracted DNA from skeletal remains in dedicated

clean rooms, built DNA libraries and enriched for DNA fragments overlapping 1.24 million

single nucleotide polymorphisms (SNPs), then sequenced the product and restricted to

libraries with evidence of authentic ancient DNA.[7,10,14] We filtered out individuals with fewer

than 15,000 SNPs covered by at least one sequence, that had unexpected ancestry for their

archaeological context and were not directly dated. We report, but do not analyze, nine

individuals that were first-degree relatives of others in the dataset, resulting in an analysis

dataset of 214 individuals. We analyzed these data together with 274 previously reported

ancient individuals,[9-11,15-27] 799 present-day individuals genotyped on the Illumina "Human

Origins" array,[23] and 300 high coverage genomes from the Simons Genome Diversity Project

(SGDP).[28] We used principal component analysis (PCA; Figure 1B, Extended Data Figure 1),

supervised and unsupervised ADMIXTURE (Figure 1D, Extended Data Figure 2),[29] *D*-

statistics, *qpAdm* and *qpGraph,*[30] along with archaeological and chronological information to

cluster the individuals into populations and investigate the relationships among them.

We described the individuals in our dataset in terms of their genetic relatedness to a

hypothesized set of ancestral populations, which we refer to as their genetic ancestry. It has

previously been shown that the great majority of European ancestry derives from three

distinct sources.[23] First, there is "hunter-gatherer-related" ancestry that is more closely related

to Mesolithic hunter-gatherers from Europe than to any other population, and that can be

further subdivided into "Eastern" (EHG) and "Western" (WHG) hunter-gatherer-related

ancestry.[7] Second, there is "NW Anatolian Neolithic-related" ancestry related to the

Neolithic farmers of northwest Anatolia and tightly linked to the appearance of agriculture.[9,10]

The third source, "steppe-related" ancestry, appears in Western Europe during the Late

Neolithic to Bronze Age transition and is ultimately derived from a population related to

Yamnaya steppe pastoralists.[7,15] Steppe-related ancestry itself can be modeled as a mixture of

EHG-related ancestry, and ancestry related to Upper Palaeolithic hunter-gatherers of the

Caucasus (CHG) and the first farmers of northern Iran.[19,21,22]

Hunter-Gatherer substructure and transitions

Of the 214 new individuals we report, 114 from Paleolithic, Mesolithic and eastern European

Neolithic contexts have almost entirely hunter-gatherer-related ancestry (in eastern Europe,

unlike western Europe, "Neolithic" refers to the presence of pottery,[31-33] not necessarily to

farming). These individuals form a cline from WHG to EHG that is correlated with geography

(Figure 1B), although it is neither geographically nor temporally uniform (Figure 2, Extended

Data Figure 3), and there is also substructure in phenotypically important variants

(Supplementary Information Note 2).

From present-day Ukraine, our study reports new genome-wide data from five Mesolithic

individuals from ~9500-6000 BCE, and 31 Neolithic individuals from ~6000-3500 BCE. On the

cline from WHG- to EHG-related ancestry, the Mesolithic individuals fall towards the East,

intermediate between EHG and Mesolithic hunter-gatherers from Sweden (Figure 1B).[7] The

Neolithic population has a significant difference in ancestry compared to the Mesolithic

(Figures 1B, Figure 2), with a shift towards WHG shown by the statistic D(Mbuti, WHG,

Ukraine_Mesolithic, Ukraine_Neolithic); Z=8.9 (Supplementary Information Table 2).

Unexpectedly, one Neolithic individual from Dereivka (I3719), which we directly date to

4949-4799 BCE, has entirely NW Anatolian Neolithic-related ancestry.

The pastoralist Bronze Age Yamnaya complex originated on the Eurasian steppe and is a

plausible source for the dispersal of steppe-related ancestry into central and western Europe

around 2500 BCE.[13] All previously reported Yamnaya individuals were from Samara[7] and

Kalmykia[15] in southwest Russia, and had entirely steppe-related ancestry. Here, we report

three Yamnaya individuals from further West – from Ukraine and Bulgaria – and show that

while they all have high levels of steppe-related ancestry, one from Ozera in Ukraine and one

from Bulgaria (I1917 and Bul4, both dated to ~3000 BCE) have NW Anatolian Neolithic-

related admixture, the first evidence of such ancestry in Yamnaya –associated individuals

(Figure 1B,D, Supplementary Data Table 2). Two Copper Age individuals (I4110 and I6561,

Ukraine_Eneolithic) from Dereivka and Alexandria dated to ~3600-3400 BCE (and thus

preceding the Yamnaya complex) also have mixtures of steppe- and NW Anatolian Neolithic-

related ancestry (Figure 1D, Supplementary Data Table 2).

At Zvejnieki in Latvia (17 newly reported individuals, and additional data for 5 first reported

in Ref. 34) we observe a transition in hunter-gatherer-related ancestry that is the opposite of

that seen in Ukraine. We find (Supplementary Data Table 3) that Mesolithic and Early

Neolithic individuals (Latvia_HG) associated with the Kunda and Narva cultures have

ancestry intermediate between WHG (~70%) and EHG (~30%), consistent with previous

reports.[34-36] We also detect a shift in ancestry between the Early Neolithic and individuals

associated with the Middle Neolithic Comb Ware Complex (Latvia_MN), who have more

EHG-related ancestry (we estimate 65% EHG, but two of four individuals appear almost

100% EHG in PCA). The most recent individual, associated with the Final Neolithic Corded

Ware Complex (I4629, Latvia_LN), attests to another ancestry shift, clustering closely with

Yamnaya from Samara,[7] Kalmykia[15] and Ukraine (Figure 2).

We report new Upper Palaeolithic and Mesolithic data from southern and western Europe.[17]

Sicilian (I2158) and Croatian (I1875) individuals dating to ~12,000 and 6100 BCE cluster with

previously reported western hunter-gatherers (Figure 1B&D), including individuals from

Loschbour[23] (Luxembourg, 6100 BCE), Bichon[19] (Switzerland, 11,700 BCE), and Villabruna[17]

(Italy 12,000 BCE). These results demonstrate that WHG populations[23] were widely

distributed from the Atlantic seaboard of Europe in the West, to Sicily in the South, to the

Balkan Peninsula in the Southeast, for at least six thousand years.

A particularly important hunter-gatherer population that we report is from the Iron Gates

region that straddles the border of present-day Romania and Serbia. This population

(Iron_Gates_HG) is represented in our study by 40 individuals from five sites. Modeling Iron

Gates hunter-gatherers as a mixture of WHG and EHG (Supplementary Table 3) shows that

they are intermediate between WHG (~85%) and EHG (~15%). However, this *qpAdm* model

does not fit well (p=0.0003, Supplementary table 3) and the Iron Gates hunter-gatherers carry

mitochondrial haplogroup K1 (7/40) as well as other subclades of haplogroups U (32/40) and

H (1/40). This contrasts with WHG, EHG and Scandinavian hunter-gatherers who almost all

carry haplogroups U5 or U2. One interpretation is that the Iron Gates hunter-gatherers have

ancestry that is not present in either WHG or EHG. Possible scenarios include genetic contact

between the ancestors of the Iron Gates population and Anatolia, or that the Iron Gates

population is related to the source population from which the WHG split during a re-

expansion into Europe from the Southeast after the Last Glacial Maximum.[17,37]

A notable finding from the Iron Gates concerns the four individuals from the site of Lepenski

Vir, two of whom (I4665 & I5405, 6200-5600 BCE), have entirely NW Anatolian Neolithic-

related ancestry. Strontium and Nitrogen isotope data[38] indicate that both these individuals

were migrants from outside the Iron Gates, and ate a primarily terrestrial diet (Supplementary

Information section 1). A third individual (I4666, 6070 BCE) has a mixture of NW Anatolian

Neolithic-related and hunter-gatherer-related ancestry and ate a primarily aquatic diet, while a

fourth, probably earlier, individual (I5407) had entirely hunter-gatherer-related ancestry

(Figure 1D, Supplementary Information section 1). We also identify one individual from

Padina (I5232), dated to 5950 BCE that had a mixture of NW Anatolian Neolithic-related and

hunter-gatherer-related ancestry. These results demonstrate that the Iron Gates was a region of

interaction between groups distinct in both ancestry and subsistence strategy.

Population transformations in the first farmers

Neolithic populations from present-day Bulgaria, Croatia, Macedonia, Serbia and Romania

cluster closely with the NW Anatolian Neolithic farmers (Figure 1), consistent with

archaeological evidence.[39] Modeling Balkan Neolithic populations as a mixture of NW

Anatolian Neolithic and WHG, we estimate that 98% (95% confidence interval [CI]; 97-

100%) of their ancestry is NW Anatolian Neolithic-related. A striking exception is evident in

8 out of 9 individuals from Malak Preslavets in present-day Bulgaria.[40] These individuals

lived in the mid-6[th] millennium BCE and have significantly more hunter-gatherer-related

ancestry than other Balkan Neolithic populations (Figure 1B,D, Extended Data Figures 1-3,

Supplementary Tables 2-4); a model of 82% (CI: 77-86%) NW Anatolian Neolithic-related,

15% (CI: 12-17%) WHG-related, and 4% (CI: 0-9%) EHG-related ancestry is a fit to the data.

This hunter-gatherer-related ancestry with a ~4:1 WHG:EHG ratio plausibly represents a

contribution from local Balkan hunter-gatherers genetically similar to those of the Iron Gates.

Late Mesolithic hunter-gatherers in the Balkans were likely concentrated along the coast and

major rivers such as the Danube,[41] which directly connects the Iron Gates with Malak

Preslavets. Thus, early farmer groups with the most hunter-gatherer-related ancestry may

have been those that lived close to the highest densities of hunter-gatherers.

In the Balkans, Copper Age populations (Balkans_Chalcolithic) harbor significantly more

hunter-gatherer-related ancestry than Neolithic populations as shown, for example, by the

statistic D(Mbuti, WHG, Balkans_Neolithic, Balkans_Chalcolithic); Z=4.3 (Supplementary

Data Table 2). This is roughly contemporary with the "resurgence" of hunter-gatherer

ancestry previously reported in central Europe and Iberia[7,10,42] and is consistent with changes

in funeral rites, specifically the reappearance around 4500 BCE of the Mesolithic tradition of

extended supine burial – in contrast to the Early Neolithic tradition of flexed burial.[43] Four

individuals associated with the Copper Age Trypillian population have ~80% NW Anatolian-

related ancestry (Supplementary Table 3), confirming that the ancestry of the first farmers of

present-day Ukraine was largely derived from the same source as the farmers of Anatolia and

western Europe. Their ~20% hunter-gatherer ancestry is intermediate between WHG and

EHG, consistent with deriving from the Neolithic hunter-gatherers of the region.

We also report the first genetic data associated with the Late Neolithic Globular Amphora

Complex. Individuals from two Globular Amphora sites in Poland and Ukraine form a tight

cluster, showing high similarity over a large distance (Figure 1B,D). Both Globular Amphora

Complex groups of samples had more hunter-gatherer-related ancestry than Middle Neolithic

groups from Central Europe[7] (we estimate 25% [CI: 22-27%] WHG ancestry, similar to

Chalcolithic Iberia, Supplementary Data Table 3). In east-central Europe, the Globular

Amphora Complex preceded or abutted the Corded Ware Complex that marks the appearance

of steppe-related ancestry,[7,15] while in southeastern Europe, the Globular Amphora Complex

bordered populations with steppe-influenced material cultures for hundreds of years[44] and yet

the individuals in our study have no evidence of steppe-related ancestry, providing support for

the hypothesis that this material cultural frontier was also a barrier to gene flow.

The movements from the Pontic-Caspian steppe of individuals similar to those associated

with the Yamnaya Cultural Complex in the 3rd millennium BCE contributed about 75% of the

ancestry of individuals associated with the Corded Ware Complex and about 50% of the

ancestry of succeeding material cultures such as the Bell Beaker Complex in central

Europe.[7,15] In two directly dated individuals from southeastern Europe, one (ANI163) from

the Varna I cemetery dated to 4711-4550 BCE and one (I2181) from nearby Smyadovo dated

to 4550-4450 BCE, we find far earlier evidence of steppe-related ancestry (Figure 1B,D).

These findings push back the first evidence of steppe-related ancestry this far West in Europe

by almost 2,000 years, but it was sporadic as other Copper Age (~5000-4000 BCE) individuals

from the Balkans have no evidence of it. Bronze Age (~3400-1100 BCE) individuals do have

steppe-related ancestry (we estimate 30%; CI: 26-35%), with the highest proportions in the

four latest Balkan Bronze Age individuals in our data (later than ~1700 BCE) and the least in

earlier Bronze Age individuals (3400-2500 BCE; Figure 1D).

A novel source of ancestry in Neolithic Europe

An important question about the initial spread of farming into Europe is whether the first

farmers that brought agriculture to northern Europe and to southern Europe were derived from

a single population or instead represent distinct migrations. We confirm that Mediterranean

populations, represented in our study by individuals associated with the Epicardial Early

Neolithic from Iberia[7], are closely related to Danubian populations represented by the

Linearbandkeramik (LBK) from central Europe[7,45] and that both are closely related to the

Balkan Neolithic population. These three populations form a clade with the NW Anatolian

Neolithic individuals as an outgroup, consistent with a single migration into the Balkan

peninsula, which then split into two (Supplementary Information Note 3).

In contrast, five southern Greek Neolithic individuals (Peloponnese_Neolithic) – three (plus

one previously published[26]) from Diros Cave and one from Franchthi Cave – are not

consistent with descending from the same source population as other European farmers. *D*-

statistics (Supplementary Information Table 2) show that in fact, these "Peloponnese

Neolithic" individuals dated to ~4000 BCE are shifted away from WHG and towards CHG,

relative to Anatolian and Balkan Neolithic individuals. We see the same pattern in a single

Neolithic individual from Krepost in present-day Bulgaria (I0679_d, 5718-5626 BCE). An

even more dramatic shift towards CHG has been observed in individuals associated with the

Bronze Age Minoan and Mycenaean cultures,[26] and thus there was gene flow into the region

from populations with CHG-rich ancestry throughout the Neolithic, Chalcolithic and Bronze

Age. Possible sources are related to the Neolithic population from the central Anatolian site of

Tepecik Ciftlik,[21] or the Aegean site of Kumtepe,[11] who are also shifted towards CHG relative

to NW Anatolian Neolithic samples, as are later Copper and Bronze Age Anatolians.[10,26]

Sex-biased admixture between hunter-gatherers and farmers

We provide the first evidence for sex-biased admixture between hunter-gatherers and farmers

in Europe, showing that the Middle Neolithic "resurgence" of hunter-gatherer-related

ancestry[7,42] in central Europe and Iberia was driven more by males than by females (Figure

3B&C, Supplementary Data Table 5, Extended Data Figure 4). To document this we used

qpAdm to compute ancestry proportions on the autosomes and the X chromosome; since

males always inherit their X chromosome from their mothers, differences imply sex-biased

mixture. In the Balkan Neolithic there is no evidence of sex bias (Z=0.27 where a positive Z-

score implies male hunter-gatherer bias), nor in the LBK and Iberian_Early Neolithic (Z=-

0.22 and 0.74). In the Copper Age there is clear bias: weak in the Balkans (Z=1.66), but

stronger in Iberia (Z=3.08) and Central Europe (Z=2.74). Consistent with this, hunter-gatherer

mitochondrial haplogroups (haplogroup U)[46] are rare and within the intervals of genome-wide

ancestry proportions, but hunter-gatherer-associated Y chromosomes (haplogroups I, R1 and

C1)[17] are more common: 7/9 in the Iberian Neolithic/Copper Age and 9/10 in Middle-Late

Neolithic Central Europe (Central_MN and Globular_Amphora) (Figure 3C).

No evidence that steppe-related ancestry moved through southeast Europe into Anatolia

One version of the Steppe Hypothesis of Indo-European language origins suggests that Proto-

Indo-European languages developed north of the Black and Caspian seas, and that the earliest

known diverging branch – Anatolian – was spread into Asia Minor by movements of steppe

peoples through the Balkan peninsula during the Copper Age around 4000 BCE.[47] If this were

correct, then one way to detect evidence of it would be the appearance of large amounts of

steppe-related ancestry first in the Balkan Peninsula, and then in Anatolia. However, our data

show no evidence for this scenario. While we find sporadic examples of steppe-related

ancestry in Balkan Copper and Bronze Age individuals, this ancestry is rare until the late

Bronze Age. Moreover, while Bronze Age Anatolian individuals have CHG-related

ancestry,[26] they have neither the EHG-related ancestry characteristic of all steppe populations

sampled to date,[19] nor the WHG-related ancestry that is ubiquitous in Neolithic southeastern

Europe (Extended Data Figure 2, Supplementary Data Table 2). An alternative hypothesis is

that the ultimate homeland of Proto-Indo-European languages was in the Caucasus or in Iran.

In this scenario, westward movement contributed to the dispersal of Anatolian languages, and

northward movement and mixture with EHG was responsible for the formation of a "Late

Proto-Indo European"-speaking population associated with the Yamnaya Complex.[13] While

this scenario gains plausibility from our results, it remains possible that Indo-European

languages were spread through southeastern Europe into Anatolia without large-scale

population movement or admixture.

Discussion

Our study shows that southeastern Europe consistently served as a genetic contact zone.

Before the arrival of farming, the region saw interaction between diverged groups of hunter-

gatherers, and this interaction continued after farming arrived. While this study has clarified

the genomic history of southeastern Europe from the Mesolithic to the Bronze Age, the

processes that connected these populations to the ones living today remain largely unknown.

An important direction for future research will be to sample populations from the Bronze

Age, Iron Age, Roman, and Medieval periods and to compare them to present-day

populations to understand how these transitions occurred.

Methods

Ancient DNA Analysis

We extracted DNA and prepared next-generation sequencing libraries in four different dedicated ancient DNA laboratories (Adelaide, Boston, Budapest, and Tuebingen). We also prepared samples for extraction in a fifth laboratory (Dublin), from whence it was sent to Boston for DNA extraction and library preparation (Supplementary Table 1).

Two samples were processed at the Australian Centre for Ancient DNA, Adelaide, Australia, according to previously published methods[7] and sent to Boston for subsequent screening, 1240k capture and sequencing.

Seven samples were processed[27] at the Institute of Archaeology RCH HAS, Budapest, Hungary, and amplified libraries were sent to Boston for screening, 1240k capture and sequencing.

Seventeen samples were processed at the Institute for Archaeological Sciences of the University of Tuebingen and at the Max Planck Institute for the Science of Human History in Jena, Germany. Extraction[48] and library preparation[49,50] followed established protocols. We performed in-solution capture as described below ("1240k capture") and sequenced on an Illumina HiSeq 4000 or NextSeq 500 for 76bp using either single- or paired-end sequencing.

The remaining 197 samples were processed at Harvard Medical School, Boston, USA. From about 75mg of sample powder from each sample (extracted in Boston or University College Dublin, Dublin, Ireland), we extracted DNA following established methods[48] replacing the column assembly with the column extenders from a Roche kit.[51] We prepared double barcoded libraries with truncated adapters from between one ninth and one third of the DNA extract. Most libraries included in the nuclear genome analysis (90%) were subjected to partial ("half") Uracil-DNA-glycosylase (UDG) treatment before blunt end repair. This treatment reduces by an order of magnitude the characteristic cytosine-to-thymine errors of ancient DNA data[52], but works inefficiently at the 5' ends,[50] thereby leaving a signal of characteristic damage at the terminal ends of ancient sequences. Some libraries were not UDG-treated ("minus"). For some samples we increased coverage by preparing additional libraries from the existing DNA extract using the partial UDG library preparation, but replacing the MinElute column cleanups in between enzymatic reactions with magnetic bead cleanups, and the final PCR cleanup with SPRI bead cleanup.[53,54]

We screened all libraries from Adelaide, Boston and Budapest by enriching for the

mitochondrial genome plus about 3,000 (50 in an earlier, unpublished, version) nuclear SNPs

using a bead-capture[55] but with the probes replaced by amplified oligonucleotides synthesized

by CustomArray Inc. After the capture, we completed the adapter sites using PCR, attaching

dual index combinations[56] to each enriched library. We sequenced the products of between

100 and 200 libraries together with the non-enriched libraries (shotgun) on an Illumina

NextSeq500 using v2 150 cycle kits for 2x76 cycles and 2x7 cycles.

In Boston, we performed two rounds of in-solution enrichment ("1240k capture") for a

targeted set of 1,237,207 SNPs using previously reported protocols.[7,14,23] For a total of 34

individuals, we increased coverage by building one to eight additional libraries for the same

sample. When we built multiple libraries from the same extract, we often pooled them in

equimolar ratios before the capture. We performed all sequencing on an Illumina NextSeq500

using v2 150 cycle kits for 2x76 cycles and 2x7 cycles. We attempted to sequence each

enriched library up to the point where we estimated that it was economically inefficient to

sequence further. Specifically, we iteratively sequenced more and more from each individual

and only stopped when we estimated that the expected increase in the number of targeted

SNPs hit at least once would be less than about one for every 100 new read pairs generated.

After sequencing, we trimmed two bases from the end of each read and aligned to the human

genome (b37/hg19) using *bwa*.[57] We then removed individuals with evidence of

contamination based on mitochondrial DNA polymorphism[58] or difference in PCA space

between damaged and undamaged reads[59], a high rate of heterozygosity on chromosome X

despite being male[59,60], or an atypical ratio of X-to-Y sequences. We also removed individuals

that had low coverage (fewer than 15,000 SNPs hit on the autosomes). We report, but do not

analyze, data from nine individuals that were first-degree relatives of others in the dataset

(determined by comparing rates of allele sharing between pairs of individuals).

After removing a small number of sites that failed to capture, we were left with a total of

1,233,013 sites of which 32,670 were on chromosome X and 49,704 were on chromosome Y,

with a median coverage at targeted SNPs on the 214 newly reported individuals of 0.90

(range 0.007-9.2; Supplementary Table 1). We generated "pseudo-haploid" calls by selecting

a single read randomly for each individual at each SNP. Thus, there is only a single allele

from each individual at each site, but adjacent alleles might come from either of the two

haplotypes of the individual. We merged the newly reported data with previously reported

data from 274 other ancient individuals[9-11,15-27], making pseudo-haploid calls in the same way

at the 1240k sites for individuals that were shotgun sequenced rather than captured.

Using the captured mitochondrial sequence from the screening process, we called

mitochondrial haplotypes. Using the captured SNPs on the Y chromosome, we called Y

chromosome haplogroups for males by restricting to sequences with mapping quality ≥30 and

bases with base quality ≥30. We determined the most derived mutation for each individual,

using the nomenclature of the International Society of Genetic Genealogy

(http://www.isogg.org) version 11.110 (21 April 2016).

Population genetic analysis

To analyze these ancient individuals in the context of present day genetic diversity, we

merged them with the following two datasets:

1. 300 high coverage genomes from a diverse worldwide set of 142 populations

sequenced as part of the Simons Genome Diversity Project[28] (SGDP merge).

2. 799 West Eurasian individuals genotyped on the Human Origins array[23], with

597,573 sites in the merged dataset (HO merge).

We computed principal components of the present-day individuals in the HO merge and

projected the ancient individuals onto the first two components using the "*lsqproject: YES*"

option in *smartpca (v15100)*[61] (https://www.hsph.harvard.edu/alkes-price/software/).

We ran *ADMIXTURE (v1.3.0)* in both supervised and unsupervised mode. In supervised mode

we used only the ancient individuals, on the full set of SNPs, and the following population

labels fixed:

- *Anatolia_Neolithic*
- *WHG*
- *EHG*
- *Yamnaya*
493

For unsupervised mode we used the HO merge, including 799 present-day individuals. We

flagged individuals that were genetic outliers based on PCA and ADMIXTURE, relative to

other individuals from the same time period and archaeological culture.

We computed *D*-statistics using *qpDstat (v710)*. *D*-statistics of the form D(A,B,X,Y) test the

null hypothesis of the unrooted tree topology ((A,B),(X,Y)). A positive value indicates that

either A and X, or B and Y, share more drift than expected under the null hypothesis. We

quote *D*-statistics as the *Z*-score computed using default block jackknife parameters.

We fitted admixture proportions with *qpAdm (v610)* using the SGDP merge. Given a set of

outgroup ("right") populations, *qpAdm* models one of a set of source ("left") populations (the

"test" population) as a mixture of the other sources by fitting admixture proportions to match

the observed matrix of f_4-statistics as closely as possible. We report a p-value for the null

hypothesis that the test population does not have ancestry from another source that is

differentially related to the right populations. We computed standard errors for the mixture

proportions using a block jackknife. Importantly, *qpAdm* does not require that the source

populations are actually the admixing populations, only that they are a clade with the correct

admixing populations, relative to the other sources. Infeasible coefficient estimates (i.e.

outside [0,1]) are usually a sign of poor model fit, but in the case where the source with a

negative coefficient is itself admixed, could be interpreted as implying that the true source is a

population with different admixture proportions. We used the following set of seven

populations as outgroups or "right populations":

• *Mbuti.DG*
• *Ust_Ishim_HG_published.DG*
• *Mota.SG*
• *MA1_HG.SG*
• *Villabruna*
• *Papuan.DG*
• *Onge.DG*
• *Han.DG*

For some analyses where we required extra resolution (Extended Data Table 4) we used an

extended set of 14 right (outgroup) populations, including additional Upper Paleolithic

European individuals[17]:

• *ElMiron*
• *Mota.SG*
• *Mbuti.DG*
• *Ust_Ishim_HG_published.DG*
• *MA1_HG.SG*
• *AfontovaGora3*
• *GoyetQ116-1_published*
• *Villabruna*
• *Kostenki14*
• *Vestonice16*
• *Karitiana.DG*
• *Papuan.DG*
• *Onge.DG*
• *Han.DG*
542

We also fitted admixture graphs with *qpGraph (v6021)*[30] (https://github.com/DReichLab/

AdmixTools, Supplementary Information, section 3). Like *qpAdm*, *qpGraph* also tries to

match a matrix of *f*-statistics, but rather than fitting one population as a mixture of other,

specified, populations, it fits the relationship between all tested populations simultaneously,

potentially incorporating multiple admixture events. However, *qpGraph* requires the graph

relating populations to be specified in advance. We tested goodness-of-fit by computing the

expected *D*-statistics under the fitted model, finding the largest *D*-statistic outlier between the

fitted and observed model, and computing a *Z*-score using a block jackknife.

For 116 individuals with hunter-gatherer-related ancestry we estimated an effective migration

surface using the software *EEMS* (https://github.com/dipetkov/eems)[62]. We computed

pairwise differences between individuals using the *bed2diffs2* program provided with *EEMS*.

We set the number of demes to 400 and defined the outer boundary of the region by the

polygon (in latitude-longitude co-ordinates) [(66,60), (60,10), (45,-15), (35,-10), (35,60)]. We

ran the MCMC ten times with different random seeds, each time with one million burn-in and

four million regular iterations, thinned to one in ten thousand.

To analyze potential sex bias in admixture, we used *qpAdm* to estimate admixture proportions

on the autosomes (default option) and on the X chromosome (option "*chrom: 23*"). We

computed Z-scores for the difference between the autosomes and the X chromosome as $Z =$

$\frac{p_A - p_X}{\sqrt{\sigma_A^2 + \sigma_X^2}}$ where p_A and p_X are the hunter-gatherer admixture proportions on the autosomes and

the X chromosome, and σ_A and σ_X are the corresponding jackknife standard deviations. Thus,

a positive Z-score means that there is more hunter-gatherer admixture on the autosomes than

on the X chromosome, indicating that the hunter-gatherer admixture was male-biased.

Because X chromosome standard errors are high and *qpAdm* results can be sensitive to which

population is first in the list of outgroup populations, we checked that the patterns we observe

were robust to cyclic permutation of the outgroups. To compare frequencies of hunter-

gatherer uniparental markers, we counted the individuals with mitochondrial haplogroup U

and Y chromosome haplogroups C2, I2 and R1, which are all common in Mesolithic hunter-

gatherers but rare or absent in Anatolian Neolithic individuals. The Iron Gates hunter-

gatherers also carry H and K1 mitochondrial haplogroups so the proportion of haplogroup U

represents the minimum maternal hunter-gatherer contribution. We computed binomial

confidence intervals for the proportion of haplogroups associated with each ancestry type

using the Agresti-Coull method[63,64] implemented in the *binom* package in *R*.

Given autosomal and X chromosome admixture proportions, we estimated the proportion of

male and female hunter-gatherer ancestors by assuming a single-pulse model of admixture. If

the proportions of male and female ancestors that are hunter-gatherer-related are given by *m*

and *f,* respectively, then the proportions of hunter-gatherer-related ancestry on the autosomes

and the X chromosome are given by $\frac{m+f}{2}$ and $\frac{m+2f}{3}$. We approximated the sampling error in

the observed admixture proportions by the estimated jackknife error and computed the

likelihood surface for *(m,f)* over a grid ranging from (0,0) to (1,1).

Direct AMS ^{14}C Bone Dates

We report 113 new direct AMS ^{14}C bone dates for 112 individuals from multiple AMS

radiocarbon laboratories. In general, bone samples were manually cleaned and demineralized

in weak HCl and, in most cases (PSU, UCIAMS, OxA), soaked in an alkali bath (NaOH) at

room temperature to remove contaminating soil humates. Samples were then rinsed to

neutrality in Nanopure H_2O and gelatinized in HCL.[65] The resulting gelatin was lyophilized

and weighed to determine percent yield as a measure of collagen preservation (% crude

gelatin yield). Collagen was then directly AMS ^{14}C dated (Beta, AA) or further purified using

ultrafiltration (PSU, UCIAMS, OxA, Poz, MAMS).[66] It is standard in some laboratories

(PSU/UCIAMS, OxA) to use stable carbon and nitrogen isotopes as an additional quality

control measure. For these samples, the %C, %N and C:N ratios were evaluated before AMS

^{14}C dating.[67] C:N ratios for well-preserved samples fall between 2.9 and 3.6, indicating good

collagen preservation.[68] For 94 new samples, we also report δ^{13}C and δ^{15}N values

(Supplementary Table 6).

All ^{14}C ages were δ^{13}C-corrected for mass dependent fractionation with measured ^{13}C/^{12}C

values[69] and calibrated with OxCal version 4.2.3[70] using the IntCal13 northern hemisphere

calibration curve.[70] For hunter-gatherers from the Iron Gates, the direct ^{14}C dates tend to be

overestimates because of the freshwater reservoir effect (FRE), which arises because of a diet

including fish that consumed ancient carbon, and for these individuals we performed a

correction (Supplementary Information Note 1),[71] assuming that 100% FRE = 545±70 yr, and

δ^{15}N values of 8.3% and 17.0% for 100% terrestrial and aquatic diets, respectively.

Acknowledgments

We thank David Anthony, Iosif Lazaridis, and Mark Lipson for comments on the manuscript,

Bastien Llamas and Alan Cooper for contributions to laboratory work, Richard Evershed for

contributing ^{14}C dates and Friederike Novotny for assistance with samples. Support for this

project was provided by the Human Frontier Science Program fellowship LT001095/2014-L

to I.M.; by DFG grant AL 287 / 14-1 to K.W.A.; by Irish Research Council grant

GOIPG/2013/36 to D.F.; by the NSF Archaeometry program BCS-1460369 to DJK (for AMS

^{14}C work at Penn State); by MEN-UEFISCDI grant, Partnerships in Priority Areas Program –

PN II (PN-II-PT-PCCA-2013-4-2302) to C.L.; by Croatian Science Foundation grant IP-

2016-06-1450 to M.N.; by European Research Council grant ERC StG 283503 and Deutsche

Forschungsgemeinschaft DFG FOR2237 to K.H.; by ERC starting grant ADNABIOARC

(263441) to R.P.; and by US National Science Foundation HOMINID grant BCS-1032255,

US National Institutes of Health grant GM100233, and the Howard Hughes Medical Institute

to D.R.

Author Contributions

SAR, AS-N, SVai, SA, KWA, RA, DA, AA, NA, KB, MBG, HB, MB, ABo, YB, ABu, JB,

SC, NC, RC, MC, CC, DD, NE, MFr, BGal, GG, BGe, THa, VH, KH, THi, SI, IJ, IKa, DKa,

AK, DLa, MLa, CL, MLe, KL, DLV, DLo, IL, MMa, FM, KM, HM, MMe, PM, VM, VP,

TDP, ASi, LS, MŠ, VS, PS, ASt, TS, MT-N, CT, IV, FVa, SVas, FVe, SV, EV, BV, CV, JZ,

SZ, PWS, GC, RK, DC, GZ, BGay, MLi, AGN, IP, AP, DB, CB, JK, RP & DR assembled

and interpreted archaeological material. CP, AS-N, NR, NB, FC, OC, DF, MFe, BGam, GGF,

WH, EH, EJ, DKe, BK-K, IKu, MMi, AM, KN, MN, JO, SP, KSi, KSt & SVai performed

laboratory work. IM, CP, AS-N, SM, IO, NP & DR analyzed data. DJK, ST, DB, CB

interpreted [14]C dates. JK, RP & DR supervised analysis or laboratory work. IM & DR wrote

the paper, with input from all co-authors.

Figure 1: Geographic locations and genetic structure of newly reported individuals. **A**: Location and groupings of newly reported individuals. **B**: Individuals projected onto axes defined by the principal components of 799 present-day West Eurasians (not shown in this plot for clarity, but shown in Extended Data Figure 1). Projected points include selected published individuals (faded colored circles, labeled) and newly reported individuals (other symbols; outliers shown by additional black circles). Colored polygons indicate the individuals that had cluster memberships fixed at 100% for the supervised admixture analysis in **D**. **C**: Estimated age (direct or contextual) for each sample. Approximate chronology used in southeastern Europe shown to the right **D**: Supervised ADMIXTURE plot, modeling each ancient individual (one per row), as a mixture of populations represented by clusters containing Anatolian Neolithic (grey), Yamnaya from Samara (yellow), EHG (pink) and WHG (green). Dates indicate approximate range of individuals in each population. Map data in **A** from the *R* package *mapdata*.

651 **Figure 2:** Structure and population change in European populations with hunter-gatherer-

652 related ancestry. This figure shows inferred ancestry proportions for populations modeled as a

653 mixture of WHG, EHG and CHG (Supplementary Table S3.1.3). Dashed lines show

654 populations from the same geographic region. Standard errors range from 1.5-8.3%

655 (Supplementary Table S3.1.3).

Figure 3: Structure and population change in European populations with NW Anatolian
Neolithic-related ancestry. **A**: each population is modeled as a mixture of NW Anatolia
Neolithic, WHG, and EHG. Dashed lines show temporal relationships between populations
from the same geographic areas with similar ancestries. Standard errors range from 0.7-6.0%
(Supplementary Table S3.2.2). **B**: Z-scores for the difference in hunter-gatherer-related
ancestry on the autosomes compared to the X chromosome when populations are modeled as
a mixture of NW Anatolia Neolithic and WHG. A positive score indicates that there is more
hunter-gatherer-related ancestry on the autosomes and therefore the hunter-gatherer-related
ancestry is male-biased. **C**: Hunter-gatherer-related ancestry proportions on the autosomes, X
chromosome, mitochondrial DNA (i.e. mt haplogroup U), and the Y chromosome (i.e. Y
chromosome haplogroups I2, R1 and C2). Bars show approximate 95% confidence intervals.
"Combined" populations merge all individuals from different times from a geographic area.

671 **Extended Data Figures**

674 **Extended Data Figure 1:** PCA of all ancient individuals, projected onto principal

675 components defined by 799 present-day West Eurasian individuals. (This differs from Figure

676 1B in that the plot is not cropped and the present-day individuals are shown.)

Extended Data Figure 2: Unsupervised ADMIXTURE plot from k=4 to 12, on a dataset
consisting of 1099 present-day individuals and 476 ancient individuals. We show newly
reported ancient individuals and some previously published individuals for comparison.

Extended Data Figure 3: Spatial structure in hunter-gatherers. Estimated effective migration

surface (EEMS).[62] This fits a model of genetic relatedness where individuals move (in a

random direction) from generation to generation on an underlying grid so that genetic

relatedness is determined by distance. The migration parameter m defines the local rate of

migration, varies on the grid and is inferred. This plot shows log_{10} m, scaled relative to the

average migration rate (which is arbitrary). Thus $log10(m)=2$, for example, implies that the

rate of migration at this point on the grid is 100 times higher than average. To restrict as much

as possible to hunter-gatherer structure, the migration surface is inferred using data from 116

individuals from populations that date earlier than ~5000 BCE and have no NW Anatolian-

related ancestry. Though the migration surface is sensitive to sampling, and fine-scale

features may not be interpretable, the migration "barrier" (region of low migration) running

north-south and separating populations with primarily WHG from primarily EHG ancestry

seems to be robust, and consistent with inferred admixture proportions. This analysis suggests

that Mesolithic hunter-gatherer population structure was clustered and not smoothly clinal, in

the sense that genetic differentiation did not vary consistently with distance. Superimposed

on this background, pies show the WHG, EHG and CHG ancestry proportions inferred for

populations used to construct the migration surface (another way of visualizing the data in

show in Figure 2, Supplementary Table 3.1.3 – we use two population models if they fit with

p>0.1, and three population models otherwise). Pies with only a single color have been fixed

to be the source populations.

703
704 **Extended Data Figure 4**: log-likelihood surfaces for the proportion of female (x-axis) and
705 male (y-axis) ancestors that are hunter-gatherer-related for the combined populations
706 analyzed in Figure 3C, and the two populations with the strongest evidence for sex-bias. Log-
707 likelihood scale ranges from 0 to -10, where 0 is the feasible point with the highest likelihood.
708
709

Supplementary Tables

References

Tringham, R. E. in *The Transition to Agriculture in Prehistoric Europe* (ed D. Price) 19-56 (Cambridge University Press, 2000).

Bellwood, P. *First Farmers: The Origins of Agricultural Societies*. 2nd edn, (Wiley-Blackwell, 2004).

Golitko, M. in *Ancient Europe, 8000 B.C. to A.D. 1000: An Encyclopedia of the Barbarian World* (eds P. Bogucki & P.J. Crabtree) 259-266 (Charles Scribners & Sons, 2003).

Vander Linden, M. in *Investigating Archaeological Cultures: Material Culture, Variability, and Transmission* (eds B.W. Roberts & M. Vander Linden) 289-319 (Springer, 2012).

Bramanti, B. *et al.* Genetic discontinuity between local hunter-gatherers and central Europe's first farmers. *Science* **326**, 137-140 (2009).

Skoglund, P. *et al.* Origins and genetic legacy of Neolithic farmers and hunter-gatherers in Europe. *Science* **336**, 466-469 (2012).

Haak, W. *et al.* Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* **522**, 207-211 (2015).

Cassidy, L. M. *et al.* Neolithic and Bronze Age migration to Ireland and establishment of the insular Atlantic genome. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 368-373 (2016).

Hofmanova, Z. *et al.* Early farmers from across Europe directly descended from Neolithic Aegeans. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 6886-6891 (2016).

Mathieson, I. *et al.* Genome-wide patterns of selection in 230 ancient Eurasians. *Nature* **528**, 499-503 (2015).

Omrak, A. *et al.* Genomic Evidence Establishes Anatolia as the Source of the European Neolithic Gene Pool. *Curr. Biol.* **26**, 270-275 (2016).

Müller, J., Rassmann, K. & Videiko, M. *Trypillia Mega-Sites and European Prehistory: 4100–3400 BCE.*, (Routledge, 2016).

Anthony, D. W. *The horse the wheel and language*. (Princeton University Press, 2007).

Fu, Q. *et al.* An early modern human from Romania with a recent Neanderthal ancestor. *Nature* (2015).

Allentoft, M. E. *et al.* Population genomics of Bronze Age Eurasia. *Nature* **522**, 167-172 (2015).

Fu, Q. *et al.* Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* **514**, 445-449 (2014).

Fu, Q. *et al.* The genetic history of Ice Age Europe. *Nature* **534**, 200-205 (2016).

Gallego Llorente, M. *et al.* Ancient Ethiopian genome reveals extensive Eurasian admixture in Eastern Africa. *Science* **350**, 820-822 (2015).

Jones, E. R. *et al.* Upper Palaeolithic genomes reveal deep roots of modern Eurasians. *Nature communications* **6**, 8912 (2015).

758	20	Keller, A. *et al.* New insights into the Tyrolean Iceman's origin and phenotype as
759		inferred by whole-genome sequencing. *Nature communications* **3**, 698 (2012).
760	21	Kilinc, G. M. *et al.* The Demographic Development of the First Farmers in Anatolia.
761		*Curr. Biol.* (2016).
762	22	Lazaridis, I. *et al.* Genomic insights into the origin of farming in the ancient Near
763		East. *Nature* **536**, 419-424 (2016).
764	23	Lazaridis, I. *et al.* Ancient human genomes suggest three ancestral populations for
765		present-day Europeans. *Nature* **513**, 409-413 (2014).
766	24	Olalde, I. *et al.* Derived immune and ancestral pigmentation alleles in a 7,000-year-
767		old Mesolithic European. *Nature* **507**, 225-228 (2014).
768	25	Raghavan, M. *et al.* Upper Palaeolithic Siberian genome reveals dual ancestry of
769		Native Americans. *Nature* **505**, 87-91 (2014).
770	26	Lazaridis, I. *et al.* Genetic origins of the Minoans and Mycenaeans. *Nature* **548**, 214-
771		218 (2017).
772	27	Lipson, M. *et al.* Parallel ancient genomic transects reveal complex population
773		history of early European farmers. *bioRxiv*, https://doi.org/10.1101/114488 (2017).
774	28	Mallick, S. *et al.* The Simons Genome Diversity Project: 300 genomes from 142
775		diverse populations. *Nature* **538**, 201-206 (2016).
776	29	Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry
777		in unrelated individuals. *Genome Res.* **19**, 1655-1664 (2009).
778	30	Patterson, N. *et al.* Ancient admixture in human history. *Genetics* **192**, 1065-1093
779		(2012).
780	31	Gronenborn, D. & Dolukhanov, P. in *The Oxford Handbook of Neolithic Europe*
781		(eds C. Fowler, J. Harding, & D. Hofmann) 195-214 (Oxford University Press,
782		2015).
783	32	Telegin, D. Ya. Neolithic cultures of Ukraine and their chronology. *Journal of World*
784		*Prehistory* **I**, 307-331 (1987).
785	33	Telegin, D. Ya. & Potekhina, I. D. *Neolithic cemeteries and populations in the*
786		*Dnieper Basin.* (British Archaeological Reports, 1987).
787	34	Jones, E. R. *et al.* The Neolithic Transition in the Baltic Was Not Driven by
788		Admixture with Early European Farmers. *Curr. Biol.*, 2185–2193 (2017).
789	35	Mittnik, A. *et al.* The Genetic History of Northern Europe. *bioRxiv*,
790		https://doi.org/10.1101/113241 (2017).
791	36	Saag, L. *et al.* Extensive farming in Estonia started through a sex-biased migration
792		from the Steppe. *Curr. Biol.* **27**, 2185-2193 (2017).
793	37	Maier, A. *The Central European Magdalenian: Regional Diversity and Internal*
794		*Variability.*, (Springer, 2015).
795	38	Borić, D. & Price, T. D. Strontium isotopes document greater human mobility at the
796		start of the Balkan Neolithic. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 3298-3303 (2013).
797	39	Krauß, R., Marinova, E., De Brue, H. & Weninger, B. The rapid spread of early
798		farming from the Aegean into the Balkans via the Sub-Mediterranean-Aegean
799		Vegetation Zone. *Quaternary International* **XXX**, 1-18 (2017).
800	40	Bacvarov, K. in *Moments in time: Papers Presented to Pál Raczky on His 60th*
801		*Birthday* (eds A. Anders & G Kulcsár) 29-34 (L'Harmattan, 2013).
802	41	Gurova, M. & Bonsall, C. 'Pre-Neolithic' in Southeast Europe: a Bulgarian
803		perspective. *Documenta Praehistorica* **XLI**, 95-109 (2014).
804	42	Brandt, G. *et al.* Ancient DNA reveals key stages in the formation of central
805		European mitochondrial genetic diversity. *Science* **342**, 257-261 (2013).
806	43	Borić, D. in *The Oxford Handbook of Neolithic Europe* (eds C. Fowler, J. Harding,
807		& D. Hofmann) 927–957 (Oxford University Press, 2015).
808	44	Szmyt, M. in *Transition to the Bronze Age (Archaeolingua 30)* (eds V. Heyd, G.
809		Kulcsár, & V. Szeverényi) 93-111 (Archaeolingua, 2013).
810	45	Olalde, I. *et al.* A Common Genetic Origin for Early Farmers from Mediterranean
811		Cardial and Central European LBK Cultures. *Mol. Biol. Evol.* **32**, 3132-3142 (2015).

46   Posth, C. *et al.* Pleistocene Mitochondrial Genomes Suggest a Single Major Dispersal
of Non-Africans and a Late Glacial Population Turnover in Europe. *Curr. Biol.* **26**,
827-833 (2016).

47   Anthony, D. W. & Ringe, D. The Indo-European Homeland from Linguistic and
Archaeological Perspectives. *Annual Review of Linguistics* **1**, 199-219 (2015).

48   Dabney, J. *et al.* Complete mitochondrial genome sequence of a Middle Pleistocene
cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. U. S.*
A. **110**, 15758-15763 (2013).

49   Meyer, M. & Kircher, M. Illumina sequencing library preparation for highly
multiplexed target capture and sequencing. *Cold Spring Harb Protoc* **2010**, pdb
prot5448 (2010).

50   Rohland, N., Harney, E., Mallick, S., Nordenfelt, S. & Reich, D. Partial uracil-DNA-
glycosylase treatment for screening of ancient DNA. *Philos. Trans. R. Soc. Lond. B*
Biol. Sci. **370**, 20130624 (2015).

51   Korlevic, P. *et al.* Reducing microbial and human contamination in DNA extractions
from ancient bones and teeth. *BioTechniques* **59**, 87-93 (2015).

52   Briggs, A. W. *et al.* Removal of deaminated cytosines and detection of in vivo
methylation in ancient DNA. *Nucleic Acids Res.* **38**, e87 (2010).

53   DeAngelis, M. M., Wang, D. G. & Hawkins, T. L. Solid-phase reversible
immobilization for the isolation of PCR products. *Nucleic Acids Res.* **23**, 4742-4743
(1995).

54   Rohland, N. & Reich, D. Cost-effective, high-throughput DNA sequencing libraries
for multiplexed target capture. *Genome Res.* **22**, 939-946 (2012).

55   Maricic, T., Whitten, M. & Pääbo, S. Multiplexed DNA sequence capture of
mitochondrial genomes using PCR products. *PLoS One* **5**, e14004 (2010).

56   Kircher, M., Sawyer, S. & Meyer, M. Double indexing overcomes inaccuracies in
multiplex sequencing on the Illumina platform. *Nucleic Acids Res.* **40**, e3 (2012).

57   Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler
transform. *Bioinformatics* **26**, 589-595 (2010).

58   Fu, Q. *et al.* A revised timescale for human evolution based on ancient mitochondrial
genomes. *Curr. Biol.* **23**, 553-559 (2013).

59   Skoglund, P. *et al.* Separating endogenous ancient DNA from modern day
contamination in a Siberian Neandertal. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 2229-
2234 (2014).

60   Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next
Generation Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014).

61   Price, A. L. *et al.* Principal components analysis corrects for stratification in genome-
wide association studies. *Nat. Genet.* **38**, 904-909 (2006).

62   Petkova, D., Novembre, J. & Stephens, M. Visualizing spatial population structure
with estimated effective migration surfaces. *Nat. Genet.* **48**, 94-100 (2016).

63   Brown, L., Cai, T. & DasGupta, A. Interval Estimation for a Binomial Proportion.
Statistical Science **16**, 101-133 (2001).

64   Agresti, A. & Coull, B. Approximate is better than "exact" for interval estimation of
binomial proportions. *The American Statistician* **52**, 119-126 (1998).

65   Longin, R. New method of collagen extraction for radiocarbon analysis. *Nature* **230**,
241-242 (1971).

66   Brown, T. A., Nelson, D. E., Vogel, J. S. & Southon, J. R. Improved collagen
extraction by modified longin method. *Radiocarbon* **30**, 171-177 (1988).

67   Kennett, D. J. *et al.* Archaeogenomic evidence reveals prehistoric matrilineal dynasty.
Nature communications **8**, 14115 (2017).

68   van Klinken, G. J. Bone collagen quality indicators for paleodietary and radiocarbon
measurements. *JAS* **26**, 687-695 (1999).

69   Stuiver, M. & Polach, H. A. Discussion: Reporting of [14]C data. *Radiocarbon* **19**, 355-
363 (1977).

70   Bronk Ramsey, C. *OxCal 4.23 Online Manual*,
867   <https://c14.arch.ox.ac.uk/oxcalhelp/hlp_contents.html> (2013).
868   71   Cook, G. T. *et al.* A freshwater diet-derived [14]C reservoir effect at the Stone Age sites
869   in the Iron Gates gorge. *Radiocarbon* **43**, 453-460 (2001).
870