Evolutionary rescue over a fitness landscape

Yoann Anciaux[1]; Luis-Miguel Chevin[2] ; Ophélie Ronce[1]; Guillaume Martin[1]

[1] ISEM, Université de Montpellier, CNRS, IRD, EPHE, Montpellier, France

[2] CEFE UMR 5175, CNRS - Université de Montpellier, Université Paul-Valéry Montpellier, EPHE, 1919 route de Mende, 34293 Montpellier, CEDEX 5, France

Contact : yoann.anciaux@umontpellier.fr

Keywords : Antibiotic resistance; Evolutionary rescue; Fisher's Geometric Model; Fitness landscape; Mutation

ABSTRACT

Evolutionary rescue describes a situation where adaptive evolution prevents the extinction of a population facing a stressing environment. Models of evolutionary rescue could in principle be used to predict the level of stress beyond which extinction becomes likely for species of conservation concern, or conversely the treatment levels most likely to limit the emergence of resistant pests or pathogens. Stress levels are known to affect both the rate of population decline (demographic effect) and the speed of adaptation (evolutionary effect), but the latter aspect has received less attention. Here, we address this issue using Fisher's Geometric Model of adaptation. In this model, the fitness effects of mutations depend both on the genotype and the environment in which they arise. In particular, the model introduces a dependence between the level of stress, the proportion of rescue mutants, and their costs before the onset of stress. We obtain analytic results under a strong-selection-weak-mutation regime, which we compare to simulations. We show that the effect of the environment on evolutionary rescue can be summarized into a single composite parameter quantifying the effective stress level, which is amenable to empirical measurement. We describe a narrow characteristic stress window over which the rescue probability drops from very likely to very unlikely as the level of stress increases. This drop is sharper than in previous models, as a result of the decreasing proportion of stress-resistant mutations as stress increases. We discuss how to test these predictions with rescue experiments across gradients of stress.

INTRODUCTION

Understanding the persistence or decline to extinction of populations facing environmental
stress is a crucial challenge both for the conservation of biodiversity and the eradication of
pests or pathogens (Gonzalez *et al.* 2013; Carlson *et al.* 2014; Alexander *et al.* 2014; Bell 2017).
In evolutionary biology, environmental stress describes any conditions in the environment that
induces a reduction in individual fitness (Koehn and Bayne 1989; Bijlsma and Loeschcke 2005).
Here, we will focus on the case where environmental stress causes a reduction of population
mean fitness that is harsh enough to trigger a decline in abundance (Hoffmann and Parsons
1997). In such a stressful environment, if heritable variation in fitness is available or arises by
mutation, adaptive evolution may allow the population to escape extinction. This phenomenon
has been called evolutionary rescue (ER) (Gomulkiewicz and Holt 1995). Evolutionary rescue is
of particular importance for understanding the emergence of genetic resistance to drugs or
treatments in medicine and agronomy (Davies and Davies 2010).

Empirical evidence supports the idea that stress levels critically determine ER probabilities
(Samani and Bell 2010; Moser and Bell 2011; Lindsey *et al.* 2013). For example, the probability
that bacteria evolve antibiotic resistance (that is, the probability of avoiding antibiotic-induced
extinction through ER) typically declines sharply, in a strongly non-linear way, with increasing
drug concentration (Drlica 2003). Evolutionary rescue thus shifts from being highly likely to
highly unlikely over a narrow window of stress levels. This critical range of stress depends on
the strain, especially on its evolutionary history with respect to exposure to the stress (Gonzalez
and Bell 2013). Stress level, as controlled by drug concentration, has also been shown to affect
the genetic basis of resistance (e.g. Harmand *et al.* 2017), with a wider diversity of genes and
alleles conferring resistance at low than at high doses. However, the underlying causes for the
relationship between stress level and ER are still poorly understood. Our aim here is to derive
new analytical predictions for this relationship. In particular, we want to predict the critical
window of stress levels above which ER is very unlikely, allowing direct comparison with
experimental data.

In the theoretical literature (reviewed in Alexander *et al.* 2014), most ER models predict that
ER probability decreases with increasing stress level, measured by the decay rate of the
stressed population. Indeed, a faster decay of the population leaves less time for adaptation to
occur before extinction (e.g. Gomulkiewicz and Holt 1995). But beyond this direct demographic

effect, stress level may also have indirect effects on ER. Indeed, a stressful environment may

not only affect the demographic properties of the population, but also its rate of adaptation,

by modifying the determinants of genetic variance in fitness (Hoffmann and Parsons 1997; De

Visser and Rozen 2005; Agrawal and Whitlock 2010). First, the rate of mutations and the

distribution of their effects on fitness change across environments (Martin and Lenormand

2006b; Wang *et al.* 2009; Agrawal and Whitlock 2010; Wang *et al.* 2014). In particular, the

fraction of beneficial mutations was found to increase in stressful environments (Remold and

Lenski 2001, 2004). Standing genetic variation for quantitative traits (notably fitness

components), also frequently depends on the environment (Hoffmann and Merilä 1999; Sgrò

and Hoffmann 2004; Charmantier and Garant 2005). Finally, the initial frequency of preexisting

variants able to rescue the population from extinction in a stressful environment may depend

on their selective cost in the past environment. In light of this empirical evidence, it seems clear

that progress towards understanding and predicting ER across stress levels requires addressing,

in a quantitative way, the joint effect of stress on the demography and genetic variation in

fitness of a population exposed to stressful conditions. This is our goal in the present article.

To do so, we develop a model that is a hybrid between two modeling traditions in ER theory,

summarized by Alexander *et al.* (2014): discrete genetic models, and quantitative genetic

models. Discrete genetic models assume a narrow genetic basis for adaptation (and ER),

whereby a single beneficial mutation can rescue an otherwise monomorphic population (Orr

and Unckless 2008; Martin *et al.* 2013; Uecker *et al.* 2014; Orr and Unckless 2014; Uecker and

Hermisson 2016). This approach was initially proposed for ER by Gomulkiewicz and Holt (1995),

and later extended to account for (i) evolutionary and demographic stochasticity (e.g. Orr and

Unckless 2008), and (ii) variation in the selection coefficients of mutations that may cause

rescue, with an arbitrary distribution of fitness effects (Martin *et al.* 2013). However, such

models do not predict how the distribution of fitness effects of mutations vary along gradients

of stress level. For this reason, they make it difficult to jointly address the two fundamental

components of stress mentioned above. On the contrary, quantitative genetics models of ER

inherently address the influence of stress on the rate of adaptation by assuming that adaptation

(and ER) is caused by evolution of a quantitative trait whose optimum changes with the

environment (Lynch *et al.* 1991; Burger and Lynch 1995; Gomulkiewicz and Holt 1995). In these

models, both the rate of population decline and the rate of adaptation under stress depend on

the distance between the phenotypic optima in the past and present environments. However,

analytical predictions are derived assuming a broad, polygenic basis for adaptation with a stable

genetic variance of the quantitative trait. The population genetic processes underlying

adaptation are not explicitly modelled, and the stochasticity involved in fixation and

establishment of mutations neglected. These complications are only explored by simulations

(e.g. Gomulkiewicz *et al.* 2010).

In order to take the best of both approaches, we rely on Fisher's (1930) Geometrical Model

(hereafter "FGM"). Fitness variation in the FGM is assumed to emerge from variation in multiple

putative phenotypic traits undergoing stabilizing selection that depends on the environment.

This model is analytically tractable, while retaining various aspects of realism (reviewed in

Tenaillon 2014). In particular, it accurately predicts how fitness effects of mutations change

across environments (Martin and Lenormand 2006b; Hietpas *et al.* 2013; Harmand *et al.* 2017)

or genetic backgrounds (Martin *et al.* 2007; MacLean *et al.* 2010; Trindade *et al.* 2012). The

FGM naturally relates environmental stress to (i) the rate of population decline, (ii) the rate and

effect of rescue mutants, and (iii) their potential costs in the past environment. Here, we

combine this FGM with population dynamic approaches that account for demographic and

evolutionary stochasticity (Martin *et al.* 2013), in a regime where selection is strong relative to

the rate of mutation. We consider rescue in asexual populations, stemming either from *de novo*

mutations or standing genetic variance. Interestingly, we show that all effects of stress on

demography and on the distribution of the fitness effects of mutations can be summarized into

a single composite measure of effective stress level. Evolutionary rescue shifts abruptly from

very likely to very unlikely over a narrow window of effective stress level, which can be

predicted from empirically measurable quantities.

METHODS

We here detail the ecological (environmental), genetic, and demographic assumptions of

the model, and the approximations used for its mathematical analysis.

Abrupt environmental shift: We define two environments: (1) a non-stressful one, denoted

as "previous environment", in which the population has a positive mean growth rate, and a

large enough population size that demographic stochasticity can be ignored; and (2) a stressful

one, denoted "new environment", in which the population initially has a negative mean growth

rate, and the population size is subject to demographic stochasticity. Conditions shift abruptly

from the previous to the new environment at $t = 0$, at which time the population size is N_0.

Eco-evolutionary dynamics: Extinction or rescue ultimately depends on details of the

stochastic population dynamics of each genotype. These are assumed to be mutually

independent (no density or frequency-dependence, see Chevin 2011), and sufficiently 'smooth'

(moderate growth or decay) that they can be approximated by a Feller diffusion (Feller 1951),

following Martin *et al.* (2013). This approximation reduces all the complexity of the life cycle

into two key parameters for each genotype i: its expected growth rate r_i (our fitness here),

and its variance in reproductive output σ_i. Our simulations below are performed for discrete

generations with Poisson offspring distributions. In this case, $\sigma_i = 1 + r_i \approx 1$ for any genotype,

as long as their growth rate is not too large ($r_i \ll 1$ per-generation, see **Appendix section I**

subsection 1 and 2 and Martin *et al.* (2013)). Note that the approximation extends to various

other forms of reproduction (see Martin *et al.* 2013).

To cause a rescue, a resistant mutant ($r_i > 0$) must establish, by avoiding extinction when

rare. The probability that this happens, for a lineage with growth rate $r_i > 0$ starting from a

single copy is $1 - e^{-2 r_i}$ (still assuming $r_i \ll \sigma_i$, with $\sigma_i \approx 1$ in the example used in simulations).

The number of individuals from which such mutations can arise declines in time, and we ignore

stochasticity in these decay dynamics. This is accurate as long as the population has large initial

size, of order $N_0 \gg 1$ (Martin *et al.* 2013).

Finally, we assume that mutation rates per capita per unit time are constant over time. This

is exact in models with discrete generations. In continuous-time models, where mutations

occur during birth events, mutation rates vary between genotypes with different birth rates,

and over time as these genotypes change in frequency. However, the constant mutation rate

model can still be approximately valid (see Martin *et al.* 2013).

ER from standing variance versus *de novo* mutation: At the onset of stress ($t = 0$), the
population either consists of a single ancestral clone, or is polymorphic at mutation-selection
balance in the previous environment. In the first case, we must derive the distribution of fitness
effects, in the new environment, of mutants arising from the ancestral clone. In the second
case, we must also describe the potential rescue variants already present in the previous
environment.

Mutations under Fisher's geometrical model (FGM): We assume that the expected growth
rate of a given genotypic class (its Malthusian fitness, or log-multiplicative fitness in discrete-
time models), is a quadratic function of its phenotype for n quantitative (continuous) traits.
Denoting as $\mathbf{z} \in \mathbb{R}^n$ the vector of breeding values (heritable components) for all traits, and as
$\mathbf{o}$ the single optimum phenotype with maximal growth r_{max}, the expected growth rate is

$$r(\mathbf{z}) = r_{max} - \|\mathbf{z} - \mathbf{o}\|^2/2 \,, \tag{1}$$

while the stochastic variance in reproductive success is assumed constant across genotypes.

The key assumption of our model is that the optimum depends on the environment. Without
loss of generality, we set the phenotypic origin at the optimum in the new environment, in
which $\mathbf{o} = \mathbf{0}$. In the previous environment, the optimum coincides with the mean phenotype
of the ancestral population ('A'): $\mathbf{o} = \mathbf{z}_A = \mathbb{E}(\mathbf{z})$, which implies that the ancestral population
was well-adapted in its original environment. The fitness of the mean ancestral phenotype $\mathbf{z}_A$
in the new environment is thus $r(\mathbf{z}_A) = r_{max} - \|\mathbf{z}_A\|^2/2 = -r_D < 0$, where r_D is its rate of
decay, and the phenotypic magnitude of the stress-induced shift of the optimum phenotype
(from $\mathbf{o} = \mathbf{z}_A$ to $\mathbf{o} = \mathbf{0}$) is $\|\mathbf{z}_A\| = \sqrt{2(r_D + r_{max})}$.

Mutations occur as a Poisson process with rate U per unit time per capita, constant over
time and across genotypes, but potentially variable across environments. Each mutation adds
a random perturbation $\mathbf{dz}$ to the phenotype, drawn from an unbiased and isotropic
multivariate Gaussian distribution $\mathbf{dz} \sim N(\mathbf{0}, \lambda \, \mathbf{I}_n)$, where $\mathbf{I}_n$ is the identity matrix in n
dimensions and λ is a scale parameter. Note that, since traits are not our main interest here,
we choose to measure mutation effects on them in units that directly relate to their fitness
effects. Therefore, λ can be understood as the variance of mutational effects on traits,
standardized by the strength of selection (see **Appendix section II subsection 1** for more details).

Note that mutation effects are additive on *phenotypes* (no epistasis), but not on *fitness*,

because $r(\mathbf{z})$ is nonlinear (Martin *et al.* 2007) .

Fig.1 illustrates the rescue process in the FGM. At the onset of stress ($t = 0$), the optimum

shifts abruptly to a new position, such that the mean growth rate becomes negative with $-r_D <$

0 (**Fig.1.C**). Meanwhile, the population size starts to drop from an initial value N_0 (**Fig.1.C**), facing

extinction in the absence of evolution. However, one or several mutants or pre-existing variants

may be close enough to the new optimum to have a positive growth rate ("resistant

genotypes", **Fig.1.A, 1.B**). These may then establish, and ultimately rescue the population

("rescue genotypes" **Fig.1.A, 1.B**).

Figure 1: Evolutionary rescue in Fisher's geometric model. In all panels, black refers to deleterious and neutral mutations ($-r_D \geq r$), blue to beneficial but not resistant mutations ($-r_D < r \leq 0$) and orange to resistant mutations ($r > 0$), around the dominant genotype of the ancestral population with phenotype $z_A \neq 0$. (A) Fitness landscape (FGM) with growth rate r (z-axis) determined by two phenotypic traits z_1 and z_2 . Dots represent the distribution of random mutant phenotypes around the dominant genotype of the ancestral population. The growth rate of this dominant genotype, in the stressful environment, is $-r_D$, and r_{max} is the maximal fitness at the

phenotypic optimum. (B) Distribution of growth rates among random mutants arising from the dominant genotype (distribution of mutation effects on fitness) for two decay rates $r_D = 0.1$ (left) and $r_D = 0.5$ (right). (C) Dynamics of the population size N_t and mean fitness $\bar{r}_t$ of a population starting from a clone at $-r_D = -0.083$ at size $N_0 = 10^5$. The black line represents the case without fixation of a beneficial mutation, the blue line the case with extinction in spite of the fixation of a beneficial, but non-resistant, mutation, and the orange line the case of a rescue. Parameters for the simulations are $r_{max} = 1.5$, $U = 2 * 10^{-5}$, $n = 4$ and $\lambda = 5 * 10^{-3}$.

Within the context of the FGM, increasing stress level may have different effects, also

discussed in Harmand *et al.* (2017). First, stronger stress may cause a larger shift in the position

of the optimum phenotype, resulting in a larger initial drop in fitness (higher r_D), as assumed in

most models of adaptation to a changing environment (Kopp and Matuszewski 2014). In

addition, the maximal possible fitness r_{max} may also be lower in the new than in the previous

environment (reduced environmental quality). Moreover, the mutational parameters (U and

λ) may change with stress, causing shifts in evolvability. Note that a change in λ may reflect a

change in the phenotypic effects of mutations, of the strength of stabilizing selection, or both.

For instance, higher stress may release cryptic genetic variance on underlying phenotypic traits

(Scharloo 1991; Hermisson and Wagner 2004), or cause increased mutation rates via SOS

responses in bacteria (Foster 2007). Finally, although less easy to conceptualize, some

environments may change the effective dimensionality of the landscape. However, in the

present paper, we only consider such changes in dimensionality in the context of rescue from

de novo mutations (where it can readily be handled by studying the effect of the parameter n).

As we will see, all our results can be expressed in terms of five parameters

($N_0 U$, r_D , r_{max} , λ, n). **Table 1** summarizes all notations in the article.

Notation	Description	Formula
N_t, N_0	N_t: population size at time t after the onset of the stress. N_0: initial population size at the onset of the stress.	
U	Mutation rate *per* individual *per* unit time.	
n	Number of traits under stabilizing selection, or phenotypic dimensionality.	
λ	Variance of mutational effects: variance of the phenotypic effects of mutations, per trait, in a trait space scaled by the strength of stabilizing selection	$\mathbf{dz} \sim N(\mathbf{0}, \lambda\, \mathbf{I}_n)$

U_c	Critical mutation rate below which the SSWM regime is valid.	$U_C = n^2 \lambda/4$
$\mathbf{z}$	n-dimensional vector $\mathbf{z} \in \mathbb{R}^n$ of (breeding values for) phenotype	
$\mathbf{z}_A, \mathbf{o}$	$\mathbf{o}$: optimal phenotype in a given environment $\mathbf{z}_A$: average phenotype of the ancestral population (before the onset of stress)	$\mathbf{o} = \mathbf{0}$: new environment $\mathbf{o} = \mathbf{z}_A$: previous environment
r, σ	Growth rate (r) and reproductive variance (σ) of a given genotype, in the new environment.	Eq.(1)
r_{max}	Maximum possible growth rate in the new environment.	$r(\mathbf{0}) = r_{max}$
r_D	Rate of decay of the ancestral phenotype $\mathbf{z}_A$ in the new environment.	$r(\mathbf{z}_A) = r_{max} - \|\mathbf{z}_A\|^2/2 = -r_D$
c	Cost of a mutation: selective disadvantage of the mutant, relative to the optimal phenotype, in the previous environment.	$c = \|\mathbf{z} - \mathbf{z}_A\|^2/2$
$c_H(y)$	Harmonic mean of the cost $c\|y$ among *de novo* mutations with scaled growth rate y in the new environment	Eq.(3)
y	Growth rate of a genotype, in the new environment, scaled by r_{max}.	$y = r/r_{max} \in [-\infty, 1]$
$f_y(y)$	Probability density function of y among random single step mutations	Eq.(2)
y_D	Rate of decay scaled by r_{max}.	$y_D = r_D/r_{max}$
ψ_D	Alternative measure of y_D	$\psi_D = 2\left(\sqrt{1 + y_D} - 1\right)$ (Eq.(6))
α	Effective stress level	Eq.(6)
α_c, r_D^c	Characteristic stress level (α_c) or decay rate (r_D^c) beyond which ER probability drops below $1/2$.	Eq.(8)
$g(\alpha)$	Function driving the dependence of rescue probabilities on stress levels.	Eq.(7)
$\omega_{DN}, \omega_{DN}^*$	ω_{DN}: rate of rescue from *de novo* mutations scaled by N_0 ω_{DN}^*: corresponding approximation when $\lambda \ll r_{max}$	Eqs.(5), (7)
$\omega_{SV}, \omega_{SV}^*$	ω_{SV}: rate of rescue from standing variance scaled by N_0 ω_{SV}^*: corresponding approximation when $\lambda \ll r_{max}$	Eqs.(5),(10)
P_R	Probability of rescue.	$P_R = 1 - e^{-N_0 \omega_R}$
ϕ_{SV}	Proportion of rescue events caused by standing variants	Eq. (10)

Table 1: Notations

Strong selection and weak mutation (SSWM) regime: The FGM described in the previous section, produces epistasis for fitness between different mutations which makes the problem highly intractable in general. To make analytical progress, we assume a regime of strong selection and weak mutation (SSWM, Gillespie 1983), which allows neglecting multiple mutants and epistasis. This regime arises when mutation rates are small relative to their typical fitness effect (as detailed below). In our context, this assumption implies that most rescue variants (pre-existing or *de novo*) are only one mutational step away from the ancestral genotype, allowing for two key simplifications. First, with a purely clonal ancestral populations, we can ignore ER by genotypes that have accumulated multiple *de novo* mutations. Second, in populations initially at mutation-selection balance, we can consider that all mutations arise from a single dominant genotype, optimal in the previous environment. Indeed, in the SSWM regime at mutation-selection balance, most segregating phenotypes remain within a narrow neighborhood of the optimum (relative to the magnitude of mutation effects on traits), so the mutation-selection balance is well-approximated by assuming that all mutations originate from the optimum phenotype. This is essentially the House-of-Cards approximation (Turelli 1984) extended to the FGM of arbitrary dimensionality (Martin and Roques 2016).

Overall, the SSWM assumption implies that the evolutionary aspects of ER are entirely determined by a single joint distribution of fitness, in the previous and new environment. This distribution corresponds to that of mutants arising from the optimal genotype of the previous environment. We thus apply the results of Martin *et al.* (2013), to this particular distribution.

Note that the SSWM approximations used in this article should apply even when multiple single-step mutants co-segregate (generating "soft selective sweeps", as detailed in Wilson *et al.* 2017). Indeed, the probability of ER, as computed for example in Orr and Unckless (2008) or Martin *et al.* (2013) and used here, is one minus the probability that no *single* mutant arises that ultimately causes ER. This means that we ignore ER requiring multiple mutational steps, but allow several single-step rescue mutations to co-segregate. Consistently, our simulations did not show any particular deviation from the theory at very mild stress, where such co-segregation of several single-step mutants is expected.

Maximal mutation rate for the SSWM regime: We conjecture that the SSWM approximation should be accurate below some threshold mutation rate U_C, i.e. whenever $U < U_C = n^2 \lambda/4$.

Indeed, Martin and Roques (2016) found that as long as $U \leq U_C$, the fitness distribution at

mutation-selection balance corresponds exactly to that expected under the House of cards

approximation (with a dominant optimal genotype plus its deleterious mutants). Whether this

same condition is sufficient for most rescue events to stem from single-step mutations is not

justified theoretically, and was simply tested by extensive stochastic simulations.

Supplementary Fig.1 further explores the range of validity of this approximation. It shows, in a

rescued population, the proportion of wild-type, single mutant, double mutant, and so on, as a

function of the mutation rate.

Distribution of single-step mutation fitness effects in the new environment: Let s be the

selection coefficient (difference in growth rate), in the new environment, of a random mutant

(with phenotype **z**) relative to its ancestor (with phenotype $\mathbf{z}_A$). The distribution of $s = r(\mathbf{z}) -$

$r(\mathbf{z}_A) = r + r_D$ among random mutants has a known exact form in the isotropic FGM (Martin

2014; Martin and Lenormand 2015), from which the distribution of growth rates ($r = s - r_D$)

in the new environment is readily obtained. It proves simpler and sufficient (see **Appendix**

section II subsection 2) to consider the scaled (and unitless) growth rate $y = r/r_{max} \in [-\infty, 1]$,

such that $y_D = r_D/r_{max} \in [0, +\infty]$ is the decay rate of the ancestor scaled to the maximum

possible growth rate. The scaled growth rates $y = r/r_{max}$ have the following probability

density function:

$$f_y(y) =$$

$$\exp\left(-\frac{r_{max}(2 + y_D - y)}{\lambda}\right)\left(\frac{r_{max}}{\lambda}\right)^{n/2}(1-y)^{n/2-1}\frac{{}_0F_1\left(\frac{n}{2}, \left(\frac{r_{max}}{\lambda}\right)^2(1+y_D)(1-y)\right)}{\Gamma(n/2)}, \qquad (2)$$

$$y \in]-\infty, 1]$$

where ${}_0F_1(.,.)$ is the confluent hypergeometric function and $\Gamma(z)$ is the gamma function. In

the SSWM regime, this probability density function approximately describes *de novo* mutations

produced after the onset of stress by the whole population, be it initially clonal or at mutation-

selection balance.

Fitness cost of single-step pre-existing mutants in the previous environment: Consider the

subset of random mutations, among those that arise from the dominant genotype of the

ancestral population, that have a scaled growth rate within the infinitesimal class $[y, y + dy]$

in the new environment. We introduce the conditional random variable $c|y$, which is the cost,

in the previous environment, of a random mutant within this subset (thus, conditional on y).

This cost is equal to the negative of the selection coefficient of the mutation relative to the

dominant genotype with phenotype $\mathbf{z}_A$. More precisely, the cost of a mutant with phenotype $\mathbf{z}$

is $c = \|\mathbf{z} - \mathbf{z}_A\|^2/2$ (using eq.(1), with $\mathbf{o} = \mathbf{z}_A$ for the previous environment). Note that,

because the mutation-selection balance in the previous environment is fully characterized by

relative fitnesses, which do not depend on the maximal growth rate in this environment, the

latter may differ from r_{max} without impacting the distribution of the costs $c|y$ and our results.

Importing results from Martin *et al.* (2013) for the SSWM regime, the total number of pre-

existing variants within the class $[y, y + dy]$ is Poisson distributed with mean $N_0 \, U \, f_y(y)/$

$c_H(y)dy$, where $c_H(y) = 1/\mathbb{E}_c(1/c|y)$ is the harmonic mean of the cost $c|y$ among

mutations with effect y in the new environment. This conditional harmonic mean depends on

the joint distribution of mutation effects on fitness (c, y) across two environments in the FGM

(given in Martin and Lenormand 2015). In our context, the dominant genotype of the ancestral

population is optimal in the previous environment and far from the optimum in the new

environment. In this case, using Eq.(9) in Martin and Lenormand (2015), the resulting

conditional harmonic mean $c_H(y)$ takes a tractable form (see Eq.(A6) for $n \geq 2$ and (A8) for

$n = 1$ in **Appendix section II subsection 4 and 5**):

$$c_H(y) = 1/\mathbb{E}_c\left(\frac{1}{c}\Big|y\right) = \begin{cases} \lambda \, v(y), & n = 1 \\ \dfrac{\lambda}{e^{v(y)} E_{(n-1)/2}(v(y))}, & n \geq 2 \end{cases}, \qquad (3)$$

$$\text{with } v(y) = \frac{r_{max}}{\lambda}\left(2 + y_D - 2\sqrt{(1 + y_D)(1 - y)} - y\right)$$

where $E_k(z) = \int_1^\infty e^{-z\,t}/t^k \, dt$ is the exponential integral function. In most of the article we

focus on the case $n \geq 2$, when considering ER from standing variance. The distributions of

mutation effects on fitness in both the previous (Eq.(3)) and the new environment (Eq.(2)) can

then be integrated to yield the probability of ER, as we show next.

General expression and assumptions for the rescue probability: Extinction occurs when no

resistant mutation manages to establish (i.e. to avoid stochastic loss). For compactness, we

define a rate of rescue ω per individual present at the onset of stress (i.e., scaled by N_0), such

that, following Martin *et al.* (2013), ER probabilities take the general form (similar to that in

e.g. Orr and Unckless 2008):

$$P_R = 1 - e^{-N_0\,\omega}\,. \tag{4}$$

The rate of rescue from *de novo* mutations alone is ω_{DN} ('DN' for *de novo*), while that from

pre-existing variance alone is ω_{SV} ('SV' for standing variants). For a purely clonal population,

the rate of rescue is $\omega = \omega_{DN}$, while for a population initially at mutation-selection balance, it

is $\omega = \omega_{DN} + \omega_{SV}$ in the SSWM regime assumed here (Martin *et al.* 2013). Applied to the

context of the FGM using Eqs.(2) and (3), the rates ω_{DN} and ω_{SV} are given by (see **Appendix**

Eq.(A5) and (A7)):

$$
\begin{aligned}
\omega_{DN} &= \frac{U}{r_D} \int_0^1 \pi(y) f_y(y)\,dy \\
\omega_{SV} &= U \int_0^1 \frac{\pi(y)}{c_H(y)} f_y(y)\,dy \\
&\text{with } \pi(y) = 1 - e^{-2\,y\,r_{max}/\sigma}
\end{aligned}
\tag{5}
$$

where $c_H(y)$, and $E_{(n-1)/2}(.)$ are defined in Eq.(3) and $\pi(y)$ is the probability of establishment

of a resistant genotype with scaled growth rate $y > 0$ in the new environment. ω_{DN} in Eq.(5)

is simply the average establishment probability of *de novo* resistant mutants times the genomic

mutation rate, divided by the rate of decay. In previous ER models (e.g. Orr and Unckless 2008;

Martin *et al.* 2013), which we denote "context-independent", the probability of rescue takes

the exact same form as Eq.(4). The expressions for the rates of rescue per capita also take a

form similar to Eq.(5): for *de novo* mutations, $\omega_{DN} = U\,q_R\,/\,r_D$, and for standing variance,

$\omega_{SV} = U\,q_R\,/\,c_H$, where q_R is the proportion of rescuers among random mutations ($q_R =$

$\int_0^1 \pi(y) f_y(y)\,dy$ in Eq.(5)) and c_H is again the harmonic mean of the cost of rescue mutations.

The important difference is that in previous models, q_R and c_H do not depend on r_D, while the

corresponding quantities in Eq.(5) do depend on the rate of decay, through its effect on $f_y(y)$

and $c_H(y)$.

The linearity of ER rates with the mutation rate U ($\omega \propto U$) arises here because of the SSWM

regime, where multiple mutations are ignored: it might not hold at higher mutation rates

(when $U > U_c$). As such, Eq.(5) makes no further assumption than the SSWM regime ($U < U_c$);

it can easily be evaluated numerically to provide a general testable theory for rescue

probabilities across stress levels, in the FGM. Yet, in order to gain more quantitative/intuitive

insight into the effects of stress, we study approximate closed forms for the rates in Eq.(5).

Small mutational effects approximation (SME): Although selection is assumed to be strong

relative to mutation ($U < U_c$, SSWM regime), it is still fairly realistic to assume that mutation

effects on traits (and thus fitness) are weak relative to the maximal growth rate in the new

environment, namely that $\lambda \ll r_{max}$. Taking a limit where $\lambda / r_{max} \to 0$, simpler expressions for

Eq. (5) are derived in the **Appendix section III**.

With this approximation, single-step resistance mutations are still rare and of large phenotypic

effect, in that they pertain to the tail of the mutant phenotype distribution. However, even

resistance mutations typically remain far from the optimum in the new environment, so that

their scaled growth rate is small: $y = r / r_{max} \ll 1$. Overall, mutation effects must fall within

the range: $4U/n^2 < \lambda \ll r_{max}$ for both the SSWM and the SME (small mutational effects)

approximation to apply (see **Appendix section III**). In the appendix, we study the convergence,

as λ / r_{max} decreases, of the results from Eq.(5) to their asymptotic limit (**Supplementary Figs.3**

and **4**).

Stochastic simulations of a discrete-time model: We checked the robustness of our

assumptions and approximations using stochastic simulations, where we tracked the

population size and genetic composition of a population across discrete, non-overlapping

generations. The size N_{t+1} of population at generation $t + 1$ was drawn as a Poisson number

$N_{t+1} \sim Poisson(N_t \overline{W})$, with $\overline{W} = \overline{e^r}$ the mean multiplicative fitness ($W = e^r$) and N_t the

population size, in the previous generation. The genotypes forming this new generation were

then sampled with replacement from the previous one with weight $W_i = e^{r_i}$. This is faster and

exactly equivalent to drawing independent Poisson reproductive outputs for each individual, or

genotype. Because of the underlying assumptions of the simulations, the corresponding

analytical approximation for the stochastic reproductive variance in Eq.(1) is $\sigma_i = \sigma \approx 1$

(assuming small growth rates $r_i \ll 1$). Mutations occurred according to a Poisson process, with

a constant rate U per capita per generation. Mutation phenotypic effects were drawn from a

multivariate normal distribution $N(\mathbf{0}, \lambda\, \mathbf{I_n})$, with multiple mutants having additive effects on

phenotype, and their fitness computed according to the FGM (Eq.(1)).

Rescue probability was estimated by running 1000 replicate simulations until either

extinction or rescue occurred. A population was considered rescued when it reached a

population size N_t and mean growth rate $\bar{r}_t$ such that its ultimate extinction probability, if it

were monomorphic, would lie bellow 10^{-12} ($\exp(-2\, N_t\, \bar{r}_t) < 10^{-12}$). This is a conservative

criterion: once $\bar{r}_t$ has become positive, we expect it to remain so, yielding further increases in

population size and thus further decreasing the probability of future extinction. We checked on

a subset of simulations that the above procedure gave the same rescue probabilities as

obtained in simulations performed until the population rebounded back to its (large) initial

size N_0.

For rescue from populations at mutation-selection balance, 8 replicate initial equilibrium

populations were generated, each by starting from an optimal clone and running the same

algorithm with fixed population size ($N_t = 10^6$) until the mean growth rate had visually

stabilized to a fixed value (close to its theoretical equilibrium value $\bar{r}_{eq} = r_{max} - U$ (for $U <$

U_c) for more than 1000 generations. Then the optimum was shifted by $\sqrt{2(r_D + r_{max})}$

phenotypic units, and 1000 replicate ER simulations were performed (same algorithm as for *de*

novo rescue), from each of the 8 replicate equilibrium populations.

All simulations and mathematical derivations were performed in *MATHEMATICA v. 9.0*

(Wolfram Research 2012).

<u>RESULTS</u>

The ER rates in Eq.(5) are analytical but only implicit functions of the model parameters. In

a small mutational effects (SME) limit, they take simpler closed form (indicated by a '*'). As we

will see below, these simpler forms mostly depend on the following two compound variables,

which summarize the various effects of stress on the fitness landscape:

$$\psi_D = 2\left(\sqrt{1 + r_D/r_{max}} - 1\right)$$
$$\alpha = \frac{\psi_D^2\, r_{max}}{4\,\lambda} \qquad . \qquad\qquad (6)$$

Both ψ_D and α increase with the decay rate r_D, decrease with increasing peak height r_{max}, and

are independent of n. The parameter α further increases with decreasing variance of

mutational effects λ. We can already see how α qualitatively reflects an "effective stress level":

stress is harder to cope with if decay rate is larger, the maximum growth rate is lower, and

mutation effects are smaller.

Rescue from *de novo* mutations: Under the SME approximation and in the SSWM regime

($4\,U/n^2 < \lambda \ll r_{max}$) the rate of *de novo* rescue (Eq.(5)), converges to (Eq. (A12) in the

Appendix):

$$\omega_{DN} \underset{\lambda \ll r_{max}}{\approx} \omega_{DN}^* = U \frac{(1+\psi_D/2)^{(1-n)/2}}{1+\psi_D/4}\, g(\alpha) \qquad (7)a$$

$$\omega_{DN}^* \underset{\psi_D/2\,\to\,0}{\to} U\, g(\alpha) \qquad\qquad (7)b \;, \qquad\qquad (7)$$

$$\text{with } g(\alpha) = \frac{e^{-\alpha}}{\sqrt{\pi\,\alpha}} - \text{erfc}\!\left(\sqrt{\alpha}\right)$$

where $\text{erfc}(.)$ is the complementary error function. Eq.(7)*b.* gives the approximate closed form

of Eq.(7)*a.* for mild stress ($\psi_D/2 \to 0$). Note that this approximation converges faster (with

decreasing ψ_D) with fewer dimensions, due to the faster vanishing of the factor $(1+$

$\psi_D/2)^{(1-n)/2}$ (in the limit $n = 1$ it vanishes for all ψ_D). We now discuss the biological

implications of these expressions.

Effect of FGM parameters on rescue: The partial derivatives of ω_{DN}^* in Eq.(7) with respect to

the FGM parameters (r_D, r_{max}, λ, n) quantify the sensitivities of ER probability to each of them

(**Appendix section III subsection 4**). First, note that $g(.)$ is a strictly decreasing function of α.

When $n > 1$ and with mild stress ($\psi_D \ll 2$), Eq.(7)*b.* applies and $\omega_{DN} \approx U\,g(\alpha)$. ER then

becomes less likely with a higher decay rate r_D, a lower peak r_{max} and a smaller variance of

mutational effects λ, and is independent of dimensionality n. For stronger stress levels, Eq.(7)*a.*

applies: these qualitative dependencies to the parameters still hold, except that ER probability

now decreases with increasing dimensionality.

Sharp drop of ER probability with stress levels: **Fig.2** shows the agreement between

simulations (stochastic discrete-time demographic model, see Methods) and the analytical

expressions in Eqs.(5) and (7), over a wide range of stress levels (quantified by r_D), and for two

values of r_{max} and U (**Supplementary Fig.3** further explores the range of validity of the

approximation). Interestingly, ER probability drops sharply with stress levels (with decay rate

r_D here), which is well captured by the term $g(\alpha)$ alone (Eq.(7)*b.*, dashed red lines in **Fig.2**).

This drop is much more pronounced than in a context-independent model (gray lines in **Fig.2**),

where stress does not affect the distribution of mutation effects. The difference between

context-independent models and the FGM is that, in the latter, increased stress implies both

faster decay (as in the former), and fewer and weaker resistance mutations. In the FGM, these

effects on the properties of rescue mutations are the main drivers of ER probabilities across

stress levels.

Figure 2: Rescue probability from *de novo* mutations. The ER probability as a function of stress levels, expressed as the initial mean decay rate of the population, is given for various values of the mutation rate $U = 10^{-3}\, U_c$ (blue) or $U = 10^{-2}\, U_c$ (orange) and the maximal fitness reachable in new the environment $r_{max} = 0.5$ (A) or 1.5 (B). Dots give the results from simulations and solid lines (blue and orange) show the corresponding theory computed numerically (Eq.(5)). The black dot-dashed (respectively red dashed) lines give the corresponding analytical approximations Eq.(7)*a.* (respectively Eq.(7)*b.*). The gray lines correspond to an equivalent theory without the FGM (named context-independent model as described in the Method section, "CI") modified from Orr and Unckless (2008). This last model was computed using a fixed proportion of resistant mutations equal to the one in Eq.(5) for a rescue probability of 0.5 (which explains why the two curves cross exactly at $P_R = 0.5$). Other parameters are $n = 4$, $N_0 = 10^5$, $\lambda = 0.005$.

Composite parameter α describing an effective stress level: The results in Eqs.(6)-(7) suggest

that a single composite parameter (α in Eq.(6)) can capture the various ways in which stress

may alter the parameters of the fitness landscape, that is, fitness peak height r_{max}, variance of

mutational effects λ, or distance to the optimum r_D. We denote this parameter the "effective

stress level". In **Fig.2**, considering the effect of stress only via α (Eq.(7)*b.*) is equally accurate as

using the more complex Eq.(7)*a.*, or the numerical computation from Eq.(5).

This simplification is further illustrated in **Fig.3**, where we use exact numerical computations

from Eq. (5) to explore different possible effects of stress. Regardless of whether stress affects

only the maladaptation of the ancestral clone (r_D, blue symbols), or also the quality of the

environment (joint change in r_D and r_{max}, orange symbols) or the variance of mutational

effects (joint change in r_D and λ, red symbols), its effect on the rescue probability is accurately

predicted by α (**Fig.3B,** black line). As predicted by Eq.(7)*b.*, the relationship between rescue

probability and α is approximately independent of dimensionality (compare circles $n = 1$ and

squares $n = 6$ in **Fig.3B**). We also note that Eq.(7)*b.* slightly overestimates the 'exact' numerical

computations of the ER probability from Eq.(5), so it provides a conservative bound when

considering the control of resistant pathogens.

Figure 3: The effective stress level. Rescue probability for clonal populations versus initial decay rate r_D (A) or the effective stress level α (B). In both panels the axes are in logarithmic scale, symbols show Eq.(5) and colors refer to different effects of increased stress level: blue symbols show only r_D increasing (with $r_{max} = 1.5$ and $\lambda = 0.005$), orange symbols r_D increasing and r_{max} decreasing linearly with r_D (according to $r_{max} = 1.5 - 5\,r_D$, with $\lambda = 0.005$) and red symbols r_D increasing and λ decreasing linearly with r_D (according to $\lambda = 0.005 - 10^{-2}\,r_D$, with $r_{max} = 1.5$). In each case, the results for both $n = 1$ (circles) and $n = 6$ (squares) are shown. The black plain line on the right panel gives the result from Eq.(7)*b.*: a single composite measure of stress (α) approximately

captures the impact of stress-induced variations in the various parameters $(r_D, r_{max}, \lambda, n)$. Other parameters are $N_0 = 10^6$ and $U = 2 * 10^{-5}$.

Characteristic stress level: **Fig.2** shows that ER drops from highly likely to highly unlikely

around a "characteristic stress level", which we can be characterized analytically (as detailed in

the **Appendix section IV subsection 1**). Consider the set of values of parameters

$(N_0, U, r_{max}, r_D, \lambda, n)$ for which the rescue probability is of given value $p \in [0,1]$. From Eq.(7)b.,

this corresponds to the set (r_{max}, r_D, λ) for which $\alpha = g^{-1}(-\log(1-p)/N_0 U)$. Using the

approximation $g(\alpha) \approx \alpha^{-3/2} e^{-\alpha}/(2\sqrt{\pi})$ (from Eq.(7) with $\alpha \gg 1$), the corresponding α can

be derived explicitly (Eq.(A17)). In particular for $p = 1/2$, the characteristic stress level α_c at

which the ER probability is $1/2$ is (Eq. (A19) in the **Appendix**):

$$\alpha_c \underset{N_0 U \gg 1}{\approx} g^{-1}\left(\frac{\log(2)}{N_0 U}\right) \underset{N_0 U \gg 1}{\approx} 0.9 \log(N_0 U) - 2.7 \,. \tag{8}$$

Under the conditions of the SME approximation (detailed in Methods), Eq.(8) applies for large

$N_0 U$ (approximately when $N_0 U \geq 5.10^4$), a necessary condition for this equation to be self-

consistent (detailed in **Appendix section IV subsection 2**).

The characteristic stress level α_c that a population can typically withstand increases only log-

linearly with population size and mutation rate. Consider the characteristic decay rate r_D^c for

which the rescue probability is $P_R = 1/2$, i.e. the decay rate that populations can overcome

half of the times. From Eq.(8) with $\alpha_c \propto (r_D^c)^2$ (Eq.(6)), this decay rate is $r_D^c \propto \sqrt{\log(N_0 U)}$ for

large $N_0 U$. For comparison, we would have $r_D^c = q_R N_0 U/\log(2)$, which is linear in $N_0 U$, in a

context-independent model where the proportion q_R of random mutations causing a rescue is

independent of r_D. The difference in the effect of $N_0 U$ on rescue probability thus stems from

the strong non-linearity (i.e. sharp drop) of rescue probability with stress level (decay rate)

under the FGM. In the FGM, overcoming a given environmental harshness requires much more

mutational input than in a context-independent model.

Characteristic stress window: It is also important to predict how sharply the ER probability

drops around the characteristic stress level. This drop can be characterized by a "characteristic

stress window" of α over which the ER probability drops from 75% to 25%. The width $\Delta\alpha$ of this

window can be scaled by the value of the characteristic stress level α_c, to get a scale-free

measure of its steepness (i.e., how sharp the drop in ER probability is, relative to the stress level

around which it occurs). This gives (from Eq.(A20) in the **Appendix**):

$$\frac{\Delta\alpha}{\alpha_c} \underset{N_0 U \gg 1}{\approx} \frac{1}{1 + 0.7\alpha_c}. \tag{9}$$

The width $\Delta\alpha$ increases with increasing $N_0 U$ until it saturates (at ~ 1.5) (see **Appendix section**

IV subsection 3 for further details). However, when scaled by the center of the window (α_c),

the width of this scaled characteristic stress window drops below 1 as α_c increases (and hence

with increasing $N_0 U$, from Eq.(8)). This result shows formally that the drop in ER probability

with increasing stress α gets proportionally sharper (relative to the position where it occurs)

as $N_0 U$ increases, but this is entirely driven at large $N_0 U$ by shifts of a window with constant

absolute width. This is illustrated in **Fig.4**, which also shows the accuracy of Eq.(9) compared to

'exact' numerical computations from Eq.(5) (as expected, the exact result deviates from Eq.(9)

for smaller $N_0 U$).

Figure 4: Scaled width of the characteristic stress window $\Delta\alpha/\alpha_c$ versus the population-scale mutation rate $N_0 U$. The dots are obtained by numerical inversion of the 'exact' Eq.(5) with two values of $r_{max} = 2$ (blue) and $r_{max} = 0.1$ (red). The orange line shows the approximate scaled width of the characteristic stress window derived in Eq.(9). Other parameters are $n = 4, \lambda = 0.005$.

Interestingly, Eq.(9) provides a scale-free measure that may be compared across

experiments, as it only depends on the genomic mutational input $N_0 U$ (via α_c). However, like

all results so far, Eq.(9) only considers ER from *de novo* mutation. We now turn to ER from

standing genetic variation.

Rate of rescue from a population at mutation-selection balance: In the SSWM regime, and for

a population at mutation-selection balance in the previous environment, each rescue event can

be tracked back to either a pre-existing variant, or a *de novo* mutation (Orr and Unckless 2008;

Martin *et al.* 2013; Orr and Unckless 2014). The proportion ϕ_{SV} of rescue events caused by

standing variance is then simply given by $\phi_{SV} = \omega_{SV}/(\omega_{SV} + \omega_{DN})$. A simple expression can

be obtained again under the SME for $n \geq 2$ (see Eq.(A21) in the **Appendix**), but the

approximation (ϕ_{SV}^*) now further requires that decay rates are not vanishingly small ($\lambda \ll$

$r_{max} \psi_D^2/4$).

$$\phi_{SV} = \frac{\omega_{SV}}{\omega_{SV} + \omega_{DN}} \underset{\lambda \ll r_{max} \psi_D^2/4}{\approx} \phi_{SV}^* = \frac{1 + \psi_D/4}{\epsilon/\psi_D + 1 + \psi_D/2} \text{ where } \epsilon = \lambda \frac{n-1}{2\, r_{max}}, \qquad (10)$$

where ψ_D is defined in Eq.(6).

Eq.(10) captures the main features of how standing variance contributes to ER across (non-

vanishing) stress levels (here, decay rate). Contrary to context-independent models, this

contribution changes non-monotically with increasing stress level (**Fig.5**). At very mild decay

rate r_D, rescue relies on mild-effect mutations. The cost of such mutations - and hence their

frequency before stress - is roughly independent of r_D (Martin and Lenormand 2015), while

their rate of production by *de novo* mutation decreases as $1/r_D$ (demographic effect), so the

contribution of standing variance to ER increases with r_D at small r_D. In contrast at large stress

levels, rescue stems from strong effect mutations. These mutations pay a substantial

"incompressible cost" before stress that increases faster than r_D (Martin and Lenormand 2015),

while their rate of *de novo* production still decreases as $1/r_D$, so the contribution of standing

variance to ER decreases with r_D at large r_D. In the limit of very large y_D, the distance between

the two optima is very large and makes most of both the cost and decay rate, so that $c_H(y) \approx$

r_D for all mutations. Hence, *de novo* mutations and standing variants contribute equally to ER

in this limit ($\phi_{SV} \to 1/2$).

Figure 5: relative contribution of standing genetic variation to ER. The proportion of ER from standing variance is shown across scaled decay rates y_D. The numerical computation for ϕ_{SV} (using Eq.(5) and (10)), for two values of $r_{max} = 0.1$ (blue) and $r_{max} = 0.7$ (orange), is compared to the approximate ϕ_{SV}^* (Eq. (10), black dashed line). Other parameters as in **Fig.2**.

These different behaviors are illustrated in **Fig.5**, showing the variation of ϕ_{SV} over a very
wide range of scaled decay rates y_D. The limit $\phi_{SV} \to \phi_{SV}^*$ in Eq.(10) provides a fairly accurate
approximation across the full range of stress levels. The limits when $y_D \to 0$ and $y_D \to \infty$ are
in fact of limited biological interest, as they correspond to stress levels where ER becomes *de*
facto certain or impossible, respectively. When focusing on the more biologically relevant range
corresponding to the characteristic stress window, which occurs near the peak in ϕ_{SV} in **Fig.5**
(see **Appendix Section IV subsection 4**), the variation of ϕ_{SV} across stress levels becomes
negligible. As illustrated in **Fig.6B**, ϕ_{SV} remains close to $1 - \sqrt{\epsilon}$ (see **Appendix** Eq.(A22)) as y_D
varies over a range where ER probabilities span several orders of magnitude (see **Appendix**
Section IV subsection 5). Note that this behavior arises when stress only shifts the optimum
(effect on r_D), but does not affect peak height (r_{max}) or the variance of mutational effects (λ).

Therefore the rate of rescue in the presence of standing variance is approximately
proportional to that with only *de novo* mutation, with proportionality constant largely
independent of the decay rate:

$$\omega = \omega_{DN} + \omega_{SV} \approx \omega_{DN}^*/\sqrt{\epsilon}, \tag{11}$$

where ϵ is defined in Eq.(10). The rough constancy of ϕ_{SV} also means that all the results
obtained previously for ER from *de novo* mutations apply in the presence of standing variance,
when stress only shifts the optima (as long as $n \geq 2$).

The ER probability profile across stress levels (shown in **Fig.6A**) is the same as that from *de*

novo mutations alone (**Fig.2**), but with a higher characteristic stress ($\alpha_c \underset{N_0 U \gg 1}{\approx} 0.9 \log(X) - 2.7$

from Eq.(8) with $X \approx N_0 U / \sqrt{\epsilon}$). Moreover, when considering the contribution from standing

variance, the difference between the FGM and context-independent models (gray line on

Fig.6A) is striking. Indeed in the latter, the ER rate from standing variance (ω_{SV}) is independent

of the decay rate, hence P_R saturates with stress to a constant value $1 - \exp(- N_0 U q_R / c_H)$,

where all ER events stem from standing variance.

Figure 6: ER probability in the presence of standing genetic variation. In each panel, stress only affects the decay rate r_D (shifting optimum). In both panels, blue solid lines show the theory for *de novo* and standing variance ('DN'+'SV') computed numerically (Eq.(5)) and the gray lines correspond to an equivalent theory without the FGM (named context-independent model as described in the Method section, "CI") modified from Orr and Unckless (2008). This last model was computed using a fixed proportion of resistant mutations equal to the one in Eq.(5) for a rescue probability of 0.5 (which explains why the two curves cross exactly at $P_R = 0.5$). The dashed red line gives the simpler expression for the overall rescue rate: $\omega \approx \omega_{DN}^* / \sqrt{\epsilon}$. (Eq.(11)) with ϵ given in Eq.(10) and ω_{DN}^* by Eq.(7). (A) ER probability in the presence of standing genetic variation as a function of r_D. The dots give the results from simulations. (B) Proportion ϕ_{SV} of rescue from standing variance as a function of r_D. The black dashed-line give the approximate theory from Eq. (10) and the dashed red line $\max(\phi_{SV}^*) \approx 1 - \sqrt{\epsilon}$ from Eq.(11). The shaded area shows the range of r_D for which the ER probability drops from 0.99 to 10^{-3}. Other parameters as in **Fig.2**.

520   Finally, note that when considering rescue from preexisting variance, U may change across

521   environments, from U_P (for previous) to U_N (for new). For example, a stress-induced increase

522   in DNA copy error would yield $U_N > U_P$. Accounting for such shifts in mutation rate at the onset

523   of stress, the total ER rate simply becomes $\omega \approx \omega_{DN}^* \left(1 + U_P / U_N \left(1 - \sqrt{\epsilon}\right) / \sqrt{\epsilon}\right)$ (from

524   Eq.(11)).

DISCUSSION

Main results: We investigated the persistence of a population of asexual organisms under an abrupt environmental alteration. We assumed that this stress causes a shift in a multidimensional fitness landscape with a single peak (Fisher's Geometrical Model - FGM -), which the population must 'climb' to avoid extinction. In such a landscape, faster population decline (due to stress-induced increase in the decay rate r_D) necessarily means that resistance mutations are fewer, have lower growth rates in the presence of stress and higher costs in its absence. We believe that this constraint, not included in previous studies, adds a key element of realism to evolutionary rescue (ER) models. In our model, variation in stress levels may affect the landscape in various ways: shifting the optimum, changing the peak height, or altering the phenotypic scale of mutations (or the strength of stabilizing selection). Under a strong selection and weak mutation (SSWM) regime and assuming small mutational effects (SME), all these effects of stress on the distribution of mutation fitness effects are approximately captured by the variation, across stress levels, of a single composite parameter α, which is approximately independent of the dimensionality of the organism (number of orthogonal traits under selection). The probability of ER drops sharply with this effective stress level, more so than in previous ER models where the rate of population decline is decoupled from the input of resistant mutations. The characteristic stress window over which this drop occurs only depends on the initial population size N_0 and genomic rate of mutation U. As N_0U gets large, the characteristic stress window reaches an asymptotic width ($\Delta\alpha$ in Eq.(9)) while its center (the characteristic stress level α_c in Eq.(8)) shifts towards higher values, approximately as $\log(N_0U)$.

When standing variance is available (population at mutation-selection balance before stress), its contribution to ER is dominant, and approximately constant across a wide range of stress levels that encompasses the characteristic stress window.

In **Table 2**, we summarize how these features compare to properties of previous ER models. We consider only the situation where stress shifts the position of the optimum, affecting r_D (as in previous models), because other effects of stress we investigate here (r_{max} and λ) are not treated in previous models.

	Context-independent model	FGM rescue model
Impact of decay rate r_D	$\omega_{DN} \propto r_D^{-1}$, $\omega_{SV} = $ constant	$\omega \propto e^{-r_D^2} r_D^{-3}$ (1)
Impact of mutational input $N_0 U$	$\omega \propto N_0 U$	$\omega \propto N_0 U$
"Characteristic decay rate" (r_D^c) for DN rescue versus $N_0 U$	$r_D^c \propto N_0 U$	$r_D^c \propto \sqrt{\log(N_0 U)}$
Relative contribution of standing variance to ER	Increases with r_D $$\phi_{SV} \approx \frac{r_D}{r_D + c}$$	~ Stable with r_D $$\phi_{SV} \approx 1 - \sqrt{\epsilon}$$

Table 2: Main results of "context-independent" models (Orr and Unckless 2008; Martin *et al.* 2013; Orr and Unckless 2014) and the present model (FGM) when stress only affects r_D (only shifts the optimum position in our model). When the dependence to the parameters is given for the overall rate of rescue ω, it applies to both the rate of rescue from *de novo* mutations ω_{DN} and from pre-existing variance ω_{SV}. (1) derived from the approximate expression $g(\alpha) \approx \alpha^{-3/2} e^{-\alpha}/(2\sqrt{\pi})$.

Genetic basis of ER patterns across environments: Our model allows identification of three

ranges of stress levels that yield different eco-evolutionary patterns, despite all leading to

extinction in the absence of evolution. First at low stress levels ($\alpha \ll \alpha_c$), although evolutionary

change is required for persistence and demographic dynamics typical of ER may be observed

(decay /rebounce), extinctions are *de facto* undetectable (ER is pervasive, $P_R \approx 1$). In this

regime, we expect several resistance mutations to establish and co-segregate (frequent "soft

sweeps" as in Wilson *et al.* 2017). Their number is predictable ($\approx N_0 \omega_R$), but the ultimate

composition of the population in asexuals will depend on more complex clonal interference

dynamics. Second, at intermediate stress levels ($\alpha = O(\alpha_c)$), small variation in stress

conditions has large impact on the probability of population survival. Over this range, $P_R \approx 1/2$

so the expected overall number of rescue mutations in the population is less than one ($N_0 \omega_R \approx$

$-\log(P_R) = 0.7$). Therefore, "hard sweeps" (including from standing variation) should be the

most frequent: a single mutation typically establishes and rescues the population. Finally, at

higher stress levels ($\alpha \gg \alpha_c$), very few populations overcome the imposed stress, and when

they do it is typically through a hard sweep ($N_0 \omega_R \ll 1$).

Estimating parameters and testing the model: Studies on the emergence of resistance to

controlled stress (e.g. antibiotics, fungicides, chemotherapy in cancer), especially in microbes,

can generate a set of estimates of P_R (the probability of resistance emergence), across stress

levels. In general, to test (or use) predictions from ER models, it is critical to empirically relate

physical measures of stress level (e.g. concentrations, temperatures, salinities, etc.) with

demographic measures (e.g. decay rates). If we assume that the main effect of stress is to shift

optimum positions, given a set of measurements of r_D ("dose-kill curves", Regoes *et al.* 2004),

the change in ER probability with stress can be predicted via Eq.(7). This simple scenario of

optimum shifting is the one that is considered by most of the literature on evolutionary ecology

across environmental gradients, so it would seem natural to test it first. Furthermore, this

scenario has received empirical support from an analysis of a few experimental studies of

distribution of mutation effects on fitness across stress levels (Martin and Lenormand 2006b).

However, these studies used mild stresses, which reduce growth without causing population

decay. A more recent study on bacteria facing lethal doses of antibiotics, i.e. in the presence of

decay (Harmand *et al.* 2017), suggests that factors other than the position of the optimum may

also change with stress (λ, r_{max}, n). Estimating these extra parameters across environments can

be challenging. The variance of mutational effects λ and the dimensionality n can be estimated

by fitting the distributions of single random mutation effects on fitness (Martin and Lenormand

2006b; Perfeito *et al.* 2014), in a single environment if it is to be assumed constant, or in each

environment otherwise. The maximal growth rate in the stress could be measured on lines well-

adapted to the environment considered.

The effective stress level α is also amenable to empirical measurement and circumvents the

issue of measuring joint changes in (r_D, r_{max}, λ) with stress. Consider a set of P_R estimates

across a range of empirically controlled stress levels, and some knowledge of the genomic

(non–neutral) mutation rate (e.g. as estimated by mutation accumulation experiments) of the

species and environment under study (U). The initial population size N_0 is easily controlled by

the experimenter. Then Eq.(7)b. suggests a simple estimator of α in each environment: $\hat{\alpha} =$

$g^{-1}(-\log(1 - P_R)/N_0 U)$.

Finally, it is also possible to circumvent the problem of stress-induced variation in the

parameters of the fitness landscape by considering multiple genetic backgrounds, in a single

environment. Each background would have a given measurable decay rate r_D, and other

parameters (λ, r_{max}, n, U) would be held fixed: while λ, n and U may change with the genetic

background, this seems less likely than with the environment. The isotropic FGM assumes a

strict equivalence between shifts in optima (multiple environments) from a given ancestor

phenotype (single genetic background), and shifts in ancestor phenotypes (multiple genetic

backgrounds) with a fixed optimum (single environment). The model could thus be applied and

tested in this context. This could yield useful insights into the effect of epistasis (background

dependence) on resistance emergence, an issue of notable importance when considering the

fate of horizontally transferred resistance or multidrug resistance (as discussed in Wong 2017)

Potential implications for resistance management: Our results suggest that stress levels have

a strongly nonlinear impact on ER probabilities (**Fig.2**), at least in the context of abrupt

environmental changes, in asexuals. This context may be particularly relevant to the chemical

treatment of pathogens (cancer therapy, antivirals, antibiotics, fungicides, herbicides, etc.). In

particular, the non-linear impact of stress on ER, if empirically confirmed, can provide insights

regarding the optimization of treatment regimens or the quantification of the effect of poor

treatment adherence on resistance emergence (e.g. in HIV, Harrigan *et al.* 2005). Our results

point to the risk that even a slight lowering of drug doses (bellow prescribed treatment levels)

could radically change the outcome of the treatment (making it *de facto* inefficient). On the

contrary, a slight increase in prescribed doses could sometimes prove sufficient to allow

efficient eradication.

Limits and possible extensions :

<u>Density-dependence and competitive release:</u> Our model ignores density dependence, but

some form may be easily introduced by considering a single density dependence coefficient

(common to all genotypes) and using logistic diffusion approximations (Lambert 2005). This

would potentially allow for "competitive release" (Read *et al.* 2011), whereby higher stresses

may favor the emergence of resistance by rapidly depleting the sensitive wild-type population,

thus releasing limiting resources for resistant genotypes. Previous models on competitive

release assumed that the number of standing resistant mutants is independent of stress level

(Read *et al.* 2011; Day and Read 2016). In this case, stress mostly limits *de novo* rescue

mutation, with limited impact on the contribution from standing variance. On the contrary, the

FGM imposes a similar drop, with stress, in the rate of rescue from *de novo* and preexisting

mutants. The positive effects of competitive release on ER probability may thus be less

important, in the FGM, than predicted from these previous models. Note however that more
generally, the effect of density dependence on ER is more safely investigated by accounting
explicitly for the effect of stress on the density-independent intrinsic rate of increase on the
one hand (as done here in a density-independent model), versus on the competition
component on growth (e.g. carrying capacity) on the other hand, than by compounding their
effects into an overall density-dependent decay rate (e.g. Chevin and Lande 2010). This would
require modeling the effect of stress on both the intrinsic rate of increase and the competition
component, possibly through a landscape with two fitness functions, describing each
component.

Anisotropy and parallel evolution in drug resistance: The present model is isotropic: all
directions in phenotype space are equivalent (in terms of mutation and selection). In contrast,
module-dependent anisotropy (where particular genes mutate along favored directions) can
lead to substantial parallel evolution in the FGM (Chevin *et al.* 2010). Parallel evolution of
resistance, whereby some (portions or sets of) genes contribute most of the resistance
mutations, is often observed among drug-resistance alleles, and can increase with stress
(Harmand *et al.* 2017), contrary to parallel evolution in a growing population, which is expected
to decrease with increasing maladaptation (Chevin *et al.* 2010). Although not explored here,
we conjecture that our model may accommodate mild anisotropy. Indeed, mild anisotropy
(even environment-dependent) might have limited impact. If mutational covariances between
traits merely "turn", "shrink" or "expand" the phenotypic mutant cloud, this would
approximately amount to a mere change in λ in an equivalent isotropic landscape (Martin and
Lenormand 2006a; Martin 2014). However, a particular form of strong anisotropy may also
arise where mutant phenotypes (in a given module) spread along a single favored direction
(Martin 2014). Only this level of anisotropy would generate clear parallel evolution, and it will
likely require implementing a fully anisotropic model.

High mutation rates: Our results relied on a strong selection and weak-mutation (SSWM)
approximation. When the mutation rate is higher (e.g. viruses or mutator bacterial strains),
multiple mutants must be accounted for as a source of ER. These can in principle be introduced
in the framework used here (Martin *et al.* 2013) but, especially when applied to the FGM, the
results quickly become intractable. Alternative population genetics assumptions would then
have to be used, but this is beyond the scope of the present work.

Conclusion: Recently, the FGM has received renewed interest for its ability to provide

testable, quantitative, and often accurate predictions regarding patterns of mutation effects

on fitness, across various species and contexts (Tenaillon 2014). The present model is an

attempt to extend its scope to model the evolution of resistance to stress. We hope that future

experimental tests will evaluate its accuracy and potential to tackle various pressing applied

issues.

Acknowledgments: We thank the editor Joachim Hermisson and the reviewers for their help

in improving this manuscript. In particular we thank Michael Kopp for extensive and very useful

comments on our manuscript. This work benefited from discussions with S. Gandon, R.

Gomulkiewicz, O. Tenaillon, F. Débarre and L. Roques. This work was supported by the French

Ministry of Higher Education, Research and Innovation (MESRI allocation doctorale to Y.A.),

Agence Nationale de la Recherche (ANR-13-ADAP-0016 "Silentadapt" to G.M. and ANR-13-

ADAP-0006 "MeCC" to O.R.) and European Research Council (ERC-2015-STG-678140-FluctEvol

to LMC). This is ISEM publication ISEM 2018-032.

BIBLIOGRAPHY

Agrawal A. F., Whitlock M. C., 2010 Environmental duress and epistasis: how does stress affect the strength of selection on new mutations? Trends Ecol. Evol. 25: 450–458.

Alexander H. K., Martin G., Martin O. Y., Bonhoeffer S., 2014 Evolutionary rescue: linking theory for conservation and medicine. Evol. Appl. 7: 1161–1179.

Bell G., 2017 Evolutionary Rescue. Annu. Rev. Ecol. Evol. Syst. 48: 605–627.

Bijlsma R., Loeschcke V., 2005 Environmental stress, adaptation and evolution: an overview. J. Evol. Biol. 18: 744–749.

Burger R., Lynch M., 1995 Evolution and Extinction in a Changing Environment: A Quantitative-Genetic Analysis. Evolution 49: 151.

Carlson S. M., Cunningham C. J., Westley P. A. H., 2014 Evolutionary rescue in a changing world. Trends Ecol. Evol. 29: 521–530.

Charmantier A., Garant D., 2005 Environmental quality and evolutionary potential: lessons from wild populations. Proc. R. Soc. B Biol. Sci. 272: 1415–1425.

Chevin L.-M., Lande R., 2010 When Do Adaptive Plasticity and Genetic Evolution Prevent Extinction of a Density-Regulated Population? Evolution 64: 1143–1150.

Chevin L.-M., Martin G., Lenormand T., 2010 Fisher's Model and the Genomics of Adaptation: Restricted Pleiotropy, Heterogenous Mutation, and Parallel Evolution. Evolution 64: 3213–3231.

Chevin L.-M., 2011 On measuring selection in experimental evolution. Biol. Lett. 7: 210–213.

Davies J., Davies D., 2010 Origins and Evolution of Antibiotic Resistance. Microbiol. Mol. Biol. Rev. 74: 417–433.

Day T., Read A. F., 2016 Does High-Dose Antimicrobial Chemotherapy Prevent the Evolution of Resistance? PLoS Comput. Biol. 12: e1004689.

De Visser J. a. G. M., Rozen D. E., 2005 Limits to adaptation in asexual populations. J. Evol. Biol. 18: 779–788.

Drlica K., 2003 The mutant selection window and antimicrobial resistance. J. Antimicrob. Chemother. 52: 11–17.

Feller W., 1951 Diffusion processes in genetics. In: *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*,

Foster P. L., 2007 Stress-Induced Mutagenesis in Bacteria. Crit. Rev. Biochem. Mol. Biol. 42: 373–397.

Gillespie J. H., 1983 Some Properties of Finite Populations Experiencing Strong Selection and Weak Mutation. Am. Nat. 121: 691–708.

Gomulkiewicz R., Holt R. D., 1995 When does Evolution by Natural Selection Prevent Extinction? Evolution 49: 201.

Gomulkiewicz R., Holt R. D., Barfield M., Nuismer S. L., 2010 Genetics, adaptation, and invasion in harsh environments. Evol. Appl. 3: 97–108.

Gonzalez A., Bell G., 2013 Evolutionary rescue and adaptation to abrupt environmental change depends upon the history of stress. Philos. Trans. R. Soc. Lond. B Biol. Sci. 368: 20120079.

Gonzalez A., Ronce O., Ferriere R., Hochberg M. E., 2013 Evolutionary rescue: an emerging focus at the intersection between ecology and evolution. Philos. Trans. R. Soc. B Biol. Sci. 368: 20120404–20120404.

Harmand N., Gallet R., Jabbour-Zahab R., Martin G., Lenormand T., 2017 Fisher's geometrical model and the mutational patterns of antibiotic resistance across dose gradients. Evolution 71: 23–37.

Harrigan P. R., Hogg R. S., Dong W. W. Y., Yip B., Wynhoven B., *et al.*, 2005 Predictors of HIV drug-resistance mutations in a large antiretroviral-naive cohort initiating triple antiretroviral therapy. J. Infect. Dis. 191: 339–347.

Hermisson J., Wagner G. P., 2004 The Population Genetic Theory of Hidden Variation and Genetic Robustness. Genetics 168: 2271–2284.

Hietpas R. T., Bank C., Jensen J. D., Bolon D. N. A., 2013 Shifting Fitness Landscapes in Response to Altered Environments. Evolution 67: 3512–3522.

Hoffmann A. A., Parsons P. A., 1997 *Extreme environmental change and evolution*. Cambridge University Press, Cambridge; New York.

Hoffmann A. A., Merilä J., 1999 Heritable variation and evolution under favourable and unfavourable conditions. Trends Ecol. Evol. 14: 96–101.

Koehn R. K., Bayne B. L., 1989 Towards a physiological and genetical understanding of the energetics of the stress response. Biol. J. Linn. Soc. 37: 157–171.

Kopp M., Matuszewski S., 2014 Rapid evolution of quantitative traits: theoretical perspectives. Evol. Appl. 7: 169–191.

Lambert A., 2005 The branching process with logistic growth. Ann. Appl. Probab. 15: 1506–1535.

Lindsey H. A., Gallie J., Taylor S., Kerr B., 2013 Evolutionary rescue from extinction is contingent on a lower rate of environmental change. Nature 494: 463–467.

Lynch M., Gabriel W., Wood A. M., 1991 Adaptive and demographic responses of plankton populations to environmental change. Limnol. Oceanogr. 36: 1301–1312.

MacLean R. C., Perron G. G., Gardner A., 2010 Diminishing Returns From Beneficial Mutations and Pervasive Epistasis Shape the Fitness Landscape for Rifampicin Resistance in Pseudomonas aeruginosa. Genetics 186: 1345–1354.

Martin G., Lenormand T., 2006a A General Multivariate Extension of Fisher's Geometrical Model and the Distribution of Mutation Fitness Effects Across Species. Evolution 60: 893–907.

Martin G., Lenormand T., 2006b The Fitness Effect of Mutations Across Environments: A Survey in Light of Fitness Landscape Models. Evolution 60: 2413–2427.

Martin G., Elena S. F., Lenormand T., 2007 Distributions of epistasis in microbes fit predictions from a fitness landscape model. Nat. Genet. 39: 555–560.

Martin G., Aguilée R., Ramsayer J., Kaltz O., Ronce O., 2013 The probability of evolutionary rescue: towards a quantitative comparison between theory and evolution experiments. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 368: 20120088.

Martin G., 2014 Fisher's geometrical model emerges as a property of complex integrated phenotypic networks. Genetics 197: 237–255.

Martin G., Lenormand T., 2015 The fitness effect of mutations across environments: Fisher's geometrical model with multiple optima. Evolution 69: 1433–1447.

Martin G., Roques L., 2016 The Non-stationary Dynamics of Fitness Distributions: Asexual Model with Epistasis and Standing Variation. Genetics: genetics.116.187385.

Moser C., Bell G., 2011 Genetic correlation in relation to differences in dosage of a stressor. J. Evol. Biol. 24: 219–223.

Orr H. A., Unckless R. L., 2008 Population extinction and the genetics of adaptation. Am. Nat. 172: 160–9.

Orr H. A., Unckless R. L., 2014 The Population Genetics of Evolutionary Rescue. PLOS Genet. 10: e1004551.

Perfeito L., Sousa A., Bataillon T., Gordo I., 2014 Rates of Fitness Decline and Rebound Suggest Pervasive Epistasis. Evolution 68: 150–162.

Read A. F., Day T., Huijben S., 2011 The evolution of drug resistance and the curious orthodoxy of aggressive chemotherapy. Proc. Natl. Acad. Sci. 108: 10871–10877.

Regoes R. R., Wiuff C., Zappala R. M., Garner K. N., Baquero F., *et al.*, 2004 Pharmacodynamic Functions: a Multiparameter Approach to the Design of Antibiotic Treatment Regimens. Antimicrob. Agents Chemother. 48: 3670–3676.

Remold S. K., Lenski R. E., 2001 Contribution of individual random mutations to genotype-by-environment interactions in Escherichia coli. Proc. Natl. Acad. Sci. U. S. A. 98: 11388–11393.

Remold S. K., Lenski R. E., 2004 Pervasive joint influence of epistasis and plasticity on mutational effects in Escherichia coli. Nat. Genet. 36: 423–426.

Samani P., Bell G., 2010 Adaptation of experimental yeast populations to stressful conditions in relation to population size. J. Evol. Biol. 23: 791–796.

Scharloo W., 1991 Canalization: genetic and developmental aspects. Annu. Rev. Ecol. Syst. 22: 65–93.

Sgrò C. M., Hoffmann A. A., 2004 Genetic correlations, tradeoffs and environmental variation. Heredity 93: 241–248.

Tenaillon O., 2014 The Utility of Fisher's Geometric Model in Evolutionary Genetics. Annu. Rev. Ecol. Evol. Syst. 45: 179–201.

Trindade S., Sousa A., Gordo I., 2012 Antibiotic Resistance and Stress in the Light of Fisher's Model. Evolution 66: 3815–3824.

Turelli M., 1984 Heritable genetic variation via mutation-selection balance: Lerch's zeta meets the abdominal bristle. Theor. Popul. Biol. 25: 138–193.

Uecker H., Otto S. P., Hermisson J., 2014 Evolutionary Rescue in Structured Populations. Am. Nat. 183: E17–E35.

Uecker H., Hermisson J., 2016 The Role of Recombination in Evolutionary Rescue. Genetics 202: 721–732.

Wang A. D., Sharp N. P., Spencer C. C., Tedman-Aucoin K., Agrawal A. F., 2009 Selection, Epistasis, and Parent-of-Origin Effects on Deleterious Mutations across Environments in *Drosophila melanogaster*. Am. Nat. 174: 863–874.

Wang A. D., Sharp N. P., Agrawal A. F., 2014 Sensitivity of the Distribution of Mutational Fitness Effects to Environment, Genetic Background, and Adaptedness: A Case Study with Drosophila. Evolution 68: 840–853.

Wilson B. A., Pennings P. S., Petrov D. A., 2017 Soft Selective Sweeps in Evolutionary Rescue. Genetics 205: 1573–1586.

Wolfram Research I., 2012 *Mathematica*. Wolfram Research, Inc, Champaign, Illinois.

Wong A., 2017 Epistasis and the Evolution of Antimicrobial Resistance. Front. Microbiol. 8.