Full title:

Prominent features of the amino acid mutation landscape in cancer

Short title:

R>H and E>K mutations are prominent features in many cancers

Zachary A. Szpiech[1,*], Nicolas B. Strauli[1,2], Katharine A. White[3], Diego Garrido Ruiz[4],

Matthew P. Jacobson[1,4], Diane L. Barber[3], Ryan D. Hernandez[1,5,6,*]

[1]Department of Bioengineering and Therapeutic Sciences, University of California, San
Francisco.

[2]Biomedical Sciences Graduate Program, University of California, San Francisco.

[3]Department of Cell and Tissue Biology, University of California, San Francisco.

[4]Department of Pharmaceutical Chemistry, University of California, San Francisco.

[5]Quantitative Biosciences Institute, University of California, San Francisco.

[6]Institute for Human Genetics, University of California, San Francisco.
16
17
18
19
20   *Corresponding authors
21
Ryan D Hernandez, PhD
UCSF MC 2530
Byers Hall Room 308
1700 4th Street
San Francisco, CA 94143
Email: ryan.hernandez@ucsf.edu
28
Zachary A Szpiech, PhD
UCSF MC 2530
Byers Hall Room 308
1700 4th Street
San Francisco, CA 94143
Email: zachary.szpiech@ucsf.edu
35

36
ABSTRACT

Cancer can be viewed as a set of different diseases with distinctions based on

tissue origin, driver mutations, and genetic signatures. Accordingly, each of these

distinctions have been used to classify cancer subtypes and to reveal common features.

Here, we present a different analysis of cancer based on amino acid mutation signatures.

Non-negative Matrix Factorization and principal component analysis of 29 cancers

revealed six amino acid mutation signatures, including four signatures that were

dominated by either arginine to histidine (Arg>His) or glutamate to lysine (Glu>Lys)

mutations. Sample-level analyses reveal that while some cancers are heterogeneous,

others are largely dominated by one type of mutation. Using a non-overlapping set of

samples from the COSMIC somatic mutation database, we validate five of six mutation

signatures, including signatures with prominent arginine to histidine (Arg>His) or

glutamate to lysine (Glu>Lys) mutations. This suggests that our classification of cancers

based on amino acid mutation patterns may provide avenues of inquiry pertaining to

specific protein mutations that may generate novel insights into cancer biology.

52
INTRODUCTION

Cancers have been described as open, complex, and adaptive systems [1].

Reflecting this, cancer progression is determined in part by genetic diversification and

clonal selection within complex tissue landscapes and with changing tumor properties

and microenvironment features [2, 3]. Genetic sequencing of tumor samples has been

critical in developing the evolutionary theory of cancer. While cancers traditionally have

been—and continue to be—classified by tissue of origin, genetic sequencing has allowed

for classification based on driver mutations [4] or nucleotide mutation signatures [5].

However, cancer cell adaptation is mediated by changes at the protein level that alter cell

biology and enable cancer cell behaviors such as increased proliferation and cell survival.

Existing cancer classifications by nucleotide mutation signatures lack a link between the

underlying genetic landscape and effects on cancer cell phenotypes. Analysis of cancers

by amino acid mutations could provide important connections between cancer evolution

and adaptive biological phenotypes as well as provide insight into how specific classes of

amino acid mutations may generally alter the function of the proteins in which they are

found. There have been some studies to examine amino acid mutations across cancers [6-

8], but these have relied on simple mutation counting methods.

Here we take a machine-learning approach to analyze amino acid mutations

across 29 cancers in order to identify characteristic amino acid mutation signatures. Our

analyses reveal that some cancer types have mutation signatures dominated by arginine to

histidine (Arg>His) mutations, some have signatures dominated by glutamate to lysine

(Glu>Lys), and others have more complex signatures that lack a single dominant amino

acid mutation. These signatures were further validated in a non-overlapping set of

samples from the COSMIC somatic mutation database. Importantly, this approach

identifies not only which amino acid mutations are prevalent among cancers but also

which amino acid mutations tend to occur together. For example, cancers with strong

Arg>His signatures will also frequently have many Ala>Thr mutations but are unlikely to

have many Glu>Lys mutations (despite all of these amino acid transitions resulting from

a G>A nucleotide mutation).

RESULTS

83   **Several cancers are enriched for R>H and E>K amino acid mutations**

84   Multiple studies have interrogated nucleotide mutation biases by analyzing

85   somatic variation across a wide range of cancers [4, 5]. However, in protein coding

86   regions of the genome (i.e. the exome), it is essential to study patterns of amino acid

87   variation to reveal information about potential functional effects at the protein level. We

88   characterized the global properties of amino acid mutations encoded by somatic

89   mutations across a range of cancers by analyzing a tumor-normal paired mutation

90   database [5] consisting of 6,931 samples across 29 cancer types. We applied filtering to

91   remove sequencing artifacts and restricted mutation data to nonsynonymous amino acid

92   mutations (see Materials and Methods, S1 Table and S2 Table for details).

93   Using this amino acid mutation database, we performed an unbiased

94   characterization of mutation signatures across cancer types using Non-negative Matrix

95   Factorization (NMF), which has proven to be a useful tool for pattern discovery in cancer

96   tissue mutation datasets [5] and other biological systems [9]. Applying NMF to the

97   pooled mutation data reveals six mutation signatures at the amino acid level (S1G Fig),

98   including two with strong Arg>His components and two with strong Glu>Lys

99   components (Fig 1A, S1 Fig). Although the cancers are comprised of a mixture of the

100   signatures identified, ten cancers (AML, colorectal, esophageal, low grade glioma,

101   kidney chromophobe, medulloblastoma, pancreatic, prostate, stomach, and uterine) have

102   majority contributions from Arg>His-prominent mutation signatures (R>H and

103   A>T/R>H). We also identify four cancers (bladder, cervix, head and neck, and

104   melanoma) that have majority contributions from Glu>Lys-prominent mutation

105   signatures (E>K and E>K/E>Q). Additionally, there are two complex signatures not

dominated by any particular amino acid mutation. Glioblastoma, kidney papillary, liver,

and thyroid cancers have majority contribution from the Complex 1 signature, and lung

adenocarcinoma, small cell lung, squamous cell lung, and neuroblastoma cancers all have

majority contribution from the Complex 2 signature. Finally, seven cancers from a

variety of tissues (ALL, breast, CLL, clear cell kidney, B-cell lymphoma, myeloma, and

ovarian) have heterogeneous mutation signature contributions.

Fig 1. Arg>His and Glu>Lys mutations define mutation signatures of a subset of

cancers.

(A) Heatmap representation of six-component NMF clustering. Of the six amino acid

mutation signatures identified, four have prominent charge-changing mutations: Arg>His

(R>H), Glu>Lys (E>K), or Glu>Gln (E>Q). Two complex signatures were also

identified. Color scale represents scaled contribution of each signature for a given cancer

type. Signature and NMF fit details can be found in S1 Fig. (B) Principal component

analysis of nonsynonymous amino acid mutations. PC1 separates cancers with high R>H

from cancers with high E>K; PC2 separates cancers with complex signatures. Colors

represent the greatest mutation signature contributing to a given cancer. Individual PC

loadings can be found in S2 Fig.

Visualizing Amino Acid Mutation Properties with Principal Component Analysis

To alternatively visualize the amino acid mutation spectrum, we use principal

component analysis to reveal cancers clustering by dominant mutation classes (Fig 1B).

We find that PC1 separates Arg>His dominant cancers from Glu>Lys dominant cancers

and that PC2 separates cancers with more complex signatures (S2 Fig). This result

reinforces our observation that Arg>His and Glu>Lys mutations are characteristic

signatures of several cancers.

Individual Cancer Samples Recapitulate Amino Acid Mutation Patterns

We also analyze samples individually with NMF and find that Arg>His and

Glu>Lys features continue to dominate (Fig 2A and S3 Fig). For many cancer subtypes

(melanoma, bladder, uterine, colorectal, low-grade glioma, cervix, neuroblastoma, and

the three different lung cancers), individual patients within each

cancer exhibit consistent amino acid signatures (Fig 2B). This is true even within

clinically diverse cancers such as bladder, uterine, colorectal, and lung cancer, which all

have multiple identified driver mutations. This suggests that the amino acid signatures we

identified may be independent of underlying driver mutations and may instead be a

consequence of common features of the cancer, tumor microenvironment, or selective

pressures, all of which may be targeted therapeutically.

Fig 2. Amino acid mutation signatures for individual samples.

(A) A heatmap representation of the six-component NMF clustering results for individual

cancer samples (only those with >10 total nonsynonymous mutations). Samples with the

same maximum signature component were grouped and sorted. Four amino acid

mutation signatures identified (R>H, E>K, E>K/E>Q, Complex 2) overlap with

signatures in Fig 1A. Color scale represents scaled contribution of each signature for a

given sample. Signature and NMF fit details can be found in S3 Fig. (B) Bars show the

total fraction of individual samples with a majority of a particular signature within

each cancer. Within cancers, a large fraction of individual samples tend to have similar

signature components.

As NMF decomposes a sample into a mixture of characteristic signatures, we can

further visualize the normalized mixture coefficients from the individual-level NMF

along the three mutation signatures with dominant Arg>His or Glu>Lys components

(R>H, E>K, and E>K/E>Q signatures; Fig 3) to determine whether samples tend to be an

equal mixture of several signatures or whether they tend to be exclusively composed of a

single signature. Indeed, Fig 3 shows a clear separation of samples with a high proportion

of Glu>Lys from other signatures.

Fig 3. Normalized NMF mixture coefficients for individual samples.

Plot of the normalized mixture coefficients across the three mutation signatures with high

R>H or E>K components for every individual sample. Colors represent the greatest

contributing mutation signature for each sample based on the full individual-level NMF

analysis. Here we see a dramatic separation of samples in the E>K component to the near

exclusion of other signatures.

Mutation Signature Validation

We validated the NMF signatures with an orthogonal data set (see Materials and

Methods) from the COSMIC database [10]. The six mutation signatures identified from

COSMIC (S4 Fig) overlap substantially with previously identified signatures (S3 Fig).

We calculated correlation coefficients between all COSMIC Data signatures and each

Alexandrov Data signature. When the correlations are very high, this indicates that NMF

has identified the same general mutation signature in the two different data sets. Indeed,

we found high correlation between the COSMIC signatures and our initially identified

signatures for five of the six (Fig 4): R>H, E>K, E>K/E>Q, Complex 2, and Complex A

are replicated. The Complex B signature does not replicate as a separate signature, but

appears to be largely incorporated into the other complex signatures. Interestingly, a new

R>Q/R>W signature is identified as a separate component in the COSMIC data. On

inspection of the Alexandrov R>H component we identified, we see that R>Q and R>W

are prominent components. Increased sample size in our replicate data set likely enabled

NMF to discriminate between these two signatures in COSMIC.

186
Fig 4. Correlations of COSMIC and Alexandrov mutation signatures. For each

COSMIC mutation signature we calculated the correlation with each Alexandrov

mutation signature. Five of six Alexandrov mutation signatures replicate in the COSMIC

data for k = 6 mutation signatures. Alexandrov Signature R>H is replicated by COSMIC

Signature 1, although a subset of mutation types that clustered in the original R>H are

identified in the larger COSMIC data set as Signature 2. Alexandrov Signature E>K is

replicated by COSMIC Signature 4. Alexandrov Signature E>K, E>Q is replicated by

COSMIC Signature 5, although it is also correlated with COSMIC Signature 4 as each of

these signatures share mutations. Alexandrov Signature Complex 2 is replicated by

COSMIC Signature 6. Alexandrov Signature Complex A is replicated by COSMIC

Signature 3. Alexandrov Signature Complex B is not faithfully replicated by any single

COSMIC Signature, as the signal appears to be spread amongst COSMIC Signatures 3, 5

and 6.

DISCUSSION

Proteomic changes can allow cancer cells to adapt to dynamic pressures including

changes in matrix composition, oxygen and nutrient availability, intracellular

metabolism, as well as increased intracellular pH (pHi), the latter enabling tumorigenic

cell behaviors [11-15]. Our analyses reveal that a subset of all possible amino acid

mutations dominate the mutation landscape of cancers, with Glu>Lys and Arg>His

mutations being the most prominent features of identified mutation signatures.

Charge-changing mutations, whether buried or surface-exposed, can alter protein

charge, electrostatics, and conformation [16]. Electrostatics of surface residues have been

shown to play a key role in protein-protein interactions [17], protein-membrane

interactions [18, 19], and kinase substrate recognition [20]. While it is important to note

that our analyses are agnostic to the location of the mutation within the proteome and

within a protein, the strong bias towards amino acid mutations that alter charge in our

identified mutation signatures may suggest an adaptive advantage conferred by these

mutations.

Glu>Lys mutations swap a negatively charged amino acid for a positively charged

amino acid, which may in some cases affect protein function. Indeed, in some cases

buried lysine mutations can induce global protein unfolding upon charging that alters

mutant protein stability and function [21]. Furthermore, Glu>Lys mutations have been

known to affect the function of PIK3CA [22-24].

Arg>His mutations swap a positively charged amino acid for a titratable amino

acid. Whereas arginine (pKa ~12) should always be protonated, histidine (pKa ~6.5) can

titrate within the narrow physiological pH range. Indeed, the pH-sensitive function of

many wild-type proteins has been shown to be mediated by titratable histidine residues

[25-27]. Moreover, recent work has shown that some Arg>His mutations can confer pH

sensitivity to the mutant protein and alter function [28]. We predict that some Arg>His

mutations may be adaptive to increased pHi, conferring a gain in pH sensing to the

mutant protein.

From our analyses, Arg>His mutations define the mutation landscape of a diverse

set of cancers across a range of tissues including brain (low-grade glioma), digestive

(colorectal), reproductive (uterine), and blood (AML) cancers. Importantly, these cancers

do not have overlapping nucleotide mutation signatures [5], which suggests that the

amino acid mutation signatures we identified may reflect other aspects of the cancers

including distinct physiological pressures, microenvironment features, or functional

requirements. Indeed, these results may help inform studies in the emerging field of

Molecular Pathologic Epidemiology (MPE) [29, 30], which seeks to integrate knowledge

across disciplines to inform personalized approaches to cancer prevention and therapy.

Linking amino acid signatures to physiological or pathological features of the cancer

could be important for identifying selective pressures that may be driving or sustaining

the cancer as well as for limiting disease progression, particularly where targeted

approaches fail [31-33].

243
244   **MATERIALS AND METHODS**

Mutation Dataset Filtering

We validated the dataset [5] by comparing known frequencies of well-studied

cancer driver genes with observed frequencies in the dataset. Specifically, BRAF is

mutated in 40–50% of melanoma samples, and IDH1 is mutated in 75–85%, low-grade

glioma, AML, and glioblastoma samples are mutated 75-85%, 8–12%, and 1–5% of the

time, respectively. We used the p53 database (http://p53.fr/index.html) to find expected

p53 mutation frequency for various cancers: colorectal, head and neck, pancreatic,

stomach, liver, and breast cancer have 43%, 42%, 34%, 32%, 31%, and 22% p53

mutation rates, respectively. The observed mutation frequencies were consistently lower

than expected for the genes/cancers we assessed, which suggests that the dataset authors

[5] were perhaps too stringent in quality control (QC) filtering. Different levels of QC

filtering were performed, and we systematically relaxed filters in order to recapitulate the

expected mutation frequencies of the selected canonical driver genes. Applying only the

'sequencing artifact' QC filter (from [5]) most closely recapitulated expected mutation

frequencies for the canonical driver genes, and this filter alone was used for the

remainder of the bioinformatics analyses.

Mapping somatic SNPs

After filtering we used part of the PolyPhen2 [34] pipeline to map mutations to

UCSC Canonical transcripts and restricted to nonsynonymous amino acid changes. The

following cancers had reduced sample sizes after filtering and nonsynonymous mutation

restriction: AML: one sample eliminated through QC filtering, two samples eliminated

because all mutations were synonymous; low grade glioma: one sample eliminated

because after QC filtering all remaining mutations were synonymous; glioblastoma: two

samples eliminated because all mutations were synonymous. All Pilocytic Astrocytoma

samples were excluded from future analysis due to low total nonsynonymous mutations

per sample.

Mutation frequency data sets

For the individual sample data, we represent each sample as a row vector with

elements giving the mutation counts observed for each nonsynonymous mutation (e.g.

Ala>Cys, Ala>Asp, etc.) and removing all samples with <10 total observed mutations.

For the aggregated data set, we sum the mutation counts across all samples of the same

cancer type (including samples with <10 mutations), giving one row vector for each

cancer type where each element represents the total number of observed nonsynonymous

mutations across all samples. For non-negative matrix factorization and principal

component analysis, we divide each row by the row sum.

NMF is an unsupervised learning method used to decompose a data matrix into a

product of two non-negative matrices representing a set of k signals and mixture

coefficients. For example if $\mathbf{X}$ is an $m \times n$ matrix representing the nonsynonymous

mutation frequency data, then the NMF of the data is given by

$$X = WH$$

287   where $\mathbf{W}$ is an $m \times k$ matrix with the k columns representing mutation signatures and $\mathbf{H}$

288   is a $k \times n$ matrix representing the mixture coefficients that best reconstruct $\mathbf{X}$. Often it is

289   not possible to factor $\mathbf{X}$ exactly, so a typical approach to solving the decomposition will

290   optimize

$$\min_{w, H \geq 0} [D(X, WH) + R(W, H)]$$

291   where D() is a loss function (often the Frobenius norm or the Kullback-Leibler

292   divergence) and R() is a regularization function. For our NMF analyses, we utilize the R

293   package *NMF* [35] with default choices for D() and R().

Principal Component Analysis (PCA)

296   PCA is a dimension reducing learning method designed to decompose a data

297   matrix into a set of orthogonal bases defined along the major axes of variation within the

298   data. Here we compute the first two principal components from our mutation frequency

299   matrix **X**. The kth principal component is represented by a vector of loadings, $w_{(k)}$. The

300   first PC is then calculated as

$$w_{(1)} = \arg\max \left\{ \frac{w^T X^T X w}{w^T w} \right\}$$

301   and subsequent PCs are calculated as

$$w_{(k)} = \arg\max \left\{ \frac{w^T \widehat{X}_k^T \widehat{X}_k w}{w^T w} \right\}$$

302   where

$$\widehat{X}_k = X - \sum_{s}^{k-1} X w_{(s)} w_{(s)}^T.$$

303   We use the R package *prcomp* to perform all PCA analyses.

Validation of NMF Mutation Signatures

306   In order to validate the mutation signatures that we discovered in our data, we

sought an orthogonal data set in which to replicate our analysis. We used the COSMIC

v81 database of somatic mutations [10]. We first filtered all mutations that were not

marked as confirmed somatic mutations. Next, as our original data set ("Alexandrov

Data") had overlapping samples within the COSMIC database, we excluded all samples

that were included in our original analysis. Finally, we excluded samples with fewer than

10 total non-synonymous mutations. This filtering resulted in a final data set of 2,236,176

non-synonymous mutations across 15,868 samples. We named this final data set the

"COSMIC Data." We then ran NMF with $k = 6$ signatures on the matrix of individual

sample mutation frequencies as described above. Results are shown in S4 Fig. We found

that five of the six mutation signatures we originally discovered were replicated in the

COSMIC data (Fig 4).

ACKNOWLEDGEMENTS

321   This work was supported by National Institutes of Health grants CA178706 to

322 Diane L. Barber and Ryan D. Hernandez; CA197855 to Diane L. Barber, and HG007644

323 to Ryan D. Hernandez; and National Institutes of Health F32 grant CA177085 to

324 Katharine A. White.

Bibliography

1.   Gillies RJ, Gatenby RA. Metabolism and its sequelae in cancer evolution and
therapy. Cancer J. 2015;21(2):88-96. doi: 10.1097/PPO.0000000000000102. PubMed
PMID: 25815848; PubMed Central PMCID: PMCPMC4446699.
2.   Greaves M, Maley CC. Clonal evolution in cancer. Nature. 2012;481(7381):306-
13. doi: 10.1038/nature10762. PubMed PMID: 22258609; PubMed Central PMCID:
PMCPMC3367003.

3.   Nowell PC. The clonal evolution of tumor cell populations. Science.
1976;194(4260):23-8. PubMed PMID: 959840.
4.   Bignell GR, Greenman CD, Davies H, Butler AP, Edkins S, Andrews JM, et al.
Signatures of mutation and selection in the cancer genome. Nature. 2010;463(7283):893-
8. doi: 10.1038/nature08768. PubMed PMID: 20164919; PubMed Central PMCID:
PMCPMC3145113.
5.   Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV,
et al. Signatures of mutational processes in human cancer. Nature. 2013;500(7463):415-
21. doi: 10.1038/nature12477. PubMed PMID: 23945592; PubMed Central PMCID:
PMCPMC3776390.
6.   Tan H, Bao J, Zhou X. Genome-wide mutational spectra analysis reveals
significant cancer-specific heterogeneity. Sci Rep. 2015;5:12566. doi:
10.1038/srep12566. PubMed PMID: 26212640; PubMed Central PMCID:
PMCPMC4515826.
7.   Anoosha P, Sakthivel R, Michael Gromiha M. Exploring preferred amino acid
mutations in cancer genes: Applications to identify potential drug targets. Biochim
Biophys Acta. 2016;1862(2):155-65. doi: 10.1016/j.bbadis.2015.11.006. PubMed PMID:
26581171.
8.   Tsuber V, Kadamov Y, Brautigam L, Berglund UW, Helleday T. Mutations in
Cancer Cause Gain of Cysteine, Histidine, and Tryptophan at the Expense of a Net Loss
of Arginine on the Proteome Level. Biomolecules. 2017;7(3). doi:
10.3390/biom7030049. PubMed PMID: 28671612.
9.   Brunet JP, Tamayo P, Golub TR, Mesirov JP. Metagenes and molecular pattern
discovery using matrix factorization. Proc Natl Acad Sci U S A. 2004;101(12):4164-9.
doi: 10.1073/pnas.0308531101. PubMed PMID: 15016911; PubMed Central PMCID:
PMCPMC384712.
10.   Forbes SA, Beare D, Boutselakis H, Bamford S, Bindal N, Tate J, et al. COSMIC:
somatic cancer genetics at high-resolution. Nucleic Acids Res. 2017;45(D1):D777-D83.
doi: 10.1093/nar/gkw1121. PubMed PMID: 27899578; PubMed Central PMCID:
PMCPMC5210583.
11.   White KA, Grillo-Hill BK, Barber DL. Cancer cell behaviors mediated by
dysregulated pH dynamics at a glance. J Cell Sci. 2017;130(4):663-9. doi:
10.1242/jcs.195297. PubMed PMID: 28202602; PubMed Central PMCID:
PMCPMC5339414.
12.   Cardone RA, Casavola V, Reshkin SJ. The role of disturbed pH dynamics and the
Na+/H+ exchanger in metastasis. Nat Rev Cancer. 2005;5(10):786-95. doi:
10.1038/nrc1713. PubMed PMID: 16175178.
13.   Grillo-Hill BK, Choi C, Jimenez-Vidal M, Barber DL. Increased H(+) efflux is
sufficient to induce dysplasia and necessary for viability with oncogene expression. Elife.
2015;4. doi: 10.7554/eLife.03270. PubMed PMID: 25793441; PubMed Central PMCID:
PMCPMC4392478.
14.   Parks SK, Chiche J, Pouyssegur J. Disrupting proton dynamics and energy
metabolism for cancer therapy. Nat Rev Cancer. 2013;13(9):611-23. doi:
10.1038/nrc3579. PubMed PMID: 23969692.

15.   Webb BA, Chimenti M, Jacobson MP, Barber DL. Dysregulated pH: a perfect
storm for cancer progression. Nat Rev Cancer. 2011;11(9):671-7. doi: 10.1038/nrc3110.
PubMed PMID: 21833026.
16.   Zheng Y, Cui Q. Microscopic mechanisms that govern the titration response and
pKa values of buried residues in staphylococcal nuclease mutants. Proteins.
2017;85(2):268-81. doi: 10.1002/prot.25213. PubMed PMID: 27862310.
17.   Zhang Z, Witham S, Alexov E. On the role of electrostatics in protein-protein
interactions. Phys Biol. 2011;8(3):035001. doi: 10.1088/1478-3975/8/3/035001. PubMed
PMID: 21572182; PubMed Central PMCID: PMCPMC3137121.
18.   Cho W, Stahelin RV. Membrane-protein interactions in cell signaling and
membrane trafficking. Annu Rev Biophys Biomol Struct. 2005;34:119-51. doi:
10.1146/annurev.biophys.33.110502.133337. PubMed PMID: 15869386.
19.   Stahelin RV. Lipid binding domains: more than simple lipid effectors. J Lipid
Res. 2009;50 Suppl:S299-304. doi: 10.1194/jlr.R800078-JLR200. PubMed PMID:
19008549; PubMed Central PMCID: PMCPMC2674730.
20.   Ubersax JA, Ferrell JE, Jr. Mechanisms of specificity in protein phosphorylation.
Nat Rev Mol Cell Biol. 2007;8(7):530-41. doi: 10.1038/nrm2203. PubMed PMID:
17585314.
21.   Chimenti MS, Khangulov VS, Robinson AC, Heroux A, Majumdar A,
Schlessman JL, et al. Structural reorganization triggered by charging of Lys residues in
the hydrophobic interior of a protein. Structure. 2012;20(6):1071-85. doi:
10.1016/j.str.2012.03.023. PubMed PMID: 22632835; PubMed Central PMCID:
PMCPMC3373022.
22.   Carson JD, Van Aller G, Lehr R, Sinnamon RH, Kirkpatrick RB, Auger KR, et al.
Effects of oncogenic p110alpha subunit mutations on the lipid kinase activity of
phosphoinositide 3-kinase. Biochem J. 2008;409(2):519-24. doi: 10.1042/BJ20070681.
PubMed PMID: 17877460.
23.   Huang CH, Mandelker D, Gabelli SB, Amzel LM. Insights into the oncogenic
effects of PIK3CA mutations from the structure of p110alpha/p85alpha. Cell Cycle.
2008;7(9):1151-6. doi: 10.4161/cc.7.9.5817. PubMed PMID: 18418043; PubMed Central
PMCID: PMCPMC3260475.
24.   Meyer DS, Koren S, Leroy C, Brinkhaus H, Muller U, Klebba I, et al. Expression
of PIK3CA mutant E545K in the mammary gland induces heterogeneous tumors but is
less potent than mutant H1047R. Oncogenesis. 2013;2:e74. doi: 10.1038/oncsis.2013.38.
PubMed PMID: 24080956; PubMed Central PMCID: PMCPMC3816227.
25.   Choi CH, Webb BA, Chimenti MS, Jacobson MP, Barber DL. pH sensing by
FAK-His58 regulates focal adhesion remodeling. J Cell Biol. 2013;202(6):849-59. doi:
10.1083/jcb.201302131. PubMed PMID: 24043700; PubMed Central PMCID:
PMCPMC3776353.
26.   Frantz C, Barreiro G, Dominguez L, Chen X, Eddy R, Condeelis J, et al. Cofilin is
a pH sensor for actin free barbed end formation: role of phosphoinositide binding. J Cell
Biol. 2008;183(5):865-79. doi: 10.1083/jcb.200804161. PubMed PMID: 19029335;
PubMed Central PMCID: PMCPMC2592832.
27.   Webb BA, White KA, Grillo-Hill BK, Schonichen A, Choi C, Barber DL. A
Histidine Cluster in the Cytoplasmic Domain of the Na-H Exchanger NHE1 Confers pH-
sensitive Phospholipid Binding and Regulates Transporter Activity. J Biol Chem.

2016;291(46):24096-104. doi: 10.1074/jbc.M116.736215. PubMed PMID: 27650500;
PubMed Central PMCID: PMCPMC5104935.
28.   DiGiammarino EL, Lee AS, Cadwell C, Zhang W, Bothner B, Ribeiro RC, et al.
A novel mechanism of tumorigenesis involving pH-dependent destabilization of a mutant
p53 tetramer. Nat Struct Biol. 2002;9(1):12-6. doi: 10.1038/nsb730. PubMed PMID:
11753428.
29.   Ogino S, Chan AT, Fuchs CS, Giovannucci E. Molecular pathological
epidemiology of colorectal neoplasia: an emerging transdisciplinary and interdisciplinary
field. Gut. 2011;60(3):397-411. doi: 10.1136/gut.2010.217182. PubMed PMID:
21036793; PubMed Central PMCID: PMCPMC3040598.
30.   Nishi A, Milner DA, Jr., Giovannucci EL, Nishihara R, Tan AS, Kawachi I, et al.
Integration of molecular pathology, epidemiology and social science for global precision
medicine. Expert Rev Mol Diagn. 2016;16(1):11-23. doi:
10.1586/14737159.2016.1115346. PubMed PMID: 26636627; PubMed Central PMCID:
PMCPMC4713314.
31.   Alfarouk KO, Stock CM, Taylor S, Walsh M, Muddathir AK, Verduzco D, et al.
Resistance to cancer chemotherapy: failure in drug response from ADME to P-gp. Cancer
Cell Int. 2015;15:71. doi: 10.1186/s12935-015-0221-1. PubMed PMID: 26180516;
PubMed Central PMCID: PMCPMC4502609.
32.   Alfarouk KO, Verduzco D, Rauch C, Muddathir AK, Adil HH, Elhassan GO, et
al. Glycolysis, tumor metabolism, cancer growth and dissemination. A new pH-based
etiopathogenic perspective and therapeutic approach to an old cancer question.
Oncoscience. 2014;1(12):777-802. doi: 10.18632/oncoscience.109. PubMed PMID:
25621294; PubMed Central PMCID: PMCPMC4303887.
33.   Gillies RJ, Verduzco D, Gatenby RA. Evolutionary dynamics of carcinogenesis
and why targeted therapy does not work. Nat Rev Cancer. 2012;12(7):487-93. doi:
10.1038/nrc3298. PubMed PMID: 22695393; PubMed Central PMCID:
PMCPMC4122506.
34.   Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al.
A method and server for predicting damaging missense mutations. Nat Methods.
2010;7(4):248-9. doi: 10.1038/nmeth0410-248. PubMed PMID: 20354512; PubMed
Central PMCID: PMCPMC2855889.
35.   Gaujoux R, Seoighe C. A flexible R package for nonnegative matrix factorization.
BMC Bioinformatics. 2010;11:367. doi: 10.1186/1471-2105-11-367. PubMed PMID:
20598126; PubMed Central PMCID: PMCPMC2912887.
461

A

Membership Fraction

0 0.2 0.6 1.0

Individual Samples

R>H   E>K   E>K   Complex2 ComplexA ComplexB
             E>Q

B

Legend: R>H, E>K, E>K E>Q, Complex 2, Complex A, Complex B

Medulloblastoma
Glioma Low Grade
Uterus
Colorectum
Stomach
Pancreas
AML
Kidney Chromophobe
ALL
Prostate
Esophageal
Melanoma
Cervix
Bladder
Head and Neck
Breast
Myeloma
Lung Adeno
Lung Small Cell
Neuroblastoma
Lung Squamous
Glioblastoma
Liver
Thyroid
Kidney Clear Cell
Kidney Papillary
Ovary
Lymphoma B-Cell
CLL

% of samples with a given signature