1

Selection on morphological traits and fluctuating asymmetry by a fungal

parasite in the yellow dung fly

WOLF U. BLANCKENHORN

Department of Evolutionary Biology and Environmental Studies, University of Zürich,

Winterthurerstrasse 190, CH-8057 Zürich, Switzerland; Fax: +41 44 635.4780; E-mail:

wolf.blanckenhorn@ieu.uzh.ch

3 Figures; 1 Table.

Word count: Abstract 248; Text: 3426; 75 References 1718

Running title Selection by a fungal parasite in dung flies

Abstract (260 words)

Evidence for selective disadvantages of large body size remains scarce in general. Previous phenomenological studies of the yellow dung fly *Scathophaga stercoraria* have demonstrated strong positive sexual and fecundity selection on male and female size. Nevertheless, the body size of flies from a Swiss study population has declined by almost 10% from 1993 to 2009. Given substantial heritability of body size, this negative evolutionary response of an evidently positively selected trait suggests important selective factors being missed (e.g. size-selective predation or parasitism). A periodic epidemic outbreak of the fungus *Entomophthora scatophagae* allowed assessment of selection exerted by this parasite fatal to adult flies. Fungal infection varied over the season from ca. 50% in the cooler and more humid spring and autumn to almost 0% in summer. The probability of dying from fungal infection increased with adult body size. All infected females died before laying eggs, so there was no fungus impact on female fecundity beyond its impact on mortality. Large males showed the typical mating advantage in the field, but this pattern of positive sexual selection was nullified by fungal infection. Mean fluctuating asymmetry of paired appendages (legs, wings) did not affect the viability, fecundity or mating success of yellow dung flies in the field. This study demonstrates rare parasite-mediated disadvantages of large adult body size in the field. Reduced ability to combat parasites such as *Entomophthora* may be an immunity cost of large size in dung flies, although the hypothesized trade-off between fluctuating asymmetry, a presumed indicator of developmental instability and environmental stress, and immunocompetence was not found here.

Keywords: body size, developmental stability, *Entomophthora*, fecundity selection, fluctuating asymmetry, fungal parasite, insect immunity, *Scathophaga stercoraria*, sexual selection, trade-off, viability selection.

Introduction (3629 words)

Systematic quantification of selection has become one of the hallmarks of modern biological research so as to acquire a thorough understanding of the process of natural selection and its evolutionary consequences. Standardized measures of selection (Arnold & Wade, 1984a,b; Lande & Arnold, 1983; Brodie *et al*., 1995) have been available for some time and applied to many species and situations to foster multiple comparative (meta-)analyses, which greatly enhanced our understanding of the action of natural selection in the wild (e.g. Endler, 1986; Kingsolver et al., 2001; Kingsolver & Pfennig, 2004; Cox & Calsbeek, 2009). Correlative phenomenological investigations of selection also are the field method of choice to understand the evolution and population biology of single species or populations in an integrative way, and to test hypotheses about the evolution of particular traits and patterns (e.g. sexual size dimorphism: Blanckenhorn, 2007).

A common pattern is that in many species of animals fecundity selection favors large females and sexual selection large males (Roff, 1992; Andersson, 1994; Kingsolver & Pfennig, 2004). Large body size also confers a number viability benefits (e.g. a longer life), so the selective forces favoring small body size, presumably primarily related to juvenile viability costs of becoming large and/or costs of maintaining a large adult size, are often cryptic (Blanckenhorn, 2000). The widespread yellow dung fly *Scathophaga stercoraria* (Diptera: Scathophagidae) is a classic model species for studies of natural, particularly sexual selection (Parker, 1979; Borgia, 1982; Sigurjónsdóttir & Snorrason, 1995) and a typical case in point. Field studies have established temporally variable but on average very strong mating advantages of large males, as well as fecundity advantages of large females (Jann et al., 2000; Blanckenhorn et al., 2003; cf. Honek, 1993). Strong sexual selection on male body size likely is the main driver of the untypical male-biased sexual size dimorphism in this species (Fairbairn, 1997; Kraushaar & Blanckenhorn, 2002; Blanckenhorn, 2007, 2009).

Nevertheless, the body size of flies from our field population in Switzerland has declined by

almost 10% over a 15-year period from 1993 – 2009 (Blanckenhorn, 2015). Given generally

substantial heritability of body size also in this species (Mousseau & Roff, 1987;

Blanckenhorn, 2000), this negative evolutionary response of a trait that is measurably

strongly positively selected suggests that we are missing important selective factors or

episodes shaping the body size of yellow dung flies (Merilä et al., 2001; Blanckenhorn, 2015;

Gotanda et al., 2015). In general, and also for the yellow dung fly, evidence for selective

disadvantages of large body size remains scarce (Blanckenhorn, 2000, 2007).

One aspect not well studied in yellow dung flies is size-dependent survival in nature. This

is generally the case for small-bodied invertebrates, for which longitudinal field studies are

essentially impossible because individuals cannot be easily marked and followed in nature, as

is the case for larger vertebrates (Merilä & Hendry, 2014, Schilthuizen & Kellermann, 2014;

Stoks et al., 2014; Blanckenhorn, 2015). At the same time, laboratory longevity estimates

(e.g. Blanckenhorn, 1997; Reim et al., 2006; Blanckenhorn et al., 2007) generally do not well

reflect field mortality. Multiple field estimates of larval survivorship at various conditions

exist suggesting some counter-selection against large body size via the necessarily longer

development time (summarized in Blanckenhorn, 2007). However, sex- and size-specific

adult survivorship in the field was so far estimated only indirectly by Burkhard et al. (2002)

using age-grading by wing injuries. The present study assesses natural selection on

morphological traits by the fatal fungal parasite *Entomophthora spp*.

The parasitic fungus *Entomophthora scatophagae* regularly infects yellow dung fly adults

in Europe and North America (Hammer, 1941; Steinkraus & Kramer, 1988; Maitland, 1994;

Steenberg et al., 2001). Primarily at humid conditions infections can be epidemic (pers. obs.),

in which case infected dead flies can be found prominently exposed near cow pastures on

flowers, long grass or fences in a characteristic posture presumably effectively disseminating

fungal spores and/or attracting other flies (Maitland, 1994; Møller, 1993). Spore transmission

likely also occurs via physical contact, e.g. during copulation (Møller, 1993). The fungus is

fierce and effective at infecting and killing adult flies within hours or days and can be

manipulated, such that related, often species-specific such fungi are being employed for

biological control of some insect pests (e.g. Steenberg et al., 2001; Nielsen & Hayek, 2006).

This study took advantage of a fungus epidemic at our field population near Zürich,

Switzerland, in 2002.

I here assess viability, fecundity and sexual selection on morphology and fluctuating

asymmetry. Morphological traits reflecting body size are often assessed in selection studies

(Kingsolver et al., 2001; Kingsolver & Pfennig, 2004). Body size is one of the most important

quantitative traits of an organism, as it strongly affects most physiological and fitness traits

(Calder, 1984; Schmidt-Nielsen, 1984; Roff, 1992) and exhibits several prominent

evolutionary patterns in many organisms (Rensch, 1950; Fairbairn 1997; Blanckenhorn, 2000;

Blanckenhorn & Demont, 2004; Kingsolver & Pfennig, 2004). Depending on the taxon,

diverse traits are typically used as surrogates of body size, which are usually highly correlated

(i.e. integrated) within individuals due to pleiotropy, epistasis or gene linkage. Nevertheless,

for functional reasons selection on various body parts may differ (e.g. Preziosi & Fairbairn,

2000; Fairbairn, 2005), producing responses in correlated traits and thus generally prompting

a multivariate approach (Lande & Arnold, 1983). I focused on paired appendages (legs,

wings), so I could also assess fluctuating asymmetry (FA; Palmer & Strobeck, 1986). Small

and random deviations from the a priori perfect symmetry in bilaterally symmetric organisms,

i.e. FA, are presumed to reflect heritable developmental instability, such that individuals with

good genes and/or living in good conditions can produce more symmetric bodies in the face

of environmental stress, ultimately augmenting their fitness. Symmetric individuals

consequently should have greater survival prospects (viability selection), and should be more

successful at acquiring mates (sexual selection: Møller & Swaddle, 1997). However,

especially the latter notion, and the evidence, remain controversial (Møller & Thornhill, 1997;

Palmer, 2000; Polak, 2003; Klingenberg, 2003; Van Dongen, 2006; Knierim et al., 2007). In

yellow dung flies, Liggett et al. (1993) and Swaddle (1997) found a negative relationship

between FA and mating and foraging success, respectively, while Floate & Coughlin (2010)

found no evidence for FA being a useful biomarker of environmental stress exerted by toxic

livestock medications (ivermectin). Our previous studies of this species found that FA is not

heritable (Blanckenhorn & Hosken, 2003), that it does not increase with inbreeding (or

homozygosity: Hosken et al., 2000), and that FA does not affect male mating success in the

field, while nevertheless being negatively related to energy reserves (Blanckenhorn et al.,

2003). Of key relevance here is the postulated link between FA and immunocompetence,

predicting that more symmetric, but likely also larger individuals are better at fending off

internal parasites such as *Entomophthora* (see e.g. Rantala et al., 2000, 2004, 2007 and

Yourth et al., 2002, for various insect species), which was tested here.

Material and Methods

Study species

Yellow dung flies occur throughout the northern hemisphere and are particularly common

around cow pastures in central Europe. The species prefers cooler climates. In lowland central

Europe, each year has a spring (March – June) and an autumn season (September –

November), while during the hot midsummer (July and August) the flies largely disappear

from the pastures due to their heat sensitivity (Blanckenhorn, 2009).

Adult *S. stercoraria* are sit-and-wait predators of small flying insects, from which they

extract protein to produce sperm and eggs (Foster, 1967). Females spend most of their time

foraging for nectar and prey in the vegetation surrounding pastures. About once a week they

1 visit fresh cattle (or other) dung to deposit a clutch of eggs. Larvae feed on and develop in the

2 dung. Multiple males typically wait at the dung pat to mate with incoming females.

3 Copulation usually takes place in the surrounding grass or on the dung pat; during the ensuing

4 oviposition the male guards the female against other competitors (Parker, 1970). Competition

5 among males for females is very strong as the operational sex ratio is highly male biased

6 (Jann et al., 2000). Larvae face unpredictable spatio-temporal variation in local temperatures,

7 dung (i.e. food) quality and quantity, intra- and inter-specific competition, and dung drying,

8 all factors that ultimately largely determine their phenotypic adult body size. Towards the end

9 of the season the flies have to reach the overwintering pupal stage before the first winter frost

10 (Blanckenhorn, 2009).

11
12 **Fly sampling**
13
14 I sampled our population in Fehraltorf near Zurich (N47º52', E8º44') roughly once a month

15 between April and November 2002 (8 seasonal samples; total of $N = 541$ flies). Each time I

16 sampled one randomly selected but otherwise representative fresh dung pat, collecting all

17 single and paired flies on and ca. 20 cm around the pat to bring them alive to the laboratory.

18 As virtually no unpaired females occur at a pat in the field because competition for mates is

19 so intense, the number of pairs corresponds to the number of females present, and the

20 proportion of paired males corresponds to the operational sex ratio (females / males).

22 **Laboratory procedures**

23 Although dead flies infected with the fungus were occasionally found on and around the

24 pasture, these were too rare and haphazard to be sampled systematically. Instead, all collected

25 flies were kept alive in the laboratory in single 100 ml bottles with sugar and water for up to

26 two weeks. Infected flies would develop the fungus within few days and eventually die; non-

27 infected, recovered or resistant flies would not (acknowledging some random background

mortality). Females received dung to oviposit one clutch of eggs, which was counted, which

yields a good estimate of body size dependent clutch size (Honek, 1993; ca. 30 – 90 eggs:

Jann et al., 2000; Blanckenhorn, 2007).

In the end, work study students measured left and right wing length as well as fore, mid

and hind tibia length of each fly. Mean values for these paired traits were subsequently

calculated, as well as signed FA as (L – R), unsigned FA as (|L – R|) (both in mm) and

unsigned, size-corrected FA as (|L – R|) / mean(L, R) in %, as suggested by Palmer &

Strobeck (1986). Paired traits were measured twice blindly by the same person to estimate

measurement error relative to fluctuating asymmetry (Palmer & Strobeck, 1986), and to

calculate the repeatability of all trait measurements (Becker, 1992). All measurements were

taken with a binocular microscope at 16x magnification.

Statistical analysis

For each monthly sample, and overall, I calculated standardized viability selection

differentials (= gradients) for both sexes (binary variable: dead/alive = infected/uninfected),

sexual selection differentials for males (binary: mated/unmated), and fecundity selection

gradients for females based on their clutch size, using standard methods (Lande & Arnold,

1983; Arnold & Wade, 1984a,b; Brodie *et al*., 1995). It turns out that almost all females that

developed the fungus and eventually died in the laboratory did not lay any eggs, so fecundity

selection coefficients only refer to healthy (uninfected) females. Selection coefficients were

calculated for each trait (4 morphological and 4 asymmetry traits) separately for females and

males, and additionally for the first principal component (PC) of all (mean) appendages

signifying a compound index of body size, as such compound indices are often preferred by

some researchers as proxies of overall body size even though they are redundant in this

context.

Because the sizes of all appendages were highly positively correlated, and because FA and

size are mathematically related (see formulae above), we calculated only univariate linear

(β_{uni}) and corresponding non-linear (γ_{uni}) selection coefficients. To do so, for each seasonal

sample we produced standardized z-scores for trait x by subtracting the sample mean from

each value and dividing by the standard deviation: $z_i = (x_i - \bar{x})/SD_x$. Relative survival or

male pairing success was computed as absolute survival or pairing success (1 or 0) divided by

the sample proportion of survived flies or mated males, respectively (Brodie & Janzen, 1996),

and relative clutch size as absolute clutch size divided by mean clutch size. We used the

univariate model of relative fitness on standardized body size $w' = c + \beta_{uni}z$ to estimate the

linear selection intensities, and the corresponding quadratic model $w' = c + \beta_1 z + 0.5\gamma_{uni}z^2$ to

estimate univariate non-linear selection intensities, γ_{uni}. These linear coefficients (gradients)

reflect the combined effects of direct and indirect selection on body size (Endler, 1986). The

difference of the regression coefficients from a slope of zero (the null hypothesis of no

selection) was tested. For estimation of the coefficients least-squares regression was applied,

but for tests of significance logistic regression was used in case our measures of success were

binary (viability and mating success: Brodie *et al.,* 1995; cf. Janzen & Stern, 1998). A full

model with sex and season (including interactions) was performed to test for variation in

selection across these main factors.

Results

Fungus prevalence varied over the season and between the sexes. Infections (as high as 50%)

were most common during the cooler and more humid periods at the beginning (spring) and

the end of the season (autumn), whereas they were rare during the hotter summer (nearly 0%);

females were more affected than males (significant sex by season interaction: $\chi^2 = 29.80$; $P <$

0.001; Fig. 1). The probability of dying by fungal infection was unaffected by mean FA ($\chi^2 =$

1.99; $P = 0.159$) but increased with fly body size ($\chi^2 = 12.56$; $P < 0.001$; Table 1), an effect

that however varied among seasonal samples ($\chi^2 = 12.55$; $P < 0.001$) but not between the

sexes ($\chi^2 = 0.27$; $P = 0.602$).

Fecundity selection (based on clutch size) on female body size was significantly positive,

as is typical in this species (Jann et al., 2000; Kraushaar & Blanckenhorn, 2002). The

intensity of fecundity selection, i.e. the slope relating relative clutch size to standardized body

size (PC), varied significantly but unsystematically over the season (Table 1; interaction test:

$F_{6,109} = 3.50$, $P = 0.03$). These estimates refer only to uninfected flies because all females

infected with the fungus died before laying eggs, and therefore do not estimate fecundity

selection exerted by the fungus beyond the parasite's effect on adult mortality.

As usual in yellow dung flies, larger males had a mating advantage (Jann et al., 2000;

Kraushaar & Blanckenhorn, 2002; Blanckenhorn et al., 2003; main effect of body size (PC):

$\chi^2 = 14.23$; $P < 0.001$), while mean FA of all paired appendages did not affect male mating

success ($\chi^2 = 1.17$; $P = 0.279$; Table 1; Fig. 2). Except for one seasonal sample on 29 May, the

large male advantage was consistent throughout the season such that sexual selection intensity

did not significantly vary across the season (body size by season interaction: $\chi^2 = 0.11$; $P =$

0.920; Table 1; Fig. 2). Interestingly, this pattern of positive sexual selection was nullified by

fungal infection in those 28 (of a total of 47) infected males (of a total of 370 males, of which

148 had a mate) found mating in the field that later succumbed to the fungus (main effect of

fungal infection: $\chi^2 = 7.61$; $P = 0.006$; fungus by size interaction: $\chi^2 = 3.75$; $P = 0.053$; Fig. 3).

Table 1 gives directional (linear) selection coefficients, β, for the overall sample.

Nonlinear (quadratic) selection coefficients, γ, were mostly low and not significant and are

therefore not presented in Table 1. The only exception was female fecundity selection on

body size, for which $\gamma = 0.056 \pm 0.025$ was significantly positive, signifying accelerating

selection (which has been reported before: Blanckenhorn, 2007, 2009). Leg and wing lengths

were expectedly highly correlated in both sexes (range of bivariate correlations: $r = 0.887$ to

0.972), whereas FA of the legs and wings were largely uncorrelated (range: $r = -0.024$ to

+0.215). Measurement of all paired traits was generally repeatable using our methods ($R =$

0.83 – 0.97), which was also true for asymmetry (R = 0.53 – 0.61), so that FA could be

discerned from measurement error (all side by individual interactions P <0.01), fulfilling the

criteria of proper FA assessment (Palmer & Strobeck, 1986; Knierim et al., 2007).

Discussion

At our Swiss study population, the entomophagous fungal parasite *Entomophthora*

scatophagae, which has been described as a specific parasite of adult yellow dung flies at

several sites in Europe and North America (Hammer, 1941; Steinkraus & Kramer, 1988;

Maitland, 1994; Steenberg et al., 2001), shows high and generally fatal infection rates of up to

50% during the cooler and more humid periods of the year. I here documented that this

fungus exerts relatively strong and consistent negative viability selection on female and male

adult body size in *S. stercoraria* in those parts of the year, when dung flies are most common

(Table 1). Fungal infection further nullifies the usual large male mating advantage in sexual

selection (Borgia, 1982; Jann *et al*. 2000; Blanckenhorn et al., 2003; Fig. 3), but does not

affect female fecundity beyond its impact on mortality. This represents the first evidence

demonstrating viability disadvantages of large yellow dung flies mediated by a parasite,

which is generally rare in animals and particularly invertebrates (Blanckenhorn, 2000;

Kingsolver & Pfennig, 2004; Gotanda et al., 2015). These results complement previous

evidence of viability disadvantages of large flies at the juvenile stage, and lend further

credence to the notion that the male-biased sexual size dimorphism of yellow dung flies is

indeed roughly at evolutionary equilibrium (Blanckenhorn, 2007).

This study is merely phenomenological, meaning that mechanisms were not assessed.

Nonetheless, I speculate that the reduced parasite resistance of large flies signifies a trade-off

between body size and immunity (Rantala & Roff, 2005; Schwarzenbach &Ward, 2006;

Cotter et al., 2008). Based on age grading by wing injuries, Burkhard et al. (2002) found that

adult age (i.e. longevity) of yellow dung flies in the field tends to be positively related to body

size. Energy reserves also scale positively with body size (Reim et al., 2006; Blanckenhorn et

al., 2007) and positively influence mating success (Blanckenhorn et al., 2003). Adult

longevity under most environmental circumstances, including complete starvation, can

therefore generally be expected to increase with body size on physiological grounds. One

possible mechanism selecting against large body size is size-selective predation and/or

parasitism (Blanckenhorn, 2000). However, beyond expectations of a rough positive

correlation between predator and prey size (Brose et al., 2006; Vucic-Pestic et al., 2010),

evidence for systematic size-selectivity of predators is generally weak at best, also for yellow

dung flies (Blanckenhorn, 2000; Blanckenhorn et al., unpublished data). Size-selective

parasitism has been reported in some parasitoids (McGregor & Roitberg, 2000), but otherwise

few data exist (Zuk & Kolluru, 1998; Blanckenhorn, 2000). Rather than invoking selective

attraction of the parasite by larger hosts, I suspect that the ability to combat the parasite is

compromised in larger flies due to their generally greater absolute energy demands in a

stressful environment (trade-off hypothesis: Rantala & Roff, 2005; Reim et al., 2006;

Schwarzenbach & Ward, 2006, 2007; Cotter et al., 2008). Females were infected more by the

parasite here (Fig. 1), probably related to their generally greater reproductive burden (i.e. the

cost of producing expensive eggs rather than cheap sperm; cf. Nunn et al., 2009; but see e.g.

Rantala et al., 2007, for opposite results). Nevertheless, the standard sex differences in

reproductive (energetic) costs should be somewhat offset by the male-biased sexual size

dimorphism of yellow dung flies, which implies relatively greater costs when producing and

maintaining the larger and condition-dependent male body (Blanckenhorn, 2000, 2007), and

might explain why size-dependent viability selection here turned out to be steeper in males

(Table 1). Yellow dung fly females indeed produce higher heritable levels than males of

phenoloxidase (PO; Schwarzenbach et al., 2005), one of the central mediators of insect

immunity (Schmid-Hempel, 2005; Rolff & Reynolds, 2009; González-Santoyo &

CórdobaAguilar, 2012), and higher PO levels decrease adult longevity in this species,

demonstrating a trade-off (Schwarzenbach & Ward, 2006). However, higher PO levels did not

lead to greater resistance against mites or another fungus (Schwarzenbach & Ward, 2007),

and PO is also unrelated to body size in *S. stercoraria* (Schwarzenbach et al., 2005). Overall,

therefore, the evidence in favor of immunity mediating the higher mortality of large-bodied

dung flies here remains limited.

In contrast to body size, fluctuating asymmetry (FA) of legs and wings influenced none of

the fitness components investigated here (contrary to Liggett et al., 1993, but confirming

Blanckenhorn et al's, 2003, earlier sexual selection study). I had expected, based on evidence

in other animals (Rantala et al., 2000, 2004), that low FA would be a signal of greater

immunocompetence augmenting resistance against parasites, but this was not found. It is not

unlikely that FA is a bad indicator of developmental stability in general, as various reviews

have revealed no clear verdict based on the available evidence on this question, so the entire

concept remains controversial (Møller & Thornhill, 1997; Møller & Swaddle, 1997; Palmer,

2000; various articles in Polak, 2003; Van Dongen, 2006; Knierim et al., 2007). In yellow

dung flies, beyond Liggett et al. (1993) the evidence for a role of FA in sexual selection is

close to nil (Blanckenhorn et al., 2003; Blanckenhorn & Hosken, 2003; this study). What

remains is that FA reliably indicates at least hot temperature stress in this species (Hosken et

al., 2000; Blanckenhorn & Hosken, further unpublished data), even though Floate & Coughlin

(2010) concluded that FA is no good biomarker of toxic chemical residues in cattle dung. As

the survival of flies infected with the fierce entomophagous fungus *Entomophthora*

scatophagae here was related to the size (discussed above) but not the fluctuating asymmetry

of wings and legs, I conclude that FA is no good indicator of immunocompetence in yellow

dung flies either (Rantala et al., 2000, 2004, 2007; Rantala & Roff, 2005; Yourth et al., 2002;

Schwarzenbach &Ward, 2006; Cotter et al., 2008).

Acknowledgements

I thank the University of Zurich and the Swiss National Fund for financial support over the

years, and Ana, Lukas, Nils and Patricia for doing the measurements.

References (77; 1753 words)

Andersson. M. 1994. *Sexual Selection*. Princeton University Press, Princeton, NJ.

Arnold, S. J., & Wade, M.J. 1984a. On the measurement of natural and sexual selection: applications. *Evolution* **38**: 720-734.

Arnold, S. J., & Wade, M.J. 1984b. On the measurement of natural and sexual selection: theory. *Evolution* **38**: 709-719.

Becker, W.A. 1992. *Manual of Quantitative Genetics*. *5th edition*. Academic Enterprises, Pullman.

Blanckenhorn, W.U. 1997. Altitudinal life history variation in the dung flies *Scathophaga stercoraria* and *Sepsis cynipsea*. *Oecologia* **109**: 342-352.

Blanckenhorn, W.U. 2000. The evolution of body size: what keeps organisms small? *Quart. Rev. Biol.* **75**: 385-407.

Blanckenhorn, W.U. 2007. Case studies of the differential equilibrium hypothesis of sexual size dimorphism in dung flies. Pages 106-114 in: D.J. Fairbairn, W.U. Blanckenhorn, T. Székely, eds. SEX, SIZE AND GENDER ROLES. Evolutionary Studies of Sexual Size Dimorphism. Oxford University Press, Oxford.

Blanckenhorn, W.U. 2009. Causes and consequences of phenotypic plasticity in body size: the case of the yellow dung fly *Scathophaga stercoraria* (Diptera: Scathophagidae). Pages 369-422 in: D.W. Whitman, T.N. Ananthakrishnan, eds. Phenotypic plasticity of insects: Mechanism and consequences. Science Publishers, Enfield NH, USA.

Blanckenhorn, W.U. 2015. Yellow dung fly body size evolution in the field: response to climate change? *Evolution* **69**: 2227-2234.

Blanckenhorn, W.U. & M. Demont. 2004. Bergmann and converse Bergmann latitudinal clines in arthropods: two ends of a continuum? *Integr. Comp. Biol.* **44**: 413-424.

Blanckenhorn, W.U., Kraushaar, U. & Reim, C. 2003. Sexual selection on morphological and physiological traits and fluctuating asymmetry in the yellow dung fly. *J. Evol. Biol.* **16**: 903-913.

Blanckenhorn, W.U., Fanti, J. & Reim, C. 2007. Size-dependent energy reserves, energy utilization and longevity in the yellow dung fly. Physiological Entomology 32: 372-381.

Blanckenhorn, W.U. & Hosken, D.J. 2003. The heritability of three condition measures in the yellow dung fly. *Behav. Ecol.* **14***:* 612-618.

Borgia, G. 1982. Experimental changes in resource structure and male density: size-related
differences in mating success among male *Scathophaga stercoraria*. *Evolution* **36**: 307-
315.
Brodie III., E.D. & Janzen, F.J. 1996. On the assignment of fitness values in statistical
analyses of selection. *Evolution* **50**: 437-442.
Brodie III., E.D., Moore, A.J. & Janzen, F.J. 1995. Visualizing and quantifying natural
selection. *Trends Ecol. Evol.* **10**: 313-318.
Brose U, et al. 2006. Consumer-resource body-size relationships in common food webs.
Ecology **87**: 2411-2417.

Burkhard, D.U., Ward, P.I. & Blanckenhorn, W.U. 2002. Using age-grading by wing injuries
to estimate size-dependent adult survivorship in the field: a case study of the yellow dung
fly *Scathophaga stercoraria*. *Ecol. Entomol.* **27**: 514-520.
Calder, W.A. 1984. *Size, Function, and Life History*. Harvard University Press, Cambridge.
Cotter, S., Myatt, J., Benskin, C. & Wilson, K. 2008. Selection for cuticular melanism reveals
immune function and life-history trade-offs in Spodoptera littoralis. *J. Evol. Biol.*, **21**:
1744-1754.
Cox, R.M. & Calsbeek, R. 2009. Sexually antagonistic selection, sexual dimorphism and the
resolution of intralocus sexual conflict. *Am. Nat.* **173**: 176-187.
Endler, J.A. 1986. *Natural Selection in the Wild*. Princeton University Press, Princeton.
Fairbairn, D.J. 1997. Allometry for sexual size dimorphism: Pattern and process in the
coevolution of body size in males and females. *Ann. Rev. Ecol. Syst.* **28**: 659–687.
Fairbairn, D.J. 2005. Allometry for sexual size dimorphism: testing two hypotheses for
Rensch's rule in the water strider *Aquarius remigis*. *Am. Nat.* **166**: S69--84
Floate, K.D. & Coghlin, P. 2010. No support for fluctuating asymmetry as a biomarker of
chemical residues in livestock dung. *Can. Entomol.* **152**: 354-368.
Foster, W. 1967. Hormone-mediated nutritional control of sexual behavior in male dung flies.
Science **158**: 1596-1597.
González-Santoyo, I. & Córdoba-Aguilar, A. 2012. Phenoloxidase: a key component of the
insect immune system. *Entomo. Exp. Appl.* **142**: 1-16.
Gotanda, K., Correa, C., Turcotte, M.M., Rolshausen, G. & Hendry, A.P. 2015. Linking
macro-trends and micro-rates: re-evaluating micro-evolutionary support for Cope's rule.
Evolution **69**: 1345-1354.

Hammer. O. 1941. Biological and ecological investigations on flies associated with pasturing cattle and their excrement. *Vidensk. Medd. Dansk Naturh. Foren.* **105**: 140-393.

Honek, A. 1993. Instraspecific variation in body size and fecundity in insects: a general relationship. *Oikos* **66**: 483-492.

Hosken, D.J., Blanckenhorn, W.U. & Ward, P.I. 2000. Developmental stability in the yellow dung fly (*Scathophaga stercoraria*): fluctuating asymmetry, heterozygosity and environmental stress. *J. Evol. Biol.* **13**: 919–926.

Jann, P., Blanckenhorn, W. U. & Ward, P. I. 2000. Temporal and microspatial variation in the intensities of natural and sexual selection in the yellow dung fly *Scathophaga stercoraria*. *J. evol. Biol.* **13**: 927-938.

Kingsolver, J.G., Hoekstra, H.E., Hoekstra, J.M., Berrigan, D., Vignieri, S.N., Hill, C.E., Hoang, A., Gibert, P. & Beerli, P. 2001. The strength of phenotypic selection in natural populations. *Am. Nat.* **157**: 245-261.

Kingsolver, J.G. & D.W. Pfennig. 2004. Individual-level selection as a cause of Cope's rule of phyletic size increase. *Evolution* **58**:1608-1612.

Knierim, U., Van Dongen, S., Forkman, B., Tuyttens, F.A.M., Špinka, M., Campo, J.L., & Weissengruber, G.E. 2007. Fluctuating asymmetry as an animal welfare indicator — A review of methodology and validity. *Physiol Behav.* **92**: 398-421.

Kraushaar, U. & Blanckenhorn, W.U. 2002. Population variation in sexual selection and its effect on body size allometry in two species of flies with contrasting sexual size dimorphism. *Evolution* **56**: 307-321.

Lande, R. & Arnold, S.J. 1983. The measurement of selection on correlated characters. *Evolution* 37: 1210-1226.

Liggett, A. C., Harvey, I.F. & Manning, J.T. 1993. Fluctuating asymmetry in *Scathophaga stercoraria* L.: successful males are more symmetrical. *Anim. Behav.* **45**: 1041-1043.

Maitland, D.P. 1994. A parasitic fungus infecting yellow dungflies manipulates host perching behaviour. *Proc. Roy. Soc. Lond. B* **258**: 187-193.

McGregor, R. R. & Roitberg, B.D, 2000. Size-selective oviposition by parasitoids and the

evolution of life-history timing in hosts: Fixed preferences vs. frequency-dependent host

selection. *Oikos* **89**: 305-312.

Merilä, J. & Hendry, A.P. 2014. Climate change, adaptation, and phenotypic plasticity: the

problem and the evidence. *Evol. Appl.* **7**: 1-14.

Merilä, J., Sheldon, B.C. & Kruuk, L.E.B. 2001. Explaining stasis: microevolutionary studies

in natural populations. *Genetica* **112-113**: 199-222.

Møller, A. P. 1993. A fungus infecting domestic flies manipulates sexual behaviour of its

host. *Behav. Ecol. Sociobiol.* **33**: 403-407.

Møller, A.P. & Swaddle, J.P. 1997. *Asymmetry, Developmental Stability and Evolution.*

Oxford University Press, Oxford.

Møller, A.P. & Thornhill, R. 1997. A meta-analysis of the heritability of developmental

stability. *J. evol. Biol.* **10**: 1-16.

Mousseau, T.A. & Roff, D.A. 1987. Natural selection and the heritability of fitness

components. *Heredity* **59**: 181-197.

Nielsen, C. & Hayek, A.E. 2006. Diurnal pattern of death and sporulation in *Entomophaga*

maimaiga-infected *Lymantria dispar*. *Entomol. Exp. Appl.* **118**: 237-243.

Nunn, C.L., Lindenfors, P., Pursall, E.R. & Rolff, J. 2009. On sexual dimorphism in immune

function. *Phil. Trans. Roy. Soc. Lond. B* **364**: 61-69.

Palmer, A.R. 2000. Quasi-replication and the contract of error: lessons from sex ratios,

heritabilities and fluctuating asymmetry. *Ann. Rev. Ecol. Syst.* **31**: 441-480.

Palmer, A. R. & Strobeck, C. 1986. Fluctuating asymmetry: measurement, analysis, patterns.

Ann. Rev. Ecol. Syst. **17**: 391-421.

Parker, G.A. 1970. The reproductive behaviour and the nature of sexual selection in

Scathophaga stercoraria L. (Diptera: Scathophagidae) II. The fertilization rate and the

spatial and temporal relationships of each sex around the site of mating and oviposition. *J.*

Anim. Ecol. **39**: 205-228.

Parker, G.A. 1979. Sexual selection and sexual conflict. In: *Sexual Selection and*

Reproductive Competition in Insects (N. S. Blum & N. A. Blum, eds), pp. 123-166.

Academic Press, New York.

Preziosi, R.F. & Fairbairn, D.J. 2000. Lifetime selection on adult body size and components

of body size in a waterstrider: opposing selection and maintenance of sexual size

dimorphism. *Evolution* **54**: 558-566.

Polak, M. (ed.) 2003. *Developmental instability. Causes and consequences*. Oxford: Oxford University Press.

Rantala, M.J. & Roff, D.A. 2005. An analysis of trade-offs in immune function, body size and development time in the Mediterranean field cricket, *Gryllus bimaculatus*. *Funct. Ecol.* **19**: 323-330.

Rantala, M.J., Koskimäki, J., Taskinen, J., Tynkkynen, K. & Suhonen, J. 2000. Immunocompetence, developmental stability and wing spot size in the damselfly *Calopteryx splendens* L. *Proc. Roy. Soc. Lond. B* **267**: 2453–2457.

Rantala, M.J., Ahtiainen, J.J. & Suhonen, J. 2004. Fluctuating asymmetry and immune function in a field cricket. *Oikos* **109**: 479–484.

Rantala, M.J., Roff, D.A. & Rantala, L.M. 2007. Forcepts size and immune function in the earwig *Forficula auricularia*. *Biol. J. Linn. Soc.* **90**: 509-516.

Reim, C., Teuschlm Y, & Blanckenhorn, W.U. 2006. Size-dependent effects of larval and adult food availability on reproductive energy allocation in the yellow dung fly. *Funct. Ecol.* **20**: 1012-1021.

Rensch, B. 1950. Die Abhängigkeit der relativen Sexualdifferenz von der Körpergrösse. *Bonn. Zoolog. Beitr.* **1**: 58-69.

Roff, D.A. 1992. *The Evolution of Life Histories*. Chapman & Hall, New York.

Rolff, J. & Reynolds, S. 2009. *Insect Infection and Immunity: Evolution, Ecology, and Mechanisms*. Oxford University Press.

Schilthuizen, M. & Kellermann, V. 2014. Contemporary climate change and terrestrial invertebrates: evolutionary versus plastic changes. *Evol. Appl.* **7**: 56–67.

Schmid-Hempel, P. 2005. Evolutionary ecology of insect immune defenses. *Annu. Rev. Entomol.* **50**: 529-551.

Schwarzenbach, G.A., Hosken, D.J. & Ward, P.I. 2005. Sex and immunity in the yellow dung fly *Scathophaga stercoraria*. *J. evol. Biol.* **18**: 455-463.

Schwarzenbach, G.A. & Ward, P.I. 2006. Responses to selection on phenoloxidase activity in yellow dung flies. *Evolution* **60**: 1612-1621.

Schwarzenbach, G.A. & Ward, P.I. 2007. Phenoloxidase activity and pathogen resistance in the yellow dung fly *Scathophaga stercoraria*. *J. evol. Biol.* **20**: 2192-2199.

Schmidt-Nielsen, K. 1984. *Scaling. Why is animal size so important?* Cambridge University

Press, Cambridge.

Sigurjónsdóttir, H. & Snorrason, S.S. 1995. Distribution of male yellow dungflies around

oviposition sites: the effect of body size. *Ecol. Entomol.* **20**: 84-90.

Steenberg, T., Jespersen, J.B., Jensen, K.-M.V., Nielsen, B.O. & Humber, R. A. 2001.

Entomopathogenic fungi in flies associated with pastured cattle in Denmark. *J. Invert.*

Pathol. 77: 186-197.

Steinkraus, D.C. & Kramer, J.P. 1988. *Entomophthora scatophagae*, a fungal pathogen of the

yellow dung fly, *Scatophaga stercoraria. Mycotaxon* **32**: 105-113.

Stoks, R., Geerts, A.N. & de Meester, L. 2014. Evolutionary and plastic responses of

freshwater invertebrates to climate change: realized patterns and future potential. *Evol.*

Appl. **7**: 42-55.

Swaddle, J.P. 1997. Developmental stability and predation success in an insect predator-prey

system. *Behav. Ecol.* **8**: 433-436.

Van Dongen, S.2006. Fluctuating asymmetry and developmental instability in evolutionary

biology: past, present and future. *J. Evol. Biol.* **19**: 1727–43.

Vucic-Pestic, O., Rall, B.C., Kalinkat, G. & Brose, U. 2010. Allometric functional response

model: Body masses constrain interaction strengths. *J. Anim. Ecol.* **79**: 249-256.

Yourth, C.P., Forbes, M. & Smith, B.P. 2002. Immune expression in a damselfly is related to

time of season, not to fluctuating asymmetry or host size. *Ecol. Entomol.* **27**: 123–128.

Zuk, M. & Kolluru, G.R. 1998. Exploitation of sexual signals by predators and parasitoids.

Quart. Rev. Biol. **73**: 415-438.

Table 1: Overall intensities ($\beta \pm$ 95% CI) of female and male adult viability selection ($N_f =$

171 & $N_m =$ 370) exerted by the fungus *Entomophthora*, female fecundity selection (clutch

size; $N_f =$ 126), and male sexual selection (pairing success; $N_m =$ 370) for one Swiss

population of yellow dung flies (*Scathophaga stercoraria*) over the season 2002. Significant

coefficients are in bold (*P < 0.001*).

Trait	Adult viability				Female fecundity		Male mating success	
	β_f	95% CI	β_m	95% CI	β	95% CI	β	95% CI
Hind tibia length	-0.158	0.305	**-0.306**	0.252	**0.185**	0.027	**0.259**	0.149
Mid tibia length	-0.089	0.310	**-0.328**	0.277	**0.179**	0.029	**0.228**	0.150
Fore tibia length	-0.196	0.304	-0.221	0.251	**0.174**	0.029	**0.226**	0.159
Wing length	-0.254	0.307	**-0.324**	0.248	**0.179**	0.029	**0.267**	0.147
Overall PC size	-0.099	0.292	**-0.317**	0.302	**0.184**	0.029	**0.253**	0.136
Hind tibia FA	-0.020	0.333	-0.086	0.258	-0.023	0.045	0.051	0.151
Mid tibia FA	0.204	0.231	0.003	0.219	-0.026	0.041	0.010	0.173
Fore tibia FA	-0.037	0.299	0.224	0.283	-0.024	0.045	0.040	0.180
Wing FA	-0.187	0.321	0.046	0.279	0.037	0.045	-0.041	0.168
Mean FA	0.035	0.284	0.097	0.327	-0.021	0.040	0.071	0.134

1 **Figure 1:** Proportion of male (filled squares) and female (open circles) flies infected by the

2 fungus *Entomophthora* over the season 2002.

1 **Figure 2:** Body size (top; here exemplified by wing length) and mean percentage of

2 fluctuating asymmetry (FA; bottom) of all traits for unpaired (filled squares) and paired males

3 (open squares) over the season.

1 **Figure 3**: Body size (here exemplified by hind tibia length) of unpaired (filled squares) and

2 paired males (open squares) when they were infected by the fungus or not (all seasonal

3 samples combined).