Title: Adaptation in a fibronectin binding autolysin of *Staphylococcus saprophyticus*

Authors: Tatum D. Mortimer[1,2], Douglas S. Annis[3], Mary B. O'Neill[1,4], Lindsey L. Bohr[1,2], Tracy
M. Smith[1,5], Hendrik N. Poinar[6,7,8,9], Deane F. Mosher[3], Caitlin S. Pepperell[1,5]

Affiliations:

[1]Department of Medical Microbiology and Immunology, School of Medicine and Public Health,
University of Wisconsin-Madison, Madison, United States

[2]Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, United
States

[3]Department of Biomolecular Chemistry, School of Medicine and Public Health, University of
Wisconsin-Madison, Madison, WI, USA

[4]Laboratory of Genetics, University of Wisconsin-Madison, Madison, Wisconsin, United States

[5]Department of Medicine (Infectious Diseases), School of Medicine and Public Health,
University of Wisconsin-Madison, Madison, United States

[6]McMaster Ancient DNA Centre, Department of Anthropology, McMaster University, Hamilton,
Canada

[7]Department of Biology, McMaster University, Hamilton, Canada

[8]Michael G. DeGroote Institute for Infectious Disease Research, McMaster University, Hamilton,
Canada

[9]Humans and the Microbiome Program, Canadian Institute for Advanced Research, Toronto,
Canada

Abstract:

Human-pathogenic bacteria are found in a variety of niches, including free-living, zoonotic, and microbiome environments. Identifying bacterial adaptions that enable invasive disease is an important means of gaining insight into the molecular basis of pathogenesis and understanding pathogen emergence. *Staphylococcus saprophyticus*, a leading cause of urinary tract infections, can be found in the environment, food, animals, and the human microbiome. We identified a selective sweep in the gene encoding the Aas adhesin, a key virulence factor that binds host fibronectin. We hypothesize that the mutation under selection (*aas*_2206A>C) facilitates colonization of the urinary tract, an environment where bacteria are subject to strong shearing forces. The mutation appears to have enabled emergence and expansion of a human pathogenic lineage of *S. saprophyticus.* These results demonstrate the power of evolutionary genomic approaches in discovering the genetic basis of virulence and emphasize the pleiotropy and adaptability of bacteria occupying diverse niches.

Importance:

Staphylococcus saprophyticus is an important cause of urinary tract infections (UTI) in women, which are common, can be severe, and are associated with significant impacts to public health. In addition to being a cause of human UTI, *S. saprophyticus* can be found in the environment, in food, and associated with animals. After discovering that UTI strains of *S. saprophyticus* are for the most part closely related to each other, we sought to determine whether these strains are specially adapted to cause disease in humans. We found evidence suggesting that a mutation in the gene *aas* is advantageous in the context of human infection. We hypothesize that the mutation allows *S. saprophyticus* to survive better in the human urinary tract. These results show how bacteria found in the environment can evolve to cause disease.

Introduction.

Urinary tract infections (UTI) are a global health problem of major significance, with an estimated annual incidence of 150-250 million and a lifetime risk of 50% among women (1–3). The associated costs to individuals and health care systems are substantial, with recent estimates from the United States numbering in the billions per year (4). UTIs are also associated with severe complications such as pyelonephritis, sepsis and premature labor (4). *Staphylococcus saprophyticus* is second only to *Escherischia coli* as a cause of UTI in reproductive aged women (5, 6).

S. saprophyticus can be found in diverse niches including the environment, foods, livestock, and as a pathogen and commensal of humans. Several features of the epidemiology of *S. saprophyticus* suggest that infections leading to UTIs are acquired from the environment, rather than as a result of person-to-person transmission (7). This implies that adoption of the pathogenic niche by *S. saprophyticus* has not entailed a trade-off in its ability to live freely in the environment. A recent PCR-based survey of virulence factors in clinical and animal associated isolates showed that *dsdA*, a gene encoding D-serine deaminase that is important for survival in urine (8), and *uafA* and *aas*, adhesins that mediate binding to uroepithelium (9, 10), were present in all isolates surveyed (11), suggesting an underlying pleiotropy, with these virulence factors playing important roles in the diverse environments occupied by *S. saprophyticus*.

The human urinary tract could represent an evolutionary dead end for *S. saprophyticus* with "virulence factors" such as DsdA, UafA and Aas serving an essential function in the primary environmental niche and enabling invasion of the urinary tract as an accidental by-product of this unknown primary function. In this case, we would expect urinary isolates to be interspersed throughout a phylogeny of isolates from the primary niche(s). However, our previous research (7) indicated that human urinary tract infections are associated with a specific lineage of *S. saprophyticus*. Invasion of the human urinary tract enables *S. saprophyticus* to grow to large numbers in urine, isolated from competing bacterial species, before being re-deposited in the environment. This is analogous to *Vibrio cholerae*, which cycles through human and environmental niches and grows to high abundance in the human gut before being deposited in the environment via stool (12, 13). Based on our previous observations and the example provided by other human pathogens that cycle through the environment, we hypothesized that the human urinary tract is an ecologically important niche for *S. saprophyticus* and sought to identify genetic signatures of adaptation to this niche.

The increased availability of sequencing data has enabled comparative genomic approaches that have led to identification of changes in gene content in association with pathogen emergence and shifts in host association. Several notable human pathogens, including *Mycobacterium tuberculosis*, *Yersinia pestis*, and *Francisella tulerensis*, are the product of a single emergence characterized by gene loss and horizontal acquisition of virulence factors (14–16). Similarly, genomic analysis of *Enterococcus faecium* revealed gene gains and losses affecting metabolism and antibiotic resistance in the emergence of a hypermutable hospital-adapted clade that coincided with the profound shift in hospital ecology caused by development

of antibiotics (17). Gene gains via recombination have also allowed *Staphylococcus aureus*
ST71 to emerge into a bovine-associated niche (18).

Using contemporary and ancient genomic data from strains of *S. saprophyticus*, we found
previously that UTI-associated lineages of *S. saprophyticus* were not attendant with specific
gene gains or losses; the evolutionary genetic processes underlying *S. saprophyticus'* adoption
of the human-pathogenic niche are likely more subtle than what has been described for
canonical pathogens (7). Here we have identified one of the mechanisms underlying *S.*
saprophyticus' adaptation to the uropathogenic niche: a selective sweep in the Aas adhesin,
which is associated with an apparently large-scale expansion into the human-pathogenic niche.
This is, to our knowledge, the first identification of a single nucleotide sweep in a bacterium.

Results.

We reconstructed the phylogeny of *S. saprophyticus* isolates (Supplementary Table 1) from a
whole genome alignment using maximum likelihood inference implemented in RAxML (Figure
1). The bacterial isolates are separated into two clades, which we have previously named
Clades P and E (7). In both clades, human associated lineages are nested among isolates from
diverse sources, including food (cheese rind, ice cream, meat), indoor and outdoor
environments, and animals. Interestingly, cheese rinds harbor diverse strains of *S.*
saprophyticus, which cluster with both human- and animal-pathogenic strains.

Thirty-three of 37 modern, human-pathogenic isolates are found within a single lineage (that we
term lineage U, for UTI-associated) to which bovine-pathogenic (mastitis), food-associated
isolates, and an ancient genome are basal. Given the association between this lineage and
illness in humans, we were curious about its potential adaptation to the human pathogenic
niche. The placement of the 800-year old strain between bovine and human-associated
lineages suggests it could represent a generalist intermediate between human-adapted and
bovid-adapted strains.

Core genome analysis of the 58 isolates of *S. saprophyticus* in our sample showed substantial
variability in gene content; the core genome is composed of 1798 genes, and there are an
additional 7110 genes in the pan genome. We found previously that uropathogenic isolates of *S.*
saprophyticus were not associated with any unique gene content (7). Given the variability in
accessory gene content among this larger sample of isolates, we decided to test for relative
differences in accessory gene content between human clinical isolates and other isolates using
Scoary (19), which performs a genome wide association study (GWAS) using gene presence
and absence. We did not identify any genes that were significantly associated with the human
pathogenic niche after correction for multiple hypothesis testing using the Bonferroni method.

In addition to the variability observed in gene content, analyses of the core genome also
indicated relatively frequent recombination among *S. saprophyticus*. We identified recombinant
regions with Gubbins (20), which identifies regions with high densities of substitutions. These
results indicated that 70% of sites in the *S. saprophyticus* alignment have been affected by
recombination. Recombination can affect bacterial evolution both by introducing novel
polymorphisms from outside the population and by reshuffling alleles without increasing overall

diversity. Considering sites that are reshuffled within the *S. saprophyticus* sample as
recombinant, we estimate a ratio of recombinant to non-recombinant SNPs of 3.4. When
considering only the SNPs that introduce novel diversity as recombinant, our estimate of the
ratio of recombinant SNPs to non-recombinant SNPs is 0.51. The mean r/m of branches in the
phylogeny is 0.82 as estimated by Gubbins (range: 0-7.6). Removal of recombinant SNPs did
not affect the topology of the maximum likelihood phylogeny (Supplementary Figure 1). We
observed regional patterns in the amount of recombination inferred, and, as expected,
recombination appears frequent at mobile elements such as the staphylococcal cassette
chromosomes (SCC$_{15305RM}$ and SCC$_{15305cap}$) and *v*Ss15305 (10).

Adaptation to a new environment may be facilitated by advantageous mutations that quickly rise
in frequency, leaving a characteristic genomic imprint: reduced diversity at the target locus and
nearby linked loci (i.e. selective sweep, (21, 22)). In order for positive selection to be evident as
a local reduction in diversity, there must be sufficient recombination that the target locus is
unlinked from the rest of the genome; for this reason, scans for sweeps have been used
primarily for sexually reproducing organisms (23–26). As described above and in prior work (7),
we found evidence of frequent recombination among *S. saprophyticus*. We hypothesized that *S.*
saprophyticus' transition to the uropathogenic niche may have been driven by selection for one
or more mutations that were advantageous in the new environment, and that levels of
recombination have been sufficient to preserve the signature of a selective sweep at loci under
positive selection. We therefore used a sliding window analysis of diversity along the *S.*
saprophyticus alignment as an initial screen for positive selection. We identified a marked
regional decrease in nucleotide diversity (π) and Tajima's D (TD) that is specific to lineage U
(Figure 2, Supplementary Figure 2). The region with decreased π /TD corresponds to
1,760,000-1,820,000 bp in *S. saprophyticus* ATCC 15305 and has the lowest values of π and
TD in the entire alignment. We investigated the sensitivity of our sliding window analyses to
sampling by randomly subsampling lineage U isolates to the same size as Clade E (n = 10); we
found the results to be robust to changes in sampling scheme and size (Supplementary Figure
3).

To complement the sliding window analysis and pinpoint candidate variants under positive
selection, we used an approach based on allele frequency differences between bacterial
isolates from different niches. We calculated Weir and Cockerham's F$_{ST}$ (27) for single
nucleotide polymorphisms (SNPs) in the *S. saprophyticus* genome using human association
and non-human association to define populations. The region of low π /TD included three non-
synonymous variants in the top 0.05% of F$_{ST}$ values (Supplementary Table 2). One of these
variants was fixed among human-associated isolates in lineage U (position 1811777 in ATCC
15305, F$_{ST}$ = 0.48) and distinct from the ancestral allele found in basal lineages of Clade P,
including the ancient strain of *S. saprophyticus* Troy. This suggests that the variant may have
been important in adaptation to the human urinary tract. To assess the significance of the F$_{ST}$
value for this variant, we performed permutations by randomly assigning isolates as human
associated, and we did not achieve F$_{ST}$ values higher than 0.28 in 100 permutations.

Selective sweeps may be evident as a longer than expected haplotype block, since neutral
variants linked to the adaptive mutation will also sweep to high frequencies (28). Given the

evidence suggesting there was a selective sweep at this locus, we used haplotype based
statistics to test for such a signature in the *S. saprophyticus* alignment. Haplotype-based
methods are hypothesized to not be applicable to bacteria due to differences between crossing
over and bacterial patterns of recombination (29), but the methods had not been tested in a
scenario akin to a classical sweep, in which local changes in diversity and the SFS have been
observed. We found that the variant at position 181177 did show a signature of a sweep using
the extended haplotype homozygosity (EHH) statistic (28) (Supplementary Figure 4). However,
the variant did not have an extreme value of nS_L, which compares haplotype homozygosity for
ancestral and derived alleles (30), after normalization by the allele frequency.

The variant of interest (*aas*_2206A>C) causes a threonine to proline change in the amino acid
sequence of Aas*,* a bifunctional autolysin with a fibronectin binding domain (Figure 4, (31)).
There are 8 additional nonsynonymous polymorphisms in the fibronectin binding domain;
however, none are as highly associated with human pathogenic isolates (Supplementary Table
2). Adhesins such as Aas are important in the pathogenesis of *S. saprophyticus* urinary tract
infections, and this gene has been previously implicated as a virulence factor (9, 31, 32).

The Aas variant is in a region known to bind fibronectin (Figure 5, (31)) and may be under
selection because it affects adhesion to this host protein. We used ELISAs to investigate
potential effects of *aas*_2206A>C on binding to fibronectin and thrombospondin-1, which binds
to this region of the homologous AtlE amidase from *Staphylococcus epidermidis* (33).
Staphylococcal autolysins contains 3 C-terminal repeats (R1-R3), which can each be divided
into two subunits (a and b) based on structural information (34). We confirmed that Aas R1a1b
binds fibronectin and discovered that it also binds thrombospondin (Supplementary Figure 5);
there was no detectable difference between the ancestral and derived R1a1b alleles in binding
to fibronectin or thrombospondin (human or bovine).

Interestingly, we observed several instances of recombination of the *aas* variant. In each case,
the recombination event reinforced the association of the derived allele with human infection.
Two of the non-human-associated bacterial isolates in lineage U – an isolate from a pig and a
second from cheese rind – had evidence of a recombination event at the *aas* locus resulting in
acquisition of the ancestral allele. Conversely, one of the human UTI isolates in Clade E (for
which the ancestral allele is otherwise fixed) acquired the derived *aas* variant.

Several human pathogens appear to have undergone recent population expansion (35–38). We
wondered whether the uropathogenic lineage of *S. saprophyticus* might also have undergone a
recent change in its effective size. The genome wide estimate of TD for lineage U was negative
(-0.58), which is consistent with population expansion. We used the methods implemented in
∂a∂i (39) to identify the demographic model that best fit the observed synonymous SFS of
lineage U (Figure 5).

The synonymous SFS showed an unexpected excess of high frequency derived alleles, which
we hypothesized were the result of gene flow from populations with ancestral variants. Within
population recombination has been shown to have no effect on SFS-based methods of
demographic inference in bacteria (40). However, external sources of recombination were not
modeled in previous studies. We used SimBac (41) to simulate bacterial populations with a

range of internal and external recombination rates. Similar to previous studies, we did not find
that within population recombination had an effect on the value of Tajima's D. However, we
found that recombinant tracts from external sources resulted in positive values of Tajima's D
(Supplementary Figure 6). Positive values of Tajima's D are also associated with population
bottlenecks and balancing selection.

We used fastGEAR (42) to identify recombinant tracts that originated outside of lineage U, and
these sites were removed from the analysis prior to demographic inference. We compared five
demographic models (constant size, instantaneous population size change, exponential
population size change, instantaneous population size change followed by exponential, and two
instantaneous population size changes , Figure 6) and used bootstrapping to estimate the
uncertainty of the parameters and to adjust the composite likelihoods using the Godambe
Information Matrix implemented in ∂a∂i (43). We found significant evidence for an expansion in
all models (Table 1). The best fitting model was an instantaneous contraction followed by an
instantaneous expansion, in which the population underwent a tight bottleneck followed by a 15-
fold expansion without recovering to its ancestral size ($v = N_e/N_{anc}$, τ = generations/N_{anc}, v_A: 2.9 x
10^{-2}, v_B: 4.5 x 10^{-1}, τ_A: 1.2 x 10^{-1}, τ_B: 3.1 x 10^{-3}).

Recombination and positive selection are known to confound the inference of bacterial
demography (40), so we used simulations to investigate their effects on our demographic
inference for uropathogenic *S. saprophyticus.* We used SFS_CODE (44) to simulate positive
selection (with a range of recombination rates) and evaluate its effects on the accuracy of
demographic inference with ∂a∂i. The method implemented in ∂a∂i relies on inference from the
synonymous SFS, but it's possible for synonymous variation to be affected by selection,
particularly at low rates of recombination (40, 45). Neutral simulations with gene conversion did
not affect demographic inference. We did find that positive selection can affect the synonymous
SFS, resulting in inference of population size changes. In simulations of positive selection in a
population of constant size, we found the spurious inference to be a bottleneck rather than an
expansion. This suggests that the observed synonymous SFS of lineage U has been affected
both by positive selection and by demographic expansion.

237 Discussion.

A central question in the population biology of infectious diseases is how and why pathogenic
traits emerge in microbes. Addressing this question is important for understanding novel
disease emergence and for identifying the genetic basis of virulence. Here we present evidence
suggesting that a mutation in *S. saprophyticus*' *aas*, which binds host matrix proteins, is under
positive selection and has enabled emergence and spread of a human pathogenic, UTI-
associated lineage of this bacterium.

S. saprophyticus is familiar to medical microbiologists and clinicians as a common cause of
UTIs (46), which are associated with significant morbidity, economic costs, and severe
complications (4). Despite its strong association with UTIs in humans, *S. saprophyticus* can also
be isolated from diverse environments including livestock, food and food processing plants, and
the environment (47, 48). Our previous research suggested that pathogenicity to humans is a
derived trait in the species (7).

This pattern is replicated here, where phylogenetic analyses link human UTI with two lineages
of *S. saprophyticus* that are nested among isolates from diverse, non-human niches (i.e. free
living, food- and animal-associated) . The *aas* mutation arose in lineage U, which contains most
of the UTI isolates. Two lineages are basal to lineage U: one is bovine-associated, and the other
contains an ancient bacterial sequence from a pregnancy-related infection in Late Byzantine
Troy. The Troy bacterium has the ancestral, bovine-associated *aas* allele, and we have
previously hypothesized (7) that this lineage could be associated with human infections in
regions where humans have close contacts – e.g. shared living quarters– with livestock, as they
did at Troy during this time.

A second cluster of UTI isolates appears in Clade E. One isolate has acquired the derived *aas*
allele, which parallels our finding that two non-human isolates in lineage U acquired the
ancestral variant; all recombination events that we observed at this locus reinforced the
association between *aas_2206A>C* and human infection.

Several UTI isolates in Clade E do not have the derived *aas* allele and the clustering of UTI
isolates suggests there may be a distinct adaptive path to virulence in this clade. Larger and
more comprehensive samples will be needed to investigate this hypothesis and to identify the
factors shaping the separation of clades P & E.

The *aas* mutation has characteristics associated with a classical selective sweep driven by
positive selection, namely a regional reduction in diversity (21) and Tajima's D (22, 49). With the
exception of the interesting allelic replacements noted above, there was also relatively little
recombination at this locus, consistent with it being functionally important. To our knowledge,
this is the first description of a single nucleotide sweep in a bacterium.

Depending on the strength of selection and recombination rate, positive selection in bacteria
has been observed to affect the entire genome, resulting in clonal replacements, or to only
affect specific regions of the genome (50). For example, multiple clonal replacements have
occurred in *Shigella sonnei* populations in Vietnam due to acquisition of resistance to
antimicrobials and environmental stress (51). Recurrent clonal replacements have also been
observed within single hosts during chronic infection of cystic fibrosis patients by *Pseudomonas*
aeruginosa (52). Environmental bacterial populations can also be subject to clonal
replacements: a metagenomic time course study of Trout Bog found evidence of clonal
replacement occurring in natural bacterial populations but not gene or region specific sweeps
(53). However, large regions of low diversity were also observed, suggesting gene-specific
selective sweeps had occurred prior to the start of the study. Shapiro et al. identified genomic
loci that differentiated *Vibrio cyclitrophicus* associated with distinct niches but that had limited
diversity within niches; they concluded that differentiation of these populations had been
enabled by recombination events that reinforced the association of alleles with the niche in
which they were advantageous (54).

The *aas_2206A>C* mutation is within a group of genetic variants that differentiates bacteria
associated with human-pathogenic versus other niches (i.e. F_{ST} outlier). SNPs associated with
specific clinical phenotypes were described recently in the pathogen *Streptococcus pyogenes*
(55), which is consistent with our finding that clinical phenotypes can represent distinct niche

spaces preferentially occupied by sub-populations of bacteria. There is also precedent for a
single nucleotide polymorphism to affect host tropism of bacteria (56).

In sexually reproducing organisms, haplotype based statistics are frequently used to identify
selective sweeps because positively selected alleles will also increase the frequency of nearby
linked loci faster than recombination can disrupt linkage, producing longer haplotypes for
selected alleles (28, 57). We found that $aas\_2206A>C$ had a longer haplotype than the
ancestral variant, but this difference was not extreme relative to other regions of the genome
(assessed with the nS_L statistic). Haplotype based statistics have been found to perform poorly
in purebred dogs, where linkage across the genome is high (58). Relatively low levels of
recombination may also contribute to a lack of sensitivity when haplotype-based detection
methods are applied to bacteria; linkage of sites is also likely to be disrupted in a less
predictable way by bacterial gene conversion than by crossing over (29). Based on our findings,
we conclude that screening for regional decreases in diversity and distortions of the SFS (i.e.
sliding window analyses) and identification of genetic variants with extreme differences in
frequency between niches can be useful in identifying candidate sites of positive selection in
bacteria.

S. saprophyticus encodes a number of adhesins including UafA, UafB, SdrI, and Aas. UafA and
Aas are found in all isolates, suggesting that they play important roles in the diverse niches
occupied by *S. saprophyticus*. Aas has autolytic, fibronectin binding, and haemagluttinating
functions (9, 31, 32, 59). We identified a single, non-synonymous polymorphism as a target of
selection in the fibronectin binding repeats of *Aas*. This variant is predicted to affect the repeat's
structure, as proline has a more rigid structure than other amino acids. Adhesins are plausible
candidates for adaptation to the uropathogenic niche, as they are known to be important
virulence factors in pathogens causing urinary tract infections (60). Fibronectin binding proteins
including Aas have been identified as virulence factors in *S. saprophyticus* and *Enterococcus*
faecalis (32, 61, 62). Adhesion to the uroepithelium is essential for uropathogens to establish
themselves in the bladder, where they are subject to strong shear stress (63): we hypothesize
that *S. saprophyticus* with the derived *aas* variant are better able to colonize the human bladder.

Invasion of the human urinary tract may provide a fitness advantage by allowing relative
enrichment of *S. saprophyticus* in a site with little competition from other bacterial species and
by providing a mechanism of dispersal in the environment. In analyses of selection in
Escherichia coli, another bacterium occupying diverse niches, residues in the adhesin FimH
were found to be subject to positive selection in uropathogenic strains (64–66). FimH binds
mannose, providing protection from shear stress through a catch bond mechanism (67).
Interestingly, *Borrelia burgdorferi*'s vascular adherence and resistance to shear stress were
recently found to be enabled by interactions between a bacterial adhesin and host fibronectin
that also use a catch bond mechanism (68). There are also precedents in *Staphylococcus*
aureus for polymorphisms in bacterial fibronectin-binding adhesins to affect the strength of
binding, and for these polymorphisms to associate with specific clinical phenotypes (69).

Further experiments are needed to investigate the effects of variation in Aas on *S.*
saprophyticus biology. In our preliminary investigations of binding using ELISA assays of

recombinant bacterial peptides, we did not detect differences between ancestral and derived
alleles in binding of the R1a1b repeat to fibronectin. The variant could still affect fibronectin
binding by altering conformation of the protein in a manner analogous to FimH in *E. coli* (66). It's
also possible that variants in the peptide affect binding under specific conditions that we did not
test. Another possibility is that the variant affects autolysis or other as yet undescribed functions
of Aas. The roles of adhesins and other virulence factors in *S. saprophyticus*'s colonization of
niches in livestock and the environment are also interesting topics for further study.

Our demographic analysis of the uropathogenic lineage of *S. saprophyticus* showed evidence of
a population bottleneck and subsequent expansion. Bottlenecks and expansion of drug resistant
clones have previously been shown to affect the population structure of *Streptococcus*
agalactiae (70), demonstrating the effects of positive selection on the demographic trajectories
of bacterial sub-populations. However, previous work has also shown that selection - and
recombination - can produce spurious results from demographic inference in bacteria (40, 71).
We used an SFS-based method to reconstruct the demographic history of *S. saprophyticus*: the
accuracy of demographic inference using these methods has been shown to be unaffected by
within-population recombination (40) and this was confirmed in our analyses of simulated data.
We found that recombination from external sources may result in an excess of intermediate
frequency variants, which is also a signature of population bottlenecks, so we masked externally
imported sites. However, the frequency of synonymous variants could still be affected by
selection on linked non-synonymous sites, including the selective sweep in *aas* that we have
described. We performed simulations to address these potential confounders and aid in the
interpretation of our demographic inferences. Simulation of a single site under positive selection
resulted in the inference of a bottleneck (N_e/N_a = 0.01-0.42), indicating that, at the recombination
rates we simulated, diversity was lost from neutrally evolving sites due to their linkage to the site
under selection. In inferences from our observed data, a bottleneck was followed by a 15-fold
expansion, suggesting that lineage U has undergone both a selective sweep and demographic
expansion.

Here we have described adaptation of *S. saprophyticus* that may have enabled its expansion
into a human pathogenic niche. Mutation of a single nucleotide within the *aas* adhesin appears
to have driven a selective sweep, and allele frequency differences at the locus are consistent
with niche-specific adaptation. Lateral gene transfer events in *aas* reinforced the association of
the positively selected allele with human infection. These results provide new insights into the
emergence of virulence in bacteria and outline an approach for discovering the molecular basis
of adaptation to the human pathogenic niche.

Methods.

DNA extraction. After overnight growth in TSB at 37°C in a shaking incubator, cultures were
pelleted and resuspended in 140 µL TE buffer. Cells were incubated overnight with 50 units of
mutanolysin. We used the MasterPure Gram Positive DNA Purification Kit (EpiCentre) for DNA

extraction.For DNA precipitation we used 1 mL 70% ethanol and centrigured at 4°C for 10
minutes. We additionally used a SpeedVac for 10 minutes to ensure pellets were dry before re-
suspending the pellet in 50 µL water.

Library preparation and sequencing. For SSC01, SSC02, and SSC03, library prep was
performed using a modified Nextera protocol as decribed by Baym et al. (72) with a
reconditioning PCR with fresh primers and polymerase for an additional 5 PCR cycles to
minimize chimeras and a two-step bead based size selection with target fragment size of 650 bp
and sequenced on an Illumina HiSeq 2500 (paired-end, 150 bp). For 43, SSC04, SCC05, and
SSMast, DNA was submitted to the University of Wisconsin-Madison Biotechnology Center for
library preparation and were prepared according the TruSeq Nano DNA LT Library Prep Kit
(Illumina Inc., San Diego, California, USA) with minor modifications. A maximum of 200 ng of
each sample was sheared using a Covaris M220 Ultrasonicator (Covaris Inc, Woburn, MA,
USA). Sheared samples were size selected for an average insert size of 550 bp using Spri bead
based size exclusion. Quality and quantity of the finished libraries were assessed using an
Agilent DNA High Sensitivity chip (Agilent Technologies, Santa Clara, CA) and Qubit dsDNA HS
Assay Kit, respectively. Libraries were standardized to 2 µM. Paired-end, 150 bp sequencing
was performed using v2 SBS chemistry on an Illumina MiSeq sequencer. Images were
analyzed using the Illumina Pipeline, version 1.8.2.

Reference guided mapping. We mapped reads to ATCC 15305 using via a pipeline (available at
https://github.com/pepperell-lab/RGAPepPipe). Briefly, read quality was assessed and trimmed
with TrimGalore! v 0.4.0 (www.bioinformatics.babraham.ac.uk/projects/trim_galore), which runs
both FastQC (www.bioinformatics.babraham.ac.uk/projects/fastqc) and cutadapt. Reads were
mapped to using BWA-MEM v 0.7.12 (73) and sorted using Samtools v 1.2 (74). We used
Picard v 1.138 (picard.sourceforge.net) to add read group information and removed duplicates.
Reads were locally realigned using GATK v 2.8.1 (75). We identified variants using Pilon v 1.16
(76) (minimum read depth: 10, minimum mapping quality: 40, minimum base quality: 20).

Assembly. We used the iMetAMOS pipeline for *de novo* assembly (77). We chose to compare
assemblies from SPAdes (78), MaSurCA (79), and Velvet (80). KmerGenie (81) was used to
select kmer sizes for assembly. iMetAMOS uses FastQC, QUAST (82), REAPR (83), LAP (84),
ALE (85), FreeBayes (86), and CGAL (87) to evaluate the quality of reads and assemblies. We
also used Kraken (88) to detect potential contamination in sequence data. For all newly
assembled isolates (43, SSC01-05, SSMast), the SPAdes assembly was the highest quality.
Assembly statistics are reported in Supplementary Table 3.

Annotation and gene content analyses. We annotated the de novo assemblies using Prokka v
1.11 (89) and used Roary (90) to identify orthologous genes in the core and accessory
genomes. To look for associations between accessory gene content and human association, we
used Scoary (19). For the analysis, we used human association as our trait, and we adjusted
the p-value for multiple comparisons using the Bonferroni method.

Alignment. When short read data for reference guided mapping were unavailable, whole
genome alignment of genomes to ATCC 15305 was performed using Mugsy v 2.3 (91).

Repetitive regions in the reference genome greater than 100 bp were identified using nucmer,
and these regions were masked in the alignment used in downstream analyses.

Maximum likelihood phylogenetic analysis. Maximum likelihood phylogenetic trees were inferred
using RAxML 8.0.6 (92). We used the GTRGAMMA substitution model and performed
bootstrapping using the autoMR convergence criteria. Tree visualizations were created in ggtree
(93).

Population genetics statistics. To calculate π and Tajima's D, we used EggLib v 2.1.10 (94), a
Python package for population genetic analyses. A script to perform the sliding window analysis
is available at https://github.com/tatumdmortimer/popgen-stats/slidingWindowStats.py. We used
vcflib (https://github.com/vcflib/vcflib) to calculate F_{ST} and EHH and selscan v 1.1.0b (95) to
calculate nS_L.

Recombination analyses. To identify recombinant regions in the *S. saprophyticus* alignment, we
used Gubbins v 2.1.0 (20). fastGEAR (42) was used with the recommended input specifications
to identify recombination events between major lineages of *S. saprophyticus*. We used Circos
(96) for visualization of recombinant tracts.

Site frequency spectrum. We used SNP-sites v 2.0.3 (97) to convert the alignment of *S.*
saprophyticus isolates to a multi-sample VCF. SnpEff v 4.1j (98) was used to annotate variants
in this VCF as synonymous, non-synonymous, or intergenic. Using the Troy genome as an
outgroup, we calculated an unfolded site frequency spectrum (SFS) for lineage U for each
category of sites. To reduce the impact of lateral gene transfer on the SFS, we removed sites
where the origin was outside lineage U based on results of fastGEAR.

Ancestral reconstruction of aas_2206A>C. We used TreeTime (99) to reconstruct the
evolutionary history of the variant in *aas* using the maximum likelihood phylogeny inferred using
RAxML.

Demography. We performed demographic inference with the synonymous SFS using ∂a∂i (39).
Models tested were the standard neutral model, expansion, and exponential growth.
Parameters, *v* (N_e/ N_a) and *τ* (time scaled by 2), were optimized for both the expansion and
exponential growth models. Significance of the expansion and exponential growth model
compared to the standard neutral model was evaluated using a likelihood ratio test. The scripts
used to perform this analysis are available at https://github.com/tatumdmortimer/popgen-stats.

Simulations. Simulations were performed in SimBac (41) to evaluate the effect of external
recombination on the SFS. Populations were simulated with sample size and θ equivalent to our
sample of lineage U (*n* = 44, θ = 0.003). The length of internal recombinant tracts was 6500 bp
(median of Gubbins output), and the length of external recombination events was 3000 bp
(median of fastGEAR output). Internal recombination was simulated at rates ranging from 0 to
0.03 (r/m = 10). External recombination was simulated at rates ranging from 0 to 0.003. The
lower bound of difference for external recombination was 0, and the upper bound was simulated
at ranges from 0.25-1.0. Simulations were performed in SFS_CODE (release date 9/10/2015)
(44) to evaluate the power of ∂a∂i to accurately estimate demographic parameters in the

presence of gene conversion and selection. We simulated a locus of length 100 kb with theta
0.003, gene conversion tract length of 1345 bp, and a range of recombination/mutation ratios
(0.0002-2.0). In addition to neutral simulations with gene conversion, we also performed
simulations with a single site under selection (γ = 10-1000) with the same parameters as neutral
simulations.

Expression of R1ab. Human-associated and ancestral strain R1ab were cloned into the
expression vector pET-ELMER (LM Maurer 2010, JBC), transformed into BL21(DE3)cells
(EMD, Gibbstown, NJ) for expression and induced with 1mM IPTG. Bacteria were lysed in 100
mM NaH_2PO_4, 10 mM Tris, 8M urea, 1 mM β-mercapto-ethanol, 5 mM imidazole, pH 8.0 (lysis
buffer). The cleared lysate was incubated overnight with nickel-nitrilotriacetic acid agarose
(Qiagen), washed and eluted in lysis buffer pH 7.0 plus 300 mM imidazole.

ELISA. Antigen was diluted to 10 µg/ml in TBS (10 mM Tris, 150 mM, pH 7.4) and used to coat
96-well microtiter plates (Costar 3590 high binding, Corning Inc., Corning, NY) with 50 µl per
well, for 16 h at 4□C. The plates were blocked with 1% BSA in TBS plus 0.05% Tween-20
(TBST) for 1 h. After washing three times with TBST, purified plasma fibronectin (100) or
platelet-derived thrombospondin-1 (101) diluted to 10, 3, 1, or 0.3µg/ml in TBST plus 0.1% BSA,
were added to the plates and incubated for 2 h. Plates were washed four times with TBST.
Rabbit anti-fibronectin and rabbit anti-thrombospondin antibodies diluted in TBST plus 0.1%
BSA were added to the appropriate wells and incubated for 1 h. Plates were washed four times
with TBST. Peroxidase-conjugated secondary antibody was incubated with the plates for 1 h.
Plates were washed four times with TBST and 50 µl per well of SureBlue TMB peroxidase
substrate (KLP) was added to each well. Color development was monitored for 10-30 min, 50 µl
of TMB stop solution (KLP) was added, followed by measurement of absorbance at 450nm.

Funding Information.

This material is based upon work supported by the National Science Foundation Graduate
Research Fellowship Program under Grant No. DGE-1256259 to TDM and MBO. Any opinions,
findings, and conclusions or recommendations expressed in this material are those of the
author(s) and do not necessarily reflect the views of the National Science Foundation. TDM is
also supported by National Institutes of Health National Research Service Award (T32
GM07215). CSP is supported by National Institutes of Health (R01AI113287).

Acknowledgments.

We thank Andrew Kitchen (University of Iowa), Jeniel Nett (UW-Madison), JD Sauer (UW-
Madison), and Rod Welch (UW-Madison) for their helpful input on this study. We also thank the
University of Wisconsin Biotechnology Center DNA Sequencing Facility for providing library
preparation and sequencing facilities and services.

References.

1. Stamm WE, Norrby SR. 2001. Urinary Tract Infections: Disease Panorama and
Challenges. J Infect Dis 183:S1–S4.

2. Ronald AR, Nicolle LE, Stamm E, Krieger J, Warren J, Schaeffer A, Naber KG, Hooton
TM, Johnson J, Chambers S, Andriole V. 2001. Urinary tract infection in adults: research
priorities and strategies. Int J Antimicrob Agents 17:343–348.

3. Barber AE, Norton JP, Spivak AM, Mulvey MA. 2013. Urinary Tract Infections: Current and
Emerging Management Strategies. Clin Infect Dis 57:719–724.

4. Foxman B. 2014. Urinary tract infection syndromes: occurrence, recurrence, bacteriology,
risk factors, and disease burden. Infect Dis Clin North Am 28:1–13.

5. Wallmark G, Arremark I, Telander B. 1978. Staphylococcus saprophyticus: A Frequent
Cause of Acute Urinary Tract Infection among Female Outpatients. J Infect Dis 138:791–
797.

6. Kahlmeter G. 2003. An international survey of the antimicrobial susceptibility of pathogens
from uncomplicated urinary tract infections: the ECO·SENS Project. J Antimicrob
Chemother 51:69–76.

7. Devault AM, Mortimer TD, Kitchen A, Kiesewetter H, Enk JM, Golding GB, Southon J,
Kuch M, Duggan AT, Aylward W, Gardner SN, Allen JE, King AM, Wright G, Kuroda M,
Kato K, Briggs DE, Fornaciari G, Holmes EC, Poinar HN, Pepperell CS. 2017. A molecular
portrait of maternal sepsis from Byzantine Troy. eLife 6:e20983.

8. Gatermann S, John J, Marre R. 1989. Staphylococcus saprophyticus urease:

characterization and contribution to uropathogenicity in unobstructed urinary tract infection

of rats. Infect Immun 57:110–116.

9. Meyer HG, Wengler-Becker U, Gatermann SG. 1996. The hemagglutinin of

Staphylococcus saprophyticus is a major adhesin for uroepithelial cells. Infect Immun

64:3893–3896.

10. Kuroda M, Yamashita A, Hirakawa H, Kumano M, Morikawa K, Higashide M, Maruyama A,

Inose Y, Matoba K, Toh H, Kuhara S, Hattori M, Ohta T. 2005. Whole genome sequence of

Staphylococcus saprophyticus reveals the pathogenesis of uncomplicated urinary tract

infection. Proc Natl Acad Sci U S A 102:13272–13277.

11. Kleine B, Gatermann S, Sakinc T. 2010. Genotypic and phenotypic variation among

Staphylococcus saprophyticus from human and animal isolates. BMC Res Notes 3:163.

12. Nelson EJ, Harris JB, Glenn Morris J, Calderwood SB, Camilli A. 2009. Cholera

transmission: the host, pathogen and bacteriophage dynamic. Nat Rev Microbiol 7:693–

702.

13. Boucher Y, Orata FD, Alam M. 2015. The out-of-the-delta hypothesis: dense human

populations in low-lying river deltas served as agents for the evolution of a deadly

pathogen. Front Microbiol 6.

14. Veyrier FJ, Dufort A, Behr MA. 2011. The rise and fall of the Mycobacterium tuberculosis

genome. Trends Microbiol 19:156–161.

15. McNally A, Thomson NR, Reuter S, Wren BW. 2016. "Add, stir and reduce": Yersinia spp.

as model bacteria for pathogen evolution. Nat Rev Microbiol 14:177–190.

16.   Larsson P, Elfsmark D, Svensson K, Wikström P, Forsman M, Brettin T, Keim P,

531   Johansson A. 2009. Molecular Evolutionary Consequences of Niche Restriction in

532   Francisella tularensis, a Facultative Intracellular Pathogen. PLoS Pathog 5:e1000472.

533   17.   Lebreton F, Schaik W van, McGuire AM, Godfrey P, Griggs A, Mazumdar V, Corander J,

534   Cheng L, Saif S, Young S, Zeng Q, Wortman J, Birren B, Willems RJL, Earl AM, Gilmore

535   MS. 2013. Emergence of Epidemic Multidrug-Resistant Enterococcus faecium from Animal

536   and Commensal Strains. mBio 4:e00534-13.

537   18.   Fitzgerald JR, Nutbeam-Tuffs S, Richardson E, Lee CY, Gupta RK, Richards AC, Black

538   NS, Corander J, McAdam PR, Spoor LE, O'Gara JP, Mendonca C, Wilson GJ. 2015.

539   Recombination-mediated remodelling of host-pathogen interactions during Staphylococcus

540   aureus niche adaptation. Microb Genomics.

541   19.   Brynildsrud O, Bohlin J, Scheffer L, Eldholm V. 2016. Rapid scoring of genes in microbial

542   pan-genome-wide association studies with Scoary. Genome Biol 17:238.

543   20.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris

544   SR. 2015. Rapid phylogenetic analysis of large samples of recombinant bacterial whole

545   genome sequences using Gubbins. Nucleic Acids Res 43:e15–e15.

546   21.   Smith JM, Haigh J. 1974. The hitch-hiking effect of a favourable gene. Genet Res.

547   22.   Braverman JM, Hudson RR, Kaplan NL, Langley CH, Stephan W. 1995. The hitchhiking

548   effect on the site frequency spectrum of DNA polymorphisms. Genetics 140:783–796.

549   23.   Harr B, Kauer M, Schlötterer C. 2002. Hitchhiking mapping: A population-based fine-

550   mapping strategy for adaptive mutations in Drosophila melanogaster. Proc Natl Acad Sci

551   99:12949–12954.

24. Nair S, Williams JT, Brockman A, Paiphun L, Mayxay M, Newton PN, Guthmann J-P,

Smithuis FM, Hien TT, White NJ, Nosten F, Anderson TJC. 2003. A Selective Sweep

Driven by Pyrimethamine Treatment in Southeast Asian Malaria Parasites. Mol Biol Evol

20:1526–1536.

25. Wright SI, Gaut BS. 2005. Molecular Population Genetics and the Search for Adaptive

Evolution in Plants. Mol Biol Evol 22:506–519.

26. de Simoni Gouveia JJ, da Silva MVGB, Paiva SR, de Oliveira SMP. 2014. Identification of

selection signatures in livestock species. Genet Mol Biol 37:330–342.

27. Weir BS, Cockerham CC. 1984. Estimating F-Statistics for the Analysis of Population

Structure. Evolution 38:1358–1370.

28. Sabeti PC, Reich DE, Higgins JM, Levine HZP, Richter DJ, Schaffner SF, Gabriel SB,

Platko JV, Patterson NJ, McDonald GJ, Ackerman HC, Campbell SJ, Altshuler D, Cooper

R, Kwiatkowski D, Ward R, Lander ES. 2002. Detecting recent positive selection in the

human genome from haplotype structure. Nature 419:832–837.

29. Shapiro BJ. 2014. Signatures of Natural Selection and Ecological Differentiation in

Microbial Genomes, p. 339–359. *In* Landry, CR, Aubin-Horth, N (eds.), Ecological

Genomics. Springer Netherlands.

30. Ferrer-Admetlla A, Liang M, Korneliussen T, Nielsen R. 2014. On Detecting Incomplete

Soft or Hard Selective Sweeps Using Haplotype Structure. Mol Biol Evol 31:1275–1291.

31. Hell W, Meyer H-GW, Gatermann SG. 1998. Cloning of aas, a gene encoding a

Staphylococcus saprophyticus surface protein with adhesive and autolytic properties. Mol

Microbiol 29:871–881.

32. Gatermann S, Meyer HG. 1994. Staphylococcus saprophyticus hemagglutinin binds

fibronectin. Infect Immun 62:4556–4563.

33. Kohler TP, Gisch N, Binsker U, Schlag M, Darm K, Völker U, Zähringer U, Hammerschmidt

S. 2014. Repeating Structures of the Major Staphylococcal Autolysin Are Essential for the

Interaction with Human Thrombospondin 1 and Vitronectin. J Biol Chem 289:4070–4082.

34. Zoll S, Schlag M, Shkumatov AV, Rautenberg M, Svergun DI, Götz F, Stehle T. 2012.

Ligand-Binding Properties and Conformational Dynamics of Autolysin Repeat Domains in

Staphylococcal Cell Wall Recognition. J Bacteriol 194:3789–3802.

35. He M, Sebaihia M, Lawley TD, Stabler RA, Dawson LF, Martin MJ, Holt KE, Seth-Smith

HMB, Quail MA, Rance R, Brooks K, Churcher C, Harris D, Bentley SD, Burrows C, Clark

L, Corton C, Murray V, Rose G, Thurston S, Tonder A van, Walker D, Wren BW, Dougan

G, Parkhill J. 2010. Evolutionary dynamics of Clostridium difficile over short and long time

scales. Proc Natl Acad Sci 107:7527–7532.

36. Nübel U, Dordel J, Kurt K, Strommenger B, Westh H, Shukla SK, Žemličková H, Leblois R,

Wirth T, Jombart T, Balloux F, Witte W. 2010. A Timescale for Evolution, Population

Expansion, and Spatial Spread of an Emerging Clone of Methicillin-Resistant

Staphylococcus aureus. PLOS Pathog 6:e1000855.

37. Pepperell CS, Casto AM, Kitchen A, Granka JM, Cornejo OE, Holmes EC, Birren B,

Galagan J, Feldman MW. 2013. The Role of Selection in Shaping Diversity of Natural M.

tuberculosis Populations. PLoS Pathog 9:e1003543.

38. Davies MR, Holden MT, Coupland P, Chen JHK, Venturini C, Barnett TC, Zakour NLB, Tse

H, Dougan G, Yuen K-Y, Walker MJ. 2015. Emergence of scarlet fever Streptococcus

pyogenes emm12 clones in Hong Kong is associated with toxin acquisition and multidrug

resistance. Nat Genet 47:84–87.

39. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. 2009. Inferring the Joint

Demographic History of Multiple Populations from Multidimensional SNP Frequency Data.

PLoS Genet 5:e1000695.

40. Lapierre M, Blin C, Lambert A, Achaz G, Rocha EPC. 2016. The impact of selection, gene

conversion, and biased sampling on the assessment of microbial demography. Mol Biol

Evol msw048.

41. Brown T, Didelot X, Wilson DJ, De Maio N. 2016. SimBac: simulation of whole bacterial

genomes with homologous recombination. Microb Genomics 2.

42. Mostowy R, Croucher NJ, Andam CP, Corander J, Hanage WP, Marttinen P. 2016.

Analysis of recent and ancestral recombination reveals high-resolution population structure

in Streptococcus pneumoniae. bioRxiv 059642.

43. Coffman AJ, Hsieh PH, Gravel S, Gutenkunst RN. 2016. Computationally Efficient

Composite Likelihood Statistics for Demographic Inference. Mol Biol Evol 33:591–593.

44. Hernandez RD. 2008. A flexible forward simulator for populations subject to selection and

demography. Bioinformatics 24:2786–2787.

45. Neher RA, Hallatschek O. 2013. Genealogies of rapidly adapting populations. Proc Natl

Acad Sci 110:437–442.

46. Widerstrom M, Wistrom J, Ferry S, Karlsson C, Monsen T. 2007. Molecular Epidemiology

of Staphylococcus saprophyticus Isolated from Women with Uncomplicated Community-

Acquired Urinary Tract Infection. J Clin Microbiol 45:1561–1564.

47.   Hedman P, Ringertz O, Lindström M, Olsson K. 1993. The origin of Staphylococcus

saprophyticus from cattle and pigs. Scand J Infect Dis 25:57–60.

48.   Kastman EK, Kamelamela N, Norville JW, Cosetta CM, Dutton RJ, Wolfe BE. 2016. Biotic

Interactions Shape the Ecological Distributions of Staphylococcus Species. mBio

7:e01157-16.

49.   Coton E, Desmonts M-H, Leroy S, Coton M, Jamet E, Christieans S, Donnio P-Y, Lebert I,

Talon R. 2010. Biodiversity of Coagulase-Negative Staphylococci in French cheeses, dry

fermented sausages, processing environments and clinical samples. Int J Food Microbiol

137:221–229.

50.   Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA

polymorphism. Genetics 123:585–595.

51.   Shapiro BJ, David LA, Friedman J, Alm EJ. 2009. Looking for Darwin's footprints in the

microbial world. Trends Microbiol 17:196–204.

52.   Holt KE, Nga TVT, Thanh DP, Vinh H, Kim DW, Tra MPV, Campbell JI, Hoang NVM, Vinh

NT, Minh PV, Thuy CT, Nga TTT, Thompson C, Dung TTN, Nhu NTK, Vinh PV, Tuyet

PTN, Phuc HL, Lien NTN, Phu BD, Ai NTT, Tien NM, Dong N, Parry CM, Hien TT, Farrar

JJ, Parkhill J, Dougan G, Thomson NR, Baker S. 2013. Tracking the establishment of local

endemic populations of an emergent enteric pathogen. Proc Natl Acad Sci 110:17522–

17527.

53.   Caballero JD, Clark ST, Coburn B, Zhang Y, Wang PW, Donaldson SL, Tullis DE, Yau

YCW, Waters VJ, Hwang DM, Guttman DS. 2015. Selective Sweeps and Parallel

Pathoadaptation Drive Pseudomonas aeruginosa Evolution in the Cystic Fibrosis Lung.

mBio 6:e00981-15.

54. Bendall ML, Stevens SL, Chan L-K, Malfatti S, Schwientek P, Tremblay J, Schackwitz W,

Martin J, Pati A, Bushnell B, Froula J, Kang D, Tringe SG, Bertilsson S, Moran MA, Shade

A, Newton RJ, McMahon KD, Malmstrom RR. 2016. Genome-wide selective sweeps and

gene-specific sweeps in natural bacterial populations. ISME J 10:1589–1601.

55. Shapiro BJ, Friedman J, Cordero OX, Preheim SP, Timberlake SC, Szabó G, Polz MF,

Alm EJ. 2012. Population Genomics of Early Events in the Ecological Differentiation of

Bacteria. Science 336:48–51.

56. Bao Y-J, Shapiro BJ, Lee SW, Ploplis VA, Castellino FJ. 2016. Phenotypic differentiation of

Streptococcus pyogenes populations is induced by recombination-driven gene-specific

sweeps. Sci Rep 6:36644.

57. Viana D, Comos M, McAdam PR, Ward MJ, Selva L, Guinane CM, González-Muñoz BM,

Tristan A, Foster SJ, Fitzgerald JR, Penadés JR. 2015. A single natural nucleotide

mutation alters bacterial pathogen host tropism. Nat Genet advance online publication.

58. Vitti JJ, Grossman SR, Sabeti PC. 2013. Detecting Natural Selection in Genomic Data.

Annu Rev Genet 47:97–120.

59. Schlamp F, van der Made J, Stambler R, Chesebrough L, Boyko AR, Messer PW. 2016.

Evaluating the performance of selection scans to detect selective sweeps in domestic

dogs. Mol Ecol 25:342–356.

60. Meyer H-GW, Müthing J, Gatermann SG. 1997. The hemagglutinin of Staphylococcus

saprophyticus binds to a protein receptor on sheep erythrocytes. Med Microbiol Immunol

(Berl) 186:37–43.

61.   Flores-Mireles AL, Walker JN, Caparon M, Hultgren SJ. 2015. Urinary tract infections:

epidemiology, mechanisms of infection and treatment options. Nat Rev Microbiol advance

online publication.

62.   King NP, Beatson SA, Totsika M, Ulett GC, Alm RA, Manning PA, Schembri MA. 2011.

UafB is a serine-rich repeat adhesin of Staphylococcus saprophyticus that mediates

binding to fibronectin, fibrinogen and human uroepithelial cells. Microbiology 157:1161–

1175.

63.   Torelli R, Serror P, Bugli F, Sterbini FP, Florio AR, Stringaro A, Colone M, Carolis ED,

Martini C, Giard J-C, Sanguinetti M, Posteraro B. 2012. The PavA-like Fibronectin-Binding

Protein of Enterococcus faecalis, EfbA, Is Important for Virulence in a Mouse Model of

Ascending Urinary Tract Infection. J Infect Dis 206:952–960.

64.   Wright KJ, Hultgren SJ. 2006. Sticky fibers and uropathogenesis: bacterial adhesins in the

urinary tract. Future Microbiol 1:75–87.

65.   Chen SL, Hung C-S, Xu J, Reigstad CS, Magrini V, Sabo A, Blasiar D, Bieri T, Meyer RR,

Ozersky P, Armstrong JR, Fulton RS, Latreille JP, Spieth J, Hooton TM, Mardis ER,

Hultgren SJ, Gordon JI. 2006. Identification of genes subject to positive selection in

uropathogenic strains of Escherichia coli: A comparative genomics approach. Proc Natl

Acad Sci 103:5977–5982.

66.   Chen SL, Hung CS, Pinkner JS, Walker JN, Cusumano CK, Li Z, Bouckaert J, Gordon JI,

Hultgren SJ. 2009. Positive selection identifies an in vivo role for FimH during urinary tract

infection in addition to mannose binding. Proc Natl Acad Sci 106:22439–22444.

67.   Schwartz DJ, Kalas V, Pinkner JS, Chen SL, Spaulding CN, Dodson KW, Hultgren SJ.

2013. Positively selected FimH residues enhance virulence during urinary tract infection by

altering FimH conformation. Proc Natl Acad Sci 110:15530–15537.

68.   Yakovenko O, Sharma S, Forero M, Tchesnokova V, Aprikian P, Kidd B, Mach A, Vogel V,

Sokurenko E, Thomas WE. 2008. FimH Forms Catch Bonds That Are Enhanced by

Mechanical Force Due to Allosteric Regulation. J Biol Chem 283:11596–11605.

69.   Niddam AF, Ebady R, Bansal A, Koehler A, Hinz B, Moriarty TJ. 2017. Plasma fibronectin

stabilizes Borrelia burgdorferi–endothelial interactions under vascular shear stress by a

catch-bond mechanism. Proc Natl Acad Sci 114:E3490–E3498.

70.   Messina JA, Thaden JT, Sharma-Kuinkel BK, Jr VGF. 2016. Impact of Bacterial and

Human Genetic Variation on Staphylococcus aureus Infections. PLOS Pathog

12:e1005330.

71.   Da Cunha V, Davies MR, Douarre P-E, Rosinski-Chupin I, Margarit I, Spinali S, Perkins T,

Lechat P, Dmytruk N, Sauvage E, Ma L, Romi B, Tichit M, Lopez-Sanchez M-J, Descorps-

Declere S, Souche E, Buchrieser C, Trieu-Cuot P, Moszer I, Clermont D, Maione D,

Bouchier C, McMillan DJ, Parkhill J, Telford JL, Dougan G, Walker MJ, Devani

Consortium, Holden MTG, Poyart C, Glaser P. 2014. Streptococcus agalactiae clones

infecting humans were selected and fixed through the extensive use of tetracycline. Nat

Commun 5.

72.   Hedge J, Wilson DJ. 2014. Bacterial Phylogenetic Reconstruction from Whole Genomes Is

Robust to Recombination but Demographic Inference Is Not. mBio 5:e02158-14.

73. Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. 2015.

Inexpensive Multiplexed Library Preparation for Megabase-Sized Genomes. PLOS ONE

10:e0128036.

74. Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. 1303.3997. arXiv e-print.

75. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin

R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25:2078–

2079.

76. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del

Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kernytsky AM, Sivachenko AY,

Cibulskis K, Gabriel SB, Altshuler D, Daly MJ. 2011. A framework for variation discovery

and genotyping using next-generation DNA sequencing data. Nat Genet 43:491–498.

77. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,

Wortman J, Young SK, Earl AM. 2014. Pilon: An Integrated Tool for Comprehensive

Microbial Variant Detection and Genome Assembly Improvement. PLOS ONE 9:e112963.

78. Koren S, Treangen TJ, Hill CM, Pop M, Phillippy AM. 2014. Automated ensemble

assembly and validation of microbial genomes. BMC Bioinformatics 15:126.

79. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,

Alekseyev MA, Pevzner PA. 2012. SPAdes: A New Genome Assembly Algorithm and Its

Applications to Single-Cell Sequencing. J Comput Biol 19:455–477.

80.   Zimin A, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA

genome Assembler. Bioinformatics btt476.

81.   Zerbino DR, Birney E. 2008. Velvet: Algorithms for de novo short read assembly using de

Bruijn graphs. Genome Res 18:821–829.

82.   Chikhi R, Medvedev P. 2014. Informed and automated k-mer size selection for genome

assembly. Bioinformatics 30:31–37.

83.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for

genome assemblies. Bioinforma Oxf Engl 29:1072–1075.

84.   Hunt M, Kikuchi T, Sanders M, Newbold C, Berriman M, Otto TD. 2013. REAPR: a

universal tool for genome assembly evaluation. Genome Biol 14:R47.

85.   Ghodsi M, Hill CM, Astrovskaya I, Lin H, Sommer DD, Koren S, Pop M. 2013. De novo

likelihood-based measures for comparing genome assemblies. BMC Res Notes 6:334.

86.   Clark SC, Egan R, Frazier PI, Wang Z. 2013. ALE: a generic assembly likelihood

evaluation framework for assessing the accuracy of genome and metagenome assemblies.

Bioinformatics 29:435–443.

87.   Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing.

ArXiv12073907 Q-Bio.

88.   Rahman A, Pachter L. 2013. CGAL: computing genome assembly likelihoods. Genome

Biol 14:R8.

89.   Wood DE, Salzberg SL. 2014. Kraken: ultrafast metagenomic sequence classification

using exact alignments. Genome Biol 15:R46.

90.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics btu153.

91.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, Fookes M, Falush D,

748   Keane JA, Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis.

749   Bioinformatics 31:3691–3693.

750   92.   Angiuoli SV, Salzberg SL. 2011. Mugsy: fast multiple alignment of closely related whole

751   genomes. Bioinformatics 27:334–342.

752   93.   Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

753   large phylogenies. Bioinformatics 30:1312–1313.

754   94.   Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2016. ggtree: an R package for visualization

755   and annotation of phylogenetic trees with their covariates and other associated data.

756   Methods Ecol Evol n/a-n/a.

757   95.   Mita SD, Siol M. 2012. EggLib: processing, analysis and simulation tools for population

758   genetics and genomics. BMC Genet 13:27.

759   96.   Szpiech ZA, Hernandez RD. 2014. selscan: An Efficient Multithreaded Program to Perform

760   EHH-Based Scans for Positive Selection. Mol Biol Evol 31:2824–2827.

761   97.   Krzywinski M, Schein J, Birol İ, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA.

762   2009. Circos: An information aesthetic for comparative genomics. Genome Res 19:1639–

763   1645.

764   98.   Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, Harris SR. 2016. SNP-

765   sites: rapid efficient extraction of SNPs from multi-FASTA alignments. biorxiv;038190v1.

99.   Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM.

2012. A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff. Fly (Austin) 6:80–92.

100. Sagulenko P, Puller V, Neher R. 2017. TreeTime: maximum likelihood phylodynamic

analysis. bioRxiv 153494.

101. Maurer LM, Tomasini-Johansson BR, Mosher DF. 2010. Emerging roles of fibronectin in

thrombosis. Thromb Res 125:287–291.

102. Annis DS, Murphy-Ullrich JE, Mosher DF. 2006. Function-blocking antithrombospondin-1

monoclonal antibodies. J Thromb Haemost 4:459–468.

Table 1. Results of demographic inference. $v = N_e/N_{ancestral}$, $\tau = $ generations$/ N_{ancestral}$

Model	Optimized parameters (standard deviation)	Log Likelihood	p-value (comparison model)
constant size		-562	
instantaneous change	v: 9.9×10^4 (6.0×10^4) τ: 4.4×10^{-2} (4.7×10^{-3})	-455	0.0 (constant size)
exponential change	v: 9.7×10^4 (9.8×10^4) τ: 4.1×10^{-2} (4.9×10^{-3})	-455	0.0 (constant size)
instantaneous change followed by exponential change	v_A: 2.1×10^{-2} (2.9×10^{-2}) v_B: 1.1 (3.9×10^{-2}) τ: 1.3×10^{-1} (1.6×10^{-1})	-404	1.6×10^{-4} (exponential change)
two instantaneous size changes	v_A: 2.9×10^{-2} (1.2×10^{-2}) v_B: 4.5×10^{-1} (2.1×10^{-1}) τ_A: 1.2×10^{-1} (3.4×10^{-2}) τ_B: 3.1×10^{-3} (5.3×10^{-3})	-393	4.6×10^{-7} (instantaneous change)

Figures.

Figure 1. Maximum likelihood phylogeny of *S. saprophyticus*. Maximum likelihood
phylogenetic analysis was performed in RAxML (92) using a whole genome alignment with
repetitive regions masked. The phylogeny is midpoint rooted, and nodes with bootstrap values
less than 90 are labelled. Branch lengths are scaled by substitutions per site. Tips are colored
based on the isolation source (pink- human, blue- animal, green- food, orange- environment).
Tips are labeled with isolate name and detailed source information. *S. saprophyticus* contains
two major clades (Clade P and Clade E). Within Clade P, there is a lineage enriched in human
pathogenic isolates (lineage U, branch labeled 'U').

Figure 2. Sliding window analysis of diversity and neutrality statistics. Population genetic
statistics were calculated for lineage U using EggLib (94). Windows were 50 kb in width with a
step size of 10 kb. A) Tajima's D. B) π (green) and θ (blue). The lowest values for Tajima's D
and π are found in the same window (1,760,000-1,820,000 bp, arrow).

Figure 3. Recombination in *S. saprophyticus*. Recombinant regions in the whole genome
alignment of *S. saprophyticus* were identified using Gubbins (20). Mobile genetic elements are
highlighted in blue on the outer rim. The window with low Tajima's D and π is highlighted in pink.
Few recombination events are inferred within this region.

Figure 4. Non-synonymous variant in Aas fibronectin binding repeat. Top- Domains of Aas
protein adapted from Hell et al. 1998. R1ab is the peptide used in the fibronectin and
thrombospondin binding experiments. Bottom- Alignment of a portion of R1 showing amino acid
sequence in Aas from selected *S. saprophyticus* strains. Amino acids are colored based on
their propensity to form beta strands (light green=high propensity, light blue=low propensity).
The alignment visualization was created in JalView.

Figure 5. Site frequency spectrum of lineage U. The ancient genome (Troy) was used as the outgroup to determine the ancestral state. Synonymous, nonsynonymous, and intergenic sites were identified with SnpEff (98). The observed synonymous SFS contain an excess of singletons and high frequency derived variants. Both the observed SFS and the SFS predicted by the best fitting model have an excess of singletons compared to the SFS expected under the standard neutral model with no population size change.

Figure 6. Cartoon of fitted demographic models. The observed synonymous SFS was fit to 5
demographic models including constant size, instantaneous population size change,
exponential population size change, instantaneous population size change followed by
exponential, and two instantaneous population size changes**.** Parameters for the instantaneous
and exponential models are the magnitude of the population size change ($\nu = N_e/N_{ancestral}$) and
the timing of the change (τ = generations/ $N_{ancestral}$**).** For models with two population size
changes, magnitudes are reported as $\nu_A = N_b/N_{ancestral}$ and $\nu_b = N_e/N_{ancestral.}$