Active mode of excretion across digestive tissues predates the origin of excretory organs

Carmen Andrikou*, Daniel Thiel, Juan A. Ruiz-Santiesteban, Andreas Hejnol*

Sars International Centre for Marine Molecular Biology, University of Bergen,

Thormøhlensgate 55, 5006 Bergen, Norway

Most bilaterian animals excrete toxic metabolites through specialized organs, such as

nephridia and kidneys, which share morphological and functional correspondences. In

contrast, the excretory mechanisms in non-nephrozoans are largely unknown, and

therefore the reconstruction of ancestral excretory mechanisms is problematic. Here, we

investigated the excretory mode of members of the Xenacoelomorpha, the sister group

to Nephrozoa, and Cnidaria, the sister group to Bilateria. By combining gene

expression, inhibitor experiments and exposure to varying environmental ammonia

conditions we show that both, Xenacoelomorpha and Cnidaria, are able to excrete

across digestive-associated tissues. Based on these results we propose that digestive-

associated tissues functioned as excretory sites before the evolution of specialized organs

in nephrozoans. We conclude that diffusion was likely the ancestral mode of excretion,

whilst the emergence of a compact, multiple-layered bilaterian body plan necessitated

the evolution of active transport excretory mechanisms that was later recruited into the

specialized excretory organs.

Introduction

Excretory organs are specialized organs that remove toxic metabolic waste products and

control water and ion balance in animals based on the principles of ultrafiltration, active

transport and passive transport/diffusion[1]. They are only present in Nephrozoa

(Deuterostomia + Protostomia)[2] (Fig. 1a) and based on morphological correspondences can

be grouped into two major architectural units: the protonephridia, only found in Protostomia,

and the metanephridia, present in both Deuterostomia and Protostomia[3,4]. Both organs are

organized into functionally similar compartments: the terminal cells of protonephridia and the

podocytes associated to metanephridial systems conduct ultrafiltration, and the tubule and

duct cells modify the filtrate through a series of selective reabsorption and secretion, via

passive and active transport mechanisms[5] (Fig. 1b). There exist also other, taxon-specific

excretory organs and excretory sites, which perform either ultrafiltration (such as the
nephrocytes of insects[6] and the rhogocytes of gastropods[7]), or absorption and secretion (such
as the malpighian tubules of various tardigrades, arachnids and insects[8], the excretory cells of
nematodes[9], the gills of fish, shore crabs and annelids[10,11], and the epidermis of planarians[12]).
Molecular studies have shown that a suite of orthologous genes is involved in the excretory
mechanisms of different nephrozoan species, regardless whether they possess specialized
excretory organs or not[9-30] (see also Supplementary Table 1). The passive ammonia
transporters Rhesus/AMTs, the active transporter $Na^+/K^+[NH_4^+]$ ATPase (NKA), the
hyperpolarization-activated cyclic nucleotide-gated $K^+[NH_4^+]$ channels (HCN), the vacuolar
H^+-ATPase (v-ATPase subunits A and B), members of the Carbonic Anhydrase A (CA) and
the water/glycerol/ammonia channels (aquaporins) are commonly used for excreting
ammonia, the most toxic metabolite (Fig. 1c) (summarized in[5,31]). Also, a group of
orthologous slit diaphragm structural components (nephrin/kirre, cd2ap, zo1,
stomatin/podocin), whose function is associated with the maintenance of the ultrafiltration
apparatus by interacting with the actin cytoskeleton and forming tight junctions[32], is localized
at the ultrafiltration site of the podocytes of the rodent kidney[33] (Fig. 1b) as well as in the
Drosophila nephrocytes[34] and the rhogocytes of gastropods[7]. Finally, a number of ion
transporters (solute carrier transporters (SLCs)) are spatially expressed in the corresponding
compartments of protonephridia of planarians and metanephridia (e.g. kidneys) of
vertebrates[35-37] (Fig. 1b).

The excretory sites and mechanisms in non-nephrozoans, however, are largely unknown. It is
commonly stated that excretion is presumably occurring via diffusion across the body wall
due to the loose (e.g. sponges) or single-epithelial (cnidarians and ctenophores) cellular
organization of these animals[1,38,39] (Fig. 1a) (herein stated as "diffusion hypothesis"). Based
on this idea, it was hypothesized that the emergence of the first excretory organs coincided
with the evolution of multilayered, solid parenchymes and increased body sizes due to the
need of more elaborate excretory mechanisms[40,41]. However, since excretion in non-
nephrozoans was never investigated in detail, the ancestral mechanisms of excretion and the
evolutionary origin of excretory organs remain unresolved[1,40-45].

An important animal group for our understanding nephrozoan evolution is their bilaterian
sister group[2,46,47], the Xenacoelomorpha (*Xenoturbella* + (Nemertodermatida + Acoela)).
These small, worm-like animals exhibit a bilateral symmetric, multilayered body plan, but
except for a special cell type with a putative excretory function (dermonephridia)[48] that seems
unique to the acoel *Paratomella*, xenacoelomorphs lack excretory organs and no defined

excretory sites have yet been described. To understand the excretory mechanisms outside
Nephrozoa and to gain insights into ancient excretory mechanisms, we therefore investigated
the excretory mode in two xenacoelomorph species and compared our findings with the non-
bilaterian cnidarian *Nematostella vectensis*.

Figure 1. Traditional diffusion hypothesis, ammonia transport mechanism and structural and
functional correspondences of protonephridial and metanephridial systems (**a**) Illustrated
phylogenetic relationship between Nephrozoa, Xenacoelomorpha and non-bilaterians[47]. Excretory
organs or specialized excretory cells/tissues using active transport (at) and ultrafiltration (ul) are so far
only reported in the group of Nephrozoa. (**b**) Cartoon depiction of the structural components of
metanephridia (podocyte, duct, tubule) in comparison to protonephridia (terminal cell, duct, tubule) and
summary of the expression domains of orthologous selected genes in relation to their components.
Animal illustrations are taken from phylopic.org (CC BY 3.0). (**c**) Ammonia cellular transport. Ammonia
(NH_3) is secreted into the lumen fluid via parallel H^+ and ammonia transport. This involves passive
diffusion through the cell membrane (dashed lines), facilitated diffusion via the Rhesus glycoproteins
(Rh), active transport via the $Na^+/K^+[NH4^+]$ ATPase (NKA), the hyperpolarization-activated cyclic
nucleotide- gated $K^+[NH4^+]$ channel (HCN) and aquaporin transporters (AQ) as well as the generation
of H^+ gradient by a vacuolar H^+-ATPase proton pump (v-ATPase) and the carbonic anhydrase (CA),
which transforms CO_2 into H^+ and HCO_3^-. Vesicular ammonia-trapping mechanism is also illustrated.

Results

Most genes involved in excretion in Nephrozoa are already present in non-nephrozoan and non-bilaterian animals

To get an overview of the presence of excretion-related genes in xenacoelomorphs and non-bilaterian animals we first searched for orthologous sequences of 20 nephrozoan candidate genes in the available transcriptomes and draft genomes of 13 xenacoelomorph species as well as in representatives of cnidarians, placozoans and sponges (Supplementary Fig. 1a and 2). We found that most of these genes were present in almost all groups with the exceptions of *slc5* (a sodium glucose cotransporter) that was only present in Cnidaria and Bilateria, and the ultrafiltration component *nephrin/kirre* that was only present in Bilateria (Supplementary Table 1 and Supplementary Fig. 1a). This analysis also revealed that the last common ancestor of placozoans, cnidarians and bilaterians already had at least two paralogs of *amts* (*amt2/3* and *am1/4*) and one *rhesus*, with independent duplications of one or both of these in various animal lineages (Supplementary Fig. 2). To identify potential excretory sites in xenacoelomorphs we examined the expression of the entire set of these candidate excretion-related genes in the acoel *Isodiametra pulchra* and the nemertodermatid *Meara stichopi*[49], which differ in their digestive system composition (*I. pulchra* has a syncytial, lumen-less gut, whereas *M. stichopi* has an epithelia-lined, cellular gut[50]) (Supplementary Fig. 1b).

Genes encoding solute carrier transporters and slit diaphragm-related components are expressed broadly in *Isodiametra pulchra* and *Meara stichopi*

Genes encoding SLCs were broadly expressed within neural (brain and nerve cords), parenchymal/subepidermal, digestive and gonadal-associated cells in both animals (Fig. 2 and 3). In particular, the three *slc1* paralogs isolated from *I. pulchra* were expressed in the brain (Fig. 2A1, A3) and around the male gonopore (Fig. 2A2, A3) and the single copy found in *M. stichopi* was restricted to scattered cells, resembling neurons (Fig. 3A). The two *slc5* paralogs found in *I. pulchra* had diverse expression patterns, with one copy being expressed around the male gonopore (Fig. 2B1) and the other one labeling brain and parenchymal cells (Fig. 2B2), whilst the *slc5* copy of *M. stichopi* was expressed in scattered subepidermal cells and two posterior lateral rows of cells (Fig. 3B). The single *slc13* copy of *I. pulchra* was marking brain and parenchymal cells (Fig. 2C) and in the same fashion all four paralogs of *M. stichopi* were expressed in scattered cells, resembling neurons (Fig. 3C1-C3). The three

slc4 paralogs isolated from *I. pulchra* were labeling individual cells at the anterior tip (Fig.

2D1, D3), brain (Fig. 2D2) and parenchymal cells (Fig. 2D2-D3), whilst in *M. stichopi* the

three *slc4* paralogs were broadly expressed including gut-wrapping cells and the mouth (Fig.

3D1), scattered subepidermal cells (Fig. 3D2-D3), gonads (Fig. 3D3) and nerve cords (Fig.

3D2 and Supplementary Fig. 3E). *Slc8* was restricted in a cup-shape region of the anterior tip

in *I. pulchra* (Fig. 2E) and in *M. stichopi* was expressed in the gut epithelium (including the

mouth) and two posterior lateral rows of cells (Fig. 3E). *Slc9* was marking the mouth in *I.*

pulchra (Fig. 2F) and in *M. stichopi* was expressed in two proximal lateral rows of cells,

resembling neurite bundles (Fig. 3F). Both *slc12* paralogs found in *I. pulchra* were labeling

neural (brain) and parenchymal cells (Fig. 2G1-G2); whilst in *M. stichopi* the expression of

the two paralogs investigated was associated with the male and female gonad region (Fig.

3G1-G2). One *slc26* copy did not reveal any expression in *I. pulchra* (Fig. 2H) although in

M. stichopi the two copies were expressed differently, with one gene being restricted in the

female gonads (Fig. 3H1) and the second one to be labeling gut-wrapping cells (Fig. 3H2).

Genes related to ultrafiltration sites had also broad expression patterns in both animals

covering neural, parenchymal/subepidermal cells as well as digestive and gonadal-associated

cells (Supplementary Fig. 4). All three *nephrin/kirre* paralogs were labeling neural cells both

in *I. pulchra* and *M. stichopi* (Supplementary Fig.4A). *Nephrin/kirre 1* was expressed in the

brain and around the male gonopore in *I. pulchra* (Supplementary Fig. 4A1) whilst

nephrin/kirre 2 and *3* were expressed in scattered cells, resembling neurons (Supplementary

Fig. 4A2-3). In *M. stichopi,* all three paralogs were also expressed in proximal lateral rows of

cells, resembling neurite bundles (Supplementary Fig. 4A1'-3'). *Cd2ap* was expressed in few

cells posterior to the statocyst and in scattered parenchymal cells, resembling neural cells in *I.*

pulchra (Supplementary Fig. 4B), whilst in *M. stichopi* this gene had a broad expression

pattern including subepidermal cells, the mouth and two posterior lateral rows of cells

(Supplementary Fig. 4B'). *Zo1* was marking anterior cells (probably the brain), the mouth

and scattered cells of the parenchyme, resembling neurons in *I. pulchra* (Supplementary Fig.

4C), while in *M. stichopi* was restricted in gut-affiliated cells and the mouth (Supplementary

Fig. 4C'). Finally, from the three *stomatin/podocin* paralogs found in *I. pulchra,* two were

expressed in the brain (Supplementary Fig. 4D1, D3) and the third one in cells of the

digestive syncytium (Supplementary Fig. 4D2) and in *M. stichopi* the five paralogs were

mainly expressed in proximal lateral rows of cells, resembling neural bundles

(Supplementary Fig. 4D1', D5'), scattered subepidermal cells (Supplementary Fig. 4D3'-

D4') and the mouth (Supplementary Fig. 4D2').

The broad expression of solute carrier transporters and slit diaphragm-related components in
acoelomorphs suggests that they are not part of defined excretory domains, as in
nephrozoans.
159

160
Figure 2. Expression of solute carrier transporters related to excrete modification in *I. pulchra.*
Whole mount in situ hybridization (WMISH) of genes encoding *slc1, slc5* and *slc13* (proximal tubule
related), *slc4, slc8, slc9, slc12* and *slc26* (distal tubule and duct related). The inset in panel A3 shows
a different focal plane. Illustrations with colored gene expression correspond to the animals shown in
the previous column. Anterior is to the left. The depicted expression patterns are for guidance and not
necessarily represent exact expression domains. Drawings are not to scale. BG indicates background
staining. Abbreviations: mo, mouth; mg, male gonopore; st, statocyst.

Figure 3. Expression of solute carrier transporters related to excrete modification in *M. stichopi.* Whole mount in situ hybridization (WMISH) of genes encoding *slc1, slc5* and *slc13* (proximal tubule related), *slc4, slc8, slc9, slc12* and *slc26* (distal tubule and duct related). The columns next to the panels show higher magnifications of the indicated domains, except of C1 that shows a different view form the animal (side view). Illustrations with colored gene expression correspond to the animals shown in the previous column. Anterior is to the left. The depicted expression patterns are for guidance and not necessarily represent exact expression domains. Drawings are not to scale. Abbreviations: mo, mouth.

The expression of a number of ammonia excretion-related genes and aquaporins suggests digestive-associated domains as putative excretion sites in *Isodiametra pulchra* and *Meara stichopi*

Genes involved in ammonia excretion (*rhesus/amts, nka, v-ATPase B, ca, hcn*) were mainly demarcating neural, digestive-associated and other parenchymal/subepidermal cells, as well

as epidermal cells (Fig. 4 and 5). In particular, the ammonia transporter *rhesus* was expressed

in cells of the anterior tip, gut-affiliated cells and the posterior ventral epidermis both in *I.*

pulchra and *M. stichopi* (Fig. 4A, 5A). The six *amt* paralogs found in *I. pulchra* were broadly

expressed including individual cells of the anterior tip (Fig. 4B5), scattered parenchymal cells

resembling neurons (Fig. 4B4), the brain (Fig. 4B1, B5), a cup-shape region of the anterior

tip (Fig. 4B1, B2), the mouth (Fig. 4B1), parenchymal and gut-affiliated cells (Fig. 4B3, B5),

whist the only *amt* present in *M. stichopi* was restricted to the whole epidermis (Fig. 5B).

Both *nka* paralogs of *I. pulchra* were strongly expressed in a sac-shaped gut-wrapping

domain (Fig. 4C). To check whether the expression was extending to the male gonadal

domain double FISH with markers of the gonads (*piwi* and *vasa*) were conducted and showed

that the expression of *nka* is restricted in a defined gut-wrapping domain, adjacent to the male

gonads (testes) (Supplementary Fig. 3B). Similarly, the single *nka* copy of *M. stichopi* was

labeling gut-affiliated cells (Fig. 5C). Nine genes belonging to the family of alpha-Carbonic

Anhydrases were found in *I. pulchra* and six in *M. stichopi*. *Ca a* was expressed in few distal

cells at the anterior tip, cells at the posterior tip adjacent to the male gonopore, two rows of

proximal parenchymal cells, resembling gut wrapping cells, and the mouth (Fig. 4D1), *ca b,*

ca c and *ca h* were all expressed in scattered parenchymal cells resembling neurons (Fig.

4D2-D3, D5), *ca x* was labeling the brain (Fig. 4D6), while *ca d* was expressed in a cup-

shape region of the anterior tip and parenchymal cells in *I. pulchra* (Fig. 4D4). In *M. stichopi*

ca a and *ca c* were marking gut-affiliated (Fig. 5D1, D3) and *ca b* was expressed in gut-

wrapping cells (Fig. 5D2). *Ca d* was expressed in scattered cells resembling neurons and cells

of the posterior epidermis (Fig. 5D4). *Hcn* was expressed in the brain in *I. pulchra* (Fig. 4E

and Supplementary Fig. 3C)*,* whilst in *M. stichopi* was labeling gut-affiliated cells (Fig. 5E).

V-ATPase B was strongly expressed in the digestive syncytium in *I. pulchra* (Fig. 4F and

Supplementary Fig. 3A) and in gut-affiliated cells in *M. stichopi* (Fig. 5F and Supplementary

Fig. 3D). In *M. stichopi v-ATPase B1* was additionally expressed in two posterior proximal

lateral rows of cells, resembling neurite bundles (Fig. 5F2) and *v-ATPase B2* in scattered

subepidermal cells (Fig. 5F1).

Figure 4. Expression of ammonia excretion-related genes in *I. pulchra.* Whole mount in situ
hybridization (WMISH) of genes encoding rh, v-ATPase B, nka, ca A 2 & 4, hcn and amts. The inset in
panel B6 and D1 show different focal planes. Illustrations with colored gene expression correspond to
the animals shown in the previous column. Anterior is to the left. The depicted expression patterns are
for guidance and not necessarily represent exact expression domains. Drawings are not to scale.
Abbreviations: mo, mouth.

Figure 5. Expression of ammonia excretion-related genes in *M. stichopi.* Whole mount in situ
hybridization (WMISH) of genes encoding rh, v-ATPase B, nka, ca A 2 & 4, hcn and amts. The
columns next to the panels show higher magnifications of the indicated domains. Illustrations with
colored gene expression correspond to the animals shown in the previous column. Anterior is to the
left. The depicted expression patterns are for guidance and not necessarily represent exact
expression domains. Drawings are not to scale.

The majority of genes encoding *aquaporins* were labeling digestive-associated domains and
neural domains. From the seven paralogs found in *I. pulchra, aq c, aq d, aq f* and *aq g* were
expressed in gut-wrapping cells (Fig. 6C-F) and *aq b* was labeling the digestive syncytium
(Fig. 6B). *Aq g* was additionally expressed in individual cells lining the gonads, whilst *aq a*
was expressed in the brain (Fig. 6A) and *aq e* was labeling parenchymal cells (Fig. 6G). In *M.*
stichopi six paralogs were found. *Aq d* and *aq f* were marking gut-wrapping cells (Fig. 7D,
F), while *aq a* and *aq c* were labeling proximal lateral rows of cells, resembling neurite
bundles (Fig. 7A, C). Finally, *aq b* was localized in individual subepidermal cells and *aq e*
was expressed in cells of the anterior tip and the ventral epidermis (Fig. 7E).
The broad expression of the ammonia excretion-related genes and *aquaporins* shows that
these genes do not label defined excretory domains, as already seen with the *slcs* and
ultrafiltration genes. However, since a considerable number of them (*rhesus,* members of *amt*
(only in *I. pulchra), nka,* members of *ca, hcn* (only in *M. stichopi*), *v-ATPase B* as well as
several *aquaporins)* were expressed in association with the digestion system, including
different types of gut-wrapping and gut-affiliated (syncytial in *I. pulchra* and gut epithelial in
M. stichopi) cells, the possibility that digestive-associated tissues could act as excretory sites
was raised.

Figure 6. Expression of aquaporins in *I. pulchra*. Whole mount in situ hybridization (WMISH) of aquaporin genes. The inset in panel B shows a different focal plane. Illustrations with colored gene expression correspond to the animals shown in the previous column. Anterior is to the left. The depicted expression patterns are for guidance and not necessarily represent exact expression domains. Drawings are not to scale.

253
Figure 7. Expression of aquaporins in *M. stichopi.* Whole mount in situ hybridization (WMISH) of aquaporin genes. The inset in panel E shows a different focal plane (side view). The columns next to the panels show higher magnifications of the indicated domains except of B that shows a different view form the animal (side view). Illustrations with colored gene expression correspond to the animals shown in the previous column. Anterior is to the left. The depicted expression patterns are for guidance and not necessarily represent exact expression domains. Drawings are not to scale.

High environmental ammonia exposure indicates a diffusion mechanism in *I. pulchra*

To reveal the excretory mechanism in Xenacoelomorphs, we conducted high environmental ammonia (HEA) incubation experiments, as previously performed in a large array of animals (summarized in[5,51]), using *I. pulchra* because of its availability in sufficient numbers. We first measured the pH of incubation mediums with different HEA concentrations (up to 1 mM) and found no difference in pH, which could otherwise have influenced any excretion rates. We then exposed animals to different HEA concentrations for a short period (2 hours) and measured the ammonia excretion during the following two hours after bringing them back into normal conditions, to test excretion via diffusion. The ammonia excretion rates of exposed animals remained unchanged after exposure to 50 and 100 µM NH_4Cl, compared to the control conditions, but increased gradually after exposure to NH_4Cl concentrations of 200 and 500 µM NH_4Cl (Fig. 8a). The increase in ammonia excretion rate could be explained by a concentration-dependent ammonia uptake during the HEA exposure and a subsequent release in normal conditions. These results suggest that ammonia excretion is concentration-dependent, which is indicative of a diffusion mechanism.

HEA exposure influences the expression of some excretion-related genes in *I. pulchra*

To test for a possible involvement of the excretion-related genes in the excretory mechanism of xenacoelomorphs, we tested for alteration of mRNA expression levels in chronically HEA exposed animals by quantitative relative expression experiments (qPCR) in *I. pulchra*. We first exposed animals to 1mM HEA for 7 days, similar to conditions used in previous studies (summarized in[5,51]), and measured the ammonia excretion over 2 hours after bringing the animals into normal conditions. As expected, the ammonia excretion rates were strongly increased, in line with the short termed HEA exposure experiments (Fig. 8a). When we tested the expression level of excretion-related genes, we found that the expression of the passive ammonia transporters *rhesus* and three *amts,* as well as the active ammonia transporter *nka* altered significantly (Fig. 8b). Other differentially expressed genes were four *aquaporins,* the *v-ATPase*, three *cas* and four *slcs* (Fig. 8b and Supplementary Fig. 5). These results indicate a putative role of these genes in ammonia excretion and suggest that acoels might not only excrete by diffusion and via passive transporters (*rhesus, amts*), but also by an alternative active transport mechanism (*nka*).

Inhibitor experiments support an active excretion mechanism via NKA transporter as well as a passive vesicular transport mode, possibly mediated by Rhesus transporter

We further tested the involvement of NKA, v-ATPase A/B and CA proteins in excretion, as
well as a possible involvement of a vesicular transport mechanism, by conducting
pharmacological inhibitor assays in *I. pulchra*, as previously demonstrated in other animals
(summarized in[5,51]). Inhibition of the CA by Azetazolamide, did not show any significant
change in ammonia excretion. Inhibition of the V-ATPase by Concanamycin C seemed to
lead to a decrease in ammonia excretion, although a 2-tailed t-test did not support a
significant change. In contrast, when perturbing the function of NKA with Quabain, the
ammonia excretion dropped significantly (Fig. 8c), which further support an active excretion
mechanism via NKA, similar to what is described for many nephrozoans[10-13,16,18,19,24-26,30,52-
54]. Interference with the vesicular transport using Colchicine also led to a significant decrease
in ammonia excretion, indicating a possible vesicular ammonia-trapping excretion mode (Fig.
8c), as demonstrated in the midgut epithelium of the tobacco hornworm[25], the gills of the
shore crab[26] and the integument of the nematode[13]. To test whether vesicular transport might
occur through Rhesus transporter as shown in other studies (summarized in[5,51]), we revealed
Rhesus protein localization by immunohistochemistry (Fig. 8d). The protein localization
mimicked the gene expression and revealed, apart of cells at the anterior tip and cells of the
posterior epidermis, individual parenchymal cells affiliated with the digestive syncytium that
extend ventrally. Higher magnification showed that the transporter was present in
cytoplasmic vesicles and not on the cellular membrane (Fig. 8d). This further indicated the
presence of a vesicular transport mechanism, in which cellular ammonia moves via Rhesus
transporters into vesicles and gets transferred to the membrane through the microtubule
network[55] (Fig. 1c). The antibody specificity was confirmed by an alignment of the epitope
and the endogenous protein, as a well as a western blot analysis (Supplementary Fig. 6).
Similar vesicular protein localization was also observed in *M. stichopi,* suggesting a similar
cytoplasmic-vesicular role of Rhesus transporter in gut-affiliated cells, also in
nemertodermatids (Fig. 8d). These data further supported the involvement of NKA and
Rhesus transporters in ammonia excretion and also indicated the presence of a putative
vesicular transport mode in *I. pulchra.*
323

Figure 8. Excretion in acoelomorphs. (**a**) Ammonia excretion rates of *I. pulchra* before (control) and after exposure for 2 hours to 50, 100, 200 and 500 µM and after exposure for 7 days in 1mM NH$_4$Cl (boxplot). Excretion was measured over two hours following the HEA treatments in at least three independent biological replicates, each divided into two separate samples (six measurements in total). Bold horizontal bar in boxes indicate the median, lower and upper box border indicate lower and upper quartile, whiskers indicate minimum and maximum. Asterisks label significant changes (p < 0.02 in an unpaired, 2-tailed t-test with unequal variance). (**b**) Quantitative relative expression of *rhesus, nka, v-ATPase B, amts, aquaporins,* and *ca* after 7 days exposure in HEA (1 mM NH$_4$Cl). Each circle indicates the average of three independent biological replicates, each with four technical replicates. Error bars indicate minimum and maximum of the biological replicates (averaged technical replicates). One-fold change represents no change; ≥ 2 indicates significantly increased expression level; ≤ 0.5 indicates significantly decreased expression level (red labels). (**c**) Effects of different inhibitors on ammonia excretion rates in *I. pulchra (boxplot, with illustration and replicates similar to Figure 4a).* The concentrations used were 5 µM Concanamycin C as a v-ATPase A/B inhibitor, 1 mM Azetazolamide as an inhibitor of the CA, 1 mM Quabain as a NKA inhibitor and 2 mM Colchicine for inhibiting the microtubule network. Concanamycin C was diluted in 0,5% DMSO for which we used an appropriate control with 0,5% DMSO. (**d**) Protein localization of Rhesus in *I. pulchra* and *M. stichopi*. Syncytium and gut are indicated in grey and the magenta staining of the lumen in *M. stichopi* is false positive staining of the gut content. Fluorescent pictures are projections of merged confocal stacks. The

nervous system is stained green with tyrosinated tubulin. (**e**) Double fluorescent WMISH of *v-ATPase* and *nka, aquaporin c* and *nka, v-ATPase* and *aquaporin b* and *v-ATPase* and *rhesus* in *I. pulchra.* White areas in the first panel are the result of merged stacks and not of overlapping expression. Nuclei are stained blue with 4',6-diamidino-2-phenylindole (DAPI). Anterior is to the left. Abbreviations: ctrl, control; ds, digestive syncytium; gwc, gut-wrapping cells. Scale bars, 50 μm for *I. pulchra* and 100 μm for *M. stichopi.*

Double fluorescent WMISH of differentially expressed genes shows similar spatial arrangement in gut-associated domains in *I. pulchra* and *M. stichopi*

To obtain a better resolution and understanding of the relative topology of the differentially expressed genes, double fluorescent WMISH were conducted for *v-ATPase, nka* and *aquaporins b* and *c* and *rhesus* (Fig. 8e). *Nka* and *v-ATPase* were expressed in a mutually exclusive manner, with *v-ATPase* to be restricted in the ventral region digestive syncytium and *nka* in an adjacent parenchymal, distal sac-shaped gut-wrapping domain. *Aquaporin c* was coexpressed with *nka* and *aquaporin b* was partially overlapping with *v-ATPase,* with *aquaporin b* expression extending into the posterior region of the digestive syncytium. Finally, *rhesus* was partly coexpressed with *v-ATPase* in the ventral region of the digestive syncytium. Similar coexpression analysis of the orthologous genes was also conducted for *M. stichopi* and revealed striking similarities in their spatial arrangement to *I. pulchra* (Supplementary Fig. 7). *V-ATPase* expression was not overlapping with *nka,* as *v-ATPase* was restricted to the gut epithelium and in two proximal lateral rows of subepidermal cells whilst *nka* was limited to cells lining the distal part of the epithelial branches of the gut extending towards the subepidermis. *V-ATPase* was partly coexpressed with *rhesus* in ventral gut-affiliated cells. Overall, these data revealed a similar spatial arrangement in gut-associated domains in both animals, which seems to be unrelated to the presence of an epithelial gut or a syncytium.

Taken together, our findings suggest that *I. pulchra* uses different mechanisms for ammonia excretion that are also known from nephrozoans; an active ammonia excretion mechanism via NKA through the digestive system, as suggested by in situ hybridization, and a passive vesicular transport mechanism likely mediated by Rhesus through digestive and likely also epidermal tissues. Given the commonalities in the expression of the involved genes in both animals, these excretory mechanisms could be plesiomorphic for Xenacoelomorpha.

HEA experiments suggest a diffusion mechanism also in the cnidarian *Nematostella vectensis*

Because our results showed the involvement of active and passive transport mechanisms
across digestive tissues outside Nephrozoa, we also investigated a non-bilaterian species, the
cnidarian *Nematostella vectensis* (Supplementary Fig. 1b), to test whether this excretion
mode might also be present outside Bilateria. The only available excretion studies in
cnidarians are few morphological studies, which suggested that the septa filaments of the
anthozoan mesenteries and the radial canals of medusozoans could serve as putative
excretory sites[56], as well as some isotopic exchange experiments on *Hydra oligactis* that have
shown that the gastrodermis seems to be involved in osmoregulation[57]. Moreover, there is
evidence that Rh and AMT genes are generally involved in ammonium excretion in corals[58],
but localization studies that would suggest excretion sites are missing.

We first tested whether *N. vectensis* excretes via diffusion by exposing early juvenile animals
to HEA for 2 hours and measuring their ammonia excretion rates afterwards, similar to the
experiments performed with *I. pulchra* (Fig. 9a). We found that also in *N. vectensis* ammonia
excretion increased significantly after HEA exposure starting at 200 and 500 µM NH_4Cl (Fig.
9a). Measurements of the pH of each incubation medium showed that the pH dropped by 0.2
when the medium contained 500 µM NH_4Cl. However, when we measured the excretion of
animals over 2 hours in a medium with an accordingly lowered pH we found that a difference
of 0.2 did not change the excretion rates (Supplementary Table 2). Therefore, the increase in
ammonia excretion rates at 200 and 500 µM NH_4Cl indicates that ammonia excretion is
concentration-dependent, supporting a diffusion mechanism also in *N. vectensis*.

Quantitative gene expression and inhibitor experiments indicates an involvement of
passive but not active transport mechanisms in *N. vectensis*
We then exposed animals to 1mM HEA for 7 days and tested the expression of the
orthologous genes altered in *I. pulchra* treatments (*rh/amt, nka, v-ATPase B* and *ca*) by
qPCR. As expected from the short-term HEA exposure experiment, specimens exposed for 7
days in HEA condition showed increased ammonia excretion rates (Fig. 9a). Also, the
expression of the passive transporters *rhesus1, rhesus2, amt6* and *amt7* as well as *v-ATPase*
altered significantly (Fig. 9b). In contrast to *I. pulchra*, none of the two *nka* transporters
showed a significant change in expression. To test whether Rhesus acts via a vesicular
transport mechanism we conducted the same pharmacological experiment as in *I. pulchra*.
Contrary to the results from acoels, we found that inhibition of vesicular transport did not
alter the ammonia excretion (Fig. 9e). We also inhibited the excretory function of v-ATPase
and CA proteins and found that none of them showed any significant change in ammonia

excretion rates (Fig. 9e). Finally, when we perturbed the function of NKA the ammonia
excretion rates did not alter (Fig. 9e), confirming the qPCR results and further supporting the
non-involvement of this transporter in excretion (Fig. 9a). These results suggest that the
ammonia excretion of *N. vectensis* is likely mediated via the passive Rhesus and AMT
transporters but neither relies on active transport mechanism mediated by NKA or on
vesicular ammonia-trapping excretion mode (Fig. 9e).
419

Gene expression of excretion related genes reveals the gastrodermis as excretory site in
N. vectensis
To understand whether these genes were expressed in gastrodermal or epidermal cells, we
revealed the spatial expression of *rhesus, amts, nka* and *v-ATPase B* by WMISH in feeding
primary polyps. All genes were mainly demarcating gastrodermal domains, such as the
endodermal body wall, the directive mesenteries, septal filaments and the pharynx (Fig. 9c
and Supplementary Fig. 8). *Rhesus 1* was additionally expressed in the tentacular ectoderm
(Fig. 9c). Protein localization of Rh, NKA and V-ATPase B revealed by
immunohistochemistry reflected the transcript expression patterns (Fig. 9d and
Supplementary Fig. 9). High magnification of Rhesus antibody staining further revealed that
the transporter was not expressed in cytoplasmic vesicles supporting a non-vesicular transport
mechanism, in agreement to the inhibitor experiments. Also, it showed that Rhesus was
localized in individual cells of the tentacular ectoderm with clumped structures at the tentacle
surface, which resembled gland cells[59] (Fig. 9d). The NKA antibody was localized in
endodermal neurons and individual cells of the mesenteries, likely neural precursors
(Supplementary Fig. 9), thus suggesting a non-excretory function of this transporter, as
expected from the qPCR and inhibitor experiments. These data imply that gastrodermis
affiliated domains likely serve as excretory sites in *N. vectensis*.
438

Nematostella vectensis (Cnidaria)

Figure 9. Excretion in *Nematostella vectensis*. (**a**) Ammonia excretion rates of *N. vectensis* before

(control) and after exposure for 2 hours to 50, 100, 200 and 500 µM and after exposure for 7 days in

1mM NH_4Cl (boxplot). Excretion was measured over two hours following the HEA treatments in at

least 3 independent biological replicates, each divided into two separate samples (6 measurements in

total). Bold horizontal bar in boxes indicate the median, lower and upper box border indicate lower and

upper quartile, whiskers indicate minimum and maximum. Asterisk labels significant changes.

Significance, $p < 0.02$ (unpaired t-test with unequal variance). (**b**) Quantitative relative expression of

rhesus, nks, v-ATPase B, amts and *ca* after exposure for 7 days in HEA (1 mM NH_4Cl). Each circle

represents the average of five independent biological replicates, each with three technical replicates.

One fold change represents no change; ≥ 2 indicates increased expression level significantly; ≤ 0.5

indicates decreased expression level significantly (red labels). (**c**) Whole mount in situ hybridization of

rh 1, rh 2, rh 3, v-ATPase B and *amt6* in feeding primary polyps. Anterior is to the top. (**d**) Protein

localization of Rh and v-ATPase in *N. vectensis* early juvenile polyps. The muscle filaments are

labeled green with phalloidin and the nervous system is stained cyan with tyrosinated tubulin. Every

picture is a full projection of merged confocal stacks. Nuclei are stained blue with 4',6-diamidino-2-

phenylindole (DAPI). The regions shown are indicated with dashed boxes in the illustrated animal. (**e**)

Effects of different inhibitors on ammonia excretion rates in *N. vectensis (boxplot, with illustration and*

replicates similar to Figure 5a). The concentrations used were 5-15 µM Concanamycin C as a v-

ATPase A/B inhibitor, 1-3 mM Azetazolamide as an inhibitor of the CA, 1-5 mM Quabain as a NKA

inhibitor and 2-10 mM Colchicine for inhibiting the microtubule network. Quabain was diluted in 0,5%

DMSO for which we used an appropriate control with 0,5% DMSO. N=3 for all treatments.

Abbreviations: ctrl, control; ebw, endodermal body wall; mes, mesenteries; ph, pharynx; sf, septal

filament; ten, tentacles.

Discussion

Overall, our findings show that acoelomorphs use, in addition to diffusion, also active

transport mechanisms, in contrast to what has been previously assumed for non-

nephrozoans[1,38,39]. Our results also suggest that excretion takes place across digestive tissues

and likely also across the epidermis, as indicated from the *rhesus* (both animals) and *amt*

(only in *M. stichopi*) expression. *N. vectensis* also seem to use gastrodermal tissues as

excretory sites but we were not able to detect any active transport mechanism. However, we

don't know whether the absence of active transport is common to all cnidarian excretion or

just a feature of *N. vectensis.* More studies in other members of Cnidaria are necessary to

elucidate this issue. Interestingly, digestive tissues with additional or assigned excretory roles

have also been reported in several nephrozoans (e.g. vertebrates, annelids, insects,

nematodes, tunicates, chaetognaths)[15,25,30,60-66]. In the light of our results this excretion

mechanism likely reflects an ancient mechanism, before the evolution of specialized organs,

such as nephridia (Supplementary Fig. 10). The molecular spatial arrangement of the

excretion sites in non-nephrozoans, however, is not sharing topological arrangements with

common nephridial domains of nephrozoans (Fig. 1b), suggesting that they are evolutionary

unrelated to nephridia. It still remains unclear whether these domains are multifunctional or

consist of specialized excretory subdomains, however a degree of cell sub-functionalization

seems to be present, as indicated by the localized gene expression in different groups of gut

wrapping and gut epithelial cells. We can however exclude the presence of ultrafiltration

sites, in agreement with previous ultrastructure studies in acoelomorphs[49], since the

homologous essential molecular components of the ultrafiltration sites of nephridia and

nephrocytes are mostly expressed in neural domains in acoelomorphs, and are absent in non-

bilaterians (nephrins), suggesting their later recruitment in the nephrozoan filtration

apparatus' (Fig. 10).

To conclude, our study shows that active transport mechanism and excretion through
digestive tissues predates the evolution of specialized excretory systems. If this is based on a
convergent recruitment, or if it reflects an ancestral state for Bilateria remains unclear.
However, if the latter is true, it correlates with the emergence of multilayered body plans and
solid internal parenchymes that separate the body wall from their digestive tract, as seen in
xenacoelomorphs and nephrozoans. We thus propose that diffusion mechanisms were the
major excretory modes present in animals with single-layered epithelial organization (Fig.
10). The emergence of more complex, multilayered body plan necessitated an active transport
of excretes, which was later recruited in specific compartments of the complex excretory
organs in the lineage of Nephrozoa.
499

Figure 10. Evolution of excretory mechanisms. Illustration of the proposed direction of fluxes in Cnidaria and Xenacoelomorpha, and evolution of active ammonia transport and ultrafiltration mechanisms. Cnidaria excrete across their intestinal epithelium via diffusion whilst in Xenacoelomorphs excretion occurs both via diffusion across the epidermis and gut-associated tissues and via active transport across gut-associated tissues. Ultrafiltration mechanism originated within Nephrozoa. Abbreviations: m, mesoglea; c, cuticle.

Methods

No statistical methods were used to predetermine sample size. The experiments were not

randomized. The investigators were not blinded to allocation during experiments and outcome

assessment.

Gene cloning and orthology assignment

Putative orthologous sequences of genes of interest were identified by tBLASTx search

against the transcriptome (SRR2681926) of *Isodiametra pulchra*, the transcriptome

(SRR2681155) and draft genome of *Meara stichopi* and the genome of *Nematostella*

vectensis (http://genome.jgi.doe.gov). Additional transcriptomes of Xenacoelomorpha species

investigated were: *Childia submaculatum* (Acoela) (SRX1534054), *Convolutriloba*

macropyga (Acoela) (SRX1343815), *Diopisthoporus gymnopharyngeus* (Acoela)

(SRX1534055), *Diopisthoporus longitubus* (Acoela) (SRX1534056), *Eumecynostomum*

macrobursalium (Acoela) (SRX1534057), *Hofstenia miamia* (Acoela) (PRJNA241459),

Ascoparia sp. (Nemertodermatida) (SRX1343822), *Nemertoderma westbladi*

(Nemertodermatida) (SRX1343819), *Sterreria* sp. (Nemertodermatida) (SRX1343821),

Xenoturbella bocki (*Xenoturbella*) (SRX1343818) and *Xenoturbella profunda* (Xenoturbella)

(SRP064117). Sequences for the placozoan *Trichoplax adhaerens,* the sponge *Amphimedon*

queenslandica, the ctenophore *Mnemiopsis leidy,* the protist *Capsaspora owczarzaki,* the

amoeba *Dictyostelium discoideum* and the nephrozoans *Homo sapiens, Saccoglossus*

kowalevskii, Strongylocentrotus purpuratus, Xenopus laevis, Branchiostoma lanceolatum,

Capitella teleta, Crassostrea gigas, Lottia gigantea, Schmidtea mediterranea, Tribolium

castaneum, Caenorhabditis elegans and *Drosophila melanogaster* were obtained from

Uniprot and NCBI public databases. Gene orthology of genes of interest identified by

tBLASTx was tested by reciprocal BLAST against NCBI Genbank and followed by

phylogenetic analyses. Amino acid alignments were made with MUSCLE[67]. RAxML (version

8.2.9)[68] was used to conduct a maximum likelihood phylogenetic analysis. Fragments of the

genes of interest were amplified from cDNA of *I. pulchra*, *M. stichopi* and *N. vectensis* by

PCR using gene specific primers. PCR products were purified and cloned into a pGEM-T

Easy vector (Promega, USA) according to the manufacturer' s instruction and the identity of

inserts confirmed by sequencing. Gene accession numbers of the gene sequences are listed in

the Supplementary Table 3.

Data Availability

All newly determined sequences have been deposited in GenBank under accession numbers

XXX-XXX. Primer sequences are available on request.

Animal systems

Adult specimens of *Isodiametra pulchra* (Smith & Bush, 1991), *Meara stichopi* Westblad,

1949 and *Nematostella vectensis* Stephenson, 1935 were kept and handled as previously

described[69-72].

Whole Mount In Situ Hybridization

Animals were manually collected, fixed and processed for *in situ* hybridization as

described[73,74]. Labeled antisense RNA probes were transcribed from linearized DNA using

digoxigenin-11-UTP (Roche, USA), or labeled with DNP (Mirus, USA) according to the

manufacturer's instructions. For *I. pulchra* and *M. stichopi*, colorimetric Whole Mount In

Situ Hybridization (WMISH) was performed according to the protocol outlined in[73]. For *N.*

vectensis, we followed the protocol described by[75]. Double Fluorescent In Situ Hybridization

(FISH) was performed as the colorimetric WMISH with the following modifications: After

the post-hybridization steps, animals were incubated overnight with peroxidase-conjugated

antibodies at 4°C (anti-DIG-POD, Roche 1:500 dilution and anti-DNP, Perkin Elmer, 1:200

dilution) followed by the amplification of the signal with fluorophore-conjugated tyramides

(1:100 TSA reagent diluents, Perkin Elmer TSA Plus Cy3 or Cy5 Kit). Residual enzyme

activity was inhibited via 45-minute incubation in 0.1% hydrogen peroxide in PTW followed

by four PTW washes prior to addition and development of the second peroxidase-conjugated

antibody[76].

Whole Mount Immunohistochemistry

Animals were collected manually, fixed in 4% paraformaldehyde in SW for 50 minutes,

washed 3 times in PBT and incubated in 4% sheep serum in PBT for 30 min. The animals

were then incubated with commercially available primary antibodies (anti-RhAG (ab55911)

rabbit polyclonal antibody, dilution 1:50 (Abcam, USA), anti-Na$^+$/K$^+$ ATPase (a1 subunit) rat

monoclonal antibody, dilution 1:100 (Sigma-Aldrich, USA) and anti-v-ATPase B1/2 (L-20)

goat polyclonal antibody, dilution 1:50 (Biotechnology, Santa Cruz, USA) overnight at 4°C,

washed 10 times in PBT, and followed by incubation in 4% sheep serum in PBT for 30 min.

Specimens were then incubated with a secondary antibody (anti rabbit-AlexaFluor 555,

Invitrogen or anti rat-AlexaFluor 555 and anti-goat-AlexaFluor 555) diluted 1:1000 overnight

at 4°C followed by 10 washes in PBT. Nuclei were stained by incubation of
animals in DAPI 1:1000 and f-actin was stained by incubation in BODIPY labeled
phallacidin 5 U/ml overnight.

Inhibitor and HEA experiments

For excretion experiments approximately 300 *I. pulchra* (the number varied slightly between
the biological replica, but was similar in the corresponding controls and treatments) and 10 *N.*
vectensis were placed into glass vials with 2 ml UV sterilized natural seawater (1:4 diluted
with distilled water for *N. vectensis*) containing the appropriate inhibitor or ammonia
concentration. Animals were given 10 minutes to adjust to the medium before the solution
was exchanged with 2 ml of fresh medium with the same appropriate condition. For the
inhibitor experiment the medium was removed after 2 hours and stored at -80°C for later
measurements. Animals from the short-term high environmental ammonia (HEA)
experiments were incubated for 2 hours, then rinsed 5 times over 20-30 minutes and
incubated for another 2 hours in fresh medium without additional ammonia, after which the
medium was removed and frozen at -80°C. We tested different inhibitor concentrations that
were used in previous studies in other invertebrates[10,12,13,18,26]. The concentrations of 5-15 µM
Concanamycin C for inhibiting v-ATPase A/B, 1-3 mM Azetazolamide as an inhibitor of the
CA, 1-5 mM Quabain to inhibit the NKA and 2-10 mM Colchicine for inhibiting the
microtubule network were selected, as no other effects like shrinking or obvious changes in
morphology or behavioral were observed. After the inhibitor incubations, the animals were
washed several times and monitored in normal conditions for several days to ensure that the
inhibitors did not cause any unspecific permanent effects. Concanamycin C was diluted in
DMSO with a final concentration of 0.5% DMSO per sample for which we used an
appropriate control with 0.5% DMSO. For the HEA experiments we enriched seawater with
NH_4Cl to the final ammonia concentrations of 50 µM, 100 µM, 200 µM, 500 µM, and 1 mM.
We also measured the pH of both incubation mediums (HEA and control) and we found no
difference. All experiments were independently repeated at least three times at different time
points, and each repeat was divided into two samples.

Determination of ammonia excretion

Ammonia concentrations were measured with an ammonia sensitive electrode (Orion,
Thermo Scientific) according to[53]. Samples were diluted 1:4 with distilled water to prevent
salt precipitation (900 µl sample + 2.7 ml water) and total ammonia was transformed into

gaseous NH_3 by adding 54 µl ionic strength adjusting solution (1.36 ml/l trisodiumcitrate

dihydrate, 1 M NaOH). Due to the small ammonia concentrations the electrode-filling

solution was diluted to 10% with distilled water, as suggested in the electrode manual. In

control conditions we determined an average excretion of 44 pmol per adult animal per hour,

although the excretion varied between different biological replicates from different

generations (min. 32 pmol/animal/hour, max. 52 pmol/animal/hour), possibly due to slightly

fluctuating conditions during long-term animal rearing. Solutions with defined concentrations

of NH_4Cl for the standard curves were made together with the solutions used in the

experiments and stored in a similar way at -80°C. The differences in excretion rates were

tested for significance with an unpaired, 2-tailed t-test with unequal variance and a p value <

0.02 was seen as significant. Boxplots were created with 'R'.

Quantitative Gene Expression

100 treated *I. pulchra* and 5 *N. vectensis* were collected after 7 days of incubation in HEA

conditions (1 mM NH_4Cl) and tested for quantitative gene expression using the BIORAD

CFX96 Real time PCR detection system. ddCt values were calculated between treated and

control animals and converted to fold differences. All experiments were repeated three to five

times with different specimens (three biological replicates for *I. pulchra* and five biological

replicates for *N. vectensis*) and two to dour technical replicates were tested for each biological

replicate (four biological replicates for *I. pulchra* and three biological replicates for *N.*

vectensis). Fold changes were calculated using polyubiquitin, actin and 18S as references for

I. pulchra and *ATPsynthase* and *EF1b* as references for *N. vectensis*[77], and a threshold of two-

fold difference was chosen as a significant change. The Ct values are provided in

Supplementary Table 4 and the primer sequences used are provided in Supplementary Table

5.

Western blot

Whole animal extracts (50 *I. pulchra* adults and 5 *N. vectensis* juveniles) were fractionated by

SDS-PAGE, loaded on Mini-PROTEAN® TGX Stain-Free Precast Gels (Bio-Rad) and

transferred to a nitrocellulose membrane using a transfer apparatus according to the

manufacturer's protocols (Bio-Rad). After incubation with 5% nonfat milk in TBST (10 mM

Tris, pH 8.0, 150 mM NaCl, 0.5% Tween 20) for 60 min, the membrane was washed once

with TBST and incubated with antibodies against Rhesus (1:1000) and NKA (1: 500) at 4 °C

for 12 h. Membranes were washed three times for 10 min and incubated with a 1:5000

dilution of horseradish peroxidase-conjugated anti-mouse or anti-rabbit antibodies for 2 h.

Blots were washed with TBST three times and developed with the ECL system (Amersham

Biosciences) according to the manufacturer's protocols.

Documentation

Colorimetric WMISH specimens were imaged with a Zeiss AxioCam HRc mounted on a

Zeiss Axioscope A1 equipped with Nomarski optics and processed through Photoshop CS6

(Adobe). Fluorescent-labeled specimens were analyzed with a SP5 confocal laser microscope

(Leica, Germany) and processed by the ImageJ software version 2.0.0-rc-42/1.50d (Wayne

Rasband, NIH). Figure plates were arranged with Illustrator CS6 (Adobe).

References

1   Schmidt-Rhaesa, A. *The evolution of organ systems*. (Oxford University Press, 2007).
2   Jondelius, U., Ruiz-Trillo, I., Baguñà, J. & Riutort, M. The Nemertodermatida are
basal bilaterians and not members of the Platyhelminthes. *Zool Scr* **31**, 201-215,
doi:10.1046/J.1463-6409.2002.00090.X (2002).
3   Ichimura, K. & Sakai, T. Evolutionary morphology of podocytes and primary urine-
producing apparatus. *Anat Sci Int* **92**, 161-172, doi:10.1007/s12565-015-0317-7 (2017).
4   Ruppert, E. E. & Smith, P. R. The Functional-Organization of Filtration Nephridia.
Biological reviews of the Cambridge Philosophical Society **63**, 231-258, doi:Doi
10.1111/J.1469-185x.1988.Tb00631.X (1988).
5   Larsen, E. H. *et al.* Osmoregulation and Excretion. *Comprehensive Physiology* **4**, 405-
573 (2014).
6   Locke, M. & Russell, V. W. in *Microscopic Anatomy of Invertebrates* Vol.
11B:Insecta (ed F. W. & Locke Harrison, M.) 687-709 (Wiley-Liss, 1998).
7   Kokkinopoulou, M. *et al.* 3D-ultrastructure, functions and stress responses of
gastropod (*Biomphalaria glabrata*) rhogocytes. *PloS one* **9**, e101078,
doi:10.1371/journal.pone.0101078 (2014).
8   Wessing, A. & Eichelberg, D. Ultrastructural aspects of transport and and
accumulation of substances in the malpighian tubules. *Fortschritte der Zoologie* **23**, 148-172
(1975).
9   Adlimoghaddam, A., Weihrauch, D. & O'Donnell, M. J. Localization of K(+), H(+),
Na(+) and Ca(2)(+) fluxes to the excretory pore in Caenorhabditis elegans: application of
scanning ion-selective microelectrodes. *J Exp Biol* **217**, 4119-4122, doi:10.1242/jeb.112441
(2014).
10   Thiel, D. *et al.* Ammonia excretion in the marine polychaete *Eurythoe complanata*
(Annelida). *J Exp Biol* **220**, 425-436 (2016).
11   Weihrauch, D., Wilkie, M. P. & Walsh, P. J. Ammonia and urea transporters in gills
of fish and aquatic crustaceans. *J Exp Biol* **212**, 1716-1730, doi:10.1242/jeb.024851 (2009).
12   Weihrauch, D. *et al.* Ammonia excretion in the freshwater planarian *Schmidtea*
mediterranea. *J Exp Biol* **215**, 3242-3253, doi:10.1242/jeb.067942 (2012).
13   Adlimoghaddam, A. *et al.* Ammonia excretion in *Caenorhabditis elegans*: mechanism
and evidence of ammonia transport of the Rhesus protein CeRhr-1. *J Exp Biol* **218**, 675-683,
doi:10.1242/jeb.111856 (2015).

14   Adlimoghaddam, A. *et al.* Ammonia excretion in *Caenorhabditis elegans*:
Physiological and molecular characterization of the rhr-2 knock-out mutant. *Comparative*
biochemistry and physiology-Molecular & integrative physiology **195**, 46-54,
doi:10.1016/j.cbpa.2016.02.003 (2016).
15   Blaesse, A. K., Broehan, G., Meyer, H., Merzendorfer, H. & Weihrauch, D. Ammonia
uptake in *Manduca sexta* midgut is mediated by an amiloride sensitive cation/proton
exchanger: Transport studies and mRNA expression analysis of NHE7, 9, NHE8, and V-
ATPase (subunit D). *Comp Biochem Phys A* **157**, 364-376, doi:10.1016/J.cbpa.2010.08.004
(2010).
16   Cruz, M. J. *et al.* Cutaneous nitrogen excretion in the African clawed frog *Xenopus*
laevis: effects of high environmental ammonia (HEA). *Aquatic toxicology* **136-137**, 1-12,
doi:10.1016/j.aquatox.2013.03.002 (2013).
17   Fehsenfeld, S. & Weihrauch, D. The role of an ancestral hyperpolarization-activated
cyclic nucleotide-gated K+ channel in branchial acid-base regulation in the green crab,
Carcinus maenas. *J Exp Biol* **219**, 887-896, doi:10.1242/jeb.134502 (2016).
18   Gerber, L. *et al.* The Legs Have It: In Situ Expression of Ion Transporters V-Type
H(+)-ATPase and Na(+)/K(+)-ATPase in the Osmoregulatory Leg Organs of the Invading
Copepod *Eurytemora affinis*. *Physiological and biochemical zoology : PBZ* **89**, 233-250
(2016).
19   Hu, M. Y. *et al.* Development in a naturally acidified environment: Na+/H+-
exchanger 3-based proton secretion leads to CO2 tolerance in cephalopod embryos. *Front*
Zool **10**, 51, doi:10.1186/1742-9994-10-51 (2013).
20   Mak, D. O., Dang, B., Weiner, I. D., Foskett, J. K. & Westhoff, C. M.
Characterization of ammonia transport by the kidney Rh glycoproteins RhBG and RhCG.
American Journal of Physiology - Renal Physiology **290**, F297-305,
doi:10.1152/ajprenal.00147.2005 (2006).
21   Patrick, M. L., Aimanova, K., Sanders, H. R. & Gill, S. S. P-type Na+/K+-ATPase
and V-type H+-ATPase expression patterns in the osmoregulatory organs of larval and adult
mosquito Aedes aegypti. *J Exp Biol* **209**, 4638-4651, doi:10.1242/jeb.02551 (2006).
22   Quijada-Rodriguez, A. R. *et al.* Ammonia-independent sodium uptake mediated by
Na(+) channels and NHEs in the freshwater ribbon leech Nephelopsis obscura. *J Exp Biol*
220, 3270-3279, doi:10.1242/jeb.159459 (2017).
23   Quijada-Rodriguez, A. R., Treberg, J. R. & Weihrauch, D. Mechanism of ammonia
excretion in the freshwater leech *Nephelopsis obscura*: characterization of a primitive Rh
protein and effects of high environmental ammonia. *American journal of physiology-*
Regulatory, integrative and comparative physiology **309**, R692-705,
doi:10.1152/ajpregu.00482.2014 (2015).
24   Shih, T. H., Horng, J. L., Hwang, P. P. & Lin, L. Y. Ammonia excretion by the skin of
zebrafish (*Danio rerio*) larvae. *American Journal of Physiology - Cell Physiology* **295**,
C1625-1632, doi:10.1152/ajpcell.00255.2008 (2008).
25   Weihrauch, D. Active ammonia absorption in the midgut of the tobacco hornworm
Manduca sexta L.: Transport studies and mRNA expression analysis of a Rhesus-like
ammonia transporter. *Insect Biochem Molec* **36**, 808-821, doi:10.1016/j.ibmb.2006.08.002
(2006).
26   Weihrauch, D., Ziegler, A., Siebers, D. & Towle, D. W. Active ammonia excretion
across the gills of the green shore crab *Carcinus maenas*: participation of Na(+)/K(+)-
ATPase, V-type H(+)-ATPase and functional microtubules. *J Exp Biol* **205**, 2765-2775
(2002).

27   Weiner, I. D. & Verlander, J. W. Role of NH3 and NH4+ transporters in renal acid-
base transport. *American journal of physiology. Renal physiology* **300**, F11-23,
doi:10.1152/ajprenal.00554.2010 (2011).
28   Weiner, I. D. & Verlander, J. W. Ammonia transport in the kidney by Rhesus
glycoproteins. *American journal of physiology. Renal physiology* **306**, F1107-1120,
doi:10.1152/ajprenal.00013.2014 (2014).
29   Wood, C. M. *et al.* Rh proteins and NH4(+)-activated Na+-ATPase in the *Magadi*
tilapia (Alcolapia grahami), a 100% ureotelic teleost fish. *J Exp Biol* **216**, 2998-3007,
doi:10.1242/jeb.078634 (2013).
30   Worrell, R. T., Merk, L. & Matthews, J. B. Ammonium transport in the colonic crypt
cell line, T84: role for Rhesus glycoproteins and NKCC1. *American journal of physiology.*
Gastrointestinal and liver physiology **294**, G429-440 (2008).
31   Gomes, D. *et al.* Aquaporins are multifunctional water and solute transporters highly
divergent in living organisms. *BBA - Biomembranes* **1788**, 1213-1228 (2009).
32   Fukasawa, H., Bornheimer, S., Kudlicka, K. & Farquhar, M. G. Slit diaphragms
contain tight junction proteins. *Journal of the American Society of Nephrology : JASN* **20**,
1491-1503, doi:10.1681/ASN.2008101117 (2009).
33   Patari-Sampo, A., Ihalmo, P. & Holthofer, H. Molecular basis of the glomerular
filtration: nephrin and the emerging protein complex at the podocyte slit diaphragm. *Annals of*
medicine **38**, 483-492 (2006).
34   Weavers, H. *et al.* The insect nephrocyte is a podocyte-like cell with a filtration slit
diaphragm. *Nature* **457**, 322-326 (2009).
35   Raciti, D. *et al.* Organization of the pronephric kidney revealed by large-scale gene
expression mapping. *Genome biology* **9**, R84 (2008).
36   Scimone, M. L., Srivastava, M., Bell, G. W. & Reddien, P. W. A regulatory program
for excretory system regeneration in planarians. *Development* **138**, 4387-4398 (2011).
37   Thi-Kim Vu, H. *et al.* Stem cells and fluid flow drive cyst formation in an invertebrate
excretory organ. *eLife* **4:e07405** (2015).
38   Greenberg, J. M. Ex Bouillabaisse Lux: The Charm of Comparative Physiology and
Biochemistry. *Amer Zool*, 737-749 (1985).
39   Ruppert, E. E., Fox, S. F. & Barnes, D. R. *Invertebrate Zoology, a functional*
evolutionary approach. 7th edn, (Brooks/Cole - Thomsen Learning, 2004).
40   Wilson, R. A. & Webster, L. A. Protonephridia. *Biological reviews of the Cambridge*
Philosophical Society **49**, 127-160 (1974).
41   Willmer, E. N. *Cytology and Evolution*. 2nd edn, (Academic Press, 1970).
42   Bartolomaeus, T. & Ax, P. Protonephridia and Metanephridia - Their Relation within
the Bilateria. *Z Zool Syst Evol* **30**, 21-45 (1992).
43   Goodrich, E. S. The study of nephridia and genital ducts since 1895. *The Quarterly*
journal of microscopical science **86**, 113-301 (1945).
44   Kümmel, G. Zwei neue Formen von Cyrtocyten. Vergleich der bisher bekannten
Cyrtocyten und Eröterung des Begriffes 'Zelltyp'. *Zeitschr für Zellforsch* **57**, 172-201 (1962).
45   Ruppert, E. E. Evolutionary Origin of the Vertebrate Nephron. *Am Zool* **34**, 542-553
(1994).
46   Rouse, G. W., Wilson, N. G., Carvajal, J. I. & Vrijenhoek, R. C. New deep-sea species
of *Xenoturbella* and the position of Xenacoelomorpha. *Nature* **530**, 94-97 (2016).
47   Cannon, J. T. *et al.* Xenacoelomorpha is the sister group to Nephrozoa. *Nature* **530**,
89-93 (2016).
48   Ehlers, U. Dermonephridia--modified epidermal cells with a probable excretory
function in *Paratomella rubra* (Acoela, Plathelminthes). *Microfauna Marina* **7**, 253-264
(1992b).

49   Smith, J. P. S. & Tyler, S. in *The origins and relationships of lower invertebrates*   (ed
S Conway-Morris, George, JD, Gibson, R, Platt, HM) (Oxford University Press, 1985).
50   Smith, J. & Tyler, S. in *The origins and relationships of lower invertebrates*   (eds S.
Conway Morris, J. D. George, R. Gibson, & H. M. Platt) 123-142 (Calderon Press, 1985).
51   Weihrauch, D. & Allen, G. J. P. Ammonia excretion in aquatic invertebrates: new
insights and questions. *J Exp Biol* **221**, jeb169219 (2018).
52   Wall, S. M. & Koger, L. M. NH+4 transport mediated by Na(+)-K(+)-ATPase in rat
inner medullary collecting duct. *The American journal of physiology* **267**, F660-670 (1994).
53   Weihrauch, D., Becker, W., Postel, U., Riestenpatt, S. & Siebers, D. Active excretion
of ammonia across the gills of the shore crab *Carcinus maenas* and its relation to
osmoregulatory ion uptake. *J Comp Physiol B* **168**, 364-376 (1998).
54   Weihrauch, D. & O'Donnell, M. J. Links between Osmoregulation and Nitrogen-
Excretion in Insects and Crustaceans. *Integr Comp Biol* **55**, 816-829, doi:10.1093/icb/icv013
(2015).
55   Kim, H. Y. *et al.* Basolateral expression of the ammonia transporter family member
Rh C glycoprotein in the mouse kidney. *American journal of physiology. Renal physiology*
296, F543-555, doi:10.1152/ajprenal.90637.2008 (2009).
56   Pütter, A. *Vergleichende Physiologie*. (G. Fischer, 1911).
57   Macklin, M. Osmotic regulation in *Hydra*: sodium and calcium localization and
source of electrical potential. *Journal of cellular physiology* **70**, 191-196 (1967).
58   Lehnert, E. M. *et al.* Extensive differences in gene expression between symbiotic and
aposymbiotic cnidarians. *G3* **4**, 277-295 (2014).
59   Moran, Y. *et al.* Neurotoxin localization to ectodermal gland cells uncovers an
alternative mechanism of venom delivery in sea anemones. *Proceedings. Biological sciences*
279, 1351-1358, doi:10.1098/rspb.2011.1731 (2012).
60   Burighel, P. & Cloney, R. A. in *Microscopic Anatomy of Invertebrates* Vol. 15 (eds
F.W. Harrison & E. E. Ruppert) Ch. Hemichordata, Chaetognatha, and the invertebrate
chordates, 221-347 (Wiley-Liss, 1997).
61   Butt, A. G. & Taylor, H. H. Salt and Water-Balance in the Spider, *Porrhothele*
Antipodiana (Mygalomorpha, Dipluridae) - Effects of Feeding Upon Hydrated Animals. *J*
Exp Biol **125**, 85-106 (1986).
62   Harpur, R. P. The nematode intestine and organic acid excretion: volatile acids in
Ascaris lumbricoides faeces. *Comparative biochemistry and physiology* **28**, 865-875 (1969).
63   Nijhout, H. F. Excretory Role of Midgut in Larvae of Tobacco Hornworm, *Manduca-
Sexta* (L). *J Exp Biol* **62**, 221-230 (1975).
64   Parry D. A. Habits, structure and development of Spadella cephaloptera and Sagitta
setosa. *J. Mar. Biol. Assoc.*, 16-36 (1944).
65   Phillips, J. E., Hanrahan, J., Chamberlin, M. & Thomson, B. Mechanisms and Control
of Reabsorption in Insect Hindgut. *Adv Insect Physiol* **19**, 329-422 (1986).
66   Tillinghast, E. K. Excretory pathways of ammonia and urea in the earthworm
Lumbricus   terrestris. *The   Journal   of   experimental   zoology* **166**,   295-300,
doi:10.1002/jez.1401660213 (1967).
67   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res* **32**, 1792-1797 (2004).
68   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis
of large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).
69   Børve, A. & Hejnol, A. Development and juvenile anatomy of the nemertodermatid
Meara stichopi (Bock) Westblad 1949 (Acoelomorpha). *Front Zool* **11** (2014).
70   De Mulder, K. *et al.* Characterization of the stem cell system of the acoel *Isodiametra
pulchra*. *Bmc Dev Biol* **9** (2009).

71   Fritzenwanker, J. H. & Technau, U. Induction of gametogenesis in the basal cnidarian
Nematostella vectensis (Anthozoa). *Development genes and evolution* **212**, 99-103 (2002).
72   Stephenson, T. A. *The British Sea Anemones*. Vol. II (The Ray Society, 1935).
73   Hejnol, A. & Martindale, M. Q. Acoel development indicates the independent
evolution of the bilaterian mouth and anus. *Nature* **456**, 382-386 (2008).
74   Martin-Duran, J. M. *et al.* Convergent evolution of bilaterian nerve cords. *Nature* **553**,
45-50, doi:10.1038/nature25030 (2018).
75   Genikhovich, G. & Technau, U. In situ hybridization of starlet sea anemone
(*Nematostella vectensis*) embryos, larvae, and polyps. *Cold Spring Harbor protocols* **2009**,
pdb prot5282 (2009).
76   Chiodin, M. *et al.* Mesodermal gene expression in the acoel *Isodiametra pulchra*
indicates a low number of mesodermal cell types and the endomesodermal origin of the
gonads. *PloS one* **8**, e55499 (2013).
77   Sinigaglia, C., Busengdal, H., Leclere, L., Technau, U. & Rentzsch, F. The bilaterian
head patterning gene six3/6 controls aboral domain development in a cnidarian. *PLoS biology*
11, e1001488, doi:10.1371/journal.pbio.1001488 (2013).
854

Acknowledgments

We thank Fabian Rentzsch, Patrick Steinmetz and Hanna Kraus (Sars Centre, University of

Bergen, Norway) for providing *N. vectensis* animals and cDNA. Peter Ladurner provided the

I. pulchra cultures that were kept in the animal facility at the Sars Centre. We thank all S9 lab

members for the help with the collections of *M. stichopi*. The study was supported by the core

budget of the Sars Centre, the European Research Council Community's Framework Program

Horizon 2020 (2014–2020) ERC grant agreement 648861 (EVOMESODERM) and FP7-

PEOPLE-2012-ITN grant no. 317172 (NEPTUNE) to A.H. and Coca-Cola Foundation to

J.A.R-S..

Author contributions

C.A. and A.H. designed the study. C.A., D.T. and J.A.R.-S. performed the experiments. C.A.

analyzed the data and C.A. and A.H. wrote the manuscript. All authors read and approved the

final manuscript.