Global phylogenetic analysis of *Escherichia coli* and plasmids carrying

the *mcr-1* gene indicates bacterial diversity but plasmid restriction

Sébastien Matamoros[1*], Jarne M. van Hattem[1*], Maris S. Arcilla[2], Niels Willemse[1], Damian C. Melles[2],

John Penders[3,4], Trung Nguyen Vinh [1,5,6], Ngo Thi Hoa[6,7], COMBAT consortium, Menno D. de Jong[1],

Constance Schultsz[1,5]

Department of Medical Microbiology, Academic Medical Center (AMC), Amsterdam, the

Netherlands[1]; Department of Medical Microbiology and Infectious Diseases, Erasmus University

Medical Center, Rotterdam, the Netherlands[2]; School for Public Health and Primary Care (Caphri),

Department of Medical Microbiology, Maastricht University Medical Center (MUMC), Maastricht, the

Netherlands[3]; School for Nutrition and Translational Research in Metabolism (NUTRIM), MUMC,

Maastricht, the Netherlands[4]; Department of Global Health-Amsterdam Institute for Global Health

and Development, AMC, Amsterdam, the Netherlands[5]; Oxford University Clinical Research Unit,

Centre for Tropical Medicine, Ho Chi Minh City, Vietnam[6]; Centre for Tropical Medicine, Nuffield

Department of Medicine, University of Oxford, Oxford, UK[7]

*S.M. and J.M.v.H. contributed equally to this work.

*Address correspondence to Sébastien Matamoros (s.p.matamoros@amc.nl)

Running title: Phylogenetic analysis of *mcr-1*-carrying *E. coli* isolates

Abstract

To understand the dynamics behind the worldwide spread of the *mcr-1* gene, we determined the population structure of *Escherichia coli* and of mobile genetic elements (MGEs) carrying the *mcr-1* gene. After a systematic review of the literature we included 65 *E. coli* whole genome sequences (WGS), adding 6 recently sequenced travel related isolates, and 312 MLST profiles. We included 219 MGEs described in 7 Enterobacteriaceae species isolated from human, animal and environmental samples. Despite a high overall diversity, 2 lineages were observed in the *E. coli* population that may function as reservoirs of the *mcr-1* gene, the largest of which was linked to ST10, a sequence type known for its ubiquity in human faecal samples and in food samples. No genotypic clustering by geographical origin or isolation source was observed. Amongst a total of 13 plasmid incompatibility types, the IncI2, IncX4 and IncHI2 plasmids accounted for more than 90% of MGEs carrying the *mcr-1* gene. We observed significant geographical clustering with regional spread of IncHI2 plasmids in Europe and IncI2 in Asia. These findings point towards promiscuous spread of the *mcr-1* gene by efficient horizontal gene transfer dominated by a limited number of plasmid incompatibility types.

Keywords

mcr-1; *Escherichia coli*; antimicrobial resistance; multilocus sequence typing; whole genome sequencing; plasmid; phylogeny; colistin; extended spectrum beta-lactamase (ESBL); international travel; ISA*pl1* transposon element.

Introduction

Antimicrobial resistance (AMR) represents a growing threat to global health [1]. With barely any new antimicrobial drugs in development [2], limiting the spread of AMR is key in order to maintain current treatment options [3].

Colistin is an antibiotic of the polymyxin class, discovered in 1950 and effective against Gram-negative bacteria [4]. The emergence of multidrug-resistant Gram-negative bacteria, especially those producing carbapenemases, has reintroduced colistin as a last resort antibiotic for the treatment of severe infections [5]. In contrast to its limited use in humans, colistin is widely used in food-producing animals [6]. While colistin resistance was long thought to be caused by chromosomal mutations only [7], the emergence of plasmid-mediated resistance, conferred by the mobilized colistin resistance (*mcr-1*) gene, was recently reported [8]. This gene encodes for a protein of the phosphoethanolamine transferase enzyme family, and its expression results in the addition of a phosphoethanolamine to lipid A, the target of colistin, decreasing the interaction between colistin and the bacterial lipopolysaccharide [8]. Since its discovery in 2015 in China, this gene has been described in several bacterial species that were isolated from animals, animal food products, humans and environmental samples from around the world [9–13]. Our previous study in travellers indicated acquisition of *mcr-1* carrying bacteria by healthy individuals during travel to destinations around the world, potentially related to food exposure, as well as rapid clearance after return [14]. It has been suggested that *mcr-1* has spread from food animals to humans [8,15–17], but there is a lack of comparison of *mcr-1* carrying isolates on a global level to support this hypothesis.

We studied the global population structure as well as the geographic and host distribution of *mcr-1*-carrying *Escherichia coli,* and mobile genetic elements (MGEs), to establish the population structure and to assess whether the spread of the *mcr-1* gene is linked to clonal dissemination or transmission of MGEs from animal, human, or environmental sources within geographic regions.

Results

Literature search

A systematic review of the literature on *mcr-1*, published until 1 January 2017 resulted in the

inclusion of 95 articles, representing a total of 410 entries (whole genome sequences, MLST profiles,

and/or plasmid types) for analysis (See detailed methods and results in Supplementary data,

Supplementary Figure 1 and Supplementary Table 1).

Population structure

Whole genome sequencing (WGS)

The genomes of 65 *mcr-1*-carrying *E. coli* were analysed, including 6 genomes obtained from *E. coli*

isolated from travellers that were sequenced for the purpose of the present study. Isolates

originated from Asia (n=36; 55.4%), Europe (n=20; 30.8%), North-America (n=4; 6.2%), South-America

(n=4; 6.2%) and Africa (n=1; 1.5%). 45 were of animal origin (69.2%), 19 of human origin (29.2%) and

one strain (1.5%) was isolated from water (Supplementary Table 1).

The average size of the genomes (all contigs in each assembly, representing chromosomes and

plasmids) of these 65 isolates was 4.9 Mbp, with a median number of genes identified of 4785

(ranging from 4266 to 7083), representing a pangenome of 23248 genes and a core genome (defined

by genes present in at least 99% of the isolates) of 2216 genes. An unbiased analysis of the

population structure was performed using a Bayesian approach with the BAPS software [18], based on

the nucleotide alignment of the core genome sequences. It revealed the presence of 5 distinct

phylogenetic clusters (Figure 1; Supplementary Figure 2; Supplementary Table 1). The largest cluster

(cluster 1) consisted of 26 isolates from 16 different STs (26/65; 40.0%) and the second cluster of 24

isolates from 15 different STs (36.9%). No significant relationship between clustering (BAPS) and

geographical origin or isolation source was observed (χ^2-test) (Figure 1) except that all 5 isolates that

belong to BAPS cluster 3 are from Europe. Twenty isolates showed less than 10 SNPs/Mbp difference

and were considered clonally related (Supplementary Tables 2 and 3).

 Multilocus sequence typing (MLST)

For 312 *E. coli* isolates originating from 69 studies, a MLST profile was published or could be deduced

from the corresponding WGS. Of these, 206 were isolated from animals or animal products (66.0%),

101 were isolated from humans (32.4%), including the 6 travel acquired isolates, and 5 from the

environment (1.6%). 141 Isolates from 25 studies (141/312; 45.2%) originated from Asia and 125

isolates from 25 studies (40.1%) from Europe together accounting for 85.3% of all included isolates.

The isolates represented 112 unique sequence types (STs) with ST10 being most common,

comprising 40/312 (12.8%) isolates originating from Africa, Asia, Europe and South-America.

eBURST analysis [19] was performed on all isolates included in the study, to identify their genetic

relatedness based on their MLST profiles. Three main clusters were identified, for which the

predicted founders, i.e. the ST in a cluster from which all other SLVs and DLVs in the cluster have

most likely diversified [19], were ST10, ST1114 and ST410. The largest cluster contained all 40 ST10

isolates and an additional 46 isolates in 21 STs that were single (SLV) or double locus variants (DLV) of

ST10 (86/312; 27.6%) (Supplementary Figure 3). The predicted founder of the second largest cluster

was ST1114, a SLV of ST165 and ST100, and included 19 isolates belonging to 7 different STs (5.4%),

while the third cluster was centred on ST410 and included 14 isolates from 3 different STs (4.5%).

A maximum-likelihood tree based on concatenated MLST gene sequences showed a main clade of

128 isolates (represented by blue branches in Figure 2 and Supplementary Figure 4; bootstrap value

of the main branch = 0.98), including most, but not all, isolates from the eBURST clusters of ST10 and

ST1114 (Supplementary Figure 4A). All isolates from these 2 eBURST clusters for which a WGS was

available were grouped in BAPS cluster 2. Similarly, all the isolates from the eBURST cluster ST410

grouped into BAPS cluster 1, along with 6 isolates from ST155. Seven isolates belonged to the

globally successful extra-intestinal pathogenic *E. coli* clone ST131 (Supplementary Figure 4A).

As observed in the WGS analysis, animal isolates were interspersed with isolates from humans and

the environment throughout the tree, as were isolates from different continents indicating a lack of

clustering by isolation source or geographical origin (Figure 2). Similarly, no clustering by health

status of the host was observed (Supplementary Figure 4B).

Mobile genetic elements

The plasmid incompatibility group of the *mcr-1*-carrying plasmids could be determined for 217

Enterobacteriaceae isolates from 7 different species (*Escherichia sp., Salmonella* sp., *Klebsiella* sp.,

Cronobacter sp., *Enterobacter* sp., *Kluyvera* sp. and *Shigella* sp.), representing a total of 219 plasmids

since 2 isolates carried 2 different plasmids carrying the *mcr-1* gene (Table 1). These plasmids were

described in 71 studies (1 to 33 plasmids per study, average = 3.1). In addition, the gene was

integrated in the chromosome of 6 isolates. The incompatibility group could not be determined for

27 of the 65 isolates for which WGS was available. Similarly the plasmid type was not available for

182 of the 312 isolates included in the MLST analysis. A total of 14 different plasmid incompatibility

groups were identified. 198/219 (90.4%) of the identified plasmids belonged to one of 3

incompatibility groups: IncX4 (77/219 plasmids, 35.2%), IncI2 (76/219 plasmids, 34.7%) and IncHI2

(45/219 plasmids, 20.5%). 50/76 IncI2 plasmids (65.8%) originated from Asia and 33/45 IncHI2

plasmids (73.3%) from Europe. IncX4 plasmids were more evenly distributed: 44/77 (57.1%) were

recovered from Europe, 29 from Asia (37.7%) and 4 from other regions (5.2%). Observed proportions

were significantly different from expected for IncI2 (χ^2-test, p < 0.001) and IncHI2 (p < 0.001) but not

for IncX4. The distribution of these 3 plasmid types was not significantly different from expected

between animal (χ^2-test, p = 0.24), human (p = 0.88) and environmental sources (p = 0.38) . Isolates

from the BAPS groups 1 and 2 carried plasmids from the 3 major types in similar proportions

(Supplementary Figure 5A; Supplementary Table 1). Isolates from the eBURST clusters of ST10 carried

plasmids belonging to 7 different incompatibility groups, including IncHI2, IncI2 and IncX4. No

clustering of plasmid type with MLST phylogeny was observed either (Supplementary Figure 5B;

Supplementary Table 1).

Figure 3 shows the alignment of the complete sequences or contigs from IncI2 (panel A), IncX4 (B)

and IncHI2 (C) incompatibility group plasmids. IncHI2 plasmids had the largest size, with sequence

lengths up to 267486 bp.

The ISA*pl1* transposon element situated upstream of the *mcr-1* gene was present in 7/9 (77.8%)

IncHI2 plasmids, but only in 11/29 (37.9%) of IncI2 plasmids and completely missing in all of the 24

reported IncX4 plasmids (Figure 3). In the isolates from travellers the ISA*pl1* transposon was

identified in 3 out of our 6 *mcr-1*-carrying contigs including one isolate from a traveler to Asia (ST101,

IncI2 incompatibility group), one to Africa (ST744, IncHI2) and one to South-America (ST744,

incompatibility group not identified).

Antimicrobial resistance genes

All sequences of the *mcr-1* gene collected in the present study were 100% identical to the original

sequence described by Liu *et al.* [8].

Multiple resistance genes were detected in most of the studied isolates (Supplementary Results). The

florfenicol resistance gene *floR* was present in 32 (49.2%) isolates; in 22 of 45 isolates from animals

(48.9%) and 10 of 19 isolates from humans (52.6%). The *baeR* and *baeS* genes, encoding novobiocin

resistance, were found in 64 (98.5%) and 65 (100%) isolates, respectively.

Plasmid analysis from the WGS data showed that 4 of the 29 *mcr-1*-carrying IncI2 plasmids (13.8%)

contained an additional ESBL gene. IncHI2 plasmids (n=9) carried between 0 and 12 additional AMR

genes. In particular, 4 plasmids carried CTX-M ESBL genes and 2 carried the *floR* gene. In 4 out of the

9 IncHI2 plasmids analysed in this study, the *mcr-1* gene was shown to be integrated alongside a

large multi-drug resistance (MDR) gene cassette (Figure 3C). None of the IncX4 plasmids carried

additional AMR genes.

Discussion

Analysis of all reported WGS of *mcr-1*-carrying isolates shows that the population of *E. coli* is highly

diverse, but is dominated by 2 large groups of related isolates. Most of the isolates from BAPS group

2 grouped into a MLST cluster centred on ST10. An overrepresentation of isolates related to ST10

and ST165 (a SLV of ST1114) in *mcr-1*-carrying *E. coli* isolates was previously reported at a smaller

scale in isolates from European farm animals [20]. *E. coli* ST10 and closely related STs are frequently

recovered from food and human intestinal samples and studies have shown a higher prevalence of

plasmid-carried AMR genes in ST10, including CTX-M ESBL genes, compared to other STs [21–24].

The second BAPS group of interest in our study included isolates belonging to ST155. This ST has

been described as a major vector of spread of ESBL genes from animals to humans [25]. It is thus

possible that zoonotic transmission lead to the spread of the *mcr-1* gene, as has been suggested in

studies from China and Vietnam [16,17], notably through the 2 main phylogenetic clusters identified in

this study.

Additionally, we found clonally related isolates, including some belonging to ST744, a SLV of ST10,

carrying the *mcr-1* gene on different plasmid backbones and recovered from different continents

(see Supplementary Results). These results point towards a worldwide dissemination *of mcr-1* driven

mainly by highly promiscuous plasmids rather than the worldwide spread of one or more *mcr-1*-

carrying clones. We hypothesize that several populations of *E. coli* isolates, notably those related to

ST10 or ST155, acquired the *mcr-1* gene due to their intrinsic ability of acquiring AMR genes and their

high prevalence in humans and food animals. These populations of commensal isolates then may

play a crucial role as a reservoir for this gene, which can explain their over-representation in the

present study.

In the timeframe of our literature search, 3 *E. coli* strains carrying the *mcr-2* gene were isolated from

animals in Belgium. These isolates belonged to ST10 (2 isolates) and ST167 which is a SLV of ST10 and

carried the gene on an IncX4-type plasmid. No WGS data was available from these isolates [26]. More

data about the *mcr-2* gene is needed to assess its spread and determine if *E. coli* ST10 plays a similar

role in its dissemination as it does for *mcr-1*.

More than 90% of published plasmid types carrying *mcr-1* genes belonged to either IncI2, IncX4 or

IncHI2. Almost 75% of the isolates carrying an IncHI2 plasmid originated from Europe: 26 from

animals and 7 from humans (Table 1 and Supplementary Table 1). In a traveller's isolate acquired in

Tunisia, the *mcr-1*-carrying plasmid was identified as an IncHI2-type backbone of the ST4 pMLST

subtype which co-carried a CTX-M-1 ESBL gene (Figure 3C). This traveller reported consumption of

beef, chicken and eggs during travel to Tunisia which can potentially be the source for the acquisition

of the *mcr-1* positive isolate. When investigating the presence of the *mcr-1* gene in cephalosporin

resistant *E. coli* isolates from chicken farms in Tunisia, Grami *et al.* [27] found that all 37 *mcr-1*-carrying

plasmids also belonged to the IncHI2-type, ST4 subtype and harboured CTX-M-1 genes. PFGE typing

of the isolates harbouring this plasmid showed various bacterial genetic backgrounds. Interestingly,

these chickens were all imported from France, either as adults or chicks. Other studies showed the

presence of this IncHI2, CTX-M-1 and *mcr-1* combination in *Salmonella enterica* Typhimurium isolates

from meat samples in Portugal from 2011 [28,29] and diarrhoeic veal calves in France [30]. The IncHI2

subtype ST4 was also detected in an *E. coli* isolate from retail chicken breast in Germany [31] and the

faecal sample of a veal calve from the Netherlands [32], suggesting widespread dissemination of this

particular plasmid in European farm animals and possible transmission to humans.

The high prevalence of novobiocin *baeR* and *baeS* and florfenicol *floR* resistance genes [33,34] in the

genomes of isolates of human and animal origin together with the fact that florfenicol and

novobiocin are used almost exclusively in veterinary medicine further supports the potential role of

food animals as an important reservoir of *mcr-1* containing bacteria and MGEs [15].

In contrast with the IncHI2 plasmids, 65.8% of all IncI2 plasmids recovered so far originated from

Asia, with a much lower prevalence in *mcr-1* carrying Enterobacteriaceae from other regions. Taken

together, these elements point toward a more regional circulation and dissemination of the *mcr-1*-

carrying plasmids IncI2 and IncHI2.

We found the IS*Apl1* transposon element associated with the *mcr-1* gene, as originally described by

Liu *et al.* [8] to be present in a minority of studied plasmids and contigs. However, since some of the

mcr-1-carrying contigs were obtained by assembly of Illumina short reads from WGS data, we cannot

exclude that some of these gaps are explained by an incomplete assembly of (plasmid) sequences.

The ISApl1 transposon element is considered to be the main driver of horizontal gene transfer of the

mcr-1 gene and has been shown to be highly unstable in IncI2 plasmids [35–37]. The absence of the

ISApl1 transposon element in mcr-1-carrying IncX4 plasmids as described here has recently been

proposed to be essential for the maintenance of the mcr-1 gene in this particular backbone, but the

exact mechanism still requires further investigation [38].

WGS analysis provided in-depth information about the mcr-1-carrying E. coli isolates and their

phylogenetic relationship, but the number of available genomes was limited. On the other hand,

whilst MLST data have a lower resolution, the higher number of available profiles allowed analysis of

the isolates' origin (geographical, source of isolation, diseased status of the host, etc.).

A limitation of our study is the potential for bias. The overrepresentation of isolates originating from

Asia and Europe could be explained by a higher prevalence of mcr-1 genes on these continents, but

the effect of publication bias cannot be excluded. Isolates from North-America only represented

2.2% of the collection. Noteworthy, colistin, except for ophthalmic ointment, has never been

marketed for use in animals in the United States [39,40].

Sampling bias should also be considered when several isolates with an identical ST are presented

from a single study, as is the case for ST100 and ST752. Additionally, in the absence of a control

population of mcr-1-negative isolates obtained from similar sources as the mcr-1 positive isolates,

results of analysis of population structures should be interpreted with caution. Because many studies

screened existing collections of (resistant) isolates for colistin resistance or presence of mcr-1,

selection bias has probably been introduced.

The findings in this study suggests that the mcr-1 gene has locally and globally disseminated through

MGEs that are mainly IncHI2, IncI2 and IncX4 plasmids and provides additional support for the

hypothesis of the animal reservoir, that is driven by the use of colistin in livestock, as a source of mcr-

1 in humans. A global ban of colistin use in animals to preserve colistin for use in human medicine

seems therefore justified.

Material and Methods

Selection of isolates for whole genome sequencing

250 We subjected 6 *mcr-1* positive isolates that were collected as part of a prospective study (COMBAT

251 study) aimed at studying acquisition of extended-spectrum β-lactamase (ESBL) -producing

252 Enterobacteriaceae during travel to whole-genome sequencing [14,41]. Additionally, we included 22

253 whole genome sequences of isolates from Vietnamese chickens and humans that were still

254 unpublished when performing our literature search [17].

Literature search

Relevant papers that published on *mcr-1* and *mcr-2* were identified in PubMed, Web of Science,

Scopus, ScienceDirect and Google Scholar using the query 'mcr-1 OR mcr1 OR mcr-2 OR mcr2 OR

(mcr AND colistin)' (see Supplementary Material for full search strategies).To be able to study the

associations between phylogeny, geographic distribution and isolation source we only included

sequences from papers that provided sufficient metadata. As a consequence, plasmids and

sequences that were deposited in online databases without metadata were not included in the

analysis.

Whole genome sequencing of *mcr-1*-positive *E. coli* isolates

Bacterial DNA was extracted from fresh pure cultures using the Qiagen DNeasy Blood and Tissue kit

(Qiagen, Hilden, Germany). Library preparation was done according to manufacturer's instruction

(Illumina, San Diego, CA, USA) and sequenced using Illumina MiSeq technology with 150nt paired-end

settings. Sequences have been deposited in the European Nucleotide Archive under the accession

numbers SAME104030441 to SAME104030446 .

Bio-informatic analysis

MLST analysis

For each *mcr-1*-carrying *E. coli* isolate for which the ST or the whole genome sequence was available,

the sequences of the corresponding alleles were downloaded from the *E. coli* MLST genes repository

of the University of Warwick

(http://mlst.warwick.ac.uk/mlst/dbs/Ecoli/handlers/getFileData/home/cbailster/mlst/zope/Extensio

ns/gadfly/Ecoli/DB/) and concatenated. When STs of isolates were not described in literature, the ST

was determined from available whole genomes using the online service provided by the Center for

Genomic Epidemiology (https://cge.cbs.dtu.dk/services/MLST/) according to the Achtman MLST

scheme [42,43]. MLST clusters (STs and their single locus or double locus variants) were defined using e-

burst V3 (http://eburst.mlst.net/v3/enter_data/single/) [19] and goeBURST v1.2.1 [44] using only profiles

from this study.

WGS and plasmid analysis

Raw sequence reads in fastq format or pre-assembled sequences in fasta format were downloaded

from online databases for all available isolates (Supplementary Table 1). Additional sequences not

yet deposited in online databases were requested from their respective authors. The quality of the

raw sequence reads was checked using fastqc

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/), quast [45] and KmerFinder 2.0

(https://cge.cbs.dtu.dk/services/KmerFinder/) (see Supplementary Methods for more details). Reads

were trimmed using Trimmomatic V0.33 [46]. *De-novo* genome assembly was performed with SPAdes

3.9 [47] for Illumina short reads and with Canu v1.3 for PacBio long reads [48]. Contigs of less than 500 bp

long were removed from the genomes to improve the overall quality of the assembly. Size of the

genomes was calculated by adding the length of all remaining contigs. Identification of open reading

frames (ORFs) and gene contents in the assembled genomes (*de-novo* assemblies and pre-assembled

sequences) was performed using Prokka v1.11 [49]. Core genome analysis was performed with Roary

v3.6.8 [50]. Clustering of isolates was performed using the hierBAPS module of the Bayesian Analysis of

Population Structure (BAPS) software v6.0 [18]. The core genome alignment output provided by Roary

was used as input for BAPS with 2 levels of hierarchy and a maximum number of cluster (K) of 10. The

estimated number of clusters was 5 for both levels of hierarchy.

Sequences (concatenated MLST loci or concatenated core genes) were aligned using mafft v6.864b

[51]. The resulting alignment was used as input for calculation of distances and tree building using

RAxML v8.1.6 [52]. TMLST and WGS trees were visualized using iTOL v3.3.1 (http://itol.embl.de/) [53].

Identification of plasmid incompatibility group and typing of IncHI2 plasmids were performed on

assembled sequences (*de-novo* or pre-assembled) via the CGE online services PlasmidFinder v1.3

(https://cge.cbs.dtu.dk/services/PlasmidFinder/) and pMLST v1.4

(https://cge.cbs.dtu.dk/services/pMLST/) [54]. Alignment and visualization of plasmids was performed

with BRIG v0.95 [55]. The majority of the isolates and plasmids described in this study were sequenced

using a short read technology (Illumina). This technology does not allow for a high quality assembly

of the plasmids due to the high number of repeat regions present in these MGEs. Therefore no

phylogenetic analysis of the *mcr-1*-carrying plasmids was conducted in this study.

Two different databases were used for identification of other antibiotic resistance genes: ResFinder

(https://cge.cbs.dtu.dk/services/ResFinder/ [56]) was used to detect acquired resistance genes

commonly located on mobile genetic elements (MGEs) and CARD Resistance Gene Identifier

(https://card.mcmaster.ca/analyze/rgi [57]) was used to detect chromosomal genes.

Statistics
The distribution of isolates and plasmids from different geographical origins and isolation sources

was determined by a χ^2-test comparing the expected distribution (proportions of the total studied

population) to the observed proportions using GraphPad Prism6 (La Jolla, CA, USA).

References

1.   Appelbaum, P. C. 2012 and beyond: Potential for the start of a second pre-antibiotic era? *J.*

Antimicrob. Chemother. **67,** 2062–2068 (2012).

2.   Brown, E. D. & Wright, G. D. Antibacterial drug discovery in the resistance era. *Nature* **529,**

336–343 (2016).

3.   Laxminarayan, R. *et al.* Antibiotic resistance-the need for global solutions. *Lancet Infect. Dis.*

13, 1057–1098 (2013).

4.   Koyama, Y., Kurosasa, A., Tsuchiya, A. & Takakuta, K. A new antibiotic 'colistin' produced by

spore-forming soil bacteria. *J Antibiot* **3,** 457–458 (1950).

5.   Lim, L. M. *et al.* Resurgence of colistin: a review of resistance, toxicity, pharmacodynamics,

and dosing. *Pharmacotherapy* **30,** 1279–91 (2010).

6.   Catry, B. *et al.* Use of colistin-containing products within the European Union and European

Economic Area (EU/EEA): development of resistance in animals and possible impact on human

and animal health. *Int. J. Antimicrob. Agents* **46,** 297–306 (2015).

7.   Olaitan, A. O., Morand, S. & Rolain, J. Mechanisms of polymyxin resistance : acquired and

intrinsic resistance in bacteria. *Front. Microbiol.* **5,** 1–18 (2014).

8.   Liu, Y.-Y. *et al.* Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in

animals and human beings in China: a microbiological and molecular biological study. *Lancet*

Infect. Dis. **3099,** (2015).

9.   Skov, R. L. & Monnet, D. L. Plasmid-mediated colistin resistance (mcr-1 gene): three months

later, the story unfolds. *Eurosurveillance* **21,** (2016).

10.   Schwarz, S. & Johnson, A. P. Transferable resistance to colistin: a new but old threat. *J.*

Antimicrob. Chemother. **71,** 2066–2070 (2016).

11.   Leangapichart, T. *et al.* Acquisition of mcr-1 Plasmid-Mediated Colistin Resistance in

Escherichia coli and Klebsiella pneumoniae during Hajj 2013 and 2014. *Antimicrob. Agents*

Chemother. **60,** 6998–6999 (2016).

12.   Vading, M. *et al.* Frequent acquisition of low-virulence strains of ESBL-producing Escherichia

coli in travellers. *J. Antimicrob. Chemother.* **71,** 3548–3555 (2016).

13.   Bernasconi, O. J. *et al.* Travelers Can Import Colistin-Resistant Enterobacteriaceae, Including

Those Possessing the Plasmid-Mediated mcr-1 Gene. *Antimicrob. Agents Chemother.* **60,**

5080–5084 (2016).

14.   Arcilla, M. S. *et al.* Dissemination of the mcr-1 colistin resistance gene. *Lancet*

Infect. Dis. **16,** 147–149 (2016).

15.   Poirel, L. & Nordmann, P. Emerging plasmid-encoded colistin resistance: the animal world as

the culprit? *J. Antimicrob. Chemother.* **71,** 2326–2327 (2016).

16.   Wang, Y. *et al.* Comprehensive resistome analysis reveals the prevalence of NDM and MCR-1

in Chinese poultry production. *Nat. Microbiol.* **2,** 16260 (2017).

17.   Trung, N. V. *et al.* Zoonotic Transmission of mcr-1 Colistin Resistance Gene from

Small-Scale Poultry Farms, Vietnam. *Emerg. Infect. Dis. J.* **23,** 529 (2017).

18.   Cheng, L., Connor, T. R., Sirén, J., Aanensen, D. M. & Corander, J. Hierarchical and spatially

explicit clustering of DNA sequences with BAPS software. *Mol. Biol. Evol.* **30,** 1224–1228

(2013).

19.   Feil Edward J an d Li, B. C., Aanensen, D. M., Hanage, W. P. & Spratt, B. G. eBURST: inferring

patterns of evolutionary descent among clusters of related bacterial genotypes from

multilocus sequence typing data. *J. Bacteriol.* **186,** 1518–1530 (2004).

20.   El Garch, F. *et al.* mcr-1 is borne by highly diverse Escherichia coli isolates since 2004 in food-

producing animals in Europe. *Clin. Microbiol. Infect.* **23,** 51e1-51e4 (2017).

21.   Manges, A. R. & Johnson, J. R. Food-Borne Origins of Escherichia coli Causing Extraintestinal

Infections. *Clin. Infect. Dis.* **55,** 712–719 (2012).

22.   Oteo, J. *et al.* Extended-spectrum β-lactamase-producing Escherichia coli in Spain belong to a

large variety of multilocus sequence typing types, including ST10 complex/A, ST23 complex/A

and ST131/B2. *Int. J. Antimicrob. Agents* **34,** 173–176 (2009).

23.   Fam, N. *et al.* CTX-M-15-producing Escherichia coli clinical isolates in Cairo (Egypt), including

isolates of clonal complex ST10 and clones ST131, ST73, and ST405 in both community and

hospital settings. *Microb. Drug Resist.* **17,** 67–73 (2011).

24.   Manges, A. R. *et al.* Clonal distribution and associated characteristics of Escherichia coli

clinical and surveillance isolates from a military medical center. *Diagn. Microbiol. Infect. Dis.*

87, 382–385 (2017).

25.   Skurnik, D. *et al.* Emergence of antimicrobial-resistant Escherichia coli of animal origin

spreading in humans. *Mol. Biol. Evol.* **33,** 898–914 (2016).

26.   Xavier, B. B. *et al.* Identification of a novel plasmid-mediated colistinresistance gene, mcr-2, in

Escherichia coli, Belgium, june 2016. *Eurosurveillance* **21,** (2016).

27.   Grami, R. *et al.* Impact of food animal trade on the spread of mcr-1-mediated colistin

resistance, tunisia, july 2015. *Eurosurveillance* **21,** 1–5 (2016).

28.   Tse, H. & Yuen, K.-Y. Dissemination of the mcr-1 colistin resistance gene. *Lancet Infect. Dis.*

3099, 3099 (2015).

29.   Figueiredo, R., Henriques, A., Sereno, R., Mendonça, N. & da Silva, G. J. Antimicrobial

Resistance and Extended-Spectrum β-Lactamases of Salmonella enterica Serotypes Isolated

from Livestock and Processed Food in Portugal: An Update. *Foodborne Pathog. Dis.* **12,** 110–

117 (2014).

30.   Haenni, M. *et al.* Co-occurrence of extended spectrum β lactamase and MCR-1 encoding

genes on plasmids. *Lancet Infect. Dis.* **3099,** 3099 (2016).

31.   Falgenhauer, L. *et al.* Chromosomal Locations of mcr-1 and bla(CTX-M-15) in Fluoroquinolone-

Resistant Escherichia coli ST410. *Emerg. Infect. Dis.* **22,** 1689–1691 (2016).

32.   Veldman, K. *et al.* Location of colistin resistance gene mcr-1 in Enterobacteriaceae from

livestock and meat. *J. Antimicrob. Chemother.* **71,** 2340–2342 (2016).

33.   Lilic, M., Jovanovic, M., Jovanovic, G. & Savic, D. J. Identification of the CysB-regulated gene,

hslJ, related to the Escherichia coli novobiocin resistance phenotype. *FEMS Microbiol. Lett.*

224, 239–246 (2003).

34. Baranova, N. & Nikaido, H. The BaeSR Two-Component Regulatory System Activates

Transcription of the yegMNOB (mdtABCD) Transporter Gene Cluster in Escherichia coli and

Increases Its Resistance to Novobiocin and Deoxycholate. *J. Bacteriol.* **184,** 4168–4176 (2002).

35. Stoesser, N., Mathers, A. J., Moore, C. E., Day, N. P. & Crook, D. W. Colistin resistance gene

mcr-1 and pHNSHP45 plasmid in human isolates of Escherichia coli and Klebsiella

pneumoniae. *Lancet Infect. Dis.* **3099,** 3099 (2016).

36. Falgenhauer, L. *et al.* Colistin resistance gene mcr-1 in extended-spectrum β-lactamase-

producing and carbapenemase-producing Gram-negative bacteria in Germany. *Lancet Infect.*

Dis. **3099,** 3099 (2016).

37. Snesrud, E. *et al.* Analysis of Serial Isolates of mcr-1- positive Escherichia coli Reveals a Highly

Active ISApl1 Transposon. *Antimicrob. Agents Chemother.* (2017). doi:10.1128/AAC.00056-17

38. Sun, J. *et al.* Genetic Analysis of the IncX4 Plasmids: Implications for a Unique Pattern in the

mcr-1 Acquisition. *Sci. Rep.* **7,** 424 (2017).

39. FDA. *2015 Summary Report On Antimicrobials Sold or Distributed for Use in Food-Producing*

Animals. (2016). at

<https://www.fda.gov/downloads/ForIndustry/UserFees/AnimalDrugUserFeeActADUFA/UCM

534243.pdf>

40. European Medicines Agency. *Updated advice on the use of colistin products in animals within*

the European Union: development of resistance and possible impact on human and animal

health. **44,** (2016).

41. Arcilla, M. S. *et al.* Import and spread of extended-spectrum beta-lactamase-producing

Enterobacteriaceae by international travellers (COMBAT study): a prospective, multicentre

cohort study. *Lancet Infect. Dis.* **17,** 78–85 (2017).

42. Larsen, M. V. *et al.* Multilocus sequence typing of total-genome-sequenced bacteria. *J. Clin.*

Microbiol. **50,** 1355–1361 (2012).

43.   Wirth, T. *et al.* Sex and virulence in Escherichia coli: An evolutionary perspective. *Mol.*

Microbiol. **60,** 1136–1151 (2006).

44.   Francisco, A. P., Bugalho, M., Ramirez, M. & Carriço, J. a. Global optimal eBURST analysis of

multilocus typing data using a graphic matroid approach. *BMC Bioinformatics* **10,** 152 (2009).

45.   Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for genome

assemblies. *Bioinformatics* **29,** 1072–1075 (2013).

46.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence

data. *Bioinformatics* **30,** 2114–2120 (2014).

47.   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications to

Single-Cell Sequencing. *J. Comput. Biol.* **19,** 455–477 (2012).

48.   Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer weighting

and repeat separation. *bioRxiv* (2017). at

<http://biorxiv.org/content/early/2017/01/04/071282.abstract>

49.   Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30,** 2068–2069

(2014).

50.   Page, A. J. *et al.* Roary: Rapid large-scale prokaryote pan genome analysis. *Bioinformatics* **31,**

3691–3693 (2015).

51.   Katoh, K. & Toh, H. Recent developments in the MAFFT multiple sequence alignment

program. *Brief. Bioinform.* **9,** 286–298 (2008).

52.   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30,** 1312–1313 (2014).

53.   Letunic, I. & Bork, P. Interactive tree of life (iTOL) v3: an online tool for the display and

annotation of phylogenetic and other trees. *Nucleic Acids Res.* **44,** W242–W245 (2016).

54.   Carattoli, A. *et al.* In Silico Detection and Typing of Plasmids using PlasmidFinder and Plasmid

Multilocus Sequence Typing. *Antimicrob. Agents Chemother.* **58,** 3895–3903 (2014).

55.   Alikhan, N.-F., Petty, N. K., Ben Zakour, N. L. & Beatson, S. A. BLAST Ring Image Generator

(BRIG): simple prokaryote genome comparisons. *BMC Genomics* **12,** 402 (2011).

56.   Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J. Antimicrob.*

Chemother. **67,** 2640–2644 (2012).

57.   McArthur, A. G. & Wright, G. D. Bioinformatics of antimicrobial resistance in the age of

molecular epidemiology. *Curr. Opin. Microbiol.* **27,** 45–50 (2015).

Acknowledgments

The authors would like to thank the team of the Department of Genome analysis of the AMC (in particular Dr. F. Baas and L. Koster) for their support in the sequencing bacterial genomes.

Financial support for this study was provided by the European COMPARE project (http://www.compare-europe.eu/) under the European Union's Horizon 2020 research and innovation programme, grant agreement No. 643476.

The VIBRE and COMBAT studies were supported by The Netherlands Organisation for Health Research and Development/The Netherlands Organisation for Scientific Research (ZonMw) under grants number 205100012 and 50-51700-98-120 respectively.

Additional information

Author contribution statement

S.M. and J.v.H. contributed equally to this work. S.M and JvH performed the systematic review, performed the experiments, interpreted the results and wrote the manuscript. N.W. contributed to bio-informatics analyses. M.A., D.M., J.P., M.d.J., the COMBAT consortium members, N.V., N.T.H. and C.S. contributed isolates and genome sequence data and associated meta data. M.d.J. and C.S. designed the study. All authors contributed to the writing of the manuscript.

Competing financial interests

The author(s) declare no competing financial interests.

The COMBAT consortium (in alphabetical order)

Martin C.J. Bootsma[8], Perry J. van Genderen[9], Abraham Goorhuis[10], Martin P. Grobusch[10], Nicky Molhoek[9], Astrid M.L. Oude Lashof[3], Ellen E. Stobberingh[3], Henri A. Verbrugh[2].

Julius Centre for Health Sciences and Primary Care, University Medical Centre Utrecht, Utrecht, the Netherlands and Department of Mathematics, Faculty of Science, Utrecht University, Utrecht, the Netherlands[8]. Department of Internal Medicine, Havenziekenhuis - Institute for Tropical Diseases, Rotterdam, the Netherlands[9]. Center of Tropical Medicine and Travel Medicine, Academic Medical Centre (AMC), Amsterdam, the Netherlands[10].

Figure legends

Figure 1: Maximum-likelihood tree based on concatenated core genome sequences of 65 *mcr-1*-carrying *E. coli* isolates.

Branch colours indicate phylogenetic clusters as determined by BAPS. Isolates from ST10, ST165 and closely related isolates are all grouped in the BAPS cluster 2 (dark blue). Leaf colours (isolates identifiers) indicate geographical region of origin. Isolation source is indicated in brackets: A = animal or meat; H = human; E = environment. The 6 travellers' isolates that were sequenced for this study are highlighted in bold and names start with CBT. Tree scale in number of substitutions per site.

*number of isolates.

Figure 2: Phylogeny of the *mcr-1*-carrying *E. coli* isolates. Maximum-likelihood tree based on concatenated MLST gene sequences, mid-point rooted.

Inner coloured circle: isolation source; outer circle: region of origin. Stars indicate the isolates from which a whole genome sequence was available. The 6 travellers' isolates that were sequenced for this study are highlighted in green. Bootstrap values between 0.9 and 1 are indicated by red triangles (size proportional to bootstrap value). The blue branches represent the main clade of 128 isolates including most isolates from ST10.Tree scale in number of substitutions per site.

See Supplementary Figure 4A for additional information on the relationship between STs, eBURST clustering and WGS BAPS clustering.

Figure 3: Alignment of *mcr-1*-containing plasmids and contigs.

Panel A: IncI2 plasmids (n = 29); panel B: IncX4 (n = 24); panel C: IncHI2 (n = 9).

Black outer ring: plasmid used as reference for the alignment; name and size of the reference indicated in the middle of each panel. Plasmid names followed by "_mcr1_contig" refer to assembled contigs from whole genome sequences. Other names refer to plasmid sequences deposited in online databases. The *mcr-1* gene and ISA-pl1 location are underlined in red.

Plasmids indicated with an asterisk are from the 6 travellers' isolates that were sequenced for this study.

Panel C: Putative MDR cassette is highlighted in orange.

	IncF	IncFI	IncFIB	IncFII	IncHI1	IncHI1A/ IncHI1B/ IncFIA	IncHI2	IncI2	IncI2/ IncX4	IncP	IncX3/ IncX4	IncX4	IncY	repB (p0111)	Total plasmids		Chromosome
Africa	0	0	1	0	0	0	4	8	0	0	0	1	0	0	**14**	6.4%	0
Asia	1	3	0	0	0	2	7	50	0	1	1	29	2	0	**96**	43.8%	2
Europe	0	0	2	1	1	0	33	5	0	3	0	44	0	1	**90**	41.1%	4
North-America	1	0	0	0	0	0	1	4	0	0	0	1	0	0	**7**	3.2%	0
Oceania	0	0	0	0	0	0	0	2	0	0	0	0	0	0	**2**	0.9%	0
South-America	0	0	0	0	0	0	0	7	1	0	0	2	0	0	**10**	4.6%	0
Total (% of Total)	**2** **(0.9)**	**3 (1.4)**	**3 (1.4)**	**1 (0.5)**	**1 (0.5)**	**2 (0.9)**	**45 (20.5)**	**76 (34.7)**	**1 (0.5)**	**4 (1.8)**	**1 (0.5)**	**77 (35.2)**	**2 (0.9)**	**1 (0.5)**	**219 (100)**	100.0%	6

Table 1: Incompatibility types of *mcr-1* carrying plasmids and distribution by geographical regions

BAPS cluster (branches)	Color key
1 (n* = 26)	
2 (n = 24)	
3 (n = 6)	
4 (n = 5)	
5 (n = 4)	

Geographical origin (leaves)	Color key
Asia (n* = 36)	
Europe (n = 20)	
North-America (n = 4)	
South-America (n = 4)	
Africa (n = 1)	

Tree scale: 0.01

A: IncI2
(n = 29)

- 210205360_pSLy1
- 210221272_plasmid_pSLy21
- A31-12_plasmid_pA31-12
- ABC149_plasmid_pABC149
- Af23_plasmid_pAF23
- BA76_plasmid_pBA76
- BA77_plasmid_pBA77
- BB1291_mcr1_contig
- CBT0637_mcr1_contig
- ※ CBT1531_mcr1_contig
- ※ CT31C.G1_mcr1_contig
- E15017_plasmid_pE15017
- EC13_plasmid_pEC13-1
- EC249_mcr1_contig
- EC5_plasmid_pEC5-1
- EG0430_plasmid_pEG430-1
- ESBL20150072_mcr1_contig
- JS-61_plasmid_pECJS-61-63
- mcr1_213_payne_mcr1_contig
- MT02C.A5_mcr1_contig
- PO169_mcr1_contig
- S2.14_plasmid_pS2.14-2
- S3_mcr1_contig
- SC23_plasmid_pSCS23
- SZ02_plasmid_pmcr1_IncI2
- VT55363_plasmid_pVT553
- WCH1410_plasmid_pMCR_1410
- WF-5-19C_plasmid_pWF-5-19C_mcr-1

B: IncX4
(n = 24)

- 014042624_mcr1_contig
- 0412016126_mcr1_contig
- 0412044854_mcr1_contig
- 0413040864_mcr1_contig
- 2012CEB2196SAL_mcr1_contig
- 2012CEB4337SAL_mcr1_contig
- A48_plasmid_pAF48
- ※ CBT1454_mcr1_contig
- ColR-40_mcr1_contig
- E15004_plasmid_pE15004
- EC702_mcr1_contig
- ECMY3_plasmid_pECMCR-1101
- Eco8314_mcr1_contig
- ENV-187_plasmid
- GD-8_plasmid_pECGD-8-33
- ICBEC72H_plasmid_pICBEC72Hmcr
- KP-6884_plasmid_pMCR1.2-IT
- mcr1_204_mcr1_contig
- MCR1_NJ_mcr1_contig
- OW3E1_plasmid_pOW3E1
- RL145_mcr1_contig
- RL158_mcr1_contig
- SZ04_plasmid_pmcr1

C: IncHI2
(n = 9)

- ※ CBT1023_mcr1_contig
- CG10C.C1_mcr1_contig
- CT41C.G2_mcr1_contig
- JS-59_plasmid_pECJS-59-244
- Sal25_plasmid
- MT31F.G1_mcr1_contig
- S38_plasmid_pS38
- S38_plasmid_pS38

Isolation source	Color key
Animal (n = 206)	
Human (n = 101)	
Environment (n = 5)	
Geographical origin	
Asia (n = 141)	
Europe (n = 125)	
Africa (n = 26)	
South-America (n = 12)	
North-America (n = 6)	
Oceania (n = 2)	

Tree scale: 0.001