Systematic genetic dissection of chitin degradation and uptake in *Vibrio cholerae*

Chelsea A. Hayes, Triana N. Dalia, and Ankur B. Dalia*

Department of Biology, Indiana University, Bloomington, Indiana, USA

*Author for correspondence: Ankur B. Dalia, email: ankdalia@indiana.edu, Tel: (812) 856-1895; Fax: (812) 855-6705.

Running Title: Genetic dissection of chitinases in *V. cholerae*

SUMMARY

Vibrio cholerae is a natural resident of the aquatic environment, where a common nutrient is the chitinous exoskeletons of microscopic crustaceans. Chitin utilization requires chitinases, which degrade this insoluble polymer into soluble chitin oligosaccharides. These oligosaccharides also serve as an inducing cue for natural transformation in *Vibrio* species. There are 7 predicted chitinase-like genes in the *V. cholerae* genome. Here, we systematically dissect the contribution of each gene to growth on chitin as well as induction of natural transformation. Specifically, we created a strain that lacks all 7 putative chitinases and from this strain, generated a panel of strains where each expresses a single chitinase. We also generated expression plasmids to ectopically express all 7 chitinases in our chitinase deficient strain. Through this analysis, we found that low levels of chitinase activity are sufficient for natural transformation, while growth on insoluble chitin as a sole carbon source requires more robust and concerted chitinase activity. We also assessed the role that the three uptake systems for the chitin degradation products GlcNAc, $(GlcNAc)_2$, and $(GlcN)_2$, play in chitin utilization and competence induction. Cumulatively, this study provides mechanistic details for how this pathogen utilizes chitin to thrive and evolve in its environmental reservoir.

INTRODUCTION

The cholera pathogen, *Vibrio cholerae*, is a natural resident of the aquatic environment. In

this niche, this bacterium forms biofilms on the chitinous shells of crustacean zooplankton.

These chitin biofilms are important for the water-borne transmission of cholera (Colwell et

al., 2003). Also, chitin, an insoluble polymer of β1,4-linked N-acetylglucosamine (GlcNAc),

serves as an important carbon and nitrogen source for *V. cholerae* in the environment (Huq

et al., 1983). To utilize this carbon source, this pathogen must degrade chitin into soluble

oligosaccharides via the action of chitinases. Subsequently, these chitin oligosaccharides

are transported across the outer membrane and into the periplasm via a chitoporin and

further broken down into mono- and di-saccharides, which can be transported across the

inner membrane by specific transporters (Meibom et al., 2004; Hunt et al., 2008).

Chitin oligosaccharides also induce the genes required for natural transformation in *V.*

cholerae, a physiological state in which cells can take up exogenous DNA and integrate it

into their chromosome by homologous recombination (Meibom et al., 2005). Therefore, the

interaction of *V. cholerae* with chitin is important for the survival and evolution of this

pathogen in its environmental reservoir as well as transmission to its human host. The

chitin utilization genes of *V. cholerae* have been identified by homology as well as by

identifying genes induced in the presence of chitin oligosaccharides (Meibom et al., 2004;

Hunt et al., 2008). For degradation, *V. cholerae* encodes 7 putative chitinase genes. These

are ChiA1 (VC1952), ChiA2 (VCA0027), VC0769, VCA0700, VC1073, VCA0140, and GbpA

(VCA0811). ChiA1 and ChiA2 have previously been implicated as the major chitinases

required for chitin degradation (Meibom et al., 2004; Watve et al., 2015; Dalia, 2016).

VC0769 and VC1073 are predicted chitinases, however, their role in chitin degradation in

V. cholerae has not formally been tested. VCA0700 is a predicted periplasmic

chitodextrinase, which further degrades soluble chitin oligosaccharides (Keyhani and

Roseman, 1996b). VCA0140 encodes a predicted spindolin-related protein, however, this

gene also contains a predicted chitin-binding domain and was therefore included as a

putative chitinase. Finally, GbpA is a GlcNAc binding protein, however, it is also predicted

to contain lytic polysaccharide monooxygenase activity (Loose et al., 2014). Chitinases have

been shown to function cooperatively in other chitinolytic organisms to promote chitin

degradation (Suzuki et al., 2002). However, a systematic analysis of chitinases has not been

performed in *V. cholerae* to assess the possibility of synergy among these enzymes or the

relative contribution of each to chitin-dependent growth and induction of natural

transformation.

Here, we systematically dissect the genes required for chitin degradation and uptake via

multiplex genome editing by natural transformation (MuGENT) (Dalia et al., 2014b). This

analysis has uncovered the chitinases that are necessary and sufficient for both chitin

utilization and chitin-induced natural transformation in *V. cholerae*.

RESULTS

Single mutants reveal that ChiA2 is critical for growth on chitin and chitin-induced natural

transformation

First, we assessed the role of each putative chitinase during growth on chitin as a sole

carbon source and chitin-induced natural transformation in single mutant strains. We find

that ChiA2 is important for both growth on chitin and chitin-induced natural

transformation (**Fig. 1A** and **1B**). This is consistent with recent reports on the importance

of ChiA2 for chitin-induced natural transformation (Mondal and Chatterjee, 2016). WT

levels of growth on chitin also required the periplasmic chitodextrinase VCA0700 (**Fig. 1A**),

however, this gene was dispensable for chitin-induced natural transformation (**Fig. 1B**).

Interestingly, while chitin-induced natural transformation was reduced in mutants lacking

ChiA2, it was not as deficient as a mutant lacking *pilA*, which fails to make the pilus

required for uptake of exogenous DNA (Seitz and Blokesch, 2013). Since soluble chitin

oligosaccharides generated by chitinases are required to induce natural competence

(Meibom et al., 2005), these data suggest that other chitinases may be able to support a low

level of chitin-induced natural transformation in the absence of ChiA2.

MuGENT for systematic genetic dissection of chitinases

While our data above highlighted that ChiA2 is important for chitin degradation, we still

observed chitin–dependent natural transformation in the absence of this enzyme. To

determine if other chitinases function cooperatively and/or in the absence of ChiA2, we

decided to generate a strain where all 7 chitinase-like genes were inactivated. To

accomplish this, we used a method we previously developed called MuGENT. This method

allows for making multiple scarless mutations simultaneously in a single step (Dalia et al.,

2014b; Hayes et al., 2017). The 7 chitinase-like genes are spread throughout both

chromosomes (**Fig. 2A**), and were targeted for inactivation by generating out-of-frame

~500bp deletions in the 5' end of each gene. Using this approach, we rapidly generated a

strain lacking all 7 putative chitinases, which we refer to as Δ7 henceforth (**Fig. 2B**). We

then created a panel of strains where each expresses only a single chitinase (i.e. are Δ6) by

systematically reverting one putative chitinase gene in each strain (**Fig. 2B**).

MuGENT of chitinases reveals that ChiA2 is sufficient for chitin-induced natural

transformation but not growth on chitin as a sole carbon source

First, we assessed our Δ7 strain for growth on chitin and chitin-induced natural

transformation. As expected, we found that this strain grew poorly on chitin (**Fig. 3A**). Also,

we find that the Δ7 strain is significantly reduced for chitin-induced natural transformation

(at the limit of detection) compared to a ChiA2 single mutant or a ChiA1 ChiA2 double

mutant (**Fig. 1B and 3B**). This is consistent with the other chitinases playing a minor role

in promoting chitin-induced natural transformation in the absence of ChiA2. Chitin

induction for natural transformation can be bypassed by overexpression of TfoX, the

master regulator of competence (Meibom et al., 2005). To confirm that natural

transformation was only attenuated in our Δ7 strain as a result of reduced chitinase

activity, we ectopically expressed TfoX in this mutant and tested natural transformation in

a chitin-independent assay. Under these conditions, as expected, we found that the Δ7

strain was as transformable as the WT (**Fig. 3C**). Also, we would predict that the Δ7 strain

would still be capable of growing on chitin degradation products. Consistent with this, we

find that this strain grows as well as the wildtype on chitobiose and the chitin monomer

GlcNAc (**Fig. S1**).

Next, we tested our panel of Δ6 mutants (each expressing a single chitinase) to determine if

any chitinase was sufficient for growth on chitin and chitin-induced natural transformation.

While no chitinase could independently support growth on chitin, the strain with just

ChiA2 could support chitin-induced natural transformation at near WT levels (**Fig. 3A** and

3B). While ChiA2 could restore chitin-induced natural transformation, none of the other

chitinases could promote this activity. These results indicate that growth on chitin and

chitin-induced natural transformation are separable phenomena since a strain only

expressing ChiA2 supported high levels of natural transformation but did not grow on

chitin as a sole carbon source. Thus, ChiA2 is both necessary and sufficient for chitin-

induced natural transformation.

One reason why ChiA2 may be the most important chitinase in *V. cholerae* is if this gene is

simply the most highly expressed chitinase under the conditions tested. Previous work has

uncovered the most highly upregulated chitinases through microarray analysis (Meibom et

al., 2004), however, these studies do not provide insight into the relative expression level

of each of the seven predicted chitinases. To assess this, we performed RNA-seq on

wildtype bacteria grown in the presence or absence of chitin hexasaccharide (GlcNAc)$_6$ to

induce the expression of chitin-regulated genes. As in prior studies (Meibom et al., 2004),

lactate was provided in both conditions as a chitin-independent carbon source. Indeed, we

find that ChiA2 is the most highly expressed chitinase under these conditions (**Fig. 4A**).

Thus, a trivial explanation for the relative importance of ChiA2 in *V. cholerae* might be that

this chitinase is the only one expressed at the levels required for efficient liberation of

chitin oligosaccharides. To test this further, we bypassed the native regulation of these

chitinases by ectopically expressing each in pMMB67EH (abbreviated pMMB), an IPTG-

inducible P$_{tac}$ expression vector that supports high levels of gene expression (Furste et al.,

1986). As expected, ectopic expression of ChiA2 restored wildtype levels of growth on

chitin to a ChiA2 single mutant (**Fig. 4B**). Ectopic expression of ChiA1, VC0769, and

VCA0700 also restored growth on chitin to a ChiA2 single mutant; however, this was not to

wildtype levels (**Fig. 4B**). Thus, this result suggests that ChiA2 expression levels alone

cannot fully account for the importance of this chitinase during growth on chitin and chitin-

induced natural transformation.

Next, we ectopically expressed each chitinase in our Δ7 mutant. This analysis uncovered

that none of these chitinases, even when overexpressed, could independently support

growth on chitin (**Fig. 4C**). However, five of these genes (*chiA1*, *chiA2*, VC0769, VCA0700, and VC1073), did enhance chitin-induced natural transformation when overexpressed (**Fig. 4D**). Together, these results suggest that low levels of chitinase activity may be sufficient to promote chitin-induced natural transformation, while robust (and possible concerted chitinase activity) is required for efficient chitin utilization. Furthermore, since chitin-induced natural transformation requires soluble chitin oligosaccharides, this analysis suggests that VC0769, VCA0700, and VC1073 possess *bona fide* chitinase activity (**Fig. 4D**), however, since they could not recover wildtype levels of growth on chitin to a *chiA2* single mutant (**Fig. 4B**), this suggests that they may have substantially lower chitinase activity than ChiA2. Conversely, VCA0140 and GbpA did not promote chitin-induced natural transformation even when overexpressed in the Δ7 background, suggesting that these genes cannot independently liberate soluble chitin oligosaccharides (**Fig. 4D**).

To confirm that chitinases were expressed to similar levels in our pMMB constructs, we generated variants of each expression vector where each chitinase had a C-terminal FLAG tag. These FLAG-tagged constructs were functional as determined by their ability to induce natural transformation in the Δ7 mutant (**Fig. S2**), and western blot analysis revealed that all 7 chitinases were secreted at similar levels when overexpressed (**Fig. 4E**). To confirm that protein in the supernatant in this experiment was the result of secretion and not cell lysis we detected the RNA polymerase alpha subunit (RpoA), which as expected, was found only in the cell pellet fraction. Thus, these results indicate that phenotypic differences observed among the distinct chitinases likely reflects differences in activity and not differences in expression / secretion.

ChiA2 works in conjunction with the periplasmic chitodextrinase VCA0700 to promote growth on chitin as a sole carbon source

Since ChiA2 was not independently sufficient to promote growth on chitin as a sole carbon source, we hypothesized that it may work in conjunction with another chitinase to promote robust chitin degradation and utilization. We hypothesized that the chitodextrinase VCA0700 would be a likely candidate for two reasons. One, when characterizing the phenotypes of single mutants, we found that loss of VCA0700 resulted in reduced growth

on chitin. Second, VCA0700 is predicted to be found at a distinct subcellular localization

and we hypothesized that the concerted action of the extracellular chitinase ChiA2 and the

predicted periplasmic chitodextrinase VCA0700 might be required for efficient chitin

utilization. To test this, we took the Δ6 strain that only expressed VCA0700 and

systematically knocked back in each of the other 6 chitinases to generate a panel of Δ5

strains where each is expressing 2 chitinases (one being VCA0700). When testing this panel

for growth on chitin, we find that the strain that contains VCA0700 and ChiA2 is capable of

wildtype levels of growth on chitin (**Fig. 5A**). Also, this strain displays wildtype levels of

chitin-induced natural transformation (**Fig. 5B**). To determine if VCA0700 could work in

conjunction with any of the other chitinases to mediate growth on chitin, we ectopically

expressed each chitinase in the Δ6 strain that only encodes VCA0700. We found that only

ChiA2 could promote robust growth in this background (**Fig. 5C**). Cumulatively, these

results indicate that ChiA2 and VCA0700 work together to efficiently degrade insoluble

chitin in *V. cholerae*.

Dissecting the role of chitin transporters during growth on chitin and chitin-induced natural

transformation

Once liberated from insoluble chitin via chitinase activity, chitin oligosaccharides are

subsequently transported into the periplasm through the action of a chitoporin (encoded

by VC0972) (Keyhani et al., 2000; Meibom et al., 2004). These oligosaccharides must then

be degraded into mono- or di-saccharides that can be transported across the inner

membrane. These are the monosaccharide GlcNAc and the disaccharides $(GlcNAc)_2$ (i.e.

chitobiose) and $(GlcN)_2$ (i.e. the unacetylated chitin disaccharide). GlcNAc and $(GlcN)_2$ are

transported via the action of two distinct PEP-dependent phosphotransferase system

transporters (VC0995 and VC1282, respectively), while $(GlcNAc)_2$ is transported via an

ABC transporter (permease encoded by VC0618 and VC0619) (Meibom et al., 2004; Hunt et

al., 2008). To assess the role of each of these transporters during growth on chitin and

chitin-induced natural transformation, we generated a panel of mutants lacking all possible

combinations of the three inner membrane transporters. We also generated a strain lacking

the outer membrane chitoporin. For growth on chitin, we find that the chitoporin VC0972

is required for wildtype levels of growth, which is consistent with previous reports (**Fig.**

6A) (Meibom et al., 2004). Also, while any one inner membrane transporter is dispensable,

loss of both the GlcNAc and (GlcNAc)$_2$ transporters resulted in lack of growth on chitin (**Fig.**

6A). Thus, this suggests that chitin is efficiently broken down to GlcNAc and (GlcNAc)$_2$,

while formation of (GlcN)$_2$ is less efficient. Indeed, in some sources of chitin, there is only 1

GlcN residue for every 6 GlcNAc residues (Meibom et al., 2004). While *V. cholerae* does

encode a putative chitin deacetylase (VC1280) adjacent to the locus required for (GlcN)$_2$

uptake, the activity or expression of this enzyme must not support robust growth on chitin

as a sole carbon source (Meibom et al., 2004; Hunt et al., 2008). Among this panel of

mutants, reduced growth on chitin as a sole carbon source directly correlated with reduced

rates of chitin-induced natural transformation (**Fig. 6A and 6B**), which suggests that

reduced rates of transformation among transporter mutants may largely be due to an

inability to grow under competence-inducing conditions.

DISCUSSION

These results suggest that ChiA2 and VCA0700 work synergistically to promote efficient

degradation of chitin in *V. cholera*e for both growth on this carbon source and induction of

natural transformation (**Fig. 7**). ChiA2 likely works extracellularly to generate chitin

oligosaccharides, while the chitodextrinase VCA0700 may work in the periplasm to

degrade these oligosaccharides further for uptake through the inner membrane. ChiA2 is

the most important extracellular chitinase under the conditions used here, since this

enzyme is necessary for growth on chitin as a sole carbon source and for chitin-induced

natural transformation. Based on expression levels, ChiA2 is the most highly expressed

chitinase in *V. cholerae*. Ectopic overexpression of other chitinases, however, did not

complement a ChiA2 mutant, which suggests that the activity (and not just expression

level) of ChiA2 may be important for the chitinolytic activity of *V. cholerae* under the

conditions tested. The importance of VCA0700 for growth on chitin is consistent with

previous reports, which suggested that the periplasmic steps of chitin degradation are

limiting for chitin utilization (Bassler et al., 1991). Loss of VCA0700 alone, however, did not

result in complete loss of growth on chitin. This suggests that in the VCA0700 mutant, the

other chitinases may work together to cleave chitin into products that can be taken up

without the need for chitodextrinase activity, albeit less efficiently.

There are a number of reasons why degradation of chitin in two stages, one extracellular

and one periplasmic, may be beneficial to the organism. One, relatively few microorganisms

can take up long chitin oligosaccharides from the extracellular environment, while many

microbes can take up chitin-derived mono- and di-saccharides. *Vibrio* species encode a

specific chitoporin (VC0972 in *V. cholerae*) to transport oligosaccharides across the outer

membrane, which could provide a competitive advantage in the environment (Suginta et

al., 2013). A second benefit to this spatially segregated degradation is that chitin

oligosaccharides serve as an important cue in the periplasm to signal upregulation of the

chitin utilization regulon (Keyhani and Roseman, 1996a; Li and Roseman, 2004) and genes

required for natural competence (Meibom et al., 2004; Dalia et al., 2014a). Thus, it is

beneficial to take up long chain oligosaccharides into the periplasm to serve as an inducing

cue prior to degradation for uptake and catabolism. Surprisingly, our analysis of VCA0700

indicated that when this chitodextrinase is ectopically expressed, some of this protein may

be secreted to the extracellular milieu and this is not a consequence of cell lysis (**Fig. 4E**).

Indeed, our results also indicate that VCA0700 can act extracellularly since overexpression

of this chitinase supported chitin-induced natural transformation in our Δ7 strain. Thus,

the concentrated action of ChiA2 and VCA0700 may be spatially segregated (extracellular

ChiA2 and periplasmic VCA0700) as previously hypothesized or it is possible that both of

these chitinases function extracellularly to efficiently degrade insoluble chitin into soluble

oligosaccharides. Further analysis of ChiA2 and VCA0700 localization and activity will shed

light on this question, which will be the focus of future work.

Induction of chitin-induced natural transformation requires ChiA2, while periplasmic

degradation via the chitobextrinase VCA0700 was largely dispensable. It was previously

shown that VCA0700 lacks detectable activity on insoluble chitin; however, it has robust

activity on soluble long chain chitin oligosaccharides (Keyhani and Roseman, 1996b). Our

previous work has shown that longer chains of chitin oligosaccharides are optimal at

inducing the activity of the chitin sensor TfoS, which is required for competence induction

(Dalia et al., 2014a). Thus, VCA0700 activity may not be required for competence induction

since this enzyme largely acts to reduce oligosaccharide chain length while playing a

limited role in liberating long chitin oligosaccharides from insoluble chitin.

Mutational analysis of the chitin transporters revealed that the outer membrane chitoporin

was important for growth on chitin as a sole carbon source and for competence induction.

The inner membrane GlcNAc and $(GlcNAc)_2$ transporters on the other hand were

genetically redundant for these activities (**Fig. 7**). Loss of chitin-induced natural

transformation in transporter mutants directly correlated with the reduced ability of these

strains to grow on chitin as a sole carbon source. A strain that only expresses the chitinase

ChiA2 (i.e. a Δ6 strain), however, is not able to grow on chitin, while it displays high rates of

natural transformation. Also, overexpression of 5 of the 7 predicted chitinases in the Δ7

strain supported natural transformation while none supported growth on chitin as a sole

carbon source. This suggests that induction of competence only requires relatively low

levels of chitinase activity and by extension only small amounts of chitin oligosaccharides,

while growth on chitin may require more robust and concerted chitinase activity. As

mentioned above, TfoS, the membrane-embedded chitin sensor required for natural

transformation, is induced by long chain chitin oligosaccharides (Dalia et al., 2014a). Thus,

loss of the chitoporin, which specifically imports long chain chitin oligosaccharides into the

periplasm (Suginta et al., 2013), may result in reduced rates of natural transformation as a

result of poor TfoS induction. Loss of natural transformation in the GlcNAc and $(GlcNAc)_2$

transporter double mutant would not be predicted to diminish TfoS induction, however,

loss of these uptake transporters may slow growth to a level where the competence

machinery is no longer efficiently expressed. Alternatively, it is possible that the

cytoplasmic chitin degradation products internalized by the GlcNAc and $(GlcNAc)_2$

transporters aid in competence induction. However, previous work has shown that

artificial activation of TfoS supports competence induction in rich medium even in the

absence of chitin (Dalia et al., 2014a; Dalia, 2016).

Chitin is the second most abundant biomolecule in nature (after cellulose) and represents

an abundant waste product of the seafood industry (Yan and Chen, 2015). The genes

defined here may represent the minimal gene set required for efficient chitin utilization,

which can be transferred to relevant non-chitinolytic microorganisms for biotech

applications. This will be a focus of future work.

In conclusion, this study systematically defines the chitinases and transporters that are

necessary and sufficient for chitin degradation and utilization in *V. cholerae*. Also, it

identifies the unique requirements for chitin-induced horizontal gene transfer by natural

transformation in this important human pathogen.

EXPERIMENTAL PROCEDURES

Bacterial strains and culture conditions

Strains were routinely grown in LB broth and on LB agar plates. When necessary, media

was supplemented with stremptomycin (100 µg/mL), spectinomycin (200 µg/mL),

kanamycin (50 µg/mL), trimethoprim (10 µg/mL), or carbenicillin (100 µg/mL). Strains in

this study are all derived from E7946 (Miller et al., 1989). All strains used in this study are

listed in **Table S1**.

For growth on chitin as the sole carbon source, we used M9 minimal medium (Difco)

supplemented with 30 µM $FeSO_4$ and ~1% chitin from shrimp shells (Sigma). To 1 mL of

M9+chitin medium, ~10^5 cells were added and grown for 48 hours at 30°C for each growth

reaction. Reactions were then plated for viable counts to assess growth. For reactions with

strains containing a pMMB plasmid, carbenicillin (20µg/mL) was added to M9+chitin

reactions to maintain the plasmid, and IPTG (100 µM) was added to induce expression.

Generating mutant strains and constructs

All mutants were generated by natural cotransformation and MuGENT as previously

described (Dalia et al., 2014b). Briefly, mutant constructs were generated by splicing-by-

overlap extension (SOE) PCR as previously described (Dalia et al., 2013). For

cotransformation and MuGENT, a selected product (i.e. one conferring resistance to an

antibiotic) was used as the transforming DNA in conjunction with an unselected product

(i.e. one that will confer the mutation of interest). By selecting for integration of the

selected product, we increase the likelihood that the unselected mutation will have

integrated into cells within a competent population. We then screen for the mutation of

interest in these cells by multiplex allele specific colony PCR (MASC-PCR) exactly as

previously described (Wang et al., 2009). All chitinase and transporter mutations were

generated using unselected SOE products. All expression constructs were generated by

traditional cloning and C-terminal FLAG tags were added by site-directed mutagenesis as

previously described (Edelheit et al., 2009). All primers used to generate mutant constructs

and plasmids are listed in **Table S2**.

Natural transformation assays

We tested chitin-induced natural transformation essentially as previously described (Dalia

et al., 2015). Briefly, $\sim10^8$ cells were incubated in a 1 mL reaction of instant ocean medium

(7 g/L; Aquarium Systems) containing $\sim$8 mg of chitin. Cells were incubated in this medium

at 30°C statically for 16-24 hours to induce competence. Then, transforming DNA (tDNA)

was added. For all natural transformation assays in this study, we used $\sim$500 ng of a PCR

product that would replace the frame-shifted transposase VC1807 with a trimethoprim

resistance cassette. Reactions were incubated for 5-24 hours with tDNA, and then 1 mL of

LB was added to outgrow reactions. The transformation efficiency was then determined by

plating reactions for viable counts on LB+Tm10 (transformants) and plain LB (total viable

counts).

For chitin-independent transformation assays, strains containing pMMB-tfoX were grown

overnight in LB with 100 µg/mL carbenicillin and 100 µM IPTG. Then, $\sim10^8$ cells were

diluted into instant ocean medium containing 100 µg/mL IPTG. Next, tDNA was added and

incubated statically at 30°C for 5-24 hours. Then, reactions were outgrown and plated as

described above to determine the transformation efficiency.

RNA-seq

RNA was prepped for sequencing on the Illumina platform exactly as previously described

(Shishkin et al., 2015). Reads obtained were mapped to the N16961 reference genome

(NC_002505 and NC_002506) and analyzed using the Tufts University Galaxy server (Afgan

et al., 2016). Reads were aggregated within ORFs and normalized for the size of the ORF to obtain normalized transcript abundance for each gene under each condition tested.

Western blot analysis

Western blots were conducted essentially as previously described (Burnette, 1981; Dalia et al., 2015). Briefly, samples were prepared for western blots by growing strains to mid-log in 20 μg/mL carbenicillin and 100μM IPTG. Cells were then spun and cell free supernatants were collected and boiled in SDS PAGE sample buffer. The cell pellets were then washed and resuspended with an equal volume of 0.5X IO and then boiled in SDS PAGE sample buffer. Samples were electrophertically separated on 10% polyacrylamide gels and transferred to a nitrocellulose membrane. FLAG-tagged proteins were probed using rabbit polyclonal α-FLAG antibodies (Sigma), while RpoA was probed using a mouse monoclonal antibody (Biolegend). Blots were developed using IRDye 800CW labeled α-rabbit or α-mouse secondary antibodies as appropriate and imaged using a LI-COR Imaging system.

ACKNOWLEDGEMENTS

We would like to thank Neil Greene for critical reading of this manuscript. Also, we thank the Malcolm Winkler and Julia van Kessel labs for reagents and advice. This work was supported by US National Institutes of Health Grant AI118863 and startup funds from the Indiana University College of Arts and Sciences to ABD.

REFERENCES

Afgan, E., Baker, D., van den Beek, M., Blankenberg, D., Bouvier, D., Cech, M. et al. (2016) The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2016 update. *Nucleic Acids Res* **44**: W3-W10.

Bassler, B.L., Yu, C., Lee, Y.C., and Roseman, S. (1991) Chitin utilization by marine bacteria. Degradation and catabolism of chitin oligosaccharides by Vibrio furnissii. *J Biol Chem* **266**: 24276-24286.

Burnette, W.N. (1981) "Western blotting": electrophoretic transfer of proteins from sodium dodecyl sulfate--polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein A. *Anal Biochem* **112**: 195-203.

Colwell, R.R., Huq, A., Islam, M.S., Aziz, K.M., Yunus, M., Khan, N.H. et al. (2003) Reduction of

cholera in Bangladeshi villages by simple filtration. *Proc Natl Acad Sci U S A* **100**: 1051-

1055.

Dalia, A.B. (2016) RpoS is required for natural transformation of Vibrio cholerae through

regulation of chitinases. *Environ Microbiol* **18**: 3758-3767.

Dalia, A.B., Lazinski, D.W., and Camilli, A. (2013) Characterization of undermethylated sites

in Vibrio cholerae. *J Bacteriol* **195**: 2389-2399.

Dalia, A.B., Lazinski, D.W., and Camilli, A. (2014a) Identification of a membrane-bound

transcriptional regulator that links chitin and natural competence in Vibrio cholerae. *MBio*

5: e01028-01013.

Dalia, A.B., McDonough, E., and Camilli, A. (2014b) Multiplex genome editing by natural

transformation. *Proc Natl Acad Sci U S A* **111**: 8937-8942.

Dalia, A.B., Seed, K.D., Calderwood, S.B., and Camilli, A. (2015) A globally distributed mobile

genetic element inhibits natural transformation of Vibrio cholerae. *Proc Natl Acad Sci U S A*

112: 10485-10490.

Edelheit, O., Hanukoglu, A., and Hanukoglu, I. (2009) Simple and efficient site-directed

mutagenesis using two single-primer reactions in parallel to generate mutants for protein

structure-function studies. *BMC Biotechnol* **9**: 61.

Furste, J.P., Pansegrau, W., Frank, R., Blocker, H., Scholz, P., Bagdasarian, M., and Lanka, E.

(1986) Molecular cloning of the plasmid RP4 primase region in a multi-host-range tacP

expression vector. *Gene* **48**: 119-131.

Hayes, C.A., Dalia, T.N., and Dalia, A.B. (2017) Systematic genetic dissection of PTS in Vibrio

cholerae uncovers a novel glucose transporter and a limited role for PTS during infection of

a mammalian host. *Mol Microbiol*.

Hunt, D.E., Gevers, D., Vahora, N.M., and Polz, M.F. (2008) Conservation of the chitin

utilization pathway in the Vibrionaceae. *Appl Environ Microbiol* **74**: 44-51.

Huq, A., Small, E.B., West, P.A., Huq, M.I., Rahman, R., and Colwell, R.R. (1983) Ecological

relationships between Vibrio cholerae and planktonic crustacean copepods. *Appl Environ*

Microbiol **45**: 275-283.

Keyhani, N.O., and Roseman, S. (1996a) The chitin catabolic cascade in the marine

bacterium Vibrio furnissii. Molecular cloning, isolation, and characterization of a

periplasmic beta-N-acetylglucosaminidase. *J Biol Chem* **271**: 33425-33432.

Keyhani, N.O., and Roseman, S. (1996b) The chitin catabolic cascade in the marine

bacterium Vibrio furnissii. Molecular cloning, isolation, and characterization of a

periplasmic chitodextrinase. *J Biol Chem* **271**: 33414-33424.

Keyhani, N.O., Li, X.B., and Roseman, S. (2000) Chitin catabolism in the marine bacterium

Vibrio furnissii. Identification and molecular cloning of a chitoporin. *J Biol Chem* **275**:

33068-33076.

Li, X., and Roseman, S. (2004) The chitinolytic cascade in Vibrios is regulated by chitin

oligosaccharides and a two-component chitin catabolic sensor/kinase. *Proc Natl Acad Sci U*

S A **101**: 627-631.

Loose, J.S., Forsberg, Z., Fraaije, M.W., Eijsink, V.G., and Vaaje-Kolstad, G. (2014) A rapid

quantitative activity assay shows that the Vibrio cholerae colonization factor GbpA is an

active lytic polysaccharide monooxygenase. *FEBS Lett* **588**: 3435-3440.

Meibom, K.L., Blokesch, M., Dolganov, N.A., Wu, C.Y., and Schoolnik, G.K. (2005) Chitin

induces natural competence in Vibrio cholerae. *Science* **310**: 1824-1827.

Meibom, K.L., Li, X.B., Nielsen, A.T., Wu, C.Y., Roseman, S., and Schoolnik, G.K. (2004) The

Vibrio cholerae chitin utilization program. *Proc Natl Acad Sci U S A* **101**: 2524-2529.

Miller, V.L., DiRita, V.J., and Mekalanos, J.J. (1989) Identification of toxS, a regulatory gene

whose product enhances toxR-mediated activation of the cholera toxin promoter. *J*

Bacteriol **171**: 1288-1293.

Mondal, M., and Chatterjee, N.S. (2016) Role of Vibrio cholerae exochitinase ChiA2 in

horizontal gene transfer. *Can J Microbiol* **62**: 201-209.

Seitz, P., and Blokesch, M. (2013) DNA-uptake machinery of naturally competent Vibrio

cholerae. *Proc Natl Acad Sci U S A* **110**: 17987-17992.

Shishkin, A.A., Giannoukos, G., Kucukural, A., Ciulla, D., Busby, M., Surka, C. et al. (2015)

Simultaneous generation of many RNA-seq libraries in a single reaction. *Nat Methods* **12**:

323-325.

Suginta, W., Chumjan, W., Mahendran, K.R., Schulte, A., and Winterhalter, M. (2013)

Chitoporin from Vibrio harveyi, a channel with exceptional sugar specificity. *J Biol Chem*

288: 11038-11046.

Suzuki, K., Sugawara, N., Suzuki, M., Uchiyama, T., Katouno, F., Nikaidou, N., and Watanabe,

T. (2002) Chitinases A, B, and C1 of Serratia marcescens 2170 produced by recombinant

Escherichia coli: enzymatic properties and synergism on chitin degradation. *Biosci*

Biotechnol Biochem **66**: 1075-1083.

Wang, H.H., Isaacs, F.J., Carr, P.A., Sun, Z.Z., Xu, G., Forest, C.R., and Church, G.M. (2009)

Programming cells by multiplex genome engineering and accelerated evolution. *Nature*

460: 894-898.

Watve, S.S., Thomas, J., and Hammer, B.K. (2015) CytR Is a Global Positive Regulator of

Competence, Type VI Secretion, and Chitinases in Vibrio cholerae. *PLoS One* **10**: e0138834.

Yan, N., and Chen, X. (2015) Sustainability: Don't waste seafood waste. *Nature* **524**: 155-

157.

FIGURE LEGENDS

Fig. 1 – *Characterizing chitinase single mutants for growth on chitin and natural*

transformation. (**A**) Growth of the indicated mutant strains in M9 minimal medium with

chitin as a sole carbon source. (**B**) Chitin-induced natural transformation of the indicated

mutant strains. All data are shown as the mean ± SD and are from at least 3 independent

biological replicates. *** = $p<0.001$.

Fig. 2 - *MuGENT for systematic inactivation of all 7 chitinase-like genes.* (**A**) Chromosomal

map of the location of the seven chitinases inactivated in this study. (**B**) MASC-PCR of the

indicated mutants. The presence of a band indicates that the gene indicated to the left is

inactivated, while the absence of a band indicates that this gene is intact.

Fig. 3 - *ChiA2 is sufficient for natural transformation, but not growth on chitin.* (**A**) Growth of

the indicated mutant strains in M9 medium with chitin as a sole carbon source. (**B**) Chitin-

induced natural transformation of the indicated mutant strains. (**C**) Chitin-independent

natural transformation assay of the indicated mutants. TfoX was induced in these

experiments with 100 μM IPTG. All data are shown as the mean ± SD and are from at least 3

independent biological replicates. * = $p<0.05$, *** = $p<0.001$, NS = not significant.

Fig. 4 – *Overexpression of single chitinases in a Δ7 strain recovers natural transformation but*

not growth on chitin. (**A**) Relative transcript abundance of the indicated genes from RNA-

seq data. (**B**) Growth of the indicated mutant strains in M9 medium containing chitin as a

sole carbon source. (**C**) Growth of the indicated mutant strains in M9 medium with chitin as

a sole carbon source. (**D**) Chitin-induced natural transformation of the indicated mutant

strains. Genes were induced in **B**, **C** and **D** with 100 μM IPTG. (**E**) Western blot analysis of

strains in the Δ7 background harboring a pMMB expression construct with the C-terminally

FLAG tagged chitinase indicated. Supernatant (S) and pellet (P) fractions were run for each

strain and probed with α-FLAG (top) and α-RpoA (bottom) antibodies. All data in **A-D** are

shown as the mean ± SD and are from at least 3 independent biological replicates. * =

$p<0.05$, ** = $p<0.01$, and *** = $p<0.001$, and NS = not significant.

Fig. 5 – *ChiA2 and VCA0700 are sufficient for growth on chitin*. (**A**) Growth of the indicated

mutant strains in M9 medium with chitin as a sole carbon source. (**B**) Chitin-induced

natural transformation of the indicated mutant strains. (**C**) Growth of the indicated strains

in M9 medium with chitin as the sole carbon source and 100 μM IPTG. All data are shown

as the mean ± SD and are from at least 3 independent biological replicates. *** = $p<0.001$,

NS = not significant.

Fig. 6 – *Role of chitin transporters for growth on chitin and chitin-induced natural*

transformation. (**A**) Growth of the indicated mutant strains in M9 medium with chitin as a

sole carbon source. (**B**) Chitin-induced natural transformation of the indicated mutant

strains. All data are shown as the mean ± SD and are from at least 3 independent biological

replicates. All statistical comparisons in **A** and **B** were made between the indicated mutant

and the WT. * = $p<0.05$, ** = $p<0.01$, and *** = $p<0.001$.

Fig. 7 – *Schematic of the chitin utilization pathway genetically dissected in this study.* First,

extracellular chitinases degrade insoluble chitin into soluble chitin oligosaccharides. While

ChiA2 is the dominant enzyme required for this process, the chitinases ChiA1, VC0769,

VC1073, and VCA0700 likely play some role. These soluble oligosaccharides are then taken

up across the outer membrane (OM) and into the periplasm via the chitoporin encoded by

VC0972. Then, these oligosaccharides are likely further broken down by the

chitodextrinase VCA0700 into $(GlcNAc)_2$ (aka chitobiose), GlcNAc, and $(GlcN)_2$, which are

taken up across the inner membrane (IM) into the cytoplasm by the transporters encoded

by VC0618-0619, VC0995, and VC1282, respectively. Our results indicate that for robust

growth on chitin, the transporters responsible for uptake of chitobiose and GlcNAc play the

largest role.

Fig. 1 – *Characterizing chitinase single mutants for growth on chitin and natural transformation*. (**A**) Growth of the indicated mutant strains in M9 minimal medium with chitin as a sole carbon source. (**B**) Chitin-induced natural transformation of the indicated mutant strains. All data are shown as the mean ± SD and are from at least 3 independent biological replicates. *** = $p < 0.001$.

Fig. 2 - *MuGENT for systematic inactivation of all 7 chitinase-like genes.* (**A**) Chromosomal map of the location of the seven chitinases inactivated in this study. (**B**) MASC-PCR of the indicated mutants. The presence of a band indicates that the gene indicated to the left is inactivated, while the absence of a band indicates that this gene is intact.

Fig. 3 - *ChiA2 is sufficient for natural transformation, but not growth on chitin.* (**A**) Growth of the indicated mutant strains in M9 medium with chitin as a sole carbon source. (**B**) Chitin-induced natural transformation of the indicated mutant strains. (**C**) Chitin-independent natural transformation assay of the indicated mutants. TfoX was induced in these experiments with 100 μM IPTG. All data are shown as the mean ± SD and are from at least 3 independent biological replicates. * = $p<0.05$, *** = $p<0.001$, NS = not significant.

Fig. 4 – *Overexpression of single chitinases in a Δ7 strain recovers natural transformation but not growth on chitin*. (**A**) Relative transcript abundance of the indicated genes from RNA-seq data. (**B**) Growth of the indicated mutant strains in M9 medium containing chitin as a sole carbon source. (**C**) Growth of the indicated mutant strains in M9 medium with chitin as a sole carbon source. (**D**) Chitin-induced natural transformation of the indicated mutant strains. Genes were induced in **B**, **C** and **D** with 100 µM IPTG. (**E**) Western blot analysis of strains in the Δ7 background harboring a pMMB expression construct with the C-terminally FLAG tagged chitinase indicated. Supernatant (S) and pellet (P) fractions were run for each strain and probed with α-FLAG (top) and α-RpoA (bottom) antibodies. All data in **A**-**D** are shown as the mean ± SD and are from at least 3 independent biological replicates. * = $p<0.05$, ** = $p<0.01$, and *** = $p<0.001$, and NS = not significant.

Fig. 5 – *ChiA2 and VCA0700 are sufficient for growth on chitin.* (**A**) Growth of the indicated mutant strains in M9 medium with chitin as a sole carbon source. (**B**) Chitin-induced natural transformation of the indicated mutant strains. (**C**) Growth of the indicated strains in M9 medium with chitin as the sole carbon source and 100 μM IPTG. All data are shown as the mean ± SD and are from at least 3 independent biological replicates. *** = $p<0.001$, NS = not significant.

Fig. 6 – *Role of chitin transporters for growth on chitin and chitin-induced natural transformation.* (**A**) Growth of the indicated mutant strains in M9 medium with chitin as a sole carbon source. (**B**) Chitin-induced natural transformation of the indicated mutant strains. All data are shown as the mean ± SD and are from at least 3 independent biological replicates. All statistical comparisons in **A** and **B** were made between the indicated mutant and the WT. * = $p<0.05$, ** = $p<0.01$, and *** = $p<0.001$.

Fig. 7 – *Schematic of the chitin utilization pathway genetically dissected in this study.* First, extracellular chitinases degrade insoluble chitin into soluble chitin oligosaccharides. While ChiA2 is the dominant enzyme required for this process, the chitinases ChiA1, VC0769, VC1073, and VCA0700 likely play some role. These soluble oligosaccharides are then taken up across the outer membrane (OM) and into the periplasm via the chitoporin encoded by VC0972. Then, these oligosaccharides are likely further broken down by the chitodextrinase VCA0700 into $(GlcNAc)_2$ (aka chitobiose), GlcNAc, and $(GlcN)_2$, which are taken up across the inner membrane (IM) into the cytoplasm by the transporters encoded by VC0618-0619, VC0995, and VC1282, respectively. Our results indicate that for robust growth on chitin, the transporters responsible for uptake of chitobiose and GlcNAc play the largest role.

Fig. S1 – *A chitinase deficient strain is still capable of growth on the chitin degradation products chitobiose and GlcNAc.* Growth curves of wildtype (black circles) and Δ7 chitinase strain (black squares) in M9 minimal medium supplemented with the carbon source indicated above each graph.

Fig. S2 – *C-terminally FLAG tagged chitinases are functional*. Natural transformation assay of the indicated strains. All strains were incubated on chitin with Carbeniciillin (20 µg/mL) and IPTG (100 µg/mL). Data are from at least three independent biological replicates and shown as the mean ± SD. * = p<0.05, ** = p<0.01, *** = p<0.001, and LOD = limit of detection.

Table S1 – *Strains used in this study*

Strain name in manuscript	Genotype and antibiotic resistances	Description	Reference / (strain#)
WT	E7946 Sm^R	Wildtype *V. cholerae* O1 El Tor strain used throughout this study – parent strain for all mutants indicated below	(Miller et al., 1989) (SAD030)
ΔChiA1	ΔChiA1	Wildtype with a 500bp deletion in the 5' end of the ChiA1 gene	This Study (CAH060 / SAD1333)
ΔChiA2	ΔChiA2	Wildtype with a 500bp deletion in the 5' end of the ChiA2 gene	This Study (CAH061 / SAD1334)
ΔVC0769	ΔVC0769	Wildtype with a 500bp deletion in the 5' end of the VC0769 gene	This Study (CAH062 / SAD1335)
ΔVCA0700	ΔVCA0700	Wildtype with a 500bp deletion in the 5' end of the VCA0700 gene	This Study (CAH064 / SAD1336)
ΔVC1073	ΔVC1073	Wildtype with a 500bp deletion in the 5' end of the VC1073 gene	This Study (CAH065 / SAD1337)
ΔVCA0140	ΔVCA0140	Wildtype with a 500bp deletion in the 5' end of the VCA0140 gene	This Study (CAH066 / SAD1338)
ΔGbpA	ΔGbpA	Wildtype with a 500bp deletion in the 5' end of the gbpA gene	This Study (CAH063 / SAD1339)
ΔChiA1 ΔChiA2	ΔChiA1 ΔChiA2	Wildtype with a 500bp deletion in the 5' end of the ChiA1 and ChiA2 genes	This Study (SAD863)
ΔpilA	ΔpilA::Spec^R	The pilA gene (VC2423) was deleted and replaced with a spectinomycin resistance cassette in the wildtype	This Study (SAD780)
WT pMMB	pMMB Carb^R	Wildtype with pMMB67EH empty vector	This Study (CAH298 / SAD1340)
ΔChiA2 pMMB	ΔChiA2 pMMB Carb^R	ΔChiA2 mutant with pMMB67EH empty vector	This Study (CAH458 / SAD1341)
ΔChiA2 pMMB-ChiA1	ΔChiA2 pMMB-ChiA1 Carb^R	ΔChiA2 mutant with pMMB67EH containing ChiA1 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH469 / SAD1342)
ΔChiA2 pMMB-ChiA2	ΔChiA2 pMMB-ChiA2 Carb^R	ΔChiA2 mutant with pMMB67EH containing ChiA2 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH463 / SAD1343)
ΔChiA2 pMMB-VC0769	ΔChiA2 pMMB-VC0769 Carb^R	ΔChiA2 mutant with pMMB67EH containing VC0769 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH464 / SAD1344)
ΔChiA2 pMMB-	ΔChiA2 pMMB-VCA0700 Carb^R	ΔChiA2 mutant with pMMB67EH	This Study

VCA0700		containing VCA0700 – cloned into the EcoRI and BamHI sites of the MCS.	(CAH465 / SAD1345)
ΔChiA2 pMMB-VC1073	ΔChiA2 pMMB-VC1073 Carb^R	ΔChiA2 mutant with pMMB67EH containing VC1073 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH467 / SAD1346)
ΔChiA2 pMMB-VCA0140	ΔChiA2 pMMB-VCA0140 Carb^R	ΔChiA2 mutant with pMMB67EH containing VCA0140 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH468 / SAD1347)
ΔChiA2 pMMB-GbpA	ΔChiA2 pMMB-GbpA Carb^R	ΔChiA2 mutant with pMMB67EH containing GbpA – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH466 / SAD1348)
Δ7	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R	All seven chitinases were inactivated by MuGENT. VC1807 was inactivated as the neutral locus throughout this process with a resistance cassette to serve as the selected product. VC1807 is a frame-shifted transposase.	This Study (CAH130 / SAD1349)
Δ6 (ChiA1 only)	ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R	The indicated chitinase was restored in the Δ7 mutant by cotransformation.	This Study (CAH165 / SAD1350)
Δ6 (ChiA2 only)	ΔChiA1, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R	The indicated chitinase was restored in the Δ7 mutant by cotransformation.	This Study (CAH166 / SAD1351)
Δ6 (VC0769 only)	ΔChiA1, ΔChiA2, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R	The indicated chitinase was restored in the Δ7 mutant by cotransformation.	This Study (CAH163 / SAD1352)
Δ6 (VCA0700 only)	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R	The indicated chitinase was restored in the Δ7 mutant by cotransformation.	This Study (CAH169 / SAD1353)
Δ6 (VC1073 only)	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R	The indicated chitinase was restored in the Δ7 mutant by cotransformation.	This Study (CAH164 / SAD1354)
Δ6 (VCA0140 only)	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔGbpA, ΔVC1807::Kan^R	The indicated chitinase was restored in the Δ7 mutant by cotransformation.	This Study (CAH167 / SAD1355)
Δ6 (GbpA only)	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔVC1807::Kan^R	The indicated chitinase was restored in the Δ7 mutant by cotransformation.	This Study (CAH168 / SAD1356)
WT pMMB-tfoX	pMMB-tfoX Carb^R	Wildtype with pMMB67EH containing the tfoX gene (VC1153) cloned into the EcoRI and BamHI sites of the MCS	This Study (SAD614)
Δ7 pMMB-tfoX	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-tfoX Carb^R	Δ7 mutant with pMMB67EH containing the tfoX gene (VC1153) cloned into the EcoRI and BamHI sites of the MCS	This Study (CAH126 / SAD1357)
Δ7 pMMB	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA,	Δ7 with pMMB67EH empty vector	This Study (CAH299 / SAD1358)

Strain	Genotype	Description	Reference
	ΔVC1807::Spec^R, pMMB Carb^R		
Δ7 pMMB-ChiA1	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-ChiA1 Carb^R	Δ7 mutant with pMMB67EH containing ChiA1 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH460 / SAD1359)
Δ7 pMMB-ChiA2	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-ChiA2 Carb^R	Δ7 mutant with pMMB67EH containing ChiA2 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH269 / SAD1360)
Δ7 pMMB-VC0769	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-VC0769 Carb^R	Δ7 mutant with pMMB67EH containing VC0769 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH270 / SAD1361)
Δ7 pMMB-VCA0700	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-VCA0700 Carb^R	Δ7 mutant with pMMB67EH containing VCA0700 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH271 / SAD1362)
Δ7 pMMB-VC1073	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-VC1073 Carb^R	Δ7 mutant with pMMB67EH containing VC1073 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH273 / SAD1363)
Δ7 pMMB-VCA0140	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-VCA0140 Carb^R	Δ7 mutant with pMMB67EH containing VCA0140 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH274 / SAD1364)
Δ7 pMMB-GbpA	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-GbpA Carb^R	Δ7 mutant with pMMB67EH containing GbpA – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH272 / SAD1365)
Δ5 (VCA0700 and ChiA1 only)	ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R	Revert the indicated chitinase into the Δ6 background that only has VCA0700	This Study (CAH191 / SAD1366)
Δ5 (VCA0700 and ChiA2 only)	ΔChiA1, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R	Revert the indicated chitinase into the Δ6 background that only has VCA0700	This Study (CAH197 / SAD1367)
Δ5 (VCA0700 and VC0769 only)	ΔChiA1, ΔChiA2, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R	Revert the indicated chitinase into the Δ6 background that only has VCA0700	This Study (CAH420 / SAD1368)
Δ5 (VCA0700 and VC1073 only)	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R	Revert the indicated chitinase into the Δ6 background that only has VCA0700	This Study (CAH421 / SAD1369)
Δ5 (VCA0700 and VCA0140 only)	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔGbpA, ΔVC1807::Spec^R	Revert the indicated chitinase into the Δ6 background that only has VCA0700	This Study (CAH422 / SAD1370)
Δ5 (VCA0700 and GbpA only)	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔVC1807::Spec^R	Revert the indicated chitinase into the Δ6 background that only has VCA0700	This Study (CAH423 / SAD1371)

Δ6 (VCA0700 only) pMMB	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R, pMMB Carb^R	Δ6 (VCA0700 only) mutant with pMMB67EH empty vector	This Study (CAH424 / SAD1372)
Δ6 (VCA0700 only) pMMB-ChiA1	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R, pMMB-ChiA1 Carb^R	Δ6 (VCA0700 only) mutant with with pMMB67EH containing ChiA1 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH462 / SAD1373)
Δ6 (VCA0700 only) pMMB-ChiA2	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R, pMMB-ChiA2 Carb^R	Δ6 (VCA0700 only) mutant with with pMMB67EH containing ChiA2 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH425 / SAD1374)
Δ6 (VCA0700 only) pMMB-VC0769	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R, pMMB-VC0769 Carb^R	Δ6 (VCA0700 only) mutant with with pMMB67EH containing VC0769 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH426 / SAD1375)
Δ6 (VCA0700 only) pMMB-VCA0700	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R, pMMB-VCA0700 Carb^R	Δ6 (VCA0700 only) mutant with with pMMB67EH containing VCA0700 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH427 / SAD1376)
Δ6 (VCA0700 only) pMMB-VC1073	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R, pMMB-VC1073 Carb^R	Δ6 (VCA0700 only) mutant with with pMMB67EH containing VC1073 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH429 / SAD1377)
Δ6 (VCA0700 only) pMMB-VCA0140	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R, pMMB-VCA0140 Carb^R	Δ6 (VCA0700 only) mutant with with pMMB67EH containing VCA0140 – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH430 / SAD1378)
Δ6 (VCA0700 only) pMMB-GbpA	ΔChiA1, ΔChiA2, ΔVC0769, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Kan^R, pMMB-GbpA Carb^R	Δ6 (VCA0700 only) mutant with with pMMB67EH containing GbpA – cloned into the EcoRI and BamHI sites of the MCS.	This Study (CAH428 / SAD1379)
ΔVC0995	ΔVC0995, ΔVC1807::Spec^R	Deleted 500bp of the 5' end of VC0995	This Study (SAD265)
ΔVC1282	ΔVC1282, ΔVC1807::Spec^R	Deleted 500bp of the 5' end of VC0995	This Study (SAD269)
ΔVC0618-19	ΔVC0618-VC0619	In-frame deletion of VC0618-19	This Study (SAD387)
ΔVC0995 ΔVC0618-19	ΔVC0995, ΔVC0618-19, ΔVC1807::Kan^R	Deleted 500bp of the 5' end of VC0995 and in-frame deletion of VC0618-19	This Study (CAH545 / SAD1380)
ΔVC0995 ΔVC1282	ΔVC0995 ΔVC1282, ΔVC1807::Kan^R	Deleted 500bp of the 5' end of VC0995 and VC1282	This Study (CAH542 / SAD1381)
ΔVC1282 ΔVC0618-19	ΔVC1282, ΔVC0618-19, ΔVC1807::Kan^R	Deleted 500bp of the 5' end of VC1282 and in-frame deletion of VC0618-19	This Study (CAH543 / SAD1382)
ΔVC0995 ΔVC1282 ΔVC0618-19	ΔVC0995, ΔVC1282, ΔVC0618-19, ΔVC1807::Kan^R	Deleted 500bp of the 5' end of VC0995 and VC1282. As well as an in-frame deletion of VC0618-19	This Study (CAH544 / SAD1383)
ΔVC0972	ΔVC0972::Spec^R	Deleted VC0972 and replaced it with a spec^R cassette	This Study (SAD115)
Δ7 pMMB-ChiA1-FLAG	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073,	Added a C-terminal FLAG tag onto the pMMB construct indicated.	This Study (CAH714 /

	ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-ChiA1-FLAG Carb^R		SAD1538)
Δ7 pMMB-ChiA2-FLAG	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-ChiA2-FLAG Carb^R	Added a C-terminal FLAG tag onto the pMMB construct indicated.	This Study (CAH715 / SAD1539)
Δ7 pMMB-VC0769-FLAG	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-VC0769-FLAG Carb^R	Added a C-terminal FLAG tag onto the pMMB construct indicated.	This Study (CAH758 / SAD1540)
Δ7 pMMB-VCA0700-FLAG	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-VCA0700-FLAG Carb^R	Added a C-terminal FLAG tag onto the pMMB construct indicated.	This Study (CAH770 / SAD1541)
Δ7 pMMB-VC1073-FLAG	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-VC1073-FLAG Carb^R	Added a C-terminal FLAG tag onto the pMMB construct indicated.	This Study (CAH769 / SAD1542)
Δ7 pMMB-VCA0140-FLAG	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-VCA0140-FLAG Carb^R	Added a C-terminal FLAG tag onto the pMMB construct indicated.	This Study (CAH762 / SAD1543)
Δ7 pMMB-GbpA-FLAG	ΔChiA1, ΔChiA2, ΔVC0769, ΔVCA0700, ΔVC1073, ΔVCA0140, ΔGbpA, ΔVC1807::Spec^R, pMMB-GbpA-FLAG Carb^R	Added a C-terminal FLAG tag onto the pMMB construct indicated.	This Study (CAH761 / SAD1544)

Table S2 - *Primers used in this study*

Primer Name	Primer Sequence (5'→3')	Description
Primers for SOE products		
BBC509	ACTACAATGTATTGATGAAGTGG	ΔChiA1 F1
BBC510	GCTAATTCAGTTTAAGCGGCCATGTTTCTCTCCTGTATTAGATG	ΔChiA1 R1
BBC511	ATGGCCGCTTAAACTGAATTAGCCAGCGTAACGCCTACTGGTAAC	ΔChiA1 F2
BBC512	TTTTGTAGTCTTGTGCTTGCAG	ΔChiA1 R2
BBC514	AAAAGCATCGCTGGAAGAGTG	ΔChiA2 F1
BBC515	GCTAATTCAGTTTAAGCGGCCATAAGTTTTCTCTCTCTTCCTTAG	ΔChiA2 R1
BBC516	ATGGCCGCTTAAACTGAATTAGCGAGTATTTATGATCGTAAGTTTACGG	ΔChiA2 F2
BBC517	TCACCGAAATTGCACCAATCAAC	ΔChiA2 R2
BBC519	CCAGAACAAACCATTGCTGATG	ΔVC0769 F1
BBC520	GCTAATTCAGTTTAAGCGGCCATGGATAAAAGTCCCTCTCTC	ΔVC0769 R1
BBC521	ATGGCCGCTTAAACTGAATTAGCAGAGTGGCAACAAGCGCTG	ΔVC0769 F2
BBC522	TTGCATGGTTCGCAAGCTTAAG	ΔVC0769 R2
BBC524	AAGTGCAGTTGGATCACTGACAC	ΔgbpA F1
BBC525	GCTAATTCAGTTTAAGCGGCCATCACAGACTCTTCTTTGTTAGC	ΔgbpA R1
BBC526	ATGGCCGCTTAAACTGAATTAGCCCACGAATGTATCGTGCCTG	ΔgbpA F2
BBC527	CTCATGCATCGTATGTGAAAGC	ΔgbpA R2
BBC529	CAGTTAATTGCTCAAAACCAGC	ΔVCA0700 F1
BBC530	GCTAATTCAGTTTAAGCGGCCATTGTTGTTCTTCCCTCAAG	ΔVCA0700 R1
BBC531	ATGGCCGCTTAAACTGAATTAGCTAAAGGGGCTGTCAGCACC	ΔVCA0700 F2
BBC532	AACGCTTTCATATCTCAGAGCG	ΔVCA0700 R2
BBC534	TTTCAGCGCCTGTCAAAGAAG	ΔVC1073 F1
BBC535	GCTAATTCAGTTTAAGCGGCCATTATTTCGAGACTTATTTTATTGAAC	ΔVC1073 R1
BBC536	ATGGCCGCTTAAACTGAATTAGCGTTCATTGAAGGCCAGACCG	ΔVC1073 F2
BBC537	CAGTGCGCTGTTTGGTATGG	ΔVC1073 R2
BBC539	AATATCAAACCCTTCCGTGACAC	ΔVCA0140 F1
BBC540	GCTAATTCAGTTTAAGCGGCCATTTCTGTTTACAAATGGCTAAC	ΔVCA0140 R1
BBC541	ATGGCCGCTTAAACTGAATTAGCACTGACGTGGGATGACTTGGAA	ΔVCA0140 F2
BBC542	AATTTGTCGAGCTTGGAAAGGAG	ΔVCA0140 R2
BBC401	ACCAGCAAAGCTAATAAAATCGAG	ΔpilA (VC2423) F1
BBC402	gtcgacggatccccggaatGAGCATATGCCTTGCTACACAAG	ΔpilA (VC2423) R1
BBC403	gaagcagctccagcctacaACTGCAGGTGCAACAATTAACTAA	ΔpilA (VC2423) F2
BBC404	CGCCATACTAACCCAATACACTC	ΔpilA (VC2423) R2
ABD927	GCAGAGAAAGGGTATCATTACTGG	ΔVC0995 F1
ABD928	GcTAATTCAGTTTAAGCGGCCATCTTAAGTTCCCCCTATAG	ΔVC0995 R1

	GATTTTTG	
ABD929	ATGGCCGCTTAAACTGAATTAgCACATCAGGTGCTTTAGGCCAATTTG	ΔVC0995 F2
ABD930	TACTCTCGTTTTTCGGCTTACTC	ΔVC0995 R2
ABD943	ATATTCTTGCGGTATTAGCCACAC	ΔVC1282 F1
ABD944	GCTAATTCAGTTTAAGCGGCCATCTTATATTTAAGATAAAGAGTTCCCTA	ΔVC1282 R1
ABD945	ATGGCCGCTTAAACTGAATTAGCATTACCATTCGTATGCCAGAGC	ΔVC1282 F2
ABD946	GCAGATGTTTCATTAAAGGGTCG	ΔVC1282 R2
BBC081	AAGCAAGTTCACGTTTGCCG	ΔVC0618-19 F1
BBC082	gtcgacggatccccggaatCATAACTTACACCTTACTCACCCAG	ΔVC0618-19 R1
BBC083	gaagcagctccagcctacaGGAGATAAATAATCATGACTACGCC	ΔVC0618-19 F2
BBC084	TAAAGTTCGCAACACGCC	ΔVC0618-19 R2
ABD800	TTTGTCGGTGGTGTTACGGTAAG	ΔVC0972 F1
ABD801	gtcgacggatccccggaatCATGGATAACTCCTAAAAATGGATATAGCTG	ΔVC0972 R1
ABD802	gaagcagctccagcctacaGTACGTGTAGGTCTGGAATACGG	ΔVC0972 F2
ABD803	AAAGCAAGATACAGAACGCGACC	ΔVC0972 R2
Primers to clone genes into pMMB67EH		
CAH0030	gataacaatttcacacaggaaacagaattcAggaggtAGAAACATGAAGCGCTATTG	ChiA1 F
CAH0031	gactctagaggatccccgggtaccgagctcTTACTGAGCATTATTCATCTGGC	ChiA1 R
CAH0032	ataacaatttcacacaggaaacagaattcAggaggtAAACTTATGAATCGAATGACTTTG	ChiA2 F
CAH0033	gactctagaggatccccgggtaccgagctcTTAATGAGTAGAACAACTCGCGGC	ChiA2 R
CAH0026	aacaatttcacacaggaaacagaattcAggaggtTTATCCATGTTTAAACTCAAACATAC	VC0769 F
CAH0027	gactctagaggatccccgggtaccgagctcTTAGCAGGACACCTTATCCCAG	VC0769 R
CAH0036	gataacaatttcacacaggaaacagaattcAggaggtACAACAATGCGTGTACTCG	VCA0700 F
CAH0037	gactctagaggatccccgggtaccgagctcTTACGCCTGAGGGCAAGTC	VCA0700 R
CAH0028	aacaatttcacacaggaaacagaattcAggaggtGAAATAATGAAAAGATCAGCATTAAC	VC1073 F
CAH0029	gactctagaggatccccgggtaccgagctcTTAGATTTTGCACACCGCTTTCC	VC1073 R
CAH0034	acaatttcacacaggaaacagaattcAggaggtATAACCATGAAATACGGATTAAAAATC	VCA0140 F
CAH0035	gactctagaggatccccgggtaccgagctcTTAGCGCCACACACCCC	VCA0140 R
CAH0038	acaatttcacacaggaaacagaattcAggaggtTCTGTGATGAAAAAACAACCTAAAATG	gbpA F
CAH0039	gactctagaggatccccgggtaccgagctcTTAACGTTTATCCCACGCCATTTC	gbpA R
BBC277	TATAGAATTCATGGATATGAATGAGCAACAG	tfoX F
BBC278	TATAGGATCCTTAACGCTGCTGACAACTTTC	tfoX R
Primers to add C-terminal FLAG tag onto pMMB chitinase expression vectors		
BBC1531	gattataaggatgacgatgacaaaTAAGAGCTCGGTACCCGG	Universal F primer for adding a C-terminal FLAG tag onto pMMB cloned chitinases

BBC1534	CTCTTAtttgtcatcgtcatccttataatcCTGAGCATTATTCATCTGGCTAAG	ChiA1 specific R
BBC1535	CTCTTAtttgtcatcgtcatccttataatcATGAGTAGAACAACTCGCGGC	ChiA2 specific R
BBC1662	CTCTTAtttgtcatcgtcatccttataatcGCAGGACACCTTATCCCAGAAC	VC0769 specific R
BBC1664	CTCTTAtttgtcatcgtcatccttataatcGATTTTGCACACCGCTTTCCATG	VC1073 specific R
BBC1666	CTCTTAtttgtcatcgtcatccttataatcCGCCTGAGGGCAAGTCACTTC	VCA0700 specific R
BBC1670	CTCTTAtttgtcatcgtcatccttataatcGCGCCACACACCCCATTCG	VCA0140 specific R
BBC1668	CTCTTAtttgtcatcgtcatccttataatcACGTTTATCCCACGCCATTTCC	GbpA specific R
Primers for MASC-PCR		
ABD969	ATGGCCGCTTAAACTGAATTAGC	F primer for all MASC-PCR reactions
BBC513	ATAAGGCTCAGAGCTATCGATC	R primer for ΔChiA1 detect ~190bp
BBC518	CAGGAAACGTTTCACAGAAGC	R primer for ΔchiA2 detect ~400bp
BBC523	TTTTGGTGCTTGTGGCGTG	R primer for VC0769 detect ~301bp
BBC528	CGCCAATCTCATACTCTTTCGC	R primer for gbpA detect ~500bp
BBC533	CTACCCCTGGCCACTCTTTACC	R primer for VCA0700 detect ~650bp
BBC538	TTTCCCACGTCATCTTGGTC	R primer for VC1073 detect ~800bp
BBC543	CCAGAACTCTTAACCACCATG	R primer for VCA0140 detect ~1000bp