1 **Inducible and reversible phenotypes in a novel mouse model of Friedreich's Ataxia**

2 Vijayendran Chandran[1, #], Kun Gao[1], Vivek Swarup[1], Revital Versano[1], Hongmei Dong[1], Maria C.
3 Jordan[3] & Daniel H. Geschwind[1,2]
4

5 [1]Program in Neurogenetics, Department of Neurology, [2]Department of Human Genetics, [3]Department
6 of Physiology, David Geffen School of Medicine, University of California, Los Angeles, CA 90095, USA.
7 [#]Present address: Department of Pediatrics, School of Medicine, University of Florida, Gainesville, FL
8 32610-0296, USA.
9 Correspondence should be addressed to D.G. (dhg@mednet.ucla.edu) or V.C. (vijayendran@ufl.edu).
10

ABSTRACT

Friedreich's ataxia (FRDA), the most common inherited ataxia, is caused by recessive mutations that

reduce the levels of frataxin (FXN), a mitochondrial iron binding protein. We developed an inducible

mouse model of *Fxn* deficiency that enabled us to control the onset and progression of disease

phenotypes by the modulation of *Fxn* levels. Systemic knockdown of *Fxn* in adult mice led to multiple

phenotypes paralleling those observed in human patients across multiple organ systems. By reversing

knockdown after clinical features appear, we were able to determine to what extent observed

phenotypes represent reversible cellular dysfunction. Remarkably, upon restoration of near wild-type

FXN levels, we observed significant recovery of function, associated pathology and transcriptomic

dysregulation even after substantial motor dysfunction and pathology were observed. This model will be

of broad utility in therapeutic development and in refining our understanding of the relative contribution

of reversible cellular dysfunction at different stages in disease.

INTRODUCTION

A guanine-adenine-adenine (GAA) trinucleotide repeat expansion within the first intron of the frataxin (*Fxn*) gene in the chromosome 9 is the major cause of Friedreich's ataxia (FRDA), the most commonly inherited ataxia(1, 2). Due to recessive inheritance of this GAA repeat expansion, patients have a marked deficiency of *Fxn* mRNA and protein levels caused by reduced *Fxn* gene transcription(2, 3). Indeed, the GAA expansion size has been shown to correlate with residual *Fxn* levels, earlier onset and increased severity of disease(4, 5). The prevalence of heterozygous carriers of the GAA expansion is between 1:60 and 1:110 in European populations, and the heterozygous carriers of a *Fxn* mutation are not clinically affected(6, 7).

FRDA is an early-onset neurodegenerative disease that progressively impairs motor function, leading to ataxic gait, cardiac abnormalities, and other medical co-morbidities, ultimately resulting in early mortality (median age of death, 35 years)(8, 9). However, identification of the causal gene led to identification of a significant number of patients with late onset, they tend to have slower progression with less severe phenotype and are associated with smaller GAA expansions(10). This shorter expansion enables residual *Fxn* expression(11), thus modifying the classical FDRA phenotype, consistent with other data indicating that *Fxn* deficiency is directly related to the FRDA phenotype(12). Extra-neurologic symptoms including metabolic dysfunction and insulin intolerance are observed in the majority and frank type I diabetes is observed in approximately 15% of patients, the severity of which is related to increasing repeat length(13-15). The mechanisms by which *Fxn* reduction leads to clinical symptoms and signs remain to be elucidated, but molecular and cellular dysfunction mediated by a critical reduction in *Fxn* levels plays a central role(16).

FXN is a nuclear-encoded mitochondrial protein involved in the biogenesis of iron-sulphur clusters, which are important for the function of the mitochondrial respiratory chain activity(17, 18). Studies in mouse have shown that *Fxn* plays an important role during embryonic development, as homozygous frataxin knockout mice display embryonic lethality(19), consistent with FXN's evolutionary

conservation from yeast to human(2, 3). Over the past several years, multiple mouse models of

frataxin deficiency, including a knock-in knockout model(20), repeat expansion knock-in model(20),

transgenic mice containing the entire *Fxn* gene within a human yeast artificial chromosome, YG8R and

YG22R(21, 22), as well as a conditional *Fxn* knockout mouse, including the cardiac-specific(23) and a

neuron specific model(23) have been generated. These existing transgenic and heterozygous knockout

FRDA animal models are either mildly symptomatic, or restricted in their ability to recapitulate the

spatial and temporal aspects of systemic FRDA pathology when they are engineered as tissue-specific

conditional knockouts(20-25). Despite advances made towards elucidating FRDA pathogenesis, many

questions remain due to the need for mouse models better recapitulating key disease features to

understand frataxin protein function, disease pathogenesis and to test therapeutic agents.

In this regard, one crucial question facing therapeutic development and clinical trials in FRDA is

the reversibility of symptoms. The natural history of the disorder has been well described(26, 27), it is

not known how clinical features such as significant motor disability relate to reversible processes (e.g.

underlying neuronal dysfunction) or reflect irreversible cell death. It is often assumed that clinically

significant ataxia and motor dysfunction reflects neurodegeneration, although this may not be the case.

This issue, while critically important for therapeutic development, is difficult to address in patients, but

we reasoned that we could begin to address this question in an appropriate mouse model.

Here we report an inducible mouse model for FRDA, the FRDAkd mouse, that permits

reversible, and yet substantial frataxin knockdown, allowing detailed studies of the temporal

progression, or recovery following restoration of frataxin expression - the latter permitting exploration of

disease reversal given optimal treatment (normalization of *Fxn* levels). We observe that *Fxn* knockdown

leads to behavioral, physiological, pathological and molecular deficits in FRDAkd mice paralleling those

observed in patients, including severe ataxia, cardiac conduction defects and increased left ventricular

wall thickness, iron deposition, mitochondrial abnormalities, low aconitase activity, and degeneration of

dorsal root ganglia, retina, as well as early mortality. We identify a signature of molecular pathway

dysfunction via genome-wide transcriptome analyses, and show reversal of this molecular phenotype

and as well as behavioral and pathological measures, even in the setting of significant disability due to

motor dysfunction in FRDAkd animals.

RESULTS

RNA interference based in vivo frataxin knockdown and rescue

To investigate the neurological and cardiac effects linked to reduced FXN levels and to create a model

for testing new therapies *in vivo*, we sought to generate mice that develop titratable clinical and

pathological features of FRDA. We employed recombinase-mediated cassette exchange for genomic

integration of a single copy shRNA transgene (doxycycline-inducible) that can mediate frataxin

silencing temporally under the control of the H1 promoter via its insertion in a defined genomic locus

that is widely expressed (rosa26 genomic locus)(28). This allowed us to circumvent the lethal effect of

organism-wide knockout, while permitting significant frataxin reduction in all tissues. Depending on the

dose of the inductor doxycycline (dox), temporal *Fxn* knockdown was achieved to control the onset and

progression of the disease (**Fig. 1a**).

Six different shRNA sequences were screened in vitro to obtain a highly efficient shRNA

targeting the mature coding sequence of frataxin (**Fig. 1b** and **Supplementary Fig. 1**). Transgenic

animals (FRDAkd) containing a single copy of this efficient shRNA transgene (**Fig. 1b)** were generated

and characterized. First, to test *Fxn* knockdown efficiency, we explored the response to dox at varying

escalating doses in drinking water. At the higher doses, we observed mortality as early as two weeks

and a 100% mortality rate by 5 to 6 weeks, not permitting extended time series analyses (Online

Methods). We found that the combination of 2 mg/ml in drinking water coupled with 5 or 10 mg/kg

intraperitoneal injection of dox twice per week, led to efficient *Fxn* knockdown within two weeks post

treatment initiation, while avoiding a high early mortality rate (Online Methods). Hence, for all

subsequent experiments we utilized this regimen to model the chronicity of this disorder in patients by

balancing the gradual appearance of clinical signs and decline in function, while limiting early demise

(**Fig. 1c**).

To determine the effect of *Fxn* deficiency in adult mice after a period of normal development

(similar to a later onset phenotype in humans), which would allow establishment of stable baselines,

and to obtain relatively homogeneous data from behavioral tests(29) we initiated dox at 3 months.

Following twenty weeks with (Tg +) or without (Tg -) doxycycline administration (**Fig. 1c;** Online

Methods), we observed highly efficient silencing of *Fxn*, reaching greater than 90% knockdown across

multiple CNS and non-CNS tissues (P < 0.05, two-way ANOVA; **Fig. 1d,e**). Using this regimen, time

series western blot analyses of 80 independent animals (wildtype with dox (Wt +) N= 24; transgenic

with dox (Tg +) N= 24; transgenic without dox (Tg -) N= 24; transgenic with dox removal (Rescue Tg ±)

N= 8, at weeks 0, 3, 8, 12, 16 and 20) confirmed efficient silencing as early as 3 weeks, and efficient

rescue, as evidenced by normal frataxin levels, post 8 weeks dox removal (P < 0.05, two-way ANOVA;

Fig. 1f,g). Together, the results indicate that FRDAkd mice treated with dox are effectively FXN

depleted in a temporal fashion and that *Fxn* expression can be reversed efficiently by dox removal,

making it suitable for studying pathological and clinical phenotypes associated with FRDA.

Frataxin knockdown mice exhibit neurological deficits

The major neurologic symptom in FRDA is ataxia, which in conjunction with other neurological deficits

including axonal neuropathy and dorsal root ganglion loss, contributes to the gait disorder and

neurological disability(8, 9). So, we first determined whether *Fxn* knockdown impacted the behavior of

Tg and Wt mice with (+) or without (-) dox, or with dox, followed by its removal (±; the "rescue"

condition), leading to a total of five groups subjected to a battery of motor behavioral tasks (each group

N = 15-30; total mice = 108; Online Methods) (**Fig. 2a-g**). Wt - and Wt + control groups were included

to access baseline and to have a control for any potential dox effect on behavior; Tg - and Tg +

transgenic groups were compared to examine the effect of genotype and *Fxn* knockdown on behavior;

The Tg ± group was included to study the effect of FXN restoration after knockdown (rescue). We

observed significant weight loss and reduced survival ratio (<90%) at 25 weeks with dox treatment in

Fxn knockdown animals (Tg +) when compared to other control groups (P < 0.05, two-way ANOVA;

Fig. 2a,b). Tg + mice exhibited a shorter distance travelled at both 12 and 24 weeks in comparison to

control animals, consistent with decreased locomotor activity (P < 0.05, two-way ANOVA; **Fig. 2c**).

Next, we assessed gait ataxia(9) using paw print analysis(30) (Online Methods). The *Fxn* knockdown

mice (Tg +) displayed reduced hind and front limb stride length when compared with Tg-, as well as the

Wt control + and - dox at 12 and 24 weeks, suggesting ataxic gait (P < 0.05, two-way ANOVA; **Fig.**

2d,e and **Supplementary Fig. 2**). Grip strength testing also showed that Tg + animals displayed defect

in their forelimb muscular strength at 12 and 24 weeks when compared with other groups (P < 0.05,

two-way ANOVA; **Fig. 2f**). Finally, motor coordination and balance were evaluated using the Rotarod

test. Whereas no significant difference in time spent on the rod before falling off was seen between Wt

+ or Wt - or Tg - and Tg ± mice after 12 weeks post dox removal (rescue), chronically treated mice (Tg

+ mice) from 12 weeks onward fell significantly faster, indicative of motor impairments (P < 0.05, two-

way ANOVA; **Fig. 2g**). These observations suggest that the knockdown of *Fxn* in mice causes motor

deficits indicative of ataxia similar to FRDA patients(9), and demonstrates the necessity of normal

levels of *Fxn* expression in adults for proper neurological function.

Frataxin knockdown leads to cardiomyopathy

Cardiac dysfunction is the most common cause of mortality in FRDA(31, 32). To examine impaired

cardiac function in FRDAkd knockdown animals, we employed electrocardiogram (ECG) and

echocardiogram analyses to measure electrical activity and monitor cardiac dimensions. The ECG of

Tg + mice displayed a significant increase in QT interval duration when compared to the other

control/comparison groups at both 12 and 24 weeks post dox treatment, suggesting abnormal heart

rate and rhythm (arrhythmia)(33) (Online Methods; P < 0.05, two-way ANOVA; **Fig. 3a,e**). However,

rescue animals (Tg ±) 12 weeks after dox removal showed normal electrocardiogram, demonstrating

that by restoring the FXN levels, the prolonged QT interval can be reversed (P < 0.05, two-way

ANOVA; **Fig. 3b,e**). We also observed that the Tg + animals at week 24 displayed absence of P-waves

suggesting an atrial conduction abnormality(34) (**Fig. 3c**), not observed in the rescued animals. Similar

abnormalities have been variably observed in many FRDA patients(35).

Progressive hypertrophic cardiomyopathy (thickening of ventricular walls) related to the severity

of frataxin deficiency(35-37) is frequently observed in FRDA patients(31). To confirm structural cardiac

abnormalities, echocardiogram analyses were performed, focusing on left ventricular function. At 12

weeks there was a non-significant trend towards increasing ventricular wall thickness. However, by 24

weeks, dox treated transgenic animals (Tg +) exhibited ventricular and posterior wall thickening,

suggesting hypertrophic cardiomyopathy in these animals when compared to other control groups (P <

0.05, two-way ANOVA; **Fig. 3d,f,g**). Together, these observations indicate that the transgenic mice

exhibit progressive cardiomyopathy due to reduced level of *Fxn*, supporting the utilization of Tg + mice

for examining the molecular mechanisms downstream of *Fxn* deficiency responsible for cardiac defects.

Cardiac pathology observed with frataxin knockdown

We next explored pathological consequences of FRDA knockdown in FRDAkd mice. In FRDA patients,

reduced frataxin induces severe myocardial remodeling, including cardiomyocyte iron accumulation,

myocardial fibrosis and myofiber disarray(9). Indeed, we observed substantially increased myocardial

iron in Tg + mice, as evidenced by increased ferric iron staining (**Fig. 4a** and **Supplementary Fig. 3**)

and the increased expression of iron metabolic proteins, ferritin and ferroportin at 20 weeks (**Fig. 4b**

and **Supplementary Fig. 3**). Cardiac fibrosis is commonly found in association with cardiac

hypertrophy and failure(38). Histological analysis by Masson's trichrome staining revealed excessive

collagen deposition in Tg + mice hearts at 20 weeks when compared to other control groups,

suggesting cardiac fibrosis (**Fig. 4c**). Further examination of cardiomyocyte ultrastructure by electron

microscopy in control mouse (Wt +) heart demonstrates normally shaped mitochondria tightly packed

between rows of sarcomeres (**Fig. 4d**). In contrast, Tg + mice demonstrate severe disorganization,

displaying disordered and irregular sarcomeres with enlarged mitochondria at 20 weeks (**Fig. 4d**). In a

minority of cases, but never in controls, we observed mitochondria with disorganized cristae and

vacuoles in Tg + mouse heart at 20 weeks, suggesting mitochondrial degeneration (**Fig. 4e**). Next, by

examining aconitase which is a Fe-S containing enzyme whose activity is reduced in FRDA patients(39,

40), activities in Tg + and other control groups, we observed decreased aconitase activity in the Tg +

mouse heart at 20 weeks. Together these observations suggest that the knockdown of *Fxn* in mice

causes cardiac pathology similar to that observed in patients(32).

Frataxin knockdown causes neuronal degeneration

In FRDA patients and mice with *Fxn* conditional knockout, a cell population that is seriously affected by

frataxin reduction is the large sensory neurons of dorsal root ganglia (DRG), which results in their

degeneration(9). As in heart tissue, lack of *Fxn* also induces mitochondrial dysfunction in the DRG(40).

Thus, we assessed whether mitochondrial abnormalities are observed in Tg + mice DRG neurons by

electron microscopy (Online Methods). As anticipated, no mitochondrial abnormalities were detected in

DRG neurons in control groups, but a significant increase ($P < 0.05$, two-way ANOVA) in condensed

mitochondria were detected in DRG neurons of Tg + mice at 20 weeks (**Fig. 5a-c,e**), but not in Tg - or

Wt + mice. In most cases we observed lipid-like empty vacuoles associated with condensed

mitochondria in DRG neurons of Tg + animals, but not in other control groups, suggesting neuronal

degeneration, similar to what is observed in FRDA patients(9) ($P < 0.05$, two-way ANOVA; Online

Methods; **Fig. 5b,c**). Previous results in human post-mortem spinal cords of FRDA patients showed a

decrease in axon size and myelin fibers(8). By analyzing more than 2000 axons per group in the lumbar

spinal cord cross-section of high-resolution electron micrographs, we observed significant reduction in

axonal size and myelin sheath thickness in the spinal cord samples of Tg + mice when compared to

other control groups at 20 weeks ($P < 0.05$, two-way ANOVA; **Fig. 5f,g**). In the cerebellum, the number

of Purkinje cells, the cerebellar granular and molecular layers were not altered due to *Fxn* knockdown

at 20 weeks (**Supplementary Fig. 4**).

Next we explored the visual system, because in FRDA patients, various visual field defects have

been reported, suggesting that photoreceptors and the retinal pigment epithelium (RPE) can be

affected(41, 42). By electron microscopic examination of the photoreceptors in the retina, which are

specialized neurons capable of phototransduction, we observed disruption in Tg + mice at 20 weeks

(Online Methods; **Fig. 5h**). Previous work has shown that the disruption of photoreceptors is related to

visual impairment(43). Similarly, we also found a significant increase in degenerating RPE cells with

vacuoles in Tg + mice, which is involved in light absorption and maintenance of visual function (**Fig. 5i**).

Together, these results suggest that *Fxn* knockdown in the CNS of adult mice results in degeneration of

retinal neurons.

Gene expression changes due to frataxin knockdown

Given the phenotypic parallels in the cardiac and nervous system abnormalities in FRDAkd mice with

chronic *Fxn* reduction following treatment with dox, we next sought to explore genome wide molecular

mechanisms and determine which pathways were affected in the heart and nervous system, and if they

were reversible. We analyzed global gene expression profiles in the heart, cerebellum and DRGs from

Tg +, Tg -, Wt + and Tg ± mice from 0, 3, 12, 16, 20 and plus 4, 8 weeks post dox treatment (n= 192).

Differential expression analyses (Online Methods) identified 1959 genes differentially expressed in Tg +

mice heart when compared to Wt + and Tg - mice (FDR < 5%, **Fig. 6a, Supplementary Table 1**).

Similarly, we observed 709 and 206 genes differentially expressed in cerebellum and DRGs of Tg +

mice. While cross tissue overlap in expression changes was significant, the majority of changes were

tissue specific; only 31% and 38% of genes that were differentially expressed in the Tg + mice

cerebellum and DRGs were also dysregulated in heart. Likewise, we observed a 19% overlap between

DRGs and cerebellum, consistent with previous observations that *Fxn* reduction causes distinct

molecular changes in different tissues(15).

Next, we analyzed these differently expressed transcripts in cardiac tissue from Tg + mice with

respect to cellular pathways. The top GO categories and KEGG pathways include chemotaxis, immune

response, lysosome and phagocytosis, vesicle transport and endocytosis, p53 signaling pathway, cell

cycle and division, protein transport and localization, nucleoside and nucleotide binding, and

mitochondrion (Benjamini corrected P-value < 0.05) (**Fig. 6a, Supplementary Table 2**). To

characterize the temporal patterns of these signaling cascades after frataxin knockdown and rescue,

we examined their time course by PCA analyses of the gene expression profiles (**Fig. 6b**). By

examining the cumulative explained variability of the first three principal components for these clusters

of genes, we show that each of these functional groups are activated as early as three weeks after dox

initiation (and remain elevated for up to 20 weeks); importantly, the aberrant expression of all of these

clusters observed in Tg + mice are largely reversed after eight weeks of rescue via *Fxn* re-expression

(**Fig. 6a,b**).

Another notable observation is that immune system activation is among the earliest pathways

regulated after *Fxn* knockdown (**Fig. 6b**). This suggests that initiation of immune responses (innate and

adaptive) is a direct consequence of *Fxn* knockdown. For example, 38 genes involved in the chemokine

signaling pathway (KEGG: mmu04062) were significantly differentially expressed due to *Fxn*

knockdown in Tg + mice heart (**Supplementary Fig. 5**). Cross validating these genes with previously

published gene expression datasets obtained from FRDA patients(44) and mouse models associated

with FRDA(20, 23), identified several genes involved in the chemokine signaling pathway (E.g.: CCL2,

3, 4, 7, CXCL1, 16, PRKCD, STAT3) consistently differentially expressed in these six independent

FRDA associated datasets (**Supplementary Fig. 6**). This implicates a vital role for chemokines and

immune response in FRDA pathology, as has been suggested for other neurodegenerative

diseases(45-48).

Shared and tissue specific gene expression changes

We next performed weighted gene co-expression network analysis (WGCNA)(49-51), a powerful

method for understanding the modular network structure of the transcriptome, to organize the frataxin

related transcriptional changes in an unbiased manner following knockdown and rescue. WGCNA

permits identification of modules of highly co-expressed genes whose grouping reflects shared

biological functions and key functional pathways, as well as key hub genes within the modules, and has

been widely applied to understanding disease related transcriptomes(52, 53). By applying WGCNA and

consensus network analysis(50) we identified 19 robust and reproducible co-expression modules

(Online Methods; **Supplementary Fig. 7 and Supplementary Table 3**) between the three tissues

datasets generated at 0, 3, 12, 16 and 20 weeks after *Fxn* knockdown and 4 and 8 weeks post dox

removal. On the basis of the module eigengene correlation with time-dependent changes after *Fxn*

knockdown, we first classified modules as up-regulated and down-regulated following knockdown

(**Supplementary Fig. 8**). Next, based on the significant module trait relationships (Wilcoxon P-value <

0.05), we identified 11 modules strongly associated with *Fxn* knockdown: three down-regulated

modules in two or more tissues after *Fxn* knockdown (yellow, lightgreen and turquoise) and three up-

regulated modules (blue, purple, and black) (**Supplementary Fig. 8**), three down-regulated modules in

heart, which were up-regulated in cerebellum (red, greenyellow and magenta) and two up-regulated

modules in heart which were down-regulated in cerebellum (cyan and pink). Although six of the gene

co-expression modules (yellow, lightgreen, turquoise, blue, purple, and black) in the heart, cerebellum

and DRG following *Fxn* knockdown are highly preserved across tissues, five modules (red,

greenyellow, magenta, cyan and pink) exhibit differential expression profiles suggesting tissue specific

molecular changes, consistent with previous observations of shared and organ specific changes(15)

(**Supplementary Fig. 8**).

As a first step toward functional annotation of the cross tissue modules, we applied GO and

KEGG pathway enrichment analyses, which showed enrichment (Benjamini-corrected p values < 0.05)

for several GO categories and pathways in the *Fxn* knockdown co-expression modules which included

several previously associated functional categories related with FRDA (**Supplementary Table 4**).

Three modules (yellow, lightgreen and turquoise) that were down-regulated in two or in all three tissues

due to *Fxn* knockdown included, nucleotide, nucleoside and ATP binding, myofibril assembly, muscle

tissue development, RNA processing, and several mitochondrial related categories: oxidative

phosphorylation, respiratory chain, NADH dehydrogenase activity, and electron transport chain. We

also observed that the genes present in turquoise module were enriched for several KEGG pathways,

namely, PPAR signaling (mmu03320; genes=14), insulin signaling (mmu04910; n=19), fatty acid

metabolism (mmu00071; n=10), cardiac muscle contraction (mmu04260; n=20), dilated

cardiomyopathy (mmu05414; n=13), and hypertrophic cardiomyopathy (mmu05410; n=14), which have

been previously associated with FRDA, reflecting the multi-systemic nature of FRDA. Similarly, three

upregulated cross tissue modules (blue, purple, and black) include, nucleotide binding, vesicle-

mediated transport, immune response (innate and adaptive), defense response, inflammatory

response, induction of apoptosis, positive regulation of cell death, cell adhesion, and skeletal system

development (**Supplementary Table 4**). These results demonstrate that unsupervised analyses can

identify groups of genes not only with shared biological functions, but also relevant to the clinical

phenotypes observed in FRDA.

Three tissue specific modules that were down-regulated in heart and up-regulated in cerebellum

(red, greenyellow and magenta) showed enrichment for, transcription regulator activity, neurological

system process, synaptic vesicle and nucleotide, nucleoside and ATP binding. Two other modules that

were up-regulated in heart and down-regulated in cerebellum (cyan and pink) were enriched for cell

cycle, cell division, mitosis and DNA replication (**Supplementary Table 4**). In summary, we observed

several metabolic functional categories that were differentially expressed (up and down) due to *Fxn*

knockdown. The modules consisting of mitochondrial and cardiac specific categories along with PPAR

signaling, insulin signaling, fatty acid metabolism pathways were down regulated in all tissues.

Likewise, the modules enriched for immune, apoptosis and cell death related categories we up-

regulated in all tissues due to *Fxn* knockdown. Synaptic and transcription regulator activity functional

categories were only up-regulated in cerebellum, whereas cell cycle, cell division, mitosis and DNA

replication related functional categories were down-regulated in cerebellum and up-regulated in the

heart. These general functional categories related to *Fxn* knockdown have been previously associated

with altered function in FRDA patients(44, 54), suggesting that genes within these modules would make

interesting candidate genes for follow up studies, because many of the genes have not been previously

associated with FRDA pathology and the disease mechanism.

Gene expression candidate biomarkers associated with frataxin knockdown

To identify candidate molecular targets and to better understand the molecular mechanism associated

with *Fxn* knockdown, we first manually combined all the GO ontology terms (see above and

Supplementary Table 4) that were enriched in the 11 modules into 26 broad functional categories

based on GO slim hierarchy (**Supplementary Table 5**) and screened for co-expressed genes within

each functional category in all three tissues (r > 0.5 and P-value < 0.05) over the time course. This

allowed us to identify critical functional sub-categories that are up or down regulated due to frataxin

knockdown and subsequently permitted us to detect differentially expressed candidate genes that are

co-expressed within each functional category (**Fig. 6d; Supplementary Fig. 9**). For example, we show

that immune, cell cycle and apoptosis related functional groups are up-regulated, whereas cardiac,

macromolecule and mitochondrial related functional groups were down-regulated (**Fig. 6d**). In the

immune category, we observed most prominent changes in complement activation pathway genes,

namely, C3, C4B, C1QB, C1QC and Serping1. Interesting, we also observed that many of these genes

were also up-regulated in peripheral blood mononuclear cells obtained from FRDA patients

(**Supplementary Fig. 10**), suggesting the potential for complement activation to act as a biomarker for

FRDA as previously suggested for other degenerative diseases(55). Similarly, we found several genes,

for example, CACNA2D1, ABCC9 and HRC involved in normal cardiac function, to be down-regulated

in heart tissue upon frataxin knockdown (**Fig. 6d**). CACNA2D1 is associated with Brugada syndrome,

also known as sudden unexpected nocturnal death syndrome, a heart condition that causes ventricular

arrhythmia(56). Mutations in ABCC9 gene can cause dilated cardiomyopathy(57) and a genetic variant

in the HRC gene has been linked to ventricular arrhythmia and sudden death in dilated

cardiomyopathy(58). These observations suggest that lower levels of frataxin causes dysregulation of

multiple genes related to arrhythmia or cardiac failure, a primary cause of death in FRDA patients.

There has been accumulating evidence suggesting that apoptosis may be an important mode of

cell death during cardiac failure(59). In agreement with this, we observed genes related to apoptosis

were up-regulated after *Fxn* knockdown in FRDAkd mice heart (**Fig. 6d**), which has been previously

associated with FRDA pathogenesis and reported in other *Fxn* deficiency models(24, 60, 61). In order

to validate our network findings, we tested CASP8 protein levels(62), observing an increase in cleaved

Caspase 8 protein levels in Tg + heart tissue compared with control mice (**Fig. 6e**). Next, we employed

the TUNEL assay to detect apoptotic cells that undergo extensive DNA degradation during the late

stages of apoptosis(63). However, we did not observe an increase in cell death in all tissues by TUNEL

staining (**Fig. 6f**).

Literature data extraction for candidate genes associated with frataxin knockdown

We next examined the phenotype-gene associations extracted by co-occurrence-based text-mining in

an attempt to link FRDA disease phenotypes with genes. For this, we screened the literature for

potential co-occurrence link association between the observed FRDAkd mice phenotypes and the

genes that are differentially expressed after *Fxn* knockdown (Online Methods). Identifying potential

biomarker candidates that are previously validated for certain phenotypes can provide insight into

disease progression, pathogenesis and extremely valuable for assessing therapeutic options(64). We

screened with the genes that are differentially expressed (FDR < 5%) and present in the co-expression

modules associated with behavioral and pathological key-terms (Eg: ataxia; **Supplementary Table 6**)

in the published literature. Interestingly, this analysis identified numerous genes in which mutations are

known to cause Mendelian forms of ataxia namely, ANO10(65, 66), CABC1(67, 68) and POLG(69, 70)

that were significantly down-regulated in the heart tissue and mildly down-regulated in the cerebellum

after *Fxn* knockdown in Tg + animals (**Fig. 6g and Supplementary Fig. 11**). We also found three other

genes that are differentially expressed due to *Fxn* knockdown, namely, CSTB(71), NHLRC1(72) and

PMM2(73) all of which are associated with other disorders when mutated causes ataxia (**Fig. 6g**).

These observations suggest that *Fxn* along with multiple other downstream candidates causes

behavioral deficits in FRDAkd mice. Similarly, for cardiac phenotypes, we identified multiple genes

related to cardiac fibrosis that were up-regulated in FRDAkd mice heart (**Fig. 6g**), including

LGALS3(74), ICAM1(75) and TIMP1(76) were up-regulated (**Fig. 6g**). Genes related to iron regulation,

included HFE(77), SLC40A1(77), HMOX1(78), TFRC(77) and GDF15(79) all of which are directly

involved in hemochromatosis and iron overload (**Fig. 6g**). We also found previously associated genes

related with muscle strength (E.g.: SRL, DCN), myelination (E.g.: PMP22, LGALS3, TLR2 and SIRT2)

and several genes related to neuronal degeneration (E.g.: GRN and APP) to be dysregulated in

FRDAkd mice (**Fig. 6g and Supplementary Fig. 11**), connecting this degenerative disorder with the

molecular signaling pathways known to be causally involved in other such disorders.

Perhaps the most prominent such known pathway associated with neurodegeneration, was

autophagy, since several autophagy-related genes (LAMP2, ATF4, TLR2, OPTN, MAPK14, SIRT2,

ICAM1, LGALS3, DCN, RCAN1, GRN) were present in these disease phenotype-associated sub

networks (**Fig. 6g**). Disruption of autophagy is also reported as altered in other *Fxn* deficiency

models(24, 60, 61) and is associated with various stress conditions including mitochondrial

dysfunction(80, 81). Autophagy is responsible for the recycling of long-lived and damaged organelles

by lysosomal degradation(82). To validate our network findings, we utilized LC3-II as a marker for

autophagy, showing autophagy activation in Tg + mice heart, but not in spinal cord tissue (**Fig. 6e, h**).

This suggests activation of apoptosis (**Fig. 6d, e**) and autophagy (**Fig. 6e, h**) may therefore potentiate

the cardiac dysfunction of Tg + mice. Next, by examining the expression levels of all these sub network

genes (**Fig. 6g**) in the datasets of other FRDA related mouse models and patient peripheral blood

mononuclear cells, we show that they are also differentially expressed in the same direction (**Fig. 6i**),

suggesting these sub network genes can be ideal candidates for molecular biomarkers in FRDA. In

summary, consistent with our behavioral, physiological and pathological findings, we show multiple

candidate genes related to key degeneration related phenotypes to be altered in FRDAkd mice.

Rescue of behavioral, pathological and molecular changes due to frataxin restoration

Our data show that mice carrying the dox inducible shRNA for *Fxn* develop normally until they are

challenged with dox, which subsequently results in substantial FXN reduction and causes multiple

behavioral and pathological features observed in patients with FRDA, including cardiac and nervous

system dysfunction (**Fig. 2-5**). *Fxn* knockdown mice displayed remarkable parallels with many of the

behavioral, cardiac, nervous system impairments along with physiological, pathological and molecular

deficits observed in patients. At a general level FRDAkd mice displayed weight loss, reduced locomotor

activity, reduced strength, ataxia and early mortality (**Fig. 2**). In the nervous system, FRDAkd mice

showed abnormal mitochondria and vacuolization in DRGs, retinal neuronal degeneration, and reduced

axonal size and myelin sheath thickness in the spinal cord(8, 9, 24, 25) (**Fig. 5**). With regards to cardiac

dysfunction, FRDAkd mice exhibited conduction defects, cardiomyopathy, evidence of iron overload,

fibrosis, and biochemical abnormalities that are commonly observed in patients(8, 9, 25) (**Fig. 3,4**).

These features correspond to a phenotype of substantial multisystem, clinical disability consistent with

moderate to severe disease after 3 months of frataxin deficiency and that lead to a 50% mortality rate

at approximately 5 months in these mice.

Given that optimal therapy in patients with FRDA could be considered replenishment of FXN

itself (e.g. via gene therapy(83) or improvement of *Fxn* transcription via small molecules(84)), we next

asked how much of the behavioral, physiological, pathological and molecular phenotype(s) observed at

this relatively severe stage of illness could be reversed following such "optimal" therapy. In this case,

optimal therapy is return of normal FXN levels under endogenous regulation through relief of

exogenous inhibition. Answering this question of reversibility is crucial for any clinical trial, whatever the

mechanism of action of the therapy, since we currently have limited information as to what represents

potentially reversible neurologic or cardiac phenotypes.

To address this, we compared two groups of mice, both transgenic and treated with dox for 12

weeks, at which time both groups show equivalent levels of substantial clinical features

(**Supplementary Fig. 12**). In one, Tg + the dox is continued for another 12 or more weeks, and in the

other Tg ± the dox is removed and the animals are followed. Restoration of FXN expression in *Fxn*

knockdown mice (Tg ±) that had reached a level of substantial clinical dysfunction led to significant

improvement in lifespan (no death until 20 weeks post dox removal) when compared to Tg + group

which resulted in 90% mortality rate by 25 week post dox treatment (**Fig. 2b**). Rescue animals (Tg ±)

also displayed rapidly improvement in: gait ataxia, body weight, muscle strength, locomotor activity, and

balance on rotarod test over the ensuing 12 week period, to a point where the treated animals were not

significantly different from controls on many tests (**Fig. 2**). Remarkably, we observed all of the six

FRDA associated clinical phenotypes tested showed significant improvement, suggesting that FRDA-

like neurological defects due to absence of the mouse *Fxn* gene can be rectified by delayed restoration

of *Fxn* (**Fig. 2**).

We next sought to determine whether the observed reversible behavioral changes in FRDAkd

mice are also accompanied by recovery of the physiological phenotype in FRDAkd mice heart, since,

changes in physiology offer attractive therapeutic targets for symptomatic and preventive treatment of

ataxia. Our results in Tg ± mice which received the dox for 12 weeks followed by 12 weeks dox removal

displayed reversal of long QT interval phenotype, when compared to Tg + mice at both 12 and 24

weeks post dox treatment (**Fig. 3a,b,e**). We observed the ventricular and posterior wall thickening only

at 24 weeks post dox treatment in Tg + animals (**Fig. 3d,f,g**), suggesting that long QT interval

phenotype is a prominent early manifestation of disease that occurs before left ventricular wall

thickness and show that correcting this aberrant physiology through activation of *Fxn* gene expression

is a potential route to therapy.

One question that intrigued us because of its striking behavioral and physiological functional

recovery is to what extent these changes represented pathological findings related to cell dysfunction

(potentially reversible) versus cell death (irreversible) recovery. Pathological and biochemical analyses

in Tg ± mice heart at post 8 weeks dox withdrawal revealed improved cardiac function displaying

reduced iron and ferritin accumulation, myocardial fibrosis, well ordered sarcomers, normal aconitase

activity and reduced mitochondrial degeneration (**Fig. 4**). Relevant to the pathogenesis of FRDA heart

and the role of iron and mitochondrial defect, it has been found that cells with these defects are

sensitized to cellular dysfunction(85, 86), and here we show this can be ameliorated by *Fxn* restoration.

In the nervous system of Tg ± mice post 8 weeks dox removal, we observed reduced vacuoles

and fewer condensed, degenerating mitochondria in DRG neurons along with several abnormal

mitochondria containing DRG neurons (**Fig. 5a,b,d,e**). However we only observed mild improvement in

myelin sheath thickness and cross section axonal size in the spinal cord of Tg ± mice during this time

period (**Fig. 5f,g**). Conversely, we observed a significant reduction in the number of vacuoles and

disrupted photoreceptors in the retina of Tg ± mice, indicating that *Fxn* restoration rescued

photoreceptor degeneration (**Fig. 5h,i**). These findings establish the principle of cellular dysfunction

reversibility in FRDAkd mouse model due to *Fxn* restoration and, therefore, raise the possibility that

some neurological and cardiac defects seen in this model and FRDA patients may not be permanent.

In line with remarkable recovery of several behavioral, physiological and pathological defects in

FRDAkd mice, we also observed that the genome-wide molecular biomarker represented by gene

expression analyses due to *Fxn* knockdown could be completely rescued after *Fxn* restoration (**Fig. 6**).

By rescuing the FXN protein levels back to the near basal level, we were able to reverse the molecular

changes completely. At post 8 weeks of dox removal after initial 12 weeks of dox treatment, we

examined the number of differentially expressed genes at false discovery rate of 5% in all tissues and

observed 100% recovery of gene expression levels in cerebellum and DRGs, and 99.95 % of reversed

gene expression levels in the heart of FRDAkd mice (**Fig. 6a,b**). These results which included several

pathways (**Fig. 6a,c,d**) that are significantly affected due to *Fxn* depletion and complete reversal (**Fig.**

6a,b) of these pathways due to *Fxn* restoration can serve as gene-expression signature for evaluation

of various therapeutic paradigm.

In summary, all six behavioral deficits in FRDAkd mice were reversed (**Fig. 2**), in the cardiac

system, the long QT interval phenotype along with various pathological mainfestations including

mitochondrial defects were reversed, in the nervous system, we observed improvement in DRGs and

photoreceptor neurons, and complete reversal of the molecular changes in all three tissues suggesting

near basal *Fxn* levels are sufficient to elevate behavioral symptoms in the preclinical FRDAkd model.

The rapidity of *Fxn* expression due to dox removal and its robust correction of various parameters, even

when restored after the onset of motor dysfunction, makes this FRDAkd mouse model an appealing

potential preclinical tool for testing various therapeutics for FRDA.

DISCUSSION

Here we report development and extensive characterization of a novel, reversible mouse model of FRDA based on knock-down of frataxin by RNA interference(28), the FRDAkd mouse. Treated transgenic mice developed abnormal mitochondria, exhibit cardiac and nervous system impairments along with physiological, pathological and molecular deficits. These abnormalities likely contributed to the behavioral phenotype of FRDAkd mice, parallel to what is observed in FRDA patients(8, 9). Importantly, restoration of *Fxn* expression, even after the development of severe symptoms and pathological defects, resulted in a drastic amelioration of the clinical phenotype, both in the cardiac and nervous systems, including motor activity. Pathology was significantly improved in the heart, DRGs and in the retina, but only mild improvement was observed in spinal cord, suggesting *Fxn* restoration for the duration of 8 weeks is not sufficient for reversal of neuronal defects in the spinal cord after *Fxn* knockdown. Interestingly, the patterns of gene expression changes due to *Fxn* knockdown in three different tissues differed(15), indicating variable response to frataxin deficiency, which could partly explain cell and tissue selectivity in FRDA(87). The onset and progression of the disease correlates with the concentration of doxycycline, and the phenotype returns to baseline after its withdrawal. The onset of transgene expression to achieve *Fxn* knockdown and robust recovery of symptoms due to restoration of *Fxn* levels in a mouse model of FRDA, even when reversed after the onset of the disease, makes this model an appealing potential preclinical tool for developing FRDA therapeutics. This approach will also enable new insights into FRDA gene function and molecular disease mechanisms.

Several models of FRDA have been developed and each have advantages and disadvantages(25). This new FRDAkd model exhibits several unique features that provide advantages for the study of FRDA pathophysiology relative to other existing models(20-25). First, induction of frataxin knockdown permits circumventing potential confounding developmental effects(19) and has the flexibility to enhance the disease onset and progression very rapidly by increasing the doxycycline

dose. Moreover, the temporal control of *Fxn* knockdown can provide further insights into the sequence

of tissue vulnerability during the disease progression(87). Second, reversibility of *Fxn* knockdown

provides a unique model to mimic the effect of an ideal therapeutic intervention. Along this line, it has

so far not been established to what extent FXN restoration after the onset of clinical motor symptoms is

successful to prevent occurrence and/or progression of FRDA. Therefore, this model will be of central

importance to gain better insights into disease pathogenesis and to test therapeutic agents. Third, the

doxycycline-dependent reduction of *Fxn* expression can be tailored to carefully determine the critical

threshold of *Fxn* levels necessary to induce selective cellular dysfunction in the nervous system (DRGs

and spinal cord) and to understand the occurrence of tissue specific dysfunction in FRDA. Thereby,

these experiments will help to understand the tissue specificity and generate clinically relevant tissue

targeted therapeutics for FRDA. Finally, the temporal control of single copy and reversible regulation of

shRNA expression against *Fxn* produces reproducible transgene expression from the well-

characterized rosa26 locus to generate the first model to exhibit and reverse several symptoms parallel

to FRDA patients. Here, we focused on the consequences of frataxin removal in an otherwise healthy

adult animal. However, we should emphasize that this FRDAkd model uniquely facilitates future studies

exploring prenatal or early post natal knockdown, and a wide variety of time course studies to

understand disease pathophysiology and to identify potential imaging, physiological, or behavioral

correlates of likely reversible or non-reversible disabilities in patients.

We show that FRDAkd animals are defective in several behaviors and exhibit weight loss,

reduced locomotor activity, reduced strength, ataxia and early mortality. All of these defects were

significantly improved following *Fxn* restoration, approaching or reaching wild-type levels. Most

importantly ataxia and survival are well-established and important clinical endpoints in FRDA(31),

readouts after *Fxn* restoration clearly improve these parameters and appear to be directly related to the

functional status of the FRDAkd mice. We conclude that *Fxn* deficient mice exhibit considerable

neurological plasticity even in a nervous system that is fully adult. Hence, utilizing these reversible

intermediate behavioral phenotypes as biomarkers will help us determine the disease progression and

test various FRDA treatment options in this model.

Hypertrophic cardiomyopathy is a common clinical feature in FRDA and approximately 60% of

patients with typical childhood onset FRDA die from cardiac failure(31). It is generally believed that

cardiac failure is caused by the loss of cardiomyocytes through activation of apoptosis(88). We

observed activation of early apoptosis pathways in heart tissue and severe cardiomyopathy

characterized by ventricular wall thickness(89). However, we did not observe TUNEL positive cells in

either heart or nervous system. This may reflect that the model is in a early phase of cell death

initiation, or rather that apoptotic cells are readily phagocytosed by neighbouring cells and are

consequently difficult to detect(90). We also observed enhanced activation of autophagy in the heart

tissue of FRDAkd mice, where autophagic cardiomyocytes are observed at a significantly higher

frequency during cardiac failure(91). These results suggest that apoptosis and autophagy together

might synergistically play a vital role in the development of cardiac defect in FRDA(92).

During *Fxn* knockdown, FRDAkd mice initially exhibited a long QT interval at 12 weeks during

electrocardiographic analyses followed by the absence of P-waves and increased ventricular wall

thickness at 24 weeks. Restoration of *Fxn* levels at 12 weeks reversed long QT interval phenotype.

However, it will be interesting to examine if the ventricular wall thickness can be restored by a more

prolonged rescue time period. Another prominent feature of *Fxn* deficiency mouse and FRDA patients

is iron accumulation and deficiency in activity of the iron-sulfur cluster dependent enzyme, aconitase, in

cardiac muscle(23, 40, 85, 86). Consistent with these observations, we observed increased iron

accumulation and reduced aconitase activity in the cardiac tissue of FRDAkd mice and we demonstrate

a marked reversal of both to a statistically significant extent, suggesting *Fxn* restoration is sufficient to

overcome and clear the iron accumulation and reverse aconitase activity(93). Our gene expression

data revealed several genes (HFE(77), SLC40A1(77), HMOX1(78), TFRC(77) and GDF15(79)) directly

involved in hemochromatosis and iron overload to be upregulated in our FRDAkd mice, all of which

were rescued to normal levels by frataxin restoration. Similarly, several downregulated genes involved

in normal cardiac function (CACNA2D1, ABCC9 and HRC) were rescued by *Fxn* restoration. Together,

these data indicate that *Fxn* restoration in symptomatic FRDAkd mice reverses the early development

of cardiomyopathy at the molecular, cellular and physiological levels.

Cellular dysfunction due to FXN deficiency is presumed to be the result of a mitochondrial

defect, since FXN localizes to mitochondria(93-95) and deficiencies of mitochondrial enzymes and

function have been observed in tissues of FRDA patients(40, 96). Here we show that FRDAkd mice

displayed accumulation of damaged mitochondria, and reduced aconitase as a direct consequence of

frataxin deficiency in heart, consistent with previous findings in conditional FRDA mouse models(23,

24), suggesting frataxin deficiency inhibits mitochondrial function leading to cellular atrophy(97).

Restoration of FXN resulted in improvement in pathological mitochondrial structure indicating that FXN

restoration prevents mitochondrial defects and may thereby enhance cell survival(93). Our observation

of mitochondrial dysfunction recovery as early as eight weeks after FXN restoration is consistent with

the highly dynamic nature of mitochondria(98).

In the nervous system, we observed higher level of condensed mitochondria in DRGs through

ultrastructural analyses. Condensed mitochondria are known to have lower respiratory control and ATP

production(99). On most occasions these condensed mitochondria in DRG neurons of FRDAkd mice

were associated with lipid-like bodies, consistent with the known close association of lipid bodies with

mitochondria observed in a variety of cell types(100). Of particular interest, it has been shown that the

junctions between these two organelles expand with an increased need for energy, suggesting the

direct flow of fatty acids from lipid bodies to the mitochondrial matrix for β-oxidation to meet the cell's

energy needs(101). We observed reduced condensed mitochondria and their association with lipid-like

bodies in rescue animals, suggesting that a substantial fraction of dysfunctioning frataxin-deficient

mitochondria containing neurons are still viable after the onset of disease and that their dysfunction can

be reversed. In the spinal cord, we observed reduction of axonal size and myelin sheath thickness in

FRDAkd animals, however after eight weeks of rescue period by FXN restoration, we observed limited

improvement, suggesting more time may be necessary for improved nervous system recovery.

Conversely, disruption of photoreceptor neurons and degenerating RPE cells in the retina displayed

robust recovery, indicating an overall morphological improvement in retinal neurons upon FXN

restoration to normal levels. These findings extend previous studies showing that many defects in

FRDA cells in vitro and cardiac function in vivo may be reversible following reintroduction of FXN(102)

by pharmacological interventions targeting aberrant signaling processes(103) or by reintroduction of

FXN by gene therapy(83).

Improved understanding of the mechanism by which FXN deficiency leads to the various

phenotypes observed in FRDA is a major goal of current research in this disorder. In this regard, gene

expression analyses identified several pathways altered, including, PPAR signaling pathway, insulin

signaling pathway, fatty acid metabolism, cell cycle, protein modification, lipid metabolism, and

carbohydrate biosynthesis, all of which have been previously associated with altered function in FRDA

patients(15, 54, 104). We also observed several immune components, namely, complement and

chemokine cascade genes being upregulated, may be implicated as a protective mechanism after *Fxn*

knockdown, and in a causal role through chronic activation of the inflammatory response(105, 106). It

will be interesting to validate these genes and pathways as potent candidate biomarkers for FRDA at

several stages during the disease progression.

Determining definitively whether the global expression changes observed in this study are

primary or secondary to *Fxn* knockdown will require further investigation by examining dense time

points immediately after *Fxn* knockdown and rescue, but our data clearly support the utility of induced

models of severe disease, whereby the consequences of gene depletion can be more cleanly

controlled to examine the molecular changes(28). Together with post-induction rescue, this study

highlights potential biomarkers and pathways of FRDA progression. For FRDA clinical trials, it will be

important to assess whether these expression changes are translatable to the human disease and

extend to other tissues that are more easily accessible than CNS or cardiac tissue. Together, these

observations suggest that this work also provides a functional genomics foundation for understanding

FRDA disease mechanism, progression and recovery.

In conclusion, our study provides multiple lines of evidence that *Fxn* knockdown in adult mice

leads to several symptoms parallel to FRDA patients. By restoring FXN levels we show reversal of

several symptoms even after significant motor defect, demonstrating the utility of this attractive model

for testing potential therapeutics, such as gene therapy(83), protein replacement therapy(102),

enhancement of mitochondrial function(107) and small molecules(108, 109). In fact, our findings

suggest that such approaches may not only enhance FXN expression and rescue downstream

molecular changes, but may too alleviate pathological and behavioral deficits associated with FRDA,

depending on the stage of the disorder.

METHODS

In vitro frataxin knockdown.

Six different shRNA sequence against the frataxin mRNA (**Supplementary Fig. 1**) were cloned into the exchange vector (proprietary material obtained from TaconicArtemis GmbH) as described in detail previously(28). N2A cells (ATCC) were transduced with exchange vectors containing the shRNA against frataxin using Fugene 6 (Promega, Madison, WI). After 24 h, cells were replated in media containing neomycin and were validated for efficient frataxin knockdown by RT-PCR and Western blotting. Mouse LW-4 (129SvEv) embryonic stem cells were cultured and the rosa26 targeting vector (from TaconicArtemis GmbH) were electroporated according to protocol described before(12). The exchange vector containing validated and selected shRNA sequence (GGATGGCGTGCTCACCATTAA) against the Fxn mRNA were electroporated to obtain positive ES cells containing shRNA expression cassette integrated into the ROSA26 locus. Correctly targeted embryonic stem cell clones were utilized to generate frataxin knockdown mice (see below).

Frataxin knockdown transgenic mice generation.

Transgenic mice were generated at UCLA Transgenic Core facility using proprietary materials obtained from TaconicArtemis GmbH (Köln, Germany). In brief, mouse LW-4 (129SvEv) embryonic stem cells with recombinase-mediated cassette exchange (RMCE) acceptor site on chromosome 6 were used for targeting insertion of distinct Tet-On frataxin shRNA expression cassettes into the ROSA26 locus(28) as depicted in **Fig. 1**. Correctly targeted embryonic stem cell clones were identified by Southern DNA blot analysis and tested for frataxin mRNA knockdown at the embryonic stem cell stage (not shown). One embryonic stem cell clone that gave acceptable mRNA knockdown were microinjected into C57BL/6J blastocysts from which chimeric mice were derived. Frataxin knockdown mice were backcrossed six generations into C57BL/6 mice.

Genotyping.

Mouse tail biopsies were collected and DNA was extracted in boiling buffer (25 mM sodium hydroxide, 0.2 mM EDTA) at 98°C for 60 min. Extracted DNA was neutralized in Tris/HCl buffer (pH5.5) and PCR was performed under the following conditions with BioMix Red (Bioline). Four primers were used in the reaction: 5'-CCATGGAATTCGAACGCTGACGTC-3', 5'-TATGGGCTATGAACTAATGACCC-3', to amplify shRNA; 5'-GAGACTCTGGCTACTCATCC-3', 5'- CCTTCAGCAAGAGCTGGGGAC-3', as genomic control. The cycling conditions for PCR amplification were: 95°C for 5 min; 95°C for 30 s, 60°C for 30 s, 72°C for 1 min, (35 cycles); 72°C for 10 min. PCR products were analyzed by gel electrophoresis using 1.5% agarose and visualized by Biospectrum imaging system (UVP).

Animal and study design.

Experiments were approved by the Animal Research Committee (ARC) of University of California, Los Angeles and were in accordance with ARC regulation. Extensive neurological and neuropsychological tests (body weight, poorly groomed fur, bald patches in the coat, absence of whiskers, wild-running, excessive grooming, freezing, hunched body posture when walking, response to object [cotton-tip swab test], visual cliff behavior analysis) were performed to ensure all animals included in the studies were healthy. Age and sex were matched between wild type (Wt) and transgenic (Tg) groups to eliminate study bias. The average age of the animals at the start of experiments was 3-4 months. Three different study cohorts were implemented: behavior, pathology and gene expression. Animals were randomly assigned to different experimental groups with in these three different study cohorts before the start of experiments. Due to high mortality rate of FRDAkd animals, we doubled the Tg + animal number (a sample size of 30 animals per Tg + treatment group for behavioral analyses (60 total)) in order to have sufficient power for statistical analyses. All the group size for our experiments were determined by statistical power analysis. The values utilized were: the power = 0.9, alpha = 0.05, Coefficient of

determination = 0.5, effect size = 0.70. Effect size and power calculations were based on our pilot

experiments.

In vivo frataxin knockdown.

Animals were divided into the following groups: Wt treated with doxycycline (Dox) (Wt +), Wt without

Dox (Wt -), Tg treated with Dox (Tg +), Tg without Dox (Tg -), Tg Dox rescue (Tg ±). First, we examined

the Tg + mice for frataxin knockdown utilizing higher dose of dox (4 and 6mg/ml), we observed mortality

as early as two weeks and a 100% mortality rate by 5 to 6 weeks (not shown). To avoid early mortality

and to have slow and steady state of disease progression we followed 2 mg/ml in drinking water

coupled with intraperitoneal injection of dox (5 or 10 mg/kg) twice per week. Doxycycline (2mg/mL) was

added to the drinking water of all treatment animals which was changed weekly. In addition, animals

were injected intraperitoneally (IP) with Dox (5 mg/kg body weight) twice a week for 10 weeks followed

by 10 mg Dox/kg body weight twice a week for 2 weeks. Animals in Tg Dox rescue (Tg ±) group were

given untreated water and not injected with Dox after week 12. All animals were weighed weekly.

Behavior cohort.

Animal information. A total of 108 animals (Wt n=32, Tg n=76) were included in this study with equal

numbers of male and female animals. For all tests, investigators were blinded to genotype and

treatment. For all behavioral tests, the variance between all the groups for that specific behavioral test

were observed to be initially not statistically significant.

Accelerating Rotarod. To measure motor function, rotarod analysis was performed weekly at the start of

Dox treatment using an accelerating rotorod (ROTOMEX-5, Columbus Instruments, Columbus, OH).

Mice were assessed for 36 weeks. Briefly, after habituation, a mouse was placed on the rotarod

rotating at 5 rpm for one min and then the rotorod was accelerated at 0.09 rpm/sec^2. The latency to fall

from a rotating rod after acceleration was recorded. Each mouse was subjected to 3 test trials within

the same day with a 15 min inter-trial interval. The average latency normalized by the mouse body

weight in the test week was used for data analysis.

Grip strength test. The grip strength was measured at week 0, 12 and 24 weeks using a digital force

gauge (Chatillon Force Measurement Systems, AMETEK TCI Division, Largo, FL). Briefly, the mouse

was allowed to grasp the steel wired grid attached to the force gauge with only forepaws. The mouse

was then pulled back from the gauge. The force applied to the grip immediately before release of the

grip is recorded as the "peak tension" and is a measurement of forepaw strength. The same

measurement was repeated with the mouse grasping the grid with all paws for whole grip strength.

Each mouse was subjected to three forepaw and whole strength measurements. The hindpaw grip

strength was calculated as the average whole grip strength minus average forepaw strength. The value

of hindpaw grip strength normalized by the mouse body weight in the test week was used for data

analysis.

Gait analyses. Gait analysis was performed at week 0, 12 and 24 weeks by allowing the animals to

walk through a 50-cm-long, 10-cm-wide runway that was lined with blank index cards. After a period of

habituation (walking through the runway three times), hind and fore paws were coated with nontoxic red

and purple paint respectively and the mouse was allowed to walk through the runway again. Footprints

were captured on the index cards. The index cards were scanned and the images were measured in

image J for calculating the stride length and other parameters.

Open field analyses. Open field testing was performed at week 0, 12 and 24 weeks. Mice were placed

in a clear plexiglass arena (27.5 cm by 27.5 cm) with a video monitoring system that recorded any

movements of the mouse within the chamber for 20 min. Locomotor activity of the mice was analyzed

by TopScan (CleverSys) software.

Pathology cohort.

Animal information. A total of 57 animals (Wt n=21 (10 females, 11 males), Tg n=36 (20 females, 16

males)) were euthanized in the study. Animals were deeply anesthetized with sodium pentobarbital (40

mg/kg body weight) and perfused intracardially with 20 mL PBS (5 mL/min) followed by 20 mL freshly

made 4% paraformaldehyde (5 mL/min). Tissue from liver, lung, spleen, pancreas, kidney, heart, eye

(retina), brain, muscle, spinal cord, dorsal root ganglion (DRG) and sciatic nerve was dissected and

collected immediately after perfusion. Collected tissue was immersed in 4% paraformaldehyde

overnight at 4°C and then transferred and kept in 40% sucrose at 4°C until embedded with O.C.T.

Tissue cryosections (5 μM) were collected with a cryostat for staining.

Staining. H & E staining was performed in Translational Pathology Core Laboratory at UCLA. The

method is described briefly below. After equilibration, the tissue section was stained with Harris

Modified Hematoxylin for 10 min, then in Eosin Y for 10 min. Gomori's iron staining and Masson's

trichrome staining were performed in Histopathology lab in UCLA, the procedure is described briefly as

below. Gomori's iron staining: the tissue section was immersed for 20 min in equal parts of 20%

hydrochloric acid and 10% potassium ferrocyanide, washed thoroughly in distilled water. After

counterstaining with Nuclear Fast Red (0.1% Kernechtrot in 5% aluminum sulfate), the slides were

dehydrated through gradual ethyl alcohol solutions and mounted for imagining. Masson's trichrome

staining: the tissue section was fixed in Bouin's fixative (75 mL picric acid saturated aqueous solution,

25 mL of 40% formaldehyde, 5 mL glacial acetic acid) for 60 min at 60°C, stained with Weigert iron

hematoxylin (0.5% hematoxylin, 47.5% alcohol, 0.58% ferric chloride, 0.5% hydrochloric acid) for 10

min, Biebrich scarlet-acid fuchsin (0.9% Biebrich scarlet, 0.1% acid fuchsin, 1% glacial acetic acid) for

10 min, immersed in phosphomolybdic-phosphotungstic solution (2.5% phosphomolybdic acid, 2.5%

phosphotungstic acid) for 7 min, stained with aniline blue solution (2.5% aniline blue, 2% acetic acid) for

7 min, dehydrate and mounted for imagining. Immunofluorescence staining was performed as followed:

cryosections were equilibrated with TBS (20 mM Tris pH7.5, 150 mM NaCl) and permeabilized in TBST

(TBS with 0.2% TritonX-100). Tissue sections were incubated with primary antibody in TBST with 4%

normal goat serum (Jackson Immuno Research) (4°C, overnight), followed by incubation with

secondary antibody in TBST with 4% normal goat serum (room temperature, 2 hrs) the next day. The

tissue section was mounted with ProLong Gold with DAPI (ThermoFisher Scientific) and stored in the

dark. The following antibodies were used: anti-ferroportin-1 (LifeSpan BioSciences, LS-B1836, rabbit,

1:200), anti-ferritin (Abcam, ab69090, rabbit, 1:500), anti-LC3 (Abgent, AM1800a, mouse, 1:200), anti-

NeuN (EMD Millipore, ABN91, chicken, 1:500), anti-NeuN (EMD Millipore, ABN78, rabbit, 1:500).

Secondary antibodies conjugated with Alexa Fluor (ThermoFisher Scientific, 1:500) were used as

indicated: Alexa Fluor 488 (goat anti-Rabbit IgG, A-11008), Alexa Fluor 594 (goat anti-chicken IgG, A-

11039), and Alexa Fluor 488 (goat anti-mouse IgG, A-11029). DNA strand breaks were determined by

TUNEL assay (Roche). Briefly, sections were equilibrated with PBS, permeabilized in 0.1% Triton X-

100 in 0.1% sodium citrate (on ice, 2 min) and incubated with TUNEL reaction mixture in a humidified

atmosphere (37°C, 60 min). Sections were mounted with ProLong Gold with DAPI and stored in the

dark.

Imaging and quantification. Slides stained for hematoxylin-eosin, iron, and trichrome were scanned (20

x magnification) by Aperio ScanScope with Aperio ImageScope Software (Leica). Immunofluorescence

staining and TUNEL staining were imaged with LSM780 confocal microscope system (Zeiss). Images

were quantified by CellProfiler. For Purkinje cell quantification, the H&E images were scanned,

visualized and exported by ImageScope (Leica Biosystems) system. Exported images were utilized to

count the Purkinje cells manually using NIH ImageJ software. Counters were blinded to genotype and

treatment.

Electron microscopy analyses. A total of 29 animals (Wt n=9 (4 females, 5 males), Tg n=20 (9 females,

11 males)) were used in the study. Mice were perfused transcardially with 2.5% glutaraldehyde, 2%

paraformaldehyde in 0.1M phosphate buffer, 0.9% sodium chloride (PBS). Pieces of heart, lumbar

spinal cord, dorsal root ganglia, muscle, and eye were dissected, postfixed for 2 hours at room

temperature in the same fixative and stored at 4°C until processing. Tissues were washed with PBS,

postfixed in 1% OsO4 in PB for 1 hour, dehydrated in a graded series of ethanol, treated with propylene

oxide and infiltrated with Eponate 12 (Ted Pella) overnight. Tissues were embedded in fresh Eponate,

and polymerized at 60°C for 48 hours. Approximately 60-70 nm thick sections were cut on a RMC

Powertome ultramicrotome and picked up on formvar coated copper grids. The sections were stained

with uranyl acetate and Reynolds lead citrate and examined on a JEOL 100CX electron microscope at

60kV. Images were collected on type 4489 EM film and the negatives scanned to create digital files.

These high quality digital images were utilized to quantify the number of condensed mitochondria.

Condensed mitochondria, vacuoles with condensed mitochondria, and vacuoles alone were manually

counted with NIH ImageJ software. Animal genotype and treatment information was blinded to the

person who conducted the evaluation.

Gene expression cohort.

Sample collection. A total of 80 animals (Wt n=24 (12 females and 12 males), Tg n=56 (31 females and

25 males)) were sacrificed and tissue was collected in this study. Animals were sacrificed at week 0, 3,

8, 12, 16, 20 and plus 4, 8 weeks post dox treatment (rescue). Mouse was sacrificed by cervical

dislocation and tissue from liver, lung, spleen, pancreas, kidney, heart, eye (retina), brain, muscle,

spinal cord, dorsal root ganglion (DRG) and sciatic nerve was dissected and rinsed in cold PBS quickly

(3X) to remove blood. Tissue samples were transferred immediately into 2 mL RNase-free tubes and

immersed into liquid nitrogen. The collected tissue was stored in -80°C freezer immediately.

RNA extraction. Heart, cerebellum and DRG neuron samples from week 0, 3, 12, 16, 20 and 4, 8

weeks post dox treatment (rescue) with four biological replicates were used for expression profiling.

Samples were randomized prior to RNA extraction to eliminate extraction batch effect. Total RNA was

extracted using the miRNeasy mini kit (Qiagen) according to manufacturer's protocol and including an

on-column DNase digest (RNase free DNAse set; Qiagen). RNA samples were immediately aliquoted

and stored at -80°C. RNA concentration and integrity were later determined using a Nanodrop

Spectrophotometer (ThermoFisher Scientific) and TapeStation 2200 (Agilent Technologies),

respectively.

Transcriptome profiling by microarray. One hundred nanograms of RNA from heart and cerebellum

tissue was amplified using the Illumina TotalPrep-96 RNA Amplification kit (ThermoFisher Scientific)

and profiled by Illumina mouse Ref 8 v2.0 expression array chips. For DRG samples 16.5 ng of RNA

was amplified using the Ovation® PicoSL WTA System V2 kit (NuGEN). Only RNA with RIN greater

than 7.0 was included for the study. A total of 64 RNA samples for each tissue (n= 192 arrays) were

included and samples were randomized before RNA amplification to eliminate microarray chip batch

effect. Raw data was log transformed and checked for outliers. Inter-array Pearson correlation and

clustering based on variance were used as quality-control measures. Quantile normalization was used

and contrast analysis of differential expression was performed by using the LIMMA package(32).

Briefly, a linear model was fitted across the dataset, contrasts of interest were extracted, and

differentially expressed genes for each contrast were selected using an empirical Bayes test

statistic(32).

Construction of co-expression networks. A weighted gene co-expression network was constructed for

each tissue dataset to identify groups of genes (modules) associated with temporal pattern of

expression changes due to frataxin knockdown and rescue following a previously described

algorithm(49, 110). Briefly, we first computed the Pearson correlation between each pair of selected

genes yielding a similarity (correlation) matrix. Next, the adjacency matrix was calculated by raising the

absolute values of the correlation matrix to a power (β) as described previously(49). The parameter β

was chosen by using the scale-free topology criterion(49), such that the resulting network connectivity

distribution best approximated scale-free topology. The adjacency matrix was then used to define a

measure of node dissimilarity, based on the topological overlap matrix, a biologically meaningful

measure of node similarity(49). Next, the probe sets were hierarchically clustered using the distance

measure and modules were determined by choosing a height cutoff for the resulting dendrogram by

using a dynamic tree-cutting algorithm(49).

Consensus module analyses. Consensus modules are defined as sets of highly connected nodes that

can be found in multiple networks generated from different datasets (tissues). Consensus modules

were identified using a suitable consensus dissimilarity that were used as input to a clustering

procedure, analogous to the procedure for identifying modules in individual sets as described

elsewhere(111). Utilizing consensus network analysis, we identified modules shared across different

tissue data sets after frataxin knockdown and calculated the first principal component of gene

expression in each module (module eigengene). Next, we correlated the module eigengenes with time

after frataxin knockdown to select modules for functional validation.

Gene Ontology, pathway and PubMed analyses. Gene ontology and pathway enrichment analysis was

performed using the DAVID platform (DAVID, https://david.ncifcrf.gov/(112)). A list of differentially

regulated transcripts for a given modules were utilized for enrichment analyses. All included terms

exhibited significant Benjamini corrected P-values for enrichment and generally contained greater than

five members per category. We used PubMatrix(113) to examine each differentially expressed gene's

association with the observed phenotypes of FRDAkd mice in the published literature by testing

association with the key-words: ataxia, cardiac fibrosis, early mortality, enlarged mitochondria, excess

iron overload, motor deficits, muscular strength, myelin sheath, neuronal degeneration, sarcomeres,

ventricular wall thickness, and weight loss in the PubMed database for every gene.

Data Availability. Datasets generated and analyzed in this study are available at Gene Expression

Omnibus. Accession number: GSE98790.

Quantitative real-time PCR.

RT-PCR was utilized to measure the mRNA expression levels of frataxin in order to identify and

validate potent shRNA sequence against frataxin gene. The procedure is briefly described below: 1.5

μg total RNA, together with 1.5 μL random primers (ThermoFisher Scientific, catalog# 48190-011), 1.5

μL 10 mM dNTP (ThermoFisher Scientific, catalog# 58875) and RNase-free water up to 19.5 μL, was

incubated at 65 °C for 5 min, then on ice for 2 min; 6 μL first strand buffer, 1.5 μL 0.1 M DTT, 1.5 μL

RNaseOUT (ThermoFisher Scientific, catalog# 100000840) and 1.5 μL SuperScript III (ThermoFisher

Scientific, catalog# 56575) were added to the mixture and incubated for 5 min at 25 °C, followed by 60

min at 50 °C, and 15 min at 70 °C. The resulted cDNA was diluted 1:10 and 3 μL was mixed with 5 μL

SensiFast SYBR No-ROX reagent (Bioline, catalog# BIO-98020), 0.6 μL primer set (forward and

reverse, 10 mM) and 1.4 μL nuclease free water for real-time PCR. Each reaction was repeated three

times on LightCycler 480 II (Roche, catalog# 05015243001). The thermocycling program is as

following: 95 °C 5min; 95 °C 10 sec, 60 °C 10 sec, 72 °C 10 sec, repeat 45 cycles; 95 °C 5 sec, 65 °C 1

min, 97 °C 5 min.

Western blotting.

Prior to samples being analyzed by Western blotting, protein concentration was estimated by Bradford assay (Bio-Rad). Proteins were separated by SDS-PAGE and transferred onto PVDF membranes. The membranes were incubated with primary and secondary antibodies for protein detection. The target bands were detected with antibodies: anti-Frataxin (Santa Cruz Biotechnology, sc-25820, rabbit, 1:200), anti-Akt (Cell Signaling Technology, #4691, rabbit, 1:500), anti-β-Actin (Sigma, #A1978, mouse, 1:1000), anti-Caspase 8 (rabbit monoclonal, Cell Signaling Technology, #8592S, rabbit, 1:200), anti-LC3 (Abgent, # AM1800a, mouse, 1:200) and goat-anti-rabbit or goat-anti-mouse HRP-conjugated secondary antibodies (ThermoFisher Scientific, #31460, and #32230, respectively,1:5000). Image J was used for band quantification.

Enzyme activity assay.

Proteins from heart tissue for all genotypes at week 20 was extracted in Tris buffer (50 mM Tris-HCl pH 7.4). Around 80 mg of tissue was homogenized with 100 μL Tris buffer, and centrifuged at 800 g for 10 min at 4°C. The supernatant was transferred to a clean tube and stored at -80°C freezer immediately for later use. Protein concentration was estimated by Bradford assay. Total of 5 μg of protein was used to determine aconitase activity by aconitase assay kit (Cayman Chemical, catalog# 705502). The absorbance of the reaction mixture at 340 nm was measured every minute for 60 min at 37°C, by Synergy-2 (BioTek) plate reader. The linear phase was used to calculate the enzyme activity. 5 μg of protein was used to measure citrate synthase activity by citrate synthase assay kit (Sigma, catalog# CS0720). The absorbance of the reaction mixture at 412 nm was measured by Synergy-2 plate reader every 31 seconds for 30 min at room temperature. The linear phase was used to calculate the enzyme activity. Aconitase activity in each sample was normalized to the citrate synthase activity from the same sample for comparison.

Echocardiography.

Echocardiography was performed on all the mice with a Siemens Acuson Sequoia C256 instrument (Siemens Medical Solutions, Mountain View, California). The mice were sedated with isoflurane vaporized in oxygen (Summit Anesthesia Solutions, Bend Oregon). Left ventricular dimensions (EDD, end diastolic dimension; ESD, end systolic dimension; PWT, posterior wall thickness; VST, ventricular septal thickness) were obtained from 2D guided M-mode and was analyzed using AccessPoint software (Freeland System LLC, Santa Fe, New Mexico) during systole and diastole. Left ventricular mass was calculated as described previously(18). Heart rate, aortic ejection time, aortic velocity and mitral inflow E & A wave amplitudes were determined from Doppler flow images. Indices of contractility such as left ventricular fractional shortening (LVFS), velocity of circumferential fiber shortening (VCF) and ejection fraction (EF) was obtained from the images. During the procedure, heart rates were maintained at physiological levels by monitoring their electrocardiograms (ECG).

Surface ECG.

Electrocardiogram (ECG) were obtained in all the mice under Isoflurane anesthesia by inserting two Pt needle electrodes (Grass Technologies, West Warwick, RI) under the skin in the lead II configuration as described previously(27). The mice were studied between 10 and 30 minutes to elucidate any rhythm alteration. The ECG data were amplified (Grass Technologies) and then digitized with HEM V4.2 software (Notocord Systems, Croissy sur Seine, France). Measurements of HR, PR, RR, QRS, QT intervals were obtained for comparison and statistical analysis among all the mice groups.

Accession codes. Gene Expression Omnibus: Datasets generated and analyzed in this study are available at GEO accession: GSE98790.

ACKNOWLEDGMENTS

We gratefully acknowledge support from the Friedreich's Ataxia Research Alliance to D.H.G. and V.C.

(including a New Investigator Award to V.C.), and the Dr. Miriam and Sheldon G. Adelson Medical

Research Foundation to D.H.G.. We thank Marianne Cilluffo for assistance with electron microscopy at

the BRI Electron Microscopy Core Facility at UCLA.

AUTHOR CONTRIBUTIONS

V.C. and D.H.G. designed the experiments and wrote the manuscript. K.G., R.V., H.D., and V.C.

performed the behavioral, pathological, biochemical and molecular experiments. V.C. conducted or

coordinated all the experiments discussed in this manuscript with input from D.H.G.. V.S. performed

network analyses and V.C. performed all subsequent bioinformatic analyses. M.J. and V.C. conducted

and coordinated electrocardiogram and echocardiogram analyses. All authors discussed the results

and provided comments and revisions on the manuscript.

REFERENCES:

1.   Chamberlain S, Shaw J, Rowland A, Wallis J, South S, Nakamura Y, von Gabain A, Farrall M, Williamson R. Mapping of mutation causing Friedreich's ataxia to human chromosome 9. Nature. 1988 Jul 21;334(6179):248-50. PubMed PMID: 2899844.

2.   Campuzano V, Montermini L, Molto MD, Pianese L, Cossee M, Cavalcanti F, Monros E, Rodius F, Duclos F, Monticelli A, Zara F, Canizares J, Koutnikova H, Bidichandani SI, Gellera C, Brice A, Trouillas P, De Michele G, Filla A, De Frutos R, Palau F, Patel PI, Di Donato S, Mandel JL, Cocozza S, Koenig M, Pandolfo M. Friedreich's ataxia: autosomal recessive disease caused by an intronic GAA triplet repeat expansion. Science. 1996 Mar 8;271(5254):1423-7. PubMed PMID: 8596916.

3.   Campuzano V, Montermini L, Lutz Y, Cova L, Hindelang C, Jiralerspong S, Trottier Y, Kish SJ, Faucheux B, Trouillas P, Authier FJ, Durr A, Mandel JL, Vescovi A, Pandolfo M, Koenig M. Frataxin is reduced in Friedreich ataxia patients and is associated with mitochondrial membranes. Human molecular genetics. 1997 Oct;6(11):1771-80. PubMed PMID: 9302253.

4.   Filla A, De Michele G, Cavalcanti F, Pianese L, Monticelli A, Campanella G, Cocozza S. The relationship between trinucleotide (GAA) repeat length and clinical features in Friedreich ataxia. American journal of human genetics. 1996 Sep;59(3):554-60. PubMed PMID: 8751856. Pubmed Central PMCID: 1914893.

5.   Montermini L, Richter A, Morgan K, Justice CM, Julien D, Castellotti B, Mercier J, Poirier J, Capozzoli F, Bouchard JP, Lemieux B, Mathieu J, Vanasse M, Seni MH, Graham G, Andermann F, Andermann E, Melancon SB, Keats BJ, Di Donato S, Pandolfo M. Phenotypic variability in Friedreich ataxia: role of the associated GAA triplet repeat expansion. Annals of neurology. 1997 May;41(5):675-82. PubMed PMID: 9153531.

6.   Andermann E, Remillard GM, Goyer C, Blitzer L, Andermann F, Barbeau A. Genetic and family studies in Friedreich's ataxia. The Canadian journal of neurological sciences Le journal canadien des sciences neurologiques. 1976 Nov;3(4):287-301. PubMed PMID: 1000412.

7.   Harding AE. Friedreich's ataxia: a clinical and genetic study of 90 families with an analysis of early diagnostic criteria and intrafamilial clustering of clinical features. Brain : a journal of neurology. 1981 Sep;104(3):589-620. PubMed PMID: 7272714.

8.   Koeppen AH, Mazurkiewicz JE. Friedreich ataxia: neuropathology revised. Journal of neuropathology and experimental neurology. 2013 Feb;72(2):78-90. PubMed PMID: 23334592. Pubmed Central PMCID: 3817014.

9.   Koeppen AH. Friedreich's ataxia: pathology, pathogenesis, and molecular genetics. Journal of the neurological sciences. 2011 Apr 15;303(1-2):1-12. PubMed PMID: 21315377. Pubmed Central PMCID: 3062632.

10.   Bhidayasiri R, Perlman SL, Pulst SM, Geschwind DH. Late-onset Friedreich ataxia: phenotypic analysis, magnetic resonance imaging findings, and review of the literature. Archives of neurology. 2005 Dec;62(12):1865-9. PubMed PMID: 16344344.

11.   Li Y, Lu Y, Polak U, Lin K, Shen J, Farmer J, Seyer L, Bhalla AD, Rozwadowska N, Lynch DR, Butler JS, Napierala M. Expanded GAA repeats impede transcription elongation through the FXN gene and induce transcriptional silencing that is restricted to the FXN locus. Human molecular genetics. 2015 Dec 15;24(24):6932-43. PubMed PMID: 26401053. Pubmed Central PMCID: 4654050.

12.   Seibler J, Kuter-Luks B, Kern H, Streu S, Plum L, Mauer J, Kuhn R, Bruning JC, Schwenk F. Single copy shRNA configuration for ubiquitous gene knockdown in mice. Nucleic acids research. 2005 Apr 14;33(7):e67. PubMed PMID: 15831785. Pubmed Central PMCID: 1079974.

13.   Martinez AR, Moro A, Abrahao A, Faber I, Borges CR, Rezende TJ, Martins CR, Jr., Moscovich
M, Munhoz RP, Segal SL, Arruda WO, Saraiva-Pereira ML, Karuta S, Pedroso JL, D'Abreu A, Jardim
LB, Lopes-Cendes I, Barsottini OG, Teive HA, Franca MC, Jr. Nonneurological Involvement in Late-
Onset Friedreich Ataxia (LOFA): Exploring the Phenotypes. Cerebellum. 2016 Jan 11. PubMed PMID:
26754264.
14.   Galea CA, Huq A, Lockhart PJ, Tai G, Corben LA, Yiu EM, Gurrin LC, Lynch DR, Gelbard S,
Durr A, Pousset F, Parkinson M, Labrum R, Giunti P, Perlman SL, Delatycki MB, Evans-Galea MV.
Compound heterozygous FXN mutations and clinical outcome in friedreich ataxia. Annals of neurology.
2016 Mar;79(3):485-95. PubMed PMID: 26704351.
15.   Coppola G, Marmolino D, Lu D, Wang Q, Cnop M, Rai M, Acquaviva F, Cocozza S, Pandolfo
M, Geschwind DH. Functional genomic analysis of frataxin deficiency reveals tissue-specific alterations
and identifies the PPARgamma pathway as a therapeutic target in Friedreich's ataxia. Human molecular
genetics. 2009 Jul 01;18(13):2452-61. PubMed PMID: 19376812. Pubmed Central PMCID: 2694693.
16.   Pandolfo M. The molecular basis of Friedreich ataxia. Advances in experimental medicine and
biology. 2002;516:99-118. PubMed PMID: 12611437.
17.   Abrahao A, Pedroso JL, Braga-Neto P, Bor-Seng-Shu E, de Carvalho Aguiar P, Barsottini OG.
Milestones in Friedreich ataxia: more than a century and still learning. Neurogenetics. 2015
Jul;16(3):151-60. PubMed PMID: 25662948.
18.   Tanaka N, Dalton N, Mao L, Rockman HA, Peterson KL, Gottshall KR, Hunter JJ, Chien KR,
Ross J, Jr. Transthoracic echocardiography in models of cardiac disease in the mouse. Circulation. 1996
Sep 01;94(5):1109-17. PubMed PMID: 8790053.
19.   Cossee M, Puccio H, Gansmuller A, Koutnikova H, Dierich A, LeMeur M, Fischbeck K, Dolle
P, Koenig M. Inactivation of the Friedreich ataxia mouse gene leads to early embryonic lethality without
iron accumulation. Human molecular genetics. 2000 May 1;9(8):1219-26. PubMed PMID: 10767347.
20.   Miranda CJ, Santos MM, Ohshima K, Smith J, Li L, Bunting M, Cossee M, Koenig M,
Sequeiros J, Kaplan J, Pandolfo M. Frataxin knockin mouse. FEBS letters. 2002 Feb 13;512(1-3):291-7.
PubMed PMID: 11852098.
21.   Al-Mahdawi S, Pinto RM, Ruddle P, Carroll C, Webster Z, Pook M. GAA repeat instability in
Friedreich ataxia YAC transgenic mice. Genomics. 2004 Aug;84(2):301-10. PubMed PMID: 15233994.
22.   Al-Mahdawi S, Pinto RM, Varshney D, Lawrence L, Lowrie MB, Hughes S, Webster Z, Blake J,
Cooper JM, King R, Pook MA. GAA repeat expansion mutation mouse models of Friedreich ataxia
exhibit oxidative stress leading to progressive neuronal and cardiac pathology. Genomics. 2006
Nov;88(5):580-90. PubMed PMID: 16919418. Pubmed Central PMCID: 2842930.
23.   Puccio H, Simon D, Cossee M, Criqui-Filipe P, Tiziano F, Melki J, Hindelang C, Matyas R,
Rustin P, Koenig M. Mouse models for Friedreich ataxia exhibit cardiomyopathy, sensory nerve defect
and Fe-S enzyme deficiency followed by intramitochondrial iron deposits. Nature genetics. 2001
Feb;27(2):181-6. PubMed PMID: 11175786.
24.   Simon D, Seznec H, Gansmuller A, Carelle N, Weber P, Metzger D, Rustin P, Koenig M, Puccio
H. Friedreich ataxia mouse models with progressive cerebellar and sensory ataxia reveal autophagic
neurodegeneration in dorsal root ganglia. The Journal of neuroscience : the official journal of the
Society for Neuroscience. 2004 Feb 25;24(8):1987-95. PubMed PMID: 14985441.
25.   Perdomini M, Hick A, Puccio H, Pook MA. Animal and cellular models of Friedreich ataxia.
Journal of neurochemistry. 2013 Aug;126 Suppl 1:65-79. PubMed PMID: 23859342.
26.   Metz G, Coppard N, Cooper JM, Delatycki MB, Durr A, Di Prospero NA, Giunti P, Lynch DR,
Schulz JB, Rummey C, Meier T. Rating disease progression of Friedreich's ataxia by the International

Cooperative Ataxia Rating Scale: analysis of a 603-patient database. Brain : a journal of neurology. 2013 Jan;136(Pt 1):259-68. PubMed PMID: 23365101. Pubmed Central PMCID: 3624678.

27.   Nakashima Y, Yanez DA, Touma M, Nakano H, Jaroszewicz A, Jordan MC, Pellegrini M, Roos KP, Nakano A. Nkx2-5 suppresses the proliferation of atrial myocytes and conduction system. Circulation research. 2014 Mar 28;114(7):1103-13. PubMed PMID: 24563458.

28.   Seibler J, Kleinridders A, Kuter-Luks B, Niehaves S, Bruning JC, Schwenk F. Reversible gene knockdown in mice using a tight, inducible shRNA expression system. Nucleic acids research. 2007;35(7):e54. PubMed PMID: 17376804. Pubmed Central PMCID: 1874634.

29.   Crawley JN. What's wrong with my mouse? : behavioral phenotyping of transgenic and knockout mice. 2nd ed. Hoboken, N.J.: Wiley-Interscience; 2007. xvi, 523 p. p.

30.   Dellon ES, Dellon AL. Functional assessment of neurologic impairment: track analysis in diabetic and compression neuropathies. Plastic and reconstructive surgery. 1991 Oct;88(4):686-94. PubMed PMID: 1896540.

31.   Tsou AY, Paulsen EK, Lagedrost SJ, Perlman SL, Mathews KD, Wilmot GR, Ravina B, Koeppen AH, Lynch DR. Mortality in Friedreich ataxia. Journal of the neurological sciences. 2011 Aug 15;307(1-2):46-9. PubMed PMID: 21652007.

32.   Smyth GK. limma: Linear Models for Microarray Data. In: Gentleman R, Carey VJ, Huber W, Irizarry RA, Dudoit S, editors. Bioinformatics and Computational Biology Solutions Using R and Bioconductor. New York, NY: Springer New York; 2005. p. 397-420.

33.   Surawicz B, Knoebel SB. Long QT: good, bad or indifferent? Journal of the American College of Cardiology. 1984 Aug;4(2):398-413. PubMed PMID: 6145737.

34.   German DM, Kabir MM, Dewland TA, Henrikson CA, Tereshchenko LG. Atrial Fibrillation Predictors: Importance of the Electrocardiogram. Annals of noninvasive electrocardiology : the official journal of the International Society for Holter and Noninvasive Electrocardiology, Inc. 2016 Jan;21(1):20-9. PubMed PMID: 26523405. Pubmed Central PMCID: 4728003.

35.   Dutka DP, Donnelly JE, Nihoyannopoulos P, Oakley CM, Nunez DJ. Marked variation in the cardiomyopathy associated with Friedreich's ataxia. Heart. 1999 Feb;81(2):141-7. PubMed PMID: 9922348. Pubmed Central PMCID: 1728941.

36.   Isnard R, Kalotka H, Durr A, Cossee M, Schmitt M, Pousset F, Thomas D, Brice A, Koenig M, Komajda M. Correlation between left ventricular hypertrophy and GAA trinucleotide repeat length in Friedreich's ataxia. Circulation. 1997 May 06;95(9):2247-9. PubMed PMID: 9142000.

37.   Durr A, Cossee M, Agid Y, Campuzano V, Mignard C, Penet C, Mandel JL, Brice A, Koenig M. Clinical and genetic abnormalities in patients with Friedreich's ataxia. The New England journal of medicine. 1996 Oct 17;335(16):1169-75. PubMed PMID: 8815938.

38.   Conrad CH, Brooks WW, Hayes JA, Sen S, Robinson KG, Bing OH. Myocardial fibrosis and stiffness with hypertrophy and heart failure in the spontaneously hypertensive rat. Circulation. 1995 Jan 1;91(1):161-70. PubMed PMID: 7805198.

39.   Bradley JL, Blake JC, Chamberlain S, Thomas PK, Cooper JM, Schapira AH. Clinical, biochemical and molecular genetic correlations in Friedreich's ataxia. Human molecular genetics. 2000 Jan 22;9(2):275-82. PubMed PMID: 10607838.

40.   Rotig A, de Lonlay P, Chretien D, Foury F, Koenig M, Sidi D, Munnich A, Rustin P. Aconitase and mitochondrial iron-sulphur protein deficiency in Friedreich ataxia. Nature genetics. 1997 Oct;17(2):215-7. PubMed PMID: 9326946.

41.   Seyer LA, Galetta K, Wilson J, Sakai R, Perlman S, Mathews K, Wilmot GR, Gomez CM, Ravina B, Zesiewicz T, Bushara KO, Subramony SH, Ashizawa T, Delatycki MB, Brocht A, Balcer LJ,

Lynch DR. Analysis of the visual system in Friedreich ataxia. Journal of neurology. 2013 Sep;260(9):2362-9. PubMed PMID: 23775342.

42. Fortuna F, Barboni P, Liguori R, Valentino ML, Savini G, Gellera C, Mariotti C, Rizzo G, Tonon C, Manners D, Lodi R, Sadun AA, Carelli V. Visual system involvement in patients with Friedreich's ataxia. Brain : a journal of neurology. 2009 Jan;132(Pt 1):116-23. PubMed PMID: 18931386.

43. Saxena S, Srivastav K, Cheung CM, Ng JY, Lai TY. Photoreceptor inner segment ellipsoid band integrity on spectral domain optical coherence tomography. Clinical ophthalmology. 2014;8:2507-22. PubMed PMID: 25525329. Pubmed Central PMCID: 4266419.

44. Coppola G, Burnett R, Perlman S, Versano R, Gao F, Plasterer H, Rai M, Sacca F, Filla A, Lynch DR, Rusche JR, Gottesfeld JM, Pandolfo M, Geschwind DH. A gene expression phenotype in lymphocytes from Friedreich ataxia patients. Annals of neurology. 2011 Nov;70(5):790-804. PubMed PMID: 22162061. Pubmed Central PMCID: 3646419.

45. Cartier L, Hartley O, Dubois-Dauphin M, Krause KH. Chemokine receptors in the central nervous system: role in brain inflammation and neurodegenerative diseases. Brain research Brain research reviews. 2005 Feb;48(1):16-42. PubMed PMID: 15708626.

46. Leszek J, Barreto GE, Gasiorowski K, Koutsouraki E, Avila-Rodrigues M, Aliev G. Inflammatory Mechanisms and Oxidative Stress as Key Factors Responsible for Progression of Neurodegeneration: Role of Brain Innate Immune System. CNS & neurological disorders drug targets. 2016;15(3):329-36. PubMed PMID: 26831258.

47. Fung TC, Olson CA, Hsiao EY. Interactions between the microbiota, immune and nervous systems in health and disease. Nature neuroscience. 2017 Jan 16. PubMed PMID: 28092661.

48. Andreasson KI, Bachstetter AD, Colonna M, Ginhoux F, Holmes C, Lamb B, Landreth G, Lee DC, Low D, Lynch MA, Monsonego A, O'Banion MK, Pekny M, Puschmann T, Russek-Blum N, Sandusky LA, Selenica ML, Takata K, Teeling J, Town T, Van Eldik LJ. Targeting innate immunity for neurodegenerative disorders of the central nervous system. Journal of neurochemistry. 2016 Sep;138(5):653-93. PubMed PMID: 27248001.

49. Zhang B, Horvath S. A general framework for weighted gene co-expression network analysis. Statistical applications in genetics and molecular biology. 2005;4:Article17. PubMed PMID: 16646834.

50. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. BMC bioinformatics. 2008;9:559. PubMed PMID: 19114008. Pubmed Central PMCID: 2631488.

51. Geschwind DH, Konopka G. Neuroscience in the era of functional genomics and systems biology. Nature. 2009 Oct 15;461(7266):908-15. PubMed PMID: 19829370. Pubmed Central PMCID: 3645852.

52. Zhang B, Gaiteri C, Bodea LG, Wang Z, McElwee J, Podtelezhnikov AA, Zhang C, Xie T, Tran L, Dobrin R, Fluder E, Clurman B, Melquist S, Narayanan M, Suver C, Shah H, Mahajan M, Gillis T, Mysore J, MacDonald ME, Lamb JR, Bennett DA, Molony C, Stone DJ, Gudnason V, Myers AJ, Schadt EE, Neumann H, Zhu J, Emilsson V. Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. Cell. 2013 Apr 25;153(3):707-20. PubMed PMID: 23622250. Pubmed Central PMCID: 3677161.

53. Parikshak NN, Swarup V, Belgard TG, Irimia M, Ramaswami G, Gandal MJ, Hartl C, Leppa V, Ubieta LT, Huang J, Lowe JK, Blencowe BJ, Horvath S, Geschwind DH. Genome-wide changes in lncRNA, splicing, and regional gene expression patterns in autism. Nature. 2016 Dec 05. PubMed PMID: 27919067.

54.   Haugen AC, Di Prospero NA, Parker JS, Fannin RD, Chou J, Meyer JN, Halweg C, Collins JB,
Durr A, Fischbeck K, Van Houten B. Altered gene expression and DNA damage in peripheral blood
cells from Friedreich's ataxia patients: cellular model of pathology. PLoS genetics. 2010 Jan
15;6(1):e1000812. PubMed PMID: 20090835. Pubmed Central PMCID: 2799513.
55.   Aiyaz M, Lupton MK, Proitsi P, Powell JF, Lovestone S. Complement activation as a biomarker
for Alzheimer's disease. Immunobiology. 2012 Feb;217(2):204-15. PubMed PMID: 21856034.
56.   Risgaard B, Jabbari R, Refsgaard L, Holst AG, Haunso S, Sadjadieh A, Winkel BG, Olesen MS,
Tfelt-Hansen J. High prevalence of genetic variants previously associated with Brugada syndrome in
new exome data. Clinical genetics. 2013 Nov;84(5):489-95. PubMed PMID: 23414114.
57.   Bienengraeber M, Olson TM, Selivanov VA, Kathmann EC, O'Cochlain F, Gao F, Karger AB,
Ballew JD, Hodgson DM, Zingman LV, Pang YP, Alekseev AE, Terzic A. ABCC9 mutations identified
in human dilated cardiomyopathy disrupt catalytic KATP channel gating. Nature genetics. 2004
Apr;36(4):382-7. PubMed PMID: 15034580. Pubmed Central PMCID: 1995438.
58.   Singh VP, Rubinstein J, Arvanitis DA, Ren X, Gao X, Haghighi K, Gilbert M, Iyer VR, Kim
DH, Cho C, Jones K, Lorenz JN, Armstrong CF, Wang HS, Gyorke S, Kranias EG. Abnormal calcium
cycling and cardiac arrhythmias associated with the human Ser96Ala genetic variant of histidine-rich
calcium-binding protein. Journal of the American Heart Association. 2013 Oct 14;2(5):e000460.
PubMed PMID: 24125847. Pubmed Central PMCID: 3835262.
59.   Gonzalez A, Fortuno MA, Querejeta R, Ravassa S, Lopez B, Lopez N, Diez J. Cardiomyocyte
apoptosis in hypertensive cardiomyopathy. Cardiovascular research. 2003 Sep 01;59(3):549-62. PubMed
PMID: 14499856.
60.   Huang ML, Becker EM, Whitnall M, Suryo Rahmanto Y, Ponka P, Richardson DR. Elucidation
of the mechanism of mitochondrial iron loading in Friedreich's ataxia by analysis of a mouse mutant.
Proceedings of the National Academy of Sciences of the United States of America. 2009 Sep
22;106(38):16381-6. PubMed PMID: 19805308. Pubmed Central PMCID: 2752539.
61.   Bolinches-Amoros A, Molla B, Pla-Martin D, Palau F, Gonzalez-Cabo P. Mitochondrial
dysfunction induced by frataxin deficiency is associated with cellular senescence and abnormal calcium
metabolism. Frontiers in cellular neuroscience. 2014;8:124. PubMed PMID: 24860428. Pubmed Central
PMCID: 4026758.
62.   Muzio M, Chinnaiyan AM, Kischkel FC, O'Rourke K, Shevchenko A, Ni J, Scaffidi C, Bretz JD,
Zhang M, Gentz R, Mann M, Krammer PH, Peter ME, Dixit VM. FLICE, a novel FADD-homologous
ICE/CED-3-like protease, is recruited to the CD95 (Fas/APO-1) death--inducing signaling complex.
Cell. 1996 Jun 14;85(6):817-27. PubMed PMID: 8681377.
63.   Kyrylkova K, Kyryachenko S, Leid M, Kioussi C. Detection of apoptosis by TUNEL assay.
Methods in molecular biology. 2012;887:41-7. PubMed PMID: 22566045.
64.   Trugenberger CA, Walti C, Peregrim D, Sharp ME, Bureeva S. Discovery of novel biomarkers
and phenotypes by semantic technologies. BMC bioinformatics. 2013 Feb 13;14:51. PubMed PMID:
23402646. Pubmed Central PMCID: 3605201.
65.   Vermeer S, Hoischen A, Meijer RP, Gilissen C, Neveling K, Wieskamp N, de Brouwer A,
Koenig M, Anheim M, Assoum M, Drouot N, Todorovic S, Milic-Rasic V, Lochmuller H, Stevanin G,
Goizet C, David A, Durr A, Brice A, Kremer B, van de Warrenburg BP, Schijvenaars MM, Heister A,
Kwint M, Arts P, van der Wijst J, Veltman J, Kamsteeg EJ, Scheffer H, Knoers N. Targeted next-
generation sequencing of a 12.5 Mb homozygous region reveals ANO10 mutations in patients with
autosomal-recessive cerebellar ataxia. American journal of human genetics. 2010 Dec 10;87(6):813-9.
PubMed PMID: 21092923. Pubmed Central PMCID: 2997370.

66.   Miskovic ND, Domingo A, Dobricic V, Max C, Braenne I, Petrovic I, Grutz K, Pawlack H,
Tournev I, Kalaydjieva L, Svetel M, Lohmann K, Kostic VS, Westenberger A. Seemingly dominant
inheritance of a recessive ANO10 mutation in romani families with cerebellar ataxia. Movement
disorders : official journal of the Movement Disorder Society. 2016 Oct 27. PubMed PMID: 27787937.
67.   Mollet J, Delahodde A, Serre V, Chretien D, Schlemmer D, Lombes A, Boddaert N, Desguerre I,
de Lonlay P, de Baulny HO, Munnich A, Rotig A. CABC1 gene mutations cause ubiquinone deficiency
with cerebellar ataxia and seizures. American journal of human genetics. 2008 Mar;82(3):623-30.
PubMed PMID: 18319072. Pubmed Central PMCID: 2427298.
68.   Gerards M, van den Bosch B, Calis C, Schoonderwoerd K, van Engelen K, Tijssen M, de Coo R,
van der Kooi A, Smeets H. Nonsense mutations in CABC1/ADCK3 cause progressive cerebellar ataxia
and atrophy. Mitochondrion. 2010 Aug;10(5):510-5. PubMed PMID: 20580948.
69.   Schicks J, Synofzik M, Schulte C, Schols L. POLG, but not PEO1, is a frequent cause of
cerebellar ataxia in Central Europe. Movement disorders : official journal of the Movement Disorder
Society. 2010 Nov 15;25(15):2678-82. PubMed PMID: 20803511.
70.   Synofzik M, Srulijes K, Godau J, Berg D, Schols L. Characterizing POLG ataxia: clinics,
electrophysiology and imaging. Cerebellum. 2012 Dec;11(4):1002-11. PubMed PMID: 22528963.
71.   Pennacchio LA, Bouley DM, Higgins KM, Scott MP, Noebels JL, Myers RM. Progressive
ataxia, myoclonic epilepsy and cerebellar apoptosis in cystatin B-deficient mice. Nature genetics. 1998
Nov;20(3):251-8. PubMed PMID: 9806543.
72.   Singh S, Ganesh S. Lafora progressive myoclonus epilepsy: a meta-analysis of reported
mutations in the first decade following the discovery of the EPM2A and NHLRC1 genes. Human
mutation. 2009 May;30(5):715-23. PubMed PMID: 19267391.
73.   Matthijs G, Schollen E, Pardon E, Veiga-Da-Cunha M, Jaeken J, Cassiman JJ, Van Schaftingen
E. Mutations in PMM2, a phosphomannomutase gene on chromosome 16p13, in carbohydrate-deficient
glycoprotein type I syndrome (Jaeken syndrome). Nature genetics. 1997 May;16(1):88-92. PubMed
PMID: 9140401.
74.   Sygitowicz G, Tomaniak M, Filipiak KJ, Koltowski L, Sitkiewicz D. Galectin-3 in Patients with
Acute Heart Failure: Preliminary Report on First Polish Experience. Advances in clinical and
experimental medicine : official organ Wroclaw Medical University. 2016 Jul-Aug;25(4):617-23.
PubMed PMID: 27629834.
75.   Salvador AM, Nevers T, Velazquez F, Aronovitz M, Wang B, Abadia Molina A, Jaffe IZ, Karas
RH, Blanton RM, Alcaide P. Intercellular Adhesion Molecule 1 Regulates Left Ventricular Leukocyte
Infiltration, Cardiac Remodeling, and Function in Pressure Overload-Induced Heart Failure. Journal of
the American Heart Association. 2016 Mar 15;5(3):e003126. PubMed PMID: 27068635. Pubmed
Central PMCID: 4943280.
76.   Polyakova V, Loeffler I, Hein S, Miyagawa S, Piotrowska I, Dammer S, Risteli J, Schaper J,
Kostin S. Fibrosis in endstage human heart failure: severe changes in collagen metabolism and
MMP/TIMP profiles. International journal of cardiology. 2011 Aug 18;151(1):18-33. PubMed PMID:
20546954.
77.   Del-Castillo-Rueda A, Moreno-Carralero MI, Cuadrado-Grande N, Alvarez-Sala-Walther LA,
Enriquez-de-Salamanca R, Mendez M, Moran-Jimenez MJ. Mutations in the HFE, TFR2, and SLC40A1
genes in patients with hemochromatosis. Gene. 2012 Oct 15;508(1):15-20. PubMed PMID: 22890139.
78.   Song W, Zukor H, Lin SH, Liberman A, Tavitian A, Mui J, Vali H, Fillebeen C, Pantopoulos K,
Wu TD, Guerquin-Kern JL, Schipper HM. Unregulated brain iron deposition in transgenic mice over-

expressing HMOX1 in the astrocytic compartment. Journal of neurochemistry. 2012 Oct;123(2):325-36. PubMed PMID: 22881289.

79.   Cui R, Gale RP, Zhu G, Xu Z, Qin T, Zhang Y, Huang G, Li B, Fang L, Zhang H, Pan L, Hu N, Qu S, Xiao Z. Serum iron metabolism and erythropoiesis in patients with myelodysplastic syndrome not receiving RBC transfusions. Leukemia research. 2014 May;38(5):545-50. PubMed PMID: 24598841. Pubmed Central PMCID: 4439195.

80.   Pavel M, Rubinsztein DC. Mammalian autophagy and the plasma membrane. The FEBS journal. 2016 Oct 18. PubMed PMID: 27758042.

81.   Bento CF, Renna M, Ghislat G, Puri C, Ashkenazi A, Vicinanza M, Menzies FM, Rubinsztein DC. Mammalian Autophagy: How Does It Work? Annual review of biochemistry. 2016 Jun 02;85:685-713. PubMed PMID: 26865532.

82.   Gustafsson AB, Gottlieb RA. Autophagy in ischemic heart disease. Circulation research. 2009 Jan 30;104(2):150-8. PubMed PMID: 19179668. Pubmed Central PMCID: 2765251.

83.   Perdomini M, Belbellaa B, Monassier L, Reutenauer L, Messaddeq N, Cartier N, Crystal RG, Aubourg P, Puccio H. Prevention and reversal of severe mitochondrial cardiomyopathy by gene therapy in a mouse model of Friedreich's ataxia. Nature medicine. 2014 May;20(5):542-7. PubMed PMID: 24705334.

84.   Gottesfeld JM, Rusche JR, Pandolfo M. Increasing frataxin gene expression with histone deacetylase inhibitors as a therapeutic approach for Friedreich's ataxia. Journal of neurochemistry. 2013 Aug;126 Suppl 1:147-54. PubMed PMID: 23859350. Pubmed Central PMCID: 3766837.

85.   Delatycki MB, Camakaris J, Brooks H, Evans-Whipp T, Thorburn DR, Williamson R, Forrest SM. Direct evidence that mitochondrial iron accumulation occurs in Friedreich ataxia. Annals of neurology. 1999 May;45(5):673-5. PubMed PMID: 10319894.

86.   Michael S, Petrocine SV, Qian J, Lamarche JB, Knutson MD, Garrick MD, Koeppen AH. Iron and iron-responsive proteins in the cardiomyopathy of Friedreich's ataxia. Cerebellum. 2006;5(4):257-67. PubMed PMID: 17134988.

87.   Lynch DR, Deutsch EC, Wilson RB, Tennekoon G. Unanswered questions in Friedreich ataxia. Journal of child neurology. 2012 Sep;27(9):1223-9. PubMed PMID: 22832776. Pubmed Central PMCID: 3674553.

88.   Fujita T, Ishikawa Y. Apoptosis in heart failure. -The role of the beta-adrenergic receptor-mediated signaling pathway and p53-mediated signaling pathway in the apoptosis of cardiomyocytes. Circulation journal : official journal of the Japanese Circulation Society. 2011;75(8):1811-8. PubMed PMID: 21747195.

89.   Bennett MR. Apoptosis in the cardiovascular system. Heart. 2002 May;87(5):480-7. PubMed PMID: 11997428. Pubmed Central PMCID: 1767116.

90.   Ravichandran KS. Beginnings of a good apoptotic meal: the find-me and eat-me signaling pathways. Immunity. 2011 Oct 28;35(4):445-55. PubMed PMID: 22035837. Pubmed Central PMCID: 3241945.

91.   Martinet W, Knaapen MW, Kockx MM, De Meyer GR. Autophagy in cardiovascular disease. Trends in molecular medicine. 2007 Nov;13(11):482-91. PubMed PMID: 18029229.

92.   Eisenberg-Lerner A, Bialik S, Simon HU, Kimchi A. Life and death partners: apoptosis, autophagy and the cross-talk between them. Cell death and differentiation. 2009 Jul;16(7):966-75. PubMed PMID: 19325568.

93.   Tan G, Chen LS, Lonnerdal B, Gellera C, Taroni FA, Cortopassi GA. Frataxin expression
rescues mitochondrial dysfunctions in FRDA cells. Human molecular genetics. 2001 Sep
15;10(19):2099-107. PubMed PMID: 11590127.
94.   Koutnikova H, Campuzano V, Foury F, Dolle P, Cazzalini O, Koenig M. Studies of human,
mouse and yeast homologues indicate a mitochondrial function for frataxin. Nature genetics. 1997
Aug;16(4):345-51. PubMed PMID: 9241270.
95.   Foury F, Cazzalini O. Deletion of the yeast homologue of the human gene associated with
Friedreich's ataxia elicits iron accumulation in mitochondria. FEBS letters. 1997 Jul 14;411(2-3):373-7.
PubMed PMID: 9271239.
96.   Lodi R, Cooper JM, Bradley JL, Manners D, Styles P, Taylor DJ, Schapira AH. Deficit of in
vivo mitochondrial ATP production in patients with Friedreich ataxia. Proceedings of the National
Academy of Sciences of the United States of America. 1999 Sep 28;96(20):11492-5. PubMed PMID:
10500204. Pubmed Central PMCID: 18061.
97.   Gonzalez-Cabo P, Palau F. Mitochondrial pathophysiology in Friedreich's ataxia. Journal of
neurochemistry. 2013 Aug;126 Suppl 1:53-64. PubMed PMID: 23859341.
98.   Chan DC. Fusion and fission: interlinked processes critical for mitochondrial health. Annual
review of genetics. 2012;46:265-87. PubMed PMID: 22934639.
99.   Desagher S, Martinou JC. Mitochondria as the central control point of apoptosis. Trends in cell
biology. 2000 Sep;10(9):369-77. PubMed PMID: 10932094.
100.   Goodman JM. The gregarious lipid droplet. The Journal of biological chemistry. 2008 Oct
17;283(42):28005-9. PubMed PMID: 18611863. Pubmed Central PMCID: 2568941.
101.   Tarnopolsky MA, Rennie CD, Robertshaw HA, Fedak-Tarnopolsky SN, Devries MC, Hamadeh
MJ. Influence of endurance exercise training and sex on intramyocellular lipid and mitochondrial
ultrastructure, substrate use, and mitochondrial enzyme activity. American journal of physiology
Regulatory, integrative and comparative physiology. 2007 Mar;292(3):R1271-8. PubMed PMID:
17095651.
102.   Vyas PM, Tomamichel WJ, Pride PM, Babbey CM, Wang Q, Mercier J, Martin EM, Payne RM.
A TAT-frataxin fusion protein increases lifespan and cardiac function in a conditional Friedreich's ataxia
mouse model. Human molecular genetics. 2012 Mar 15;21(6):1230-47. PubMed PMID: 22113996.
Pubmed Central PMCID: 3284115.
103.   Rufini A, Cavallo F, Condo I, Fortuni S, De Martino G, Incani O, Di Venere A, Benini M,
Massaro DS, Arcuri G, Serio D, Malisan F, Testi R. Highly specific ubiquitin-competing molecules
effectively promote frataxin accumulation and partially rescue the aconitase defect in Friedreich ataxia
cells. Neurobiology of disease. 2015 Mar;75:91-9. PubMed PMID: 25549872. Pubmed Central PMCID:
4358773.
104.   Coppola G, Choi SH, Santos MM, Miranda CJ, Tentler D, Wexler EM, Pandolfo M, Geschwind
DH. Gene expression profiling in frataxin deficient mice: microarray evidence for significant expression
changes without detectable neurodegeneration. Neurobiology of disease. 2006 May;22(2):302-11.
PubMed PMID: 16442805. Pubmed Central PMCID: 2886035.
105.   Shen Y, McMackin MZ, Shan Y, Raetz A, David S, Cortopassi G. Frataxin Deficiency Promotes
Excess Microglial DNA Damage and Inflammation that Is Rescued by PJ34. PloS one.
2016;11(3):e0151026. PubMed PMID: 26954031. Pubmed Central PMCID: 4783034.
106.   Lu C, Schoenfeld R, Shan Y, Tsai HJ, Hammock B, Cortopassi G. Frataxin deficiency induces
Schwann cell inflammation and death. Biochimica et biophysica acta. 2009 Nov;1792(11):1052-61.
PubMed PMID: 19679182. Pubmed Central PMCID: 3563672.

107.   Strawser CJ, Schadt KA, Lynch DR. Therapeutic approaches for the treatment of Friedreich's
ataxia. Expert review of neurotherapeutics. 2014 Aug;14(8):949-57. PubMed PMID: 25034024.
108.   Sandi C, Sandi M, Anjomani Virmouni S, Al-Mahdawi S, Pook MA. Epigenetic-based therapies
for Friedreich ataxia. Frontiers in genetics. 2014;5:165. PubMed PMID: 24917884. Pubmed Central
PMCID: 4042889.
109.   Kearney M, Orrell RW, Fahey M, Brassington R, Pandolfo M. Pharmacological treatments for
Friedreich ataxia. The Cochrane database of systematic reviews. 2016 Aug 30(8):CD007791. PubMed
PMID: 27572719.
110.   Oldham MC, Horvath S, Geschwind DH. Conservation and evolution of gene coexpression
networks in human and chimpanzee brains. Proceedings of the National Academy of Sciences of the
United States of America. 2006 Nov 21;103(47):17973-8. PubMed PMID: 17101986. Pubmed Central
PMCID: 1693857.
111.   Langfelder P, Horvath S. Eigengene networks for studying the relationships between co-
expression modules. BMC systems biology. 2007;1:54. PubMed PMID: 18031580. Pubmed Central
PMCID: 2267703.
112.   Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists
using DAVID bioinformatics resources. Nature protocols. 2009;4(1):44-57. PubMed PMID: 19131956.
113.   Becker KG, Hosack DA, Dennis G, Jr., Lempicki RA, Bright TJ, Cheadle C, Engel J. PubMatrix:
a tool for multiplex literature mining. BMC bioinformatics. 2003 Dec 10;4:61. PubMed PMID:
14667255. Pubmed Central PMCID: 317283.
288
289
290

Figure 1

Figure 1: Efficient temporal in vivo frataxin knockdown and rescue. (**a**) Schematic representation of the inducible vector system to mediated expression of shRNA expression against frataxin. The vector contains shRNA sequence against frataxin (*Fxn*) gene regulated by the H1 promoter with tet-operator sequences (tetO) and tet repressor (tetR) under the control of the CAGGS promoter. Transcription of the *Fxn* shRNA is blocked in cells expressing the tetR. Upon induction by doxycycline (dox), tetR is removed from the tetO sequences inserted into the promoter, allowing transcription of shRNA against *Fxn*. shRNA expression leads to RNAi-mediated knockdown of *Fxn* gene. (**b**) Predicted minimum free energy secondary structure of expressed shRNA targeting the *Fxn* is shown with the sequence (sense strand) and its complement sequence (antisense strand) in the duplex form along with hairpin loop. (**c**) Timeline for doxycycline treatment of mice with a double IP injection (5 and 10 mg/kg) of dox per week and in drinking water (2 mg/ml). IP injections and dox in water was withdrawn for rescue animals. Arrow signs indicated different time intervals considered for downstream analyses. (**d**) Transgenic mice without (Tg -) or with (Tg +) doxycycline treatment (see c) for 20 weeks were analyzed by Western blot for protein levels of FXN in various organs. (**e**) For quantification, FXN values were normalized to the level of b-tubulin in each lane. (**f**) Time series FXN knockdown and rescue in liver. Wild-type mice with dox (Wt +), transgenic mice with (Tg +) and without (Tg -) dox, and transgenic mice with (Tg ±) dox removal (rescue) samples were analyzed for week 0,3,8,12,16 and 20. Rescue animals (Tg ±) were given dox for 12 weeks and doxycycline was withdrawn for additional 4 and 8 weeks. (**g**) For quantification, FXN values were normalized to the level of b-tubulin in each lane. N=4 animals per group, *=p<0.01; two-way ANOVA test; n.s., not significant. Error bars represent mean ± SEM for panels e and g.

Figure 2

Figure 2: Neurological deficits due to frataxin knockdown. Body weight, survival, open field, gait analysis, grip strength and rotarod in five groups of animals; wild-type mice with dox (Wt +, n= 16(WK 0), n= 16(WK 12), n= 16(WK 24)) and without dox (Wt -, n= 16(WK 0), n= 16(WK 12), n= 16(WK 24)), transgenic mice with dox (Tg +, n= 30(WK 0), n= 21(WK 12), n= 15(WK 24)) and without dox (Tg -, n= 16(WK 0), n= 15(WK 12), n= 15(WK 24)), and transgenic mice with dox removal (Tg ± Rescue, n= 30(WK 0), n= 21(WK 12), n= 20(WK 24)). (**a**) Body weight from before 6 weeks and upto 34 weeks after dox treatment. (**b**) Survival was significantly declined in dox treated Tg + group of animals, no mortality was observed after dox withdrawal (Tg ± Rescue). (**c**) Open field test showing significant decline in total distance traveled by the dox treatment transgenic animals (Tg +) at 12 and 24 weeks when compared across all other groups. After dox withdrawal, there were no differences between the rescue group (Tg ± Rescue) and the three control groups at week 24. (**d**) Gait footprint analysis of all five groups of mice at 0, 12 and 24 weeks was evaluated for stride length. Dox treated transgenic (Tg +) animals revealed abnormalities in walking patterns displaying significantly reduced stride length, however the rescue group (Tg ± Rescue) displayed normal stride length when compared to other groups. (**e**) Representative walking footprint patterns. (**f**) Grip-strength test, dox treated transgenic (Tg +) mice had reduced forelimb grip strength at 12 and 24 weeks when compared across all other groups. After dox withdrawal, there were no significant differences between the rescue group (Tg ± Rescue) and the three control groups at week 24. (**g**) Rotarod test in mice upto 34 weeks after dox treatment. Dox treated transgenic (Tg +) animals stayed less time on the rotarod than the control groups, although after dox withdrawal, there was no significant difference between the rescue group (Tg ± Rescue) and the three control groups. Values between all five groups are shown as mean ± SME. Two-way ANOVA test *=P≤0.001; n.s., not significant.

a Wt + (Normal QT)

Tg + (Long QT)

b Tg + (12 WK – Long QT)

Tg ± Rescue (24 WK – Normal QT)

c Tg + (12 WK – P-wave)

Tg + (24 WK – No P-wave)

d Wt + (24 WK)

Tg + (24 WK)

e Observed QT interval (milliseconds)

f Ventricular septal wall thickness (mm)

g Posterior wall thickness (mm)

WK 12   WK 24

■ Wt +   ■ Tg -   ■ Tg +   ■ Tg ± Rescue

Figure 3

Figure 3: Frataxin knockdown mice exhibit signs of cardiomyopathy. (**a**) Representative traces of ECG recording in a wild-type and transgenic animal after dox treatment for 20 weeks, showing long QT intervals in Tg + animal. (**b**) ECG recording of a same dox treated transgenic (Tg +) animal at 12 and after dox withdrawal for additional 12 weeks (Tg $\pm$ Rescue), showing normal QT interval. (**c**) ECG recording of a same dox treated transgenic (Tg +) animal at 12 and 24 weeks, showing absences of P-wave only at 24 week. (**d**) Representative echocardiograms from the left ventricle of a wild-type and transgenic mouse at 24 week after dox treatment. (**e-f**) Quantification of observed QT interval (**e**), ventricular septal wall thickness (**f**) and posterior wall thickness (**g**) are shown for week 12 and 24. N = 6-8 animals per group, *=p<0.05; Student's t test. Error bars represent mean ± SEM.

Figure 4

Figure 4: Cardiopathology of frataxin knockdown mice. (**a**) Gomori's iron staining and quantification of iron deposition in dox treated transgenic (Tg +), wild-type (Wt +) and dox withdrawn transgenic (Tg ± Rescue) animals. Dox treated transgenic (Tg +) mice showing myocardial iron-overload (a) also displayed altered expression of ferritin protein (**b**) which is involved in iron storage. Both iron-overload and ferritin protein levels were significantly lower in Tg ± Rescue animals (a-b). (**c**) Masson's trichrome staining and quantification showing increased fibrosis in Tg + mice when compared to Wt + and Tg ± Rescue animals. (**d**) Electron micrographs of cardiac muscle from Wt +, Tg + and Tg ± Rescue animals at 20 week after dox treatment. Double arrow lines indicate sarcomere. m = mitochondria. Scale bars, 1 µm. Data are representative of three biological replicates per group. (**e**) Higher magnification of electron micrographs of cardiac muscle from Tg + mice, showing normal (m) and degenerating mitochondria (asterisks). (**f**) Aconitase activity was assayed in triplicate in tissues removed from 3 hearts in each group. Values represent mean ± SME. One-way ANOVA test *=P≤0.05.

Figure 5

Figure 5: Frataxin knockdown mice exhibit neuronal degeneration. (**a**) Electron microscopic analysis of Wt +, Tg + and Tg ± Rescue animal DRG neurons at 20 week after dox treatment. Arrows indicate condensed mitochondria with lipid-like empty vacuoles. Insert in Tg + panel shows higher magnification of electron micrographs of condensed mitochondria with lipid-like empty vacuoles in DRG neurons. Insert in Tg ± Rescue panel shows higher magnification of abnormal mitochondria (asterisks). Nu= nucleus. (**b**) Higher magnification of electron micrographs of Tg + and Tg ± Rescue animal DRG neurons at 20 week after dox treatment. Tg + panel shows degenerating mitochondria and condensed mitochondria with lipid-like empty vacuoles in DRG neurons. Tg ± Rescue panel shows two DRG neurons, one consisting of normal mitochondria (insert) and the other neuron with abnormal mitochondria (insert). (**c**) Higher magnification of Tg + animals showing condensed mitochondria with lipid-like empty vacuoles in DRG neurons. (**d**) Higher magnification of Tg ± rescue animals showing two DRG neurons, one consisting of normal mitochondria (normal DRG) and the other neuron with abnormal mitochondria (affected DRG). NM= normal mitochondria, CM= condensed mitochondria, EV= lipid-like empty vacuoles, AM= abnormal mitochondria. (**e**) Quantification of altered mitochondria in DRG neurons. Data are from multiple images from three biological replicates per group. Values represent mean ± SME. Two-way ANOVA test *=P≤0.05. (**f**) Electron micrographs of spinal cord axon cross-section, displaying reduced myelin sheath thickness and axonal cross-section area in Tg + and Tg ± Rescue animals. Bottom panel shows representative area utilized for quantification. (**g-h**) Quantification of myelin sheath thickness and axonal cross-section area in the spinal cord. Data are from 2000 or more axons per group in the lumbar spinal cord cross-section of high-resolution electron micrographs from three biological replicates per group. Values represent mean ± SME. One-way ANOVA test *=P≤0.05. (**i**) Electron micrographs of rod and cone photoreceptor cells, showing their disruption in Tg + animals (asterisks). (**j**) Retinal pigment epithelium cell layer showing the presences of large vacuoles (arrows) in Tg + animals along with disruption in their photoreceptor cells (asterisks).

Figure 6

Figure 6: Gene expression analysis of frataxin knockdown mice. (**a**) Heat map of significantly up- and down-regulated genes (rows) in heart tissue of Tg + mice from 0, 3, 12, 16, 20 and plus 4, 8 weeks post dox treatment relative to controls are grouped into 13 functional categories. (**b**) Summary of differentially expressed genes during *Fxn* knockdown and rescue in heart, cerebellum and DRG tissues from four biological replicates. (**c**) Cumulative percent of variability in Tg + gene expression data explained by the first three principal component for each functional category. (**d**) Networks highlighting differentially expressed genes due to *Fxn* knockdown in Tg + mice for selected functional categories. Nodes represents genes and edges are present between nodes when their gene expression correlation is greater than 0.5. Node size and color (red= up-regulation and green= down-regulation) denotes extent of differential expression. (**e**) Western blot analyses of Caspase 8, FXN and LC3 following 20 weeks with or without dox treatment in heart. (**f**) TUNEL assay of heart, spinal cord and DRG neurons at 20 weeks after dox treatment in Tg + mice. (**g**) Literature associated gene networks highlighting differentially expressed genes due to *Fxn* knockdown in Tg + mice for selected key-terms. Nodes represents genes which has pairwise correlation greater than 0.5 with *Fxn* gene and edge size represents strength of their gene expression correlation. Node size correspond to number of PubMed hits with co-occurrence of gene and its corresponding key-term. Node color represents up-regulated (red) and down-regulated (green). (**h**) Representative images and quantification of LC3 staining in heart tissue at 20 weeks after dox treatment. Values represent mean from three biological replicates per group ± SME. One-way ANOVA test *=P≤0.05. (**i**) Heat map depicting expression of key genes (rows) across samples (columns) for seven groups (red corresponds to gene up-regulation and blue to down-regulation). Seven groups represents, (i) ataxia, (ii) cardiac fibrosis, (iii) excess iron overload, (iv) muscular strength, (v) myelin sheath, (vi) neuronal degeneration and (vii) autophagy related genes. Column represents, seven independent datasets obtained from, FRDAsh mice, cardiac specific knockout mice(23), knock-in knockout mice(20), knock-in mice(20) and patient peripheral blood mononuclear cells(44).