The authors wish to be identified to the reviewers.

Understanding the dynamics of laboratory populations of *Drosophila melanogaster*:

long-term experiments meet individual-based modelling

Authors: Sudipta Tung[a], M. Rajamani[b,c], Amitabh Joshi[b] and Sutirth Dey[a]*

Affiliations:

[a]Population Biology Laboratory, Biology Division, Indian Institute of Science Education and
Research-Pune, Dr. Homi Bhabha Road, Pune, India, 411 008

[b]Evolutionary Biology Laboratory, Evolutionary and Organismal Biology Unit, Jawaharlal
Nehru Centre for Advanced Scientific Research, Jakkur P.O., Bengaluru, India, 560 064

[c]Present Address: Clintus Network Ltd., No. 205/1 & 2, PCR Warehouse Complex, Soukya
Road, Koralur Village, Hoskote, Bengaluru 562 114

15
*Name and address of the corresponding author:
Sutirth Dey
Associate Professor, Biology Division
Indian Institute of Science Education and Research (IISER) Pune
Dr. Homi Bhabha Road, Pune, India 411 008
Tel: +91-20-25908054

Email: s.dey@iiserpune.ac.in

Running head: Modeling *Drosophila* population dynamics
25

Key words: fluctuation index; stability; constancy; persistence; minimum critical size;

time-series; stage-structured model.

ABSTRACT

How do the amount of resources available to different life-stages affect the dynamics of

stage-structured populations? To investigate this issue, we present analyses of census data

from a 49-generation experiment on replicate laboratory populations of the common fruit

fly, *Drosophila melanogaster*, subjected to four different combinations of larval and adult

nutritional levels. We also elucidate the mechanistic underpinning of the dynamics through a

stage-structured individual-based model that incorporates life-history parameters common to

many holometabolous insect populations. The model captures the dynamics of all the

nutritional regimes, resolves an observed discrepancy between data from an earlier empirical

study and our results, and is able to successfully reproduce the dynamics from an independent

Drosophila experiment. We also compare our study with experiments and models on the

dynamics of various other species to understand which aspects of dynamics are generalizable.

We then use this model to explore how the interaction between nutrition levels and various

facets of unequal number of males and females affect population dynamics. We show that the

effects of unequal sex-ratio and sex-specific culling are greatly influenced by fecundity but

not by levels of juvenile nutrition. We also demonstrate that the efficiency of a widely-used

pest control method (Sterile Insect Technique) depends on a complex interaction between the

levels of juvenile nutrition and the density-independent adult fecundity.

1. INTRODUCTION

The laboratory ecology of *Drosophila melanogaster* has been investigated for more than half a century, leading to considerable understanding of various density-dependent effects on the population dynamics of laboratory cultures of this species (reviewed by Mueller 1985; Mueller and Joshi 2000). Briefly, three density-dependent feedback loops ― effects of larval crowding on larval survivorship and subsequent adult fecundity, and effects of adult crowding on adult fecundity ― are thought to be the primary drivers of the dynamics of discrete generation cultures of *D. melanogaster* (reviewed in Mueller and Joshi 2000). Several recursions incorporating one or more of these density-dependent feedback mechanisms have also been proposed to model the dynamics of laboratory cultures of *D. melanogaster* (Bakker 1961; De Jong 1976; Nunney 1983; Prout and McChesney 1985; Rodriguez 1989). Mueller (Mueller 1988) explicitly incorporated all three density-dependent feedback mechanisms into a single recursion as: $n_{t+1} = ½ . G(N_t). F(Vn_t). W(Vn_t). V.n_t,$ where n_t and N_t represent the number of eggs and adults in generation t, respectively, V is the density-independent probability of larval viability, $W(Vn_t)$ and $F(Vn_t)$ are the functions representing the effects of larval density on larval survivorship and adult fecundity, respectively, and $G(N_t)$ is the function reflecting the effect of adult density on adult fecundity. This model remains the most detailed abstraction of *D. melanogaster* dynamics in the literature, and gave rise to several interesting predictions that were subsequently verified empirically (Mueller and Huynh 1994; Mueller and Joshi 2000; Sheeba and Joshi 1998).

Despite this rich body of work, several aspects of *Drosophila* population dynamics in the laboratory still remain poorly understood. For example, although we know about the role of larval and adult nutrition in affecting population stability, it is still not clear if and how

various life-history-related parameters, like egg hatchability and critical minimum size for

pupation, interact with these nutritional regimes. Moreover, there is little theoretical or

empirical understanding of how these nutritional regimes affect the various aspects of the

population size distribution (mean, skewness and the position of the various quantiles) in

Drosophila. There are two primary reasons for these lacunae. First, the absence of empirical

datasets of sufficient length (although see Mueller et al. 2000) over multiple nutritional

regimes precludes meaningful investigation of population size distributions. Second, most

models of *Drosophila* dynamics are deterministic, which rules out a theoretical exploration of

the population size distributions, except in the chaotic regime. This has limited the study of

the dynamics of *Drosophila* populations to stability properties (coefficient of variation,

autocorrelations) and average population sizes, thus missing out on several interesting aspects

of the dynamics with potentially large explanatory power.

In order to fill in some of these gaps in our understanding of *Drosophila* population

dynamics, we conducted a 49-generation long experiment to describe the main features of the

dynamics of laboratory populations of *D. melanogaster* subjected to four different nutritional

regimes. We also developed and simulated a stochastic model of *Drosophila* population

dynamics to generate time series data similar to our experiments. We then compared the

experimental data with our simulation results to show that our model is able to capture

various qualitative and quantitative aspects of *Drosophila* population dynamics in the

laboratory. We then demonstrated the usefulness of our model for understanding the

dynamics of laboratory populations of *D. melanogaster* in three ways. First, we used it to

resolve a discrepancy between observations from an earlier study and our results. Second, we

used it to generate clear predictions about how various life-history-related parameters affect

the dynamics of the populations under the various nutritional regimes. Third, we used data

from a previous experimental study to validate some of these predictions.

After establishing the bonafides of our model as a descriptor of *Drosophila* dynamics, we

briefly compared its various features and predictions with those arising from models and

experiments on other species. We then used our model to investigate various facets of a major

question, namely the effects of unequal numbers of males and females on population

dynamics. Simple models of population dynamics often assume that the entire dynamics can

be understood in terms of the number of females in the population (May 1976). Over the last

few decades, the consequences of relaxing this assumption has been extensively investigated

both theoretically and experimentally (reviewed in Rankin and Kokko 2007). This line of

study helps us understand the dynamics of those populations in which the two sexes face very

different rates of mortality due to natural or anthropogenic reasons (Berec et al. 2016;

Halvorsen et al. 2017; Perlman et al. 2015). Moreover, one of the highly successful pest

eradication strategies, the Sterile Insect Technique (SIT), explicitly depends on altering the

sex ratio of the population (Dyck et al. 2005). In nature, the amount of food available would

obviously vary across space and time and this, in principle, can modulate the effects of sex

ratio on the dynamics. Yet, the interaction of food and sex ratio in shaping the dynamics in

general and stability in particular, has remained relatively poorly understood. Our simulations

suggested that this interaction is best understood in conjunction with the density-independent

fecundity of the organisms. In a nutshell, the effects of changing the sex ratio on population

stability and average size are dependent on certain aspects of the provided nutrition and not

others. However, when it comes to SIT, the efficacy of the program depends more critically

on the food provided over a large and biologically relevant parameter range.

2. MATERIALS AND METHODS

2.1 Experiment

The details of this experiment have been reported in the PhD thesis of one of the authors (Dey 2007). Here we present a brief account and refer the readers to the SOM for more details of *Drosophila* laboratory ecology (Text S1) and the experiments (Text S2).

The experiment consisted of thirty-two populations of *D. melanogaster*, each represented by a single vial culture. These 32 populations were randomly allotted to one of four nutritional regimes, such that there were eight populations per regime. Following established norms (Mueller and Huynh 1994; Mueller et al. 2000) these regimes were called HH, HL, LH, LL — where the first letter indicates the quantity of larval food and the second letter represents the status of adult nutrition. In case of larval food, H and L denoted ~6 mL and ~2 mL of banana-jaggery medium per vial, respectively, whereas in the case of adult nutrition, H and L referred, respectively, to the presence and absence of live yeast paste supplement to banana-jaggery medium. Thus, for example, HL denotes a nutritional regime comprising of ~6 mL medium per vial for the larvae, but no live yeast paste supplement for the adults, and so on. The experiment was started with 8 adults (4 males and 4 females) in each population in the first generation, and the number of adults in each successive generation were noted. The experiment was continued for 49 discrete generations and yielded 32 time-series (see Figure S3 for representative time series).

2.2 Statistical analyses

Distributional properties of the experimental time series of adult populations were assessed using the mean, median, 5^{th}, 10^{th}, 25^{th}, 75^{th}, 90^{th} and 95^{th} percentiles in box plots (Zar 1999).

140   The constancy stability (Grimm and Wissel 1997) of the populations was measured as the

141   fluctuation index (*FI*, Dey and Joshi 2006) which is given as:

142
$$FI = \frac{1}{T \times \overline{N}} \times \sum_{t=0}^{t=T\text{-}1} \left| N_{t+1} - N_t \right|$$

where $\overline{N}$ is the mean population size over *T* generations and N_{t+1} and N_t are the population

sizes at the $(t+1)^{\text{th}}$ and t^{th} generations, respectively. This formulation suggests that *FI*

increases when the variance of the time series is increased, or the mean or the strength of

autocorrelation of the time series is decreased.

In order to investigate the effects and interaction of larval and adult nutritional levels on

constancy stability, the *FI* data were subjected to a two-way analysis of variance (ANOVA)

with larval nutrition (fixed factor, two levels: Low and High) crossed with adult nutrition

(fixed factor, two levels: Low and High). All statistical analyses were performed using

STATISTICA® v5 (StatSoft. Inc., Tulsa, Oklahoma).

2.3 The model and simulations

2.3.1 *Model formulation*

Our individual-based, stochastic model (see Text S3 for the details of model formulation) is

based on a previous deterministic model (Mueller 1988) with a few modifications (see

section S5). The biological justifications for the various functional forms used and the model-

calibration process has been described in detail in the SOM (see Text S4). Here we present a

brief account to highlight the salient features of the model. A schematic diagram of the model

processes has been included as Figure S2 for visualizing the sequence in which model

processes were executed. Terms in italics denote parameters in the model, whereas state

variables are in the `Courier New` font. For a given generation, the model starts with the

number of eggs (`numegg`), a fixed fraction (*hatchability*) of which hatch to form

`numlarva`, i.e. number of larvae:

$$\texttt{numlarva} = hatchability \times \texttt{numegg} \qquad \qquad \text{...eq 1}$$

Each larva is then assigned a body-size (in arbitrary units of length) through a random draw

from a normal distribution whose mean (mean_size) is positively related to the amount of

food (*food*) in the culture but negatively related to `numlarva`:

$$\text{mean\_size} = x_1 \times (1 - 1/(x_2 + \exp(-x_3 \times \texttt{numlarva} + food))) \qquad \text{...eq 2}$$

where x_1, x_2 and x_3 are parameters (see Table 1 for descriptions) and exp denotes the

exponential function. The standard deviation of the normal distribution is assumed to be a

constant (=0.45).

Table 1: List of parameters used in the model

Parameter	Description	Value
food	Amount of larval food present	1.76 (LL and LH) and 2.56 (HL and HH)
adnut	Quality of adult nutrition/fecundity booster	1 (LL and HL), 1.29 (HH) and 1.49 (LH)
hatchability	Egg-to-larval viability	0.98
m_c	Critical mass *i.e.* the minimum mass/size required to become a viable adult	1.1 (JB) and 1 (FEJ)
sen_adden	The coefficient of sensitivity of female-fecundity to adult density	0.17
sen_adsize	The coefficient of sensitivity of female-fecundity to adult size	1.7
sigma_size	Standard deviation of larval body size distribution	0.45
x_1	Parameter	2.5
x_2	Parameter	1

x_3	Parameter	0.009
x_4	Parameter	1
x_5	Parameter	85
x_6	Parameter	2

Larvae which are larger than a critical minimum size (m_c) form the adults of the generation,

while the remaining larvae are assumed dead. The adults are randomly assigned to be males

or females and it is assumed that all females are fertilized irrespective of the number of males

in the population (this assumption is relaxed in the simulations discussed in section 3.9). In

Drosophila, as in many insects, fecundity of females is known to vary positively with body

size (Honěk 1993). We modeled this positive relationship using a parameter for sensitivity of

fecundity to adult body size (*sen_adsize*). Another factor that affects the density-independent

fecundity of adult flies is the presence of protein supplement in the form of live yeast paste

which boosts the fecundity of the female flies. This effect was modeled through an adult

nutrition parameter called *adnut* such that a value of *adnut* =1 would represent the fecundity

of the flies in the absence of yeast paste, while *adnut* >1 represented the fecundity-boosting

effects of the yeast paste (see Text S4 for model calibration). Taken together, the adult

density-independent fecundity of the i^{th} female was given as:

$$\text{addens\_ind\_fec}_i = adnut \times x_5 \times \ln(x_6 + sen\_adsize \times \text{size\_adult}_i) \qquad \ldots \text{eq 3}$$

where size_adult$_i$ is the size of the i^{th} adult and x_5 and x_6 are parameters.

In *Drosophila*, adult fecundity is negatively affected by adult density (Mueller and Huynh

1994; Rodriguez 1989). We modeled this effect using a hyperbolic function

$$\text{addens\_eff} = (1/(1 + sen\_adden \times \texttt{numadult})) \qquad \ldots \text{eq 4}$$

where the parameter *sen_adden* determines the sensitivity of female-fecundity to adult

density (`numadult`). The number of eggs laid by each female is computed as the product of

eq 3 and eq 4, and summed up over all females to form the `numegg` for the next generation.

2.3.2 *Simulations:*

To investigate the population size distributions for each nutritional regime (LH, HH, HH or

HL), we simulated eight replicate runs of the model with 49 generations in each replicate, to

keep parity with the experimental data. However, none of our conclusions changed when the

length of the time series was increased (see Text S9 for discussion). Every simulation run

started with 18 eggs. When there was extinction in any generation (i.e. `numadult` = 0), the

time series was reset with four females with body size $= 2 \times m_c$. Following previous studies

(Sah et al. 2013; Tung et al. 2014), we also incorporated additional demographic stochasticity

in the model by considering a 50% chance of extinction, whenever population size went

below eight. If extinction occurred due to demographic stochasticity, the population was reset

in the same way as mentioned above.

We also explored the effects of wide ranges of life-history related parameters (*hatchability*,

m_c, *sen_adden*, *sen_adsize*) on population stability. For each value of a given parameter, we

took an arithmetic mean of *FI* measured over 100 replicate time-series, each of which was

100-generations long. All other conditions of the simulations were identical to those in the

previous paragraph.

Our empirical data revealed that the HL regime had a greater average population size and

lower *FI* compared to the HH populations (see section 3.5 for details) whereas an earlier

study (Mueller and Huynh 1994) had shown that the population size of HH was greater than

that of HL and the two regimes had similar constancy stability. In order to investigate this

discrepancy between the two results, we simulated our model with five different values of

larval food ranging from 3.0 to 7.0 in step size of 1.0. Each value of larval food level (*food*)

was crossed with two values of *adnut* – 1.0 and 1.29- which represented the presence and

absence respectively of yeast for the adults. For each *food* × *adnut* combination, we simulated

eight 49-generation long time-series, and obtained the corresponding population size

distributions, and *FI* and egg-to-adult viability values. All other parameter values were

identical to the earlier simulations (Table 1).

2.4 36-generation simulation and experiment

To validate one of the predictions arising from our model, we compared our model output

with the dynamics of four *Drosophila* populations selected for faster development and early

reproduction (henceforth called FEJ_{1-4}) for ~125 generations (Dey et al. 2008; Prasad et al.

2003). The FEJ_{1-4} lines were derived from four ancestral populations (JB_{1-4}), which served as

controls in that experiment. Incidentally, one of these JB populations is the ancestor for the

32 populations used in the present study. For each FEJ_i or JB_i ($i \in 1$-4) population

(represented by single vial cultures), there were four replicates each under HL and LH

regimes. Thus, there were 16 FEJ populations and 16 JB populations, that experienced the

LH regime and similarly 16+16 that experienced the HL regime. The maintenance details of

this 36-generation long experiment are given elsewhere (Dey et al. 2008) and are similar to

the experimental protocol of the present study.

The rationale and the details of the model recalibration are presented in the SOM (Text S7). It

should be noted here that the empirical *FI* values are those reported in Figure 2a of the earlier

study (Dey et al. 2008) and are being re-plotted here only to facilitate comparison with the

simulation results. The population size distribution data from these experiments is being

reported for the first time here.

2.5 Simulating the dynamics of sex-biased populations

This part of the study consisted of three inter-related questions on the consequences of

systematic departures from equal number of males and females in the population. We

mention the simulations briefly here, with the details being relegated to Text S8.

a) Distortion of sex ratio and sex-specific culling: Here we simulated the behaviour of *FI* and

average population size for two different values of fecundity values (high: $x_5 = 85$ and low: x_5

$= 17$) crossed with two values of *food* (high = 1.76 and low = 2.56). For sex ratio distortion,

we varied the probability of an individual being female from 0.1 to 0.9, while for sex-specific

culling, a fixed percentage (0% - 95%) of males or females were removed after the

assignment of sex.

b) Sterile insect technique (SIT): In this method, a fixed number or proportion of sterile males

are introduced into the population every generation. Following an established modelling

strategy (Knipling 1959), we assumed the proportion of females producing eggs for the next

generation to be a linear function of the fertile/sterile male ratio. Thus, proportion of egg

laying females = min(($num\_fertile$/($\beta \times num\_sterile$)), 1), where $num\_fertile$ and $num\_sterile$

are the number of fertile and sterile males in the population, β is the competitive ability of the

sterile males and min() is the minimum function. We studied the interaction of density-

independent female fecundity with the amount of juvenile food available to determine the

efficacy of SIT. The latter was estimated as the minimum number of sterile males required to

ensure extinction of the population within an arbitrary number of generations (here 10) with a

reasonably high success rate (here 90%). For this, we explored a large range of parameter

values for density-independent fecundity (i.e. $adnut \times x_5$, range = 1 - 150, step size = 5) and

larval food (*food*, range = 0.5 - 3.0, step size = 0.2). To avoid confounding extinctions due to

intrinsic reasons with extinctions due to SIT, the efficacy of SIT was estimated only for those

parameter combinations where the probability of an extinction event within first 10

generations in the absence of SIT is below 0.1 (Figure S10, see section 3.9 for discussion).

3. RESULTS AND DISCUSSION

3.1 Experiments: Larval and adult nutritional regimes interact to shape the adult

dynamics

There was a significant interaction between the larval and adult nutrition ($F_{1, 28}$= 17.92, *p*=

0.0002) suggesting that enhancing the fecundity of flies (through a supply of yeast) causes a

much greater increase in *FI* when the amount of larval food is limiting (i.e. LL and LH) than

when it not limiting (i.e. HL and HH) (Figure S1). This is because although both LL and LH

experience substantial larval crowding, the greater fecundity of the LH flies leads to higher

larval crowding even with moderate adult population sizes which, in turn, causes regular

population crashes. On the other hand, even when there are population crashes, the greater

fecundity of the LH flies ensures a high population size in the next generation. Together,

these two effects ensure large amplitude oscillations in LH population sizes, and a

substantially larger *FI* than the LLs (Tukey's HSD *p* = 0.00016). On the other hand, although

the fecundity of females in the HH populations is larger than those in the HLs, the non-

limiting amount of larval food ensures that the population crashes are only marginally more

severe in the former. This leads to a much lower (although statistically significant; Tukey's

HSD *p* = 0.00017) increase in *FI* in the HH populations, compared to the HL populations

(Figure S1).

3.2 Experiments: The differences in the dynamics of the populations are reflected in their population size distributions and *FI*

Figure 1. Population size distribution and constancy stability of experimental and simulated time-series. (A) Descriptive statistics of the population size distributions. Red dashed lines = means, thin black lines = medians, edges of the boxes=25th and 75th percentiles, whiskers=10th and 90th percentiles and the circles outside = 5th and 95th percentiles of the distributions. White boxes represent experimental data while grey shaded boxes are for simulated time-series. **(B)** Average (± SEM) *FI* of the experimental and simulated time-series in the four regimes. Both plots suggested a good agreement between the experiments and the simulations. The populations in the HL regime were the most stable with highest average population size while those in the LH regime were the least stable with lowest population size.

We began our investigation by examining the distribution of population size which is ultimately related to the temporal dynamics of populations (Figure S3). Both larval and adult nutritional levels were found to affect the population size distributions (Figure 1A, the white boxes). Specifically, when larval food is low, population size distributions have lower values of mean, median, 25th percentiles and 75th percentiles, compared to when larval food is high (*cf* LH with HH and LL with HL in Figure 1A). Interestingly, irrespective of the level of larval nutrition, providing yeast to the adults reduced the population sizes (*cf* LH with LL and HH with HL in Figure 1A). Moreover, in the LH and HH regimes (Figure 1A), the population size distributions are much more skewed to the left (i.e. median < mean), which is indicative of crashes in population numbers from various medium to high population sizes (see also

figure 4 in Dey and Joshi 2013). All these observations are due to the fact that low levels of

larval food or increased adult fecundity increase the larval crowding by reducing the per-

capita food available to the larvae. Consequently, fewer larvae are able to acquire a body size

greater than m_c, which reduces the egg-to-adult survivorship, and hence the adult population

sizes. Interestingly, the mean and the median population sizes were very close for the HL

populations, but not so for the other three (Figure 1A). This showed that the population size

distributions of HL had little or no skew, while the other three regimes exhibited positive

skewness. This implied that in spite of having a larger average population size compared to

the other three regimes, the HL populations exhibited lower amplitude fluctuations relative to

their own mean population size. Thus, not surprisingly, the HL populations were found to

have the lowest *FI* (Figure 1B) amongst the four regimes.

Post-hoc tests (Tukey's HSD) on *FI*s in the four nutritional regimes revealed all pair-wise

differences to be significant with the rank order: LH > HH > LL > HL (Figure 1B, the white

bars). Although these four regimes have never been studied together till date, subsets of them

have been studied in all kinds of combinations. Thus, it has been shown that in terms of

constancy stability LH< HL~HH (Mueller and Huynh 1994), LH< HL and LH< LL (Dey

and Joshi 2013). Our results (Figure 1B) are in excellent agreement with all these studies

except those of Mueller and Huynh (1994) who showed theoretically and empirically, that the

constancy stability of HL and HH were not different. Moreover, Mueller and Huynh (Mueller

and Huynh 1994) also predicted the average population size of the HH regime to be much

larger than that of the HL regime, which also did not match our observations (Figure 1A). We

resolve this issue later in this study (section 3.5) using our individual-based model of

Drosophila dynamics.

3.3 **Simulations: High level of correspondence between the experimental data and the**

model output

The simulation results (grey bars) matched the various salient features of population size

distribution (Figure 1A) and population stability (Figure 1B) in the empirical observations in

all four nutritional regimes. To the best of our knowledge, there are no models of *Drosophila*

dynamics whose predictions have been verified in this detail with experimental data. This is

more a reflection on the paucity of good quality long time series data, rather than any

shortcoming on the part of the modelers. In fact, in the context of dynamics of laboratory

populations of *D. melanogaster,* our 49 generation data-set is perhaps the second-longest in

the literature in terms of number of generations.

Although the model did an excellent job in capturing the various aspects of the experimental

data (*cf* Figure S3 and S4; also Figure S5), these details (and therefore the parameter values

that lead to them) are obviously experiment-specific and shall vary across studies. Therefore,

the usefulness of our model is more in terms of the mechanistic understanding that it

generates about how the dynamics is affected by the interaction of various life-history and

environmental variables. That was our next object of investigation.

3.4 **Simulations: The effects of various life-history related traits on dynamics**

3.4.1 *Hatchability (hatchability) and critical size (m_c)*:

Our model predicted that with decreasing hatchability of the eggs up to moderate values,

population *FI* decreases (i.e. constancy stability increases) in all four regimes (Figure 2A).

This is because a reduced hatchability in generation *t* is conceptually equivalent to reduced

fecundity in generation *t*-1, which is known to be a stabilizing factor (Mueller 1988). As

expected, the destabilizing effects of increasing hatchability are more pronounced where the

larval crowding is already very high (LH) and are mildest where larval crowding is the lowest

(i.e. HL). Interestingly when hatchability values are very low (~.35 or less), further

reductions lead to a decline in the average population size without greatly affecting the

variation in population sizes over time (Figure S7). Since the expression for *FI* includes the

average population size in the denominator, this leads to a minor increase in the *FI* values for

very low values of hatchability.

Figure 2. Effect of varying life-history related parameters of the model on constancy stability. Each point represents average (± SEM) fluctuation index of 100 replicates of 100-gen long simulated time series. Error bars are too small to be visible. (**A**) In all four regimes, as *hatchability* reduces, larval density also reduces and thus populations become more stable. (**B**) As critical size increases, the populations become more destabilized. (**C**) Increasing the sensitivity of female-fecundity to adult density (*sen_adden*) increases constancy stability in all regimes except LH. (**D**) Increasing the sensitivity of female-fecundity to adult body size (*sen_adsize*), reduces constancy stability in all regimes except LH. See section 3.4.2 for explanations for the anomalous behaviors in the LH regime.

Like hatchability, increasing larval critical size (m_c) also has a negative effect on constancy

stability (Figure 2B). This works in two ways. First, all else being equal, increasing m_c means

that fewer larvae would be able to attain m_c, which would reduce larval survivorship. This is

analogous to reducing survivorship through reduced larval food amount, which is a

destabilizing factor. Secondly, increasing m_c means on an average, the surviving adults would

have a greater body size, which would translate into larger fecundity and thus, destabilize the

dynamics. Conversely, decreasing m_c is always expected to stabilize the dynamics (Mueller

1988): a prediction that we return to in section 3.6.

3.4.2 *Sensitivity of female fecundity to adult density (sen_adden) and adult body size*

(sen_adsize):

Increasing adult density is known to negatively affect female fecundity in *Drosophila*

melanogaster (Mueller and Huynh 1994). In our model, *sen_adden* determines the strength of

this effect, such that for same adult density, greater *sen_adden* results in lesser fecundity.

This in turn enhances larval survivorship, by increasing the amount of food available per

capita, which has a stabilizing effect on the dynamics. On the other hand, *sen_adsize*

determines the strength of the positive correlation between body size and fecundity, such that

increasing *sen_adsize* will increase fecundity, thereby reducing larval survivorship,

ultimately leading to destabilized dynamics. In a nutshell, an increase in *sen_adden* and

decrease in *sen_adsize* is expected to lead to a stabilization of the population dynamics. Our

simulation results agreed with this prediction in all the regimes except LH (Figure 2C and

2D). In the LH regime, both *sen_adden* and *sen_adsize* seemed to have little effect on *FI*,

even though, reducing *sed-adden* and increasing *sen_adsize* caused the total egg number to

go up (Figure S8A and S8C). The reason for this unintuitive behaviour was revealed when we

investigated the effect of these two parameters on the egg-to-adult survivorship. Increasing *sen_adden* (Figure S8B), or decreasing *sen_adsize* (Figure S8D), hardly affected the egg-to-adult viability in the LH regime. This is because the reduced levels of larval food ensured that even with low fecundity, there was substantial larval crowding in this regime (as evidenced by the low levels of egg-to-adult viability) so that there was almost no effect of changing *sen_adden* or *sen_adsize* on larval mortality. As a result, the destabilizing effect of increasing fecundity was not seen in the LH populations. Another interesting observation from Fig 2C is that at low values (< 0.14) of *sen_adden*, HH has a much larger *FI* than LH, despite having similar levels of average egg number (Fig S8A). This is because for low values of *sen_adden*, the egg-to-adult viability of LH is lower than that of HH (Fig S8B), due to the fact that the amount of larval food for the former regime is much less than that of the latter. Consequently, greater number of adults can be supported by HH, which makes higher amplitude fluctuations in population sizes more likely, thereby increasing the values of *FI*.

The primary insight from the above observations is that even in the highly simplified dynamics under laboratory conditions, the environment can interact with the life-history related traits of the organisms to lead to very counter-intuitive effects on the dynamics.

3.5 Simulations and Experiments: Population dynamics is shaped jointly by the quality and quantity of nutrition

As stated already, one of our empirical results did not match the observations of an earlier study (Mueller and Huynh 1994). We found that HL populations had greater constancy stability and average size than the HH populations whereas Mueller and Huynh (Mueller and Huynh 1994) reported that the HH populations had similar constancy stability but much greater average size than the HL populations. The primary difference between the two

experiments was in terms of the amount of food given to the larvae. In the experiment of

Mueller and Huynh (Mueller and Huynh 1994), the HL and HH larva got 40 mL of food in a

250 mL bottle while in our experiment the corresponding larva got ~6 mL food in a 37 mL

vial. Consequently, the adult population sizes in the HL and HH regime varied in the range of

~40-240 in our experiment, but ~ 400-1600 in the previous experiment.

Figure 3. Simulations on effects of varying larval nutrition on population dynamics.
(**A**) Population size distributions for the simulated time-series under low *adnut* (cyan) and
high *adnut* (orange) conditions for different levels of larval food amount (*food*). White
dotted lines = means, thin black lines = medians and the circles outside = 5[th] and 95[th]
percentiles of the distributions. The relative positions of the population size distribution of
low *adnut* and high *adnut* regimes reverses as the larval food amount increases. (**B**)
Average (± SEM) fluctuation index of the low *adnut* and high *adnut* regimes become
comparable, when the level of larval food is high. (**C**) Although the low *adnut* regimes
have greater average (± SEM) egg-to-adult viability than the high *adnut* regime for low
values of *food*, the viabilities become comparable as *food* increases. Error bars are too
small to be visible here.

To investigate whether the differences in the larval food amount could explain the observed

discrepancies, we simulated the HH and HL regime for different levels of larval food,

keeping all other parameters the same as in the earlier simulation. We found that as the level

of larval food increased, the relationship between the population size distribution of HH and

HL reversed (Figure 3A). Furthermore, with increasing value of larval food, the *FI* in the HH

regime reduced and approached the value seen in the HL regime (Figure 3B). The underlying

mechanisms behind these observations can be understood as follows.

Due to the availability of yeast paste to the adults in the HH regime, the per-capita fecundity

of the females is very high. Consequently, when the amount of larval food is relatively less

(as in our experiment) there is larval crowding which reduces the survivorship in the HH

regime. Therefore, with increasing levels of food, the survivorship increases (as in Figure

3C), which is manifested as increased population size in the HH regime (Figure 3A, orange

boxes). The HL populations also face some amount of larval crowding at lower levels of

food. However, since they do not have increased fecundity at high adult population sizes (due

to the absence of yeast), the increase in population size plateaus off at a much lower level of

food (Figure 3A, cyan boxes).

In order to visualize the effects of increased food amount on constancy stability, we need to

appreciate that reduced larval crowding has two opposing effects on the dynamics. First, it

stabilizes the dynamics by reducing larval mortality. At the same time, it can destabilize the

dynamics by increasing the body size of the females at eclosion. As the larval food level

increases, both these factors come into play. However, as there are upper bounds to both

survivorship (=1) and the body size of the flies (= the physiological limit of body size),

beyond a certain amount of larval food, both these factors cease to play a major role, and the

FI in both regimes become similar. This can be clearly seen in Figure 3B and explains why in

the presence of large amount of larval food, HL and HH populations have similar constancy,

as reported previously (Mueller and Huynh 1994). When the amount of larval food is

relatively small (while still being high, compared to LH or LL regimes), the destabilizing

effect of reduced survivorship overpowers the stabilizing effect of diminished fecundity due

to reduced body size. This is because there is a minimum value for the body size (= m_c),

which automatically places a lower bound on the fecundity of the flies irrespective of the

level of larval crowding. Since the HH populations experience greater larval crowding than

the HL populations, they are expected to exhibit lower constancy, as seen in our experiments

(Figure 1B, white bars) and simulations (Figure 1B, grey bars).

In the *Drosophila* population dynamics literature, labels like LH and HL have typically been

used as qualitative descriptors to signify the levels of larval crowding (highly crowded *versus*

un-crowded) and state of adult nutrition (yeasted *versus* un-yeasted). As described in the

Introduction section, this categorization has broad explanatory power in terms of the nature of

the dynamics: LH leads to high amplitude oscillations while HL leads to relatively stable

dynamics. However, the above comparison between the HL and HH regimes from the two

different studies shows that changing just one environmental parameter (here, the absolute

quantity of otherwise 'high' larval food) can lead to a rich array of dynamics. This again

highlights how the actual values of the environmental parameters interact with life-history

related traits in determining population dynamics.

3.6 Simulations and Experiments: Reduction in m_c is one way for population stability to

evolve

One of the predictions of our model is that decreasing m_c should lead to stabilization of the

dynamics (Figure 2B and Section 3.3.1). This prediction is consistent with earlier theoretical

studies (Mueller 1988) and has been empirically validated using laboratory populations of *D.*

melanogaster (Dey et al. 2008; Prasad et al. 2003). These earlier experiments used a

population of flies (FEJs) that had evolved reduced m_c as a correlated response to selection

for faster development and early reproduction. Consequently, they were found to have

reduced *FI* compared to the corresponding ancestral control populations (JBs). In order to see

whether our model was capable of recovering the other features of the dynamics from the

earlier experiment (Dey et al. 2008), we set a slightly lower value of m_c for the FEJs and kept

all other parameters the same as in the previous simulations (Table 1). Our model was again

able to capture the trends in the distributional properties (Figure 4A) and the *FI* values

(Figure 4B and 4C) of JB and FEJ populations in both regimes. The empirical data in Figure

1 and Figure 4 are from two completely independent experiments done at different times. The

fact that our model is able to predict the major features of the latter data-set based on

parameterizations done for a subset of the former shows that our parameterization was robust.

However, we again emphasize here that the main focus of this part of the study was to gather

insights about how the various life-history and environmental parameters interact, and the

excellent quantitative match between the data and the model is essentially a secondary,

though greatly encouraging, result.

Figure 4. Validating model predictions on JB and FEJ populations. (**A**) Descriptive statistics of the population size distributions of experimental and simulated JB and FEJ populations. Red dashed lines = means, thin black lines = medians, edges of the boxes=25th and 75th percentiles, whiskers=10th and 90th percentiles and the circles outside = 5th and 95th percentiles of the distributions. White boxes represent experimental data while grey shaded boxes denote simulated time-series. Average (± SEM) *FI* of JB and FEJ populations corresponding to the experimental and simulated time-series under (**B**) LH and (**C**) HL regimes. Experimental data shows that in both the regimes, FEJs have lower *FI* than the JBs, as predicted by the model. Simulated FEJ populations capture well the empirical trends for population size distribution and constancy stability.

3.7 Comparison with experiments and models on other species

Since our model is based on fairly generic life-history features (see Text S3), it is reasonable

to assume that it would be a good predictor of the dynamics across a wide range of taxa. For

example, in another dipteran, the pitcher-plant mosquito (*Wyeomyia smithii*) increased larval

food amounts reduces larval mortality and increases adult fecundity (Istock et al. 1975).

Furthermore, in the same species, adult fecundity is positively and negatively affected by

increased levels of adult food and adult crowding respectively (Istock et al. 1975). Since

these are exactly the processes that govern the dynamics of *Drosophila* laboratory

populations, it can be safely predicted that the dynamics of *W. smithii*, under different

nutritional levels, will be well captured by our model. Unfortunately, we could not locate any

studies on the population dynamics of this species to verify this prediction.

Paucity of studies in the literature was not the problem with the widely-studied crustacean

genus *Daphnia*. In this system, increasing food concentration increases size at maturity and

fecundity (Boersma and Vijverberg 1995; Rinke and Vijverberg 2005), as well as the slope of

the body length-clutch size relationship (Rinke and Vijverberg 2005), which is equivalent to

increasing the value of *sen_adsize* in our model. An IBM incorporating these relationships

predicted increased amplitude of fluctuations in population sizes with increasing

concentration of food (Vanoverbeke 2008), which agrees well with experimental

observations (McCauley et al. 1999) and our results (Figure 2D). Simulation studies on

another crustacean (*Armadillidium vulgare*) showed that increased level of food also

increases the population size (Rushton and Hassall 1987), which is consistent with our

observations (Figure 3A). It should be noted here, though, that all these *Daphnia* experiments

and models deal with continuous culture systems, as opposed to the discrete generation

systems investigated in our study. Moreover, in these systems, the adults and the juveniles

share the same food, whereas in our model, the effects of the larval food (determined by the

parameter *food*) is different from the effects of the adult nutrition (modeled using *adnut*). In

spite of these non-trivial differences, the primary mechanism by which food amounts affect

the life history, and thereby the dynamics, remain alike.

Moving away from the arthropods, not surprisingly, leads to more serious departures from the

insights gathered from our study, although some broad patterns are still discernible. For

example, an IBM for the fish yellow perch (*Perca flavescens*) showed that increased food

(forage fish) amount leads to larger body size, higher fecundity and lower adult abundance

(Rose et al. 1999). However, in the same study, when the amount of food was increased for

another fish, walleye (*Sander vitreus*), the model predicted larger body size, lower fecundity

and higher adult abundance. This difference was partly due to the fact that the life-history

parameters and the ecological food-bases of the two fishes were very different. Although the

authors did not report the effects of these food manipulations on the population dynamics, it

is clear that changing the levels of food can affect the population size, which appears to be a

fairly robust observation in this context. In a comprehensive review covering 138 species,

including reptiles, birds and small mammals, Boutin reported that adding food to the

environment typically leads to 2-3 fold increase in population density but little or no change

in the population dynamics (Boutin 1990; although see Klenner and Krebs 1991 and

references therein). More importantly, litter or clutch sizes are hardly affected by food

supply, except when there is severe food shortage due to natural reasons or severe

overpopulation (Boutin 1990).

There can be several reasons for the differences between the observations of these vertebrate

studies and ours. First, there is no reason to expect that a model based on arthropod life

history can accurately capture the population dynamics of vertebrates. Second, it is very

difficult to estimate the parameter space for the vertebrate systems that is analogous to the

parameter space used in our model. Third, all the vertebrate studies quoted here are on natural

populations with multiple sources of food and other ecological interactions, at least some of

which can interact with the effects of food manipulation in non-intuitive ways (e.g. Klenner

and Krebs 1991).

The above account suggests that there are substantial differences between how food levels

affect the dynamics of insects / crustaceans on one hand and higher vertebrates on the other.

Not surprisingly, our model is a better descriptor of the former. With that knowledge, we

were in a position to go beyond laboratory populations of *D. melanogaster* and investigate

the general consequences of various demographic processes for population dynamics. For the

purpose of this study, we chose to investigate the effects of unequal numbers of males and

females, and particularly its interaction with the amount of food available, on population size

and stability.

3.8 Simulations: The effects of sex-ratio and sex-specific culling are greatly influenced

by fecundity but not by levels of juvenile food

Theoretical studies have typically neglected the effects of sex ratio on the stability of

population dynamics (although see Johnson 1994), an important lacuna that we address here.

We found that when the baseline fecundity is high, the *FI* of the population is highest when

the sex ratio is close to 1:1 and reduces when the sex ratio becomes more skewed (closed

circles in Figure 5A). This is consistent with the observation that population growth rate

typically maximizes either when both sexes are present in equal proportions or when the

proportion of females is slightly higher (Jenouvrier et al. 2010; Miller and Inouye 2013).

However, although equal or female-biased sex ratios are also expected to maximize the

average population size (Rankin and Kokko 2007; Schmickl and Karsai 2010), our model

showed that the maximum population size was reached when the proportion of females in the

population was ~15%, i.e. when the population was male-biased (closed circles in Figure

5B). This general pattern was not altered irrespective of the amount of food given to the

larvae (*cf* Figure 5 with Figure S9).

Figure 5. Effect of varying sex-ratio and sex-specific culling on population dynamics for low nutrient scenario. Figure 5. Effect of varying sex-ratio and sex-specific culling on population dynamics under low nutrient levels. **A)** Fluctuation index and **B)** Average population size as a function of the expected fraction of females in the population. **C)** Fluctuation index and **D)** Average population size as a function of the percentage of females culled. **E)** Fluctuation index and **F)** Average population size as a function of the percentage of males culled. Closed and open circles denote high and low levels of density-independent adult fecundity. See section 3.8 for explanations. Each point represents average (± SEM) fluctuation index or population size of 100 replicates of 100-gen long simulated time series. Error bars are too small to be visible.

To explain this observed discrepancy, we note that many of the previous studies on effects of

sex ratio assume the mating success of the females to be a function of the number of available

males (Jenouvrier et al. 2010; Rankin and Kokko 2007; Schmickl and Karsai 2010).

Although applicable for strictly monogamous species, this assumption need not always be

true for insects like *D. melanogaster* where both males and females mate multiple times

(although see Vahl et al. 2013), and a single mating can sustain high lifetime fecundity of a

female, unless the proportion of males is extremely low. That is the reason our model

assumed that all females contributed to the egg numbers of the next generation, irrespective

of the number of males in the population (we relax this assumption in the next section).

Consequently, male numbers can only have negative effects on the number of eggs in the

next generation through the density-dependent reduction in fecundity. On the other hand,

each female in the population has a positive effect on the number of eggs in the next

generation through its fecundity, and a negative effect through its contribution to the density-

dependent reduction in fecundity (eq. 4). The sex ratio leading to the maximum population

size is thus determined as an interaction between these two opposing effects. When female

fecundity is high and not male-limited, this number is much lower than what is expected

when these two conditions are not met. This explains why, in our model, the maximum

population size is attained under male-biased conditions. When the number of females in the

population is increased beyond this threshold, the number of eggs in the next generation

increases, which, in turn, results in more intense larval crowding. Conceptually, this is similar

to boosting the fecundity by supplying yeast to the adults which reduces the average

population size (Figure 1A), as can be seen by comparing LL with LH and HL with HH in

Figure 1A.

When the density-independent fecundity was low (open circles in Figure 5A and 5B) there

was monotonic increase and decrease in average population size (Figure 5B) and *FI* (Figure

5A), respectively, with increasing proportion of females in the population. This trend was

clearly different from the high fecundity case (*cf* open circles and closed circles in Figure 5A

and Figure 5B), indicating that fecundity itself can alter the effects of sex ratio on population

dynamics. To the best of our knowledge, this effect has previously not been reported in the

literature and can be understood as follows. When the per capita female fecundity is low, the

number of eggs in the next generation is reduced and the average size of the population is

therefore limited more by the total number of eggs produced than the larval density-

dependent mortality. Moreover, under such circumstances, the populations are also more

vulnerable to demographic stochasticity, which leads to an increase in *FI*. Consequently, with

increases in proportion of females, both the average size of the population and the population

stability increases.

Sex-specific mortality or culling of the adult individuals: In a population, sexes often suffer

unequal mortality rates. This can happen due to, *inter alia*, sex-biased predation (Boukal et al.

2008) or human preference for a given sex for commercial or other purposes. We modeled

this phenomenon using our model by explicitly culling a fixed percentage of a sex in each

generation and, not surprisingly, found that female (Figure 5C-D) and male (Figure 5E-F)

culling have very different effects on the dynamics. The patterns observed in Figure 5C-F can

be explained by noting that female culling increases the ratio of males in the population and

vice versa, and the subsequent effects on *FI* and average population size can be deduced

based on Figure 5A-B. One interesting point to note is that when the female fecundity is high,

culling of females can increase the average population size, a phenomenon that is termed as

the Hydra effect in the population dynamics literature (Abrams 2009). However, when female

fecundity is low, increasing the fraction of females culled reduces population size, which is

consistent with recent theoretical and empirical data on the beetle *Callosobruchus maculatus*

(Snyder et al. 2014). Moreover, although changing the amount of juvenile food led to

changes in the numerical values of the average population size and *FI*, the corresponding

patterns remained unaltered (*cf* Figure 5C-F and Figure S9C-F), suggesting that the effects of

changes in sex-ratio on population dynamics are not altered by juvenile nutrition.

To summarize, different amounts of juvenile food affects the interaction of sex ratio and

population dynamics quantitatively but not qualitatively. However, in species like

Drosophila, where adult fecundity can be increased by giving external food (equivalent to

increasing *adnut*), adult food levels will play a role in determining the qualitative and

quantitative effects of unequal numbers of sexes on the dynamics.

612 3.9 Simulations: Fecundity and nutrition levels interact to determine the efficacy of

613 Sterile Insect Technique (SIT)

One of the major applications of skewed sex-ratios has been in the context of SIT, wherein a

large number of sterile individuals (typically males) are released into an insect pest

population (Klassen 2005). These sterile individuals compete with the fertile individuals, thus

reducing their reproductive fitness, which ultimately leads to effective suppression of

population density or complete eradication of the pests (Knipling 1955). This method has

been successfully used to eradicate several pest species like melon fly, Tsetse fly,

screwworm, Mexican fruit fly and West Indian fruit fly (see Dyck et al. 2005 for a

comprehensive review). Not surprisingly, the consequences of releasing large number of

sterile males have been extensively investigated in the theoretical population dynamics

literature (reviewed in Barclay 1980) primarily using simple non-linear models (e.g.

Berryman et al. 1973). These studies have typically not incorporated the details of life-history

or environmental variables like food amount on the efficacy of SIT. In fact, we could locate

very few individual or agent-based models on SIT in the literature (Isidoro et al. 2009; Lin et

al. 2015; Stone 2013). Simple models predict that when the fecundity of females is high,

introducing sterile males can increase the pest population size (Berryman et al. 1973),

whereas the present study shows that changes in larval food amount can affect the dynamics

of populations. Therefore, we used our model to explore the interaction of the amount of food

and fecundity of the pests on the expected number of individuals needed for a successful pest

eradication using SIT.

Figure 6. Effect of varying food and density-independent fecundity of the model on sterile male technique. For each combination of *food* and density-independent fecundity, this graph denotes the minimum number of sterile males needed to induce extinction within 10 generations with 90% certainty. Each point here is an average value obtained from 100 replicate simulations and plotted according to the adjacent colour index. Broadly, when fecundity is low, required number of sterile males does not depend on the amount of available food. But, for higher fecundity values, food and fecundity together determine the required number of sterile males needed for the desired level of extinction. For this analysis, we have considered only those food × fecundity combinations, where the basal probability of extinction is <10%.

Following prior studies (Knipling 1959), we assumed that the number of females that get

fertilized and lay eggs is a function of the ratio of fertile to sterile males in the population (see

section 2.5). As noted earlier, neither our fly populations nor our model-derived time series

underwent too many extinctions. However, for this set of simulations, we needed to consider

parameter values of food and density-independent fecundity (i.e. *adnut* $\times x_5$ in equation 3)

that were very different (greater as well as lesser) than what was applicable for our fly

populations. Consequently, the extinction rates no longer remained negligible which in turn

meant that sterile male induced extinctions needed to be distinguished from endogenous

extinctions. Therefore, we first simulated the probability of extinction over 10 generations for

various combinations of fecundity and food values (Figure S10). From this set, we used only

those fecundity-food combinations which had less than 10% probability of going extinct

without any perturbation to estimate the minimum number of sterile insects to be introduced

that could lead to population extinction within 10 generations with a 90% probability (Figure

6). Thus, the lower the number of males needed to induce extinction, the higher was the

efficiency of SIT. We found that when density-independent fecundity is $<\sim40$, changing the

amount of food had no effects on SIT efficiency. However, beyond this range, the efficiency

of SIT reduced with the amount of food available, and the effect became more prominent as

fecundity increased. The reason for this observation can be understood when we consider the

effects of increasing the amounts of food for different fecundity levels. When fecundity is

low, larval crowding is less and therefore, increasing the amount of food does not increase

the population size. Consequently, the number of sterile males needed to bring extinction is

also unaffected. However, when the fecundity is high, there is considerable larval crowding.

Consequently, increasing the amount of food increases the average population size (as seen in

Figure 3A) which in turn reduces the efficiency of SIT. Thus we find that the efficiency of

SIT depends on an interaction between the amount of food available to the larva and its

density independent fecundity (which can be altered by environmental factors). To the best of

our knowledge, this interaction has not hitherto been reported. It should be noted here, that

for our flies, the values of density independent fecundity (i.e. *adnut* $\times x_5$) considered were 85

and ~125 for un-yeasted and yeasted conditions respectively. Therefore, the observed

interaction between food and fecundity happens in a parameter range for fecundity ($>\sim40$)

which is very relevant biologically. Another point regarding these results is the order of

magnitude of the population sizes. In nature, pest population sizes would typically be in

orders of 10^5-10^7 whereas in our simulations, the population sizes were in orders of 10^1-10^2.

This is because it is known that SIT is typically much more effective at low population sizes

and therefore any application of SIT in the field is typically preceded by a chemical treatment

that greatly reduces the population size (Klassen 2005). Thus, although the population sizes

in our simulations are smaller than what would be experienced in a real-world application of

SIT, they are off by no more than 1-2 orders of magnitude. Nevertheless, we believe that our

results are still relevant to SIT application since the purpose of our study was not to model

the application of SIT to a particular real world population of a specific insect pest species,

but rather to derive insights about how outcomes of sterile male release could possibly be

affected by nutrition × fecundity interactions.

4. CONCLUSION

Mathematical modeling of the dynamics of laboratory populations has a long and venerable

history (Kingsland 1995; Mueller and Joshi 2000) and has been extensively done for several

model systems like *Tribolium* (Costantino et al. 1997), *Callosobruchus* (Tuda and Shimada

2005), protists (Holyoak et al. 2000), mites (Benton and Beckerman 2005) etc. Depending on

the objectives of their investigation, these studies have employed different kinds of modeling

tools, ranging from simple deterministic difference equations, to coupled differential

equations and individual-based models (reviewed in Mueller and Joshi 2000). The value of

our study is first in the close correspondence between empirical observations and simulation

results, and second in terms of the insights gained regarding the interaction of the

environmental factors (larval and adult food level) with life-history related traits to determine

population dynamics and stability. Our model was able to replicate several predictions about

the dynamics of insects and crustaceans. Given that these two taxa represent about 70% of all

animal species on earth (Zhang 2011), one can be reasonably confident that the general

insights derived from this study are broadly applicable.

ACKNOWLEDGEMENTS

We thank N. G. Prasad, Mallikarjun Shakarad and M. Rajanna for help in conducting the experiments, and Neelesh Dahanukar for helpful discussion regarding data analysis. For financial support, ST thanks JNCASR, Bengaluru, for a Project Oriented Biological Education (POBE) fellowship, and the Council for Scientific and Industrial Research, Government of India, for a Senior Research Fellowship. AJ thanks the Science and Engineering Research Board of the Department of Science and Technology, Government of India, for support through a J. C. Bose National Fellowship. This study was partly sponsored by an extra-mural grant from the Council of Scientific and Industrial Research, Government of India, and in-house funding from IISER-Pune and JNCASR, Bengaluru.

REFERENCES

Abrams, P. A. 2009. When does greater mortality increase population size? The long history
and diverse mechanisms underlying the hydra effect. Ecology Letters 12:462-474.
Bakker, K. 1961. An analysis of factors which determine success in competition for food
among larve of *Drosophila melanogaster*. Archives Neerlandaises de Zoologie
14:200-281.
Barclay, H. J. 1980. The sterile insect release method on species with two-stage life cycles.
Researches on Population Ecology 21:165-180.
Benton, T. G., and A. P. Beckerman. 2005. Population dynamics in a noisy world: lessons
from a mite experimental system. Advances in Ecological Research 37:143-181.
Berec, L., D. Maxin, and V. Bernhauerová. 2016. Male-killing bacteria as agents of insect
pest control. Journal of Applied Ecology 53:1270-1279.
Berryman, A. A., T. P. Bogyo, and L. C. Dickmann. 1973. Computer simulation of
population reduction by release of sterile insects: II. The effects of dynamic survival
and multiple mating, Pages 31-43 Computer models and application of the sterile
male technique. Vienna, International Atomic Energy Agency.
Boersma, M., and J. Vijverberg. 1995. Synergistic effects of different food species on life-
history traits of *Daphnia galeata*. Hydrobiologia 307:109-115.
Boukal, D. S., L. Berec, and V. Křivan. 2008. Does sex-selective predation stabilize or
destabilize predator-prey dynamics? . PLoS One 3:e2687.
Boutin, S. 1990. Food supplementation experiments with terrestrial vertebrates: patterns,
problems, and the future. Canadian Journal of Zoology/Revue Canadienne de
Zoologie 68:203-220.
Costantino, R. F., R. A. Desharnais, J. M. Cushing, and B. Dennis. 1997. Chaotic dynamics
in an insect population. Science 275:389-391.
De Jong, G. 1976. A model of competition for food. I. Frequency-dependent viabilities.
American Naturalist 110:1013-1027.
Dey, S. 2007. Experimental and theoretical investigation of the dynamics and stability of
single populations and metapopulations of *Drosophila melanogaster* in the laboratory.
Doctoral thesis, Jawaharlal Nehru Centre for Advanced Scientific Research,
Bangalore.
Dey, S., and A. Joshi. 2006. Stability via asynchrony in *Drosophila* metapopulations with
low migration rates. Science 312:434-436.
Dey, S., and A. Joshi. 2013. Effects of constant immigration on the dynamics and persistence
of stable and unstable *Drosophila* populations. Scientific Reports 3:1405.
Dey, S., N. G. Prasad, M. Shakarad, and A. Joshi. 2008. Laboratory evolution of population
stability in Drosophila: constancy and persistence do not necessarily coevolve.
Journal of Animal Ecology 77:670-677.
Dyck, V. A., J. Hendrichs, and A. S. Robinson. 2005. Sterile insect technique: principles and
practice in area-wide integrated pest management. Dordrecht, The Netherlands,
Springer.
Grimm, V., and C. Wissel. 1997. Babel, or the ecological stability discussions: an inventory
and analysis of terminology and a guide for avoiding confusion. Oecologia 109:323-
334.
Halvorsen, K. T., T. K. Sørdalen, L. A. Vøllestad, A. B. Skiftesvik, S. H. Espeland, and E. M.
Olsen. 2017. Sex- and size-selective harvesting of corkwing wrasse (*Symphodus
melops*)—a cleaner fish used in salmonid aquaculture. ICES Journal of Marine
Science 74:660-669.

Holyoak, M., S. Lawler, P., and P. H. Crowley. 2000. Predicting extinction: Progress with an individual-based model of protozoan predator and prey. Ecology 81:3312-3329.

Honěk, A. 1993. Intraspecific variation in body size and fecundity in insects: a general relationship. Oikos 66:483-492.

Isidoro, C., N. Fachada, F. Barata, and A. Rosa. 2009. Agent-Based Model of *Aedes aegypti* Population Dynamics, Pages 53-64 *in* L. S. Lopes, N. Lau, P. Mariano, and L. M. Rocha, eds. Progress in Artificial Intelligence: 14th Portuguese Conference on Artificial Intelligence, EPIA 2009, Aveiro, Portugal, October 12-15, 2009. Proceedings. Berlin, Heidelberg, Springer Berlin Heidelberg.

Istock, C. A., S. S. Wasserman, and H. Zimmer. 1975. Ecology and evolution of the pitcher-plant mosquito: 1. population dynamics and laboratory responses to food and population density. Evolution 29:296-312.

Jenouvrier, S., H. Caswell, C. Barbraud, and H. Weimerskirch. 2010. Mating behavior, population growth, and the operational sex ratio: a periodic two-sex model approach. American Naturalist 175:739-752.

Johnson, S. D. 1994. Sex ratio and population stability. Oikos 69:172-176.

Kingsland, S. 1995, Modeling nature: episodes in the history of population ecology. Chicago, University of Chicago Press.

Klassen, W. 2005. Area-wide integrated pest management and the sterile insect technique *in* V. A. Dyck, J. Hendrichs, and A. S. Robinson, eds. Sterile insect technique: principles and practice in area-wide integrated pest management. Dordrecht, The Netherlands, Springer.

Klenner, W., and C. J. Krebs. 1991. Red squirrel population dynamics I. The effect of supplemental food on demography. Journal of Animal Ecology 60:961-978.

Knipling, E. F. 1955. Possibilities of insect control or eradication through the use of sexually sterile males. Journal of Economic Entomology 48:459-462.

Knipling, E. F. 1959. Sterile-male method of population control. Science 130:902-904.

Lin, S., M. H. DeVisser, and J. P. Messina. 2015. An agent-based model to simulate tsetse fly distribution and control techniques: A case study in Nguruman, Kenya. Ecological Modelling 314:80-89.

May, R. M. 1976. Simple mathematical models with very complicated dynamics. Nature 261:459-467.

McCauley, E., R. M. Nisbet, W. Murdoch, A. M. DeRoos, and W. S. C. Gurney. 1999. Large-amplitude cycles of Daphnia and its algal prey in enriched environments. Nature 402:653-656.

Miller, T. E., and B. D. Inouye. 2013. Sex and stochasticity affect range expansion of experimental invasions. Ecology Letters 16:354-361.

Mueller, L. D. 1985. The evolutionary ecology of *Drosophila*. Evolutionary Biology 19:37-98.

—. 1988. Density-dependent population growth and natural selection in food limited environments: The *Drosophila* model. American Naturalist 132:786-809.

Mueller, L. D., and P. T. Huynh. 1994. Ecological determinants of stability in model populations. Ecology 75:430-437.

Mueller, L. D., and A. Joshi. 2000, Stability in model populations. Princeton, Princeton University Press.

Mueller, L. D., A. Joshi, and D. J. Borash. 2000. Does population stability evolve? Ecology 81:1273-1285.

Nunney, L. 1983. Sex differences in larval competition in *Drosophila melanogaster*: the testing of a competition model and its relevance to frequency-dependent selection. American Naturalist 121:67-93.

Perlman, S. J., C. N. Hodson, P. T. Hamilton, G. P. Opit, and B. E. Gowen. 2015. Maternal
transmission, sex ratio distortion, and mitochondria. Proceedings of the National
Academy of Sciences, USA 112:10162-10168.
Prasad, N. G., S. Dey, M. Shakarad, and A. Joshi. 2003. The evolution of population stability
as a by-product of life-history evolution. Proceedings of the Royal Society of London,
Series B: Biological Sciences (Supplement: Biology Letters) 270:S84-86.
Prout, T., and F. McChesney. 1985. Competition among immatures affect their fertility:
population dynamics. American Naturalist 126:521-558.
Rankin, D. J., and H. Kokko. 2007. Do males matter? The role of males in population
dynamics. Oikos 116:335-348.
Rinke, K., and J. Vijverberg. 2005. A model approach to evaluate the effect of temperature
and food concentration on individual life-history and population dynamics of
Daphnia. Ecological Modelling 186:326-344.
Rodriguez, D. J. 1989. A model of population dynamics for the fruit fly *Drosophila*
melanogaster with density dependence in more than one life stage and delayed
density effects. Journal of Animal Ecology 58:349-365.
Rose, K. A., E. S. Rutherford, D. S. McDermot, J. L. Forney, and E. L. Mills. 1999.
Individual-Based Model Of Yellow Perch And Walleye Populations In Oneida Lake.
Ecological Monographs 69:127-154.
Rushton, S. P., and M. Hassall. 1987. Effects of food quality on isopod population dynamics.
Functional Ecology 4:359-367.
Sah, P., J. P. Salve, and S. Dey. 2013. Stabilizing biological populations and metapopulations
by Adaptive Limiter Control Journal of Theoretical Biology 320:113-123.
Schmickl, T., and I. Karsai. 2010. The interplay of sex ratio, male success and density-
independent mortality affects population dynamics. Ecological Modelling 221:1089-
1097.
Sheeba, V., and A. Joshi. 1998. A test of simple models of population growth using data from
very small populations of *Drosophila melanogaster*. Current science 75:1406-1410.
Snyder, K. T., N. A. Freidenfelds, and T. E. Miller. 2014. Consequences of sex-selective
harvesting and harvest refuges in experimental meta-populations. Oikos 123:309-314.
Stone, C. M. 2013. Transient population dynamics of mosquitoes during sterile male releases:
modelling mating behaviour and perturbations of life history parameters. PLoS One
8:e76228.
Tuda, M., and M. Shimada. 2005. Complexity, evolution, and persistence in host-parasitoid
experimental systems with *Callosobruchus* beetles as the host Advances in Ecological
Research 37:37-75.
Tung, S., A. Mishra, and S. Dey. 2014. A comparison of six methods for stabilizing
population dynamics. Journal of Theoretical Biology 356:163-173.
Vahl, W. K., G. Boiteau, M. E. de Heij, P. D. MacKinley, and H. Kokko. 2013. Female
fertilization: effects of sex-specific density and sex ratio determined experimentally
for Colorado potato beetles and *Drosophila* fruit flies. PLoS One 8:e60381.
Vanoverbeke, J. 2008. Modeling individual and population dynamics in a consumer–resource
system: behavior under food limitation and crowding and the effect on population
cycling in *Daphnia*. Ecological Modelling 216:385-401.
Zar, J. H. 1999, Biostatistical Analysis. New Jersey, Prentice Hall.
Zhang, Z. Q. 2011. Animal biodiversity: An introduction to higher-level classification and
taxonomic richness. Zootaxa 3148:7-12.

Supplementary online material

for

Understanding the dynamics of laboratory populations of *Drosophila melanogaster*: long-term experiments meet individual-based modelling

Authors: Sudipta Tung[a], M. Rajamani[b,c], Amitabh Joshi[b] and Sutirth Dey[a]*

Affiliations:

[a]Population Biology Laboratory, Biology Division, Indian Institute of Science Education and Research-Pune, Dr. Homi Bhabha Road, Pune, India, 411 008

[b]Evolutionary Biology Laboratory, Evolutionary and Organismal Biology Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Jakkur P.O., Bengaluru, India, 560 064

[c]Present Address: Clintus Network Ltd., No. 205/1 & 2, PCR Warehouse Complex, Soukya Road, Koralur Village, Hoskote, Bengaluru 562 114

*Name and address of the corresponding author:
Sutirth Dey
Associate Professor, Biology Division
Indian Institute of Science Education and Research (IISER) Pune
Dr. Homi Bhabha Road, Pune, India 411 008
Tel: +91-20-25908054
Email: s.dey@iiserpune.ac.in

Text S1. Laboratory ecology of *Drosophila melanogaster*:

In laboratory cultures of D. melanogaster, if the larval crowding is high, the mean amount of
food available per larva is reduced. As a result, a large proportion of larvae are unable to
attain the critical body mass needed for successful pupation, thus increasing larval mortality
(Bakker 1961). Since the body size of the adults depends mainly on the amount of resources
gathered during the larval stage, the adults emerging out of crowded cultures are generally
small in size (Marks 1982) and exhibit low fecundity (Chiang and Hodson 1950). Adult
fecundity is also reduced with increasing density of adults in a culture and this is generally
attributed to increased interference with egg laying (Pearl 1932). Interestingly, this negative
effect of adult density on fecundity can be ameliorated to a substantial degree by supplying
the adults with excess amount of live yeast paste (Mueller and Huynh 1994). Since survival
and fecundity are the major factors affecting the growth rate of a population, it seems
plausible that these three density-dependent feedback loops — effects of larval crowding on
larval survivorship and subsequent adult fecundity, and effects of adult crowding on adult
fecundity — play a major role in determining the dynamics and stability of D. melanogaster
populations in the laboratory (Mueller and Joshi 2000).

Text S2. Experiment:

The experiment was comprised of thirty-two populations of *D. melanogaster*, each represented by a single vial (9 cm h × 2.4 cm dia.) culture. These populations were derived from a long standing, large outbred population (JB_1), maintained on a 21-day discrete generation cycle. Details of the ancestry and maintenance protocol of the JB populations can be found elsewhere (Sheeba et al. 1998), and are not germane to this study. These 32 populations were randomly allotted to one of four nutritional regimes, such that there were eight populations per regime. Following established norms (Mueller and Huynh 1994; Mueller et al. 2000) these regimes were called HH, HL, LH, LL — where the first letter indicates the quantity of larval food and the second letter represents the status of adult nutrition. In case of larval food, H and L denoted ~6 mL and ~2 mL of banana-jaggery medium per vial, respectively, whereas in the case of adult nutrition, H and L referred, respectively, to the presence and absence of live yeast paste supplement to banana-jaggery medium. Thus, for example, HL denotes a nutritional regime comprising of ~6 mL medium per vial for the larvae, but no live yeast paste supplement for the adults, and so on.

Each population was initiated (generation 0) with eight male and eight female flies, and from this point onwards (except for extinction) there was no direct control on the number of adults in a vial. After oviposition in the vial for 24 hours (day 0), the adults were counted and discarded and the eggs formed the next generation. Once the adults started eclosing in these vials, they were transferred to adult collection vials every day with a change of medium every alternate day. Strict vial-to-vial correspondence was maintained between the egg vials and their corresponding adult collection vials. The process of adult collection continued until 18 days after day 0, after which the flies were conditioned for three days in the presence / absence of live yeast paste. The live yeast paste is known to boost the fecundity of the females (Chippindale et al. 1993) and reduce the effect of adult density on adult fecundity (Mueller and Huynh 1994). On day 21 after egg collection, the adults were transferred to fresh food vials containing ~2 mL or ~6 mL of banana-jaggery medium for a duration of 24 hours. After this, the adults were counted and discarded, while the eggs laid during this period formed the next generation. If there were no adults in a population, then an extinction event was recorded and the population was rescued by allowing four female flies from the ancestral JB_1 population to lay eggs for 24 hours. All extinction were recorded in the adult time series as 4 individuals (i.e. the number of rescuing females).

The complete details of this experiment have been reported in the PhD thesis of one of the authors (Dey 2007).

Text S3. Model formulation:

The model can be divided into two modules: pre-adult and adult. For a given generation t, the pre-adult module takes the number of eggs and the total amount of larval food as input and computes the number of viable adults and the body size of each of those adults as an output. The output of the pre-adult module and the nature of the adult food available act as inputs for the adult module and the output is the total number of eggs produced that form the input for the pre-adult module in generation $t + 1$. Thus, although our model produces the adult numbers in each generation, structurally it is an egg-to-egg recursion. This modelling strategy has been employed earlier (Mueller 1988), and is preferred over an adult-to-adult recursion. This is because, due to density-dependent mortality, for a given amount of larval food, the relationship between adult numbers and the corresponding number of eggs from which they have arisen is single-humped (Chiang and Hodson 1950). Consequently a given number of adults can arise from very different number of eggs (Prout and McChesney 1985). Thus, for example, assuming say 10% mortality at low crowding, 10 eggs will always lead to ~9 adults. However, if one sees 9 adults, this could have arisen from 10 eggs (assuming 10% mortality at low crowding) or 100 eggs (assuming say 91% mortality at high crowding). Thus, tracking the adult numbers is never sufficient for the purpose of modelling *Drosophila* dynamics (Prout and McChesney 1985).

S3.1 *Pre-adult module*:

S3.1.1 *Step 1: Obtaining number of larva (*`numlarva`*) from the number of eggs (*`numegg`*):*

This module starts with a given number of eggs (`numegg`) and assumes that only a fixed fraction (*hatchability*) of them will hatch into larvae, due to density-independent mortality. Thus, the number of viable larvae is given by

$$\texttt{numlarva} = hatchability \times \texttt{numegg} \quad \ldots\ldots\ldots\ldots\ldots\ldots..(1)$$

where $0 \leq hatchability \leq 1$. Reasons like intrinsic poor viability of the eggs, environmental stresses (say some toxins in the environment) or other ecological/random stochastic factors, can make *hatchability* < 1. However, under normal laboratory conditions, hatchability generally remains above 0.9 (Bakker 1961). Therefore, we have taken *hatchability* = 0.98 and kept it same for all the simulations throughout the study (unless explicitly mentioned otherwise). Note that here we consider *hatchability* as a density-independent parameter because there is no experimental evidence to suggest that hatchability is affected by egg density. However, *hatchability* can be easily made a function of egg or adult density, without affecting the other parts of the model.

S3.1.2 *Step 2: Obtaining the body size of each larva*

In a *Drosophila* culture, the newly hatched larvae eat the larval food provided and grow in size and body mass. Although in the strict sense, size and mass are different quantities, for

the sake of simplicity, we use them interchangeably to indicate biological growth. Due to
among-individual variation in traits like larval feeding rate, food-to-biomass conversion
efficiency etc., a distribution of larval body sizes ensues at the end of the larval growth period
(Bakker 1961). When the number of larvae in the food is increased, the amount of food
available per larva, on an average, is reduced which, in turn, reduces the average body-size
attained at the end of the larval stage (Chiang and Hodson 1950; Miller and Thomas 1958).
Following a previous study (Bakker 1961) we assumed the distribution of larval body sizes at
the end of feeding to be normal with a mean (mean_size) that was an increasing function of
the total amount of larval food (*food*), but a decreasing function of the number of larvae
(`numlarva`). Specifically,

mean_size = $x_1 \times (1 - 1/(x_2 + \exp(-x_3 \times$ `numlarva` $+ food)))$ …..……(2)

where x_1, x_2 and x_3 are parameters (non-negative constants) and exp is the exponential
function. The function is a logistic function of '*food* - $x_3 \times$ `numlarva`', which has an upper
limit x_1 and lower limit $x1(1-1/x_2)$. Thus, theoretically, x_1 could be defined as the maximum
possible average body size (theoretically attained when number of larvae is very small and
amount of food is close to infinite) and $1/x_2$ as the maximum fractional reduction in average
body size (theoretically attained when the number of larvae is infinitely large and the amount
of food is vanishingly small). We arbitrarily assigned $x_1 = 2.5$ and $x_2 = 1$. The parameter x_3
determines the rate of decline of mean_size with increasing the number of larvae for a given
value of *food*. The values of x_3 and low-high values of *food* were calibrated systematically
(see Text S4) to match the observed empirical pattern of larval viability and body size for
different egg densities with a fixed amount of larval food (Chiang and Hodson 1950;
Rodriguez 1989). Note that here our intention was to adopt a function that can mimic the
general qualitative features of the relationship between mean_size*, food* and `numlarva`.
Equation 2 is just one function that fulfills this criterion and potentially many others would
have also served our purpose (for example, logistic function of *food/*`numlarva` or variants
thereof).

For the sake of simplicity, standard deviation (*sigma_size*) of the body-size distribution was
kept as a density-independent constant (=0.45). Computationally, once `numlarva` is
calculated from equation 1, each larva is assigned a body size value by drawing random
numbers from a normal N(mean_size, *sigma_size^2*) distribution. The absolute function
(abs(x) = -x if $x < 0$ else x) is used to avoid any negative value of body size.

S3.1.3 *Step 3: Larval body size assignment, critical mass cut off and adult body size*

In order to complete metamorphosis and become an adult, *Drosophila* larvae, like those of
many other holometabolous insect species (Davidowitz et al. 2003), need to attain a critical
minimum size before pupation (Bakker 1961; Robertson 1963). To incorporate this
phenomenon into our model, we considered a (deterministic) cut off, critical size (m_c), to be a
density-independent constant (following Mueller 1988) and compared the size of each larvae

against it. All larvae whose body sizes were less than m_c were considered to have failed in
becoming adults. The number of remaining larvae was considered to be the adult population
size (numadult) of the current generation.

1002

Empirical studies indicate a positive correlation between larval and adult body size in
Drosophila (Bakker 1961). Therefore, we considered adult body size to be a linear function
of larval body size, i.e.

$$\text{size\_adult}_i = x_4 \times \text{size\_larva}_i \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots(3)$$

where size_adult$_i$ and size_larva$_i$ denote the body size of the i^{th} larva and the corresponding
adult respectively (size_larva$_i > m_c$) and x_4 is a parameter (= 1.0 for the sake of simplicity),
which maps larval size to adult size. Note that x_4 is necessarily a positive quantity as it relates
the body size of larvae to those of adults (both of which are positive). The nature of equation
3 ensures that heavier larvae lead to heavier adults, irrespective of the value of x_4.

1012

1013 Thus, the pre-adult module takes a life-history variable (numegg) as an input and gives two
1014 life-history related variables, number of adults (numadult) and the distribution of the adult
1015 body sizes (size_adult$_i$), as output.

1016

Recently, it has been discovered that *Drosophila* larvae can exhibit cannibalism under
conditions of extreme food deprivation (Vijendravarma et al. 2013). However, we chose not
to incorporate this phenomenon in the model since the extent of cannibalism among the
larvae under the level of crowding found in our populations is still not known. More
critically, there is no evidence till date to indicate that this is a density-dependent
phenomenon. If we assume larval cannibalism to be density-independent, then this
phenomenon can be easily incorporated into our model by multiplying numadult with
another constant.

1025

S3.2 *Adult module*:

S3.2.1 *Step 1: Assigning a gender to each individual*

The first task in this module is to assign a gender to each adult individual. In this study
(unless explicitly mentioned otherwise), sex-ratio was considered to be independent of adult
numbers and was always taken to be 1:1, so each adult was assigned to be female with
probability 0.5 and male otherwise. However, due to the inherent stochasticity of the process,
the realized sex ratio could deviate slightly from 1:1, which is biologically realistic,
particularly in small populations of the kind that we are studying. *Drosophila* is a sexually
dimorphic species with the females being significantly larger than the males. Therefore,

ideally, only the heaviest individuals should have been designated as females. However, we
ignore this complication in our model and assign sex irrespective of adult body size.

1037

S3.2.2 *Step 2: Calculating the total number of eggs produced by all females*

After the assignment of sex, fecundity of the females is computed based on their body size
and current adult density. In many holometabolous insects, including *Drosophila*, fecundity
or egg laying ability is positively correlated with the body size of the females (Honěk 1993).
Prior studies have assumed that the fecundity of *Drosophila* females scale linearly on a ln-ln
scale (Mueller 1988), which implies that slight increases in body size values in the upper
range can lead to large increases in fecundity. This is somewhat unrealistic as larger flies
would also require to expend substantial amount of energy in body maintenance and therefore
it is more likely that the rate of increase of fecundity would eventually slow down with
increasing body size. This relationship could be modelled in many ways and we chose to use
the logarithmic function to represent the effects of female body size on adult density-
independent fecundity. Finally, live yeast paste is known to boost female fecundity
irrespective of the density (Mueller and Huynh 1994). This phenomenon, is incorporated by
adding a density-independent constant (*adnut*) which denotes the fecundity boost due to yeast
supplement to the adults (for not yeasted, *adnut* = 1; for yeasted, *adnut* >1). Taken together,
the adult density-independent component of fecundity of the i^{th} female can be represented as:

$\qquad$ addens_ind_fec$_i$ = $adnut \times x_5 \times \ln(x_6 + sen\_adsize \times \text{size\_adult}_i)$……………(4)

where *sen_adsize* is the strength of relationship between female-fecundity and adult body size
and x_5 and x_6 are scaling parameters. It should be noted here that in the above formulation,
two of the constants (*adnut* and x_5) can easily be combined to create a single constant.
However, we refrain from that in order to retain the ease of biological interpretation.

1059

Another important factor that reduces per capita female fecundity in insects is adult density
(Mueller 1988; Rich 1956). Following an earlier study (Mueller 1988) we modelled this
relationship as:

$\qquad$ addens_eff = $(1/(1 + sen\_adden \times \text{numadult}))$…………………..(5)

where *sen_adden* is the sensitivity of female-fecundity to adult density. This hyperbolic
function denotes negative feedback of adult density on adult fecundity, a relationship which
is found not only in *Drosophila* (Mueller and Huynh 1994; Rodriguez 1989) but in many
other species as well (Băncilă et al. 2016; Michael and Bundy 1989). This negative feedback
automatically makes population growth negatively density dependent and such negative
feedback is known to be a necessary condition for population regulation (Turchin 1999). The
values of x_5, x_6, *sen_adsize* and *sen_adden* were determined by an extensive search over the
parameter space in order to match the experimentally observed fecundity of a healthy female
fly as well as to satisfy a previously observed pattern of female fecundity versus adult density

for both yeasted and un-yeasted conditions (Fig 2 in Mueller and Huynh 1994; Fig 6.5 in

Mueller and Joshi 2000).

Combining equations 4 and 5, the fecundity of the i^{th} female is given as:

$$fec_i = addens\_ind\_fec_i \times addens\_eff……………(6)$$

such that the number of eggs in the next generation,

$$numegg_{t+1} = \Sigma_i\ fec_i……………………………(7)$$

Thus, the adult module takes two life-history related parameters from the output of the pre-

adult module and returns the number of eggs in the next generation as the output. This output

then serves as the input for the pre-adult module for the next generation, and thus the

iterations continue.

An earlier version of this model, and its correspondence with the experimental data, has been

reported in the Master's Thesis of one of the authors (Tung 2012).

 Text S4. Model calibration:

The calibration of the individual-based model was attained in three sequential steps.

S4.1 *Step 1. Calibrating the pre-adult module*:

Previous studies have shown that the value of m_c is around half the maximum larval size that
is physiologically attainable (reviewed in Prasad and Joshi 2003). Since we had already fixed
the maximum value of the mean larval size (i.e. *mean_size*) at 2.5, we started with an initial
value of $m_c = 1.25$. Further, we arbitrarily started with $x_3 = 0.01$. We then simulated steps 1
and 2 combined (see Text S3.1) to obtain the effects of `numegg` (range 1-1500) on larval
viability and larval body size, for different amounts of *food* (range 0.5-6.0). The upper limits
on the ranges of `numegg` and *food* roughly correspond respectively to the upper ranges in
egg numbers typically seen and the amount of food (in mL) typically used in *Drosophila* vial
populations in the laboratory (Dey et al. 2008). This set of simulations allowed us to narrow
down the values of *food* to a range that roughly corresponded to the patterns and values of
larval survivorship (Bakker 1961; Mueller and Joshi 2000; Rodriguez 1989) and body size
(Bakker 1961; Chiang and Hodson 1950; Rodriguez 1989) in *Drosophila* laboratory
populations. The correspondence between simulation results and prior experimental data
were judged by visual pattern-matching as per standard methodological recommendations
(Railsback and Grimm 2011).

1103

S4.2 *Step 2. Calibrating the adult module*:

We next fine-tuned the parameters of the adult module, such that the fecundity of the female
flies matched those of our flies (Dey et al. 2008). For the un-yeasted condition, we fixed
adnut = 1, so that addens_ind_fec$_i$ in the LHS of eq 4 represented the adult density-
independent fecundity of females. We arbitrarily fixed $x_6 = 2$ and simulated eq 4 across a
range of values for *size_adult*. This range for *size_adult* was obtained from the larval module
(after the calibration described above) for different values of `numegg`. Our aim was to obtain
a value for x_5 and range for *sen_adsize* that that led to a matching of the empirically observed
qualitative patterns of addens_ind_fec with increasing *size_adult* (Chiang and Hodson 1950;
Robertson 1957; Rodriguez 1989) (also see Text S3.2 above). We then calibrated the
sen_adden parameter in eq 5 by visually matching the simulation results to the existing
empirical patterns of adult density-dependent female fecundity decline (Fig 2 in Mueller and
Huynh 1994, Fig 6.5 in Mueller and Joshi 2000).

At this stage, we had ranges of parameter values based on prior studies that roughly matched
various life-history patterns from different prior single-generation experiments. We now used
these ranges to calibrate the model for the various population dynamics indices for our
experimental dataset.

S4.3 *Calibrating for our dataset:*

Using the ranges of the various parameter values obtained above, we simulated 49-generation
long time series experiments to fine tune the parameter values for *food*, *adnut*, *sen_adden* and
sen_adsize. For this, we crossed the ranges of these parameter values to obtain a total of
129,920 parameter combinations (i.e. 8 values of *food* × 58 *sen_adden* × 20 *sen_adsize* × 14
adnut) and for each combination, we ran 50 replicate simulations. From these 50 replicate
time series for each parameter combination, we computed the means of the average
population size, *FI*, coefficient of variation (CV), and first and second autocorrelation lags of
the simulation output. We then obtained the corresponding indices for the eight replicates
each of LL and HH regimes of the experimental time series. The match between the
simulation and experimental data was judged by computing the absolute of the percentage
deviation between the simulation mean and experimental mean for each index and summing
them up over the five indices. This led to a set of parameter values for which the sum over all
deviations were minimized separately for the LL and the HH regimes. We then simulated the
HH and LL time series regimes for these parameter values and did minor heuristic
adjustments to obtain better matches (judged visually as per standard recommendations
Railsback and Grimm 2011) with the various indices of the dynamics of the experimental
populations.

The above calibration process relied on fixing some parameters (e.g. x_1, x_2 etc.) arbitrarily as
constants, some based on prior experimental results (e.g. m_c), and tuning the values of the
other parameters to best match certain properties of the experimental data. Thus, the values of
the best-fit parameters could be potentially very different if a different set of properties of the
experimental data were chosen or the arbitrary constants were set to some other values. This
is an inherent feature of pattern-oriented modelling. Since the main aim of this study was to
vary these parameters and observe the corresponding effects on the dynamics, our calibration
goal was not to arrive at impeccable parameter estimates but to derive a set of values that
enabled our model to reasonably describe multiple facets of the empirical dynamics.

It should be noted here that once the values of *food* were obtained for the LL and HH
regimes, we used the same values for the LH and HL regime. In other words, the LL and HH
regime were equivalent to "training" datasets while the LH and HL regimes were equivalent
to "prediction" datasets. This allowed us to avoid the issue of circularity in terms of
parameterization and judging model performances. The final values of all the parameters
used are presented in Table 1.

Text S5. Comparisons with a previous model:

Our model is similar to a previous model of the population dynamics of *Drosophila melanogaster* (Mueller 1988), with three major differences. First, the previous model was fully deterministic, while ours is stochastic and individual-based (for larvae and adults). This feature allowed us to study the various properties of the population size distributions and compare them with the experimental data, which would not have been possible with a deterministic model (except perhaps for chaotic dynamics). This also allowed us to account for certain biological features that can have a major impact on population dynamics. For example, allotting the sex of every individual using a uniform distribution allowed us to account for demographic stochasticity in the number of females, even though the expected sex ratio was 1:1. The previous model, being deterministic, assumed that a fixed fraction of the individuals in the population were females.

Second, we considered female fecundity to be a logarithmic function of female body size, whereas in the previous study, it was modelled as a power law. This implies that in the previous study, when body size was large, small differences in size translated into large differences in fecundity, which was not the case with our model.

The third difference relates to the way effects of adult density on adult fecundity was modelled. In the previous model, low nutrition for adults refers to a condition where a dilute solution of yeast is provided (Mueller and Huynh 1994), as a result of which, yeast is not limiting at low densities and the fecundity of the flies is as high as the case when yeast is provided (see Fig 2 of Mueller and Huynh 1994). For this reason, in the previous model, the effects of adult nutrition are conceptualized solely as a density-dependent reduction of fecundity, assuming that when densities are low, the fecundities are the same. However, in our study, low food for adults means no yeast at all and consequently, the density-independent fecundity of our non-yeasted flies is much lower. That is why, we model the effects of adding yeast using the parameter *adnut* (which determines how much the fecundity is boosted due to yeast supplement) and assume that the density-dependent reduction in fecundity is similar in magnitude for both yeasted and non-yeasted flies. The latter condition is not a property of our model but gets imposed because we chose to use the same value of *sen_adden* for both LL and HH. Thus, this difference is simply a result of modelling strategies and the data used for model formulation and calibration.

Overall, as stated above, our model is more appropriately considered as a consequential extension of the earlier model (Mueller 1988), rather than a completely new model.

Text S6. Role of extinction in the present study:

Persistence, or the ability to resist extinction, is an important aspect of the dynamics of any population (Grimm and Wissel 1997). Prior empirical studies have suggested that extinction plays a crucial role in determining the dynamics of laboratory populations of *Drosophila melanogaster* (Dey and Joshi 2006; Dey and Joshi 2007). However, most of these studies have been conducted on the LH regime, under extremely low levels of larval food which leads to high extinction frequencies. For example, Dey and Joshi (Dey and Joshi 2006) provided ~ 1mL of larval food and observed a per generation extinction probability of ~0.37. However, the current empirical study used ~2mL of food in the low food regimes, which led to a per generation extinction frequency of 0.06 ($\pm$ 0.0067 SEM) in LH and 0.03 ($\pm$ 0.01 SEM) in LL populations. The two other regimes did not suffer any extinctions at all. Therefore, there was no way for us to compare the regimes in terms of extinctions, which is why we chose to neglect extinction in this study, including in the process of model calibration. Interestingly, even though we neglected extinction during model calibration, the observed values of extinction frequency from the simulations for parameter values mentioned in Table 1, were fairly close to the experimental values (LH: 0.069 ± 0.013 SEM; LL: 0.003 ± 0.003 SEM; HL and HH: 0). We note here that these values refer to the extinction probabilities under parameter values that were close to our experimental conditions (i.e. Figure 1). When we studied the effects of changing the parameter values (Figures 2-6), it is entirely possible that the amount of extinctions observed were very different. Therefore, in principle, for every figure showing the effects of changing a parameter on constancy stability (here, *FI*), a corresponding graph on persistence stability (in the form of extinction probabilities or some related measure) can also be investigated. Although important in its own right, such an investigation is clearly out of scope of this paper which focuses primarily on dynamics in terms of constancy stability.

 Text 7. Model Recalibration for the dynamics of the FEJ-JB populations:

To use the time-series data from an earlier experiment (Dey et al. 2008), we first re-calibrated
our model by reducing the value of m_c of FEJs from 1.1 to 1.0. This is because it has been
suggested that due to selection for faster pre-adult development, the FEJs had a lower value
of m_c (Prasad et al. 2001). Moreover, to keep parity with the experimental data, we used 16
replicates each of FEJ and JB in both HL and LH nutritional regimes, and each replicate was
simulated for 36 generations. Every other detail of the parameter values and the simulation
were identical to those mentioned above. We then compared the population size distributions
and *FI* values of the simulations against those observed from the empirical data.

Text S8. Simulating the dynamics of sex biased populations:

We investigated the dynamics and stability of populations under three scenarios where there could be unequal numbers of males and females, namely systematic distortion of sex ratio, sex-specific culling and the sterile insect technique (SIT). In all these cases, relevant alterations were made in the adult module while the pre-adult module was kept unchanged.

a) Distortion of sex ratio: To simulate sex ratio distortions, we varied the probability of a given adult to be a female in the adult module, while keeping the rest of the model unchanged. We studied the effects of these various sex ratios at two different levels of density-independent basal fecundity, namely high [($adnut \times x_5$)=1.49×85] and low [($adnut \times x_5$) = 1.49×17)]. For each sex ratio × fecundity combination, we also studied the effects of low and high levels of food to the juvenile stage ($food$ = 1.76 and 2.56 respectively). For each sex ratio × fecundity × food combination, we ran 100 replicate time series, each of which was 100 generations long. The mean (±SEM) of average population size and fluctuation index over these 100 replicates was plotted.

b) Sex-specific culling: This was similar to the above case except that in every generation, after the gender was assigned to the adults, we explicitly removed a fixed percentage of males or females from the population. The effect of adult crowding on adult fecundity was computed after the culling.

c) Sterile insect technique (SIT): This is a popular pest control technique (Dyck et al. 2005), wherein large numbers of sterile males of the pest species are introduced into the population. These sterile males compete with the fertile ones for potential mating partners, thereby bringing down the number of successful matings which can lead to offspring. The number of females that mate with fertile males is directly proportional to the ratio of number of fertile: sterile males and inversely proportional to the competitive ability of the sterile males (Knipling 1955). For the sake of simplicity, we considered the proportion of females producing eggs for the next generation as a linear function of the fertile/sterile male ratio. Thus, proportion of egg laying females = min((num_fertile/($\beta \times$num_sterile)), 1), where num_fertile and num_sterile are the number of fertile and sterile males in the population, β is the competitive ability of the sterile males and min() is the minimum function. In this section, we studied the interaction of density-independent female fecundity with the amount of juvenile food available in determining the efficacy of SIT.

In our model, low values of $food$ or high female fecundity can lead to extinctions due to larval overcrowding. This creates a problem in terms of judging the efficacy of introducing sterile males to cause extinction. Therefore, we first investigated the extinction probabilities over a large number of combinations of larval food ($food$, range = 0.5 - 3.0, step size = 0.2) and density-independent fecundity (i.e. $adnut \times x_5$, range = 1 - 150, step size = 5) in the absence of any introduced sterile males. For each $food$ × fecundity combination, we computed the probability of extinction within 10 generations over 10 replicates and plotted the average extinction probability (Figure S10). We then limited our explorations on the efficacy of SIT to only those $food$ × fecundity combinations where the probability of an extinction event within the first 10 generations was less than 0.1. For each $food$ × fecundity

combination, we sought to compute the minimum number of sterile males that would lead to
an extinction within 10 generations. Therefore, we increased the number of sterile males in
step size of 1 and simulated 10 generation long time-series for each *food* × fecundity × sterile
male number combination till there was an extinction. For a given *food* × fecundity
combination, this step was repeated 100 times and the average number of sterile males that
were needed to cause extinction was recorded.

Text S9. No evidence of transients in our experiments and simulations:

Due to logistic constraints, most observed population dynamics time series tend to be short. However, it is well known that the transient behavior of population dynamics models can be very different from the equilibrium behaviors (Hastings 2004; Hastings and Higgins 1994). In this study, to keep parity with our experiments, we had limited the length of each simulated time series to 49. To investigate if the long-term behavior of these time series was any different from the short-term behavior, we simulated the dynamics in each regime for 1000 generations, and computed all the quantities represented in Figure 1 for the last 49 generations (Figure S6). Comparing generations 1-49 with generations 952-1000 revealed no major differences in either the population-size distributions or *FI*. This suggests that the transient dynamics in our model are almost indistinguishable from the longer-term dynamics.

Although their absence is re-assuring from a modelling perspective, transients are very much expected from a biological standpoint. This is because experimental evolution studies suggest that in *Drosophila melanogaster*, even 10-15 generations is often sufficient for noticeable divergence in life-history related traits that can affect the dynamics (for examples see Prasad and Joshi 2003). Therefore, all else being equal, one would expect at least some of the stability determining parameters to evolve during the course of the experiment, which in turn is expected to lead to transient dynamics in a long time-series. Yet, we did not incorporate any evolution in our model, which meant that the various life-history parameters detailed in Table 1, remained constant in a particular simulation run. This was because it has been previously shown that at least over 45 generations, there are no observable changes in stability determining demographic parameters in laboratory populations of *D. melanogaster* (Mueller et al. 2000). Thus, we felt that it was safe to ignore changes in life-history related parameters in the context of our empirical data and, therefore, did not incorporate their evolution in our model. However, we note that the structure of our model is such that it can be very easily extended to incorporate the evolution of stability-determining parameters and the effects of such evolutionary change on population dynamics.

Figure S1. Effects of larval and adult nutrition on constancy stability. Interaction of larval and adult nutrition to determine constancy stability of population is statistically significant. High adult nutrition destabilizes population more when larval nutrition is low. Error bars = 95% CI.

Figure S2. Schematic diagram of the model processes.

Figure S3. Representative empirical time-series of population size over 49-generations for LH, HL, LL and HH regimes respectively.

Figure S4. Representative simulation time-series of population size over 49-generations for LH, HL, LL and HH regimes respectively.

Figure S5. Autocorrelation coefficients of empirical and simulation time-series. For four regimes, we have computed first five autocorrelation lags, averaged over eight replicates for both (**A**) empirical and (**B**) simulation time-series. Error bars indicate SEM.

Figure S6. Population size distribution and constancy stability of short and long term dynamics. Blue boxes and bars represent the data for long term dynamics (generation 951-1000), where transients are excluded, whereas the grey shaded boxes and bars represents short term dynamics (generation 1-49). (**A**) Descriptive statistics of the population size distributions for long and short term dynamics in four regimes. Red dashed lines = means, thin black lines = medians, edges of the boxes=25th and 75th percentiles, whiskers=10th and 90th percentiles and the circles outside = 5th and 95th percentiles of the distributions. (**B**) Average (± SEM) fluctuation index of the population size distributions for long and short term dynamics in four regimes. In both panels, there are no systematic differences in population size distributions and constancy stability between the short- and long-term dynamics.

Figure S7. Effect of varying *hatchability* on population size distribution in four regimes. Red squares = means, thin red lines = medians, edges of the boxes=25th and 75th percentiles of the distributions.

Figure S8. Effect of varying *sen_adden* and *sen_adsize* on the average egg number and egg-to-adult viability. Each point represents average (± SEM) fluctuation index of 100 replicates of 100-generation long simulated time series. Error bars are too small to be visible. Effects of sensitivity to adult density (*sen_adden*) on **A.** Average egg number and **B.** Egg-to-adult viability. Effects of sensitivity to adult size (*sen_adsize*) on **C.** Average egg number and **D.** Egg-to-adult viability. Note that in B and D LH is the least affected by increases in the parameter values. See text (section 3.4.2) for explanation.

Figure S9. Effect of varying sex-ratio and sex-specific culling on population dynamics for high nutrient scenario. Fluctuation index and average population size were computed as the proportion of females (**A** and **B**), percentage of female culling (**C** and **D**) and male culling (**E** and **F**) increases in the population respectively. Each cases were investigated at two levels of density-independent fecundity. Here, each point represents average (± SEM) fluctuation index or population size of 100 replicates of 100-gen long simulated time series. Error bars are too small to be visible.

Figure S10. Probability of extinction within 10 generations in *food* × density-independent fecundity space. For each combination of food and fecundity values, we computed average probability of extinction within first 10 generations over 10 replicates and plotted according to the adjacent colour index.

REFERENCES

Bakker, K. 1961. An analysis of factors which determine success in competition for food among larve of *Drosophila melanogaster*. Archives Neerlandaises de Zoologie 14:200-281.

Băncilă, R. I., A. Ozgul, T. Hartel, T. Sos, and B. R. Schmidt. 2016. Direct negative density-dependence in a pond-breeding frog population. Ecography 39:449–455.

Chiang, H. C., and A. C. Hodson. 1950. An analytical study of population growth in *Drosophila melanogaster*. Ecological Monographs 20:173-206.

Chippindale, A., A. M. Leroi, S. B. Kim, and M. R. Rose. 1993. Phenotypic plasticity and selection in *Drosophila* life-history evolution. 1. Nutrition and the cost of reproduction. Journal of Evolutionary Biology 6:171-193.

Davidowitz, G., L. J. D'Amico, and H. F. Nijhout. 2003. Critical weight in the development of insect body size. Evolution and Development 5:188-197.

Dey, S. 2007. Experimental and theoretical investigation of the dynamics and stability of single populations and metapopulations of *Drosophila melanogaster* in the laboratory. Doctoral thesis, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore.

Dey, S., and A. Joshi. 2006. Stability via asynchrony in *Drosophila* metapopulations with low migration rates. Science 312:434-436.

Dey, S., and A. Joshi. 2007. Local perturbations do not affect stability of laboratory fruitfly metapopulations. PLoS One 2:e233.

Dey, S., N. G. Prasad, M. Shakarad, and A. Joshi. 2008. Laboratory evolution of population stability in Drosophila: constancy and persistence do not necessarily coevolve. Journal of Animal Ecology 77:670-677.

Dyck, V. A., J. Hendrichs, and A. S. Robinson. 2005. Sterile insect technique: principles and practice in area-wide integrated pest management. Dordrecht, The Netherlands, Springer.

Grimm, V., and C. Wissel. 1997. Babel, or the ecological stability discussions: an inventory and analysis of terminology and a guide for avoiding confusion. Oecologia 109:323-334.

Hastings, A. 2004. Transients: the key to long-term ecological understanding? Trends in Ecology & Evolution 19:39-45.

Hastings, A., and K. Higgins. 1994. Persistence of transients in spatially structured ecological models. Science 263:1133-1136.

Honěk, A. 1993. Intraspecific variation in body size and fecundity in insects: a general relationship. Oikos 66:483-492.

Knipling, E. F. 1955. Possibilities of insect control or eradication through the use of sexually sterile males. Journal of Economic Entomology 48:459-462.

Marks, R. W. 1982. Genetic variability for density sensitivity of three components of fitness in *Drosophila melanogaster*. Genetics 101:301-316.

Michael, E., and D. A. P. Bundy. 1989. Density dependence in establishment, growth and worm fecundity in intestinal helminthiasis: the population biology of *Trichuris muris* (Nematoda) infection in CBA/Ca mice. Parasitology 98:451-458.

Miller, R. S., and J. L. Thomas. 1958. The effects of larval crowding and body size on the longevity of adult *Drosophila melanogaster*. Ecology 39:118-125.

Mueller, L. D. 1988. Density-dependent population growth and natural selection in food limited environments: The *Drosophila* model. American Naturalist 132:786-809.

Mueller, L. D., and P. T. Huynh. 1994. Ecological determinants of stability in model populations. Ecology 75:430-437.

Mueller, L. D., and A. Joshi. 2000, Stability in model populations. Princeton, Princeton
University Press.
Mueller, L. D., A. Joshi, and D. J. Borash. 2000. Does population stability evolve? Ecology
81:1273-1285.
Pearl, R. 1932. The influence of density of population upon egg production in *Drosophila*
melanogaster. Journal of Experimental Zoology, Part A: Comparative Experimental
Biology 63:57-84.
Prasad, N. G., and A. Joshi. 2003. What have two decades of laboratory life-history evolution
studies on *Drosophila melanogaster* taught us? Journal of Genetics 82:45-76.
Prasad, N. G., M. Shakarad, D. Anitha, M. Rajamani, and A. Joshi. 2001. Correlated
responses to selection for faster development and early reproduction in *Drosophila*
melanogaster: the evolution of larval traits. Evolution 55:1363-1372.
Prout, T., and F. McChesney. 1985. Competition among immatures affect their fertility:
population dynamics. American Naturalist 126:521-558.
Railsback, S. F., and V. Grimm. 2011, Agent-based and individual-based modeling: a
practical introduction. Princeton, Princeton University Press.
Rich, E. R. 1956. Egg cannibalism and fecundity in Tribolium. Ecology 37:109-120.
Robertson, F. W. 1957. Studies in quantitaive inheritance. XI. Genetic and environmental
correlation between body size and egg production in *Drosophila melanogaster*.
Journal of Genetics 55:428-443.
—. 1963. The ecological genetics of growth in Drosophila 6. The genetic correlation between
the duration of the larval period and body size in relation to larval diet. Genetical
Research 4:74-92.
Rodriguez, D. J. 1989. A model of population dynamics for the fruit fly *Drosophila*
melanogaster with density dependence in more than one life stage and delayed
density effects. Journal of Animal Ecology 58:349-365.
Sheeba, V., A. Madhyastha, and A. Joshi. 1998. Oviposition preference for novel versus
normal food resources in laboratory populations of *Drosophila melanogaster*. Journal
of Biosciences 23:93-100.
Tung, S. 2012. Insights into the mechanistic underpinnings for the evolution of population
stability in *Drosophila*, based on modeling of laboratory populations, Indian Institute
of Science Education and Research- Kolkata, Kolkata.
Turchin, P. 1999. Population regulation: a synthetic view. Oikos 84:153-159.
Vijendravarma, R. K., S. Narasimha, and T. J. Kawecki. 2013. Predatory cannibalism in
Drosophila melanogaster larvae. Nature Communications 4:1789.