An FGF-driven feed-forward circuit patterns the cardiopharyngeal

mesoderm in space and time

Florian Razy-Krajka, Basile Gravez, Nicole Kaplan, Claudia Racioppi, Wei Wang and

Lionel Christiaen*

Center for Developmental Genetics, Department of Biology, College of Arts and Science,

New York University, New York, NY, USA

* author for correspondence: email: lc121@nyu.edu, twitter: @lionlchristiaen, phone: +1

212 992 8695

Abstract

In embryos, multipotent progenitors divide to produce distinct progeny and express

their full potential. In vertebrates, multipotent cardiopharyngeal progenitors produce

second-heart-field-derived cardiomyocytes, and branchiomeric skeletal head muscles.

However, the mechanisms underlying these early fate choices remain largely elusive.

The tunicate *Ciona* emerged as an attractive model to study early cardiopharyngeal

development at high resolution: through two asymmetric and oriented divisions,

defined cardiopharyngeal progenitors produce distinct first and second heart

precursors, and pharyngeal muscle (aka atrial siphon muscle, ASM) precursors. Here,

we demonstrate that differential FGF-MAPK signaling distinguishes between heart and

ASM precursors. We characterize a feed-forward circuit that promotes the successive

activations of essential ASM determinants, *Hand-related*, *Tbx1/10* and *Ebf*. Finally, we

show that coupling FGF-MAPK restriction and cardiopharyngeal network deployment

with cell divisions defines the timing of gene expression and permits the emergence of

diverse cell types from multipotent progenitors.

Introduction

Developmental genetics knowledge guided progress towards driving mammalian stem cells into forming pure cultures of selected cell types *in vitro* (e.g. (Kattman et al., 2011; Mazzoni et al., 2011; Peljto and Wichterle, 2011). By contrast, in the embryo, pluripotent cells generate diverse cell types in defined proportions, as they divide before individual daughter cells adopt distinct fates.

Subsets of the heart and head/neck myocytes recently emerged as related derivatives of multipotent progenitors located in the cardiopharyngeal mesoderm (Diogo et al., 2015a; Tzahor, 2009; Tzahor and Evans, 2011). Early lineage tracing, transplantations and controlled explant culture experiments demonstrated that the anterior splanchnic/pharyngeal mesoderm of amniote embryos can produce either skeletal muscles or heart tissue, depending upon exposure to growth factors and signaling molecules (Nathan et al., 2008; Tirosh-Finkel et al., 2006; Tzahor et al., 2003; Tzahor and Lassar, 2001). Clonal analyses in the mouse further revealed the existence of common *Mesp1*-expressing progenitors for subsets of the second heart field-derived cardiomyocytes and branchiomeric facial, jaw, neck and even œsophageal muscles (Gopalakrishnan et al., 2015; Lescroart et al., 2014; Lescroart et al., 2015; Lescroart et al., 2010; Lescroart et al., 2012). In pluripotent stem cells, controlled *Mesp1* expression can drive mesodermal progenitors towards cardiac and/or skeletal muscle fates (Bondue et al., 2008; Chan et al., 2016; Chan et al., 2013). Proper development of the pharyngeal apparatus and second heart field derivatives require shared inputs from Tbx1, Nkx2.5 and Islet1 transcription factors (e.g. (Cai et al., 2003; George et al., 2015; Jerome and Papaioannou, 2001; Kelly et al., 2004; Merscher et al., 2001; Mosimann et al., 2015; Nevis et al., 2013; Prall et al., 2007; Tzahor and Evans, 2011; Vitelli et al., 2002a; Watanabe et al., 2012; Witzel et al., 2017; Yagi et al., 2003; Zhang et al., 2006)). Taken together, this growing body of evidence points to the existence of a mesodermal

field of multipotent progenitors capable of producing either SHF-derived cardiomyocytes or branchiomeric skeletal muscles in early vertebrate embryos (Diogo et al., 2015b; Mandal et al., 2017). However, the mechanisms that distinguish fate-restricted heart and head muscle precursors remain largely elusive.

The tunicate Ciona, which is among the closest living relatives to the vertebrates (Delsuc et al., 2006; Putnam et al., 2008), has emerged as a simple chordate model to characterize multipotent cardiopharyngeal progenitors and the mechanisms that initiate heart vs. pharyngeal muscle fate choices (Kaplan et al., 2015; Razy-Krajka et al., 2014; Stolfi et al., 2010; Tolkin and Christiaen, 2016; Wang et al., 2013). Ciona tailbud embryos possess two multipotent cardiopharyngeal progenitors on either side. Like their vertebrate counterparts, these cells emerge from *Mesp+* progenitors towards the end of gastrulation; they are induced by FGF-MAPK signaling and have been termed *trunk ventral cells* (aka TVCs; (Christiaen et al., 2008; Davidson and Levine, 2003; Davidson et al., 2006; Davidson et al., 2005; Satou et al., 2004; Stolfi et al., 2010)). TVCs activate conserved cardiac markers, including *Hand*, *Gata4/5/6* and *Nk4/Nkx2-5*, and migrate as polarized pairs of cells, until the left and right pairs meet at the ventral midline and begin to divide asymmetrically along the mediolateral axis (Figure 1A; (Christiaen et al., 2008; Davidson et al., 2005; Satou et al., 2004; Stolfi et al., 2010)). The first oriented asymmetric divisions produce small median first heart precursors (FHPs), and large lateral second trunk ventral cells (STVCs), which specifically activate *Tbx1/10* (Davidson et al., 2005; Stolfi et al., 2010; Wang et al., 2013). STVCs later divide again to produce small median second heart precursors (SHPs), and large lateral atrial siphon muscle founder cells (ASMFs), which activate *Ebf* (aka *COE*; (Razy-Krajka et al., 2014; Stolfi et al., 2010; Stolfi et al., 2014c)). The transcription factors Hand-related (Hand-r)/Notrlc, which is expressed in the TVCs and maintained in the STVCs and ASMFs after each division, and Tbx1/10 are required for *Ebf* activation in the ASMFs,

whereas Nk4/Nkx2.5 represses *Tbx1/10* and *Ebf* expression in the second heart

precursors (SHPs)(Razy-Krajka et al., 2014; Tolkin and Christiaen, 2016; Wang et al.,

2013). Conversely, Tbx1/10 and Ebf inhibit cardiac markers, and likely determinants,

such as *Gata4/5/6* and *Hand* (Razy-Krajka et al., 2014; Stolfi et al., 2010; Stolfi et al.,

2014a; Wang et al., 2013). These regulatory cross-antagonisms underlie the transition

from transcriptionally primed multipotent progenitors to separate fate-restricted

precursors, by limiting the deployment of the heart- and pharyngeal-muscle-specific

programs to their corresponding specific precursors (Kaplan et al., 2015).

Here, we identify regulatory mechanisms ensuring the emergence of diverse fate-

restricted precursors from multipotent progenitors. We show that differential FGF-

MAPK signaling, feed-forward regulatory circuits and coupling with the cell cycle

control the spatially restricted activation of *Tbx1/10* and *Ebf*, successively, thus

permitting the emergence of both first and second heart precursors, and

ASM/pharyngeal muscle precursors from common multipotent progenitors.

Results

MAPK signaling is active in the multipotent cardiopharyngeal progenitors and progressively restricted to the pharyngeal muscle precursors.

During the earliest stages of cardiopharyngeal development in ascidians, multipotent progenitors display multilineage transcriptional priming, (Razy-Krajka et al., 2014; Stolfi et al., 2014b), and subsequent regulatory cross-antagonisms segregate distinct cardiopharyngeal programs to their corresponding fate-restricted progenitors (Stolfi et al., 2010; Wang et al., 2013); reviewed in (Kaplan et al., 2015)). For instance, the ASM-specific factor Ebf is necessary and sufficient to terminate the heart program and impose a pharyngeal muscle fate (Razy-Krajka et al., 2014; Stolfi et al., 2010). However, the symmetry-breaking events leading to cardiopharyngeal mesoderm patterning and ASM-specific expression of *Ebf* remain unknown. We surmised that differential signaling inputs determine the stereotyped spatio-temporal patterning of early cardiopharyngeal progenitors.

The Ciona homologs of specific FGF-MAPK pathway components, including *FGF receptor substrate 2/3* (*Frs2/3*; (Gotoh et al., 2004))*, Ets.b*, and *Fgf4/5/6*, are preferentially expressed in the TVCs, in the STVCs and in the ASMFs as cells transition from multipotent progenitor to distinct heart vs. ASM fate-restricted states (Razy-Krajka et al., 2014). These patterned expressions of MAPK effector genes prompted us to evaluate a role for FGF-MAPK pathway in cardiopharyngeal fate decisions.

We first used an antibody specific to the dual phosphorylated form of Extracellular Regulated Kinase (dpERK) to monitor Mitogen Activated Protein Kinase (MAPK) activity in the cardiopharyngeal mesoderm. We detected dpERK staining in the newly born TVCs, marked by the B7.5-lineage-specific *Mesp>H2B::mCherry* transgene (Figure S1), as previously observed (Davidson et al, 2006). We also detected weaker but

persistent dpERK staining in the TVCs during migration (Figures 1 and S1). Following

the first and second asymmetric divisions of the TVCs and STVCs, dpERK staining was

successively restricted to the more lateral STVCs and ASMFs, respectively (Figures 1A,

B; S1).

The canonical FGF/Ras/MEK/ERK pathway is necessary and sufficient to

promote pharyngeal muscle specification in the cardiopharyngeal lineage.

This exclusion of MAPK activity from the medial first and second heart precursors

opened the possibility that differential ERK activity is required for proper STVC and

ASMF vs. heart precursors fate decisions. In Ciona, signaling through the sole FGF

receptor (FGFR) governs ERK activity in several developmental processes, including

neural induction (Bertrand et al., 2003; Hudson et al., 2003) and central nervous

system patterning (Haupaix et al., 2014; Racioppi et al., 2014; Stolfi et al., 2011; Wagner

et al., 2014), early endomesoderm and notochord fate specification (Imai et al., 2002;

Picco et al., 2007; Shi and Levine, 2008; Shi et al., 2009; Yasuo and Hudson, 2007).

Notably, FGF-MAPK signaling is active in the *Mesp+* cardiogenic B7.5 blastomeres

(Imai et al., 2006; Shi and Levine, 2008), where targeted misexpression of a dominant

negative form of FGFR (dnFGFR) blocks TVC induction (Christiaen et al., 2008;

Davidson et al., 2006). We used a TVC-specific *Foxf* enhancer (*Foxf(TVC):bpFog-*

1>dnFGFR::mCherry, hereafter called *Foxf>dnFGFR*; (Beh et al., 2007)), to bypass

early effects and achieve later misexpression of dnFGFR in the TVCs and their progeny.

Importantly, although TVC fate specification and the onset of *Foxf* expression require

FGF-MAPK signaling (Beh et al., 2007; Davidson et al., 2006), we confirmed that this

perturbation altered neither initial TVC induction, nor the expression of the *Foxf* driver

(Figure S2). Consistent with proper TVC induction, *Foxf>dnFGFR* prevented neither

TVC migration nor asymmetric divisions, but it abolished the expression of both

Tbx1/10 in the STVCs and *Ebf* in the ASMFs (Figure 1C). This data indicate that FGF-

MAPK signaling is required in the cardiopharyngeal progenitors and/or their progeny

for ASM fate specification, beyond the initial TVC induction.

Upon FGF-MAPK-dependent induction, the TVCs express *Hand-related/Hand-r*

(renamed after *Notrlc/Hand-like*; (Christiaen et al., 2008; Davidson and Levine, 2003;

Davidson et al., 2006; Satou et al., 2004; Stolfi et al., 2014c; Woznica et al., 2012)),

which encodes a basic helix-loop-helix (bHLH) transcription factor necessary for *Ebf*

expression in the ASMFs (Razy-Krajka et al, 2014). Moreover, the *Hand-r* TVC

enhancer contains putative Ets1/2 binding sites, which are necessary for reporter gene

expression, and presumably mediate the transcriptional inputs of FGF-MAPK (Woznica

et al., 2012). Since *Hand-r* and *Foxf* expressions start at approximately the same time in

newborn TVCs, we used *Foxf>dnFGFR*, which did not alter the onset of *Hand-r*

expression in the TVCs (Figure S2), to test whether the maintenance of *Hand-r*

expression in migratory TVCs requires prolonged FGF-MAPK inputs. *Foxf>dnFGFR*

inhibited *Hand*-r expression in late TVCs (Figure 1C), indicating that sustained *Hand-r*

expression requires continuous FGF-MAPK signaling, as did TVC-expressed FGF-

MAPK pathway components (Figure S2).

To test whether the spatial restriction of MAPK activity explains the patterned

expressions of *Hand-r*, *Tbx1/10* and *Ebf* following asymmetric cell divisions, we used

gain-of-function perturbations to force FGF-MAPK activity throughout the

cardiopharyngeal mesoderm and assayed gene expression (Figure 2). We focused on the

canonical FGF-MAPK pathway where signal transduction involves Ras, Raf, MEK and

ERK downstream of FGFR and upstream of transcriptional effectors (Lemmon and

Schlessinger, 2010). We first used M-Ras^{G22V}, a defined constitutively active form of M-

Ras, which mediates FGF signaling in *Ciona*, where other classical *Ras* genes are

missing (Keduka et al., 2009), and focused on *Htr7* and *Tbx1/10* expression following

the first asymmetric TVC division in 15 hours post-fertilization (hpf) embryos. *Htr7*

encodes a *trans*-membrane G-protein coupled receptor and, like *Hand-r*, its expression

and maintenance in the TVCs require MAPK activity (Figure S2; (Razy-Krajka et al.,

2014)), and become restricted to the lateral STVC following asymmetric division.

However, *Htr7* mRNAs are cleared more rapidly from the FHPs, making the patterned

expression easier to analyze than that of *Hand-r* (Figures 2 and 3D; (Razy-Krajka et al.,

2014)). Importantly, misexpression of M-Ras^{G22V} using the TVC-specific *Foxf* enhancer

altered cell division asymmetry and/or orientation in under 50% of the embryos, still

allowing us to identify large lateral and small median cells in a small majority of

embryos (Figure S3). Compared to control embryos overexpressing wild-type M-Ras

(M-Ras^{WT}), TVC-specific gain of M-Ras function caused both persistent *Htr7* expression

and ectopic activation of *Tbx1/10* in the small median cells following asymmetric

divisions (Figure 2B,C). Similarly, *Foxf>M-Ras^{G22V}*-expressing 18hpf larvae displayed

ectopic *Ebf* activation throughout the cardiopharyngeal mesoderm (Figure 2B,C). This

cannot be simply accounted for by general disruption of cell division patterns at this

later stage (Figure S3), since similar disruptions can be caused by *Foxf>dnFGFR*, which

inhibits *Ebf* expression (Figure 1C). These results indicated that forced M-Ras activity is

sufficient to upregulate STVC and ASMF markers ectopically. This is consistent with the

idea that spatially defined signaling upstream of M-Ras restricts MAPK activity, thus

localizing STVC- and ASM-specific gene activities.

To further probe the signal transduction pathway, we engineered a constitutively

active version of the Ciona Mek1/2 protein by introducing phosphomimetic mutations

of two conserved serine residues in the catalytic domain, as previously shown for the

mammalian homolog (Cowley et al., 1994; Mansour et al., 1994). Early misexpression of

Mek^{S220E,S216D} in the B7.5 lineage using the *Mesp* enhancer caused ectopic TVC

induction, mimicking the effects of gain of Ets1/2 function (Figure S4; (Davidson et al.,

2006)). As seen with M-Ras^{G22V}, TVC-specific misexpression of Mek^{S220E,S216D} using the

Foxf enhancer also caused ectopic expression of *Htr7* and *Tbx1/10*, and *Ebf* in 15 and

18hpf larvae, respectively (Figure 2B, C). Taken together, these results indicate that

activity of the canonical FGF-Ras-MEK-ERK pathway is progressively restricted to the

STVC and ASMF, and is both necessary and sufficient to promote STVC- and ASMF-

specific gene expressions.

Continuous FGF-MAPK activity is required for the successive activations of

***Tbx1/10* and *Ebf*.**

FGF-MAPK signaling is sufficient and necessary to maintain *Hand-r* expression in

late TVCs (Figure 1), and *Hand-r* is necessary for *Ebf* expression in the ASMF (Razy-

Krajka et al., 2014). Therefore, it is possible that later FGF-MAPK signaling is

dispensable for *Tbx1/10* and *Ebf* activation and ASM specification, as long as STVC and

ASMF cells inherit sustained levels of *Hand-r* mRNAs and/or proteins. To disentangle

late from early requirements of FGF-MAPK signaling, we incubated embryos at

different stages with the MEK/MAPKK inhibitor U0126, which abolishes dual ERK

phosphorylation and the initial MAPK-dependent TVC induction in Ciona embryos

(Figure S1; (Davidson et al., 2006; Hudson et al., 2003)). MEK inhibition during TVC

migration (i.e. between 9.5 and 12.5 hpf, Figure 3A) blocked the expression of *Hand-r*

and *Htr7* in late TVCs (Figure 3B, E). U0126 treatments in late TVCs, and through the

first asymmetric division (i.e. between 12 and 15 hpf, Figure 3A) did not alter TVC

division patterns (Figure S3C), but it blocked both the maintenance of *Hand-r* and

Htr7, and the activation of *Tbx1/10* in the STVCs (Figure 3C, D, F, G). Finally, MEK

inhibition in late STVCs and through asymmetric divisions (i.e. between 15 and 18 hpf)

also did not alter STVC divisions (Figure S3G), but it blocked the ASMF-specific

expression of *Ebf* (Figure 3H). These results indicate that continuous MEK activity is

required throughout cardiopharyngeal development to successively activate TVC-,
STVC-, and ASMF-expressed genes.

Since *Ebf* expression is maintained for several days in the ASMF derivatives as they
differentiate into body wall and siphon muscles (Razy-Krajka et al., 2014), we tested
whether continued MEK activity is also required for the maintenance of *Ebf* expression
past its onset and cells' commitment to an ASM fate. Using both regular and intron-
specific antisense probes, which specifically detect nascent transcripts (Wang et al.,
2013), we showed that later MEK inhibition (i.e. U0126 incubation between 17 and 20
hpf) did not block the maintenance of *Ebf* transcription in the ASMPs (Figure 3I, J).
This indicates that sustained MEK activity is required until the onset of *Ebf* expression,
but not beyond, and the maintenance of *Ebf* expression during ASM development is
independent of MAPK.

Since U0126 treatments affect the whole embryo, we sought to further confirm the
later roles for FGF-MAPK signaling specifically in the cardiopharyngeal mesoderm. To
this aim, we used an STVC-specific enhancer from the *Tbx1/10* locus (termed *T6*; Figure
3K, L; Figure S5; (Tolkin and Christiaen, 2016); Racioppi et al., in preparation) to drive
expression of either dnFGFR or the constitutively active M-Ras^{G22V} starting at ~14hpf,
and assayed *Ebf* expression at 18hpf (Figure 3K, L). These perturbations minimally
affected the cell division patterns (Figure S3), such that cells corresponding to FHP,
SHP and ASMF could be identified by their position relative to the midline in many
embryos (Figure 3K). M-Ras^{G22V} misexpression caused conspicuous ectopic *Ebf*
expression in the SHPs, whereas dnFGFR-mediated inhibition of MAPK activity blocked
Ebf activation in the lateral ASMFs. These results support the notion that localized FGF-
MAPK activity is necessary and sufficient for ASMF-specific expression of *Ebf*.

Coherent feed-forward circuits for cardiopharyngeal mesoderm patterning and ASM fate specification.

The above results indicate that *Hand-r*, *Tbx1/10* and *Ebf* require ongoing FGF-MAPK activity for their successive activations in the TVCs, STVCs and ASMFs, respectively. We previously showed that RNAi and/or CRISPR-mediated inhibition of either *Hand-r* or *Tbx1/10* function blocks *Ebf* activation in the ASMFs, where both *Hand-r* and *Tbx1/10* expressions are maintained (Razy-Krajka et al., 2014; Tolkin and Christiaen, 2016; Wang et al., 2013). We used epistasis assays to systematically test whether early regulators mediate the effects of FGF-MAPK on later gene expression and ASM fate specification, or whether FGF-MAPK signaling acts both upstream and in parallel to early regulators in a more complex regulatory circuit.

We first revisited the regulatory relationships between FGF-MAPK, *Hand-r* and *Tbx1/10* in late TVCs and early STVCs. We validated single guide RNAs (sgRNAs) for CRISPR/Cas9-mediated mutagenesis of *Hand-r* (Table S1; (Gandhi et al., 2017a)), and determined that *Hand-r* function is necessary for *Tbx1/10* activation in the STVCs (Figure 4A). Co-expression of a CRISPR-resistant *Hand-r* cDNA (Hand-r^{PAMmis}) rescued *Tbx1/10* expression in the STVCs, indicating that *Tbx1/10* down-regulation in this CRISPR "background" is specifically due to *Hand-r* loss-of-function (Figure 4A). To further probe if *Hand-r* activity is necessary for FGF-MAPK-dependent *Tbx1/10* expression, we used gain of M-Ras function in a *Hand-r* CRISPR "background". Whereas, misexpression of the constitutively active M-Ras^{G22V} caused ectopic *Tbx1/10* expression, concomitant loss of *Hand-r* function diminished both endogenous and ectopic *Tbx1/10* expression in the STVC and FHP, respectively (Figure 4A). Although, remaining ectopic activation could still be observed, possibly because M-Ras^{G22V} could boost *Hand-r* expression in heterozygous cells where CRISPR/Cas9 disrupted only one

copy of the gene. This data indicate that Hand-r function is necessary for FGF-MAPK-

induced activation of *Tbx1/10*.

To further probe the epistatic relationships between *Hand-r* and MAPK signaling

upstream of *Tbx1/10*, we attempted to rescue *Tbx1/10* expression in U0126-treated

embryos, by over-expressing *Hand-r* with the TVC-specific *Foxf* enhancer. Neither did

Hand-r over-expression cause ectopic *Tbx1/10* activation (in the FHPs), nor was it

sufficient to rescue *Tbx1/10* expression in 15hpf STVCs (Figure 4B). Taken together,

these data indicate that both Hand-r and MAPK activities are required to activate

Tbx1/10 in the STVCs. These results also imply that MAPK signaling is restricted to the

STVC independently of Hand-r activity, which suffice to explain the STVC-specific

activation of *Tbx1/10*. Finally, we isolated a minimal STVC-specific enhancer from the

Tbx1/10 locus and identified conserved putative Ets binding sites, which were necessary

for reporter gene expression (Figure S5). This suggests that the FGF-MAPK-Ets

pathway directly regulates *Tbx1/10* expression in the STVCs.

Next, we investigated the epistatic relationship between FGF-MAPK, *Hand-r*, and

Tbx1/10 upstream of *Ebf* in the ASMFs. We first used previously validated

CRIPSR/Cas9 reagents targeting the *Tbx1/10* coding region (Tolkin and Christiaen,

2016), to confirm that B7.5-lineage-specific loss of *Tbx1/10* function inhibited *Ebf*

activation, and verified that this effect could be rescued by over-expression of a

CRISPR/Cas9-resistant *Tbx1/10* cDNA, expressed with a minimal TVC-specific *Foxf*

enhancer (Figure 4C; Tbx1/10^PAMmis^). In these rescue experiments, we observed ectopic

Ebf activation in the SHP, as previously described when driving Tbx1/10 expression

with a TVC-specific *Foxf* enhancer (Wang et al., 2013). As explained below, this ectopic

activation could be attributed to a precocious expression of *Ebf* in the STVCs (Figure

4E). To test whether *Tbx1/10* was also required for ectopic *Ebf* expression in response to

MAPK activation (see Figure 2), we combined CRISPR/Cas9-mediated *Tbx1/10*

knockout with constitutive MAPK activation using the M-Ras^{G22V} mutant and observed

a significant inhibition of both endogenous and ectopic *Ebf* expression in the 18hpf

ASMF and SHP, respectively (Figure 4C). Taken together, these results show that

Tbx1/10 function is necessary for FGF-MAPK-induced expression of *Ebf* in the ASMFs.

To further test whether *Tbx1/10* acts in parallel and/or downstream of MAPK to

activate *Ebf*, we combined gain of *Tbx1/10* function with perturbations of FGF-MAPK

signaling and assayed *Ebf* expression. We realized that *Foxf*-driven misexpression of

Tbx1/10 caused precocious *Ebf* activation in 15hpf STVCs (Figure 4D, E). This

precocious expression remained remarkably patterned, suggesting that STVC-restricted

FGF-MAPK activity prevented *Ebf* expression in the dpERK-negative, small median

FHPs (Figures 1B, 4E, S1). Indeed, co-expression of Tbx1/10 and M-Ras^{G22V} caused both

precocious and ectopic *Ebf* expression in the 15hpf medial and lateral TVC derivatives,

which would be FHPs and STVCs in control embryos, respectively. This data confirms

that Tbx1/10 misexpression does not suffice to cause ectopic *Ebf* expression in the

FHPs, because the latter presumably lack FGF-MAPK activity, as is the case in control

embryos.

U0126-mediated MEK inhibition from 12 to 15hpf, i.e. after the onset of

Foxf>Tbx1/10 misexpression, further confirmed that MAPK activity is required in

parallel to Tbx1/10 for precocious *Ebf* activation in 15hpf STVCs (Figure 4D, E). Taken

together, these results indicate that Tbx1/10 and MAPK are both required to activate

Ebf in the cell cycle following that of *Tbx1/10* onset.

Since *Hand-r* expression is maintained in the ASMF, and CRISPR/Cas9- or RNAi-

mediated *Hand-r* knockdown blocked both *Tbx1/10* (Figure 4A) and *Ebf* expression

([Razy-Krajka et al., 2014](#)), we reasoned that Hand-r could also act both upstream and in

parallel to Tbx1/10 for *Ebf* activation. To test this possibility, we assayed *Ebf* expression

in 18hpf ASMF following defined perturbations of Hand-r and Tbx1/10. As expected,

CRISPR/Cas9-mediated *Hand-r* mutagenesis strongly inhibited *Ebf* expression, and

this effect could be rescued by a CRISPR-resistant *Hand-r* cDNA (Figure 4F). To test

whether this effect was mediated by a loss of *Tbx1/10* expression, we attempted to

rescue the Hand-r loss-of-function by over-expressing Tbx1/10 using the *Foxf* enhancer.

As explained above, *Foxf*-mediated Tbx1/10 misexpression caused precocious and

ectopic *Ebf* expression in larvae co-electroporated with control sgRNAs (Figure 4D, E,

F). By contrast, combining loss of Hand-r function with Tbx1/10 misexpression

inhibited both the endogenous and ectopic *Ebf* expression (Figure 4F), indicating that

Hand-r is also required in parallel to Tbx1/10 for *Ebf* activation in the ASMFs.

Taken together, these analyses suggest that coherent feed-forward circuits govern

the sequential activation of *Hand-r*, *Tbx1/10* and *Ebf* in response to continuous but

progressively restricted FGF-MAPK inputs (Figure 4G), thus linking spatial patterning

to the temporal deployment of the regulatory cascade leading to localized *Ebf* activation

and pharyngeal muscle specification.

The cell cycle entrains the temporal deployment of the cardiopharyngeal

gene regulatory network.

In principle, the feed-forward circuit described above is sufficient to explain the

successive activations of *Hand-r*, *Tbx1/10* and *Ebf*. However, *Tbx1/10* and *Ebf* do not

turn on until after TVC and STVC divisions, respectively. Notably, even when we

misexpressed Tbx1/10 in the TVCs, *Ebf* was activated only after cell division and in the

lateral-most cells, where FGF-MAPK signaling is normally maintained (Figures 1B, 4E).

This sequence of events -divisions followed by gene activation- is paramount as it

permits the birth of first and second heart precursors, whose fates are antagonized by

Tbx1/10 and Ebf (Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et al., 2013).

Therefore, we sought to investigate the role(s) of the cell cycle in controlling the timing

of *Tbx1/10* and *Ebf* activation.

We first evaluated the effects of cytochalasin B, a classic inhibitor of cytokinesis

widely used to study cell fate specification in ascidians (Figure 5A; (Whittaker, 1973)).

Treatments starting before TVC divisions (12 hpf) did not block *Tbx1/10* or *Ebf*

expression in embryos fixed after their normal onset at either 16 or 19hpf, respectively

(Figure 5B). Similarly, treatment starting between the first and second asymmetric

divisions (15hpf) did not block localized *Ebf* expression at 19hpf (Figure 5B). This

indicates that *Tbx1/10* and *Ebf* activations occur by default in the absence of

cytokinesis, most likely because FGF-MAPK signaling persists throughout the shared

cytoplasm. This data thus illustrates how the spatial restriction of FGF-MAPK signaling,

following cell divisions, leads to the localized activations of *Tbx1/10* and *Ebf*.

Cytochalasin treatments usually lead to the formation of polynucleated cells (e.g.

Figure 5B, middle panel), because the cell cycle and nucleokinesis continue. To alter cell

cycle progression more comprehensively, and specifically in the cardiopharyngeal

lineage, we used genetically encoded inhibitors of cell cycle transitions: Cdkn1b.a and

Cdkn1b.b (also known as Noto16), the ortholog of which is a potent inhibitor of the G1/S

transition in the ascidian species *Halocynthia roretzi* (Kuwajima et al., 2014), and the

G2/M inhibitor Wee1 (Dumollard et al., 2017). We used the TVC-specific *Foxf* enhancer

to misexpress these negative regulators of cell cycle progression, monitored cell

divisions and assayed *Tbx1/10* expression at 15hpf, when control TVCs have divided and

the lateral-most STVCs normally express *Tbx1/10*. Each perturbation efficiently

inhibited TVC divisions, such that only two cells were visible on either side of the

embryos (Figure 5D). In these delayed TVCs, *Tbx1/10* expression was strongly reduced

compared to control STVCs (Figure 5D). However, 20 to 40% of the delayed TVCs

expressed *Tbx1/10* to variable extents. This suggests that the cardiopharyngeal regulatory network can qualitatively unfold independently of cell cycle progression, but the latter is necessary for *Tbx1/10* expression to its wild-type levels.

We next used the STVC-specific *Tbx1/10 T6* enhancer (Figure S5), to misexpress Cdkn1b.a, Noto16 and Wee1, and assayed *Ebf* expression at later stages. Inhibitors of the G1/S transition failed to block STVC divisions (data not shown), most likely because *Tbx1/10(T6)*-driven products did not accumulate quickly enough to interfere with the G1/S transition in STVCs, since this cell cycle lasts only ~2 hours compared to ~6 hours for the TVC interphase. Therefore, we focused the analyses of *Ebf* response to cell cycle perturbations on misexpression of the G2/M inhibitor Wee1. Analyses of 18hpf larvae, fixed approximately 2 hours after the documented onset of *Ebf* expression in ASMFs (Razy-Krajka et al., 2014), indicated that *Ebf* can turn on in arrested STVCs that failed to divide upon Wee1 misexpression (Figure 5E).

Because ~30% of the embryos showed variable expression, as was the case for *Tbx1/10* in the previous experiment, we reasoned that perturbations of the G2/M transition could alter the dynamics of *Ebf* upregulation. We investigated this possibility using embryos fixed every 30 minutes between 15.5hpf and 18hpf, when cells transition from a late *Tbx1/10+*; *Ebf-* STVC state to a committed *Ebf+*, *Mrf+* ASMF state (Razy-Krajka et al., 2014; Wang et al., 2013). First, we scored the proportion of embryos with delayed STVCs or conspicuous ASMFs at each time point and showed that Wee1 misexpression strongly delays cell cycle progression, blocking cell divisions in a substantial fraction of embryos (Figure S6).

The proportion of *Ebf+* ASMFs in control embryos progressively increased from ~20% at 15.5hpf to >90% by 18hpf, revealing an under-appreciated dynamic at the onset of *Ebf* expression, which appears to take at least one hour to be "strongly" expressed in >75% of newborn ASMFs (Figure S6).

To evaluate the impact of Wee1-induced mitosis inhibition on *Ebf* accumulation, we

focused on undivided STVCs at each time point (hence the lower numbers in Figure S6A

compare to Figure S6B). By 17hpf, wee1-expressing delayed STVCs showed "strong" *Ebf*

expression in comparably high proportions of embryos. However, these proportions

were significantly lower at 16 and 16.5hpf (Chi-square tests, *P*=0.002 and *P*=0.0003,

respectively), with ~1.5 and ~1.2 times less "strongly" expressing cells than in the

control distributions (hypergeometric tests, *P*=0.0005 and *P*=0.0001, respectively).

These semi-quantitative analyses suggest that the cardiopharyngeal network can

eventually unfold independently of cell divisions, leading to high levels of *Ebf*

expression, albeit with a delay. This suggests that STVC division entrains *Ebf*

upregulation in early ASMFs.

Finally, we reasoned that Wee1-expressed delayed STVC that activate *Ebf* would not

turn on heart markers (Wang et al., 2017). Indeed, delayed STVCs failed to activate the

pan-cardiac *Lrp4/8*, and the SHP-markers *Kirr* and *Dach* (Figure 5F; (Wang et al.,

2017). This indicated that coupling STVC division with localized FGF-MAPK activity

and timed *Ebf* upregulation permits the localized activation of *Ebf* and the emergence of

cardiac progenitors.

424   Transition from a MAPK-dependent to a MAPK-independent and

425   autoregulative mode of *Ebf* expression in early ASMFs.

We sought to further probe the mechanisms that regulate the initiation of *Ebf*

expression in early ASMFs, and their biological significance for fate specification. Since

we observed a progressive accumulation of *Ebf* mRNAs, and a transition from a MAPK-

dependent onset to a MAPK-independent maintenance of *Ebf* transcription (Figure 3I,

J), we reasoned that the window of MAPK-dependence might coincide with the

accumulation of *Ebf* mRNAs between 16 and 17hpf. To test this possibility, we treated

embryos with the MEK inhibitor U0126 at successive time points, assayed ongoing

transcription using intronic probes and counted the numbers of *Ebf* transcribing cells

(Figure 6A). This analysis revealed that *Ebf* transcription gradually lost its sensitivity to

MAPK inhibition between 16 and 17hpf, i.e. during the first hour of the ASMF cycle

when *Ebf* mRNAs normally accumulate (Figure S6A,B).

Because *Ebf* transcription becomes independent from MAPK by the time *Ebf* mRNA

have accumulated to "high" levels, and because *Ebf* expression lasts for several days in

the progeny of the ASMFs, we reasoned that autoregulation might suffice to maintain

high levels of *Ebf* mRNA past the MAPK-dependent onset. To test this possibility, we

misexpressed the Ebf coding sequence using the STVC-specific *T6* enhancer as

described (Tolkin and Christiaen, 2016). Assaying endogenous *Ebf* transcription with

intronic probes demonstrated that, in addition to its normal expression in the ASMFs,

Ebf misexpression caused precocious and ectopic activation of the endogenous locus in

the STVCs, and in the MAPK-negative SHPs, respectively (Figure 6C-F). This result

suggests that *Ebf* transcription bypasses both requirements for cell-division coupling

and MAPK inputs if high levels of *Ebf* gene products are present in the cell.

We reasoned that, if high levels of *Ebf* expression can promote its own transcription

independently of MAPK signaling, then Ebf misexpression should be sufficient to rescue

a chemical inhibition of MAPK at a critical stage. We tested this possibility by

combining Ebf misexpression using the STVC-specific *T6* enhancer and U0126

treatments starting at 16hpf, which normally block *Ebf* expression (Figure 6A, D-F). We

observed that transcription of the endogenous *Ebf* locus became independent of early

MAPK activity upon misexpression of an Ebf cDNA, further supporting the notion that

high levels of *Ebf* expression suffice to maintain *Ebf* transcription independently of

MAPK activity.

A potentially important implication of this transient MAPK requirement is to render

Ebf expression initially reversible. For instance, *Ebf* occasionally turns on precociously

in the STVCs of a small proportion of embryos (Figure S6C). Given the powerful anti-

cardiogenic effects of Ebf (Razy-Krajka et al., 2014; Stolfi et al., 2010), persistent *Ebf*

expression would have dramatic consequences for SHP development (Wang et al.,

2013). However, because MAPK activity is excluded from the SHPs, and the early phase

of *Ebf* expression depends upon continuous MAPK activity, we surmise that *Ebf* cannot

be maintained in the SHPs. For instance, when embryos from the same electoporated

batch were fixed at the time of early U0126 treatment (i.e. 15.75 and 16.25hpf) and ~4

hours later, at 20hpf, and assayed for *Ebf* transcription using intronic probes, initially

wild-type patterns of *Ebf* transcription could not be maintained (Figure S7A). This

suggests that, although *Ebf* can be activated precociously in a MAPK-dependent

manner, its expression shuts off in the SHPs upon MAPK inhibition following STVC

division.

We further addressed the interplay between cell division, MAPK signaling and *Ebf*

expression. We reasoned that, if cell divisions entrain Ebf accumulation and the

transition to a MAPK-independent autoregulative mode, then delaying STVC divisions

should extend the period of MAPK-dependent *Ebf* transcription. We tested this

possibility by expressing Wee1 under the control of the STVC-specific *T6* enhancer, and

treated embryos with U0126 at 17hpf, which inhibited the maintenance of *Ebf*

transcription in only 15% to 20% of the control embryos (Figures 6A, S7B). The

proportion of embryos showing U0126-sensitive *Ebf* transcription increased to almost

50% upon *T12>Wee1* expression (Figure S7B), which is consistent with our hypothesis

that inhibiting the G2/M transition delays the accumulation of *Ebf* gene products, thus

postponing the transition from a low level/MAPK-dependent to an high level/MAPK-

independent and self-activating mode of *Ebf* regulation.

We propose a model for *Ebf* regulation whereby Hand-r, Tbx1/10, ongoing MAPK

signaling and cell-cycle-regulated transcriptional input(s) govern the onset and initial

accumulation of *Ebf* gene products during the first hour of the ASMF cycle, whereas the

maintenance of *Ebf* expression relies primarily on MAPK-independent autoactivation,

following initial accumulation (Figure 7).

Discussion

Here, we demonstrated that the progressive restriction of FGF-MAPK signaling

follows oriented and asymmetric cell divisions of multipotent progenitors and patterns

the ascidian cardiopharyngeal mesoderm in space and time. Dynamic FGF-MAPK

activity patterns lead to the localized expression of *Hand-r*, *Tbx1/10* and *Ebf* in fate-

restricted pharyngeal muscle precursors, and their concomitant exclusion for first and

second heart precursors. We show that coherent feedforward circuits encode the

successive activations of *Hand-r*, *Tbx1/10* and *Ebf*, whereas cell divisions entrain the

progression of this regulatory sequence and thus define the timing of gene expression.

Finally, we provide evidence that the initiation of *Ebf* expression depends on MAPK

activity in early ASMF, until Ebf accumulation permits MAPK-independent auto-

activation. Given the potent anti-cardiogenic, and pro-pharyngeal muscle effects of Ebf

(Razy-Krajka et al., 2014; Stolfi et al., 2010), we surmise that the latter switch

corresponds to the transition from a cardiopharyngeal multipotent state to a committed

pharyngeal muscle identity.

Spatial patterning by localized maintenance of FGF-MAPK signaling.

Our results demonstrate that MAPK signaling is maintained only in the lateral-most

daughter cells following each asymmetric division of multipotent cardiopharyngeal

progenitors - the TVCs and STVCs. This asymmetric maintenance is necessary and

sufficient for the progressive and localized deployment of the pharyngeal muscle

network. Notably, the TVCs themselves are initially induced by similar polarized FGF-

MAPK signaling coincidental to asymmetric cell divisions of their mother cells, the B8.9

and B8.10 founder cells (Davidson et al., 2006). At this stage, asymmetrical

maintenance of sustained FGF-MAPK signaling involves intrinsic Cdc42-dependent

polarity of the founder cells, which promotes polarized cell-matrix adhesion of the

prospective TVC membrane to the ventral epidermis. The latter differential integrin-

mediated adhesion promotes localized MAPK activation, leading to TVC induction

(Cooley et al., 2011; Norton et al., 2013). It has been proposed that adhesion- and

caveolin-dependent polarized FGFR recycling during mitosis accounts for the localized

activation of MAPK in the prospective TVCs (Cota and Davidson, 2015). Whereas

similar mechanisms could in principle account for asymmetric maintenance of FGF-

MAPK signaling in STVCs and ASMFs, this has not been formally tested and there are

notable differences opening the possibility that other mechanisms may be at work:

during TVC induction, MAPK signaling is maintained in the smaller daughter cell that

contacts the epidermis, whereas in the following divisions, MAPK activity persists in the

larger daughter cells and all cells maintain contact with the epidermis (Kaplan et al., in

preparation). Moreover, using fusion proteins as in previous studies, we could not

observed a marked polarized distribution of FGFR molecules to the lateral-most cells

(the STVCs and ASMFs; Kaplan et al., in preparation). However, the fact that

constitutively active forms of M-Ras and Mek1/2 were sufficient to bypass the loss of

MAPK activity, and impose pharyngeal muscle specification, indicates that differential

FGF-MAPK activity is regulated upstream of M-Ras. Further work is needed to

elucidate the cellular and molecular mechanisms governing the spatiotemporal patterns

of FGF-MAPK signaling in the cardiopharyngeal mesoderm. In particular, it will be

important to disentangle the relative impacts of extrinsic (i.e. tissues, contacts) vs.

intrinsic (i.e. asymmetric cell division) effects onto FGF-MAPK signaling and the

downstream transcriptional inputs.

Our preliminary analyses indicate that perturbations of the FGF-Ras-MEK pathway

can alter cardiopharyngeal cell division patterns (Figure S3). While these effects did not

account for the observed changes in gene expression, future studies will unravel FGF-

MAPK-dependent and cardiopharyngeal-specific mechanisms governing the orientation

and asymmetry of progenitor cell division (Kaplan et al., in preparation).

Transcriptional consequences of differential FGF-MAPK signaling.

Differential FGF-MAPK signaling rapidly impacts cell-specific gene expression, we

thus surmise that transcriptional effectors are dynamically regulated. Even though we

have not formally identified the downstream DNA-binding transcription factor (see

discussion below), it is conceivable that the phosphorylated forms of either

transcriptional effector could persist through cell division upon maintenance of FGF-

MAPK activity. However, we have shown that continuous MAPK activity is needed

following each division. Therefore, we must invoke elusive phosphatase activities, such

as dual-specificity phosphatases (DUSPs; (Patterson et al., 2009), which would reset

transcriptional effectors to a dephosphorylated state, thus rendering steady-state FGF-

Ras-MAPK inputs necessary.

Systematic dephosphorylation of FGF-MAPK transcriptional effectors is particularly

important for heart fate specification. As shown in our companion paper (Wang et al.,

2017), whole genome analyses indicate that heart-specific *de novo* gene expression

requires MAPK inhibition (Wang et al., 2017). The molecular mechanisms remain

elusive, but one simple possibility is that, lest fate-restricted heart precursors inhibit

MAPK activity, they will activate *Tbx1/10* and *Ebf*, which will block the cardiac program

(Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et al., 2013).

Finally, we previously proposed that repressor inputs from Nk4/Nkx2-5 are needed

in the second heart precursors to avoid ectopic activation of *Ebf* (Wang et al., 2013). The

observation that *Nk4* transcripts are detected in all cardiopharyngeal cells opened the

question as to how *Ebf* would escape repression by Nk4 in the ASMFs. Differential

MAPK activity offers an intriguing possibility: for instance, Nk4/Nkx2.5-mediated

repression in other species involves the co-repressor Groucho/TLE (Choi et al., 1999),

which is strongly expressed in the cardiopharyngeal mesoderm (Razy-Krajka et al.,

2014); and, in flies, MAPK-mediated phosphorylation of Groucho inhibits its repressor

function (Cinnamon et al., 2008; Cinnamon and Paroush, 2008; Hasson et al., 2005).

Therefore, it is possible that persistent MAPK signaling dampens Groucho/TLE-

mediated repressive inputs on cell-specific regulatory genes like *Ebf*. Future studies will

determine whether such mechanisms provide bistable switches underlying MAPK-

dependent fate choices in the cardiopharyngeal mesoderm.

Temporal deployment of the pharyngeal muscle network

The localized and successive activation of *Tbx1/10* and *Ebf* in STVCs, and ASMFs,

respectively, are important features of the cardiopharyngeal network that permit the

emergence of diverse cell fates: first and second heart precursors, and atrial siphon

muscle   precursors.   Experimental   misexpression   of   Ebf   throughout   the

cardiopharyngeal mesoderm suffice to inhibit heart development (Razy-Krajka et al.,

2014; Stolfi et al., 2010), illustrating the importance of *Ebf* restriction to the ASMF for

the emergence of first and second heart precursors.

Our analyses indicate that the sequential activations of *Hand-r*, *Tbx1/10* and *Ebf* is

encoded in the feed-forward structure of this sub-circuit, whereas the continuous

requirement for MAPK inputs and their progressive exclusion from heart progenitors

restrict the competence to activate *Tbx1/10* and *Ebf* to the most lateral cells, after each

division. Our model implies that each gene may directly respond to transcriptional

inputs from MAPK signaling. We have not formally identified the transcription

factors(s) that mediate the transcriptional response to FGF-MAPK signaling. However,

multipotent cardiopharyngeal progenitors express *Ets1/2* and *Elk*, two common

transcriptional effectors of FGF/MAPK signaling in *Ciona* (Bertrand et al., 2003;

Christiaen et al., 2008; Davidson et al., 2006; Gainous et al., 2015), and Ets1/2 has been

implicated in the initial FGF-MAPK-dependent induction of multipotent TVCs

(Christiaen et al., 2008; Davidson et al., 2006), and its expression is also progressively

restricted to the lateral-most progenitors following each division (Razy-Krajka et al.,

2014). Taken together, Ets1/2 and, to some extend, Elk are intriguing candidate

transcriptional effectors of FGF/MAPK signaling in cardiopharyngeal development.

The binding preferences of Ets-family factors have been extensively studied in

Ciona, and they do not depart markedly from conserved Ets-family binding sites with a

GGAW core (Bertrand et al., 2003; Farley et al., 2015; Farley et al., 2016; Gueroult-

Bellone et al., 2017; Khoueiry et al., 2010). Putative Ets-family binding sites in the TVC-

specific *Hand-r* enhancer are conserved between *Ciona intestinalis* and its sibling

species *C. robusta* and *C. savignyi*, and necessary for its activity in reporter assays

(Woznica et al., 2012). Similarly, minimal STVC and ASM enhancers for *Tbx1/10* and

Ebf, respectively, contain conserved putative Ets-family binding sites, which contribute

to proper reporter gene expression in transfection assays (Figure S5; (Razy-Krajka et al.,

2014; Wang et al., 2013) and data not shown). Taken together, these observations

suggest that the proposed feed-forward sub-circuit involves direct transcriptional inputs

from FGF-MAPK-regulated Ets-family factors on the cardiopharyngeal enhancers of

Hand-r, *Tbx1/10* and *Ebf*.

Whereas the regulatory architecture of the MAPK; Hand-r; Tbx1/10; Ebf sub-circuit

explains the sequence of activation events, it is also crucial for its correct deployment,

and the generation of diverse cell identities, that genes are not fully activated before

successive cell divisions. While divisions are not absolutely required for *Ebf* to

eventually turn on, cell cycle progression appears to entrain the deployment of this

network, especially for *Tbx1/10* and *Ebf* activation in STVCs and ASMFs, respectively.

These observations imply that the intrinsic dynamic of the network is slower than observed. This allows first and second heart precursors to be born prior to the onset of *Tbx1/10* and *Ebf*, respectively. The latter sequence is essentially for the heart progenitors to escape the anti-cardiogenic effects of Tbx1/10 (Wang et al., 2013), and Ebf (Razy-Krajka et al., 2014).

Initial *Ebf* expression in early ASMFs is also labile and MAPK-dependent for approximately one hour. This continued requirement for MAPK inputs ensures that, in the rare instances when *Ebf* expression starts in the multipotent STVC progenitors and/or expands to the nascent SHPs, inhibition of MAPK shuts off *Ebf* expression before it reaches the levels needed for commitment to an ASM fate. Indeed, our results indicate that, once *Ebf* mRNAs have accumulated to high levels, its expression becomes auto-regulative and MAPK-independent. We surmise that this transition coincides with a fundamental switch from a multipotent cardiopharyngeal state to a committed pharyngeal muscle identity.

From this standpoint, the observed entrainment of *Ebf* expression by the cell cycle can be seen as acceleration of the transition to commitment following asymmetric division of multipotent progenitors. Although the mechanisms remain elusive, it is likely that this requires the M/G1 transition, as the G1 phase has been shown to be particularly conducive to the expression of fate-specific regulators in mammalian pluripotent stem cells (Dalton, 2015; Pauklin et al., 2016; Pauklin and Vallier, 2013; Soufi and Dalton, 2016).

Conserved dual effects of FGF-MAPK signaling on heart development in chordates

Previous studies highlighted how FGF-MAPK signaling is necessary along side Mesp during early cardiac development in *Ciona* (Christiaen et al., 2008; Davidson, 2007;

Davidson et al., 2006), and how this early requirement also exists in vertebrates (Abu-
Issa et al., 2002; Alsan and Schultheiss, 2002; Barron et al., 2000; Brand, 2003; Reifers
et al., 2000; Zaffran and Frasch, 2002). We now know that these early FGF-MAPK
inputs induce and maintain multipotent cardiopharyngeal states in *Ciona*, including the
Tbx1/10+ multipotent progenitors that eventually produce the second heart lineage
((Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et al., 2013), (Wang et al., 2017) and
this study). Similarly, in vertebrates, regulatory interplay between Fgf8 and Fgf10
signaling and Tbx1 is required for development of both pharyngeal arch and second
heart field derivatives, presumably in part by maintaining an undifferentiated and
proliferative state (Abu-Issa et al., 2002; Aggarwal et al., 2006; Brown et al., 2004;
Chen et al., 2009; Hu et al., 2004; Ilagan et al., 2006; Kelly and Papaioannou, 2007;
Park et al., 2006; Park et al., 2008; Vitelli et al., 2002b; Watanabe et al., 2010;
Watanabe et al., 2012). In fish, FGF signaling is necessary for cardiomyocyte addition at
the arterial pole, in a manner reminiscent of its role in second heart field development
(de Pater et al., 2009; Lazic and Scott, 2011). Notably, FGF signaling acts in successive
phases, and its inhibition is necessary for final myocardial specification and
differentiation (Hutson et al., 2010; Marques et al., 2008; Tirosh-Finkel et al., 2010;
van Wijk et al., 2009). Conversely, continued FGF signaling beyond the multipotent
mesodermal progenitor stages was shown to promote smooth muscle and epicardial
differential in the heart (Hutson et al., 2010; van Wijk et al., 2009), and also myoblast
specification and/or skeletal muscle differentiation in the head, with the expression of
FGF ligands being maintained in the pharyngeal arches (Bothe et al., 2011; Buckingham
and Vincent, 2009; Michailovici et al., 2015; Michailovici et al., 2014; von Scheven et
al., 2006). Taken together, these and our data suggest that FGF-MAPK signaling plays
evolutionary conserved roles during chordate cardiopharyngeal development, by
promoting the specification of successive mesodermal and *Tbx1+* multipotent states,

and a fate-restricted non-cardiac muscle identity, while MAPK inhibition is required for

myocardial specification and differentiation in the first and second heart field,

successively.

Material and methods

Animals, electroporations, and chemical treatments

Gravid wild *Ciona intestinalis* type A, now called *Ciona robusta* ([Pennati et al., 2015](#)),

were obtained M-REP (Carlsbad, CA, USA), and kept under constant light to avoid

spawning. Gametes from several animals were collected separately for *in vitro* cross-

fertilization followed by dechorionation and electroporation as previously described

([Christiaen et al., 2009a](#), [b](#)). Different quantities of plasmids were electroporated

depending on the constructs. Typically, 50 µg of DNA was electroporated for NLS::lacZ

or plain mCherry driving constructs but only 15 µg for *Mesp-1>H2B::mCherry*. For

perturbation constructs, 70 µg were usually electroporated, except for

Mesp>NLS::Cas9::NLS (30 µg) and pairs of U6>sgRNA plasmids (25 µg each). U0126

(Cell Signaling Technology, Danvers, MA) was used at 5µM in artificial seawater from a

stock solution of 20mM in DMSO. Cytochalasin B (Sigma, Saint Louis, MO) was used at

~3 µg/mL from a 10 mg/mL stock solution in DMSO as previously performed ([Jeffery et](#)

[al., 2008](#)). Control embryos were incubated in parallel with corresponding

concentrations of DMSO alone.

In situ **hybridization**

In situ hybridizations were carried out essentially as described previously ([Christiaen et](#)

[al., 2009c](#); [Razy-Krajka et al., 2014](#)), using DIG labeled riboprobes, anti-DIG-POD Fab

fragments (Roche, Indianapolis, IN), and Tyramide Amplification Signal coupled to

Fluorescein (Perkin Elmer, MA). Reporters expressed in the lineage of interest were

marked using anti-ß-galactosidase monoclonal mouse antibody (1:1000; Promega,

Fitchburg, WI) or anti-mCherry rabbit polyclonal antibody (1:500; BioVision 5993-

100), respectively targeted with anti-mouse or anti-rabbit secondary antibody coupled

with Alexa 648 (1:500; Invitrogen, Carlsbad, CA). The different probes used in this

study were described previously (Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et

al., 2013).

dpERK/mcherry double fluorescent immunostaining

Samples were fixed, as for *in situ* hybridizations, in MEM-PFA with Tween 20 (0.05%)

but only for 30 minutes at room temperature, washed three times in PBSt (Tween 20

0.01%) for 10 minutes, gradually dehydrated every 10 minutes in Ethanol/PBS series

(33%, 50%, 80%) and Methanol 100%. Samples were then gradually rehydrated every

10 minutes in Methanol/PBSt series, rinsed three times in PBSt, permeabilized with

PBS Triton-100 (0.2%) for 30 minutes and incubated for 2 hours at room temperature

with anti-dpERK mouse monoclonal antibody (1:200; Sigma, Saint Louis, MO) and

anti-mCherry polyclonal antibody from rabbit (1:500; Biovision, Milpitas, CA) in PBS

0.01% Triton-100 (T-Pbs) supplemented with 2% normal goat serum. Samples were

then washed three times in T-PBS and incubated in anti-mouse and anti-rabbit

antibodies (1 :500 each), respectively coupled with Alexa 488 and Alexa 568

(Invitrogen, Carlsbad, CA), overnight at 4°C or for 2 hours at room temperature.

Finally, samples were rinsed three times in T-PBS for 15 minutes and mounted in

Prolong Gold (Molecular Probes, Eugene, OR).

Molecular cloning
719
Coding sequences for wild-type M-Ras (KH.L172.2), Mek1/2 (KH.L147.22), Cdkn1b.a

(Cdkn1b, KH.C14.564), and Cdkn1b.b (Noto16, KH.S643.6) were PCR-amplified from

cDNA libraries prepared by reverse transcription of total RNA from mixed

developmental stages. Insertion of the products into expressing vectors was performed

using regular restriction/ligation or In-fusion (Clontech, Mountain View, CA)

procedure. Oligonucleotide directed mutagenesis or two-step overlap PCRs were used to

generate the point mutated forms M-Ras[G22V] and Mek[S220E,S216D] from the corresponding

wild-type sequences. We also used oligonucleotide directed mutagenesis to generate

mismatches in the PAM sequences adjacent to the sgRNA targets for Hand-r (153C>T

574C>T for Hand-r[PAMmis]) and Tbx1/10 (325G>A and 579G>A for Tbx1/10[PAMmis]). Due to

the absence of a correct PAM sequence (NGG, (reverse complement CCN)),

overexpressed Hand-r[PAMmis] and Tbx1/10[PAMmis] are resistant to the Cas9 nuclease

activity. We also used oligonucleotide directed mutagenesis to generate the mutations in

two putative Ets binding sites from the corresponding wild-type sequence of the

Tbx1/10 enhancer element: -3646TC>CT -3638GA>AG upstream from the initiation

codon (E1 and E2 in Figure S5 C, respectively). Primer sequences are listed in

Supplementary Table 1.

738
CRISPR/Cas9-mediated loss of Hand-r function

The pair of single guide RNA (sgRNA) targeting Tbx1/10 (sgTbx1/10) has been validated

previously (Tolkin and Christiaen, 2016). Rescue of the Tbx1/10 loss-of-function was

achieved by TVC-specific overexpression of Tbx1/10[PAMmis] driven by a *Foxf* enhancer

(Foxf-1>Tbx1/10[PAMmis]). For Hand-r loss of function, sgRNAs were first designed to

avoid genomic off-targets and tested as described (Gandhi et al., 2017b). In short,

sgRNA expressing cassettes (U6>sgRNA) were assembled by single step overlap PCR.

Individual PCR products (~25 μg) were electroporated with EF1a>NLS::Cas9::NLS

(30μg) , Myod905>Venus (50 μg), driving ubiquitous expression of Cas9 and a widely

expressed fluorescent reporter construct, respectively. Efficient electroporation was

confirmed by observation of fluorescence before genomic DNA extraction around 16 hpf

(18ºC) using QIAamp DNA Micro kit (Qiagen, German Town, MD). Mutagenesis

efficacy of individual sgRNAs, as a linear function of Cas9-induced indel frequency, was

estimated from electrophoregrams following Singer sequencing of the targeted regions

amplified from extracted genomic DNA by PCR. Result of the relative quantification of

the indel frequency ("corrected peakshift" of 22% and 24%) was considered high enough

for both sgRNAs targeting Hand-r, which were finally selected. The corresponding

cassettes were cloned into plasmid for repeated electroporations to study the loss of

function of Hand-r. Rescue of Hand-r loss-of-function was achieved by overexpression

of Hand-r[PAMmis] driven by a Foxf TVC specific enhancer (Foxf-1>Hand-r[PAMmis]). In order

to control the specificity of the CRISPR/Cas9 system, sgRNAs targeting *Neurogenin*, a

gene not expressed in the TVC and their progeny, was electroporated in parallel.

Sequences of the DNA targets and oligonucleotides used for the sgRNAs are listed in

Supplementary Table 1.

***Tbx1/10* enhancer and *cis*-regulatory analysis**

To isolate the minimal STVC-specific element of *Tbx1/10*, we used conserved non

coding sequences between *Ciona robusta* and *Ciona savignyi* as a guide for molecular

dissection (Figure S5 A, http://genome.lbl.gov/vista/index.shtml; (Frazer et al., 2004)).

We cloned a full-length *cis*-regulatory DNA construct (~7kbp) that was analyzed by

introducing large deletions to map the functional elements. We found a fragment of

1264 bp, that we called T6, located at -4682 bp upstream from the initiation codon that

was sufficient for STVC expression as well as in the mesenchyme and endoderm of the

reporter gene (Figure S5 A, B). 5' deletions of the T6 element show that the main *cis*-

regulatory elements required exclusively for STVC expression map in a 575 bp element,

which we called T12, at -4116 bp upstream from the initiation codon sufficient (Figure

S5A). The sequence of this element, which we called *T12*, reveals the presence of

putative Ets1/2 binding sites (Figure S5C) that were predicted using JASPAR (Khan et

al., 2017) and CisBP (Weirauch et al., 2014) databases. The minimal *Tbx1/10* STVC-

specific enhancer was further analyzed using point mutations of the candidate Ets1/2

sites with higher predicted scores (Figure S5D).

Observation and imaging

Samples were usually scored under a DM2500 epifluorescent microscope (Leica

Microsystems, Wetzlar, Germany). Imaging was performed using a TCS SP8 X inverted

confocal microscope equipped with a white light laser, AOBS and HyD detectors (Leica

Microsystems).

Acknowledgement

We thank Robert Kelly (Université Aix-Marseille, CNRS, France) for feedbacks on the

manuscript. We are grateful to Alberto Stolfi for collaborative inputs throughout the

project. We thank Farhana Salek and Kristyn Millan for technical support. This project

was funded by NIH/NHLBI R01 award HL108643, and trans-Atlantic network of

excellence award 15CVD01 from the Leducq Foundation to L.C.

Figures

Figure 1. Spatio-temporal restriction of ERK activity reflects FGF requirement for the specification of cardiopharyngeal progenitors. (**A**) Schematic of *Ciona* development showing asymmetric cell divisions and resulting cell fates of the cardiopharyngeal mesoderm (CPM). Embryonic and larval stages (St) according to (Hotta et al., 2007) with hours post fertilization (hpf) at 18°C. Anterior tail muscle (ATM, gray), trunk ventral cell (TVC, green), secondary TVC (STVC, green), first heart precursor (FHP, red), second heart precursor (SHP, orange), atrial siphon founder cell (ASMF, blue). Black bars link sister cells. Dashed lines: ventral midline. The first stage presents a quasi-lateral view while the second and third stages present quasi-ventral views. Anterior is to the left. Scale bar, 50 μm. (**B**) ERK activity visualized by anti-dpERK antibody (green). TVCs and their progeny are marked by mCherry driven by *Mesp* and revealed by anti-mCherry antibody (red). H$_2$B::mCherry and hCD4::mCherry accumulate in the nuclei and at the cell membrane, respectively. Arrowheads indicate STVCs and ASMFs at 14 and 16 hpf, respectively. Arrows indicate FHPs and open arrowheads mark SHPs. Anterior to the left. Scale bar, 10 μm. See also Figure S1 for broader time series of dpERK immunostaining in the B7.5 lineage. (**C, D**) TVC-specific overexpression of dnFGFR induces loss of expression of key lateral CPM markers visualized by *in situ* hybridization. (**C**) Representative expression patterns of key CPM genes (*Hand-related*, *Tbx1/10*, *Ebf*) in control embryos (Ctrl, electroporated with Foxf(TVC):bpFOG-1>Venus) and TVC-specific dnFGFR expression (electroporated with *Foxf(TVC):bpFOG-1>dnFGFR::mCherry*) individuals. TVCs and progeny are marked with *Mesp>NLS::lacZ* (red). Loss of expression in half of the TVC progeny, as presented for *Ebf*, is assumed to be due to left-right mosaicism. Arrowheads mark the ASMFs. Anterior is to the left. Scale bar, 10 μm. (**D**) Corresponding histograms with the phenotype proportions. For simplicity, loss of gene expression in half or all of the TVCs and their progeny were combined in the same category. "n" corresponds to the number of individual halves documented per condition.

821
Figure 2. Constitutively active M-Ras and MEK are sufficient to impose a pharyngeal muscle
fate in the cardiopharyngeal lineage. (**A**) Diagram of the FGF/MAPK transduction pathway with
constitutive activation by M-Ras^{G22V} and MEK^{S216D,S220E} mutants. (**B**) Expression patterns of markers of the
lateral TVC progeny, *Htr7* (a, b, c,), *Tbx1/10* (d, e, f) and *Ebf* (g, h, i), visualized by *in situ* hybridization
following TVC-specific over-expression of M-Ras^{WT} (as control), M-Ras^{G22V} and MEK^{S216D,S220E}. M-Ras^{WT}
overexpression (a, d, g) does not alter the wild-type spatial expression patterns of *Htr7*, *Tbx1/10* and *Ebf* in
lateral TVC derivatives (STVC and ASMF) and excluded from the median heart precursors. TVC-specific
over-expression of M-Ras^{G22V} (b, e, h) or MEK^{S216D,S220E} (c, f, i) induces ectopic expression of STVC and/or
ASMF markers (*Htr7*, *Tbx1/10* and *Ebf*) in the more median cells, that normally form cardiac precursors.
Arrowheads indicate STVCs and ASMFs at 15 and 18 hpf, respectively. Arrows indicate FHPs and open
arrowheads mark SHPs. At 18 hpf, the FHPs start dividing or have divided into 4 cells. Anterior to the left.
Scale bar, 10 μm. (**C**) Corresponding histograms: Larvae with TVC-specific over-expression of MEK^{WT}
retain the wild-type expression patterns. For simplicity, ectopic expressions in half to all of the cardiac
precursors were combined in the same phenotype category. "n" corresponds to the number of embryo
halves documented per condition. See also Figure S2.
837

A

1st asymmetric division

2nd asymmetric division

U0126 (5µM) incubation times

12-15 hpf

17-20 hpf

9.5-12.5 hpf

15-18 hpf

04   06   08   10   12   14   16   18   20   hpf (18°C)

B7.5 ⋯ B8.10 ATM TVC migration STVC ASMF oASMP iASMP SHP FHP

B7.5 B8.9 ATM TVC FHP

B n=200 n=200

Hand-r 12.5 hpf

☐ downregulation
■ WT pattern

DMSO U0126
9.5>12.5 hpf

C n=100 n=100

Hand-r 15 hpf

☐ downregulation
▨ WT pattern
■ FHP ectopic expression

DMSO U0126
12>15 hpf

D DMSO 12-15 hpf | U0126 12-15 hpf

15 hpf *Hand-r*

E n=100 n=100

Htr7 12.5 hpf

☐ downregulation
■ WT pattern

DMSO U0126
9.5>12.5 hpf

F n=100 n=100

Htr7 15 hpf

☐ downregulation
▨ WT pattern
■ FHP ectopic expression

DMSO U0126
12>15 hpf

G n=64 n=100

Tbx1/10 16 hpf

☐ downregulation
▨ WT pattern
■ ectopic expression

DMSO U0126
12>16 hpf

H n=132 n=128

Ebf 18 hpf

☐ downregulation
▨ WT pattern
■ ectopic expression

DMSO U0126
15>18 hpf

I n=65 n=92

Ebf transcription 20 hpf

☐ downregulation
▨ WT pattern
■ ectopic expression

DMSO U0126
17>20 hpf

J DMSO 17-20 hpf | U0126 17-20 hpf

20 hpf *intronic Ebf*

K Tbx1/10(T6) >M-Ras^WT | Tbx1/10(T6) >M-Ras^G22V | Tbx1/10(T6) >dnFGFR

18 hpf *Ebf*

L n=27 n=19 n=37 n=12

Ebf 18 hpf

☐ downregulation
▨ WT pattern
■ SHP ectopic expression

T6> Venus M-Ras^WT M-Ras^G22V dnFGFR

Figure 3. Temporal requirement for MAPK activity permits the progressive deployment of the cardiopharyngeal regulatory program. (**A**) Summary of the CPM cell lineage showing the different U0126 treatments with regard to the timing of the cell divisions. Abbreviations and color codes as in Figure 1. (**B, C**) Proportions of embryo halves with wild-type or downregulated expression of *Hand-r* at 12.5 hpf (B) and 15 hpf (C) following 3-hour incubations in U0126 (with DMSO as control treatment). (**D**) *Hand-r* expression visualized by *in situ* hybridization at 15 hpf in control (DMSO treated) and U0126 treated embryos. In control embryos, *Hand-r* remains expressed in the STVCs and downregulated in the FHPs. In U0126 (12-15 hpf) treated embryos, downregulation of *Hand-r* expression is observed throughout the TVC progeny (STVCs and FHPs), suggesting inhibition of transcription and inheritance of remnant

transcripts following TVC divisions. (**E, F**) Proportions of embryo halves with wild-type or downregulated
expression of *Htr7* at 12.5 hpf (E) and 15 hpf (F) following 3-hour incubations in U0126 (with DMSO as
control treatment). (**G**) Proportions of larvae with wild-type expression or downregulated expression of
Tbx1/10 at 16 hpf following 4-hour incubation in U0126 (with DMSO as control). (**H**) Proportions of larvae
with wild-type or downregulated expression of *Ebf* at 18 hpf following a three hour incubation in U0126
(with DMSO as control). (**I**) Proportions of larvae with wild-type or downregulated transcription of *Ebf* at
18 hpf following a three hour incubation in U0126 (DMSO as vehicle control). (**J**) Pattern of nascent *Ebf*
transcripts visualized by *in situ* hybridization with intronic probes (green) at 20 hpf. The nuclear dots
reveal the active transcription sites in the four ASMPs per side in larvae, both control/DMSO- and U0126-
treated from 17 to 20 hpf. (**K**) *Ebf* expression (green) in 18hpf larvae expressing control M-Ras^{WT},
constitutively active M-Ras^{G22V} or dominant negative dnFGFR under the control of the T12 element, an
STVC-specific *Tbx1/10* enhancer. Arrows: first heart precursors (FHP); open arrowhead: second heart
precursors (SHPs); closed arrowheads: ASM founder cells (ASMFs); dotted line: midline. (**L**) Proportions of
larvae with wild-type or downregulated expression of *Ebf* at 18 hpf in larvae with Venus (control), M-Ras^{WT},
M-Ras^{G22}, or dnFGFR driven by *Tbx1/10 cis*-regulatory sequence and overexpressed in the STVCs. "n" :
number of individual halves documented per condition.
865

867
Figure 4. M-Ras/MAPK-driven feed-forward subcircuits control the successive activations of
Hand-r, Tbx1/10 and Ebf. (**A**) Proportions of embryo halves with indicated *Tbx1/10* expression patterns
following TVC-specific CRISPR/Cas9-mediated mutagenesis of *Neurogenin/Neurog* as a control (*sgCtrl*),
and Hand-r (*sgHand-r*). TVC-specific overexpression of a CRISPR/Cas9-resistant form of Hand-r with
mutation in the PAM sequence (*Hand-r^PAMmis*) rescued *Tbx1/10* expression in the *sgHand-r* "background".
TVC-specific overexpression of a constitutively active M-Ras mutant (M-Ras^G22) (control: M-Ras^WT) was
sufficient to induce ectopic expression of *Tbx1/10* in the FHPs in *sgCtrl* embryos but not in *sgHand-r*
embryos indicating that Hand-r is necessary for M-Ras-dependent activation of *Tbx1/10* transcription. (**B**)
Proportions of embryo halves with indicated *Tbx1/10* expression patterns following TVC-specific
overexpression of Hand-r or a neutral reporter (Venus) and treated from 12 to 15hpf with the MEK
inhibitor U0126 (+) or with DMSO (-) as control. Hand-r overexpression is not sufficient to rescue loss of
Tbx1/10 expression due to MAPK inhibition indicating that M-Ras/MAPK activity is required in parallel of
Hand-r expression to activate Tbx1/10 transcription in the TVC progeny. (**C**) Tbx1/10 is necessary
downstream of M-Ras/MAPK activity to activate *Ebf* transcription in the TVC progeny. Shown are
proportions of *Ebf* expression phenotypes following TVC-specific CRISPR/Cas9-mediated loss of *Tbx1/10*
function (*sgTbx1/10*), with *Neurog*-targeting sgRNA as control (*sgCtrl*). Specificity of Tbx1/10 loss of
function was validated through rescue of *Ebf* expression with TVC-specific overexpression of a
CRISPR/Cas9 resistant form of *Tbx1/10* (Tbx1/10^PAMmis). Ectopic *Ebf* expression in SHPs in Tbx1/10^PAMmis
larvae is explained by precocious misexpression of Tbx1/10 in the TVC as described in Wang et al, 2013.
TVC-specific overexpression of M-Ras^G22 (M-Ras^G22), with wild type M-Ras (M-Ras^WT) as control, was
sufficient to induce ectopic expression of *Ebf* in the cardiac precursors in *sgCtrl* embryos but not in
sgTbx1/10 embryos indicating that Tbx1/10 is necessary for M-Ras-dependent activation of *Ebf*
transcription. (**D, E**) Proportions (D) and examples (E) of 15hpf larvae halves showing indicated *Ebf*
expression phenotypes in *sgCtrl* and *sgHand-r* CRISPR/Cas9 conditions combined with TVC-specific
overexpression of a neutral reporter (Venus), Hand-r^PAMmis, or Tbx1/10, and with MEK inhibition by U0126
(+) or not (DMSO control (-)). Arrowhead: STVCs, Arrows: FHPs, dotted line: ventral midline (**F**) Loss of
Hand-r function impaired the ability of Tbx1/10 to induce ectopic *Ebf* expression. For simplicity, ectopic
expressions in half to all of the cardiac precursors were combined in the same phenotype category. "n=":
number of individual halves documented per condition. (**G**) Summary model of the temporal deployment of
FGF/MAPK-driven feed-forward sub-circuits leading to the sequential activations of *Tbx1/10* and *Ebf* in the
STVCs and ASMFs, respectively.

Figure 5. Temporal deployment of the cardiopharyngeal network is partially coupled with cell cycle progression. (**A**) Schematic representation of the canonical eukaryotic cell cycle, and actions of the perturbations used in this study. (**B,C**) Tbx1/10 and Ebf expression at indicated time points, and following inhibition of cytokinesis by cytochalasin B treatment at indicated time points. Note that 15 to 19hpf treatment is applied AFTER the first division and birth the FHPs, which do not activate *Ebf* at 19hpf (right panel, arrows). (**D**) Inhibition of G1/S or G2/M blocks TVC division, and reduces *Tbx1/10* expression. Pictures shows TVCs that have divided in controls but not in experimental cells, with one cell occasionnally turning on *Tbx1/10*, but not the other. Left: the proportions of embryos showing strong *Tbx1/10* expression is substantially reduced compared to control embryos (e.g. Figure 1, and (Wang et al., 2013)). (**E**) Inhibition of G2/M in the STVCs by misexpression of Wee1 using the *Tbx1/10 T6* enhancer inhibits STVC division, and has a mild impact on *Ebf* expression at 18hpf. Open arrows indicate STVCs that have not divided, but express high (middle) or low (right) levels of *Ebf*. Left: control larva showing high *Ebf* expression in the ASMF (closed arrowheads), but neither in the SHPs (open arrowheads) nor in the FHPs (Arrows). (**F**) Misexpression of Wee1 using the *Tbx1/10 T6* enhancer (*Tbx1>Wee1*) inhibits STVC division. Right: the proportions of embryos showing strong *Lrp4/8, Kirr1* or *Dach* expression into late STVCs is reduced compared to SHPs in control embryos (T6>NLS::LacZ) at 18 hpf. Notably, the pan-cardiac marker *Lrp4/8* is still expressed in FHPs (arrows). Nuclei are marked in red with Mesp>NLS::lacZ, membranes in blue with Mesp>hCD4::mCherry, FHP labeled with arrows and SHP with arrowheads. 'n=', number of individual halves scored per condition. Scale bar, 5 μm. In all image panels, dotted line : ventral midline.

Figure 6. *Ebf* **regulation transitions from MAPK-dependent to autoregulative during the early phase of ASMF cycle.** (**A**) Proportions of 20hpf larva halves showing the indicated number of Ebf-expressing cells following U0126 treatments started at the indicated time points. This indicates that, by 17hpf, Ebf expression, which started at ~16hpf, has become largely insensitive to loss of MAPK activity. (**B**) Summary lineage diagram and time scale indicating the approximate stages for U0126 and DMSO (control) treatments for the results shown in (C, D). (C) Control (Ctrl) and Ebf-misexpressing embryos fixed at 16hpf, prior to chemical treatments, and stained for nascent transcripts with an intronic *Ebf* probe. In controls, the ASMFs (solid arrowhead), but neither the SHPs (open arrowheads) nor the FHPs (arrows), actively transcribe *Ebf* (green nuclear dots). In Larvae misexpressing the Ebf cDNA under the control of the STVC-specific Tbx1/10 enhancer, divisions are delayed and STVCs (solid arrowheads) activated transcription of endogenous *Ebf* loci (green nuclear dots). (D) After 4 hours, U0126 treated ASMFs no longer transcribe Ebf (top right image, solid arrowheads), whereas control DMSO-treated ASMFs do (top left, green nuclear dots). Upon misexpression of the Ebf cDNA in the STVCs and derivatives, ongoing *Ebf* transcription is detected at 20hpf in both DMSO and U0126-treated cells, and it persists in both ASMFs (solid arrowheads), and SHPs (open arrowheads). (**E, F**) Proportions of larvae halves showing the indicated *Ebf* transcription patterns, in indicated experimental conditions, as illustrated in C and D, respectively.

Figure 7. Summary model. (**A**) Schematic representation of cardiopharyngeal lineage cells at successive time points representing the main fate transitions. hpf: hours post-fertilization; TVC: trunk ventral cells; L: Leader T: trailer; migr.: migration; STVC: second trunk ventral cells; FHP: first heart precursors; dotted line: midline; black bars link sister cells; ASMF: atrial siphon muscle founder cells; SHP: second heart precursors; iASMP: inner atrial siphon muscle precursors; oASMP: outer atrial siphon muscle precursor (these cells correspond to stem-cell-like *Mrf*-; Notch+ precursors and *Mrf*+; Notch- differentiating myoblasts, respectively; see (Razy-Krajka et al., 2014) for details). (**B**) Lineage diagram and documented regulatory relationships between indicated genes and pathways, as showing here and in (Razy-Krajka et al., 2014; Wang et al., 2013). In TVCs, primed heart and ASM markers are coexpressed, and maintenance of the STVC and ASM markers requires ongoing FGF/MAPK signaling. Following the first oriented and asymmetric cell division, FGF-MAPK is maintained only in the STVCs, which permits the continued expression of Hand-r and the activation of *Tbx1/10*. Cell division, presumably through G1-specific inputs, contributes to *Tbx1/10* activation, and Tbx1/10 function antagonizes *Gata4/5/6* expression (Wang et al., 2013). In the FHPs, termination of FGF-MAPK signaling inhibits *Hand-r* expression and prevents *Tbx1/10* activation. Following oriented and asymmetric division of the STVCs, FGF/MAPK signaling persists only in the ASMFs, where it permits the transient maintenance of *Hand-r* and *Tbx1/10*, both of which act in parallel to FGF/MAPK to activate *Ebf* expression, together with contributions from presumed G1 inputs. Ebf activities further antagonize the cardiac program (marked by *Gata4/5/6*, *Nk4/Nkx2.5* and *Hand* expression; (Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et al., 2013)). Once Ebf expression reaches "high levels", its regulation becomes MAPK-independent and self-activating (this study). It also feeds back negatively on early activators such as Hand-r, and promotes the expression of the muscle determinant *Mrf* (Razy-Krajka et al., 2014; Tolkin and Christiaen, 2016). We propose that this transition represents commitment to an ASM fate. In the SHPs, termination of FGF/MAPK signaling prevents maintenance of Hand-r and Tbx1/10 expression, which, together with repressive inputs from Nk4/Nkx2.5, inhibits *Ebf* activation (Wang et al., 2013), and permits heart fate specification (Wang et al., submitted).

Figure S1. Detailed patterns of MAPK activity during early cardiopharyngeal development.
(**A**) MAPK activation during TVC induction. Close-up views of B7.5 lineage cells marked with
Mesp>H2B::mCherry (nuclei) and Mesp>hCD4::mCherry (membranes) and immunostained for dpERK at
indicated successive time points between 7 and 10hpf. DpERK staining was not detected in the founder cells
at 7hpf, but increased sharply and specifically in the smaller trunk ventral cells (TVCs, open arrows) at
7.5hpf, but not in the larger anterior tail muscles (ATMs). DpERK staining persisted throughout TVC
migration (see also B). (B) MAPK activation patterns during cardiopharyngeal fate diversification. DpERK
staining was clearly detected in migrating TVCs (open arrows, 11 to 13hpf); in lateral large STVCs (open
arrows, 14 to 15hpf), but not in the small median first heart precursors (FHPs, arrows, 14 to 15hpf); in the
large lateral atrial siphon muscle founder cells (ASMFs, solid arrowheads, 16 to 17hpf), but neither in the
FHPs (arrows), nor in the second heart precursors (SHPs, open arrowheads). (**C**) Treatment with the MEK
inhibitor U0126 from 12 to 15.5hpf abolished dpERK staining in the lateral STVCs, compared to a control
treatment with DMSO. Numbers of embryos showing the presented pattern out of the total numbers of
embryos are shown.

Figure S2 - part 1.
(A) Effects of perturbations on _Foxf_ TVC-specific enhancer activity. Proportions of Mesp>H2B:mCherry-positive embryos showing Foxf::bpFog-1>NLS:GFP activity (i.e. GFP+) in the indicated conditions : TVC-specific CRISPR/Cas9 mediated loss-of-function of Hand-r (sgHand-r), and corresponding control (Neurogenin/sgCtrl) at 15 hpf ; TVC-specific CRISPR/Cas9 mediated loss-of-function of Tbx1/10 (sgTbx1/10) and corresponding control (Neurogenin/sgCtrl) at 18 hpf ; TVC-targeted dnFGFR embryos (FoxF::bpFog-1>dnFGFR) and corresponding control (FoxF::bpFog-1>NLS::LacZ) at 15 hpf ; Inhibition of MAPK activity with 4-hour incubations in U0126 (DMSO as vehicle control) at indicated times. TVCS and their progeny marked with Mesp>H2B::mCherry and possible effects on _Foxf_ enhancer activity of these perturbations have been verified with TVC-specific green staining. There were no significant difference in the proportions of GFP+ embryos between each perturbations and controls. **(B-D) Other markers expressed in the TVC need continuous FGF-MAPK inputs for maintenance**. All panels show the proportions of 12.5hpf embryos halves showing expression of the indicated genes in late TVCs, following electroporation of either a Foxf(TVC)>Venus control of a Foxf(TVC)>dnFGFR construct that inhibits signaling through FGFR. Wild-type pattern were first reported in (Razy-Krajka et al., 2014).

Figure S2 - part 2.
(E-G) Foxf>dnFGFR does not inhibit TVC induction. TVC-specific expression of dnFGFR (FoxF::bpFog-1>dnFGFR) do not show difference in TVC markers expression in comparison to control (FoxF::bpFog-1>Venus) in early TVC (*Foxf* and *Hand-r* at 8 hpf, *Nkx2-5* at 10 hpf).
Arrows indicate anterior TVCs, adjacent cells on the right are ATMs, which do not express TVC markers.

Figure S3. Effects of defined FGF-MAPK signaling perturbations on cell division patterns in the cardiopharyngeal lineage. **(A)** Representative phenotype in unperturbed conditions (control plasmid: *Foxf>LacZ*) at 15hpf. TVCs divide asymmetrically and in a medio-lateral orientation. **(B)** Representative misoriented TVC division phenotype (Foxf>dnFGFR). One TVC divided improperly in the antero-posterior axis. **(C)** Proportions of embryo halves showing wild-type vs. orientation or symmetry phenotype, scored at 15hpf. For pharmacological treatments, embryos were incubated in DMSO or U0126 from 12hpf-15hpf. Proportions differ significantly between *Foxf>MRas* and *Foxf>caMRas*, and between DMSO and U0126 treatment conditions (** indicates p-value<0.01, Chi^2 test). **(D)** Representative phenotype in unperturbed conditions (FoxF>LacZ) at 18hpf. STVCs divide in a medio-lateral orientation. **(E)** Representative misoriented STVC division phenotype (FoxF>caMRas), scored as 'abnormal but can interpret.' STVCs likely divided improperly in the antero-posterior axis. **(F)** Proportions of larva halves in *Foxf*(TVC)-driven perturbations showing wild-type pattern vs. orientation and/or symmetry phenotypes, patterns scored at 18hpf as abnormal but interpretable or abnormal/can't interpret. For pharmacological treatments, larvae were incubated in DMSO or U0126 from 12hpf-18hpf. Proportions differ significantly in all conditions in comparison to the corresponding control (** indicates p-value<0.01, Chi^2 test). **(G)** Proportions of larva halves in *Tbx1*-driven perturbations showing wild-type pattern vs. orientation and/or symmetry phenotype, patterns scored at 18hpf as abnormal but interpretable or abnormal/can't interpret. For pharmacological treatments, larvae were incubated in DMSO or U0126 from 15hpf-18hpf. Proportions differ significantly only between Tbx1>MRas and Tbx1>caMRas conditions (** indicates p-value<0.01, Chi^2 test).

Figure S4. The constitutively active MEK[S216D,S220E] mutant is sufficient to impose a TVC identity to the whole B7.5 lineage. (A) Control late tailbud embryo showing the left side B7.5 lineage expressing GFP and a MEK[WT] control under the control of the *Mesp* enhancer. Two TVCs and two ATMs are normally induced, and TVCs migrated into the trunk. (B) Late tailbud embryo showing the left side B7.5 lineage expressing GFP and a MEK[S216D,S220E] mutant under the control of the *Mesp* enhancer. Four cells are observed as having migrated into the trunk, indicating that they have been induced to acquire a TVC fate and migrate, replicating FGF-MAPK gain-of-function phenotypes as described in (Davidson et al., 2006). (C) Proportions of embryo halves showing the indicated phenotypes. Extra migration is interpreted as ectopic induction of the TVC fate in all B7.5 lineage cells. Scale bar ~ 20μm.

1074
Figure S5. The Tbx1/10 enhancer has conserved putative Ets binding sites required for
reporter gene expression. (A) Alignment of *Tbx1/10* locus between *Ciona robusta/Ciona savignyi*
using VISTA. (B) *Ciona* larva (st 26) expressing GFP driven by the *T12* element, a STVC-specific *Tbx1/10*
enhancer (green) and Mesp>H2B::mCherry to track the B7.5 lineage cell nuclei (red). The *Tbx1/10*
enhancer drives the unc76::GFP reporter in the STVC progeny, including ASM precursors (ASMPs) (white
arrows) and the second heart precursors (SHPs) (white arrowheads). No expression is detected in the first
heart precursors (FHPs) (white arrowheads) and anterior tail muscle (ATMs) (orange arrows). (C)
Sequence alignment of *Tbx1/10* enhancer between *Ciona robusta/Ciona savignyi* Conserved blocks in the
orange boxes with putative Ets binding sites. (D) Proportion of larvae expressing both GFP and mCherry in
the STVC progeny when co-electroporated wild-type and mutant *Tbx1/10* reporters lacking the indicated
putative Ets binding sites and Mesp>H2B::mCherry in comparison to the control (no enhancer, only
Tbx1/10 basal promoter driving unc::76GFP). n: number of electroporated larval halves.
1087
1088

Figure S6. Dynamics of Ebf upregulation and entrainment by the cell cycle. (**A**) Proportions of larva halves fixed at successive time points and showing undivided STVCs, or ASMFs with or without ectopic *Ebf* expression in the SHPs following STVC-specific expression of the G2/M inhibitor Wee1 (+), or a control construct (-). See Figure S6E for an example of ectopic *Ebf* expression in the SHPs (grey labels). Note the sharp increase in % of larva with ASMF nbetween 15.5 and 16hpf, indicating that mitosis occurs primarily during this time window, but is delayed in a majority of larvae upon Wee1 misexpression. (**B**) Proportions of larva halves with cells showing indicated Ebf expression. The numbers (n) for cells expressing Wee1 focus on cells that have not divided (% shown in E), to estimate the dynamics of *Ebf* activation in G2/M-inhibited cells. Control cells consist mostly ASMFs after 15.5hpf as shown in (E). Wee1 and controls distributions differ significantly only at 16 and 16.5hpf (**, p<0.01 , Chi² test), suggesting that Wee1 merely delays the accumulation of Ebf transcripts. (**A**) 15.5hpf Cardiopharyngeal lineage cells

expressing Mesp>H2B::mCherry (red) and control Tbx1/10>unc76::GFP construct (not visible). **Rare precocious activation of *Ebf* transcription in STVCs.** (**C.a**) Green nuclear dot indicates nascent Ebf transcription in an STVC (open arrow), but not the other, and not in the first heart precursors (FHP; arrow). (**C.b**) left pair of nuclei shows an STVC (open arrow) and an FHP (arrow), neither of which express *Ebf*, whereas the cousin ASMF (solid arrowhead) shows nascent *Ebf* transcription (green dot). Dotted line : midline. (**D**) Proportions of STVCs and ASMFs showing indicated *Ebf* expression patterns. Note that >90% of STVCs do not express *Ebf*, which turns on almost exclusively in ASMFs. (**E**) Cardiopharyngeal lineage cells with *Ebf* expression in the ASMFs (solid arrowheads), and ectopically in the SHP (open arrowheads), but not in the FHPs (arrows), following misexpression of Wee1 using the STVC-specific *Tbx1/10 T12* enhancer. Dotted line: midline.

1113
Figure S7. MAPK signaling is necessary for *Ebf* expression only in early ASMF, and cell cycle
inputs shorten the MAPK-dependent period. (**A**) Proportions of larva halves showing the indicated
Ebf transcriptional activity (assayed using intronic probes). Batches of larvae expressing
Mesp>H2B::mCherry were split to be fixed for WMFISH or treated with U0126 at 3 successive time points
(15.75hpf, 16hpf or 16.25hpf), and the treated larvae were fixed at 20hpf. This data shows that, although all
batches expressed *Ebf* at the beginning of the experiment, only when MEK was inhibited later (16.25hpf)
did *Ebf* transcription persist in 20hpf larvae. (**B**) Proportions of larva halves showing the indicated
numbers of *Ebf*+ cells at 20hpf, following expression of the G2/M inhibitor Wee1 in the STVCs, under the
control of the *Tbx1/10 T12* enhancer (+). Negative controls (-) were electroporated with a
Tbx1/10(T12)>Venus construct. Larvae were also treated with U0126 (+) or DMSO (as negative control, (-
)), starting at 17hpf, which corresponds to the transition from a MAPK-dependent to a MAPK-independent
autoregulative mode of *Ebf* expression (see Figure 6A). Wee1-induced delays in cell cycle progression
increased the sensitivity of late *Ebf* expression to MAPK inhibition, further supporting the notion that cell
divisions accelerate the transition from MAPK-dependent to MAPK-independent self-activating regulation
of *Ebf* transcription.

REFERENCES

1129

Abu-Issa, R., Smyth, G., Smoak, I., Yamamura, K., Meyers, E.N., 2002. Fgf8 is
required for pharyngeal arch and cardiovascular development in the mouse.
Development 129, 4613-4625.

Aggarwal, V.S., Liao, J., Bondarev, A., Schimmang, T., Lewandoski, M., Locker, J.,
Shanske, A., Campione, M., Morrow, B.E., 2006. Dissection of Tbx1 and Fgf interactions
in mouse models of 22q11DS suggests functional redundancy. Human molecular
genetics 15, 3219-3228.

Alsan, B.H., Schultheiss, T.M., 2002. Regulation of avian cardiogenesis by Fgf8
signaling. Development 129, 1935-1943.

Barron, M., Gao, M., Lough, J., 2000. Requirement for BMP and FGF signaling
during cardiogenic induction in non-precardiac mesoderm is specific, transient, and
cooperative. Developmental dynamics : an official publication of the American
Association of Anatomists 218, 383-393.

Beh, J., Shi, W., Levine, M., Davidson, B., Christiaen, L., 2007. FoxF is essential for
FGF-induced migration of heart progenitor cells in the ascidian Ciona intestinalis.
Development 134, 3297-3305.

Bertrand, V., Hudson, C., Caillol, D., Popovici, C., Lemaire, P., 2003. Neural tissue in
ascidian embryos is induced by FGF9/16/20, acting via a combination of maternal
GATA and Ets transcription factors. Cell 115, 615-627.

Bondue, A., Lapouge, G., Paulissen, C., Semeraro, C., Iacovino, M., Kyba, M.,
Blanpain, C., 2008. Mesp1 acts as a master regulator of multipotent cardiovascular
progenitor specification. Cell stem cell 3, 69-84.

Bothe, I., Tenin, G., Oseni, A., Dietrich, S., 2011. Dynamic control of head mesoderm
patterning. Development 138, 2807-2821.

Brand, T., 2003. Heart development: molecular insights into cardiac specification
and early morphogenesis. Developmental biology 258, 1-19.

Brown, C.B., Wenning, J.M., Lu, M.M., Epstein, D.J., Meyers, E.N., Epstein, J.A.,
2004. Cre-mediated excision of Fgf8 in the Tbx1 expression domain reveals a critical
role for Fgf8 in cardiovascular development in the mouse. Developmental biology 267,
190-202.

Buckingham, M., Vincent, S.D., 2009. Distinct and dynamic myogenic populations in
the vertebrate embryo. Current opinion in genetics & development 19, 444-453.

Cai, C.L., Liang, X., Shi, Y., Chu, P.H., Pfaff, S.L., Chen, J., Evans, S., 2003. Isl1
identifies a cardiac progenitor population that proliferates prior to differentiation and
contributes a majority of cells to the heart. Developmental cell 5, 877-889.

Chan, S.S., Hagen, H.R., Swanson, S.A., Stewart, R., Boll, K.A., Aho, J., Thomson,
J.A., Kyba, M., 2016. Development of Bipotent Cardiac/Skeletal Myogenic Progenitors
from MESP1+ Mesoderm. Stem cell reports 6, 26-34.

Chan, S.S., Shi, X., Toyama, A., Arpke, R.W., Dandapat, A., Iacovino, M., Kang, J.,
Le, G., Hagen, H.R., Garry, D.J., Kyba, M., 2013. Mesp1 patterns mesoderm into
cardiac, hematopoietic, or skeletal myogenic progenitors in a context-dependent
manner. Cell stem cell 12, 587-601.

Chen, L., Fulcoli, F.G., Tang, S., Baldini, A., 2009. Tbx1 regulates proliferation and
differentiation of multipotent heart progenitors. Circulation research 105, 842-851.

Choi, C.Y., Lee, Y.M., Kim, Y.H., Park, T., Jeon, B.H., Schulz, R.A., Kim, Y., 1999. The
homeodomain transcription factor NK-4 acts as either a transcriptional activator or
repressor and interacts with the p300 coactivator and the Groucho corepressor. The
Journal of biological chemistry 274, 31543-31552.

Christiaen, L., Davidson, B., Kawashima, T., Powell, W., Nolla, H., Vranizan, K.,
Levine, M., 2008. The transcription/migration interface in heart precursors of Ciona
intestinalis. Science 320, 1349-1352.

Christiaen, L., Wagner, E., Shi, W., Levine, M., 2009a. Electroporation of transgenic
DNAs in the sea squirt Ciona. Cold Spring Harbor protocols 2009, pdb prot5345.

Christiaen, L., Wagner, E., Shi, W., Levine, M., 2009b. Isolation of sea squirt (Ciona)
gametes, fertilization, dechorionation, and development. Cold Spring Harbor protocols
2009, pdb prot5344.

Christiaen, L., Wagner, E., Shi, W., Levine, M., 2009c. Whole-mount in situ
hybridization on sea squirt (Ciona intestinalis) embryos. Cold Spring Harbor protocols
2009, pdb prot5348.

Cinnamon, E., Helman, A., Ben-Haroush Schyr, R., Orian, A., Jimenez, G., Paroush,
Z., 2008. Multiple RTK pathways downregulate Groucho-mediated repression in
Drosophila embryogenesis. Development 135, 829-837.

Cinnamon, E., Paroush, Z., 2008. Context-dependent regulation of Groucho/TLE-
mediated repression. Current opinion in genetics & development 18, 435-440.

Cooley, J., Whitaker, S., Sweeney, S., Fraser, S., Davidson, B., 2011. Cytoskeletal
polarity mediates localized induction of the heart progenitor lineage. Nature cell biology
13, 952-957.

Cota, C.D., Davidson, B., 2015. Mitotic Membrane Turnover Coordinates Differential
Induction of the Heart Progenitor Lineage. Developmental cell 34, 505-519.

Cowley, S., Paterson, H., Kemp, P., Marshall, C.J., 1994. Activation of MAP kinase
kinase is necessary and sufficient for PC12 differentiation and for transformation of
NIH 3T3 cells. Cell 77, 841-852.

Dalton, S., 2015. Linking the Cell Cycle to Cell Fate Decisions. Trends in cell biology
25, 592-600.

Davidson, B., 2007. Ciona intestinalis as a model for cardiac development. Seminars
in cell & developmental biology 18, 16-26.

Davidson, B., Levine, M., 2003. Evolutionary origins of the vertebrate heart:
Specification of the cardiac lineage in Ciona intestinalis. Proceedings of the National
Academy of Sciences of the United States of America 100, 11469-11473.

Davidson, B., Shi, W., Beh, J., Christiaen, L., Levine, M., 2006. FGF signaling
delineates the cardiac progenitor field in the simple chordate, Ciona intestinalis. Genes
& development 20, 2728-2738.

Davidson, B., Shi, W., Levine, M., 2005. Uncoupling heart cell specification and
migration in the simple chordate Ciona intestinalis. Development 132, 4811-4818.

de Pater, E., Clijsters, L., Marques, S.R., Lin, Y.-F., Garavito-Aguilar, Z.V., Yelon, D.,
Bakkers, J., 2009. Distinct phases of cardiomyocyte differentiation regulate growth of
the zebrafish heart. Development 136, 1633-1641.

Delsuc, F., Brinkmann, H., Chourrout, D., Philippe, H., 2006. Tunicates and not
cephalochordates are the closest living relatives of vertebrates. Nature 439, 965-968.

Diogo, R., Kelly, R.G., Christiaen, L., Levine, M., Ziermann, J.M., Molnar, J.L.,
Noden, D.M., Tzahor, E., 2015a. The Cardiopharyngeal Field and Vertebrate Evolution:
A New Heart for a New Head. Nature in press.

Diogo, R., Kelly, R.G., Christiaen, L., Levine, M., Ziermann, J.M., Molnar, J.L.,
Noden, D.M., Tzahor, E., 2015b. A new heart for a new head in vertebrate
cardiopharyngeal evolution. Nature 520, 466-473.

Dumollard, R., Minc, N., Salez, G., Aicha, S.B., Bekkouche, F., Hebras, C.,
Besnardeau, L., McDougall, A., 2017. The invariant cleavage pattern displayed by
ascidian embryos depends on spindle positioning along the cell's longest axis in the
apical plane and relies on asynchronous cell divisions. eLife 6.

Farley, E.K., Olson, K.M., Zhang, W., Brandt, A.J., Rokhsar, D.S., Levine, M.S., 2015.
Suboptimization of developmental enhancers. Science 350, 325-328.

Farley, E.K., Olson, K.M., Zhang, W., Rokhsar, D.S., Levine, M.S., 2016. Syntax
compensates for poor binding sites to encode tissue specificity of developmental
enhancers. Proceedings of the National Academy of Sciences of the United States of
America 113, 6508-6513.

Frazer, K.A., Pachter, L., Poliakov, A., Rubin, E.M., Dubchak, I., 2004. VISTA:
computational tools for comparative genomics. Nucleic acids research 32, W273-279.

Gainous, T.B., Wagner, E., Levine, M., 2015. Diverse ETS transcription factors
mediate FGF signaling in the Ciona anterior neural plate. Developmental biology 399,
218-225.

Gandhi, S., Haeussler, M., Razy-Krajka, F., Christiaen, L., Stolfi, A., 2017a.
Evaluation and rational design of guide RNAs for efficient CRISPR/Cas9-mediated
mutagenesis in Ciona. Developmental biology.

1243  Gandhi, S., Haeussler, M., Razy-Krajka, F., Christiaen, L., Stolfi, A., 2017b.
1244 Evaluation and rational design of guide RNAs for efficient CRISPR/Cas9-mediated
1245 mutagenesis in Ciona. bioRxiv.

1246  George, V., Colombo, S., Targoff, K.L., 2015. An early requirement for nkx2.5 ensures
1247 the first and second heart field ventricular identity and cardiac function into adulthood.
1248 Developmental biology 400, 10-22.

1249  Gopalakrishnan, S., Comai, G., Sambasivan, R., Francou, A., Kelly, R.G., Tajbakhsh,
1250 S., 2015. A Cranial Mesoderm Origin for Esophagus Striated Muscles. Developmental
1251 cell 34, 694-704.

1252  Gotoh, N., Laks, S., Nakashima, M., Lax, I., Schlessinger, J., 2004. FRS2 family
1253 docking proteins with overlapping roles in activation of MAP kinase have distinct
1254 spatial-temporal patterns of expression of their transcripts. FEBS letters 564, 14-18.

1255  Gueroult-Bellone, M., Nitta, K.R., Kari, W., Jacox, E., Beule Dauzat, R., Vincentelli,
1256 R., Diarra, C., Rothbacher, U., Dantec, C., Cambillau, C., Piette, J., Lemaire, P., 2017.
1257 Spacer sequences separating transcription factor binding motifs set enhancer quality
1258 and strength. bioRxiv.

1259  Hasson, P., Egoz, N., Winkler, C., Volohonsky, G., Jia, S., Dinur, T., Volk, T., Courey,
1260 A.J., Paroush, Z., 2005. EGFR signaling attenuates Groucho-dependent repression to
1261 antagonize Notch transcriptional output. Nature genetics 37, 101-105.

1262  Haupaix, N., Abitua, P.B., Sirour, C., Yasuo, H., Levine, M., Hudson, C., 2014.
1263 Ephrin-mediated restriction of ERK1/2 activity delimits the number of pigment cells in
1264 the Ciona CNS. Developmental biology 394, 170-180.

1265  Hotta, K., Mitsuhara, K., Takahashi, H., Inaba, K., Oka, K., Gojobori, T., Ikeo, K.,
1266 2007. A web-based interactive developmental table for the ascidian Ciona intestinalis,
1267 including 3D real-image embryo reconstructions: I. From fertilized egg to hatching
1268 larva. Developmental dynamics : an official publication of the American Association of
1269 Anatomists 236, 1790-1805.

1270  Hu, T., Yamagishi, H., Maeda, J., McAnally, J., Yamagishi, C., Srivastava, D., 2004.
1271 Tbx1 regulates fibroblast growth factors in the anterior heart field through a reinforcing
1272 autoregulatory loop involving forkhead transcription factors. Development 131, 5491-
1273 5502.

1274  Hudson, C., Darras, S., Caillol, D., Yasuo, H., Lemaire, P., 2003. A conserved role for
1275 the MEK signalling pathway in neural tissue specification and posteriorisation in the
1276 invertebrate chordate, the ascidian Ciona intestinalis. Development 130, 147-159.

1277  Hutson, M.R., Zeng, X.L., Kim, A.J., Antoon, E., Harward, S., Kirby, M.L., 2010.
1278 Arterial pole progenitors interpret opposing FGF/BMP signals to proliferate or
1279 differentiate. Development 137, 3001-3011.

1280  Ilagan, R., Abu-Issa, R., Brown, D., Yang, Y.P., Jiao, K., Schwartz, R.J., Klingensmith,
1281 J., Meyers, E.N., 2006. Fgf8 is required for anterior heart field development.
1282 Development 133, 2435-2445.

Imai, K.S., Levine, M., Satoh, N., Satou, Y., 2006. Regulatory blueprint for a chordate
embryo. Science 312, 1183-1187.

Imai, K.S., Satoh, N., Satou, Y., 2002. Early embryonic expression of FGF4/6/9 gene
and its role in the induction of mesenchyme and notochord in Ciona savignyi embryos.
Development 129, 1729-1738.

Jeffery, W.R., Chiba, T., Krajka, F.R., Deyts, C., Satoh, N., Joly, J.S., 2008. Trunk
lateral cells are neural crest-like cells in the ascidian Ciona intestinalis: insights into the
ancestry and evolution of the neural crest. Developmental biology 324, 152-160.

Jerome, L.A., Papaioannou, V.E., 2001. DiGeorge syndrome phenotype in mice
mutant for the T-box gene, Tbx1. Nature genetics 27, 286-291.

Kaplan, N., Razy-Krajka, F., Christiaen, L., 2015. Regulation and evolution of
cardiopharyngeal cell identity and behavior: insights from simple chordates. Current
opinion in genetics & development 32, 119-128.

Kattman, S.J., Witty, A.D., Gagliardi, M., Dubois, N.C., Niapour, M., Hotta, A., Ellis,
J., Keller, G., 2011. Stage-specific optimization of activin/nodal and BMP signaling
promotes cardiac differentiation of mouse and human pluripotent stem cell lines. Cell
stem cell 8, 228-240.

Keduka, E., Kaiho, A., Hamada, M., Watanabe-Takano, H., Takano, K., Ogasawara,
M., Satou, Y., Satoh, N., Endo, T., 2009. M-Ras evolved independently of R-Ras and its
neural function is conserved between mammalian and ascidian, which lacks classical
Ras. Gene 429, 49-58.

Kelly, R.G., Jerome-Majewska, L.A., Papaioannou, V.E., 2004. The del22q11.2
candidate gene Tbx1 regulates branchiomeric myogenesis. Human molecular genetics
13, 2829-2840.

Kelly, R.G., Papaioannou, V.E., 2007. Visualization of outflow tract development in
the absence of Tbx1 using an FgF10 enhancer trap transgene. Developmental dynamics :
an official publication of the American Association of Anatomists 236, 821-828.

Khan, A., Fornes, O., Stigliani, A., Gheorghe, M., Castro-Mondragon, J.A., van der
Lee, R., Bessy, A., Cheneby, J., Kulkarni, S.R., Tan, G., Baranasic, D., Arenillas, D.J.,
Sandelin, A., Vandepoele, K., Lenhard, B., Ballester, B., Wasserman, W.W., Parcy, F.,
Mathelier, A., 2017. JASPAR 2018: update of the open-access database of transcription
factor binding profiles and its web framework. Nucleic acids research.

Khoueiry, P., Rothbacher, U., Ohtsuka, Y., Daian, F., Frangulian, E., Roure, A.,
Dubchak, I., Lemaire, P., 2010. A cis-regulatory signature in ascidians and flies,
independent of transcription factor binding sites. Current biology : CB 20, 792-802.

Kuwajima, M., Kumano, G., Nishida, H., 2014. Regulation of the number of cell
division rounds by tissue-specific transcription factors and Cdk inhibitor during
ascidian embryogenesis. PloS one 9, e90188.

Lazic, S., Scott, I.C., 2011. Mef2cb regulates late myocardial cell addition from a
second heart field-like population of progenitors in zebrafish. Developmental biology
354, 123-133.

Lemmon, M.A., Schlessinger, J., 2010. Cell signaling by receptor tyrosine kinases.
Cell 141, 1117-1134.

Lescroart, F., Chabab, S., Lin, X., Rulands, S., Paulissen, C., Rodolosse, A., Auer, H.,
Achouri, Y., Dubois, C., Bondue, A., Simons, B.D., Blanpain, C., 2014. Early lineage
restriction in temporally distinct populations of Mesp1 progenitors during mammalian
heart development. Nature cell biology 16, 829-840.

Lescroart, F., Hamou, W., Francou, A., Theveniau-Ruissy, M., Kelly, R.G.,
Buckingham, M., 2015. Clonal analysis reveals a common origin between nonsomite-
derived neck muscles and heart myocardium. Proceedings of the National Academy of
Sciences of the United States of America 112, 1446-1451.

Lescroart, F., Kelly, R.G., Le Garrec, J.F., Nicolas, J.F., Meilhac, S.M., Buckingham,
M., 2010. Clonal analysis reveals common lineage relationships between head muscles
and second heart field derivatives in the mouse embryo. Development 137, 3269-3279.

Lescroart, F., Mohun, T., Meilhac, S.M., Bennett, M., Buckingham, M., 2012. Lineage
tree for the venous pole of the heart: clonal analysis clarifies controversial genealogy
based on genetic tracing. Circulation research 111, 1313-1322.

Mandal, A., Holowiecki, A., Song, Y.C., Waxman, J.S., 2017. Wnt signaling balances
specification of the cardiac and pharyngeal muscle fields. Mechanisms of Development
143, 32-41.

Mansour, S.J., Resing, K.A., Candi, J.M., Hermann, A.S., Gloor, J.W., Herskind,
K.R., Wartmann, M., Davis, R.J., Ahn, N.G., 1994. Mitogen-activated protein (MAP)
kinase phosphorylation of MAP kinase kinase: determination of phosphorylation sites
by mass spectrometry and site-directed mutagenesis. Journal of biochemistry 116, 304-
314.

Marques, S.R., Lee, Y., Poss, K.D., Yelon, D., 2008. Reiterative roles for FGF
signaling in the establishment of size and proportion of the zebrafish heart.
Developmental biology 321, 397-406.

Mazzoni, E.O., Mahony, S., Iacovino, M., Morrison, C.A., Mountoufaris, G., Closser,
M., Whyte, W.A., Young, R.A., Kyba, M., Gifford, D.K., Wichterle, H., 2011. Embryonic
stem cell-based mapping of developmental transcriptional programs. Nature methods
8, 1056-1058.

Merscher, S., Funke, B., Epstein, J.A., Heyer, J., Puech, A., Lu, M.M., Xavier, R.J.,
Demay, M.B., Russell, R.G., Factor, S., Tokooya, K., Jore, B.S., Lopez, M., Pandita, R.K.,
Lia, M., Carrion, D., Xu, H., Schorle, H., Kobler, J.B., Scambler, P., Wynshaw-Boris, A.,
Skoultchi, A.I., Morrow, B.E., Kucherlapati, R., 2001. TBX1 is responsible for
cardiovascular defects in velo-cardio-facial/DiGeorge syndrome. Cell 104, 619-629.

Michailovici, I., Eigler, T., Tzahor, E., 2015. Craniofacial Muscle Development.
Current topics in developmental biology 115, 3-30.

Michailovici, I., Harrington, H.A., Azogui, H.H., Yahalom-Ronen, Y., Plotnikov, A.,
Ching, S., Stumpf, M.P., Klein, O.D., Seger, R., Tzahor, E., 2014. Nuclear to cytoplasmic
shuttling of ERK promotes differentiation of muscle stem/progenitor cells.
Development 141, 2611-2620.

Mosimann, C., Panakova, D., Werdich, A.A., Musso, G., Burger, A., Lawson, K.L.,
Carr, L.A., Nevis, K.R., Sabeh, M.K., Zhou, Y., Davidson, A.J., DiBiase, A., Burns, C.E.,
Burns, C.G., MacRae, C.A., Zon, L.I., 2015. Chamber identity programs drive early
functional partitioning of the heart. Nature communications 6, 8146.

Nathan, E., Monovich, A., Tirosh-Finkel, L., Harrelson, Z., Rousso, T., Rinon, A.,
Harel, I., Evans, S.M., Tzahor, E., 2008. The contribution of Islet1-expressing
splanchnic mesoderm cells to distinct branchiomeric muscles reveals significant
heterogeneity in head muscle development. Development 135, 647-657.

Nevis, K., Obregon, P., Walsh, C., Guner-Ataman, B., Burns, C.G., Burns, C.E., 2013.
Tbx1 is required for second heart field proliferation in zebrafish. Developmental
dynamics : an official publication of the American Association of Anatomists 242, 550-
559.

Norton, J., Cooley, J., Islam, A.F., Cota, C.D., Davidson, B., 2013. Matrix adhesion
polarizes heart progenitor induction in the invertebrate chordate Ciona intestinalis.
Development 140, 1301-1311.

Park, E.J., Ogden, L.A., Talbot, A., Evans, S., Cai, C.L., Black, B.L., Frank, D.U.,
Moon, A.M., 2006. Required, tissue-specific roles for Fgf8 in outflow tract formation
and remodeling. Development 133, 2419-2433.

Park, E.J., Watanabe, Y., Smyth, G., Miyagawa-Tomita, S., Meyers, E., Klingensmith,
J., Camenisch, T., Buckingham, M., Moon, A.M., 2008. An FGF autocrine loop initiated
in second heart field mesoderm regulates morphogenesis at the arterial pole of the
heart. Development 135, 3599-3610.

Patterson, K.I., Brummer, T., O'Brien, P.M., Daly, R.J., 2009. Dual-specificity
phosphatases: critical regulators with diverse cellular targets. The Biochemical journal
418, 475-489.

Pauklin, S., Madrigal, P., Bertero, A., Vallier, L., 2016. Initiation of stem cell
differentiation involves cell cycle-dependent regulation of developmental genes by
Cyclin D. Genes & development 30, 421-433.

Pauklin, S., Vallier, L., 2013. The cell-cycle state of stem cells determines cell fate
propensity. Cell 155, 135-147.

Peljto, M., Wichterle, H., 2011. Programming embryonic stem cells to neuronal
subtypes. Current opinion in neurobiology 21, 43-51.

Pennati, R., Ficetola, G.F., Brunetti, R., Caicci, F., Gasparini, F., Griggio, F., Sato, A.,
Stach, T., Kaul-Strehlow, S., Gissi, C., Manni, L., 2015. Morphological Differences
between Larvae of the Ciona intestinalis Species Complex: Hints for a Valid Taxonomic
Definition of Distinct Species. PloS one 10, e0122879.

Picco, V., Hudson, C., Yasuo, H., 2007. Ephrin-Eph signalling drives the asymmetric
division of notochord/neural precursors in Ciona embryos. Development 134, 1491-
1497.

Prall, O.W., Menon, M.K., Solloway, M.J., Watanabe, Y., Zaffran, S., Bajolle, F.,
Biben, C., McBride, J.J., Robertson, B.R., Chaulet, H., Stennard, F.A., Wise, N., Schaft,
D., Wolstein, O., Furtado, M.B., Shiratori, H., Chien, K.R., Hamada, H., Black, B.L.,
Saga, Y., Robertson, E.J., Buckingham, M.E., Harvey, R.P., 2007. An Nkx2-
5/Bmp2/Smad1 negative feedback loop controls heart progenitor specification and
proliferation. Cell 128, 947-959.

Putnam, N.H., Butts, T., Ferrier, D.E., Furlong, R.F., Hellsten, U., Kawashima, T.,
Robinson-Rechavi, M., Shoguchi, E., Terry, A., Yu, J.K., Benito-Gutierrez, E.L.,
Dubchak, I., Garcia-Fernandez, J., Gibson-Brown, J.J., Grigoriev, I.V., Horton, A.C., de
Jong, P.J., Jurka, J., Kapitonov, V.V., Kohara, Y., Kuroki, Y., Lindquist, E., Lucas, S.,
Osoegawa, K., Pennacchio, L.A., Salamov, A.A., Satou, Y., Sauka-Spengler, T., Schmutz,
J., Shin, I.T., Toyoda, A., Bronner-Fraser, M., Fujiyama, A., Holland, L.Z., Holland,
P.W., Satoh, N., Rokhsar, D.S., 2008. The amphioxus genome and the evolution of the
chordate karyotype. Nature 453, 1064-1071.

Racioppi, C., Kamal, A.K., Razy-Krajka, F., Gambardella, G., Zanetti, L., di Bernardo,
D., Sanges, R., Christiaen, L.A., Ristoratore, F., 2014. Fibroblast growth factor signalling
controls nervous system patterning and pigment cell formation in Ciona intestinalis.
Nature communications 5, 4830.

Razy-Krajka, F., Lam, K., Wang, W., Stolfi, A., Joly, M., Bonneau, R., Christiaen, L.,
2014. Collier/OLF/EBF-Dependent Transcriptional Dynamics Control Pharyngeal
Muscle Specification from Primed Cardiopharyngeal Progenitors. Developmental cell
29, 263-276.

Reifers, F., Walsh, E.C., Leger, S., Stainier, D.Y., Brand, M., 2000. Induction and
differentiation of the zebrafish heart requires fibroblast growth factor 8
(fgf8/acerebellar). Development 127, 225-235.

Satou, Y., Imai, K.S., Satoh, N., 2004. The ascidian Mesp gene specifies heart
precursor cells. Development 131, 2533-2541.

Shi, W., Levine, M., 2008. Ephrin signaling establishes asymmetric cell fates in an
endomesoderm lineage of the Ciona embryo. Development 135, 931-940.

Shi, W., Peyrot, S.M., Munro, E., Levine, M., 2009. FGF3 in the floor plate directs
notochord convergent extension in the Ciona tadpole. Development 136, 23-28.

Soufi, A., Dalton, S., 2016. Cycling through developmental decisions: how cell cycle
dynamics control pluripotency, differentiation and reprogramming. Development 143,
4301-4311.

Stolfi, A., Gainous, T.B., Young, J.J., Mori, A., Levine, M., Christiaen, L., 2010. Early
chordate origins of the vertebrate second heart field. Science 329, 565-568.

Stolfi, A., Gandhi, S., Salek, F., Christiaen, L., 2014a. Tissue-specific genome editing
in Ciona embryos by CRISPR/Cas9. Development 141, 4115-4120.

Stolfi, A., Lowe, E.K., Racioppi, C., Ristoratore, F., Brown, C.T., Swalla, B.J.,
Christiaen, L., 2014b. Divergent mechanisms regulate conserved cardiopharyngeal
development and gene expression in distantly related ascidians. eLife 3, e03728.

Stolfi, A., Sasakura, Y., Satou, Y., Christiaen, L., Dantec, C., Endo, T., Naville, M.,
Nishida, H., Swalla, B., Volff, J.-N., Voskoboynik, A., Dauga, D., Lemaire, P., 2014c.
Guidelines for the Nomenclature of Genetic Elements in Tunicate Genomes. Genesis
under review.

Stolfi, A., Wagner, E., Taliaferro, J.M., Chou, S., Levine, M., 2011. Neural tube
patterning by Ephrin, FGF and Notch signaling relays. Development 138, 5429-5439.

Tirosh-Finkel, L., Elhanany, H., Rinon, A., Tzahor, E., 2006. Mesoderm progenitor
cells of common origin contribute to the head musculature and the cardiac outflow
tract. Development 133, 1943-1953.

Tirosh-Finkel, L., Zeisel, A., Brodt-Ivenshitz, M., Shamai, A., Yao, Z., Seger, R.,
Domany, E., Tzahor, E., 2010. BMP-mediated inhibition of FGF signaling promotes
cardiomyocyte differentiation of anterior heart field progenitors. Development 137,
2989-3000.

Tolkin, T., Christiaen, L., 2016. Rewiring of an ancestral Tbx1/10-Ebf-Mrf network
for pharyngeal muscle specification in distinct embryonic lineages. Development in
press.

Tzahor, E., 2009. Heart and craniofacial muscle development: a new developmental
theme of distinct myogenic fields. Developmental biology 327, 273-279.

Tzahor, E., Evans, S.M., 2011. Pharyngeal mesoderm development during
embryogenesis: implications for both heart and head myogenesis. Cardiovascular
research 91, 196-202.

Tzahor, E., Kempf, H., Mootoosamy, R.C., Poon, A.C., Abzhanov, A., Tabin, C.J.,
Dietrich, S., Lassar, A.B., 2003. Antagonists of Wnt and BMP signaling promote the
formation of vertebrate head muscle. Genes & development 17, 3087-3099.

Tzahor, E., Lassar, A.B., 2001. Wnt signals from the neural tube block ectopic
cardiogenesis. Genes & development 15, 255-260.

van Wijk, B., van den Berg, G., Abu-Issa, R., Barnett, P., van der Velden, S., Schmidt,
M., Ruijter, J.M., Kirby, M.L., Moorman, A.F., van den Hoff, M.J., 2009. Epicardium
and myocardium separate from a common precursor pool by crosstalk between bone
morphogenetic protein- and fibroblast growth factor-signaling pathways. Circulation
research 105, 431-441.

Vitelli, F., Morishima, M., Taddei, I., Lindsay, E.A., Baldini, A., 2002a. Tbx1
mutation causes multiple cardiovascular defects and disrupts neural crest and cranial
nerve migratory pathways. Human molecular genetics 11, 915-922.

Vitelli, F., Taddei, I., Morishima, M., Meyers, E.N., Lindsay, E.A., Baldini, A., 2002b.
A genetic link between Tbx1 and fibroblast growth factor signaling. Development 129,
4605-4611.

von Scheven, G., Alvares, L.E., Mootoosamy, R.C., Dietrich, S., 2006. Neural tube
derived signals and Fgf8 act antagonistically to specify eye versus mandibular arch
muscles. Development 133, 2731-2745.

Wagner, E., Stolfi, A., Gi Choi, Y., Levine, M., 2014. Islet is a key determinant of
ascidian palp morphogenesis. Development 141, 3084-3092.

Wang, W., Razy-Krajka, F., Siu, E., Ketcham, A., Christiaen, L., 2013. NK4
antagonizes Tbx1/10 to promote cardiac versus pharyngeal muscle fate in the ascidian
second heart field. PLoS biology 11, e1001725.

Wang, W., Xiang, S., Jullian, E., Kelly, R.G., Satija, R., Christiaen, L., 2017. A single
cell transcriptional roadmap for cardiopharyngeal fate diversification. BioRxiv.

Watanabe, Y., Miyagawa-Tomita, S., Vincent, S.D., Kelly, R.G., Moon, A.M.,
Buckingham, M.E., 2010. Role of mesodermal FGF8 and FGF10 overlaps in the
development of the arterial pole of the heart and pharyngeal arch arteries. Circulation
research 106, 495-503.

Watanabe, Y., Zaffran, S., Kuroiwa, A., Higuchi, H., Ogura, T., Harvey, R.P., Kelly,
R.G., Buckingham, M., 2012. Fibroblast growth factor 10 gene regulation in the second
heart field by Tbx1, Nkx2-5, and Islet1 reveals a genetic switch for down-regulation in
the myocardium. Proceedings of the National Academy of Sciences of the United States
of America 109, 18273-18280.

Weirauch, M.T., Yang, A., Albu, M., Cote, A.G., Montenegro-Montero, A., Drewe, P.,
Najafabadi, H.S., Lambert, S.A., Mann, I., Cook, K., Zheng, H., Goity, A., van Bakel, H.,
Lozano, J.C., Galli, M., Lewsey, M.G., Huang, E., Mukherjee, T., Chen, X., Reece-Hoyes,
J.S., Govindarajan, S., Shaulsky, G., Walhout, A.J.M., Bouget, F.Y., Ratsch, G.,
Larrondo, L.F., Ecker, J.R., Hughes, T.R., 2014. Determination and inference of
eukaryotic transcription factor sequence specificity. Cell 158, 1431-1443.

Whittaker, J.R., 1973. Segregation during ascidian embryogenesis of egg cytoplasmic
information for tissue-specific enzyme development. Proceedings of the National
Academy of Sciences of the United States of America 70, 2096-2100.

Witzel, H.R., Cheedipudi, S., Gao, R., Stainier, D.Y., Dobreva, G.D., 2017. Isl2b
regulates anterior second heart field development in zebrafish. Scientific reports 7,
41043.

Woznica, A., Haeussler, M., Starobinska, E., Jemmett, J., Li, Y., Mount, D.,
Davidson, B., 2012. Initial deployment of the cardiogenic gene regulatory network in the
basal chordate, Ciona intestinalis. Developmental biology 368, 127-139.

Yagi, H., Furutani, Y., Hamada, H., Sasaki, T., Asakawa, S., Minoshima, S., Ichida, F.,
Joo, K., Kimura, M., Imamura, S., Kamatani, N., Momma, K., Takao, A., Nakazawa, M.,
Shimizu, N., Matsuoka, R., 2003. Role of TBX1 in human del22q11.2 syndrome. Lancet
362, 1366-1373.

Yasuo, H., Hudson, C., 2007. FGF8/17/18 functions together with FGF9/16/20
during formation of the notochord in Ciona embryos. Developmental biology 302, 92-
103.

Zaffran, S., Frasch, M., 2002. Early signals in cardiac development. Circulation
research 91, 457-469.

Zhang, Z., Huynh, T., Baldini, A., 2006. Mesodermal expression of Tbx1 is necessary
and sufficient for pharyngeal arch and cardiac outflow tract development. Development
133, 3587-3595.
1529