An FGF-driven feed-forward circuit for spatiotemporal patterning of the

cardiopharyngeal mesoderm in a simple chordate

Florian Razy-Krajka, Basile Gravez and Lionel Christiaen*

Center for Developmental Genetics, Department of Biology, College of Arts and Science,

New York University, New York, NY, USA

* author for correspondence: email: lc121@nyu.edu, twitter: @lionlchristiaen, phone: +1

10   212 992 8695

Abstract

In embryos, pluripotent stem cells and multipotent progenitors must divide and produce distinct progeny to express their full developmental potential. In vertebrates, mounting evidence point to the existence of multipotent cardiopharyngeal progenitors that produce second-heart-field-derived cardiomyocytes, and branchiomeric skeletal head muscles. However, the cellular and molecular mechanisms underlying these early fate choices remain largely elusive. The tunicate *Ciona* has emerged as an attractive model to study early cardiopharyngeal development at high spatial and temporal resolution: through two asymmetric and oriented cell divisions, defined multipotent cardiopharyngeal progenitors produce distinct first and second heart precursors, and pharyngeal muscle (aka atrial siphon muscle, ASM) precursors. Here, we demonstrate that differential FGF/MAPK signaling distinguishes between MAPK-negative heart precursors, and MAPK-positive multipotent progenitors and ASM precursors. We characterize an FGF/MAPK-driven feed-forward circuit that promotes the successive activations of essential cardiopharyngeal determinants, *Tbx1/10* and *Ebf*. Finally, we show that coupling FGF/MAPK restriction and cardiopharyngeal network deployment with cell divisions permits the emergence of diverse cell types from common multipotent progenitors.

Introduction

In the past few years, studies guided by developmental genetics knowledge progressed towards driving mammalian stem cells into forming pure cultures of selected cell types *in vitro* (e.g. (Kattman et al., 2011; Mazzoni et al., 2011; Peljto and Wichterle, 2011). By contrast, in their embryonic context, pluripotent cells generate diverse cell types in defined proportions. This simple observation implies that pluripotent stem cells and multipotent embryonic progenitors must divide before individual cells among their progeny adopt distinct fates, as a result of differential exposure to inducing signals and/or inheritance of cell autonomous determinants.

Subsets of the heart and head/neck myocytes recently emerged as related derivatives of multipotent progenitors located in the mesodermal cardiopharyngeal field (Diogo et al., 2015; Tzahor, 2009; Tzahor and Evans, 2011). Specifically, early lineage tracing, transplantations and controlled explant culture experiments demonstrated that the anterior splanchnic/pharyngeal mesoderm of amniote embryos can produce either skeletal muscles or heart tissue, depending upon exposure to growth factors and signaling molecules (Nathan et al., 2008; Tirosh-Finkel et al., 2006; Tzahor et al., 2003; Tzahor and Lassar, 2001). Clonal analyses in the mouse further revealed the existence of common *Mesp1*-expressing progenitors for subsets of the second heart field-derived cardiomyocytes and branchiomeric facial, jaw, neck and even esophageal muscles (Gopalakrishnan et al., 2015; Lescroart et al., 2014; Lescroart et al., 2015; Lescroart et al., 2010; Lescroart et al., 2012). *In vitro* studies using pluripotent stem cells indicated that controlled *Mesp1* expression can drive mesodermal progenitors towards cardiac and/or skeletal muscle fates (Bondue et al., 2008; Chan et al., 2016; Chan et al., 2013). Genetic labeling and functional studies showed that proper development of the pharyngeal apparatus and second heart field derivatives require shared inputs from Tbx1, Nkx2-5 and Islet1 transcription factors (e.g. (Cai et al., 2003; George et al., 2015;

Jerome and Papaioannou, 2001; Kelly et al., 2004; Merscher et al., 2001; Mosimann et al., 2015; Nevis et al., 2013; Prall et al., 2007; Tzahor and Evans, 2011; Vitelli et al., 2002a; Watanabe et al., 2012; Witzel et al., 2017; Yagi et al., 2003; Zhang et al., 2006)). Taken together, a growing body of evidence point to the existence of a mesodermal field of multipotent progenitors capable of producing either SHF-derived cardiomyocytes or branchiomeric skeletal muscles in early amniote embryos (Diogo et al., 2015; Mandal et al., 2017). However, the mechanisms that distinguish fate-restricted heart and head muscle precursors remain largely elusive.

The tunicate *Ciona*, which is among the closest living relatives to the vertebrates (Delsuc et al., 2006; Putnam et al., 2008), has emerged as a simple chordate model to characterize multipotent cardiopharyngeal progenitors and the mechanisms that initiate heart vs. pharyngeal muscle fate choices (Kaplan et al., 2015; Razy-Krajka et al., 2014; Stolfi et al., 2010; Tolkin and Christiaen, 2016; Wang et al., 2013). *Ciona* tailbud embryos possess two multipotent cardiopharyngeal progenitors on either side. Like their vertebrate counterparts, these cells emerge from *Mesp+* progenitors towards the end of gastrulation; they are induced by FGF/MAPK signaling and have been termed *trunk ventral cells* (aka TVCs; (Christiaen et al., 2008; Davidson and Levine, 2003; Davidson et al., 2006; Davidson et al., 2005; Satou et al., 2004; Stolfi et al., 2010)). TVCs activate conserved cardiac markers, including *Hand*, *Gata4/5/6* and *Nk4/Nkx2-5*, and migrate as bilateral polarized pairs of cells, until the left and right pairs meet at the ventral midline and begin to divide asymmetrically along the mediolateral axis (Figure 1A; (Christiaen et al., 2008; Davidson et al., 2005; Satou et al., 2004; Stolfi et al., 2010)). The first oriented asymmetric divisions produce small median first heart precursors (FHPs), and large lateral second trunk ventral cells (STVCs), which specifically activate *Tbx1/10* expression (Davidson et al., 2005; Stolfi et al., 2010; Wang et al., 2013). STVCs later divide again to produce small median second heart precursors (SHPs), and large

lateral atrial siphon muscle founder cells (ASMFs), which activate *Ebf* (aka *COE*; (Razy-

Krajka et al., 2014; Stolfi et al., 2010; Stolfi et al., 2014c)). The transcription factors

Hand-related (Hand-r)/Notrlc, which is expressed in the TVCs and maintained in the

STVCs and ASMFs after each division, and Tbx1/10 are required for *Ebf* activation in the

ASMFs, whereas Nk4/Nkx2-5 represses *Tbx1/10* and *Ebf* expression in the second heart

precursors (SHPs)(Razy-Krajka et al., 2014; Tolkin and Christiaen, 2016; Wang et al.,

2013). Conversely, Tbx1/10 and Ebf inhibit cardiac markers, and likely determinants,

such as *Gata4/5/6* and *Hand* (Razy-Krajka et al., 2014; Stolfi et al., 2010; Stolfi et al.,

2014a; Wang et al., 2013). These regulatory cross-antagonisms presumably underlie the

transition from transcriptionally primed multipotent progenitors to separate fate-

restricted precursors, by limiting the deployment of the heart- and pharyngeal-muscle-

specific programs to their corresponding specific precursors (Kaplan et al., 2015).

Here, we identify regulatory mechanisms ensuring the emergence of diverse fate-

restricted precursors from multipotent progenitors. We show that differential

FGF/MAPK signaling, feed-forward regulatory mechanisms and coupling with the cell

cycle control the spatially restricted activation of *Tbx1/10* and *Ebf*, successively, thus

permitting the emergence of both first and second heart precursors, and

ASM/pharyngeal muscle precursors from common multipotent progenitors.

Results

MAPK signaling is active in the multipotent cardiopharyngeal progenitors and progressively restricted to the pharyngeal muscle precursors.

During the earliest stages of cardiopharyngeal development in ascidians, multipotent progenitors co-express early regulators of both the heart and ASM programs, a phenomenon referred to as multilineage transcriptional priming, (Razy-Krajka et al., 2014; Stolfi et al., 2014b). Subsequent regulatory cross-antagonisms lead to the segregation of these distinct cardiopharyngeal programs to their corresponding fate-restricted progenitors (Stolfi et al., 2010; Wang et al., 2013); reviewed in (Kaplan et al., 2015)). ASM-specific expression of *Ebf* is necessary and sufficient to terminate the heart program and impose a pharyngeal muscle fate (Razy-Krajka et al., 2014; Stolfi et al., 2010). Antagonistic Tbx1/10 and Nk4 activities determine ASM-specific *Ebf* activation (Wang et al., 2013); however, the symmetry-breaking events leading to cardiopharyngeal mesoderm patterning and ASM-specific expression of *Ebf* remain unknown. In particular, we surmised that differential signaling inputs determine the stereotyped spatio-temporal patterning of early cardiopharyngeal progenitors.

The *Ciona* homologs of specific FGF/MAPK pathway components, including *FGF receptor substrate 2/3* (*Frs2/3*; (Gotoh et al., 2004))*, Ets.b*, and *Fgf4/5/6*, are preferentially expressed in the TVCs, in the STVCs and in the ASMFs as cells transition from a multipotent progenitor state to distinct heart vs. ASM fate-restricted precursors (Razy-Krajka et al., 2014). This patterned expression of MAPK effector genes prompted us to evaluate a role for FGF/MAPK pathway in cardiopharyngeal fate decisions.

We first used an antibody specific to the dual phosphorylated form of Extracellular Regulated Kinase (dpERK) to monitor Mitogen Activated Protein Kinase (MAPK) activity in the cardiopharyngeal mesoderm. We detected dpERK staining in the newly

born TVCs, marked by the B7.5-lineage-specific *Mesp>H2B::mCherry* transgene, as previously observed (Davidson et al, 2006). We also detected weaker but persistent dpERK staining in the TVCs during migration (Figs. 1 and S1). Following the first and second asymmetric divisions of the TVCs and STVCs, dpERK staining was successively restricted to the more lateral STVCs and ASMFs, respectively (Figures 1A, B; S1).

The canonical FGF/Ras/MEK/ERK pathway is necessary and sufficient to promote pharyngeal muscle specification in the cardiopharyngeal lineage.

This exclusion of MAPK activity from the medial first and second heart precursors opened the possibility that differential ERK activity is required for proper STVC and ASMF vs. heart precursors fate decisions. In *Ciona*, signaling through the sole FGF receptor (FGFR) governs ERK activity in several developmental processes, including neural induction (Bertrand et al., 2003; Hudson et al., 2003) and central nervous system patterning (Haupaix et al., 2014; Racioppi et al., 2014; Stolfi et al., 2011; Wagner et al., 2014), early endomesoderm and notochord fate specification (Imai et al., 2002; Picco et al., 2007; Shi and Levine, 2008; Shi et al., 2009; Yasuo and Hudson, 2007). Notably, FGF/MAPK signaling is active in the only *Mesp+* cardiogenic B7.5 blastomeres (Imai et al., 2006; Shi and Levine, 2008), where targeted misexpression of a dominant negative form of FGFR (dnFGFR) using a B7.5-lineage-specific *Mesp* driver blocks TVC induction (Davidson et al., 2006). We used a TVC-specific *FoxF* enhancer (*FoxF(TVC):bpFOG-1>dnFGFR::mCherry*, hereafter called *FoxF>dnFGFR*; (Beh et al., 2007)), to bypass early effects and achieve later misexpression of dnFGFR in the TVCs and their progeny. *FoxF>dnFGFR* prevented neither TVC migration nor asymmetric divisions, but it abolished the expression of both *Tbx1/10* in the STVCs and *Ebf* in the ASMFs (Figure 1C). This data indicate that FGF/MAPK signaling is required in the cardiopharyngeal

progenitors and/or their progeny for ASM fate specification, beyond the initial TVC

induction.

Upon FGF/MAPK-dependent induction, the TVCs express *Hand-related/Hand-r*

(renamed after *Notrlc/Hand-like*; (Christiaen et al., 2008; Davidson and Levine, 2003;

Davidson et al., 2006; Satou et al., 2004; Stolfi et al., 2014c; Woznica et al., 2012)),

which encodes a basic helix-loop-helix (bHLH) transcription factor necessary for *Ebf*

expression in the ASMFs (Razy-Krajka et al, 2014). Moreover, the *Hand-r* TVC enhancer

contains putative Ets1/2 binding sites, which are necessary for reporter gene expression,

and presumably mediate the transcriptional inputs of FGF/MAPK (Woznica et al., 2012).

Since *Hand-r* and *FoxF* expressions start at approximately the same time in newborn

TVCs, we used *FoxF>dnFGFR* to test whether the maintenance of *Hand-r* expression in

migratory TVCs requires prolonged FGF/MAPK inputs after initial TVC induction.

FoxF>dnFGFR inhibited *Hand-r* expression in late TVCs (Figure 1C), indicating that

sustained *Hand-r* expression requires continuous FGF/MAPK signaling.

To test whether the spatial restriction of MAPK activity explains the patterned

expressions of *Hand-r*, *Tbx1/10* and *Ebf* following asymmetric cell divisions, we used

gain-of-function perturbations to force FGF/MAPK activity throughout the

cardiopharyngeal mesoderm and assayed gene expression (Figure 2). We focused on the

canonical FGF/MAPK pathway where signal transduction involves Ras, Raf, MEK and

ERK downstream of FGFR and upstream of transcriptional effectors (Lemmon and

Schlessinger, 2010). We first used M-Ras^{G22V}, a defined constitutively active form of M-

Ras, which mediates FGF signaling in *Ciona*, where other classical *Ras* genes are missing

(Keduka et al., 2009). To assay the transcriptional consequences of forced M-Ras activity

in the cardiopharyngeal lineage, we first focus on *Htr7* and *Tbx1/10* expression following

the first asymmetric TVC division in 15 hours post-fertilization (hpf) embryos. *Htr7*

encodes a *trans*-membrane G-protein coupled receptor and, like *Hand-r*, its expression

and maintenance in the TVCs require MAPK activity (Figure S2; (Razy-Krajka et al.,

2014)), and become restricted to the lateral STVC following asymmetric division.

However, *Htr7* mRNAs appear to be cleared more rapidly from the FHPs, making the

patterned expression easier to analyze than that of *Hand-r* (Figures 2 and 3D; (Razy-

Krajka et al., 2014)). Importantly, misexpression of M-Ras^{G22V} using the TVC-specific

FoxF enhancer did not alter the cell division patterns, allowing us to identify large lateral

STVCs and small median FHPs. Compared to control embryos overexpressing wild-type

M-Ras (M-Ras^{WT}), TVC-specific gain of M-Ras function caused both persistent *Htr7*

expression and ectopic activation of *Tbx1/10* in the first heart precursors following

asymmetric divisions. Similarly, *FoxF>M-Ras^{G22V}*-expressing 18hpf larvae displayed

ectopic *Ebf* activation throughout the cardiopharyngeal mesoderm (Figure 2B, C). These

results indicated that forced M-Ras activation throughout the cardiopharyngeal lineage

is sufficient to ectopically activate STVC and ASMF markers. This is consistent with the

idea that spatially defined signaling upstream of M-Ras restricts MAPK activity, thus

localizing STVC- and ASM-specific gene activities.

To further probe the signal transduction pathway, we engineered a constitutively

active version of the *Ciona* Mek1/2 protein by introducing phosphomimetic mutations of

two conserved serine residues in the catalytic domain, as previously shown for the

mammalian homolog (Cowley et al., 1994; Mansour et al., 1994). Early misexpression of

this Mek^{S220E,S216D} construct in the B7.5 blastomeres using a *Mesp* enhancer caused

ectopic TVC induction, mimicking the effects of published gain of Ets1/2 function

(Figure S3; (Davidson et al., 2006)). Mirroring the effects of M-Ras^{G22V} gain-of-function

experiments, TVC-specific misexpression of Mek^{S220E,S216D} using the *FoxF* enhancer also

caused ectopic expression of *Htr7* and *Tbx1/10*, and *Ebf* in 15 and 18hpf larvae,

respectively (Figure 2B, C). Taken together, these results indicate that activity of the

canonical FGF-Ras-MEK-ERK pathway is progressively restricted to the STVC and

ASMF, and is both necessary and sufficient to promote STVC- and ASMF-specific gene

expressions.

Continuous FGF/MAPK activity is required for the successive activations of *Tbx1/10* and *Ebf*.

FGF/MAPK signaling is sufficient and necessary to maintain *Hand-r* expression in

late TVCs (Figure 1), and *Hand-r* is necessary for *Ebf* expression in the ASMF (Razy-

Krajka et al., 2014). Therefore, it is possible that later FGF/MAPK signaling is

dispensable for *Tbx1/10* and *Ebf* activation and ASM specification, as long as STVC and

ASMF cells inherit sustained levels of *Hand-r* mRNAs and/or proteins. To disentangle

late from early requirements of FGF/MAPK signaling for TVC progeny specification, we

incubated embryos at different stages with the MEK/Mapkk inhibitor U0126, which

abolishes dual ERK phosphorylation and the initial MAPK-dependent TVC induction in

Ciona embryos (Figure S1; (Davidson et al., 2006; Hudson et al., 2003)). MEK inhibition

during TVC migration (i.e. between 9.5 and 12.5 hpf, Figure 3A) blocked the expression

of *Hand-r* and *Htr7* in late TVCs (Figure 3B, E). Similarly, U0126 treatments in late

TVCs, and through the first asymmetric division (i.e. between 12 and 15 hpf, Figure 3A)

blocked both the maintenance of *Hand-r* and *Htr7*, and the activation of *Tbx1/10* in the

STVCs (Figure 3C, D, F, G). Finally, MEK inhibition in late STVCs and through

asymmetric divisions (i.e. between 15 and 18 hpf) blocked the ASMF-specific expression

of *Ebf* (Figure 3H). These results indicate that continuous MEK activity is required

throughout cardiopharyngeal development to successively activate TVC-, STVC-, and

ASMF-expressed genes.

Since *Ebf* expression is maintained for several days in the ASMF derivatives as they

differentiate into body wall and siphon muscles (Razy-Krajka et al., 2014), we tested

whether continued MEK activity is also required for the maintenance of *Ebf* expression

past its initial onset and cells' commitment to an ASM fate. Using both regular and
intron-specific antisense probes, which specifically detect nascent transcripts (Wang et
al., 2013), we showed that later MEK inhibition (i.e. U0126 incubation between 17 and
20 hpf) did not block the maintenance of *Ebf* transcription in the ASMPs (Figure 3I, J).
This indicates that sustained MEK activity is required until the onset of *Ebf* expression,
but not beyond, the maintenance of *Ebf* expression during ASM development is
independent of MAPK.

Since U0126 treatments affect the whole embryo, we sought to further confirm the
later roles for FGF/MAPK signaling specifically in the cardiopharyngeal mesoderm. To
this aim, we used an STVC-specific enhancer from the *Tbx1/10* locus (termed *T12*; Figure
3K, L; (Tolkin and Christiaen, 2016); Racioppi et al., in preparation) to drive expression
of either dnFGFR or the constitutively active M-Ras^{G22V} starting at ~14hpf, and assayed
Ebf expression at 18hpf (Figure 3K, L). These perturbations minimally affected the cell
division patterns, such that cells corresponding to FHP, SHP and ASMF could be
identified by their position relative to the midline (Figure 3K). M-Ras^{G22V} misexpression
caused conspicuous ectopic *Ebf* expression in the SHPs, whereas dnFGFR-mediated
inhibition of MAPK activity blocked *Ebf* activation in the lateral ASMFs. These results
support the notion that localized FGF/MAPK activity is necessary and sufficient for
ASMF-specific expression of *Ebf*.

Coherent feed-forward circuits for cardiopharyngeal mesoderm patterning
and ASM fate specification.

The above results indicate that *Hand-r*, *Tbx1/10* and *Ebf* require ongoing
FGF/MAPK activity for their successive activations in the TVCs, STVCs and ASMFs,
respectively. We previously showed that RNAi and/or CRISPR-mediated inhibition of

either *Hand-r* or *Tbx1/10* function blocks *Ebf* activation in the ASMFs, where both *Hand-r* and *Tbx1/10* expressions are maintained (Razy-Krajka et al., 2014; Tolkin and Christiaen, 2016; Wang et al., 2013). Therefore, observations such as the loss of *Ebf* expression upon *FoxF>dnFGFR* electroporation could be due to an early loss of *Hand-r* and/or *Tbx1/10*. We used epistasis assays to systematically test whether early regulators mediate the effects of FGF/MAPK on later gene expression and ASM fate specification, or whether FGF/MAPK signaling acts both upstream and in parallel to early regulators in a more complex regulatory circuit.

We first revisited the regulatory relationships between FGF/MAPK, *Hand-r* and *Tbx1/10* in late TVCs and early STVCs. We validated single guide RNAs (sgRNAs) for CRISPR/Cas9-mediated mutagenesis of *Hand-r* (Table S1; (Gandhi et al., 2017)), and determined that *Hand-r* function is necessary for *Tbx1/10* activation in the STVCs (Figure 4A). Co-expression of a modified *Hand-r* cDNA containing wobble base mutations that disrupt the sgRNA protospacer adjacent motif (PAM; Hand-r[PAMmis]) rescued *Tbx1/10* expression in the STVCs, indicating that *Tbx1/10* down-regulation in this CRISPR "background" is specifically due to *Hand-r* loss-of-function (Figure 4A). To further probe if *Hand-r* activity is necessary for FGF/MAPK-dependent *Tbx1/10* expression, we used gain of M-Ras function in a *Hand-r* CRISPR "background". Whereas, misexpression of the constitutively active M-Ras[G22V] caused ectopic *Tbx1/10* expression, concomitant loss of *Hand-r* function diminished both endogenous and ectopic *Tbx1/10* expression in the STVC and FHP, respectively (Figure 4A). Although, remaining ectopic activation could still be observed, possibly because M-Ras[G22V] could boost *Hand-r* expression in heterozygous cells where CRISPR/Cas9 disrupted only one copy of the gene. This data indicate that *Hand-r* is necessary for FGF/MAPK-induced activation of *Tbx1/10*.

To further probe the epistatic relationships between *Hand-r* and MAPK signaling

upstream of *Tbx1/10*, we attempted to rescue *Tbx1/10* expression in U0126-treated

embryos, by over-expressing *Hand-r* with the TVC-specific *FoxF* enhancer. Neither did

Hand-r over-expression cause ectopic *Tbx1/10* activation (in the FHPs), nor was it

sufficient to rescue *Tbx1/10* expression in 15hpf STVCs (Figure 4B). Taken together,

these data indicate that both Hand-r and MAPK activities are required to activate

Tbx1/10 in the STVCs. These results also imply that MAPK signaling is restricted to the

STVC independently of Hand-r activity, which suffice to explain the STVC-specific

activation of *Tbx1/10*.

Next, we investigated the epistatic relationship between FGF/MAPK, *Hand-r*, and

Tbx1/10 upstream of *Ebf* in the ASMFs. We first used previously validated CRIPSR/Cas9

reagents targeting the *Tbx1/10* coding region (Tolkin and Christiaen, 2016), to confirm

that B7.5-lineage-specific loss of *Tbx1/10* function inhibited *Ebf* activation, and verified

that this effect could be rescued by over-expression of a CRISPR/Cas9-resistant *Tbx1/10*

cDNA, expressed with a minimal TVC-specific *FoxF* enhancer (Figure 4C;

Tbx1/10[PAMmis]). In these rescue experiments, we observed ectopic *Ebf* activation in the

SHP, as previously described when driving Tbx1/10 expression with a TVC-specific *FoxF*

enhancer (Wang et al., 2013). As explained below, this ectopic activation could be

attributed to a precocious expression of *Ebf* in the STVCs (Figure 4E). To test whether

Tbx1/10 was also required for ectopic *Ebf* expression in response to MAPK activation, we

combined CRISPR/Cas9-mediated *Tbx1/10* knockout with constitutive MAPK activation

using the M-Ras[G22V] mutant and observed a significant inhibition of both endogenous

and ectopic *Ebf* expression in the 18hpf ASMF and SHP, respectively (Figure 4C). Taken

together, these results show that *Tbx1/10* function is necessary for FGF/MAPK-induced

expression of *Ebf* in the ASMFs.

306 To further test whether *Tbx1/10* acts in parallel and/or downstream of MAPK to

307 activate *Ebf*, we combined gain of *Tbx1/10* function with perturbations of FGF/MAPK

308 signaling and assayed *Ebf* expression. We realized that *FoxF*-driven misexpression of

309 Tbx1/10 caused precocious *Ebf* activation in 15hpf STVCs (Figure 4D, E). This precocious

310 expression remained remarkably patterned, suggesting that STVC-restricted FGF/MAPK

311 activity prevented *Ebf* expression in the dpERK-negative, small median FHPs (Figures

312 1B, 4E, S1). Indeed, co-expression of both Tbx1/10 and M-Ras^{G22V} caused both

313 precocious and ectopic *Ebf* expression in the 15hpf medial and lateral TVC derivatives,

314 which would be FHPs and STVCs in control embryos, respectively. This data confirms

315 that Tbx1/10 misexpression does not suffice to cause ectopic *Ebf* expression in the FHPs,

316 because the latter presumably lack FGF/MAPK activity, as is the case in control embryos.

317 U0126-mediated MEK inhibition from 12 to 15hpf, i.e. after the onset of

318 *FoxF>Tbx1/10* misexpression, further confirmed that MAPK activity is required in

319 parallel to Tbx1/10 for precocious *Ebf* activation in 15hpf STVCs (Figure 4D, E). Taken

320 together, these results indicate that Tbx1/10 and MAPK are both required to activate *Ebf*

321 in the cell cycle following that of *Tbx1/10* onset.

323 Since *Hand-r* expression is maintained in the ASMF, and CRISPR/Cas9- or RNAi-

324 mediated *Hand-r* knockdown blocked both *Tbx1/10* (Figure 4A) and *Ebf* expression

325 (Razy-Krajka et al., 2014), we reasoned that Hand-r could also act both upstream and in

326 parallel to Tbx1/10 for *Ebf* activation. To test this possibility, we assayed *Ebf* expression

327 in 18hpf ASMF following defined perturbations of Hand-r and Tbx1/10. As expected,

328 CRISPR/Cas9-mediated *Hand-r* mutagenesis strongly inhibited *Ebf* expression, and this

329 effect could be rescued by a CRISPR-resistant *Hand-r* cDNA (Figure 4F). To test

330 whether this effect was mediated by a loss of *Tbx1/10* expression, we attempted to rescue

331 the Hand-r loss-of-function by over-expressing Tbx1/10 using the *FoxF* enhancer. As

explained above, *FoxF*-mediated Tbx1/10 misexpression caused precocious and ectopic

Ebf expression in larvae co-electroporated with control sgRNAs (Figure 4D, E, F). By

contrast, combining loss of Hand-r function with Tbx1/10 misexpression inhibited both

the endogenous and ectopic *Ebf* expression (Figure 4F), indicating that Hand-r is also

required in parallel to Tbx1/10 for *Ebf* activation in the ASMFs.

Taken together, these analyses of the epistatic relationships between FGF/MAPK

signaling, *Hand-r*, *Tbx1/10* and *Ebf* suggest that coherent feed-forward circuits govern

the sequential activation of *Hand-r*, *Tbx1/10* and *Ebf* in response to continuous but

progressively restricted FGF/MAPK inputs (Figure 4G), thus linking spatial patterning to

the temporal deployment of the regulatory cascade leading to localized *Ebf* activation

and pharyngeal muscle specification.

The cell cycle entrains the temporal deployment of the cardiopharyngeal

gene regulatory network.

In principle, the feed-forward circuit described above is sufficient to explain the

successive activations of *Hand-r*, *Tbx1/10* and *Ebf*. However, *Tbx1/10* and *Ebf* do not

turn on until after oriented and asymmetrical divisions of the TVCs and STVCs,

respectively. Notably, even when we misexpressed Tbx1/10 in the TVCs, *Ebf* was

activated only after TVC division and in the lateral-most cells, where FGF/MAPK

signaling is normally maintained (Figures 1B, 4E). This sequence of events -divisions

followed by gene activation- is paramount in the cardiopharyngeal mesoderm, as it

permits the birth of first and second heart precursors, whose fates are antagonized by

Tbx1/10 and Ebf (Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et al., 2013).

Therefore, we sought to investigate the role(s) of the cell cycle in controlling the timing of

Tbx1/10 and *Ebf* activations.

We first evaluated the effects of cytochalasin B, a classic inhibitor of cytokinesis

widely used to study cell fate specification in ascidians (Figure 5A; (Whittaker, 1973)).

Treatments starting before TVC divisions (12 hpf) did not block *Tbx1/10* or *Ebf*

expression in embryos fixed after their normal onset at either 16 or 19hpf, respectively

(Figure 5B). Similarly, treatment starting between the first and second asymmetric

divisions (15hpf) did not block localized *Ebf* expression at 19hpf (Figure 5B). This

indicates that *Tbx1/10* and *Ebf* activations occur by default in the absence of cytokinesis,

most likely because FGF/MAPK signaling persists throughout the shared cytoplasm.

This data thus illustrates how the spatial restriction of FGF/MAPK signaling, following

cell divisions, leads to the localized activations of *Tbx1/10* and *Ebf*, and permits the

emergence of first and second cardiac precursors.

Cytochalasin treatments usually lead to the formation of polynucleated cells (e.g.

Figure 5B, middle panel), because the cell cycle and nucleokinesis continue in these

artificial endoreplicating cells. To alter cell cycle progression more comprehensively, and

specifically in the cardiopharyngeal lineage, we used genetically encoded inhibitors of

cell cycle transitions: Cdkn1b.a and Cdkn1b.b (also known as Noto16), the ortholog of

which is a potent inhibitor of the G1/S transition in the ascidian species *Halocynthia*

roretzi (Kuwajima et al., 2014), and Wee1, a G2/M inhibitor, as previously described

(Dumollard et al., 2017). We used the TVC-specific *FoxF* enhancer to misexpress these

negative regulators of cell cycle progression, monitored cell divisions and assayed

Tbx1/10 expression at 15hpf, when control TVCs have divided and the lateral-most

STVCs normally express *Tbx1/10*. Each perturbation efficiently inhibited TVC divisions,

such that only two cells were visible on either side of the embryos (Figure 5C). In these

delayed TVCs, *Tbx1/10* expression was strongly reduced compared to control STVCs

(Figure 5C; compare to Figure 4A, B). However, approximately 40% of the delayed TVCs

expressed *Tbx1/10* to variable extents. This suggests that the cardiopharyngeal

regulatory network can qualitatively unfold independently of cell cycle progression, but

the latter is necessary for *Tbx1/10* expression to its wild-type levels.

We next used the STVC-specific *Tbx1/10 T12* enhancer, to misexpress Cdkn1b.a,

Noto16 and Wee1, and assay *Ebf* expression at later stages. Inhibitors of the G1/S

transition failed to block STVC divisions (data not shown), most likely because *T12*-

driven products did not accumulate quickly enough to interfere with the G1/S transition

in STVCs (this cell cycle lasts only ~2 hours compared to ~6 hours for the TVC

interphase), suggesting that the G1 phase is too short for *T12*-driven gene products to

accumulate before the G1/S transition. Therefore, we focused the analyses of *Ebf*

response to cell cycle perturbations on misexpression of the G2/M inhibitor Wee1.

Preliminary analyses of 18hpf larvae, fixed approximately 2 hours after the documented

onset of *Ebf* expression in ASMFs (Razy-Krajka et al., 2014), indicated that *Ebf* can turn

on in arrested STVCs that failed to divide upon Wee1 misexpression (Figure 5D).

Because ~30% of the embryos showed variable expression, as was the case for

Tbx1/10 in the previous experiment, we reasoned that perturbations of the G2/M

transition could alter the dynamics of *Ebf* upregulation. We investigated this possibility

using embryos fixed every 30 minutes between 15.5hpf and 18hpf, when cells transition

from a late *Tbx1/10+*; *Ebf-* STVC state to a committed *Ebf+*, *Mrf+* ASMF state (Razy-

Krajka et al., 2014; Wang et al., 2013). First, we observed that the proportion of embryos

with conspicuous ASMFs increased from ~20% to >90% between 15.5 and 16.5 hpf in

control embryos (Figure 4E). By contrast, Wee1-expressing cells had divided in only

~35% of the embryos by 16.5hpf, and that proportion gradually increased to ~70% by

18hpf (Figure 4E), indicating that Wee1 misexpression strongly delays cell cycle

progression, blocking cell divisions in a substantial fraction of embryos.

Focusing on ASMFs, we found that the proportion of *Ebf+* cells in control embryos

progressively increased from ~20% showing "weak" expression at 15.5hpf to >90%

showing "strong" expression by 18hpf (Figure 5F; see Figure 5D for examples of "weak"

and "strong" expression). This semi-quantitative analysis revealed an under-appreciated

dynamic at the onset of *Ebf* expression, which appears to take at least one hour to be

"strongly" expressed in >75% of newborn ASMFs (Figure 4F).

To evaluate the impact of Wee1-induced mitosis inhibition on *Ebf* accumulation, we

focused on undivided STVCs at each time point (hence the lower numbers in Figure 4F

compare to Figure 4E). By 17hpf, wee1-expressing delayed STVCs showed "strong" *Ebf*

expression in comparably high proportions of embryos. However, these proportions

were significantly lower at 16 and 16.5hpf (Chi-square tests, *P*=0.002 and *P*=0.0003,

respectively), with ~1.5 and ~1.2 times less "strongly" expressing cells than in the control

distributions (hypergeometric tests, *P*=0.0005 and *P*=0.0001, respectively). These semi-

quantitative data suggests that the cardiopharyngeal network can eventually unfold and

lead to high levels of *Ebf* expression independently of cell divisions, albeit with a delay

revealing that cell divisions probably entrain *Ebf* upregulation in early ASMFs.

Transition from a MAPK-dependent to a MAPK-independent and

autoregulative mode of *Ebf* expression in early ASMFs.

Given the semi-quantitative nature of our analysis, and the relatively subtle effects

observed on *Ebf* dynamics, we sought to further probe the mechanisms that regulate the

onset and upregulation of *Ebf* expression in early ASMFs, and the biological significance

for cell-fate specification. Since we observed a progressive accumulation of *Ebf* mRNAs,

and a transition from a MAPK-dependent onset to a MAPK-independent maintenance of

Ebf transcription (Figure 3I, J), we reasoned that the window of MAPK-dependence

might coincide with the accumulation of *Ebf* mRNAs between 16 and 17hpf. To test this

possibility, we treated embryos with the MEK inhibitor U0126 at successive time points,

assayed ongoing transcription using intronic probes and counted the numbers of *Ebf*

transcribing cells (Figure 6A). This analysis revealed that *Ebf* transcription gradually lost

its sensitivity to MAPK inhibition between 16 and 17hpf, i.e. during the first hour of the

ASMF cycle when *Ebf* mRNAs normally accumulate (as shown in Figure 5E, F).

Because *Ebf* transcription becomes independent from MAPK by the time *Ebf* mRNA

have accumulated to "high" levels, and because *Ebf* expression lasts for several days in

the progeny of the ASMFs, we reasoned that autoregulation might suffice to maintain

high levels of *Ebf* mRNA past the MAPK-dependent onset. To test this possibility, we

misexpressed the Ebf coding sequence using the STVC-specific *T12* enhancer as

described (Tolkin and Christiaen, 2016). Assaying endogenous *Ebf* transcription using

intronic probes demonstrated that, in addition to its normal expression in the ASMFs,

Ebf misexpression caused precocious and ectopic activation of the endogenous locus in

the STVCs, and in the MAPK-negative SHPs, respectively (Figure 6C-F). This result

suggests that *Ebf* transcription bypasses both requirements for cell-division coupling and

MAPK inputs if high levels of *Ebf* gene products are present in the cell.

We reasoned that, if high levels of *Ebf* expression can promote its own transcription

independently of MAPK signaling, then Ebf misexpression should be sufficient to rescue

a chemical inhibition of MAPK at a critical stage. We tested this possibility by combining

Ebf misexpression using the STVC-specific *T12* enhancer and U0126 treatments starting

at 16hpf, which normally block *Ebf* expression (Figure 6A, D-F). We observed that

transcription of the endogenous *Ebf* locus became independent of early MAPK activity

upon misexpression of an Ebf cDNA, further supporting the notion that high levels of

Ebf expression suffice to maintain *Ebf* transcription independently of MAPK activity.

A potentially important implication of this transient MAPK-dependence of is to

render *Ebf* expression initially reversible. For instance, *Ebf* occasionally turns on

precociously in the STVCs of a small proportion of embryos (Figure S4). Given the

powerful anti-cardiogenic effects of Ebf (Razy-Krajka et al., 2014; Stolfi et al., 2010),

persistent *Ebf* expression would have dramatic consequences for SHP development

(Wang et al., 2013). However, because MAPK activity is excluded from the SHPs, and the

early phase of *Ebf* expression depends upon continuous MAPK activity, we surmise that

Ebf cannot be maintained in the SHPs. For instance, when embryos from the same

electroporated batch were fixed at the time of early U0126 treatment (i.e. 15.75 and

16.25hpf) and ~4 hours later, at 20hpf, and assayed for *Ebf* transcription using intronic

probes, initially wild-type patterns of *Ebf* transcription could not be maintained (Figure

S5A). This suggests that, although *Ebf* can be activated precociously in a MAPK-

dependent manner, its expression shuts off in the SHPs upon MAPK inhibition following

STVC division.

We further addressed the interplay between cell division, MAPK signaling and *Ebf*

expression. We reasoned that, if cell divisions entrain Ebf accumulation and the

transition to a MAPK-independent autoregulative mode, then delaying STVC divisions

should extend the period of MAPK-dependent *Ebf* transcription. We tested this

possibility by expressing Wee1 under the control of the STVC-specific *T12* enhancer, and

treated embryos with U0126 at 17hpf, which inhibited the maintenance of *Ebf*

transcription in only 15% to 20% of the control embryos (Figures 6A, S5B). The

proportion of embryos showing U0126-sensitive *Ebf* transcription increased to almost

50% upon *T12>Wee1* expression (Figure S5B), which is consistent with our hypothesis

that inhibiting the G2/M transition delayed the accumulation of *Ebf* gene products thus

postponing the transition from a low level/MAPK-dependent to an high level/MAPK-

independent and self-activating mode of *Ebf* regulation.

Taken together these data lead us to propose a model for *Ebf* regulation whereby

Hand-r, Tbx1/10, ongoing MAPK signaling and cell-cycle-regulated transcriptional

input(s) govern the onset and initial accumulation of *Ebf* gene products during the first

hour of the ASMF cycle, whereas the maintenance of *Ebf* expression relies primarily on

MAPK-independent autoactivation, following initial accumulation (Figure 7).

Discussion

Here, we demonstrated that the progressive restriction of FGF/MAPK signaling

follows asymmetric cell divisions of multipotent progenitors and patterns the ascidian

cardiopharyngeal mesoderm in space and time. This leads to the localized expression of

Hand-r, *Tbx1/10* and *Ebf* in fate-restricted pharyngeal muscle precursors, and their

concomitant exclusion for first and second heart precursors. We show that coherent

feedforward circuits encode the successive activations of *Hand-r*, *Tbx1/10* and *Ebf*,

whereas cell divisions entrain the progression of this regulatory sequence and thus

define the timing of gene expression. Finally, we provide evidence that the initiation of

Ebf expression depends on MAPK activity in early ASMF, until Ebf accumulation permits

MAPK-independent auto-activation. Given the potent anti-cardiogenic, and pro-

pharyngeal muscle effects of Ebf (Razy-Krajka et al., 2014; Stolfi et al., 2010), we surmise

that the latter switch corresponds to the transition from a cardiopharyngeal multipotent

state to a committed pharyngeal muscle identity.

Spatial patterning by localized maintenance of FGF/MAPK signaling.

Our results demonstrate that MAPK signaling is maintained only in the lateral-most

daughter cells following each asymmetric division of multipotent cardiopharyngeal

progenitors - the TVCs and STVCs. This asymmetric maintenance is necessary and

sufficient for the progressive and localized deployment of the pharyngeal muscle

network. Notably, the TVCs themselves are initially induced by similar polarized

FGF/MAPK signaling coincidental to asymmetric cell divisions of their mother cells, aka

the B8.9 and B8.10 founder cells (Davidson et al., 2006). Detailed analyses have since

indicated that asymmetrical maintenance of sustained FGF/MAPK signaling involves

intrinsic Cdc42-dependent polarity of the founder cells, which promotes polarized cell-

matrix adhesion of the prospective TVC membrane to the ventral epidermis. The latter

differential integrin-mediated adhesion promotes localized MAPK activation, leading to

TVC induction (Cooley et al., 2011; Norton et al., 2013). It has been proposed that

adhesion- and caveolin-dependent polarized FGFR recycling during mitosis accounts for

the localized activation of MAPK in the prospective TVCs (Cota and Davidson, 2015).

Whereas similar mechanisms could in principle account for asymmetric maintenance of

FGF/MAPK signaling in STVCs and ASMFs, this has not been formally tested and there

are notable differences opening the possibility that other mechanisms may be at work:

during TVC induction, MAPK signaling is maintained in the smaller daughter cell that

contacts the epidermis, whereas in the following divisions, MAPK activity persists in the

larger daughter cells and all cells maintain contact with the epidermis (Nicole Kaplan

and Lionel Christiaen, data not shown). Moreover, using an FGFR::mKate2 fusion

protein similar to that used in previous studies, we could not observed a marked

polarized distribution of FGFR molecules to the lateral-most cells (the STVCs and

ASMFs; Yelena Bernadskaya and Lionel Christiaen, data not shown). However, the fact

that constitutively active forms of M-Ras and Mek1/2 were sufficient to bypass the loss of

MAPK activity, and impose pharyngeal muscle specification, indicates that differential

FGF/MAPK activity is regulated upstream of M-Ras. Further work is needed to elucidate

the cellular and molecular mechanisms governing the spatiotemporal patterns of

FGF/MAPK signaling in the cardiopharyngeal mesoderm. In particular, it will be

important to disentangle the relative impacts of extrinsic (i.e. tissues, contacts) vs.

intrinsic (i.e. asymmetric cell division) effects onto FGF/MAPK signaling and the

downstream transcriptional inputs.

Transcriptional effects of differential FGF/MAPK signaling.

Because differential FGF/MAPK signaling rapidly impacts cell-specific gene

expression, we surmise that transcriptional effectors are dynamically regulated. For

instance, even though we have not formally identified the downstream DNA-binding

transcription factor (see discussion below), it would be conceivable that the

phosphorylated forms of either transcriptional effector persist through cell division upon

maintenance of FGF/MAPK activity. However, we have shown that continuous MAPK

activity is needed following each division. Therefore, we must invoke elusive

phosphatase activities, such as dual-specificity phosphatases (DUSPs; (Patterson et al.,

2009), which would reset transcriptional effectors to a dephosphorylated state, thus

rendering steady-state FGF/Ras/MAPK inputs necessary.

Systematic dephosphorylation of FGF/MAPK transcriptional effectors is likely to be

particularly important for heart fate specification. For instance, whole genome analyses

indicate that heart-specific *de novo* gene expression requires MAPK inhibition (Wang et

al., 2017). Although the molecular mechanisms remain elusive, one simple possibility is

that, lest fate-restricted heart precursors inhibit MAPK activity, they will activate

Tbx1/10 and *Ebf*, which will block the cardiac program (Razy-Krajka et al., 2014; Stolfi et

al., 2010; Wang et al., 2013). Finally, we previously proposed that repressor inputs from

Nk4 are needed in the second heart precursors to avoid ectopic activation of *Ebf* (Wang

et al., 2013). The observation that Nk4 transcripts are detected in all cardiopharyngeal

cells opened the question as to how *Ebf* would escape repression by Nk4 in the ASMFs.

Differential MAPK activity offers an intriguing possibility: for instance, Nk4/Nkx2-5-

mediated repression in other species involves the co-repressor Groucho/TLE (Choi et al.,

1999), which is strongly expressed in the cardiopharyngeal mesoderm (Razy-Krajka et

al., 2014); and, in flies, MAPK-mediated phosphorylation of Groucho inhibits its

repressor function (Cinnamon et al., 2008; Cinnamon and Paroush, 2008; Hasson et al.,

2005). Therefore, it is possible that persistent MAPK signaling dampens Groucho/TLE-

mediated repressive inputs on cell-specific regulatory genes like *Ebf*. Future studies will

determine whether such mechanisms provide bistable switches underlying MAPK-

dependent fate choices in the cardiopharyngeal mesoderm.

Temporal deployment of the pharyngeal muscle network

The localized and successive activation of *Tbx1/10* and *Ebf* in STVCs, and ASMFs,

respectively, are important features of the cardiopharyngeal network that permit the

emergence of diverse cell fates: first and second heart precursors, and atrial siphon

muscle precursors. Experimental misexpression of Ebf throughout the cardiopharyngeal

mesoderm suffice to inhibit heart development (Razy-Krajka et al., 2014; Stolfi et al.,

2010), illustrating how important it is for *Ebf* expression to be restricted to the ASMF,

once the first and second heart precursors are born and have terminated MAPK activity.

Our analyses indicate that the sequential activations of *Hand-r, Tbx1/10* and *Ebf* is

encoded in the feed-forward structure of this sub-circuit, whereas the continuous

requirement for MAPK inputs and their progressive exclusion from heart progenitors

restrict the competence to activate *Tbx1/10* and *Ebf* to the most lateral cells, after each

division. Our model implies that each gene may directly respond to transcriptional

inputs from MAPK signaling. We have not formally identified the transcription factors(s)

that mediate the transcriptional response to FGF/MAPK signaling. However,

multipotent cardiopharyngeal progenitors express *Ets1/2* and *Elk*, two common

transcriptional effectors of FGF/MAPK signaling in *Ciona* (Bertrand et al., 2003;

Christiaen et al., 2008; Davidson et al., 2006; Gainous et al., 2015). Moreover, Ets1/2

has been implicated in the initial FGF/MAPK-dependent induction of multipotent TVCs

(Christiaen et al., 2008; Davidson et al., 2006), and it expression is also progressively

restricted to the lateral-most progenitors following each division (Razy-Krajka et al.,

2014). Taken together, Ets1/2 and, to some extend, Elk are intriguing candidate

transcriptional effectors of FGF/MAPK signaling in cardiopharyngeal development.

The binding preferences of Ets-family factors have been extensively studied in *Ciona*,

and they do not depart markedly from conserved Ets-family binding sites with a GGAW

core (Bertrand et al., 2003; Farley et al., 2015; Farley et al., 2016; Gueroult-Bellone et al.,

2017; Khoueiry et al., 2010). Putative Ets-family binding sites in the TVC-specific *Hand-*

r enhancer are conserved between *Ciona intestinalis* and its sibling species *C. robusta*

and *C. savignyi*, and necessary for its activity in reporter assays (Woznica et al., 2012).

Similarly, minimal STVC and ASM enhancers for *Tbx1/10* and *Ebf*, respectively, contain

conserved putative Ets-family binding sites, although their function has not been tested

((Razy-Krajka et al., 2014; Wang et al., 2013) and data not shown). Taken together, these

observations suggest that the proposed feed-forward sub-circuit involves direct

transcriptional   inputs   from   FGF/MAPK-regulated   Ets-family   factors   on   the

cardiopharyngeal enhancers of *Hand-r*, *Tbx1/10* and *Ebf*.

Whereas the regulatory architecture of the MAPK; Hand-r; Tbx1/10; Ebf sub-circuit

explains the sequence of activation events, it is also crucial for its correct deployment,

and the generation of diverse cell identities, that genes are not fully activated before

successive cell divisions. While divisions are not absolutely required for *Ebf* to eventually

turn on, cell cycle progression appears to entrain the deployment of this network,

especially for *Tbx1/10* and *Ebf* activation in STVCs and ASMFs, respectively. These

observations imply that, while the network can eventually unfold, its intrinsic dynamic is

slower than observed. This allows first and second heart precursors to be born prior to

the onset of *Tbx1/10* and *Ebf*, respectively. The latter sequence is essentially for the heart

progenitors to escape the anti-cardiogenic effects of Tbx1/10 (Wang et al., 2013), and Ebf

(Razy-Krajka et al., 2014).

Initial *Ebf* expression in early ASMFs is also labile and MAPK-dependent for

approximately one hour. This continued requirement for MAPK inputs ensures that, in

the rare instances when *Ebf* expression starts in the multipotent STVC progenitors

and/or expands to the nascent SHPs, inhibition of MAPK shuts off *Ebf* expression before

it reaches the levels needed for commitment to an ASM fate. Indeed, our results indicate

that, once *Ebf* mRNAs have accumulated to high levels, its expression becomes auto-

regulative and MAPK-independent. We surmise that this transition coincides with a

fundamental switch from a multipotent cardiopharyngeal state to a committed

pharyngeal muscle identity.

From this standpoint, the observed entrainment of *Ebf* expression by the cell cycle

can be seen as acceleration of the transition to commitment following asymmetric

division of multipotent progenitors. Although the mechanisms remain elusive, it is likely

that this requires the M/G1 transition, as the G1 phase has been shown to be particularly

conducive to the expression of fate-specific regulators in mammalian pluripotent stem

cells (Dalton, 2015; Pauklin et al., 2016; Pauklin and Vallier, 2013; Soufi and Dalton,

2016).

633   Conserved dual effects of FGF/MAPK signaling on heart development in

634   chordates

Previous studies highlighted how FGF/MAPK signaling is necessary along side Mesp

during early cardiac development in *Ciona* (Christiaen et al., 2008; Davidson, 2007;

Davidson et al., 2006), and how this early requirement also exists in vertebrates (Abu-

Issa et al., 2002; Alsan and Schultheiss, 2002; Barron et al., 2000; Brand, 2003; Reifers

et al., 2000; Zaffran and Frasch, 2002). We now know that these early FGF/MAPK

inputs induce and maintain multipotent cardiopharyngeal states in *Ciona*, including the

Tbx1/10+ multipotent progenitors that eventually produce the second heart lineage

((Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et al., 2013; Wang et al., 2017), and

this study). Similarly, in vertebrates, regulatory interplay between Fgf8 and Fgf10

signaling and Tbx1 is required for development of both pharyngeal arch and second

heart field derivatives, presumably in part by maintaining an undifferentiated and

proliferative state (Abu-Issa et al., 2002; Aggarwal et al., 2006; Brown et al., 2004; Chen

et al., 2009; Hu et al., 2004; Ilagan et al., 2006; Kelly and Papaioannou, 2007; Park et

al., 2006; Park et al., 2008; Vitelli et al., 2002b; Watanabe et al., 2010; Watanabe et al.,

2012). Notably, FGF signaling acts in successive phases, and its inhibition is necessary

for final myocardial specification and differentiation (Hutson et al., 2010; Marques et al.,

2008; Tirosh-Finkel et al., 2010; van Wijk et al., 2009). Conversely, continued FGF

signaling beyond the multipotent mesodermal progenitor stages was shown to promote

smooth muscle and epicardial differential in the heart (Hutson et al., 2010; van Wijk et

al., 2009), and also myoblast specification and/or skeletal muscle differentiation in the

head, with the expression of FGF ligands being maintained in the pharyngeal arches

(Bothe et al., 2011; Buckingham and Vincent, 2009; Michailovici et al., 2015;

Michailovici et al., 2014; von Scheven et al., 2006). Taken together, these and our data

suggest that FGF/MAPK signaling plays evolutionary conserved roles during chordate

cardiopharyngeal development, by promoting the specification of successive mesodermal

and *Tbx1+* multipotent states, and a fate-restricted non-cardiac muscle identity, while

MAPK inhibition is required for myocardial specification and differentiation in the first

and second heart field, successively.

Material and methods

Animals, electroporations, and chemical treatments

Gravid wild *Ciona intestinalis* type A, now called *Ciona robusta* (Pennati et al., 2015),

were obtained M-REP (Carlsbad, CA, USA), and kept under constant light to avoid

spawning. Gametes from several animals were collected separately for *in vitro* cross-

fertilization followed by dechorionation and electroporation as previously described

(Christiaen et al., 2009a, b). Different quantities of plasmids were electroporated

depending on the constructs. Typically, 50 µg of DNA was electroporated for NLS::lacZ

or plain mCherry driving constructs but only 15 µg for *Mesp-1>H2B::mCherry*. For

perturbation constructs, 70 µg were usually electroporated, except for

Mesp>NLS::Cas9::NLS (30 µg) and pairs of U6>sgRNA plasmids (25 µg each). U0126

(Cell Signaling Technology, Danvers, MA) was used at 5µM in artificial seawater from a

stock solution of 20mM in DMSO. Cytochalasin B (Sigma, Saint Louis, MO) was used at

~3 µg/mL from a 10 mg/mL stock solution in DMSO as previously performed (Jeffery et

al., 2008). Control embryos were incubated in parallel with corresponding

concentrations of DMSO alone.

***In situ* hybridization**

In situ hybridizations were carried out essentially as described previously (Christiaen et

al., 2009c; Razy-Krajka et al., 2014), using DIG labeled riboprobes, anti-DIG-POD Fab

fragments (Roche, Indianapolis, IN), and Tyramide Amplification Signal coupled to

Fluorescein (Perkin Elmer, MA). Reporters expressed in the lineage of interest were

marked using anti-ß-galactosidase monoclonal mouse antibody (1:1000; Promega,

Fitchburg, WI) or anti-mCherry rabbit polyclonal antibody (1:500; BioVision 5993-100),

respectively targeted with anti-mouse or anti-rabbit secondary antibody coupled with

Alexa 648 (1:500; Invitrogen, Carlsbad, CA). The different probes used in this study were

described previously (Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et al., 2013).

dpERK/mcherry double fluorescent immunostaining

Samples were fixed, as for *in situ* hybridizations, in MEM-PFA with Tween 20 (0.05%)

but only for 30 minutes at room temperature, washed three times in PBSt (Tween 20

0.01%) for 10 minutes, gradually dehydrated every 10 minutes in Ethanol/PBS series

(33%, 50%, 80%) and Methanol 100%. Samples were then gradually rehydrated every 10

minutes in Methanol/PBSt series, rinsed three times in PBSt, permeabilized with PBS

Triton-100 (0.2%) for 30 minutes and incubated for 2 hours at room temperature with

anti-dpERK mouse monoclonal antibody (1:200; Sigma, Saint Louis, MO) and anti-

mCherry polyclonal antibody from rabbit (1:500; Biovision, Milpitas, CA) in PBS 0.01%

Triton-100 (T-Pbs) supplemented with 2% normal goat serum. Samples were then

washed three times in T-PBS and incubated in anti-mouse and anti-rabbit antibodies

(1 :500 each), respectively coupled with Alexa 488 and Alexa 568 (Invitrogen, Carlsbad,

CA), overnight at 4°C or for 2 hours at room temperature. Finally, samples were rinsed

three times in T-PBS for 15 minutes and mounted in Prolong Gold (Molecular Probes,

Eugene, OR).

Molecular cloning

708
Coding sequences for wild-type M-Ras (KH.L172.2), Mek1/2 (KH.L147.22), Cdkn1b.a

(Cdkn1b, KH.C14.564), and Cdkn1b.b (Not016, KH.S643.6) were PCR-amplified from

cDNA libraries prepared by reverse transcription of total RNA from mixed

developmental stages. Insertion of the products into expressing vectors was performed

using regular restriction/ligation or In-fusion (Clontech, Mountain View, CA) procedure.

Oligonucleotide directed mutagenesis or two-step overlap PCRs were used to generate

the point mutated forms M-Ras^{G22V} and Mek^{S220E,S216D} from the corresponding wild-type

sequences. We also used oligonucleotide directed mutagenesis to generate mismatches in

the PAM sequences adjacent to the sgRNA targets for Hand-r (153C>T 574C>T for

Hand-r^{PAMmis}) and Tbx1/10 (325G>A and 579G>A for Tbx1/10^{PAMmis}). Due to the absence

of a correct PAM sequence (NGG, (reverse complement CCN)), overexpressed Hand-

r^{PAMmis} and Tbx1/10^{PAMmis} are resistant to the Cas9 nuclease activity. Primer sequences

are listed in Supplementary Table 1.

722
CRISPR/Cas9-mediated loss of Hand-r function

The pair of single guide RNA (sgRNA) targeting Tbx1/10 (sgTbx1/10) has been validated

previously (Tolkin and Christiaen, 2016). Rescue of the Tbx1/10 loss-of-function was

achieved by TVC-specific overexpression of Tbx1/10^{PAMmis} driven by a *FoxF* enhancer

(FoxF-1>Tbx1/10^{PAMmis}). For Hand-r loss of function, sgRNAs were first designed to

avoid genomic off-targets and tested as described (Gandhi et al., 2017). In short, sgRNA

expressing cassettes (U6>sgRNA) were assembled by single step overlap PCR. Individual

PCR products (~25 μg) were electroporated with EF1a>NLS::Cas9::NLS (30μg) ,

Myod905>Venus (50 μg), driving ubiquitous expression of Cas9 and a widely expressed

fluorescent reporter construct, respectively. Efficient electroporation was confirmed by

observation of fluorescence before genomic DNA extraction around 16 hpf (18°C) using

QIAamp DNA Micro kit (Qiagen, German Town, MD). Mutagenesis efficacy of individual

sgRNAs, as a linear function of Cas9-induced indel frequency, was estimated from

electrophoregrams following Singer sequencing of the targeted regions amplified from

extracted genomic DNA by PCR. Result of the relative quantification of the indel

frequency ("corrected peakshift" of 22% and 24%) was considered high enough for both

sgRNAs targeting Hand-r, which were finally selected. The corresponding cassettes were

cloned into plasmid for repeated electroporations to study the loss of function of Hand-r.

Rescue of Hand-r loss-of-function was achieved by overexpression of Hand-r^{PAMmis}

driven by a FoxF TVC specific enhancer (FoxF-1>Hand-r^{PAMmis}). In order to control the

specificity of the CRISPR/Cas9 system, sgRNAs targeting *Neurogenin*, a gene not

expressed in the TVC and their progeny, was electroporated in parallel. Sequences of the

DNA targets and oligonucleotides used for the sgRNAs are listed in Supplementary Table

1.

Observation and imaging

Samples were usually scored under a DM2500 epifluorescent microscope (Leica

Microsystems, Wetzlar, Germany). Imaging was performed using a TCS SP8 X inverted

confocal microscope equipped with a white light laser, AOBS and HyD detectors (Leica

Microsystems).

Acknowledgement

We thank Robert Kelly (Université Aix-Marseille, CNRS, France) for feedbacks on the

manuscript. We are grateful to Wei Wang, Nicole Kaplan, Claudia Racioppi and Alberto

Stolfi for collaborative inputs throughout the project. We thank Farhana Salek and

Kristyn Millan for technical support. This project was funded by NIH/NHLBI R01 award

HL108643, and trans-Atlantic network of excellence award 15CVD01 from the Leducq

Foundation to L.C.

Figures

Figure 1. Spatio-temporal restriction of ERK activity reflects FGF requirement for the specification of cardiopharyngeal progenitors. (**A**) Schematic of *Ciona* development showing asymmetric cell divisions and resulting cell fates of the cardiopharyngeal mesoderm (CPM). Embryonic and larval stages (St) according to (Hotta et al., 2007) with hours post fertilization (hpf) at 18°C. Anterior tail muscle (ATM, gray), trunk ventral cell (TVC, green), secondary TVC (STVC, green), first heart precursor (FHP, red), second heart precursor (SHP, orange), atrial siphon founder cell (ASMF, blue). Black bars link sister cells. Dashed lines: ventral midline. The first stage presents a quasi-lateral view while the second and third stages present quasi-ventral views. Anterior is to the left. Scale bar, 50 µm. (**B**) ERK activity visualized by anti-dpERK antibody (green). TVCs and their progeny are marked by mCherry driven by *Mesp* and revealed by anti-mCherry antibody (red). H₂B::mCherry and hCD4::mCherry accumulate in the nuclei and at the cell membrane, respectively. Arrowheads indicate STVCs and ASMFs at 14 and 16 hpf, respectively. Arrows indicate FHPs and open arrowheads mark SHPs. Anterior to the left. Scale bar, 10 µm. See also Figure S1 for broader time series of dpERK immunostaining in the B7.5 lineage. (**C, D**) TVC-specific overexpression of dnFGFR induces loss of expression of key lateral CPM markers visualized by *in situ* hybridization. (**C**) Representative expression patterns of key CPM genes (*Hand-related*, *Tbx1/10*, *Ebf*) in control embryos (control, electroporated with FoxF(TVC):bpFOG-1>Venus) and TVC-specific dnFGFR expression (electroporated with *FoxF(TVC):bpFOG-1>dnFGFR::mCherry*) individuals. TVCs and progeny are marked with *Mesp>NLS::lacZ* (red). Loss of expression in half of the TVC progeny, as presented for *Ebf*, is assumed to be due to left-right mosaicism. Arrowheads mark the ASMFs. Anterior is to the left. Scale bar, 10 µm. (**D**) Corresponding histograms with the phenotype proportions. For simplicity, loss of gene expression in half or all of the TVCs and their progeny were combined in the same category. "n" corresponds to the number of individual halves documented per condition.

787
Figure 2. Constitutively active M-Ras and MEK are sufficient to impose a pharyngeal muscle
fate in the cardiopharyngeal lineage. (**A**) Diagram of the FGF/MAPK transduction pathway with
constitutive activation by M-Ras^{G22V} and MEK^{S216D,S220E} mutants. (**B**) Expression patterns of markers of the
lateral TVC progeny, *Htr7* (a, b, c,), *Tbx1/10* (d, e, f) and *Ebf* (g, h, i), visualized by *in situ* hybridization
following TVC-specific over-expression of M-Ras^{WT} (as control), M-Ras^{G22V} and MEK^{S216D,S220E}. M-Ras^{WT}
overexpression (a, d, g) does not alter the wild-type spatial expression patterns of *Htr7*, *Tbx1/10* and *Ebf* in
lateral TVC derivatives (STVC and ASMF) and excluded from the median heart precursors. TVC-specific
over-expression of M-Ras^{G22V} (b, e, h) or MEK^{S216D,S220E} (c, f, i) induces ectopic expression of STVC and/or
ASMF markers (*Htr7*, *Tbx1/10* and *Ebf*) in the more median cells, that normally form cardiac precursors.
Solid arrowheads indicate STVCs and ASMFs at 15 and 18 hpf, respectively. Arrows indicate FHPs and open
arrowheads mark SHPs. At 18 hpf, the FHPs start dividing or have divided into 4 cells. Anterior to the left.
Scale bar, 10 μm. (**C**) Corresponding histograms: Larvae with TVC-specific over-expression of MEK^{WT} retain
the wild-type expression patterns. For simplicity, ectopic expressions in half to all of the cardiac precursors
were combined in the same phenotype category. "n" corresponds to the number of embryo halves
documented per condition. See also Figure S2.
803

804
Figure 3. Temporal requirement for MAPK activity permits the progressive deployment of
the cardiopharyngeal regulatory program. (A) Summary of the CPM cell lineage showing the

different U0126 treatments with regard to the timing of the cell divisions. Abbreviations and color codes as
in Figure 1. (**B, C**) Proportions of embryo halves with wild-type or downregulated expression of *Hand-r* at
12.5 hpf (B) and 15 hpf (C) following 3-hour incubations in U0126 (with DMSO as control treatment). (**D**)
Hand-r expression visualized by *in situ* hybridization at 15 hpf in control (DMSO treated) and U0126 treated
embryos. In control embryos, *Hand-r* remains expressed in the STVCs and downregulated in the FHPs. In
U0126 (12-15 hpf) treated embryos, downregulation of *Hand-r* expression is observed throughout the TVC
progeny (STVCs and FHPs), suggesting inhibition of transcription and inheritance of remnant transcripts
following TVC divisions. (**E, F**) Proportions of embryo halves with wild-type or downregulated expression of
Htr7 at 12.5 hpf (E) and 15 hpf (F) following 3-hour incubations in U0126 (with DMSO as control
treatment). (**G**) Proportions of larvae with wild-type expression or downregulated expression of *Tbx1/10* at
16 hpf following 4-hour incubation in U0126 (with DMSO as control). (**H**) Proportions of larvae with wild-
type or downregulated expression of *Ebf* at 18 hpf following a three hour incubation in U0126 (with DMSO
as control). (**I**) Proportions of larvae with wild-type or downregulated transcription of *Ebf* at 18 hpf following
a 3-hour incubation in U0126 (DMSO as vehicle control). (**J**) Pattern of nascent *Ebf* transcripts visualized by
in situ hybridization with intronic probes (green) at 20 hpf. The nuclear dots reveal the active transcription
sites in the four ASMPs per side in larvae, both control/DMSO- and U0126-treated from 17 to 20 hpf. (**K**)
Ebf expression (green) in 18hpf larvae expressing control M-Ras^{WT}, constitutively active M-Ras^{G22V} or
dominant negative dnFGFR under the control of the T12 element, an STVC-specific *Tbx1/10* enhancer.
Arrows: first heart precursors (FHP); open arrowhead: second heart precursors (SHPs); closed arrowheads:
ASM founder cells (ASMFs); dotted line: midline. (**L**) Proportions of larvae with wild-type or downregulated
expression of *Ebf* at 18 hpf in larvae with Venus (control), M-Ras^{WT}, M-Ras^{G22}, or dnFGFR driven by
Tbx1/10 cis-regulatory sequence and overexpressed in the STVCs. "n" : number of individual halves
documented per condition.
830

Figure 4. M-Ras/MAPK-driven feed-forward subcircuits control the successive activations of
Hand-r, *Tbx1/10* and *Ebf*. (**A**) Proportions of embryo halves with indicated *Tbx1/10* expression patterns
following TVC-specific CRISPR/Cas9-mediated mutagenesis of *Neurogenin/Neurog* as a control (*sgCtrl*),
and Hand-r (*sgHand-r*). TVC-specific overexpression of a CRISPR/Cas9-resistant form of Hand-r with
mutation in the PAM sequence (*Hand-r^PAMmis*) rescued *Tbx1/10* expression in the *sgHand-r* "background".
TVC-specific overexpression of a constitutively active M-Ras mutant (M-Ras^G22) (control: M-Ras^WT) was
sufficient to induce ectopic expression of *Tbx1/10* in the FHPs in *sgCtrl* embryos but not in *sgHand-r*
embryos indicating that Hand-r is necessary for M-Ras-dependent activation of *Tbx1/10* transcription. (**B**)
Proportions of embryo halves with indicated *Tbx1/10* expression patterns following TVC-specific
overexpression of Hand-r or a neutral reporter (Venus) and treated from 12 to 15hpf with the MEK inhibitor
U0126 (+) or with DMSO (-) as control. Hand-r overexpression is not sufficient to rescue loss of *Tbx1/10*
expression due to MAPK inhibition indicating that M-Ras/MAPK activity is required in parallel of Hand-r
expression to activate Tbx1/10 transcription in the TVC progeny. (**C**) Tbx1/10 is necessary downstream of M-
Ras/MAPK activity to activate *Ebf* transcription in the TVC progeny. Shown are proportions of *Ebf*
expression phenotypes following TVC-specific CRISPR/Cas9-mediated loss of *Tbx1/10* function (*sgTbx1/10*),
with *Neurog*-targeting sgRNA as control (*sgCtrl*). Specificity of Tbx1/10 loss of function was validated
through rescue of *Ebf* expression with TVC-specific overexpression of a CRISPR/Cas9 resistant form of
Tbx1/10 (Tbx1/10^PAMmis). Ectopic *Ebf* expression in SHPs in Tbx1/10^PAMmis larvae is explained by precocious
misexpression of Tbx1/10 in the TVC as described in Wang et al, 2013. TVC-specific overexpression of M-
Ras^G22 (M-Ras^G22), with wild type M-Ras (M-Ras^WT) as control, was sufficient to induce ectopic expression of
Ebf in the cardiac precursors in *sgCtrl* embryos but not in *sgTbx1/10* embryos indicating that Tbx1/10 is
necessary for M-Ras-dependent activation of *Ebf* transcription. (**D, E**) Proportions (D) and examples (E) of
15hpf larvae halves showing indicated *Ebf* expression phenotypes in *sgCtrl* and *sgHand-r* CRISPR/Cas9
conditions combined with TVC-specific overexpression of a neutral reporter (Venus), Hand-r^PAMmis, or
Tbx1/10, and with MEK inhibition by U0126 (+) or not (DMSO control (-)). Arrowhead: STVCs, Arrows:
FHPs, dotted line: ventral midline (**F**) Loss of Hand-r function impaired the ability of Tbx1/10 to induce
ectopic *Ebf* expression. For simplicity, ectopic expressions in half to all of the cardiac precursors were
combined in the same phenotype category. "n": number of individual halves documented per condition. (**G**)
Summary model of the temporal deployment of FGF/MAPK-driven feed-forward subcircuits leading to the
sequential activations of *Tbx1/10* and *Ebf* in the STVCs and ASMFs, respectively.

Figure 5. Temporal deployment of the cardiopharyngeal network is partially coupled with cell cycle progression. (**A**) Schematic representation of the canonical eukaryotic cell cycle, and actions of the perturbations used in this study. (**B**) Tbx1/10 and Ebf expression at indicated time points, and following inhibition of cytokinesis by cytochalasin B treatment at indicated time points. Note that 15 to 19hpf treatment is applied AFTER the first division and birth the FHPs, which do not activate *Ebf* at 19hpf (right panel, arrows). (**C**) Inhibition of G1/S or G2/M blocks TVC division, and reduces *Tbx1/10* expression. Picture shows left-right mosaic embryo, with TVCs that have not divided on the electroporated side (marked by *Mesp>H2B::mCherry*, red), one cell turned on Tbx1/10, but not the other. Left: the proportions of embryos showing strong Tbx1/10 expression is substantially reduced compared to control embryos (e.g. Figure 1, and (Wang et al., 2013)). (**D**) Inhibition of G2/M in the STVCs by misexpression of Wee1 using the *Tbx1/10 T12* enhancer inhibits STVC division, and has a mild impact on *Ebf* expression at 18hpf. Open arrows indicate STVCs that have not divided, but express high (middle) or low (right) levels of *Ebf*. Left: control larva showing high *Ebf* expression in the ASMF (closed arrowheads), but neither in the SHPs (open arrowheads) nor in the FHPs (Arrows). (**E**) Proportions of larva halves fixed at successive time points and showing undivided STVCs, or ASMFs with or without ectopic *Ebf* expression in the SHPs following STVC-specific expression of the G2/M inhibitor Wee1 (+), or a control construct (-). See Figure S4C for an example of ectopic *Ebf* expression in the SHPs (grey labels). Note the sharp increase in % of larva with ASMF between 15.5 and 16hpf, indicating that mitosis occurs primarily during this time window, but is delayed in a majority of larvae upon Wee1 misexpression. (**F**) Proportions of larva halves with cells showing indicated Ebf expression. The numbers (n) for cells expressing Wee1 focus on cells that have not divided (% shown in E), to estimate the dynamics of *Ebf* activation in G2/M-inhibited cells. Control cells consist mostly ASMFs after 15.5hpf as shown in (E). Wee1 and controls distributions differ significantly only at 16 and 16.5hpf (**, $p < 0.01$, Chi^2 test), suggesting that Wee1 merely delays the accumulation of Ebf transcripts. In all image panels, dotted line : ventral midline.

Figure 6. *Ebf* regulation transitions from MAPK-dependent to autoregulative during the early phase of ASMF cycle. (**A**) Proportions of 20hpf larva halves showing the indicated number of Ebf-expressing cells following U0126 treatments started at the indicated time points. This indicates that, by 17hpf, Ebf expression, which started at ~16hpf, has become largely insensitive to loss of MAPK activity. (B) Summary lineage diagram and time scale indicating the approximate stages for U0126 and DMSO (control) treatments for the results shown in (C, D). (C) Control (Ctrl) and Ebf-misexpressing embryos fixed at 16hpf, prior to chemical treatments, and stained for nascent transcripts with an intronic *Ebf* probe. In controls, the ASMFs (solid arrowhead), but neither the SHPs (open arrowheads) nor the FHPs (arrows), actively transcribe *Ebf* (green nuclear dots). In Larvae misexpressing the Ebf cDNA under the control of the STVC-specific Tbx1/10 enhancer, divisions are delayed and STVCs (solid arrowheads) activated transcription of endogenous *Ebf* loci (green nuclear dots). (D) After 4 hours, U0126 treated ASMFs no longer transcribe Ebf (top right image, solid arrowheads), whereas control DMSO-treated ASMFs do (top left, green nuclear dots). Upon misexpression of the Ebf cDNA in the STVCs and derivatives, ongoing *Ebf* transcription is detected at 20hpf in both DMSO and U0126-treated cells, and it persists in both ASMFs (solid arrowheads), and SHPs (open arrowheads). (**E, F**). Proportions of larvae halves showing the indicated *Ebf* transcription patterns, in indicated experimental conditions, as illustrated in C and D, respectively.

907
Figure 7. Summary model. (**A**) Schematic representation of cardiopharyngeal lineage cells at successive
time points representing the main fate transitions. hpf: hours post-fertilization; TVC: trunk ventral cells; L:
Leader T: trailer; migr.: migration; STVC: second trunk ventral cells; FHP: first heart precursors; dotted
line: midline; black bars link sister cells; ASMF: atrial siphon muscle founder cells; SHP: second heart
precursors; iASMP: inner atrial siphon muscle precursors; oASMP: outer atrial siphon muscle precursor
(these cells correspond to stem-cell-like *Mrf-*; Notch+ precursors and *Mrf*+; Notch- differentiating
myoblasts, respectively; see (Razy-Krajka et al., 2014) for details). (**B**) Lineage diagram and documented
regulatory relationships between indicated genes and pathways, as showing here and in (Razy-Krajka et al.,
2014; Wang et al., 2013). In TVCs, primed heart and ASM markers are coexpressed, and maintenance of the
STVC and ASM markers requires ongoing FGF/MAPK signaling. Following the first oriented and
asymmetric cell division, FGF/MAPK is maintained only in the STVCs, which permits the continued
expression of Hand-r and the activation of *Tbx1/10*. Cell division, presumably through G1-specific inputs,
contributes to *Tbx1/10* activation, and Tbx1/10 function antagonizes *Gata4/5/6* expression (Wang et al.,
2013). In the FHPs, termination of FGF/MAPK signaling inhibits *Hand-r* expression and prevents *Tbx1/10*
activation. Following oriented and asymmetric division of the STVCs, FGF/MAPK signaling persists only in
the ASMFs, where it permits the transient maintenance of *Hand-r* and *Tbx1/10*, both of which act in parallel
to FGF/MAPK to activate *Ebf* expression, together with contributions from presumed G1 inputs. Ebf
activities further antagonize the cardiac program (marked by *Gata4/5/6*, *Nk4/Nkx2-5* and *Hand* expression;
(Razy-Krajka et al., 2014; Stolfi et al., 2010; Wang et al., 2013)). Once Ebf expression reaches "high levels",
its regulation becomes MAPK-independent and self-activating (this study). It also feeds back negatively on
early activators such as Hand-r, and promotes the expression of the muscle determinant *Mrf* (Razy-Krajka et
al., 2014; Tolkin and Christiaen, 2016). We propose that this transition represents commitment to an ASM
fate. In the SHPs, termination of FGF/MAPK signaling prevents maintenance of Hand-r and Tbx1/10
expression, which, together with repressive inputs from Nk4/Nkx2-5, inhibits *Ebf* activation (Wang et al.,
2013), and permits heart fate specification (Wang et al., 2017).
933

Figure S1. Detailed patterns of MAPK activity during early cardiopharyngeal development.
(**A**) MAPK activation during TVC induction. Close-up views of B7.5 lineage cells marked with
Mesp>H2B::mCherry (nuclei) and Mesp>hCD4::mCherry (membranes) and immunostained for dpERK at
indicated successive time points between 7 and 10hpf. DpERK staining was not detected in the founder cells
at 7hpf, but increased sharply and specifically in the smaller trunk ventral cells (TVCs, open arrows) at
7.5hpf, but not in the larger anterior tail muscles (ATMs). DpERK staining persisted throughout TVC
migration (see also B). (B) MAPK activation patterns during cardiopharyngeal fate diversification. DpERK
staining was clearly detected in migrating TVCs (open arrows, 11 to 13hpf); in lateral large STVCs (open
arrows, 14 to 15hpf), but not in the small median first heart precursors (FHPs, arrows, 14 to 15hpf); in the
large lateral atrial siphon muscle founder cells (ASMFs, solid arrowheads, 16 to 17hpf), but neither in the
FHPs (arrows), nor in the second heart precursors (SHPs, open arrowheads). (**C**) Treatment with the MEK
inhibitor U0126 between abolished dpERK staining in the lateral STVCs, compared to a control treatment
with DMSO. Numbers of embryos showing the presented pattern out of the total numbers of embryos are
shown.

950
951 **Figure S2. Other markers expressed in the TVC need continuous FGF/MAPK inputs for**
952 **maintenance**. All panels show the proportions of 12.5hpf embryos halves showing expression of the
953 indicated genes in late TVCs, following electroporation of either a FoxF(TVC)>Venus control of a
954 FoxF(TVC)>dnFGFR construct that inhibits signaling through FGFR. Wild-type pattern were first reported
955 in (Razy-Krajka et al., 2014).
956

Figure S3. The constitutively active MEK^{S216D,S220E} mutant is sufficient to impose a TVC identity to the whole B7.5 lineage. (A) Control late tailbud embryo showing the left side B7.5 lineage expressing GFP and a MEK^{WT} control under the control of the *Mesp* enhancer. Two TVCs and two ATMs are normally induced, and TVCs migrated into the trunk. (B) Late tailbud embryo showing the left side B7.5 lineage expressing GFP and a MEK^{S216D,S220E} mutant under the control of the *Mesp* enhancer. Four cells are observed as having migrated into the trunk, indicating that they have been induced to acquire a TVC fate and migrate, replicating FGF/MAPK gain-of-function phenotypes as described in (Davidson et al., 2006). (C) Proportions of embryo halves showing the indicated phenotypes. Extra migration is interpreted as ectopic induction of the TVC fate in all B7.5 lineage cells. Scale bar ~ 20μm.

Figure S4. Rare precocious activation of *Ebf* transcription in STVCs. (**A**) 15.5hpf
Cardiopharyngeal lineage cells expressing Mesp>H2B::mCherry (red) and control Tbx1/10>unc76::GFP
construct (not visible). (**A.a**) Green nuclear dot indicates nascent Ebf transcription in an STVC (open arrow),
but not the other, and not in the first heart precursors (FHP; arrow). (A.b) left pair of nuclei shows an STVC
(open arrow) and an FHP (arrow), neither of which express *Ebf*, whereas the cousin ASMF (solid arrowhead)
shows nascent *Ebf* transcription (green dot). Dotted line : midline. (**B**) Proportions of STVCs and ASMFs
showing indicated *Ebf* expression patterns. Note that >90% of STVCs do not express *Ebf*, which turns on
almost exclusively in ASMFs. (**C**) Cardiopharyngeal lineage cells with *Ebf* expression in the ASMFs (solid
arrowheads), and ectopically in the SHP (open arrowheads), but not in the FHPs (arrows), following
misexpression of Wee1 using the STVC-specific *Tbx1/10 T12* enhancer. Dotted line: midline.

```
Primer name              primer sequence
MRAS N FWD INF bpFOG-1   cacacaaGCGGCCGCaaccATGGCGACCGTGCCGAATC
MRAS C REV INF EcoR1     gctcagctggaattcTGGCCTCTAGGTGGAGCTAC
MRAS G22V TOP            GTGGTTGGCGATGTTGGTGTCGGGAAG
MRAS G22V BOT            CTTCCCGACACCAACATCGCCAACCAC
MEK1/2 N FWD Not1        caaGCGGCCGCaaccATGCCTCCTAAACGTAAGT
MEK1/2 C REV Ecor1       ctggaattcCCCATCATATTAATCAGGTACA
MEK MUT R                CCCTACAAACTCGTTGGCCATATCGTCGATCAGTTGCCCGCTCA
MEK MUT F                CTGATCGACGATATGGCCAACGAGTTTGTAGGGACAAGATCATA
Noto16 FWD INF bpFOG-1   acacacaaGCGGCCGCaaccATGGTTCCCCCACCTTCGTAC
Noto16.2 REV INF EcoR1   GCTCAGCTGGAATTCATTTTCGTATCAATTACTTGCTTTGG
Cdkn1B FWD INF bpFOG-1   acacacaaGCGGCCGCaaccATGGCGGACAAAAAACCCCCG
Cdkn1B REV INF EcoR1     gctcagctggaattcCGTGGCACAGTATGACGTCAC
Tbx1 G325A Mut Top       CGGCTCCGTGGAGAAGAAAATGAGCG
Tbx1 G325A Mut Bot       CGCTCATTTTCTTCTCCACGGAGCCG
Tbx1 G579A Mut Top       GAAAGATTGGTGGCCGTAGAAGCAAAACTGGAAATG
Tbx1 G579A Mut Bot       CATTTCCAGTTTTGCTTCTACGGCCACCAATCTTTC
Hand-r C153T Mut Top     CTTTGCAACCGAAAATCCACACATGGTAGC
Hand-r C153T Mut Bot     GCTACCATGTGTGGATTTTCGGTTGCAAAG
Hand-r C574T Mut Top     GTCGCGTCCGAGTCATCCGTATTATCACTTC
Hand-r C574T Mut Bot     GAAGTGATAATACGGATGACTCGGACGCGAC
```

		final sgRNA
sgHandR.3.aFwd	GAGGCCCTGCTACCATGTGTGTTTAAGAGCTATGCTGGAAACAG	GAGGCCCTGCTACCATGTGT
sgHandR.3.cFwd	GAGGTTCGAAGTGATAATACGTTTAAGAGCTATGCTGGAAACAG	GAGGTTCGAAGTGATAATAC
sgHandR.3.aRev	ACACATGGTAGCAGGGCCTCatctataccatcggatgccttc	GAGGCCCTGCTACCATGTGT
sgHandR.3.cRev	GTATTATCACTTCGAACCTCatctataccatcggatgccttc	GAGGTTCGAAGTGATAATAC

Table S1. oligonucleotides sequences

REFERENCES

Abu-Issa, R., Smyth, G., Smoak, I., Yamamura, K., Meyers, E.N., 2002. Fgf8 is
required for pharyngeal arch and cardiovascular development in the mouse.
Development 129, 4613-4625.

Aggarwal, V.S., Liao, J., Bondarev, A., Schimmang, T., Lewandoski, M., Locker, J.,
Shanske, A., Campione, M., Morrow, B.E., 2006. Dissection of Tbx1 and Fgf interactions
in mouse models of 22q11DS suggests functional redundancy. Human molecular genetics
15, 3219-3228.

Alsan, B.H., Schultheiss, T.M., 2002. Regulation of avian cardiogenesis by Fgf8
signaling. Development 129, 1935-1943.

Barron, M., Gao, M., Lough, J., 2000. Requirement for BMP and FGF signaling
during cardiogenic induction in non-precardiac mesoderm is specific, transient, and
cooperative. Developmental dynamics : an official publication of the American
Association of Anatomists 218, 383-393.

Beh, J., Shi, W., Levine, M., Davidson, B., Christiaen, L., 2007. FoxF is essential for
FGF-induced migration of heart progenitor cells in the ascidian Ciona intestinalis.
Development 134, 3297-3305.

Bertrand, V., Hudson, C., Caillol, D., Popovici, C., Lemaire, P., 2003. Neural tissue in
ascidian embryos is induced by FGF9/16/20, acting via a combination of maternal GATA
and Ets transcription factors. Cell 115, 615-627.

Bondue, A., Lapouge, G., Paulissen, C., Semeraro, C., Iacovino, M., Kyba, M.,
Blanpain, C., 2008. Mesp1 acts as a master regulator of multipotent cardiovascular
progenitor specification. Cell stem cell 3, 69-84.

Bothe, I., Tenin, G., Oseni, A., Dietrich, S., 2011. Dynamic control of head mesoderm
patterning. Development 138, 2807-2821.

Brand, T., 2003. Heart development: molecular insights into cardiac specification and
early morphogenesis. Developmental biology 258, 1-19.

Brown, C.B., Wenning, J.M., Lu, M.M., Epstein, D.J., Meyers, E.N., Epstein, J.A.,
2004. Cre-mediated excision of Fgf8 in the Tbx1 expression domain reveals a critical role
for Fgf8 in cardiovascular development in the mouse. Developmental biology 267, 190-
202.

Buckingham, M., Vincent, S.D., 2009. Distinct and dynamic myogenic populations in
the vertebrate embryo. Current opinion in genetics & development 19, 444-453.

Cai, C.L., Liang, X., Shi, Y., Chu, P.H., Pfaff, S.L., Chen, J., Evans, S., 2003. Isl1
identifies a cardiac progenitor population that proliferates prior to differentiation and
contributes a majority of cells to the heart. Developmental cell 5, 877-889.

Chan, S.S., Hagen, H.R., Swanson, S.A., Stewart, R., Boll, K.A., Aho, J., Thomson,
J.A., Kyba, M., 2016. Development of Bipotent Cardiac/Skeletal Myogenic Progenitors
from MESP1+ Mesoderm. Stem cell reports 6, 26-34.

Chan, S.S., Shi, X., Toyama, A., Arpke, R.W., Dandapat, A., Iacovino, M., Kang, J., Le,
G., Hagen, H.R., Garry, D.J., Kyba, M., 2013. Mesp1 patterns mesoderm into cardiac,
hematopoietic, or skeletal myogenic progenitors in a context-dependent manner. Cell
stem cell 12, 587-601.

Chen, L., Fulcoli, F.G., Tang, S., Baldini, A., 2009. Tbx1 regulates proliferation and
differentiation of multipotent heart progenitors. Circulation research 105, 842-851.

Choi, C.Y., Lee, Y.M., Kim, Y.H., Park, T., Jeon, B.H., Schulz, R.A., Kim, Y., 1999. The
homeodomain transcription factor NK-4 acts as either a transcriptional activator or
repressor and interacts with the p300 coactivator and the Groucho corepressor. The
Journal of biological chemistry 274, 31543-31552.

Christiaen, L., Davidson, B., Kawashima, T., Powell, W., Nolla, H., Vranizan, K.,
Levine, M., 2008. The transcription/migration interface in heart precursors of Ciona
intestinalis. Science 320, 1349-1352.

Christiaen, L., Wagner, E., Shi, W., Levine, M., 2009a. Electroporation of transgenic
DNAs in the sea squirt Ciona. Cold Spring Harbor protocols 2009, pdb prot5345.

Christiaen, L., Wagner, E., Shi, W., Levine, M., 2009b. Isolation of individual cells
and tissues from electroporated sea squirt (Ciona) embryos by fluorescence-activated cell
sorting (FACS). Cold Spring Harbor protocols 2009, pdb prot5349.

Christiaen, L., Wagner, E., Shi, W., Levine, M., 2009c. Whole-mount in situ
hybridization on sea squirt (Ciona intestinalis) embryos. Cold Spring Harbor protocols
2009, pdb prot5348.

Cinnamon, E., Helman, A., Ben-Haroush Schyr, R., Orian, A., Jimenez, G., Paroush,
Z., 2008. Multiple RTK pathways downregulate Groucho-mediated repression in
Drosophila embryogenesis. Development 135, 829-837.

Cinnamon, E., Paroush, Z., 2008. Context-dependent regulation of Groucho/TLE-
mediated repression. Current opinion in genetics & development 18, 435-440.

Cooley, J., Whitaker, S., Sweeney, S., Fraser, S., Davidson, B., 2011. Cytoskeletal
polarity mediates localized induction of the heart progenitor lineage. Nature cell biology
13, 952-957.

Cota, C.D., Davidson, B., 2015. Mitotic Membrane Turnover Coordinates Differential
Induction of the Heart Progenitor Lineage. Developmental cell 34, 505-519.

Cowley, S., Paterson, H., Kemp, P., Marshall, C.J., 1994. Activation of MAP kinase
kinase is necessary and sufficient for PC12 differentiation and for transformation of NIH
3T3 cells. Cell 77, 841-852.

Dalton, S., 2015. Linking the Cell Cycle to Cell Fate Decisions. Trends in cell biology
25, 592-600.

Davidson, B., 2007. Ciona intestinalis as a model for cardiac development. Seminars
in cell & developmental biology 18, 16-26.

Davidson, B., Levine, M., 2003. Evolutionary origins of the vertebrate heart:
Specification of the cardiac lineage in Ciona intestinalis. Proceedings of the National
Academy of Sciences of the United States of America 100, 11469-11473.

Davidson, B., Shi, W., Beh, J., Christiaen, L., Levine, M., 2006. FGF signaling
delineates the cardiac progenitor field in the simple chordate, Ciona intestinalis. Genes &
development 20, 2728-2738.

Davidson, B., Shi, W., Levine, M., 2005. Uncoupling heart cell specification and
migration in the simple chordate Ciona intestinalis. Development 132, 4811-4818.

Delsuc, F., Brinkmann, H., Chourrout, D., Philippe, H., 2006. Tunicates and not
cephalochordates are the closest living relatives of vertebrates. Nature 439, 965-968.

Diogo, R., Kelly, R.G., Christiaen, L., Levine, M., Ziermann, J.M., Molnar, J.L.,
Noden, D.M., Tzahor, E., 2015. A new heart for a new head in vertebrate
cardiopharyngeal evolution. Nature 520, 466-473.

Dumollard, R., Minc, N., Salez, G., Aicha, S.B., Bekkouche, F., Hebras, C.,
Besnardeau, L., McDougall, A., 2017. The invariant cleavage pattern displayed by
ascidian embryos depends on spindle positioning along the cell's longest axis in the
apical plane and relies on asynchronous cell divisions. eLife 6.

Farley, E.K., Olson, K.M., Zhang, W., Brandt, A.J., Rokhsar, D.S., Levine, M.S., 2015.
Suboptimization of developmental enhancers. Science 350, 325-328.

Farley, E.K., Olson, K.M., Zhang, W., Rokhsar, D.S., Levine, M.S., 2016. Syntax
compensates for poor binding sites to encode tissue specificity of developmental
enhancers. Proceedings of the National Academy of Sciences of the United States of
America 113, 6508-6513.

Gainous, T.B., Wagner, E., Levine, M., 2015. Diverse ETS transcription factors
mediate FGF signaling in the Ciona anterior neural plate. Developmental biology 399,
218-225.

Gandhi, S., Haeussler, M., Razy-Krajka, F., Christiaen, L., Stolfi, A., 2017. Evaluation
and rational design of guide RNAs for efficient CRISPR/Cas9-mediated mutagenesis in
Ciona. Developmental biology 425, 8-20.

George, V., Colombo, S., Targoff, K.L., 2015. An early requirement for nkx2.5 ensures
the first and second heart field ventricular identity and cardiac function into adulthood.
Developmental biology 400, 10-22.

Gopalakrishnan, S., Comai, G., Sambasivan, R., Francou, A., Kelly, R.G., Tajbakhsh,
S., 2015. A Cranial Mesoderm Origin for Esophagus Striated Muscles. Developmental
cell 34, 694-704.

Gotoh, N., Laks, S., Nakashima, M., Lax, I., Schlessinger, J., 2004. FRS2 family
docking proteins with overlapping roles in activation of MAP kinase have distinct
spatial-temporal patterns of expression of their transcripts. FEBS letters 564, 14-18.

Gueroult-Bellone, M., Nitta, K.R., Kari, W., Jacox, E., Beule Dauzat, R., Vincentelli,
R., Diarra, C., Rothbacher, U., Dantec, C., Cambillau, C., Piette, J., Lemaire, P., 2017.
Spacer sequences separating transcription factor binding motifs set enhancer quality and
strength. bioRxiv.

Hasson, P., Egoz, N., Winkler, C., Volohonsky, G., Jia, S., Dinur, T., Volk, T., Courey,
A.J., Paroush, Z., 2005. EGFR signaling attenuates Groucho-dependent repression to
antagonize Notch transcriptional output. Nature genetics 37, 101-105.

Haupaix, N., Abitua, P.B., Sirour, C., Yasuo, H., Levine, M., Hudson, C., 2014. Ephrin-
mediated restriction of ERK1/2 activity delimits the number of pigment cells in the
Ciona CNS. Developmental biology 394, 170-180.

Hotta, K., Mitsuhara, K., Takahashi, H., Inaba, K., Oka, K., Gojobori, T., Ikeo, K.,
2007. A web-based interactive developmental table for the ascidian Ciona intestinalis,
including 3D real-image embryo reconstructions: I. From fertilized egg to hatching larva.
Developmental dynamics : an official publication of the American Association of
Anatomists 236, 1790-1805.

Hu, T., Yamagishi, H., Maeda, J., McAnally, J., Yamagishi, C., Srivastava, D., 2004.
Tbx1 regulates fibroblast growth factors in the anterior heart field through a reinforcing
autoregulatory loop involving forkhead transcription factors. Development 131, 5491-
5502.

Hudson, C., Darras, S., Caillol, D., Yasuo, H., Lemaire, P., 2003. A conserved role for
the MEK signalling pathway in neural tissue specification and posteriorisation in the
invertebrate chordate, the ascidian Ciona intestinalis. Development 130, 147-159.

Hutson, M.R., Zeng, X.L., Kim, A.J., Antoon, E., Harward, S., Kirby, M.L., 2010.
Arterial pole progenitors interpret opposing FGF/BMP signals to proliferate or
differentiate. Development 137, 3001-3011.

Ilagan, R., Abu-Issa, R., Brown, D., Yang, Y.P., Jiao, K., Schwartz, R.J., Klingensmith,
J., Meyers, E.N., 2006. Fgf8 is required for anterior heart field development.
Development 133, 2435-2445.

Imai, K.S., Levine, M., Satoh, N., Satou, Y., 2006. Regulatory blueprint for a chordate
embryo. Science 312, 1183-1187.

Imai, K.S., Satoh, N., Satou, Y., 2002. Early embryonic expression of FGF4/6/9 gene
and its role in the induction of mesenchyme and notochord in Ciona savignyi embryos.
Development 129, 1729-1738.

Jeffery, W.R., Chiba, T., Krajka, F.R., Deyts, C., Satoh, N., Joly, J.S., 2008. Trunk
lateral cells are neural crest-like cells in the ascidian Ciona intestinalis: insights into the
ancestry and evolution of the neural crest. Developmental biology 324, 152-160.

Jerome, L.A., Papaioannou, V.E., 2001. DiGeorge syndrome phenotype in mice
mutant for the T-box gene, Tbx1. Nature genetics 27, 286-291.

Kaplan, N., Razy-Krajka, F., Christiaen, L., 2015. Regulation and evolution of
cardiopharyngeal cell identity and behavior: insights from simple chordates. Current
opinion in genetics & development 32, 119-128.

Kattman, S.J., Witty, A.D., Gagliardi, M., Dubois, N.C., Niapour, M., Hotta, A., Ellis,
J., Keller, G., 2011. Stage-specific optimization of activin/nodal and BMP signaling
promotes cardiac differentiation of mouse and human pluripotent stem cell lines. Cell
stem cell 8, 228-240.

Keduka, E., Kaiho, A., Hamada, M., Watanabe-Takano, H., Takano, K., Ogasawara,
M., Satou, Y., Satoh, N., Endo, T., 2009. M-Ras evolved independently of R-Ras and its
neural function is conserved between mammalian and ascidian, which lacks classical
Ras. Gene 429, 49-58.

Kelly, R.G., Jerome-Majewska, L.A., Papaioannou, V.E., 2004. The del22q11.2
candidate gene Tbx1 regulates branchiomeric myogenesis. Human molecular genetics 13,
2829-2840.

Kelly, R.G., Papaioannou, V.E., 2007. Visualization of outflow tract development in
the absence of Tbx1 using an FgF10 enhancer trap transgene. Developmental dynamics :
an official publication of the American Association of Anatomists 236, 821-828.

Khoueiry, P., Rothbacher, U., Ohtsuka, Y., Daian, F., Frangulian, E., Roure, A.,
Dubchak, I., Lemaire, P., 2010. A cis-regulatory signature in ascidians and flies,
independent of transcription factor binding sites. Current biology : CB 20, 792-802.

Kuwajima, M., Kumano, G., Nishida, H., 2014. Regulation of the number of cell
division rounds by tissue-specific transcription factors and Cdk inhibitor during ascidian
embryogenesis. PloS one 9, e90188.

Lemmon, M.A., Schlessinger, J., 2010. Cell signaling by receptor tyrosine kinases. Cell
141, 1117-1134.

Lescroart, F., Chabab, S., Lin, X., Rulands, S., Paulissen, C., Rodolosse, A., Auer, H.,
Achouri, Y., Dubois, C., Bondue, A., Simons, B.D., Blanpain, C., 2014. Early lineage
restriction in temporally distinct populations of Mesp1 progenitors during mammalian
heart development. Nature cell biology 16, 829-840.

Lescroart, F., Hamou, W., Francou, A., Theveniau-Ruissy, M., Kelly, R.G.,
Buckingham, M., 2015. Clonal analysis reveals a common origin between nonsomite-
derived neck muscles and heart myocardium. Proceedings of the National Academy of
Sciences of the United States of America 112, 1446-1451.

Lescroart, F., Kelly, R.G., Le Garrec, J.F., Nicolas, J.F., Meilhac, S.M., Buckingham,
M., 2010. Clonal analysis reveals common lineage relationships between head muscles
and second heart field derivatives in the mouse embryo. Development 137, 3269-3279.

Lescroart, F., Mohun, T., Meilhac, S.M., Bennett, M., Buckingham, M., 2012. Lineage
tree for the venous pole of the heart: clonal analysis clarifies controversial genealogy
based on genetic tracing. Circulation research 111, 1313-1322.

Mandal, A., Holowiecki, A., Song, Y.C., Waxman, J.S., 2017. Wnt signaling balances
specification of the cardiac and pharyngeal muscle fields. Mechanisms of Development
143, 32-41.

Mansour, S.J., Resing, K.A., Candi, J.M., Hermann, A.S., Gloor, J.W., Herskind, K.R.,
Wartmann, M., Davis, R.J., Ahn, N.G., 1994. Mitogen-activated protein (MAP) kinase
phosphorylation of MAP kinase kinase: determination of phosphorylation sites by mass
spectrometry and site-directed mutagenesis. Journal of biochemistry 116, 304-314.

Marques, S.R., Lee, Y., Poss, K.D., Yelon, D., 2008. Reiterative roles for FGF signaling
in the establishment of size and proportion of the zebrafish heart. Developmental biology
321, 397-406.

Mazzoni, E.O., Mahony, S., Iacovino, M., Morrison, C.A., Mountoufaris, G., Closser,
M., Whyte, W.A., Young, R.A., Kyba, M., Gifford, D.K., Wichterle, H., 2011. Embryonic
stem cell-based mapping of developmental transcriptional programs. Nature methods 8,
1056-1058.

Merscher, S., Funke, B., Epstein, J.A., Heyer, J., Puech, A., Lu, M.M., Xavier, R.J.,
Demay, M.B., Russell, R.G., Factor, S., Tokooya, K., Jore, B.S., Lopez, M., Pandita, R.K.,
Lia, M., Carrion, D., Xu, H., Schorle, H., Kobler, J.B., Scambler, P., Wynshaw-Boris, A.,
Skoultchi, A.I., Morrow, B.E., Kucherlapati, R., 2001. TBX1 is responsible for
cardiovascular defects in velo-cardio-facial/DiGeorge syndrome. Cell 104, 619-629.

Michailovici, I., Eigler, T., Tzahor, E., 2015. Craniofacial Muscle Development.
Current topics in developmental biology 115, 3-30.

Michailovici, I., Harrington, H.A., Azogui, H.H., Yahalom-Ronen, Y., Plotnikov, A.,
Ching, S., Stumpf, M.P., Klein, O.D., Seger, R., Tzahor, E., 2014. Nuclear to cytoplasmic
shuttling of ERK promotes differentiation of muscle stem/progenitor cells. Development
141, 2611-2620.

Mosimann, C., Panakova, D., Werdich, A.A., Musso, G., Burger, A., Lawson, K.L.,
Carr, L.A., Nevis, K.R., Sabeh, M.K., Zhou, Y., Davidson, A.J., DiBiase, A., Burns, C.E.,
Burns, C.G., MacRae, C.A., Zon, L.I., 2015. Chamber identity programs drive early
functional partitioning of the heart. Nature communications 6, 8146.

Nathan, E., Monovich, A., Tirosh-Finkel, L., Harrelson, Z., Rousso, T., Rinon, A.,
Harel, I., Evans, S.M., Tzahor, E., 2008. The contribution of Islet1-expressing splanchnic
mesoderm cells to distinct branchiomeric muscles reveals significant heterogeneity in
head muscle development. Development 135, 647-657.

Nevis, K., Obregon, P., Walsh, C., Guner-Ataman, B., Burns, C.G., Burns, C.E., 2013.
Tbx1 is required for second heart field proliferation in zebrafish. Developmental
dynamics : an official publication of the American Association of Anatomists 242, 550-
559.

Norton, J., Cooley, J., Islam, A.F., Cota, C.D., Davidson, B., 2013. Matrix adhesion
polarizes heart progenitor induction in the invertebrate chordate Ciona intestinalis.
Development 140, 1301-1311.

Park, E.J., Ogden, L.A., Talbot, A., Evans, S., Cai, C.L., Black, B.L., Frank, D.U., Moon,
A.M., 2006. Required, tissue-specific roles for Fgf8 in outflow tract formation and
remodeling. Development 133, 2419-2433.

Park, E.J., Watanabe, Y., Smyth, G., Miyagawa-Tomita, S., Meyers, E., Klingensmith,
J., Camenisch, T., Buckingham, M., Moon, A.M., 2008. An FGF autocrine loop initiated
in second heart field mesoderm regulates morphogenesis at the arterial pole of the heart.
Development 135, 3599-3610.

Patterson, K.I., Brummer, T., O'Brien, P.M., Daly, R.J., 2009. Dual-specificity
phosphatases: critical regulators with diverse cellular targets. The Biochemical journal
418, 475-489.

Pauklin, S., Madrigal, P., Bertero, A., Vallier, L., 2016. Initiation of stem cell
differentiation involves cell cycle-dependent regulation of developmental genes by Cyclin
D. Genes & development 30, 421-433.

Pauklin, S., Vallier, L., 2013. The cell-cycle state of stem cells determines cell fate
propensity. Cell 155, 135-147.

Peljto, M., Wichterle, H., 2011. Programming embryonic stem cells to neuronal
subtypes. Current opinion in neurobiology 21, 43-51.

Pennati, R., Ficetola, G.F., Brunetti, R., Caicci, F., Gasparini, F., Griggio, F., Sato, A.,
Stach, T., Kaul-Strehlow, S., Gissi, C., Manni, L., 2015. Morphological Differences
between Larvae of the Ciona intestinalis Species Complex: Hints for a Valid Taxonomic
Definition of Distinct Species. PloS one 10, e0122879.

Picco, V., Hudson, C., Yasuo, H., 2007. Ephrin-Eph signalling drives the asymmetric
division of notochord/neural precursors in Ciona embryos. Development 134, 1491-1497.

Prall, O.W., Menon, M.K., Solloway, M.J., Watanabe, Y., Zaffran, S., Bajolle, F., Biben,
C., McBride, J.J., Robertson, B.R., Chaulet, H., Stennard, F.A., Wise, N., Schaft, D.,
Wolstein, O., Furtado, M.B., Shiratori, H., Chien, K.R., Hamada, H., Black, B.L., Saga, Y.,
Robertson, E.J., Buckingham, M.E., Harvey, R.P., 2007. An Nkx2-5/Bmp2/Smad1
negative feedback loop controls heart progenitor specification and proliferation. Cell 128,
947-959.

Putnam, N.H., Butts, T., Ferrier, D.E., Furlong, R.F., Hellsten, U., Kawashima, T.,
Robinson-Rechavi, M., Shoguchi, E., Terry, A., Yu, J.K., Benito-Gutierrez, E.L., Dubchak,
I., Garcia-Fernandez, J., Gibson-Brown, J.J., Grigoriev, I.V., Horton, A.C., de Jong, P.J.,
Jurka, J., Kapitonov, V.V., Kohara, Y., Kuroki, Y., Lindquist, E., Lucas, S., Osoegawa, K.,
Pennacchio, L.A., Salamov, A.A., Satou, Y., Sauka-Spengler, T., Schmutz, J., Shin, I.T.,
Toyoda, A., Bronner-Fraser, M., Fujiyama, A., Holland, L.Z., Holland, P.W., Satoh, N.,
Rokhsar, D.S., 2008. The amphioxus genome and the evolution of the chordate
karyotype. Nature 453, 1064-1071.

Racioppi, C., Kamal, A.K., Razy-Krajka, F., Gambardella, G., Zanetti, L., di Bernardo,
D., Sanges, R., Christiaen, L.A., Ristoratore, F., 2014. Fibroblast growth factor signalling
controls nervous system patterning and pigment cell formation in Ciona intestinalis.
Nature communications 5, 4830.

Razy-Krajka, F., Lam, K., Wang, W., Stolfi, A., Joly, M., Bonneau, R., Christiaen, L.,
2014. Collier/OLF/EBF-Dependent Transcriptional Dynamics Control Pharyngeal
Muscle Specification from Primed Cardiopharyngeal Progenitors. Developmental cell 29,
263-276.

Reifers, F., Walsh, E.C., Leger, S., Stainier, D.Y., Brand, M., 2000. Induction and
differentiation of the zebrafish heart requires fibroblast growth factor 8
(fgf8/acerebellar). Development 127, 225-235.

Satou, Y., Imai, K.S., Satoh, N., 2004. The ascidian Mesp gene specifies heart
precursor cells. Development 131, 2533-2541.

Shi, W., Levine, M., 2008. Ephrin signaling establishes asymmetric cell fates in an
endomesoderm lineage of the Ciona embryo. Development 135, 931-940.

Shi, W., Peyrot, S.M., Munro, E., Levine, M., 2009. FGF3 in the floor plate directs
notochord convergent extension in the Ciona tadpole. Development 136, 23-28.

Soufi, A., Dalton, S., 2016. Cycling through developmental decisions: how cell cycle
dynamics control pluripotency, differentiation and reprogramming. Development 143,
4301-4311.

Stolfi, A., Gainous, T.B., Young, J.J., Mori, A., Levine, M., Christiaen, L., 2010. Early
chordate origins of the vertebrate second heart field. Science 329, 565-568.

Stolfi, A., Gandhi, S., Salek, F., Christiaen, L., 2014a. Tissue-specific genome editing
in Ciona embryos by CRISPR/Cas9. Development 141, 4115-4120.

Stolfi, A., Lowe, E.K., Racioppi, C., Ristoratore, F., Brown, C.T., Swalla, B.J.,
Christiaen, L., 2014b. Divergent mechanisms regulate conserved cardiopharyngeal
development and gene expression in distantly related ascidians. eLife 3, e03728.

Stolfi, A., Sasakura, Y., Satou, Y., Christiaen, L., Dantec, C., Endo, T., Naville, M.,
Nishida, H., Swalla, B., Volff, J.-N., Voskoboynik, A., Dauga, D., Lemaire, P., 2014c.
Guidelines for the Nomenclature of Genetic Elements in Tunicate Genomes. Genesis
under review.

Stolfi, A., Wagner, E., Taliaferro, J.M., Chou, S., Levine, M., 2011. Neural tube
patterning by Ephrin, FGF and Notch signaling relays. Development 138, 5429-5439.

Tirosh-Finkel, L., Elhanany, H., Rinon, A., Tzahor, E., 2006. Mesoderm progenitor
cells of common origin contribute to the head musculature and the cardiac outflow tract.
Development 133, 1943-1953.

Tirosh-Finkel, L., Zeisel, A., Brodt-Ivenshitz, M., Shamai, A., Yao, Z., Seger, R.,
Domany, E., Tzahor, E., 2010. BMP-mediated inhibition of FGF signaling promotes
cardiomyocyte differentiation of anterior heart field progenitors. Development 137,
2989-3000.

Tolkin, T., Christiaen, L., 2016. Rewiring of an ancestral Tbx1/10-Ebf-Mrf network for
pharyngeal muscle specification in distinct embryonic lineages. Development in press.

Tzahor, E., 2009. Heart and craniofacial muscle development: a new developmental
theme of distinct myogenic fields. Developmental biology 327, 273-279.

Tzahor, E., Evans, S.M., 2011. Pharyngeal mesoderm development during
embryogenesis: implications for both heart and head myogenesis. Cardiovascular
research 91, 196-202.

Tzahor, E., Kempf, H., Mootoosamy, R.C., Poon, A.C., Abzhanov, A., Tabin, C.J.,
Dietrich, S., Lassar, A.B., 2003. Antagonists of Wnt and BMP signaling promote the
formation of vertebrate head muscle. Genes & development 17, 3087-3099.

Tzahor, E., Lassar, A.B., 2001. Wnt signals from the neural tube block ectopic
cardiogenesis. Genes & development 15, 255-260.

van Wijk, B., van den Berg, G., Abu-Issa, R., Barnett, P., van der Velden, S., Schmidt,
M., Ruijter, J.M., Kirby, M.L., Moorman, A.F., van den Hoff, M.J., 2009. Epicardium
and myocardium separate from a common precursor pool by crosstalk between bone
morphogenetic protein- and fibroblast growth factor-signaling pathways. Circulation
research 105, 431-441.

Vitelli, F., Morishima, M., Taddei, I., Lindsay, E.A., Baldini, A., 2002a. Tbx1 mutation
causes multiple cardiovascular defects and disrupts neural crest and cranial nerve
migratory pathways. Human molecular genetics 11, 915-922.

Vitelli, F., Taddei, I., Morishima, M., Meyers, E.N., Lindsay, E.A., Baldini, A., 2002b.
A genetic link between Tbx1 and fibroblast growth factor signaling. Development 129,
4605-4611.

von Scheven, G., Alvares, L.E., Mootoosamy, R.C., Dietrich, S., 2006. Neural tube
derived signals and Fgf8 act antagonistically to specify eye versus mandibular arch
muscles. Development 133, 2731-2745.

Wagner, E., Stolfi, A., Gi Choi, Y., Levine, M., 2014. Islet is a key determinant of
ascidian palp morphogenesis. Development 141, 3084-3092.

Wang, W., Razy-Krajka, F., Siu, E., Ketcham, A., Christiaen, L., 2013. NK4
antagonizes Tbx1/10 to promote cardiac versus pharyngeal muscle fate in the ascidian
second heart field. PLoS biology 11, e1001725.

Wang, W., Xiang, S., Jullian, E., Kelly, R.G., Satija, R., Christiaen, L., 2017. A single
cell transcriptional roadmap for cardiopharyngeal fate diversification. BioRxiv.

Watanabe, Y., Miyagawa-Tomita, S., Vincent, S.D., Kelly, R.G., Moon, A.M.,
Buckingham, M.E., 2010. Role of mesodermal FGF8 and FGF10 overlaps in the
development of the arterial pole of the heart and pharyngeal arch arteries. Circulation
research 106, 495-503.

Watanabe, Y., Zaffran, S., Kuroiwa, A., Higuchi, H., Ogura, T., Harvey, R.P., Kelly,
R.G., Buckingham, M., 2012. Fibroblast growth factor 10 gene regulation in the second
heart field by Tbx1, Nkx2-5, and Islet1 reveals a genetic switch for down-regulation in the
myocardium. Proceedings of the National Academy of Sciences of the United States of
America 109, 18273-18280.

Whittaker, J.R., 1973. Segregation during ascidian embryogenesis of egg cytoplasmic
information for tissue-specific enzyme development. Proceedings of the National
Academy of Sciences of the United States of America 70, 2096-2100.

Witzel, H.R., Cheedipudi, S., Gao, R., Stainier, D.Y., Dobreva, G.D., 2017. Isl2b
regulates anterior second heart field development in zebrafish. Scientific reports 7,
41043.

Woznica, A., Haeussler, M., Starobinska, E., Jemmett, J., Li, Y., Mount, D., Davidson,
B., 2012. Initial deployment of the cardiogenic gene regulatory network in the basal
chordate, Ciona intestinalis. Developmental biology 368, 127-139.

Yagi, H., Furutani, Y., Hamada, H., Sasaki, T., Asakawa, S., Minoshima, S., Ichida, F.,
Joo, K., Kimura, M., Imamura, S., Kamatani, N., Momma, K., Takao, A., Nakazawa, M.,
Shimizu, N., Matsuoka, R., 2003. Role of TBX1 in human del22q11.2 syndrome. Lancet
362, 1366-1373.

Yasuo, H., Hudson, C., 2007. FGF8/17/18 functions together with FGF9/16/20 during
formation of the notochord in Ciona embryos. Developmental biology 302, 92-103.

Zaffran, S., Frasch, M., 2002. Early signals in cardiac development. Circulation
research 91, 457-469.

Zhang, Z., Huynh, T., Baldini, A., 2006. Mesodermal expression of Tbx1 is necessary
and sufficient for pharyngeal arch and cardiac outflow tract development. Development
133, 3587-3595.
1370