1 **Evolutionary genetic analysis uncovers multiple species with distinct habitat**

2 **preferences and antibiotic resistance phenotypes in the *Stenotrophomonas maltophilia***

3 **complex**

6
7
8

Luz Edith Ochoa-Sánchez and Pablo Vinuesa[*]

Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Cuernavaca,
Morelos, Mexico.

*Correspondence:

Corresponding Author

vinuesa@ccg.unam.mx

Keywords: multilocus sequence analysis, species delimitation, multidrug resistance,
multispecies coalescent, population genetic structure, recombination, metallo-beta-
lactamase, multivariate statistics.

Running title: *Stenotrophomonas* evolutionary genetics and ecology

Abstract

The genus *Stenotrophomonas* (*Gammaproteobacteria*) has a broad environmental distribution. *S. maltophilia* is its best known species because it is a globally emerging, multidrug-resistant (MDR), opportunistic pathogen. Members of this species are known to display high genetic, ecological and phenotypic diversity, forming the so-called *S. maltophilia* complex (Smc). Heterogeneous resistance and virulence phenotypes have been reported for environmental Smc isolates of diverse ecological origin. We hypothesized that this heterogeneity could be in part due to the potential lumping of several cryptic species in the Smc. Here we used state-of-the-art phylogenetic and population genetics methods to test this hypothesis based on the multilocus dataset available for the genus at pubmlst.org. It was extended with sequences from complete and draft genome sequences to assemble a comprehensive set of reference sequences. This framework was used to analyze 108 environmental isolates obtained in this study from the sediment and water column of four rivers and streams in Central Mexico, affected by contrasting levels of anthropogenic pollution. The aim of the study was to identify species in this collection, defined as genetically cohesive sequence clusters, and to determine the extent of their genetic, ecological and phenotypic differentiation. The multispecies coalescent, coupled with Bayes factor analysis was used to delimit species borders, together with population genetic structure analyses, recombination and gene flow estimates between sequence clusters. These analyses consistently revealed that the Smc contains at least 5 significantly differentiated lineages: *S. maltophilia* and Smc1 to Smc4. Only *S. maltophilia* was found to be intrinsically MDR, all its members expressing metallo-β-lactamases (MBLs). The other Smc lineages were not MDR and did not express MBLs. We also obtained isolates related to *S. acidaminiphila*, *S. humi* and *S. terrae*. They were significantly more susceptible to antibiotics than *S. maltophilia*. We demonstrate that the sympatric lineages recovered display significantly differentiated habitat preferences, antibiotic resistance profiles and beta-lactamase expression phenotypes, as shown by diverse multivariate analyses and robust univariate statistical tests. We discuss our data in light of current models of bacterial speciation, which fit these data well, stressing the implications of species delimitation in ecological, evolutionary and clinical research.

1. Introduction

Bacterial species identification and delimitation are non-trivial tasks, which are critical in certain settings such as the clinic, bio-terrorism and industry. More generally, the conclusions drawn from evolutionary and ecological analyses are strongly dependent on organismal classification, as species are the relevant units of diversity (Vinuesa et al., 2005b;Koeppel et al., 2008;Shapiro et al., 2016). Proper species delimitation is a requisite to discover the species-specific phenotypic attributes underlying their ecological niche differentiation (Cadillo-Quiroz et al., 2012;Shapiro et al., 2012;Cordero and Polz, 2014).

We hypothesized that problems with species delimitations have hindered progress in systematic, taxonomic and ecological research on the ubiquitous genus *Stenotrophomonas* (Gammaproteobacteria, *Xhanthomonadales*, *Xanthomonadaceae*) (Palleroni and Bradbury, 1993), which currently comprises 12 validly described species (http://www.bacterio.net/stenotrophomonas.html). This limitation particularly affects the *S. maltophilia* species complex (Smc) (Svensson-Stadler et al., 2011), which has long been recognized to have a broad ecological distribution, being associated with humans, animals, plants and diverse anthropogenic and natural environments (Berg et al., 1999;Ryan et al., 2009;Berg and Martinez, 2015). Although different genotyping methods, particularly AFLPs (Hauben et al., 1999), rep-PCR (Adamek et al., 2011) and multilocus sequence analysis/typing (MLSA/MLST) (Kaiser et al., 2009;Vasileuskaya-Schulz et al., 2011) have clearly revealed the existence of multiple genomic groups within the Smc, proper recognition of species borders within the complex has not yet been satisfactorily achieved. This has ultimately hindered the discovery of statistically significant associations between species and traits such as habitat preferences, antibiotic resistance phenotypes and pathogenicity potential (Adamek et al., 2011;Berg and Martinez, 2015;Deredjian et al., 2016). *S. maltophilia* is an important globally emerging and multidrug-resistant (MDR) opportunistic pathogen causing difficult-to-treat infections (Chang et al., 2015). High mortality rates are reported mainly in the immunocompromised, cancer and cystic fibrosis patients, as well as those with central venous catheters or long-lasting antibiotic therapy (Looney et al., 2009;Brooke, 2012). Therefore, the identification of significant genotype-phenotype associations is critical for the safe use of particular strains from the Smc with high potential for diverse environmental biotechnologies such as bioremediation, plant growth promotion and protection (Ryan et al., 2009;Berg and Martinez, 2015).

The main objectives of this study were: *i*) to identify genetically cohesive and differentiated sequence clusters (genospecies) among a collection of environmental *Stenotrophomonas* isolates by using a combination of state-of-the-art phylogenetic and population genetic methods; *ii*) to test whether such lineages exhibit distinct phenotypic and ecological attributes, as predicted by current models of bacterial speciation. We used the multilocus dataset for the genus available at pubmlst.org (Kaiser et al., 2001;Vasileuskaya-Schulz et al., 2011), and extended it with sequences extracted from complete (Crossman et al., 2008;Lira et al., 2012;Zhu et al., 2012;Davenport et al., 2014;Vinuesa and Ochoa-Sánchez, 2015) and draft (Patil et al., 2016) genome sequences to assemble a comprehensive MLSA dataset with representative strains of 11 out of 12 validly described *Stenotrophomonas* species. We used this reference dataset to study our collection of environmental *Stenotrophomonas* isolates (*n* = 108) recovered from the sediments and water column of

several rivers with contrasting levels of contamination in the state of Morelos, Central
Mexico.

For an initial exploration of this dataset, we used thorough maximum-likelihood tree
searching. The evidence from this phylogenetic analysis was used to define diverse species
border hypotheses, which were formally evaluated in a Bayesian framework under the
multispecies coalescent (MSC) model (Rannala and Yang, 2003;Edwards et al.,
2007;Degnan and Rosenberg, 2009) by subjecting them to Bayes factor (BF) analysis (Kass
and Raftery, 1995). To the best of our knowledge, this is the first study that evaluates the
utility of this Bayesian statistical framework for bacterial species delimitation, which is
emerging as a successful and promising strategy for species delimitation in plants and
animals (Fujita et al., 2012;Aydin et al., 2014;Grummer et al., 2014). The MSC model is
independent of gene concatenation and acknowledges the very well known fact that gene
trees have independent evolutionary histories embedded within a shared species tree
(Degnan and Rosenberg, 2006;Rosenberg, 2013). The basic MSC model assumes that gene
tree discordance is solely the result of stochastic coalescence of gene lineages within a
species phylogeny. Populations, rather than alleles sampled from single individuals, are the
units to infer phylogeny in the MSC framework, effectively connecting traditional
phylogenetic inference with population genetics, providing estimates of topology,
divergence times and population sizes (Rannala and Yang, 2003;Edwards et al., 2007;Heled
and Drummond, 2010).

Current microbial speciation models predict that bacterial species-like lineages should be
identifiable by significantly reduced gene flow between them, even when recombination
levels are high within species. Such lineages should also display differentiated ecological
niches (Koeppel et al., 2008;Cadillo-Quiroz et al., 2012;Shapiro and Polz, 2014). This
study shows the power of modern phylogenetic and population genetic methods to delimit
species borders in bacteria and demonstrates that the Smc, as currently defined in
pubmlst.org, genome databases and literature, contains multiple genospecies that are
ecologically and phenotypically differentiated. We discuss our findings and approaches in
the light of current models of bacterial speciation, highlighting the practical implications
and ecological relevance of proper species delimitation.

2. Materials and Methods

Sampling sites and isolation of environmental *Stenotrophomonas* strains

Stenotrophomonas strains were recovered from the sediments and water columns at 6 sites
of 4 rivers and streams in the State of Morelos, Central Mexico (Table 1, supplementary
Fig. S1). These sites experience different levels of anthropogenic pollution, broadly
classified as low (L), intermediate (I) and high (H), based on triplicate counts of thermo-
tolerant coliforms (TTCs) on mFC agar (Oxoid). Thermotolerant *E. coli* (TTEc) counts
were obtained on modified m-TEC agar (USEPA, 2002), using the one-step membrane
filtration (0.45 µm) method (APHA, 2005). Water samples were taken in sterilized 1L
recipients at 5-20 cm depth (2 per site). Six physico-chemical parameters of the water

columns were measured using a HANNA multi-parametric HI9828 instrument operated in
continuous measurement mode, for 1 minute, along a 10 m transect (Fig. S2). Sediment
samples (3 per site, along a 3 m linear transect) were taken from the same sites in sterile
plastic cores dug 2-3 cm deep into the sediment. Samples were kept on ice until processing
within 4-8 hrs (APHA, 2005). Sampling took place at the end of the dry season (April-
May), between 2012 and 2014. Oligotrophic [NAA (Aagot et al., 2001) and R2A (Ultee et
al., 2004) agar] and rich media [LAM (Jawad et al., 1994) and MacConkey], supplemented
or not with antibiotics [trimethoprim 30 + carbenicillin 100, ciprofloxacin 4, ceftazidime 8,
cefotaxime 4, imipenem 4 (µg/ml)] were used to isolate bacteria from these samples by
plating 100 µl of serial dilutions (1 to $10e^{-4}$) in triplicate for each sample and incubation at
30°C for up to 24 hrs. Single colonies were repeatedly streaked on the same media for
strain purification. Bacteria were routinely grown on LB and stored frozen in this medium
supplemented with 20% (V/V) glycerol at -80°C.

Determination of antibiotic resistance and β-lactamase expression profiles

A total of 15 antimicrobials from 6 families and two inhibitor/β-lactamase combinations
were used to determine the resistance profiles of each strain by streaking them in parallel
on agar plates supplemented with the antibiotics and concentrations indicated in
supplementary Table S1. Double disk synergism (DDS) assays were performed to
determine the expression phenotypes of specific β-lactamase types [Ambler class A
extended spectrum beta-lactamases (ESBLs), class B metallo-beta-lactamases (MBLs) and
class C cephalosporinases (AmpC)], as detailed in the legend to Fig S10. The antibiotic
breakpoint concentrations and growth inhibition zones were interpreted according to the
26[th] edition of the Clinical and Laboratory Standards Institute (CLSI, 2016) values for
Stenotrophomonas, *Pseudomonas aeruginosa* or *Enterobacteriaceae*, when not available
for the first or second genus, respectively (cutoff values are shown in Tables S1 and S2).

PCR amplification of 16S rDNA sequences and their phylogenetic analysis

172   All strains recovered were classified at the genus level by phylogenetic analysis of the 16S
173   rRNA gene (*rrs*) sequences amplified with the universal fD1/rD1 primers (Weisburg et al.,
174   1991), as previously described (Vinuesa et al., 2005a), and detailed in the supplementary
175   material (supplementary protocol 1).

PCR amplification and maximum likelihood phylogenetic analysis of multilocus
sequence data

For multilocus sequence analysis (MLSA) of environmental *Stenotrophomonas* isolates we
used the primers and conditions reported at http://pubmlst.org/smaltophilia/, except for the
mutM_steno_6F (5'-ytdcccgaagtmgaaacyac-3') and mutM_steno_684R (5'-
gcagytcctgytcgaartarcc-3') primers, which were designed *de novo* using the Primers4Clades
server (Contreras-Moreira et al., 2009) fed with *mutM* orthologues identified using the

GET_HOMOLOGUES package (Contreras-Moreira and Vinuesa, 2013) from
Stenotrophomonas genome sequences (data not shown). PCR amplicons were purified and
commercially sequenced at both strands by Macrogen, (South Korea). Raw reads were
assembled with the phredphrap script (de la Bastide and McCombie, 2007), codon-based
multiple sequence alignments generated with an in-house Perl script, and MSA borders
stripped to match the reference pubmlst.org profiles. Individual gene alignments were
concatenated and the resulting matrix subjected to model selection with jModelTest2
(Darriba et al., 2012) for phylogenetic analysis under the maximum likelihood criterion in
PhyML3 (Guindon et al., 2010). Tree searches were initiated from 1000 random seed trees
and a BioNj phylogeny, under the BEST moves option, as previously described (Vinuesa et
al., 2008).

196 Identification of the 7 MLSA loci in genome sequences retrieved from GenBank

We selected 24 complete (Crossman et al., 2008;Lira et al., 2012;Zhu et al.,
2012;Davenport et al., 2014;Vinuesa and Ochoa-Sánchez, 2015) and draft (Patil et al.,
2016) genome sequences available in GenBank to expand our dataset with additional key
reference strains. The orthologs of the seven MLSA loci were identified using from single-
copy homologous gene clusters computed with the GET_HOMOLOGUES package
(Contreras-Moreira and Vinuesa, 2013). We found that the *gap* gene in the draft genome
sequence of strain *S. ginsengisoli* DSMC24757[T] (Acc. No. LDJM00000000) (Patil et al.,
2016) contains a thymidine insertion at position 602 that causes a frame-shift mutation and
a premature end of the gene. Consequently, the last 72 sites of this sequence were re-coded
as '?' (missing characters).

208 Sequence data availability

209 The sequences generated in this study for multilocus sequence analysis were deposited in
210 GenBank under accession numbers KX895367-KX896038.

212 Bayesian species delimitation using the multispecies coalescent (MSC) and Bayes
213 factors (BFs)

Bayesian species delimitation from multilocus data under the MSC model was performed
using the recent *BEAST2 module (version 0.7.2) for BEAST 2 (Heled and Drummond,
2010;Bouckaert et al., 2014), to evaluate a set of explicit hypotheses of species-boundaries.
*BEAST2 was run using the best fitting partitioning scheme (see supplementary protocol
2) and the TrN+G model with empirical frequencies, without rate estimation. Trees were
unlinked across partitions, setting the ploidy level to 1 for each gene tree and assuming a
constant population IO population model. A non-correlated relaxed log-normal clock
(Drummond et al., 2006) was assumed for each partition, fixing the clock rate of the first
partition and estimating the remaining ones. A non-calibrated Yule prior was set on the
species tree. The default 1/x population mean size prior was changed for a proper inverse

gamma prior (Baele et al., 2013), with shape parameter alpha = 2 and scale parameter beta
= 2 and an initial value of 0.05. The upper and lower bounds were set to 0.001 and 1000.0,
respectively. Path sampling was used to estimate the marginal likelihoods of each species
delimitation model in *BEAST2 runs for BF calculations (Lartillot and Philippe,
2006;Baele et al., 2012;Grummer et al., 2014), with the MODEL_SELECTION 1.3.1
package. Each *BEAST2 chain was run for 10^8 generations, sampling the posterior every
20000^{th}, with 10 replicate runs and the alpha value set to 0.3, applying 50% burnin. A final
triplicate *BEAST2 analysis was set up to get the final estimate of the multispecies
phylogeny under the best delimitation model with the same parameters, priors, chain length
and sampling frequency described above. Convergence and mixing of replicate runs was
checked in tracer (http://tree.bio.ed.ac.uk/software/tracer/), as well as the effective sample
size values for each parameter. The species tree corresponding to the best species-
delimitation hypothesis was visualized with densitree (Bouckaert, 2010), on combined
post-burnin (50%) species tree files generated with logcombiner. A summary tree was
generated from the latter with treeannotator and visualized with FigTree v1.4.2
http://tree.bio.ed.ac.uk/software/figtree/.

DNA polymorphism, population structure and recombination analyses

Descriptive statistics for DNA polymorphisms, population differentiation, gene flow,
diverse neutrality and population growth tests, as well as coalescent simulations, were
computed with DNAsp v.5.10.01 (Rozas et al., 2003), as previously described (Vinuesa et
al., 2005b). Bayesian analysis of population structure based on multilocus sequence data
was performed in STRUCTURE v2.3.4 under the admixture and correlated gene
frequencies models (Pritchard et al., 2000;Falush et al., 2003;2007). Twenty runs were
launched for each K value between 2 and 10, with 10^5 steps sampled after a burnin of $2x10^5$
chain iterations. The best K value was defined by the Evanno (Evanno et al., 2005) and
Pritchard (Pritchard et al., 2000) methods, as implemented in CLUMPAK (Kopelman et al.,
2015). Estimation of recombination rates of selected lineages was performed with
ClonalFrameML v1.0.9 (Didelot and Wilson, 2015), using ML trees and Ti/Tv ratios
estimated under the HKY85+G model with PhyML3 (Guindon et al., 2010).

Statistical analyses

All statistical and graphical analyses were performed with base R (R Development Core
Team, 2016) and add-on packages. Basic data manipulation, transformation and graphical
displays were achieved with functions of the tidyverse metapackage (https://CRAN.R-
project.org/package=tidyverse). Tests for normality, homoscedasticity, outliers and skew
were performed with the car (https://cran.r-project.org/package=car) and moments
(https://cran.r-project.org/package=moments) packages. Robust ANOVA and associated
post-hoc analyses (Wilcox, 2016) were performed with the WRS2 package (https://cran.r-
project.org/package=WRS2). Empirical distributions of test statistics were generated by
bootstrapping with the boot package (https://cran.r-project.org/package=boot). Multivariate
association plots for categorical data were performed with the vcd package (https://cran.r-

266   [project.org/package=vcd](https://cran.r-project.org/package=vcd)). Multiple correspondence analysis (MCA) was performed with
267   the FactoMineR (https://cran.r-project.org/package=FactoMineR) and factoextra packages
268   (https://cran.r-project.org/package=factoextra).

3. Results

3.1 Evaluation of different isolation media for the recovery of *Stenotrophomonas* from aquatic ecosystems with contrasting degrees of fecal contamination

We sampled 6 sites located in four rivers/streams of Morelos (supplementary figure S1) that were ranked into three categories based on their pollution level (low, intermediate, high), based on counts of thermotolerant fecal coliforms and *E. coli* (Table 1). Additional physicochemical parameters of each sampling site are presented in Fig. S2. Classification of the isolates at the genus level was based on the phylogenetic analysis of 16S rRNA gene sequences ($n = 697$), as shown in Fig. S3. *Stenotrophomonas* was the second most abundant genus ($n = 154$, 22.1 %) recovered in our collection, after *Pseudomonas* ($n = 239$, 34.3 %), as shown in Fig S4. The inset in Fig. S4 shows a Trellis barplot summarizing the relative efficiency of the different microbiological media tested for the recovery of *Stenotrophomonas*. The analysis reveals that environmental *Stenotrophomonas* strains can be efficiently recovered on the oligotrophic NAA medium supplemented with imipenem (8 µg/ml; > 90% recovery efficiency). The rich McConkey medium amended with imipenem is also useful (4 µg/ml; ~60%), selecting non-fermenting (whitish) colonies.

3.2 Phylogenetic structure of the genus *Stenotrophomonas* and the definition of species border hypotheses

We used intense maximum likelihood (ML) tree searching (see methods) to obtain a global hypothesis of the phylogenetic structure of the genus based on 194 non-redundant multilocus STs (Fig. 1). This dataset contains sequences retrieved from pubmlst.org (seven loci for 103 STs representative of all the "classic" genogroups and clusters defined in previous works (Kaiser et al., 2009;Vasileuskaya-Schulz et al., 2011), plus 24 selected reference strains from different species for which the genome sequences were retrieved from GenBank and the 7 loci extracted, as explained in methods. This comprehensive set of reference strains comprises 11 out of 12 validly describes species of the genus as of April 2017. Recently *S. tumulicola* (Handa et al., 2016) was added to the list (www.bacterio.net/stenotrophomonas.html; last access April 10th, 2017), which is the only species missing from our analysis, as it lacks a genome sequence or MLSA data in pubmlst.org. To this set we added the sequences generated in this study for 108 environmental isolates from Mexican rivers, comprising 63 haplotypes (distinct multilocus sequence types). Figure 1A presents the best hypothesis found among 1001 independent tree searches (the ln*L* profile of the tree search is shown in figure S5A), displaying the Smc clade in collapsed form. The ML tree was rooted using four *Xanthomonas* species as the outgroup, chosen based on the evidence of a comprehensive ML phylogeny of nearly full-length 16S rRNA gene sequences for all type strains currently described in the order *Xanthomonadales* (Fig. S6). *S. panacihumi* (Yi et al., 2010) represents the most basal lineage of the genus, which is consistent with its position on the 16S rRNA gene phylogeny (Fig. S6). However, this species was described based on the *rrs* sequence analysis of a single isolate and currently has a non-validated taxonomic status. Two large clades follow, labeled I and II on figure 1A. All species grouped in clade I are of diverse environmental origin, lacking strains reported as opportunistic pathogens. Three of our environmental

isolates tightly cluster with the type strain of *S. acidaminiphila* (Assih et al., 2002),
including strain ZAC14D2_NAIMI4_2, for which we have recently reported its complete
genome sequence (Vinuesa and Ochoa-Sánchez, 2015). The type strain of this species
(AMX19[T], LMG22073[T]) was isolated from the sludge of a lab-scale anaerobic chemical
waste water reactor in Iztapalapa, Mexico City, 1999 (Assih et al., 2002). A single isolate
of our collection is phylogenetically related to *S. humi*, while 7 others form a perfectly
supported clade with the type strain of *S. terrae* (Heylen et al., 2007). In conclusion, all
environmental isolates grouped in clade I were conservatively classified as indicated by the
labeled boxes on Fig. 1A, based on the very strong support of the monophyletic clusters
formed with the corresponding type strains.

Clade IIa groups strains of *S. chelatiphaga* and *S. rhizophila* and clade IIb groups strains of
the *S. maltophilia* species complex (Smc) (Fig. 1A). Both of them hold human
opportunistic pathogens and environmental isolates, and suffer from taxonomic problems.
The taxonomic inconsistency of classifying strains of genogroups #8 and # 10 as *S.*
maltophilia was previously recognized (Vasileuskaya-Schulz et al., 2011). They cluster
within the *S. chelatiphaga-S. rhizophila* clade, making *S. maltophilia* polyphyletic. *S.*
maltophilia strains of genogroup #8 were already recognized to belong to *S. rhizophila*, but
the taxonomic status of its sister clade, genogroup #10, holding strains labeled as *S.*
maltophilia, was not clarified (Vasileuskaya-Schulz et al., 2011).

Figure 1B shows the same phylogeny presented in Fig. 1A, but collapsing clades I and IIa,
and displaying the Smc strains grouped in cluster IIb. All terminal clades containing only
reference sequences were also collapsed to avoid excessive cluttering. Figure S5B displays
the same tree but without collapsing those clusters. The Smc clade was conservatively split
into 5 potential species (labeled boxes in Fig. 1B) based on the deep and strongly supported
branches subtending MLSA phylogroups Smc1-Smc4, and taking into account the location
of the well characterized *S. maltophilia* model strains K279a (Crossman et al., 2008), D457
(Lira et al., 2012) and the type strain of the species ATCC 13637[T] (Davenport et al., 2014).
The latter three are spread across the clade labeled as *S. maltophilia* (Fig. 1B). Each of the
phylogroups Smc1 to Smc4 currently holds ecologically coherent groups of strains: Smc1
and Smc2 contain exclusively Mexican river isolates recovered in this study, Smc3 groups
cystic fibrosis isolates and Smc4 predominately rhizosphere isolates from diverse plants
and parts of the world. In contrast, the large *S. maltophilia* clade holds heterogeneous
groups of isolates of clinical and environmental origin, including the previously defined
MLSA phylogroups A to E (Kaiser et al., 2001;Vasileuskaya-Schulz et al., 2011) and
AFLP genogroups #1 to #10 (Hauben et al., 1999). We conservatively define a new MLSA
phylogroup F, comprising isolates of clinical and environmental origin from different
continents, but dominated by river isolates reported herein (Fig. 1B). We note that *S.*
pavanii DSM 25135[T], an endophytic N_2-fixing bacterium isolated from sugarcane in Brazil
(Ramos et al., 2010), is clearly nested within genogroup #1, closely related to the reference
strain *S. maltophilia* D457 (Fig. 1B). This represents an additional taxonomic inconsistency
for the *S. maltophilia* clade not previously reported. Here we suggest that *S. pavanii* is a
late heteronym of *S. maltophilia*. *S. africana*, nested within genogroup #4, had already been
described as a later heterotypic synonym of *S. maltophilia* (Coenye et al., 2004;Kaiser et
al., 2009), the same as *Pseudomonas beteli* 962[T], related to MLSA genogroup E, and *P.*
hibiscicola ATCC19867, related to genogroup #3 (Hauben et al., 1999;Vasileuskaya-

Schulz et al., 2011). The latter two species were already recognized in 1990 to be
misclassified based on the analysis of an extensive set of phenotypic features, being
synonyms of *S. maltophilia* (Van Den Mooter and Swings, 1990). The taxonomic status of
S. tumulicola (Handa et al., 2016) could not be revised in the present work because it lacks
multilocus sequence data.

363 3.3 Bayesian species delimitation based on the multispecies coalescent (MSC) model
364 and Bayes factor (BF) analysis of marginal likelihoods

We used a recent software implementation of the MSC model (Heled and Drummond,
2010;Bouckaert et al., 2014) to test the explicit species delimitation hypotheses highlighted
in Figs. 1A and 1B by means of BF analysis of marginal likelihoods (Grummer et al., 2014)
in a formal Bayesian statistical framework. The best partitioning scheme (see
supplementary protocol 2 and Fig. S7) was used for all *BEAST2 runs. Of particular
interest to this work was the evaluation of the following five species delimitation
hypotheses within the Smc: split1 (species assignations as defined by the shaded areas on
Figs. 1A and 1B), lump_*S maltophilia*+Smc1, lump_*S. maltophilia*+Smc12, lump_*S.
maltophilia*+Smc123, lump_*S. maltophilia*+Smc1234, which successively lump the *S.
maltophilia* sequence cluster with the proposed Smc1-Smc4 genospecies (Table 2 and Fig.
1B). Analysis of the logfiles for the path sampling runs for each model and replicate had
effective sample size (ESS) values > 150 for all parameters, most of them with ESSs >>
500. As shown in Table 2, the split1 hypothesis was the favored one as it attained the
highest marginal likelihood. The BF analysis provides overwhelming evidence (Table 2)
that the Smc, as actually defined in pubmlst.org, lumps multiple cryptic species, strongly
supporting the species delimitation hypothesis presented in Fig. 2B, which conservatively
splits the complex into the 5 species: *S. maltophilia* and the four new lineages Smc1, Smc2,
Smc3 and Smc4. Smc1 and Smc2 contain only Mexican representatives of environmental
Stenotrophomonas sampled in this study. The split1 vs. lump_cluster#8+cluster#10 (Fig
1A) model was also evaluated, providing overwhelming evidence in favor of separating the
two genogroups #8 and #10 as distinct species (ln-BF > 5, Table 2). Supplementary figures
S8 and S9 show the consensus and DensiTree (see methods) species tree representations,
respectively, of the merged (3 replicate runs), post-burnin (50%) samples, for the best
hypothesis.

390 3.4 Population genetic structure analysis of Mexican environmental *S. maltophilia*
391 complex isolates

In order to challenge the results of the multispecies coalescent (MSC)-based species
delimitations presented in the previous section, we performed a Bayesian population
clustering analysis on all Mexican isolates grouped within the Smc clade (Fig. 1B) using
STRUCTURE (see methods). We included also the Smc4 lineage, identified among
pubmlst.org sequences, as an outgroup control. Figure 2 shows the optimally aligned
STRUCTURE barplots for the 20 replicate runs made for the indicated *K* values, depicting
the ancestry proportions of each individual. Evanno's delta-*K* method estimated an optimal

number of 3 clusters, while Pritchard's method suggested an optimal $K = 6$. This analysis
uncovered a strong population subdivision of the Mexican Smc isolates, which is consistent
with the MSC-based species delimitation (Fig. 1B). Given the independent evidence from
the MSC analysis, we favor a conservative $K = 4$, as it consistently resolves the same 4
clusters classified as distinct species by the phylogenetic approach (*S. maltophilia*, Smc1,
Smc2 and Smc4). Detailed inspection of the barplots reveals that already at $K = 4$ an
important substructure exists within the *S. maltophilia* and Smc1 lineages. At $K = 6$ the
Mexican *S. maltophilia* population gets subdivided into three clusters, while Smc1 is split
into two subgroups. Clear evidences of admixture exist in both clades, suggesting that gene
flow and recombination might be at play in these clusters.

3.5 McDonald-Kreitman (MK) neutrality test, genetic differentiation and gene flow
estimates between pairs of environmental lineages of the *S. maltophilia* complex

To determine the statistical support of the major clusters revealed by STRUCTURE we
computed the K^{*}_{ST} index of population genetic differentiation (Hudson et al., 1992)
between them. The results presented in Table 3 indicate that the Smc1, Smc2 and *S.*
maltophilia lineages are strongly differentiated ($p < 0.001$) based on the K^{*}_{ST} index, with
multiple fixed differences between populations (range 9 - 45) and mean population
divergences (*D*xy) > 4.5%. The high F_{ST} fixation indices (range $0.43 - 0.59$) further denote
very strong population differentiation. This is consistent with the low numbers of effective
migrants per generation estimated (*Nm* range $0.34 - 0.66$), indicating limited genetic flux
between these lineages. We applied the MK test (McDonald and Kreitman, 1991) and
computed the "neutrality index" (*NI*) (Rand and Kann, 1996) to test if the observed
polymorphisms between pairs of lineages have evolved by the accumulation of neutral
mutations by random drift, the fixation of adaptive mutations by selection, or a mixture of
the two. As shown in Table 3, the *G* and *NI* indices for pairwise species comparisons
indicate that fixed differences between species are due to nonsynonymous differences more
often than expected, suggesting that positive selection may be driving divergence of Smc1
and *S. maltophilia* from Smc2. This signal is not significant ($p = 0.077$) for the Smc1 and *S.*
maltophilia comparison.

3.6 Comparative analysis of DNA polymorphisms and recombination rates across the
S. maltophilia complex (Smc) and S. terrae lineages

Table 4 presents basic descriptive statistics of DNA polymorphisms, neutrality and
population growth tests computed for the Mexican populations/genospecies with >= 10
isolates (Smc1, Smc2, *S. maltophilia* and *S. terrae*. Based on their average nucleotide
diversity per site (π) values, the lineages sort in the following decreasing order of diversity:
S. maltophilia > Smc1 > *S. terrae* > Smc2. The high π values, high numbers of haplotypes
(*h*) and the high haplotype diversity values (*Hd*) consistently reveal that species in the
genus comprise notoriously diverse gene pools. Tajima's *D* values are all negative, but non-
significant, suggesting that the loci are either under purifying selection or that populations
are undergoing expansions. We used the R_2 population growth test statistic (Ramos-Onsins

and Rozas, 2002) to test the null hypothesis of constant population size. Since all *p*-values
were > 0.1, there is no evidence of population expansion.

Table 5 shows the estimates for *R*/*theta*, the ratio between the mean number of
recombination to mutation events. The ratios are > 1 (except for *S. terrae*) and for Smc2
and *S. maltophilia* the recombination events are estimated to be almost twice and nearly
three times the number of mutation events, respectively. This indicates that homologous
recombination events introduce significantly more polymorphisms into the
Stenotrophomonas genomes than point mutations. The inverse mean DNA import length
estimates (1/*delta*) suggest that on average rather long sequence stretches are affected by
recombination (range 375-1472 nt), with a considerable mean sequence divergence (range
0.009-0.065).

3.7 High diversity of novel STs among the Mexican Smc isolates

The allele numbers and STs were determined for each of the 77 isolates in the Smc (Smc1
= 11, Smc2 = 15 and *S. maltophilia* = 51) recovered from Mexican rivers, by comparing
them with the corresponding 177 STs and associated alleles downloaded from pubmlst.org
(as of Nov. 18[th], 2016) using an in house Perl script. A high diversity of new alleles was
discovered, as summarized in Table S4. Only the ST139, displayed by the Mexican *S.*
maltophilia isolate ESTM1D_MKCAZ16_6, was previously reported in pubmlst.org,
highlighting the novelty of the genotypes recovered in this study. A single entry is currently
found in pubmlst.org for ST139, which corresponds to the Spanish isolate S3149-CT11 (id
220), recovered in 2007 from a surgical wound of a patient treated in an hospital from
Barcelona. The Mexican Smc strains add 56 new STs (numbers 178 – 233) to those
reported in the pubmlst.org database, with ST219 being the most prevalent one (see
supplementary Tables S3, S4 and Figure S10), shared by 9 isolates recovered from the
sediments of the highly contaminated TEX site (Table 1), both on MK and NAA media.

3.8 Multivariate association mapping of species, antimicrobial resistance phenotypes
and habitat preferences by multiple correspondence analysis (MCA)

We used MCA to visualize the associations between the antibiotic resistance profiles, β-
lactamase production phenotypes (Fig. S11), habitat preferences and species assignations
(Figs. 1A and 1B) made for the Mexican *S. maltophilia*, Smc1, Smc2 and *S. terrae* isolates
(all with > 10 isolates /species). Figure 3 depicts the MCA factor-individual biplot resulting
from the analysis of 17 active variables and 4 supplementary categorical variables, listed in
the figure caption. The clouds of individuals for each species were hidden (visible in Fig.
S12) to avoid over-plotting, but the 95% confidence intervals (CIs) for species are shown as
color-coded ellipses. The first two dimensions explain 45.3% of the variance, the first
dimension accounting for > 3.8 times the variability explained by the second and following
ones, as shown in the screeplot presented in Fig. S13A. The variable plot depicted in Fig.
S13B reveals that the active variable species is most correlated one with the two first two
dimensions, indicating that their resistance profiles and β-lactamase expression phenotypes

are distinct. The variables GM, KA, MER, FEP, TET, ATM (abbreviations defined in the
legend to Fig. 3), β-lactamase expression and the MDR condition are strongly correlated
with the first component, while IMP and SM are the variables most strongly associated
with the second dimension (Fig. S13B). Figure 3 shows that the *S. maltophilia* (malt)
strains form a distinct and independent cloud that is characterized by a very strong
association with a resistance status to the following antibiotics: CAZ, CAZ.CLA, GM, FEP,
GM, KA and SM. It is also strongly associated with the MDR condition and metallo-β-
lactamase production. The latter are the most-strongly contributing variables for the
delimitation of this group, as depicted in the variable-categories MCA map presented in
Fig. S14. *S. maltophilia* shows a preference for the sediments of contaminated sites. The
resistance phenotypes and habitat preferences of the Smc1 and Smc2 lineages largely
overlap, those of *S. terrae* being more differentiated, but partially overlapping with the 95%
CI ellipse for Smc2. The Smc1 and Smc2 lineages are strongly associated with non MDR,
aminoglycoside, P.T. and Tm.Cb sensitivity, showing a preference for the water column of
clean or moderately contaminated sites (Fig 3 and Fig S13). The *S. terrae* isolates are
distinctly and strongly associated with carbapenem and ATM sensitivity (Fig 3 and Fig
S14). The statistical significance of these antibiotic resistance and habitat preference
patterns will be formally tested in the next two sections, respectively.

3.9 Only *S. maltophilia* is truly multidrug-resistant (MDR) and isolates from polluted
sites express resistance to more antibiotic families

We performed one-way ANOVA analyses to evaluate differences *i*) in the mean number of
resistances to individual antibiotics (NumR), *ii*) in the mean number of distinct drug
families (NumFam) across species, and *iii*) to determine whether *S. maltophilia* isolates
recovered from high and low pollution sites have the same NumR and NumFam. Figures
4A, 4B, 4E and 4F display violin and boxplots for the raw count data, which revealed
skewed, non-normal distributions, with a few outliers. Key assumptions (homoscedasticity
and normality) made by standard one-way ANOVA were formally tested (Tables S5, and
S6), which confirmed multiple cases of highly significant departures from normality.
Consequently, we performed robust one-way ANOVA (Wilcox, 2016) using trimmed
means ($tr = .2$) and bootstrap ($nboot = 2000$) to simulate the distribution of the trimmed
sample means and compute the appropriate critical values for the confidence intervals (95%
CIs). Figs. 4C and 4D depict mean plots with 95% CIs for the NumR and NumFam,
respectively, across the four species with > 10 isolates. These clearly show that *S.*
maltophilia strains express significantly higher mean$_{(tr=.2)}$ NumR (12.63) and NumFam
(4.66) than the other three species, only *S. maltophilia* being truly MDR. Highly significant
results for both the NumR [$F_t = 176.3447$, $p = 0$. Variance explained (σ^2) $= 0.821$; effect
size (*e.s.*) $= .906$] and NumFam ($F_t = 32.2755$, $p = 0$; $\sigma^2 = .511$; *e.s.* $= .715$) were obtained,
thus rejecting the null hypothesis of equal trimmed means for both variables across species,
and revealing huge effect sizes ($>.8$). A Wilcox robust *post-hoc* test of the trimmed mean
NumR and NumFam comparisons across pairs of species confirms that all those involving
S. maltophilia are very highly significant ($p = 0$; Figs. S15A and S15B, Tables S7 and S8),
as indicated by asterisks on the mean$_{(tr=.2)}$ plots shown in Figs. 4C and 4D.

Sample sizes for *S. maltophilia* isolates recovered from sites with high and low pollution
were large enough (Fig. 4E) to test the hypothesis of equal NumR and NumFam conditional
on pollution level. We used the robust *yuenbt*($tr = .2$, *nboot* = 998) method for independent
mean comparisons (Wilcox, 2016). The tests for NumR [$T_y = 2.1964$, $p = 0.038$; mean$_{(tr=.2)}$
difference = 1.233, $CI_{95\%} = (0.0663, 2.4004)$, *e.s.* = .43] and for NumFam [$T_y = 2.6951$, p
= 0.022; mean$_{(tr=.2)}$ difference = 0.95, $CI_{95\%} = (0.1636, 1.7364)$, *e.s.* = .5], indicating that
both variables have significantly higher values in the high-pollution sites (Figs. 4G and
4H). However, given that the 95% *CIs* of the mean NumR comparsion overlap (Fig. 4G),
and that a high number of tied (repeated) values occur (Fig. 4E), we run also Wilcox's
robust percentile bootstrap method for comparing medians [*medpb2*(*nboot* = 2000)], which
provides good control over the probability of Type I error when tied values occur. For
NumR the test could not reject the null of equal medians [$M^* = 1$, $p = .094$, $CI_{95\%} = (0,$
2.5)], but was significant for NumFam [$M^* = 1$, $p = .0495$, $CI_{95\%} = (0, 2)$].

539 3.10 *Stenotrophomonas* species differ in their β-lactamase expression patterns and
only *S. maltophilia* strains express metallo-β-lactamases

Association plots (Fig. 5) revealed a very highly significant association ($p \approx 0$) between
species and type of β-lactamases expressed (Fig. S11). Metallo-β-lactamase expression was
significantly and exclusively associated with *S. maltophilia* isolates. Most isolates from the
Smc1 and Smc2 lineages did not express any kind of β-lactamase, although expression of
extended spectrum of β-lactamases could be detected in a few isolates of these species. No
β-lactamase expression was detected in *S. terrae* isolates.

548 3.11 The prevalence of environmental *Stenotrophomonas* species recovered from
Mexican rivers is significantly associated with habitat and pollution level

We performed a multi-way association analysis to test the null hypothesis that
Stenotrophomonas species prevalence is independent of isolation habitat (sediment *vs.*
water column), the pollution level of the sampling site (Table 1) and isolation medium (Fig.
6). The test strongly rejects the null hypothesis ($p < .00001$). *S. maltophilia* was mainly
recovered on MK plates, being significantly associated with polluted sediments. The Smc1
lineage displayed a moderately significant association with clean water columns, although
some isolates could also be recovered from contaminated sediments using NAA, which is
consistent with the MCA results presented in Fig. 3. In contrast, Smc2 isolates were very
significantly overrepresented in the water columns of clean rivers and underrepresented in
sediments, suggesting a high level of ecological specialization. *S. terrae* isolates were
mainly recovered on oligotrophic NAA plates from the sediments of clean sites (Table 1).

4. Discussion

In this study we demonstrate the power of complementary phylogenetic and population genetics approaches to delimit genetically and ecologically coherent species among a diverse collection of environmental isolates of the genus *Stenotrophomonas*. Importantly, this is done based exclusively on molecular evolutionary criteria (Vinuesa et al., 2005b), without using any arbitrary sequence or phenetic similarity cut-off values, as embraced by the standard polyphasic approach that dominates current bacterial taxonomic practice (Kämpfer and Glaeser, 2012). The robustness of the species delimitations proposed here are supported by the statistically significant associations they exhibit with distinct habitat preferences, antibiotic resistance profiles, MDR status and β-lactamase expression phenotypes.

To our knowledge, this is the first study that used the multispecies coalescent (MSC) model (Rannala and Yang, 2003;Edwards et al., 2007;Degnan and Rosenberg, 2009) coupled with Bayes factor (BF) analyses (Kass and Raftery, 1995) for microbial species delimitation. The MSC model has the virtue of relaxing the implicit assumption made by the concatenation approach that the phylogeny of each gene is shared and equal to that of the species tree. Although this assumption is problematic, the concatenation approach is the current standard in microbial multilocus sequence analysis (Gevers et al., 2005;Vinuesa, 2010;Glaeser and Kampfer, 2015), including phylogenomics (Rokas et al., 2003;Wu and Eisen, 2008). It has been shown that phylogenetic estimates from concatenated datasets under coalescence are inconsistent (Kubatko and Degnan, 2007;Song et al., 2012), that is, converge to wrong solutions with higher probability as the number of concatenated loci increases. However, the impact of this inconsistency still needs to be thoroughly evaluated with clonally multiplying microbial organisms experiencing different rates of recombination (Hedge and Wilson, 2014). In our analyses, the topology of the ML phylogeny inferred from the concatenated dataset (Fig.1) is largely congruent with the Bayesian species tree inferred under the MSC (Fig S8). It is worth noting that the numbers on the branches in this type of species tree denote the estimated population sizes. That of *S. maltophilia* is about 1 order of magnitude larger than the population size estimates for the Smc1 and Smc2 genospecies (Fig. S8), which reflects the genetic heterogeneity of strains grouped in the *S. maltophilia* lineage, which includes the majority of the Mexican isolates, along with reference strains from the pubmlst.org and genome databases, isolated across the globe. Lumping this heterogeneous set of recombining sub-lineages into a single species results in coalescent events higher up in the species tree than those observed for the Smc1 and Smc2 lineages. This is consistent with the marked internal structure revealed by the Bayesian structure analysis within *S. maltophilia* (Fig. 2), which suggests that additional cryptic species may be found within the *S. maltophilia* lineage. Patil and colleagues recently proposed that the type strains of *S. africana*, *P. beteli* and *P. hibiscicola*, which are phylogenetically placed within the *S. maltophilia* clade (Fig. 1A), and have been reclassified as *S. maltophilia*, actually represent distinct species, based on their estimates of genomic average nucleotide identity values < 94% (Patil et al., 2016). In our view, these clusters represent incipient species that are still capable of recombining between them, as suggested by the admixture found in the STRUCTURE barplots (Fig. 2) and by our estimates for recombination within *S. maltophilia* (Table 5). Further investigations involving comparative and population genomics are required to identify clear signatures of

speciation within the *S. maltophilia* sub-lineages, including "speciation genes and islands"
(Shapiro et al., 2016). Despite the conservative approach taken in this study, the BF
analysis provides statistical support in favor of splitting the *S. maltophilia* complex as
currently defined in pubmlst.org into the following 5 broad evolutionary lineages: *S.*
maltophilia and the genospecies Smc1 to Smc4. Overwhelming support (ln-BF > 5) was
also obtained indicating that genogroup #10 (Vasileuskaya-Schulz et al., 2011), the sister
clade of *S. rhizophila* (genogroup #8), constitutes an independent, non-described species
(Table 2). This is of practical importance from a biotechnological perspective because it has
been argued that plant-associated *S. rhizophila* strains (Vasileuskaya-Schulz's genogroup
#8) can be safely and easily separated from *S. maltophilia* pathogens in clade IIb (Berg and
Martinez, 2015) based on 16S rRNA gene sequences and *ggpS* and *smeD* PCR-based
typing (Ribbeck-Busch et al., 2005). However, it would be important to define differences
of the former with strains in the sister genogroup #10, which holds both rape rhizosphere
and human blood and tigh bone infection isolates. In summary, our MSC-BF analysis
provided strong evidence for the existence of 5 new species in the analyzed dataset.

Considering that the MSC model implemented in *BEAST 2 was not specifically
developed for bacteria, and given that this model has been put under criticism due to
detectable model misspecification when tested on diverse empirical animal and plant
datasets using posterior predictive simulations (Reid et al., 2014), it was important to
evaluate the robustness of the Bayesian species delimitations with independent methods.
We challenged the proposed species borders within the Smc by performing well-established
population genetic analyses on our collection of environmental isolates from the sister
lineages *S. maltophilia*, Smc1 and Smc2. We focused on detecting population genetic
structure, estimating gene flow between the phylogenetically defined species, and
identifying signatures of selection. Such data and evidence are predicted by current
ecological models of bacterial speciation to reflect speciation events (Vinuesa et al.,
2005b;Koeppel et al., 2008 ;Vos, 2011;Cadillo-Quiroz et al., 2012;Shapiro and Polz, 2014).

Current models of bacterial speciation suggest that groups of closely related strains that
display some degree of resource partitioning, and consequently occupy different niches,
will be affected by independent selective sweeps caused by the gain of a beneficial gene
either by horizontal transfer or by adaptive mutation. These may sweep to fixation if
recombination levels are low in relation to selection coefficients, or form so called
"speciation islands or continents" in the genomes of highly recombining populations
(Cadillo-Quiroz et al., 2012;Shapiro et al., 2012). Such (sympatric) populations are
predicted to be discernable as sequence clusters that diverge from each other because of the
fixation of different adaptive mutations. This leads to the formation of independent
genetically and ecologically coherent units as gene flow between them gradually drops as
they diverge by means of natural selection (Vos, 2011;Shapiro and Polz, 2014). We could
show that the Mexican *S. maltophilia*, Smc1 and Smc2 lineages satisfy these predictions.
The K^*_{ST} test statistic (Hudson et al., 1992) detected highly significant genetic
differentiation between all pairs of these sympatric lineages based on DNA polymorphisms
(Table 3). This is consistent with the results from the STRUCTURE analysis. Conversely,
the number of migrants between these lineages was negligible, evidencing low levels of
gene flow between them. Additionally, the "neutrality index" (*NI*), and the results of the
MK tests suggest that positive selection, rather than drift, is the force promoting divergence

between the lineages, which is in line with predictions from the adaptive divergence model
(Vos, 2011). However, the latter interpretation needs to be considered cautiously, as the
high relative rate of between-species non-synonymous substitutions observed could also be
generated by within-species purifying selection to eliminate slightly deleterious mutations
(Hughes, 2005;Hughes et al., 2008). The latter interpretation is consistent with the observed
negative, but not significant Tajima's D values (Table 4). These are not likely to reflect a
population expansion, given the non-significant p-values of the powerful R_2 statistic for
population growth, which is well suited for small sample sizes such as those of the Smc1
and Smc2 lineages (Ramos-Onsins and Rozas, 2002). We could show that recombination is
an important force, providing genetic cohesion to these lineages, with *Rho/theta* estimates
ranging from 1.11 in Smc1 to nearly 3 in *S. maltophilia*. Since recombination events are
only detectable when a tract of multiple polymorphisms are introduced in a population, it is
clear that most of the observed polymorphisms within the analyzed populations originate
from recombination rather than point mutations. The high recombination levels detected
within the *S. maltophilia* lineage suggests that speciation within this group is an ongoing,
possibly not yet finished process, along a "spectrum" of speciation, resulting in "fuzzy"
borders between the sub-lineages (Hanage et al., 2005;Shapiro et al., 2016). However, these
can be already detected as phylogenetic and STRUCTURE clusters, even with the limited
resolving power provided by the 7 gene MLST scheme used.

As predicted by the ecological speciation models recently developed for bacteria (Koeppel
et al., 2008;Vos, 2011;Shapiro and Polz, 2014), the marked genetic differentiation detected
between the sympatric *S. maltophilia*, Smc1 and Smc2 environmental populations is
significantly associated with different habitat preferences, antibiotic susceptibility profiles
and β-lactamase expression phenotypes. These attributes strongly suggest that these
lineages have differentiated ecological niches. The significant differences in habitat
preferences could provide some micro-geographic separation between the populations
coexisting in the same river, which might partly explain the reduced gene flow measured
between them, despite of being sister lineages, contributing to their genetic differentiation.
Similar patterns have been reported for other aquatic microbes such as *Desulfolobus*
(Oakley et al., 2010), *Exiguobacterium* (Rebollar et al., 2012) and *Vibrio* (Shapiro et al.,
2012;Friedman et al., 2013). Consequently, our results support the growing body of
evidence pointing to niche partitioning as a major factor promoting evolutionary divergence
between closely related sympatric prokaryotic populations, even when they exhibit high
levels of recombination (Shapiro and Polz, 2014).

As noted before, *S. maltophilia* is well-known as an emergent opportunistic multidrug-
resistant (MDR) nosocomial pathogen, causing increasing morbidity and mortality (Looney
et al., 2009;Brooke, 2012). Comparative genomics and functional analyses have clearly
established that the MDR or extensively drug-resistant (XDR) phenotype displayed by
clinical isolates of this species is largely intrinsic, resulting from the expression of a
combination of several types of efflux pumps (RND, MATE, MFS and ABC types) and
diverse chromosomally-encoded antibiotic resistance genes [*aph (3')-IIc*, *aac(6')-lz* and
Sm*qnr*], including the metallo-beta-lactamase *blaL1* and the inducible Ambler class A beta-
lactamase *blaL2* (Crossman et al., 2008;Brooke, 2014;García-León et al., 2014;Sanchez,
2015;Youenou et al., 2015). However, contradictory results have been reported regarding
the MDR status of environmental isolates of the Smc. For example, a recent ecological

study of a large collection isolates classified as *S. maltophilia* recovered from diverse
agricultural soils in France and Tunisa concluded that they display a high diversity of
antibiotic resistance profiles, expressing resistance against 1 to 12 antibiotics, with clinical
and manure isolates expressing the highest numbers (Deredjian et al., 2016). These isolates
were vaguely classified as *S. maltophilia* based on growth on the selective VIA isolation
medium (Kerr et al., 1996) and PCR detection of the *smeD* gene (Pinot et al., 2011). We
argue that the large phenotypic variance observed in that and similar studies result from the
lack of proper species delimitation. This cannot be achieved with such coarse typing
methods, most likely resulting in the lumping of multiple species into *S. maltophilia*. In
contrast, in the present study we found a very strong statistical association between the
MDR condition and metallo-β-lactamase (MBL) production with the *S. maltophilia* lineage,
whereas the sibling Smc1 and Smc2 genospecies were found to express on average
resistance to < 3 antibiotic families (Figs. 4B and 4D and S13B), most strains not
expressing any kind of β-lactamase, and none expressing MBLs (Fig. 5). Consequently,
intrinsic MDR can only be assumed for the *S. maltophilia* strains of clinical and
environmental origin.

In conclusion, the results presented here provide the first in depth and integrative molecular
systematic, evolutionary genetic and ecological analysis of the genus *Stenotrophomonas*.
The study demonstrates that both phylogenetic and population genetic approaches are
necessary for robust delimitation of natural species borders in bacteria. Failure to properly
delimit such lineages hinders downstream ecological and functional analysis of species.
Comparative and population genomic studies are required to resolve pending issues
regarding the speciation status of the sub-lineages within the *S. maltophilia*.

5. Conflict of Interest Statement: The authors declare that the research was conducted in
the absence of any commercial or financial relationships that could be construed as a
potential conflict of interest.

6. Author Contributions

726 All authors read and approved the manuscript. LEOS and PV conceived and designed the
727 project. LEOS generated the collection of isolates, performed wet-lab experiments and
728 analyzed resistance phenotypes. PV performed bioinformatics, statistical and evolutionary
729 genetic analyses. PV wrote the manuscript.

7. Funding

This work is part of LEOS's PhD project in the Programa de Doctorado en Ciencias
Biomédicas, Universidad Nacional Autónoma de México, and was supported by a PhD
scholarship from Consejo Nacional de Ciencia y Tecnología (CONACyT-México) and
student travel scholarships from PAEP-UNAM. We gratefully acknowledge financial

support obtained from CONACYT-México (grant 179133) and DGAPA-PAPIIT/UNAM
(grant IN211814) to PV.

8. Acknowledgements

We gratefully thank Dr. Eria Rebollar and Dr. Claudia Silva for their critical reading of the
manuscript. Javier Rivera Campos is acknowledged for technical support with wet-lab
experiments. Antonio Trujillo from CCG-UNAM is thanked for support with field work.
Don Juan Alvear Gutiérrez and Ing.Norberto Bahena are gratefully acknowledged for
supporting our sampling at the natural parks Los Sauces and Las Estacas, respectively. José
Alfredo Hernández and the UATI at CCG-UNAM are acknowledged for support with
Linux server administration. Dr. Jesús Silva Sánchez from the INSP in Cuernavaca,
Mexico, is gratefully thanked for his support throughout the work, particularly regarding
the interpretation of disk-diffusion assays and for providing laboratory reagents.

9. References

Aagot, N., Nybroe, O., Nielsen, P., and Johnsen, K. (2001). An altered *Pseudomonas* diversity is
recovered from soil by using nutrient-poor *Pseudomonas*-selective soil extract media.
Appl. Environ. Microbiol. 67**,** 5233-5239. doi: 10.1128/AEM.67.11.5233-5239.2001.
Adamek, M., Overhage, J., Bathe, S., Winter, J., Fischer, R., and Schwartz, T. (2011). Genotyping of
environmental and clinical *Stenotrophomonas maltophilia* isolates and their pathogenic
potential. *PLoS One* 6**,** e27615. doi: 10.1371/journal.pone.0027615.
Apha (ed.). (2005). *Standard methods for the analysis of water and wastewater—Section 9222D.*
Thermotolerant (fecal) coliform membrane filter procedure. Washington, D.C.: American
Public Health Association.
Assih, E.A., Ouattara, A.S., Thierry, S., Cayol, J.L., Labat, M., and Macarie, H. (2002).
Stenotrophomonas acidaminiphila sp. nov., a strictly aerobic bacterium isolated from an
upflow anaerobic sludge blanket (UASB) reactor. *Int. J. Syst. Evol. Microbiol.* 52**,** 559-568.
doi: 10.1099/00207713-52-2-559.
Aydin, Z., Marcussen, T., Ertekin, A.S., and Oxelman, B. (2014). Marginal likelihood estimate
comparisons to obtain optimal species delimitations in Silene sect. Cryptoneurae
(Caryophyllaceae). *PLoS One* 9**,** e106990. 10.1371/journal.pone.0106990
PONE-D-14-10584 [pii].
Baele, G., Lemey, P., Bedford, T., Rambaut, A., Suchard, M.A., and Alekseyenko, A.V. (2012).
Improving the accuracy of demographic and molecular clock model comparison while
accommodating phylogenetic uncertainty. *Mol Biol Evol* 29**,** 2157-2167. doi:
10.1093/molbev/mss084.
Baele, G., Li, W.L., Drummond, A.J., Suchard, M.A., and Lemey, P. (2013). Accurate model selection
of relaxed molecular clocks in bayesian phylogenetics. *Mol. Biol. Evol.* 30**,** 239-243. doi:
10.1093/molbev/mss243.
Berg, G., and Martinez, J.L. (2015). Friends or foes: can we make a distinction between beneficial
and harmful strains of the *Stenotrophomonas maltophilia* complex? *Front. Microbiol.* 6**,**
241. doi: 10.3389/fmicb.2015.00241.

Berg, G., Roskot, N., and Smalla, K. (1999). Genotypic and phenotypic relationships between
clinical and environmental isolates of *Stenotrophomonas maltophilia*. *J. Clin. Microbiol.* 37**,**
3594-3600.
Bouckaert, R., Heled, J., Kuhnert, D., Vaughan, T., Wu, C.H., Xie, D., Suchard, M.A., Rambaut, A.,
and Drummond, A.J. (2014). BEAST 2: a software platform for Bayesian evolutionary
analysis. *PLoS Comput Biol* 10**,** e1003537. doi: 10.1371/journal.pcbi.1003537.
Bouckaert, R.R. (2010). DensiTree: making sense of sets of phylogenetic trees. *Bioinformatics* 26**,**
1372-1373. doi: 10.1093/bioinformatics/btq110.
Brooke, J.S. (2012). *Stenotrophomonas maltophilia*: an emerging global opportunistic pathogen.
Clin. Microbiol. Rev. 25**,** 2-41. doi: 10.1128/CMR.00019-11.
Brooke, J.S. (2014). New strategies against *Stenotrophomonas maltophilia*: a serious worldwide
intrinsically drug-resistant opportunistic pathogen. *Expert Rev. Anti Infect. Ther.* 12**,** 1-4.
doi: 10.1586/14787210.2014.864553.
Cadillo-Quiroz, H., Didelot, X., Held, N.L., Herrera, A., Darling, A., Reno, M.L., Krause, D.J., and
Whitaker, R.J. (2012). Patterns of gene flow define species of thermophilic *Archaea*. *PLoS*
Biol. 10**,** e1001265. doi: 10.1371/journal.pbio.1001265.
Clsi (2016). *Clinical and Laboratory Standards Institute (CLSI) Performance Standards for*
Antimicrobial susceptibility testing. Wayne, Pennsylvania Clinical and Laboratory Standards
Institute.
Coenye, T., Vanlaere, E., Falsen, E., and Vandamme, P. (2004). *Stenotrophomonas africana*
Drancourt et al. 1997 is a later synonym of *Stenotrophomonas maltophilia* (Hugh 1981)
Palleroni and Bradbury 1993. *Int. J. Syst. Evol. Microbiol.* 54**,** 1235-1237. doi:
10.1099/ijs.0.63093-0.
Contreras-Moreira, B., Sachman-Ruiz, B., Figueroa-Palacios, I., and Vinuesa, P. (2009).
primers4clades: a web server that uses phylogenetic trees to design lineage-specific PCR
primers for metagenomic and diversity studies. *Nucleic Acids Res.* 37**,** W95-W100. gkp377
doi: 10.1093/nar/gkp377.
Contreras-Moreira, B., and Vinuesa, P. (2013). GET_HOMOLOGUES, a versatile software package
for scalable and robust microbial pangenome analysis. *Appl. Environ. Microbiol.* 79**,** 7696-
7701. doi: 10.1128/AEM.02411-13.
Cordero, O.X., and Polz, M.F. (2014). Explaining microbial genomic diversity in light of evolutionary
ecology. *Nat. Rev. Microbiol.* 12**,** 263-273. doi: 10.1038/nrmicro3218.
Crossman, L.C., Gould, V.C., Dow, J.M., Vernikos, G.S., Okazaki, A., Sebaihia, M., Saunders, D.,
Arrowsmith, C., Carver, T., Peters, N., Adlem, E., Kerhornou, A., Lord, A., Murphy, L.,
Seeger, K., Squares, R., Rutter, S., Quail, M.A., Rajandream, M.A., Harris, D., Churcher, C.,
Bentley, S.D., Parkhill, J., Thomson, N.R., and Avison, M.B. (2008). The complete genome,
comparative and functional analysis of *Stenotrophomonas maltophilia* reveals an organism
heavily shielded by drug resistance determinants. *Genome Biol.* 9**,** R74. doi: 10.1186/gb-
2008-9-4-r74.
Chang, Y.T., Lin, C.Y., Chen, Y.H., and Hsueh, P.R. (2015). Update on infections caused by
Stenotrophomonas maltophilia with particular attention to resistance mechanisms and
therapeutic options. *Front. Microbiol.* 6**,** 893. doi: 10.3389/fmicb.2015.00893.
Darriba, D., Taboada, G.L., Doallo, R., and Posada, D. (2012). jModelTest 2: more models, new
heuristics and parallel computing. *Nat. Methods* 9**,** 772. nmeth.2109 [pii]
10.1038/nmeth.2109.
Davenport, K.W., Daligault, H.E., Minogue, T.D., Broomall, S.M., Bruce, D.C., Chain, P.S., Coyne,
S.R., Gibbons, H.S., Jaissle, J., Li, P.E., Rosenzweig, C.N., Scholz, M.B., and Johnson, S.L.

825   (2014). Complete genome sequence of *Stenotrophomonas maltophilia* type strain 810-2
826   (ATCC 13637). *Genome Announc.* 2. 2/5/e00974-14. doi: 10.1128/genomeA.00974-14.
827   De La Bastide, M., and Mccombie, W.R. (2007). Assembling genomic DNA sequences with PHRAP.
828   *Curr. Protoc. Bioinformatics* Chapter 11**,** Unit11 14. doi: 10.1002/0471250953.bi1104s17.
829   Degnan, J.H., and Rosenberg, N.A. (2006). Discordance of species trees with their most likely gene
830   trees. *PLoS Genet.* 2**,** e68.
831   Degnan, J.H., and Rosenberg, N.A. (2009). Gene tree discordance, phylogenetic inference and the
832   multispecies coalescent. *Trends Ecol. Evol.* 24**,** 332-340. doi: 10.1016/j.tree.2009.01.009.
833   Deredjian, A., Alliot, N., Blanchard, L., Brothier, E., Anane, M., Cambier, P., Jolivet, C., Khelil, M.N.,
834   Nazaret, S., Saby, N., Thioulouse, J., and Favre-Bonte, S. (2016). Occurrence of
835   *Stenotrophomonas maltophilia* in agricultural soils and antibiotic resistance properties.
836   *Res. Microbiol.* S0923-2508(16)00006-1 [pii]
837   10.1016/j.resmic.2016.01.001.
838   Didelot, X., and Wilson, D.J. (2015). ClonalFrameML: efficient inference of recombination in whole
839   bacterial genomes. *PLoS Comput. Biol.* 11**,** e1004041. doi: 10.1371/journal.pcbi.1004041.
840   Drummond, A.J., Ho, S.Y., Phillips, M.J., and Rambaut, A. (2006). Relaxed phylogenetics and dating
841   with confidence. *PLoS Biol.* 4**,** e88. doi: 10.1371/journal.pbio.0040088.
842   Edwards, S.V., Liu, L., and Pearl, D.K. (2007). High-resolution species trees without concatenation.
843   *Proc Natl Acad Sci U S A* 104**,** 5936-5941. doi: 10.1073/pnas.0607004104.
844   Evanno, G., Regnaut, S., and Goudet, J. (2005). Detecting the number of clusters of individuals
845   using the software STRUCTURE: a simulation study. *Mol. Ecol.* 14**,** 2611-2620. doi:
846   10.1111/j.1365-294X.2005.02553.x.
847   Falush, D., Stephens, M., and Pritchard, J.K. (2003). Inference of Population Structure Using
848   Multilocus Genotype Data. Linked loci and correlated allele frequencies. *Genetics* 164**,**
849   1567-1587.
850   Falush, D., Stephens, M., and Pritchard, J.K. (2007). Inference of population structure using
851   multilocus genotype data: dominant markers and null alleles. *Mol. Ecol. Notes.* 7**,** 574-578.
852   doi: 10.1111/j.1471-8286.2007.01758.x.
853   Friedman, J., Alm, E.J., and Shapiro, B.J. (2013). Sympatric speciation: when is it possible in
854   bacteria? *PLoS One* 8**,** e53539. doi: 10.1371/journal.pone.0053539.
855   Fujita, M.K., Leache, A.D., Burbrink, F.T., Mcguire, J.A., and Moritz, C. (2012). Coalescent-based
856   species delimitation in an integrative taxonomy. *Trends Ecol. Evol.* 27**,** 480-488. doi:
857   10.1016/j.tree.2012.04.012.
858   García-León, G., Salgado, F., Oliveros, J.C., Sánchez, M.B., and Martínez, J.L. (2014). Interplay
859   between intrinsic and acquired resistance to quinolones in *Stenotrophomonas maltophilia*.
860   *Environ. Microbiol.* 16**,** 1282-1296. 10.1111/1462-2920.12408.
861   Gevers, D., Cohan, F.M., Lawrence, J.G., Spratt, B.G., Coenye, T., Feil, E.J., Stackebrandt, E., Van De
862   Peer, Y., Vandamme, P., Thompson, F.L., and Swings, J. (2005). Opinion: Re-evaluating
863   prokaryotic species. *Nat. Rev. Microbiol.* 3**,** 733-739.
864   Glaeser, S.P., and Kampfer, P. (2015). Multilocus sequence analysis (MLSA) in prokaryotic
865   taxonomy. *Syst. Appl. Microbiol.* 38**,** 237-245. doi: 10.1016/j.syapm.2015.03.007.
866   Grummer, J.A., Bryson, R.W., Jr., and Reeder, T.W. (2014). Species delimitation using Bayes
867   factors: simulations and application to the Sceloporus scalaris species group (Squamata:
868   Phrynosomatidae). *Syst. Biol.* 63**,** 119-133. syt069 [pii]
869   10.1093/sysbio/syt069.
870   Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New
871   algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
872   performance of PhyML 3.0. *Syst. Biol.* 59**,** 307-321. doi: 10.1093/sysbio/syq010.

Hanage, W.P., Fraser, C., and Spratt, B.G. (2005). Fuzzy species among recombinogenic bacteria.
BMC Biol. 3**,** 6.

Handa, Y., Tazato, N., Nagatsuka, Y., Koide, T., Kigawa, R., C., S., and Sugiyama, J. (2016).
Stenotrophomonas tumulicola sp. nov., a major contaminant of the stone chamber interior
in the Takamatsuzuka Tumulus. *Int. J. Syst. Evol. Microbiol.* 66**,** 1119–1124. doi:
10.1099/ijsem.0.000843.

Hauben, L., Vauterin, L., Moore, E.R., Hoste, B., and Swings, J. (1999). Genomic diversity of the
genus *Stenotrophomonas*. *Int. J. Syst. Bacteriol.* 49 Pt 4**,** 1749-1760. doi:
10.1099/00207713-49-4-1749.

Hedge, J., and Wilson, D.J. (2014). Bacterial phylogenetic reconstruction from whole genomes is
robust to recombination but demographic inference is not. *MBio* 5**,** e02158. mBio.02158-
14. doi: 10.1128/mBio.02158-14.

Heled, J., and Drummond, A.J. (2010). Bayesian inference of species trees from multilocus data.
Mol. Biol. Evol. 27**,** 570-580. doi: 10.1093/molbev/msp274.

Heylen, K., Vanparys, B., Peirsegaele, F., Lebbe, L., and De Vos, P. (2007). *Stenotrophomonas terrae*
sp. nov. and *Stenotrophomonas humi* sp. nov., two nitrate-reducing bacteria isolated from
soil. *Int. J. Syst. Evol. Microbiol.* 57**,** 2056-2061. doi: 10.1099/ijs.0.65044-0.

Hudson, R.R., Boos, D.D., and Kaplan, N.L. (1992). A statistical test for detecting geographic
subdivision. *Mol. Biol. Evol.* 9**,** 138-151.

Hughes, A.L. (2005). Evidence for abundant slightly deleterious polymorphisms in bacterial
populations. *Genetics* 169**,** 533-538. doi: 10.1534/genetics.104.036939.

Hughes, A.L., Friedman, R., Rivailler, P., and French, J.O. (2008). Synonymous and nonsynonymous
polymorphisms versus divergences in bacterial genomes. *Mol. Biol. Evol.* 25**,** 2199-2209.
doi: 10.1093/molbev/msn166.

Jawad, A., Hawkey, P.M., Heritage, J., and Snelling, A.M. (1994). Description of Leeds
Acinetobacter Medium, a new selective and differential medium for isolation of clinically
important *Acinetobacter* spp., and comparison with Herellea agar and Holton's agar. *J.*
Clin. Microbiol. 32**,** 2353-2358.

Kaiser, O., Puhler, A., and Selbitschka, W. (2001). Phylogenetic Analysis of Microbial Diversity in
the Rhizoplane of Oilseed Rape (Brassica napus cv. Westar) Employing Cultivation-
Dependent and Cultivation-Independent Approaches. *Microb Ecol* 42**,** 136-149.

Kaiser, S., Biehler, K., and Jonas, D. (2009). A *Stenotrophomonas maltophilia* multilocus sequence
typing scheme for inferring population structure. *J. Bacteriol.* 191**,** 2934-2943. doi:
10.1128/JB.00892-08.

Kämpfer, P., and Glaeser, S.P. (2012). Prokaryotic taxonomy in the sequencing era--the polyphasic
approach revisited. *Environ. Microbiol.* 14**,** 291-317. doi: 10.1111/j.1462-
2920.2011.02615.x.

Kass, R.E., and Raftery, A.E. (1995). Bayes factors. *J. Am. Stat. Assoc.* 90**,** 773-795.

Kerr, K.G., Denton, M., Todd, N., Corps, C.M., Kumari, P., and Hawkey, P.M. (1996). A new selective
differential medium for isolation of *Stenotrophomonas maltophilia*. *Eur. J. Clin. Microbiol.*
Infect. Dis. 15**,** 607-610.

Koeppel, A., Perry, E.B., Sikorski, J., Krizanc, D., Warner, A., Ward, D.M., Rooney, A.P., Brambilla, E.,
Connor, N., Ratcliff, R.M., Nevo, E., and Cohan, F.M. (2008). Identifying the fundamental
units of bacterial diversity: a paradigm shift to incorporate ecology into bacterial
systematics. *Proc. Natl. Acad. Sci. U. S. A.* 105**,** 2504-2509. doi: 10.1073/pnas.0712205105.

Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A., and Mayrose, I. (2015). Clumpak: a
program for identifying clustering modes and packaging population structure inferences
across K. *Mol. Ecol. Resour.* 15**,** 1179-1191. doi: 10.1111/1755-0998.12387.

Kubatko, L.S., and Degnan, J.H. (2007). Inconsistency of phylogenetic estimates from concatenated
data under coalescence. *Syst. Biol.* 56**,** 17-24.
Lartillot, N., and Philippe, H. (2006). Computing Bayes factors using thermodynamic integration.
Syst. Biol. 55**,** 195-207. doi: 10.1080/10635150500433722.
Lira, F., Hernandez, A., Belda, E., Sanchez, M.B., Moya, A., Silva, F.J., and Martinez, J.L. (2012).
Whole-genome sequence of *Stenotrophomonas maltophilia* D457, a clinical isolate and a
model strain. *J. Bacteriol.* 194**,** 3563-3564. doi: 10.1128/JB.00602-12.
Looney, W.J., Narita, M., and Muhlemann, K. (2009). *Stenotrophomonas maltophilia*: an emerging
opportunist human pathogen. *Lancet Infect. Dis.* 9**,** 312-323.doi: 10.1016/S1473-
3099(09)70083-0.
Mcdonald, J.H., and Kreitman, M. (1991). Adaptive protein evolution at the Adh locus in
Drosophila. *Nature* 351**,** 652-654. doi: 10.1038/351652a0.
Oakley, B.B., Carbonero, F., Van Der Gast, C.J., Hawkins, R.J., and Purdy, K.J. (2010). Evolutionary
divergence and biogeography of sympatric niche-differentiated bacterial populations.
ISME J. 4**,** 488-497. doi: 10.1038/ismej.2009.146.
Palleroni, N.J., and Bradbury, J.F. (1993). *Stenotrophomonas*, a new bacterial genus for
Xanthomonas maltophilia (Hugh 1980) Swings et al. 1983. *Int. J. Syst. Bacteriol.* 43**,** 606-
609. doi: 10.1099/00207713-43-3-606.
Patil, P.P., Midha, S., Kumar, S., and Patil, P.B. (2016). Genome Sequence of Type Strains of Genus
Stenotrophomonas. *Front. Microbiol.* 7**,** 309. doi: 10.3389/fmicb.2016.00309.
Pinot, C., Deredjian, A., Nazaret, S., Brothier, E., Cournoyer, B., Segonds, C., and Favre-Bonte, S.
(2011). Identification of *Stenotrophomonas maltophilia* strains isolated from
environmental and clinical samples: a rapid and efficient procedure. *J. Appl. Microbiol.*
111**,** 1185-1193. doi: 10.1111/j.1365-2672.2011.05120.x.
Pritchard, J.K., Stephens, M., and Donnelly, P. (2000). Inference of population structure using
multilocus genotype data. *Genetics* 155**,** 945-959.
R Development Core Team (2016). "R: A Language and Environment for Statistical Computing.
https://www.R-project.org." (Vienna, Austria).
Ramos-Onsins, S.E., and Rozas, J. (2002). Statistical properties of new neutrality tests against
population growth. *Mol. Biol. Evol.* 19**,** 2092-2100.
Ramos, P.L., Van Trappen, S., Thompson, F.L., Rocha, R.C., Barbosa, H.R., De Vos, P., and Moreira-
Filho, C.A. (2010). Screening for endophytic nitrogen-fixing bacteria in Brazilian sugar cane
varieties used in organic farming and description of *Stenotrophomonas pavanii* sp. nov.
Int. J. Syst. Evol. Microbiol. 61**,** 926-931. doi: 10.1099/ijs.0.019372-0.
Rand, D.M., and Kann, L.M. (1996). Excess amino acid polymorphism in mitochondrial DNA:
contrasts among genes from *Drosophila*, mice, and humans. *Mol. Biol. Evol.* 13**,** 735-748.
Rannala, B., and Yang, Z. (2003). Bayes estimation of species divergence times and ancestral
population sizes using DNA sequences from multiple loci. *Genetics* 164**,** 1645-1656.
Rebollar, E.A., Avitia, M., Eguiarte, L.E., Gonzalez-Gonzalez, A., Mora, L., Bonilla-Rosso, G., and
Souza, V. (2012). Water-sediment niche differentiation in ancient marine lineages of
Exiguobacterium endemic to the Cuatro Cienegas Basin. *Environ. Microbiol.* 14**,** 2323-2333.
doi: 10.1111/j.1462-2920.2012.02784.x.
Reid, N.M., Hird, S.M., Brown, J.M., Pelletier, T.A., Mcvay, J.D., Satler, J.D., and Carstens, B.C.
(2014). Poor fit to the multispecies coalescent is widely detectable in empirical data. *Syst.*
Biol. 63**,** 322-333. doi: 10.1093/sysbio/syt057.
Ribbeck-Busch, K., Roder, A., Hasse, D., De Boer, W., Martinez, J.L., Hagemann, M., and Berg, G.
(2005). A molecular biological protocol to distinguish potentially human pathogenic

968   *Stenotrophomonas maltophilia* from plant-associated *Stenotrophomonas rhizophila.*
969   *Environ. Microbiol.* 7**,** 1853-1858. EMI928 [pii]
970 10.1111/j.1462-2920.2005.00928.x.
971 Rokas, A., Williams, B.L., King, N., and Carroll, S.B. (2003). Genome-scale approaches to resolving
972   incongruence in molecular phylogenies. *Nature* 425**,** 798-804.
973 Rosenberg, N.A. (2013). Discordance of species trees with their most likely gene trees: a unifying
974   principle. *Mol. Biol. Evol.* 30**,** 2709-2713. doi: 10.1093/molbev/mst160.
975 Rozas, J., Sánchez-Delbarrio, J.C., Messeguer, X., and Rozas, R. (2003). DnaSP, DNA polymorphism
976   analyses by the coalescent and other methods. *Bioinformatics* 19**,** 2496-2497.
977 Ryan, R.P., Monchy, S., Cardinale, M., Taghavi, S., Crossman, L., Avison, M.B., Berg, G., Van Der
978   Lelie, D., and Dow, J.M. (2009). The versatility and adaptation of bacteria from the genus
979   *Stenotrophomonas*. *Nat. Rev. Microbiol.* 7**,** 514-525. doi: 10.1038/nrmicro2163.
980 Sanchez, M.B. (2015). Antibiotic resistance in the opportunistic pathogen *Stenotrophomonas*
981   *maltophilia*. *Front. Microbiol.* 6**,** 658. doi: 10.3389/fmicb.2015.00658.
982 Shapiro, B.J., Friedman, J., Cordero, O.X., Preheim, S.P., Timberlake, S.C., Szabo, G., Polz, M.F., and
983   Alm, E.J. (2012). Population genomics of early events in the ecological differentiation of
984   bacteria. *Science* 336**,** 48-51. doi: 10.1126/science.1218198.
985 Shapiro, B.J., Leducq, J.B., and Mallet, J. (2016). What Is Speciation? *PLoS Genet.* 12**,** e1005860.
986   doi: 10.1371/journal.pgen.1005860.
987 Shapiro, B.J., and Polz, M.F. (2014). Ordering microbial diversity into ecologically and genetically
988   cohesive units. *Trends Microbiol.* 22**,** 235-247. doi: 10.1016/j.tim.2014.02.006.
989 Song, S., Liu, L., Edwards, S.V., and Wu, S. (2012). Resolving conflict in eutherian mammal
990   phylogeny using phylogenomics and the multispecies coalescent model. *Proc. Natl. Acad.*
991   *Sci. U. S. A.* 109**,** 14942-14947. doi: 10.1073/pnas.1211733109.
992 Svensson-Stadler, L.A., Mihaylova, S.A., and Moore, E.R. (2011). *Stenotrophomonas* interspecies
993   differentiation and identification by *gyrB* sequence analysis. *FEMS Microbiol. Lett.* 327**,** 15-
994   24. doi: 10.1111/j.1574-6968.2011.02452.x.
995 Ultee, A., Souvatzi, N., Maniadi, K., and Konig, H. (2004). Identification of the culturable and
996   nonculturable bacterial population in ground water of a municipal water supply in
997   Germany. *J. Appl. Microbiol.* 96**,** 560-568.
998 Usepa (2002). "Method 1603: *Escherichia coli* (*E.coli*) in Water by Membrane Filtration Using
999   Modified membrane-Thermotolerant *Escherichia coli* Agar (Modified mTEC) (September
1000   2002)"
1001 Van Den Mooter, M., and Swings, J. (1990). Numerical analysis of 295 phenotypic features of 266
1002   *Xanthomonas* strain and related strains and an improved taxonomy of the genus. *Int. J.*
1003   *Syst. Bacteriol.* 40**,** 348-369.
1004 Vasileuskaya-Schulz, Z., Kaiser, S., Maier, T., Kostrzewa, M., and Jonas, D. (2011). Delineation of
1005   *Stenotrophomonas* spp. by multi-locus sequence analysis and MALDI-TOF mass
1006   spectrometry. *Syst. Appl. Microbiol.* 34**,** 35-39. doi: 10.1016/j.syapm.2010.11.011.
1007 Vinuesa, P. (2010). "Multilocus Sequence Analysis and Bacterial Species Phylogeny Estimation," in
1008   *Molecular Phylogeny of Microorganisms,* eds. A. Oren & R.T. Papke. Caister Academic
1009   Press), 41-64.
1010 Vinuesa, P., and Ochoa-Sánchez, L.E. (2015). Complete Genome Sequencing of *Stenotrophomonas*
1011   *acidaminiphila* ZAC14D2_NAIMI4_2, a Multidrug-Resistant Strain Isolated from Sediments
1012   of a Polluted River in Mexico, Uncovers New Antibiotic Resistance Genes and a Novel
1013   Class-II Lasso Peptide Biosynthesis Gene Cluster. *Genome Announc.* 3**,** e01433-01415. doi:
1014   10.1128/genomeA.01433-15.

Vinuesa, P., Rojas-Jimenez, K., Contreras-Moreira, B., Mahna, S.K., Prasad, B.N., Moe, H., Selvaraju,
S.B., Thierfelder, H., and Werner, D. (2008). Multilocus sequence analysis for assessment
of the biogeography and evolutionary genetics of four *Bradyrhizobium* species that
nodulate soybeans on the Asiatic continent. *Appl. Environ. Microbiol.* 74**,** 6987-6996.
Vinuesa, P., Silva, C., Lorite, M.J., Izaguirre-Mayoral, M.L., Bedmar, E.J., and Martínez-Romero, E.
(2005a). Molecular systematics of rhizobia based on maximum likelihood and Bayesian
phylogenies inferred from *rrs*, *atpD*, *recA* and *nifH* sequences, and their use in the
classification of *Sesbania* microsymbionts from Venezuelan wetlands. *Syst. Appl.*
Microbiol. 28**,** 702-716.
Vinuesa, P., Silva, C., Werner, D., and Martínez-Romero, E. (2005b). Population genetics and
phylogenetic inference in bacterial molecular systematics: the roles of migration and
recombination in *Bradyrhizobium* species cohesion and delineation. *Mol. Phylogenet. Evol.*
34**,** 29-54.
Vos, M. (2011). A species concept for bacteria based on adaptive divergence. *Trends Microbiol.* 19**,**
1-7. S0966-842X(10)00187-3. doi: 10.1016/j.tim.2010.10.003.
Weisburg, W.G., Barns, S.M., Pelletie, D.A., and Lane, D.J. (1991). 16S ribosomal DNA amplification
for phylogenetic study. *J. Bacteriol.* 173**,** 697-703.
Wilcox, R. (2016). *Introduction to Robust Estimation and Hypothesis Testing, 4th ed.* New York:
Academic Press.
Wu, M., and Eisen, J.A. (2008). A simple, fast, and accurate method of phylogenomic inference.
Genome Biol. 9**,** R151.
Yi, H., Srinivasan, S., and Kim, M.K. (2010). *Stenotrophomonas panacihumi* sp. nov., isolated from
soil of a ginseng field. *J. Microbiol.* 48**,** 30-35. doi: 10.1007/s12275-010-0006-0.
Youenou, B., Favre-Bonte, S., Bodilis, J., Brothier, E., Dubost, A., Muller, D., and Nazaret, S. (2015).
Comparative Genomics of Environmental and Clinical *Stenotrophomonas maltophilia*
Strains with Different Antibiotic Resistance Profiles. *Genome Biol. Evol.* 7**,** 2484-2505.
evv161. doi: 10.1093/gbe/evv161.
Zhu, B., Liu, H., Tian, W.X., Fan, X.Y., Li, B., Zhou, X.P., Jin, G.L., and Xie, G.L. (2012). Genome
sequence of *Stenotrophomonas maltophilia* RR-10, isolated as an endophyte from rice
root. *J. Bacteriol.* 194**,** 1280-1281. doi: 10.1128/JB.06702-11.

1045
1046
1047

Table 1. Sampling sites for this study in Morelos, Mexico and pollution level based on counts of thermotolerant fecal coliforms (TTFCs) and thermotolerant *E. coli* (TTEc) colony forming units (cfu/100 ml) measured in the water column.

River	Site name	code	coordinates[a]	TTFCs[b]	TTEc[c]	Pol.level[c]
Apatlaco	Temixco	TEX	18°51'14.4"N 99°13'20.4"W	5.66e5	3.6e5	H
Apatlaco	Zacatepec	ZAC	18°38'23.4"N 99°11'44.5"W	5.34e5	3.2e5	H
Sauces	Los Sauces (river)	SAU	18°41'50.0"N 99°07'45.7"W	240	97	L
Sauces	Los Sauces (flooded soil)	SAUr	18°41'50.0"N 99°07'45.7"W	966	166	L
Estacas	Las Estacas	EST	18°43'57.5"N 99°06'48.3"W	83	17	L
Yautepec	Bonifiacio García	YAU	18°43'24.0"N 99°06'53.3"W	1.96e3	833	I

[a]See supplementary Figure S1 for maps and pictures of the sampling sites.
[b]Thermo-tolerant fecal coliform counts were performed on mFC agar using the membrane-filtration method.
[c]Thermo-tolerant *E. coli* counts were performed on mTEC agar using the membrane-filtration method.
[d]Pollution level, coded as high (H > 1000), intermediate (200 < I < 1000) and low (L < 200), based on the colony forming units (cfu) of thermo-tolerant *E. coli*.

Table 2. Bayes factor (BF) analysis for 5 species delimitation hypotheses within the *S. maltophilia* complex, plus genogroups #8 and #10, based on marginal likelihoods computed for each hypothesis by path sampling (see methods).

Species model[a]	marginal ln$L$ estimate	Model rank	ln-BF[b] (vs. best)	ln-BF[c] (vs. previous)
split1	-31347.16515	1 (best)	NA	NA
lump_Smal+Smc1	-31384.34206	2	4.308835**	4.308835** (2 vs. 1)
lump_Smal +Smc12	-31498.71633	3	5.714071***	5.432623*** (3 vs. 2)
lump_Smal +Smc123	-31521.18269	4	5.852303***	3.805166** (4 vs. 3)
lump_Smal +Smc1234	-31593.24923	5	6.19882***	4.970737** (5 vs. 4)

[a]The best (split1) model assumes that genogroup #10 is a sister clade of *S. rhizophila* (Fig. 1A) and splits the Smc into *S. maltophilia* (Smal) and four additional species clades (Smc1-Smc4), according to Fig. 1B. The following models consecutively lump the *S. maltophilia* clade with the Smc1-Smc4 clades, with lump_*S. maltophilia*+Smc1234 representing the whole *S. maltophilia* complex as a single species (Figs 1A and 1B). The marginal ln$L$ for the Lump_clades_#8+#10 model is -31465.00904, resulting in a ln-BF = 5.462508*** when compared against the split1 (best) model.

[b]The ln-Bayes factors are computed based on the marginal likelihood estimates ln-BF = $\log(2(M_0-M_1))$, where M_0 is the best model (model 1), which is compared against each of the following ones. *indicates positive support [ln(BF) is n the range of 1.1-3]; ** indicates strong support [ln(BF) is n the range of 3-5]; *** indicates overwhelming support [ln(BF) > 5].
[c]The ln-Bayes factors are computed as described above, but involve M_0 as the model preceding model M_1 from the ranked model list.
NA = not applicable.

1080
1081
1082
Table 3. McDonald-Kreitman (MK) neutrality tests, genetic differentiation and gene flow
estimates between environmental isolates of the Smc1, Smc2 and *S. maltophilia*
(Smal) lineages of the *S. maltophilia* complex recovered from Mexican rivers based
on the concatenated dataset (3591 sites). *D*xy is the interpopulation genetic distance.
K^*_{ST} is Hudson's index of population genetic differentiation. F_{ST} is the fixation
index. Nm represents the number of migrants per generation. *NI* is the neutrality
index and *G* is the likelihood ratio or *G*-test of independence.
1090

Lineages	No. Fixed diffs.	*Dxy*	[a]K^*_{ST}	F_{ST}	*Nm*	*NI*	*G*	[b]*p-value*
Smc1-Smc2	45	0.04616	0.13791***	0.57819	0.36	0.328	6.709	0.00959**
Smc1-Smal	9	0.04542	0.06654***	0.43022	0.66	0.247	3.133	0.07672
Smc2-Smal	28	0.04924	0.09508***	0.59384	0.34	0.312	6.020	0.02515*

1091 [a] The significance of the estimated statistic was computed using the permutation test with
1092 10000 iterations.
1093 [b] *p*-values for the *G*-test.
1094
1095 **Table 4.** Descriptive statistics of DNA polymorphisms, neutrality and population growth
1096 tests for environmental isolates of the Smc1, Smc2, *S. maltophilia* (Smal) and *S.*
1097 *terrae* (Sterr) lineages recovered from Mexican rivers based on the concatenated
1098 dataset (3591 sites). The *p*-values for the R_2 statistic were estimated using 10000
1099 coalescent simulations assuming an intermediate level of recombination.
1100

Species	No. Seqs	*S*	*Eta*	*π*	*h/Hd*	*Theta / site*	*Tajima's D*	R_2	*p-value*
Smc1	11	258	280	0.02502	9/0.945	0.02662	-0.29156 (NS)	0.1390	0.45
Smc2	15	173	185	0.01393	14/0.990	0.01584	-0.45427 (NS)	0.1199	0.19
Smal	51	500	560	0.02652	33/0.929	0.03470	-0.85761 (NS)	0.0903	0.17
Sterr	10	318	328	0.02388	7/0.067	0.04455	-1.30647 (NS)	0.1482	0.49

1101
1102
1103
1104
1105 **Table 5.** Recombination estimates for environmental isolates of the Smc1, Smc2, *S.*
1106 *maltophilia* (Smal) and *S. terrae* (Sterr) lineages recovered from Mexican rivers
1107 based on the concatenated dataset (3591 sites). The figures indicate the posterior
1108 mean and their variances are shown in parentheses.
1109

Lineages	R/theta	1/delta	nu
Smc1	1.10774 (0.07076)	0.0006789 (3.213e-08)	0.0152529 (7.570e-07)
Smc2	1.89989 (0.12028)	0.0009080 (3.121e-08)	0.00864951 (3.081e-7)
Smal	2.95326 (0.10044)	0.0010379 (1.302e-08)	0.0129774 (1.648e-7)
Sterr	0.87233 (0.05113)	0.0021663 (3.443e-07)	0.0650391 (1.2279e-5)

1110
1111
1112

Figure legends

Figure 1. Maximum likelihood multilocus phylogeny of the genus *Stenotrophomonas* and species delimitation hypotheses. The tree shown corresponds to the best one found out of 1001 searches under the GTR+G model and BEST moves, using the concatenated alignment (7-loci) for 194 non-redundant STs, containing all validly described species of the genus except for *S. tumulicola*, additional key reference strains and 63 haplotypes from the 108 Mexican environmental isolates analyzed in this study. A) The S. maltophilia complex (Smc) clade is collapsed in this tree. B) The Smc clade is displayed, collapsing the clades visible in B. Clades are labeled using the A-E codes of Kaiser et al. (2009) and #1-#10 of Hauben et al. (1999) for easy cross-reference. The shaded areas indicate species assignation hypotheses specifically evaluated in this study by the multispecies coalescent using Bayes factors and by population genetics analyses (Tables 2 and 3). The combined evidence of both approaches reveals that the Smc complex should be split into S. maltophilia and four new species lineages: Smc1, Smc2, Smc3 and Smc4 (Fig. 1B). We also show (Fig 1A) that genotype #10 represents a non-described species that should not be merged with *S. rhizophila* clade #8 (Table 2). Type and other key reference strains are highlighted in bold-face. The bar indicates the number of expected substitutions per site under the GTR+G model.

Figure 2. Optimally aligned STRUCTURE barplots for 20 replicate runs executed for $K = 3$ to $K = 6$ generated with CLUMPAK, showing the population genetic structure of the Mexican isolates from the Smc classified as *S. maltophilia* (Smalt) Smc1 (Sm1) and Smc2 (Sm2). The Sm4 cluster corresponds to Smc4 sequences from pubmlst.org, included in the analysis as an outgroup to the Mexican Smc strains (see Fig. 1B).

Figure 3. Multivariate correspondence analysis (MCA) factor –individuals biplot map, summarizing the associations between antibiotic resistance profiles, β-lactamase production phenotypes and species assignations. The state of 17 active variables (IMP=imipenem, MER=meropenem, CAZ=ceftazidime, CAZ.CLA=ceftazidime/clavulanate, FEP=cefepime, ATM=aztreonam, b.lactamase, species, Tm.Cb=trimethoprim+carbenicillin, CL=chloramphenicol, SM=streptomycin, GM=gentamicin, KA=kanamycin, P.T=piperacillin/tazobactam, NAL=nalidixic acid, CIP4=ciprofloxacin, TET=tetracycline) and 4 supplementary categorical variables [Habitat, Env_Qual (environmental quality), Isol_Med (isolation medium), MDR] are shown, sorted along the first two dimensions that together explain 45.3% of the total variance. Some variables like CTX4 (cefotaxime) were excluded, due to lack of variability in the observed states. Species name abbreviations are as follows: malt = *S. maltophilia*; Sm1 = Smc1; Sm2 = Smc2; terr = *S. terrae*.

Figure 4. Panels A, B, E and F display violin and boxplots for the raw count data of number of resistances to individual antimicrobials (NumR) and distinct drug families (NumFam) for Mexican environmental isolates. The big white dot shows the median and the smaller blue one the mean of the distribution of individual observations, represented as small open circles. Yellow dots indicate outlier data points. Panels C, D, G and H show meanplots for NumR and NumFam (see labels on the ordinates) for 20% trimmed means and 95% confidence intervals estimated by non-parametric bootstrap (*nboot* = 1000). Upper row panels (A-D) correspond to analyses for the Mexican isolates classified in the four species/lineages indicated on the x-axes. Lower panels (E-H) correspond to Mexican *S. maltophilia* isolates recovered from high and low pollution sites, based on the criteria indicated on Table 1. The 4 *S. maltophilia* isolates recovered from sites with intermediate contamination level were excluded, as only populations of organisms with > 10 isolates were considered. Species codes are as defined in Fig. 3.

Figure 5. Two-way association plot for the categorical variables species and β-lactamase production phenotype and *Stenotrophomonas* species containing > 10 Mexican environmental isolates. The bars on the plot represent the Pearson residuals, the color code and the height of the bars denote the significance level and magnitude of the residuals, and their widths are proportional to the sample size. The β-lactamase codes are as follows: 0 = no β-lactamase activity detected; CI = clavulanate-inducible class A or class C (AmpC) cephalosporinase; CI_M = clavulanate-inducible cephalosporinase plus metallo β-lactamase (MBL); E = extended-spectrum β-lactamase (ESBL); E_M = ESBL plus MBL; M = MBL. Species codes are as defined in Fig. 3.

Figure 6. Four-way association plot showing the results of multiway-chi-square analysis for the categorical variables species (abbreviations as defined in Fig. 3), habitat (W = water column; S = flooded soil; D = sediment), isolation medium (N = NAA; MK = MacConkey; LE = Leed's Medium) and pollution level (H = high; L = low; M = intermediate, based on counts of thermotolerant coliforms and *E. coli*, as defined in Table 1), using Friendly's residual coloring scheme to highlight the significant associations. Species codes are as defined in Fig. 3. The bars on the plot represent the Pearson residuals, the color code and the height of the bars denote the significance level and magnitude of the residuals, and their widths are proportional to the sample size.