Heart-Specific Activin Signaling Promotes Cardiomyopathy and Cardiac Aging through

Autophagy Inhibition

Kai Chang[1][§], Ping Kang[1][§], Ying Liu[1][§], Kerui Huang[1], Erika Taylor[2], Rolf Bodmer[2], Karen Ocorr[2],

Hua Bai[1]*

1 Department of Genetics, Development, and Cell Biology, Iowa State University, Ames, IA

50011

2 Development, Aging, and Regeneration Program, Sanford-Burnham-Prebys Medical

Discovery Institute, La Jolla, California 92037

*Corresponding Author:

Hua Bai

Phone: 515-294-9395

Email: hbai@iastate.edu

§ These authors contributes equally to this work.

22 **Running title:**

23 **Activin promotes cardiac aging**

Abstract

Age-dependent loss of cardiac tissue homeostasis largely impacts heart performance and contributes significantly to cardiovascular diseases later in life. Cellular quality control machinery, such as autophagy/lysosome system, plays a crucial role in maintaining cardiac health and preventing age-induced cardiomyopathy and heart failure. However, how aging alters autophagy/lysosome system to impact cardiac function remain largely unknown. Here using *Drosophila* heart as a model system, we show that activin signaling, a member of TGF-beta superfamily, negatively regulates cardiac autophagy and promotes age-dependent decline of cardiac functions. We found that cardiac-specific knockdown of *Daw*, an activin-like protein in *Drosophila*, increased cardiac autophagosome number and prevented age-induced cardiac arrhythmias and diastolic dysfunction. Conversely, cardiac-specific expressed activin type I receptor Babo results in pre-matured cardiac dysfunction at young ages. Although Daw positively modulates mTOR signaling (Mechanistic target of rapamycin, the major negative regulator for autophagy), activation of mTOR through the knockdown of *Tsc1* (Tuberous sclerosis protein 1) did not rescue the autophagy and cardiac aging phenotypes in Daw RNAi flies. On the other hand, inhibition of autophagy via chloroquine feeding or Atg1 knockdown attenuated the beneficial effects of Daw RNAi on age-related cardiac arrhythmias. Finally, reduction in cardiac activin signaling prolonged lifespan and improved climbing ability during aging. Thus, our findings highlight the emerging role of activin signaling in mTOR-independent regulation of autophagy and cardiac aging.

Key words: Activin ligand, dawdle, myoglianin, Smad2, Atg8a, mTORC1, TSC1, Autophagy flux

Introduction

Aging is associated with an exponential increase of the incidence of cardiovascular diseases (CVD) [1,2]. Age-related changes in cardiovascular structure and output have been linked to increased risk of coronary heart disease, sudden cardiac death and stroke in aging population [3]. During normal aging, the left ventricular wall of human hearts becomes thickened and the diastolic filling rate of left ventricle gradually decreases. On the other hand, the left ventricular systolic performance at rest remains less or shows no change with age [3]. Several mechanisms underlying age-associated changes in cardiovascular structure and function are proposed, for example changes in growth factor signaling, decreased cellular quality control, altered calcium handling, elevated extracellular matrix deposition or fibrosis, increased mitochondria damage, and the production of reactive oxygen species (ROS) [2,4]. Resolving the contributing mechanisms of age-dependent decline of cardiovascular function is a critical for the development of biomedical interventions for the treatment of cardiovascular diseases.

Cellular quality control systems, such as macroautophagy (hereafter autophagy), are essential to maintain tissue homeostasis during aging [5,6]. Autophagy is a highly conserved process that maintains tissue and cellular homeostasis by degradation and recycling of damaged organelles, protein aggregates and other cytoplasmic substances. The autophagic process initiated with the isolated membrane, or phagophore that elongates to form a double-membrane structure, the autophagosome. Then the autophagosome fuses with a lysosome to form an autolysosome to degrade the enclosed materials along with autophagosomal membrane [7]. Disruption of autophagy pathways by knocking out Atg5 in the mouse heart accelerates cardiac aging, including an increase in left ventricular hypertrophy and decrease in factional shortening [8]. Induction of autophagy by rapamycin inhibition of mTOR (Mechanistic target of rapamycin) protected mouse

cardiomyocytes from antimycin A-induced oxidative stress [9]. Although many longevity

interventions activating autophagy can greatly preserve cardiac function during aging [2], no

evidence has indicated that autophagy activation alone can delay the aging process in animal

hearts. Furthermore, how aging negatively impacts and modulates autophagy is largely unknown

[5]. In general, autophagy activity declines with age [8,10,11]. However, some studies suggest

that autophagy activity remains no change with age [12-15]. The age-dependent alternation of

autophagy and the vital role of autophagy in cardiac protection need to be further investigated.

Autophagy can be activated by various stress, such as dietary restriction, hypoxia, oxidative

stress, and pathogen infection [16]. Autophagy activity is tightly controlled many signaling

pathways, in particular the nutrient signaling like mTOR and AMPK (AMP-activated protein

kinase) [17]. These pathways are often involved in longevity control. We recently identified

activin, a TGF-beta family protein, as a pro-aging hormonal factor regulating autophagy and

proteostasis in fight muscle of *Drosophila* [18]. As in mammals, TGF-beta family in *Drosophila*

has two branches, bone morphogenetic protein (BMP) and activin signaling pathways [19]. In both

pathways, signaling starts with ligand binding to a receptor complex composed of type I and type

II receptor kinases, followed by phosphorylation of receptor-activated Smad (R-Smad). There are

three ligands in *Drosophila* activin signaling pathway, Activin-beta (Act-beta), Dawdle (Daw),

and Myoglianin (Myo). We showed that reduced expression of Daw, but not Act-beta and Myo,

prolongs lifespan and increases autophagosome formation [18].

It is known that TGF-beta factors promotes cardiac fibrosis during aging [20]. However, it

is unclear whether TGF-beta family protein, in particular activin, regulates cardiac homeostasis

and function through autophagy. Given that the cardiac development and aging are remarkably

conserved between *Drosophila* and mammal, as well as the advanced genetic tools and cardiac

movement detection methodologies in *Drosophila* [21], we have applied fly heart model to

investigate the role of activin signaling in the regulation of cardiac autophagy and cardiac aging.

Here we show that activin signaling acts on hearts to regulate autophagy and age-induced

cardiomyopathy in *Drosophila*. The regulation of cardiac aging by activin signaling is ligand-

dependent. RNAi against activin-like protein Daw preserved cardiac function with age, while

reduction in growth differentiation factor 11 (GDF11)/myostatin-like protein Myo promotes

cardiac aging. Reduction of activin signaling in the heart prolongs fly lifespan and improves

climbing ability during aging. In addition, we found that Daw negatively regulates cardiac

autophagy, while autophagy inhibition attenuates the positive effects of Daw RNAi on cardiac

aging. Interestingly, despite the positive relationship between Daw and mTOR, we found that

activation of mTOR through TSC1 RNAi did not block the beneficial effects of Daw knockdown

during cardiac aging. Our findings suggest that *Drosophila* activin signaling may regulate

autophagy and age-induced cardiomyopathy through novel mTOR-independent mechanisms.

Results

Heart-specific knockdown of Daw slows down cardiac aging

Our previous study demonstrated that activin-like protein Daw regulates tissue

homeostasis (especially in flight muscle) and longevity in *Drosophila* [18]. The expression of

Daw and other activin family ligands (Myo and Act-beta) are also detected in fly heart from

previous transcriptome analyses [22-24]. To verify the expression of these activin ligand in the

heart, we performed a tissue-specific ribosome profiling using Ribo-Tag technology. We found

that both Daw and Myo expressed highly in the heart compared to Act-beta (Fig. 1A). To

investigate the role of activin signaling in cardiac aging, we first knocked down the expression of

Daw specifically in fly hearts with a binary GAL4/UAS system by crossing Daw RNAi lines to a

cardiac driver (Hand-gal4). We then analyzed cardiac performances of young and old flies using

a high-speed video imaging system (semi-automatic optical heartbeat analysis, SOHA) [25] (Fig.

1B). Consistent with previous studies, cardiac arrhythmia, heart period, and diastolic interval of

control flies increase with age (Fig. 1 C, 1D, 1E). There is no significant change in systolic

intervals (Fig. 1F). In contract, flies with cardiac-specific knockdown of *Daw* attenuates age-

induced arrhythmia, heart period, and diastolic interval (Fig. 1 C, 1D, 1E). To avoid the

potential off-target effects in knockdown experiments, we tested three different Daw RNAi fly

lines and all of them produce similarly phenotypes.

Because Hand-gal4 drives gene expression in both cardiomyocytes and pericardial cells

(or nephrocytes) [26]. To test whether Daw acts on cardiomyocytes to regulate cardiac

arrhythmia, we crossed Daw RNAi into a cardiomyocyte-specific driver Tin-gal4 [27]. Similar to

the results from Hand-gal4 (Fig. 2A), cardiomyocyte-specific knockdown of *Daw* prevented age-

related increases of cardiac arrhythmia (Fig. 2B). In addition, *Daw* knockdown preserved cardiac

myofibrillar structure with age (Supplemental Figure S1A). We also tested whether Daw acts on

pericardial cells to regulate cardiac aging using a pericardial cell driver Dot-gal4 [28].

Interestingly, knockdown of *Daw* in pericardial cells prevented age-induced arrhythmia (Fig.

2C), but not diastolic interval (Fig. 2D). Thus, Daw appears to promote cardiac aging, in

particular age-induced arrhythmia and diastolic dysfunction, through cardiomyocyte-specific

regulation.

Cardiomyocyte-specific knockdown of activin receptor *Babo*, but not ligand *Myo*, delays

cardiac aging

To investigate whether Daw acts through its receptor Babo to modulate cardiac aging, we

first asked if Babo plays a similar role in preventing age-related increases of cardiomyopathy.

Consistently, we found that cardiomyocyte-specific knockdown of *Babo* blocked the age-

induced cardiac arrhythmia, heart period, and diastolic interval (Fig. 3A, 2B, 2C). Two Babo

RNAi lines were used and both gave similar results. In a reciprocal experiment, we used a

GeneSwitch heart driver to express a constitutive active form of *Babo* (Babo-Act) in adult hearts.

We found that over-expression of *Babo* significantly induced cardiac arrhythmia and diastolic

interval at young age (Fig. 3E, 3F; Supplemental Figure S1B). Interestingly, over-expression of

Babo also increased the diastolic diameter of the heart tube, suggesting a role of Babo in dilated

cardiomyopathy of the fly heart (Supplemental Figure S1C).

In *Drosophila*, there are three Babo isoforms and isoforms of Babo exhibit tissue-specific

expression pattern and response differently to different activin ligands [29]. We found that all

three Babo isoforms showed enriched expression in fly heart (about three fold) (Fig. 4A). Since

both Daw and Myo are highly expressed in the heart, we examine the differential regulation of

cardiac aging by the two activin ligands. Notably, knockdown of *Myo* did not block the age-

induced arrhythmia, rather it further increased the arrhythmia at old age (Fig. 4B). These results

suggest that Daw and Myo differentially regulate cardiac aging, and Daw may directly regulate

age-related cardiomyopathy through its receptor Babo-c.

Daw regulates autophagosome formation and promotes cardiac mTOR activities

Autophagy is one of the key mechanisms in maintaining tissue hemostasis and its activity

is thought to decline with aging [6,8]. Our previous studies showed that activin signaling

regulates autophagosome formation and Atg8a transcription [18]. Here, we examined the

potential role of Daw in the regulation of cardiac autophagy. Using immunostaining and an

antibody against *Drosophila* Atg8a protein (fly homolog of mammalian LC3/GABARAP family

proteins), we observed an increased Atg8a-positive autophagosome in the heart of Daw RNAi

flies (Fig. 5A). Interestingly, flies with heart-specific Daw knockdown maintained high levels of

cardiac autophagosome number at old age, whereas in control flies autophagosome number

decreased with age (Fig. 5A). Because autophagy is a dynamic process, the increased

autophagosome number could result from increased autophagy activities, or blockage of

lysosomal degradation [30]. To examine the role of Daw in controlling autophagy flux (the

turnover of autophagosome), we exposed semi-intact fly hearts to a V-ATPase inhibitor

bafilomycin A1 (Baf A1), which blocks both lysosomal acidification and autophagosome-

lysosome fusion [31]. Notably, both young and aged hearts in control flies maintained active

autophagy flux (or autophagosome turnover activity), as indicated by an increase of

autophagosome number after one hour of Baf A1 treatment (Fig. 5B). Daw RNAi flies showed

less changes in autophagosome number after Baf A1 treatment at both young and old ages,

although the total autophagosome numbers per heart in Daw RNAi flies are higher than control

flies (Fig. 5B). Thus, both control and Daw RNAi flies maintain active cardiac autophagy with

age and Daw knockdown resulted an induction of autophagosome formation.

It is known that nutrient-sensing pathways, such as mTOR, negatively regulate the

autophagy initiation and autophagosome formation [32]. We then tested whether Daw regulates

autophagosome formation through mTOR signaling. Cardiomyocyte-specific Daw knockdown

decreased the phosphorylation of 4E-BP, a major target of mTOR signaling (Fig. 5C). On the

other hand, incubation of conditioned media from Daw-expressing *Drosophila* Kc167 cells

induced the phosphorylation of 4E-BP, while conditioned media from Myo-expressing cells did

not increase p4E-BP levels (Fig. 5D). Taken together, these results suggest that Daw positively

regulates mTOR activity.

Activation of mTOR does not attenuate the autophagosome induction and delayed cardiac

aging in Daw RNAi flies

Given that Daw promotes mTOR signaling, a negative regulator of autophagy, we predict

that Daw negatively regulate autophagosome formation through mTOR. To test this possibility,

we first generated a double knockdown flies by combining *UAS-daw RNAi* and *UAS-Tsc1 RNAi*.

Tsc1 (Tuberous sclerosis protein 1) is a negative regulator upstream of mTOR (in particular

mTOR Complex 1, mTORC1). Cardiomyocyte-specific knockdown of *Tsc1* significantly

induced p4E-BP levels (Supplemental Figure S2A, S2B). Surprisingly, the number of Atg8a-

positive autophagosome remains no changed between Daw RNAi and Daw; Tsc1 double RNAi

flies (Fig. 6A, 6B). In addition, activation of mTOR via Tsc1 RNAi (two independent RNAi

lines) did not change the cardiac arrhythmia (Fig. 6C) and showed reduced diastolic interval at

young age (Fig. 6D), which is very different from cardiac-specific over-expression of activin

receptor *Babo* (Fig. 3E, 3F). Furthermore, Tsc1 RNAi did not attenuate the effects of Daw

knockdown on age-induced cardiac arrhythmia (Fig. 6E). Daw; Tsc1 double RNAi flies did not

show age-dependent increase of arrhythmia, similar to Daw RNAi alone. These results suggest

that Daw negatively regulates autophagosome formation and promotes cardiac aging

independent of mTOR activation.

Intriguingly, Tsc1 knockdown flies (with high mTOR activities) did not exhibit age-

induced cardiac arrhythmia (Fig. 7A) and diastolic dysfunction (Fig. 7B). This observation is

somewhat inconsistent with the notion of mTOR being a pro-aging factor. To further confirm the

unexpected role of mTOR in the regulation of cardiac aging, we tested other key factors involved

in mTOR signaling. Reptor (Repressed by TOR) is a recently identified bZIP transcription factor

downstream of mTORC1 signaling in *Drosophila* [33]. The transcriptional activities of Reptor is

negatively regulated by mTORC1 and rapamycin. Similar to Tsc1 RNAi, cardiomyocyte-specific

knockdown of *Reptor* blocked age-induced arrhythmia (Fig. 7C) and diastolic interval (Fig. 7D).

Furthermore, over-expression of Rheb (Ras Homolog Enriched in Brain), a positive regulator of

mTORC1, resulted in decreased diastolic interval, while no changes in cardiac arrhythmia (Fig.

7E, 7F). The positive role of mTORC1 signaling in preventing age-induced arrhythmia is

unexpected, and somewhat inconsistent with the pro-aging role of mTORC1 as suggested by

previous studies [34-36]. However, we should point out that mTORC1 might regulate tissue

homeostasis and cardiac arrhythmia through distinct mechanisms. Interestingly, we observed a

significant increase of cardiac arrhythmia in cardiomyocyte-specific knockdown of *PTEN*

(Phosphatase and tensin homologue deleted on chromosome ten), a major negative regulator of

PI3K/AKT signaling, suggesting activation of PI3K signaling is detrimental to cardiac function

(Fig. 7G, 7H). Taken together, we found that activation of both activin and PI3K signaling

pathways specifically in cardiomyocytes negatively impacts on heart function, while increased

mTOR activity prevents age-related increase of cardiac arrhythmia in *Drosophila*.

Autophagy inhibition accelerates cardiac aging and attenuates the effects of Daw on age-

related decline of cardiac function

Since we observed an induction of autophagosome formation in Daw knockdown flies,

we next tested whether autophagy activity is required for Daw-regulated cardiac aging. Cardiac-

specific (Hand-gal4 and Tin-gal4) knockdown of *Atg1*, a serine/threonine-protein kinase

essential for autophagy initiation, increased cardiac arrhythmia and diastolic interval at young

age (Fig. 8A, 8B), similar to what we observed in Babo over-expression flies.

To test whether autophagy plays any role in Daw-regulated cardiac aging, we first fed

control and Daw RNAi flies with lysosomal inhibitor chloroquine (CQ) for 24 hours. As

predicted, CQ treatment increased cardiac arrhythmia of Daw RNAi flies at advanced age (Fig.

8C). To further test the role of cardiac-specific autophagy in Daw-mediated cardiac aging, we

generated a double RNAi line targeting both Daw and Atg1 simultaneously. Consistent with CQ

results, cardiac-specific knockdown of both Daw and Atg1 showed age-dependent increase of

cardiac arrhythmia that is similar to control flies (Fig. 8D). Thus, these results suggest that

autophagy is required for Daw-regulated cardiac aging.

Cardiac-specific activin signaling regulates systemic aging

The important roles of several tissues (such as adipose, muscle, and hypothalamus) in

systemic aging control have been previously reported [10,37,38]. However, whether heart can

serve as an aging control center remains unclear. Here we test the effects of cardiac-specific

knockdown and over-expression of activin signaling in longevity control. Knockdown of *Daw* in

the heart using two cardiac tissue drivers (Hand-gal4 and Tin-gal4) significantly prolongs

lifespan (Fig. 9A, 9B, Supplemental Figure S3). On the other hand, knockdown of *Babo* slightly

extended median lifespan (Fig. 9C, Supplemental Figure S3), while over-expression of *Babo*

shortened median lifespan and increased mortality early in life (Fig. 9D, Supplemental Figure

S3).

To test if long-lived Daw RNAi flies show slow aging in non-heart tissues, we performed

climbing assays as an indirect functional assessment for both flight muscle and neuromuscular

junctions. Consistent with prolonged lifespan, cardiac-specific knockdown of *Daw* prevented

age-dependent decline of climbing ability (Fig. 9E) Taken together, our data suggest that

reduction in cardiac activin signaling improves cardiac function, prolongs lifespan and

healthspan in *Drosophila*.

Discussion

TGF-beta signaling plans vital roles in many human diseases, including cancer, kidney

and cardiovascular diseases [39]. Originally identified as a reproductive hormone, TGF-beta

subfamily member activin has become an emerging target of therapy for the treatment of many

human diseases [40]. In the present study, we investigated the role of activin in cardiac aging

using a *Drosophila* heart model. Similar to its role in flight muscle aging [18], reduction of

activin signaling prevents age-related decline of cardiac function (e.g. diastole interval and

arrhythmia). Activin negatively regulates cardiac autophagy and blockade of autophagy process

attenuates the effects of activin on cardiac aging, suggesting the autophagy is essential for

activin-regulated cardiac aging. Interestingly, despite its positive regulation of mTOR, fly

activin-like protein Daw controls autophagy and cardiac arrhythmia independent of mTOR

signaling. Although we did not directly examine the role of mTORC1 key components (like

Raptor) in cardiac aging and the genetic interaction between Raptor and Daw, the consistent

results from multiple mTORC1 regulators or effectors (Tsc1, Rheb, and Reptor) suggest that

mTORC1 may not contribute to Daw-mediated autophagy and cardiac aging control.

There is a positive correlation between activin A mRNA expression and heart failure

[41,42]. Activin A promotes cardiac fibrosis and myocardial damage after ischemia reperfusion

[43], while heterozygous mutant of activin receptor-like kinase 4 (ALK4) protects mouse heart

from pressure overload-induced fibrosis and dysfunction [44]. Our findings support the

protective role of reduced activin signaling in cardiomyopathy, in particular age-induced cardiac

dysfunction. Interestingly, we also found a ligand-dependent regulation of cardiac aging by two

different activin-like ligands. We show that knockdown of activin-like protein Daw delayed

cardiac aging, while reduction in GDF11/ myostatin-like protein Myo promotes cardiac aging.

Although GDF11 can bind to activin type II receptor with similar affinities to that of activin A,

two different ligands may activate very different downstream cascades and cellular processes.

Similarly, TGF-beta family protein GDF11 was identified as an anti-aging circulating factor that

regulates age-related cardiac hypertrophy [45], although the anti-aging effects of GDF11 is still

under debating [46,47].

It has been shown that activin signaling play important roles in tissue homeostasis

[40,48]. Our findings suggest that fly activin-like protein Daw regulates cardiac aging through

autophagy inhibition. It remains to be determine how Daw regulates autophagy. We found that

Daw knockdown induced the number of autophagosome, whereas the autophagy flux

(autophagosome turnover) remains no change. These results suggest that Daw regulates early

step of autophagy to promote autophagosome formation, rather than the late step (lysosome

degradation). While basal autophagy activity in fly hearts is relatively low (based on the number

of autophagosome under normal condition), we found that autophagy is still active in old fly

hearts and autophagy flux remain no change during aging. These observations are not quite

surprising, because several previous studies also found that autophagy/lysosome system remains

active in aged animals [12,13,15,49]. In addition, Daw seems to only modulate the

autophagosome formation, but not autophagy flux. Reduced Daw expression maintains high

autophagosome formation activity throughout the life. These findings raise the possibility that

the cardioprotection by Daw inhibition may be achieved by increasing the autophagosome cargo,

without modulation of the degradation/turnover rate.

It is well known that mTOR is a major autophagy regulator and it inhibits autophagosome

formation and autophagy flux by increasing the phosphorylation of the serine/threonine protein

kinase complex Atg1 (ULK1, ULK2 in mammals), the initiator of the autophagy machinery [7].

Surprisingly, mTOR activation did not repress the high autophagosome formation in Daw

knockdown flies, suggesting unknown mTOR-independent mechanisms involved in Daw-

regulated autophagy. mTOR-independent autophagy regulation has been observed in a previous

genome-wide screen [50]. This study identified an important role of type III PI3 kinases in

autophagy control. They also find that growth factor signaling pathways, such as MAPK-ERK

and AKT, negatively regulate autophagy by inhibiting the type III PI3 kinase cascade. Since it is

known that TGF-beta/actin tightly links to AKT and MAPK signaling pathways, it is possible

that Daw regulates autophagy via AKT/MAPK-regulated type III PI3 kinases.

It is known that mTOR promotes age-dependent increase of cardiac dysfunction in

Drosophila [35,36]. Cardiac-specific over-expression of mTOR increases stress-induced heart

failure, whereas over-expression of Tsc1 and Tsc2 prevents age-related increase of stress-

induced heart failure [35]. However, our studies suggest that mTOR activation prevents age-

induced cardiac arrhythmia, rather than accelerates it. The inconsistency may be due to the

distinct roles of mTOR in the regulation of cardiac arrhythmia and electrical pacing-induced

heart failure. Interestingly, the protective effects of mTOR was previously observed in a mouse

model with cardiac-specific over-expression of mTOR (mTOR-Tg) [51]. The mTOR-Tg mice

show attenuated inflammatory response and cardiomyopathy upon left ventricular pressure

overload induced by transverse aortic constriction. Thus, the specific role of mTOR in stress- and

aging-induced cardiac dysfunction remains to be further examined.

During normal aging, the heart undergoes complex phenotypic changes such as

progressive myocardial remodeling, declined myocardial contractile capacity, increased left

ventricular wall thickness and chamber size, prolonged diastole as well as increased arrhythmia

[3]. All of these biological changes can gradually alter the cardiac functions and confer

vulnerability of the heart to various cardiovascular stresses, thus increase the chance of

developing cardiovascular disease dramatically. The present study uncovered an important role

of TGF-beta/activin signaling pathway in the regulation of autophagy and cardiac aging in

Drosophila. It is likely that autophagy might not be the only process controlled by activin

signaling during cardiac aging. Therefore, future research is needed to explore the role of activin

in other key homeostatic regulatory mechanisms and stress response pathways. The advanced

understanding of TGF-beta/activin signaling pathway and its role in tissue homeostasis will help

the development of therapeutic interventions targeting activin for the treatment of age-related

cardiovascular diseases.

Materials and methods

Fly husbandry and stocks

Flies were maintained at 25 ℃, 60% relative humidity and 12-hour light/dark. Adults

were reared on agar-based diet with 0.8% cornmeal, 10% sugar, and 2.5% yeast (unless

otherwise noted). Fly stocks used in the present study are: UAS-Daw RNAi (BDSC #34974,

BDSC #50911, and VDRC #105309), UAS-Babo RNAi (BDSC #25933, BDSC #40886), Babo-

Act (or Babo-1A3) [52], UAS-Myo RNAi (BDSC #31114), UAS-Tsc RNAi (BDSC #52931,

#54034), UAS-Reptor RNAi (BDSC #25983), UAS-PTEN RNAi (BDSC #33643), UAS-Rheb

(BDSC #9688), UAS-Atg1 RNAi (BDSC #26731), Hand-gal4 (or Hand4.2-gal4) [26], and Tin-

gal4 (or TinCΔ4-gal4) [27], Dot-gal4 [28], Hand-GS-gal4 [23], UAS-RpL13A-FLAG (gift from

Dr. Pankaj Kapahi).

Daw RNAi (BDSC #34974), Babo RNAi (BDSC #25933), and Myo RNAi (BDSC

#31114) were backcrossed into yw^R background for 5~7 generations, and yw^R flies were used as

control/wild-type flies in the experiments. For other UAS-RNAi lines, $y^1 v^1$; P[CaryP]attP2

(BDSC #36303) and y^1 v^1; P[CaryP]attP40 (BDSC # 36304) were used as control/wild-type.

Female flies were used in all experiments.

Ribo-tag and quantitative RT-PCR

For Ribo-tag analysis, about 200 female flies (Hand-gal4>UAS-RpL13A-FLAG) were

homogenized in lysis buffer containing 1% NP-40 (Thermo Fisher Scientific, Waltham, MA,

USA). FLAG-tagged polyribosomes were immunoprecipitated by incubating tissue lysate (input

samples) with anti-FLAG antibodies and SureBeads Protein G Magnetic Beads (Bio-Rad,

Hercules, CA, USA). Actively translated mRNAs were extracted from bead-antibody-ribosome

complex using RNeasy Micro kit (Qiagen, Hilden, Germany). Isolated mRNA was reverse

transcribed to cDNA using iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA).

For Babo isoform expression analysis, Total RNA from 12~15 dissected adult hearts

(yw^R) was extracted using Trizol reagent (Thermo Fisher Scientific, Waltham, MA, USA).

DNase-treated total RNA was quantified and about 50~100 ng of total RNA was reverse

transcribed to cDNA using iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA). QRT-

PCR was performed with a Quantstudio 3 Real-Time PCR System (Thermo Fisher Scientific,

Waltham, MA, USA). mRNA abundance of each gene was normalized to the expression of

ribosomal protein L32 (*RpL32* or *rp49*) by the method of comparative C_T . Primer sequences are

listed in Supplementary Table S1.

Fly heartbeat analysis

To measure cardiac function parameters, semi-intact Drosophila adult fly hearts were

prepared according to previously described protocols [25]. High-speed 3000 frames movies were

taken at a rate of 100 frames per second using a Hamamatsu ORCA-Flash4.0 digital CMOS

camera (Hamamatsu Photonics, Hamamatsu, Japan) on an Olympus BX51WI microscope with a

10X water immersion lens (Olympus, Waltham, MA, USA). The live images that contain heart

tube within abdominal A3 segment were processed using HCI imaging software (Hamamatsu

Photonics, Hamamatsu, Japan). M-modes and cardiac parameters were generated using SOHA, a

MATLAB-based image application as described previously [25,53]. The M-mode provides a

snapshot of movement of heart wall over time. Four cardiac parameters were analyzed, including

heart period, diastolic interval, systolic interval, and arrhythmia index. Diastolic interval (DI) is

the duration for each heart relaxation phase (diastole). Heart period (HP) is the pause time

between the two consecutive diastole. Systolic interval (SI) is calculated as the HP minus the DI.

Arrhythmia index (AI) is calculated as the standard deviation of all HP in each fly normalized to

the median HP.

Antibodies and immunostaining

Anti-Atg8a polyclonal antibodies were generated in rat against a recombinant Atg8a

protein (1-60 aa) (Iowa State University Hybridoma Facility) and affinity purified using CNBr-

Activated Sepharose 4B (GE Healthcare Life Sciences, Marlborough, MA, USA). Other

antibodies for immunostaining included: phospho-4E-BP1 antibody (1:300) (#2855, Cell

Signaling Technology, Danvers, MA, USA), Alexa Fluor 488-conjugated Phalloidin for F-actin

staining (Thermo Fisher Scientific, Waltham, MA, USA), all fluorescence-conjugated secondary

antibodies from Jackson ImmunoResearch, West Grove, PA, USA). For immunostaining, adult

female flies were collected and dissected in PBS. Hearts were fixed in 4% paraformaldehyde for

15 min at RT. After washing in PBS with 0.1% Triton X-100 (PBST), the fixed hearted were

blocked in 5% normal goat serum diluted in PBST for 1 hour at RT. Hearts were then washed

with PBST and incubated overnight at 4 ℃ with primary antibodies diluted in 5% NGS. After

washing with PBST, the samples were incubated for 2 hours at RT with appropriate

fluorescence-conjugated secondary antibodies. Hearts were mounted in ProLong Gold anti-fade

reagent (Thermo Fisher Scientific, Waltham, MA, USA) before imaged using an

epifluorescence-equipped BX51WI microscope (Olympus, Waltham, MA, USA). For imaging

analysis, fluorescence images were first processed using the deconvolution module in Olympus

cellSens demensions package, and then the number/area of positive immunostaining was

measured with the "Measure and Count" module provided by cellSens software (Olympus,

Waltham, MA, USA).

Western blotting

Antibodies for western blot included: beta-Actin antibody (1:2000) (#4967, Cell

Signaling Technology), phospho-4E-BP1 antibody (1:1000) (#2855, Cell Signaling Technology),

and HRP-conjugated secondary antibodies (Jackson ImmunoResearch, West Grove, PA, USA).

Protein samples were homogenized in lysis buffer with leupeptin, benzamidine, antipain, PMSF

and 2-Mercaptoethonal. Supernatant was denatured at 95 ℃ for 5 min. About 30 μg of denatured

protein was separated on Mini-PROTEAN precast gels (Bio-Rad, Hercules, CA, USA) and then

transfer to 0.2 μm PVDF membrane. Following incubation with primary and secondary

antibodies, the blots were visualized with Pierce ECL Western Blotting Substrate (Thermo

Fisher Scientific, Waltham, MA, USA).

Bafilomycin A1 and Chloroquine Treatment

For Bafilomycin A1 treatment, semi-intact hearts were incubated with 100 nM of

Bafilomycin A1 (Alfa Aesar, Haverhill, MA, USA) in artificial hemolymph (buffer receipt in

[25]) for two hours at room temperature prior to the appropriate immunostaining. DMSO was

used as a control.

For chloroquine treatments, 100 μl of 20 mM chloroquine diphosphate salt (Sigma-

Aldrich, St Louis, MO, USA) was added onto the fly food. Flies were fed with chloroquine for at

least 24 hours prior to the appropriate cardiac analysis.

Cell culture and Daw activity assay

A cell-based Daw activity assay was previously described [54]. Briefly, *Drosophila*

Kc167 S2 cells were transfected with FLAG-tagged Daw for three days (Transfection with

empty vector was served as a control). Conditioned media were harvested and incubated with a

fresh flash of Kc167 cells. Three hours after incubation, cell lysate was analyzed by western

blotting.

Demography and survival analysis

Newly enclosed female flies were allowed to mate for two days, then separated from

males and assigned to replicate one-liter demography cages at a density of 100-125 flies per

cage. Three independent cages were set-up per genotype. Food was changed every two days, at

which time dead flies were removed from the cage and counted. Survival analysis was conducted

with JMP statistical software (SAS Institute, Cary, NC, USA), and data from replicate cages

were combined. Mortality distributions were compared by Log-rank test.

Climbing assay

Climbing ability was measured by tapping flies to the bottom of an empty vial and

counting (with video recording) flies that climbed 8 vertical centimeters within 20 seconds.

Thirty females (10 per vial) were scored once every other week for each genotype.

Statistical analysis

GraphPad Prism (GraphPad Software, La Jolla, CA) was used for statistical analysis. To

compare the mean value of treatment groups versus that of control, either student t-test or one-

way ANOVA was performed using Dunnett's test for multiple comparison. The effects of

mutants during aging was analyzed by two-way ANOVA, including Tukey multiple comparisons

test. In SOHA analysis, the outliers were identified using Robust regression and Outlier removal

(ROUT) method (Q=1%) prior to the data analysis.

Acknowledgements

We thank Bloomington Drosophila Stock Center, Harvard Medical School, and Drosophila

Genomics Resource Center for fly stocks and cDNA clones. We thank Michael O'Connor for the

kind advice, activin reagents, and fly lines, Dr. Pankaj Kapahi for Ribo-tag protocols and fly

lines, ISU Hybridoma Facility for Atg8a antibody production.

Author Contributions Statement

Conceived and designed the experiments: RB KO HB. Performed the experiments: KC PK YL

KH HB. Analyzed the data: KC PK YL KH ET HB. Wrote the paper: KC RB KO HB. All

authors reviewed and approved the manuscript.

Competing Interests

The authors declare that no competing interest exists.

Figure Legends

Figure 1. Heart-specific knockdown of *Daw* slows down cardiac aging. (A). Actively

translated mRNA profiling using Ribo-tag qPCR showed that all three activin ligands have

detectable expression in fly heart, while the mRNA expression of *Daw* and *Myo* are relatively

higher than *Act-beta*. Data are represented as mean $\pm$ SE of three trials. Statistical significance is

assessed by one-way ANOVA followed by Tukey multiple comparisons test (* p<0.05, **

p<0.01, *** p<0.001). (B). Representative M-mode traces (8 seconds) showing age-dependent

movement of heart wall in control (or Hang-gal4>wild-type/WT) and heart-specific Daw

knockdown flies (Hang-gal4>UAS-Daw RNAi). (C-E). Age-induced arrhythmia index, heart

period, and diastolic interval are blocked by heart-specific knockdown of *Daw*. (F). No

significant change in systolic interval between young and old control/WT and Daw RNAi flies.

Flies were cultured at 40% relative humidity. Hand-gal4 driver was used to knock down gene

expression specifically in cardiac tissues (cardiomyocytes and pericardial cells). Results from

three independent UAS-Daw RNAi lines are shown (RNAi-1: Bloomington Drosophila Stock

Center (BDSC) #34974, RNAi-2: Vienna Drosophila Resource Center (VDRC) #105309, RNAi-

3: BDSC #50911). Statistical significance is assessed by two-way ANOVA followed by Tukey

multiple comparisons test (* p<0.05, ** p<0.01, *** p<0.001, ns = not significant). N=7~20. The

interaction between genotype and age is statistically significant for heart period (P = 0.0041) and

diastolic interval (P = 0.0243).

Figure 2. Tissue-specific role of Daw in age-dependent cardiac performance. (A-B).

Knockdown of *Daw* using both Hand-gal4 (cardiomyocytes and pericardial cells) and Tin-gal4

(cardiomyocytes) prevents age-induced cardiac arrhythmia. Student t-test (* p<0.05, ** p<0.01,

*** p<0.001). N=25~31. (C-D). Knockdown of *Daw* using Dot-gal4 (pericardial cells) prevents

age-induced cardiac arrhythmia, but not diastolic interval. Student t-test (* p<0.05, ** p<0.01,

*** p<0.001). N=13~26.

Figure 3. Cardiomyocyte-specific knockdown of activin receptor *Babo* delays cardiac aging.

(A-D). Age-induced arrhythmia index, heart period, and diastolic interval, but not systolic

interval, are delayed by cardiomyocyte-specific knockdown of *Babo* (Tin-gal4). Results from

two independent UAS-Babo RNAi lines are shown (RNAi-1: BDSC #25933, RNAi-2: BDSC

#40886). One-way ANOVA followed by Tukey multiple comparisons test (*** $p < 0.001$, **

$p < 0.01$, * $p < 0.05$, ns = not significant). N=15~30. (E-F). Over-expression of constitutive activate

form of Babo (Babo-Act) induces cardiac arrhythmia and diastolic interval at young age (2-

week-old). Hand-GS-gal4 was used to induce adult-onset Babo activation. RU: RU486,

Mifepristone. Flies were cultured at 40% relative humidity. One-way ANOVA (*** $p < 0.001$, **

$p < 0.01$, * $p < 0.05$). N=6~9.

Figure 4. Babo isoform-specific expression in fly hearts and ligand-dependent regulation of

cardiac aging. (A). Ribo-tag PCR shows that all three Babo isoforms are expressed in dissected

adult hearts (1-week-old). Student t-test (* $p < 0.05$). N=3. (B). Two activin ligands, Daw and

Myo, play opposite roles in controlling cardiac arrhythmia. Knockdown of *Daw* prevents age-

induced cardiac arrhythmia, while knockdown of *Myo* promotes arrhythmia. One-way ANOVA

(*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=25~29.

Figure 5. Daw negatively regulates autophagosome formation and promotes cardiac mTOR

activities. (A). Quantification of the fluorescence signaling of mcherry-Atg8a reporter in young

and old control and cardiomyocyte-specific Daw knockdown flies. One-way ANOVA (***

$p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=4. (B). Quantification of autophagy flux

by comparing the changes in mcherry-Atg8a signaling before and after lysosomal inhibitor Baf

A1 treatment. One-way ANOVA (*** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, ns = not significant). N=4.

(C). Immunostaining of the phosphorylation of 4E-BP (p4E-BP) from the dissected

cardiomyocytes of cardiac-specific knockdown of *Daw* and *Myo* flies. Phalloidin was used to

stain F-actin of the cardiomyocytes. Scale bar: 10 μm. (D). Western blots showing p4E-BP levels

in Kc167 cells after the incubation with conditioned media containing Daw and Myo proteins. β–

Actin was used as a loading control.

Figure 6. Activation of mTOR did not attenuate the effects of Daw knockdown in

autophagosome induction and cardiac aging. (A). Immunostaining of autophagosome in

cardiomyocytes from Daw RNAi and Daw; Tsc1 double knockdown flies using Atg8a antibody.

Scale bar: 20 μm. (B). Quantification of autophagosome staining in Panel A. (C-D). Cardiac

arrhythmia and diastolic interval of cardiomyocyte-specific Tsc1 knockdown flies. Two Tsc1

RNAi used (RNAi-1: BDSC #52931, RNAi-2: BDSC #54034). One-way ANOVA (***

$p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=16~18. (E). Cardiomyocyte-specific

knockdown of *Tsc1* did not rescue Daw-regulated cardiac arrhythmia to the wild-type level. One-

way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=7~30.

Figure 7. Activation of mTOR signaling prevents cardiac aging. (A-B). Cardiomyocyte-

specific knockdown of *Tsc1* prevents age-induced arrhythmia and diastolic interval. One-way

ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=14~18. (C-D).

Cardiomyocyte-specific knockdown of *Reptor* prevents age-induced arrhythmia and diastolic

interval. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=13~28.

(E-F). Cardiomyocyte-specific over-expression of *Rheb* reduced diastolic interval, but not

arrhythmia index. Student t-test (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant).

N=21~31. (G-H). Cardiomyocyte-specific knockdown of *PTEN* reduced diastolic interval, but

not arrhythmia index. Student t-test (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant).

N=21~31.

Figure 8. Autophagy disruption accelerates cardiac aging and attenuates the effects of Daw

on age-related decline of cardiac function. (A-B). Both arrhythmia and diastolic interval are

induced in heart-specific Atg1 knockdown flies. Two driver used (Hand-gal4 and Tin-gal4). (C).

Chloroquine treatments (20 mM, 24 hours) increases cardiac arrhythmia of 6-week-old Daw

RNAi flies to the wild-type level. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns =

not significant). N=14~38. (D). Cardiac-specific knockdown of Atg1 attenuates the effects of

Daw on age-induced arrhythmia. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not

significant). N=16~35.

Figure 9. Cardiac-specific activin signaling regulates systemic aging. (A). Survival curve

showing the lifespan of heart-specific (Hand-gal4) Daw knockdown and control flies. Log-Rank

test, $p<0.01$. (B). Lifespan of cardiomyocyte-specific (Tin-gal4) Daw RNAi flies. Log-Rank test,

$p<0.01$. (C). Lifespan of cardiomyocyte-specific Babo RNAi flies. Log-Rank test, $p<0.05$. (D).

Lifespan of cardiomyocyte-specific Babo over-expression flies. Log-Rank test, $p<0.05$. (E).

Heart-specific knockdown of *Daw* prevents age-related decline of climbing ability. One-way

ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=5.

Supplemental Figure S1. (A). F-actin staining showing cardiomyocyte structure in young and

old control and cardiomyocyte-specific Daw knockdown flies. (B). M-mode of heat-specific

Babo over-expression. RU: RU486. (C). Over-expression of Babo increases diastolic diameter.

N=7~8.

Supplemental Figure S2. (A). Immunostaining of p4e-BP in cardiomyocyte-specific

knockdown of Daw or Tsc1. (B). Quantification of p4E-BP staining of Panel A.

References
546

547 1. Dai DF, Chen T, Johnson SC, Szeto H, Rabinovitch PS (2012) Cardiac aging: from molecular
548   mechanisms to significance in human health and disease. Antioxid Redox Signal 16:
549   1492-1526.
550 2. North BJ, Sinclair DA (2012) The intersection between aging and cardiovascular disease. Circ
551   Res 110: 1097-1108.
552 3. Lakatta EG, Levy D (2003) Arterial and cardiac aging: major shareholders in cardiovascular
553   disease enterprises: Part II: the aging heart in health: links to heart disease. Circulation
554   107: 346-354.
555 4. Dai DF, Santana LF, Vermulst M, Tomazela DM, Emond MJ, et al. (2009) Overexpression of
556   catalase targeted to mitochondria attenuates murine cardiac aging. Circulation 119: 2789-
557   2797.
558 5. Quarles EK, Dai DF, Tocchi A, Basisty N, Gitari L, et al. (2015) Quality control systems in
559   cardiac aging. Ageing Res Rev 23: 101-115.
560 6. Rubinsztein DC, Marino G, Kroemer G (2011) Autophagy and aging. Cell 146: 682-695.
561 7. Shibutani ST, Yoshimori T (2014) A current perspective of autophagosome biogenesis. Cell
562   Res 24: 58-68.
563 8. Taneike M, Yamaguchi O, Nakai A, Hikoso S, Takeda T, et al. (2010) Inhibition of autophagy
564   in the heart induces age-related cardiomyopathy. Autophagy 6: 600-606.
565 9. Dutta D, Xu J, Kim JS, Dunn WA, Jr., Leeuwenburgh C (2013) Upregulated autophagy
566   protects cardiomyocytes from oxidative stress-induced toxicity. Autophagy 9: 328-344.
567 10. Demontis F, Perrimon N (2010) FOXO/4E-BP signaling in Drosophila muscles regulates
568   organism-wide proteostasis during aging. Cell 143: 813-825.
569 11. Chang JT, Kumsta C, Hellman AB, Adams LM, Hansen M (2017) Spatiotemporal regulation
570   of autophagy during Caenorhabditis elegans aging. Elife 6.
571 12. Linton PJ, Gurney M, Sengstock D, Mentzer RM, Jr., Gottlieb RA (2015) This old heart:
572   Cardiac aging and autophagy. J Mol Cell Cardiol 83: 44-54.
573 13. Wohlgemuth SE, Julian D, Akin DE, Fried J, Toscano K, et al. (2007) Autophagy in the heart
574   and liver during normal aging and calorie restriction. Rejuvenation Res 10: 281-292.
575 14. Boyle AJ, Shih H, Hwang J, Ye J, Lee B, et al. (2011) Cardiomyopathy of aging in the
576   mammalian heart is characterized by myocardial hypertrophy, fibrosis and a
577   predisposition towards cardiomyocyte apoptosis and autophagy. Exp Gerontol 46: 549-
578   559.
579 15. Inuzuka Y, Okuda J, Kawashima T, Kato T, Niizuma S, et al. (2009) Suppression of
580   phosphoinositide 3-kinase prevents cardiac aging in mice. Circulation 120: 1695-1703.
581 16. Kroemer G, Marino G, Levine B (2010) Autophagy and the integrated stress response. Mol
582   Cell 40: 280-293.
583 17. He C, Klionsky DJ (2009) Regulation mechanisms and signaling pathways of autophagy.
584   Annu Rev Genet 43: 67-93.
585 18. Bai H, Kang P, Hernandez AM, Tatar M (2013) Activin signaling targeted by
586   insulin/dFOXO regulates aging and muscle proteostasis in Drosophila. PLoS Genet 9:
587   e1003941.
588 19. Upadhyay A, Moss-Taylor L, Kim MJ, Ghosh AC, O'Connor MB (2017) TGF-beta Family
589   Signaling in Drosophila. Cold Spring Harb Perspect Biol.
590 20. Biernacka A, Frangogiannis NG (2011) Aging and Cardiac Fibrosis. Aging Dis 2: 158-173.
591 21. Ocorr K, Vogler G, Bodmer R (2014) Methods to assess Drosophila heart development,
592   function and aging. Methods 68: 265-272.

22. Chintapalli VR, Wang J, Dow JA (2007) Using FlyAtlas to identify better Drosophila
melanogaster models of human disease. Nat Genet 39: 715-720.
23. Monnier V, Iche-Torres M, Rera M, Contremoulins V, Guichard C, et al. (2012) dJun and
Vri/dNFIL3 are major regulators of cardiac aging in Drosophila. PLoS Genet 8:
e1003081.
24. Cannon L, Zambon AC, Cammarato A, Zhang Z, Vogler G, et al. (2017) Expression patterns
of cardiac aging in Drosophila. Aging Cell 16: 82-92.
25. Ocorr K, Fink M, Cammarato A, Bernstein S, Bodmer R (2009) Semi-automated Optical
Heartbeat Analysis of small hearts. J Vis Exp.
26. Han Z, Olson EN (2005) Hand is a direct target of Tinman and GATA factors during
Drosophila cardiogenesis and hematopoiesis. Development 132: 3525-3536.
27. Lo PC, Frasch M (2001) A role for the COUP-TF-related gene seven-up in the diversification
of cardioblast identities in the dorsal vessel of Drosophila. Mech Dev 104: 49-60.
28. Zhang F, Zhao Y, Han Z (2013) An in vivo functional analysis system for renal gene
discovery in Drosophila pericardial nephrocytes. J Am Soc Nephrol 24: 191-197.
29. Jensen PA, Zheng X, Lee T, O'Connor MB (2009) The Drosophila Activin-like ligand
Dawdle signals preferentially through one isoform of the Type-I receptor Baboon. Mech
Dev 126: 950-957.
30. Klionsky DJ, Abdelmohsen K, Abe A, Abedin MJ, Abeliovich H, et al. (2016) Guidelines for
the use and interpretation of assays for monitoring autophagy (3rd edition). Autophagy
12: 1-222.
31. Mauvezin C, Nagy P, Juhasz G, Neufeld TP (2015) Autophagosome-lysosome fusion is
independent of V-ATPase-mediated acidification. Nat Commun 6: 7007.
32. Scott RC, Schuldiner O, Neufeld TP (2004) Role and regulation of starvation-induced
autophagy in the Drosophila fat body. Dev Cell 7: 167-178.
33. Tiebe M, Lutz M, De La Garza A, Buechling T, Boutros M, et al. (2015) REPTOR and
REPTOR-BP Regulate Organismal Metabolism and Transcription Downstream of
TORC1. Dev Cell 33: 272-284.
34. Kaeberlein M, Powers RW, 3rd, Steffen KK, Westman EA, Hu D, et al. (2005) Regulation of
yeast replicative life span by TOR and Sch9 in response to nutrients. Science 310: 1193-
1196.
35. Wessells R, Fitzgerald E, Piazza N, Ocorr K, Morley S, et al. (2009) d4eBP acts downstream
of both dTOR and dFoxo to modulate cardiac functional aging in Drosophila. Aging Cell
8: 542-552.
36. Luong N, Davies CR, Wessells RJ, Graham SM, King MT, et al. (2006) Activated FOXO-
mediated insulin resistance is blocked by reduction of TOR activity. Cell Metab 4: 133-
142.
37. Hwangbo DS, Gershman B, Tu MP, Palmer M, Tatar M (2004) Drosophila dFOXO controls
lifespan and regulates insulin signalling in brain and fat body. Nature 429: 562-566.
38. Zhang G, Li J, Purkayastha S, Tang Y, Zhang H, et al. (2013) Hypothalamic programming of
systemic ageing involving IKK-beta, NF-kappaB and GnRH. Nature 497: 211-216.
39. Akhurst RJ, Hata A (2012) Targeting the TGFbeta signalling pathway in disease. Nat Rev
Drug Discov 11: 790-811.
40. Tsuchida K, Nakatani M, Hitachi K, Uezumi A, Sunada Y, et al. (2009) Activin signaling as
an emerging target for therapeutic interventions. Cell Commun Signal 7: 15.

41. Yndestad A, Ueland T, Oie E, Florholmen G, Halvorsen B, et al. (2004) Elevated levels of
activin A in heart failure: potential role in myocardial remodeling. Circulation 109: 1379-
1385.
42. Wei Q, Wang YN, Liu HY, Yang J, Yang CY, et al. (2013) The expression and role of
activin A and follistatin in heart failure rats after myocardial infarction. Int J Cardiol 168:
2994-2997.
43. Chen Y, Rothnie C, Spring D, Verrier E, Venardos K, et al. (2014) Regulation and actions of
activin A and follistatin in myocardial ischaemia-reperfusion injury. Cytokine 69: 255-
262.
44. Li CY, Chen YH, Wang Q, Hou JW, Wang H, et al. (2016) Partial inhibition of activin
receptor-like kinase 4 attenuates pressure overload-induced cardiac fibrosis and improves
cardiac function. J Hypertens 34: 1766-1777.
45. Loffredo FS, Steinhauser ML, Jay SM, Gannon J, Pancoast JR, et al. (2013) Growth
differentiation factor 11 is a circulating factor that reverses age-related cardiac
hypertrophy. Cell 153: 828-839.
46. Schafer MJ, Atkinson EJ, Vanderboom PM, Kotajarvi B, White TA, et al. (2016)
Quantification of GDF11 and Myostatin in Human Aging and Cardiovascular Disease.
Cell Metab 23: 1207-1215.
47. Smith SC, Zhang X, Zhang X, Gross P, Starosta T, et al. (2015) GDF11 does not rescue
aging-related pathological hypertrophy. Circ Res 117: 926-932.
48. Ghosh AC, O'Connor MB (2014) Systemic Activin signaling independently regulates sugar
homeostasis, cellular metabolism, and pH balance in Drosophila melanogaster. Proc Natl
Acad Sci U S A 111: 5729-5734.
49. Chapin HC, Okada M, Merz AJ, Miller DL (2015) Tissue-specific autophagy responses to
aging and stress in C. elegans. Aging (Albany NY) 7: 419-434.
50. Lipinski MM, Hoffman G, Ng A, Zhou W, Py BF, et al. (2010) A genome-wide siRNA
screen reveals multiple mTORC1 independent signaling pathways regulating autophagy
under normal nutritional conditions. Dev Cell 18: 1041-1052.
51. Song X, Kusakari Y, Xiao CY, Kinsella SD, Rosenberg MA, et al. (2010) mTOR attenuates
the inflammatory response in cardiomyocytes and prevents cardiac dysfunction in
pathological hypertrophy. Am J Physiol Cell Physiol 299: C1256-1266.
52. Brummel T, Abdollah S, Haerry TE, Shimell MJ, Merriam J, et al. (1999) The Drosophila
activin receptor baboon signals through dSmad2 and controls cell proliferation but not
patterning during larval development. Genes Dev 13: 98-111.
53. Fink M, Callol-Massot C, Chu A, Ruiz-Lozano P, Izpisua Belmonte JC, et al. (2009) A new
method for detection and quantification of heartbeat parameters in Drosophila, zebrafish,
and embryonic mouse hearts. Biotechniques 46: 101-113.
54. Serpe M, O'Connor MB (2006) The metalloprotease tolloid-related and its TGF-beta-like
substrate Dawdle regulate Drosophila motoneuron axon guidance. Development 133:
4969-4979.
678
679

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig 7

Fig. 8

Fig. 9

Supplemental Figure S1

A

Tin>WT   Tin>Daw RNAi

F-actin

B

Hand-GS > Babo-Act

C

Hand-GS>Babo-Act

Supplemental Figure S2

Tin-gal4>

A

p4E-BP

WT   Daw RNAi   Tsc1 RNAi

B

% p4E-BP positive staining per heart

1.5

1.0

0.5

0.0

**

*

WT   Daw RNAi   Tsc1 RNAi

Tin-gal4>

Supplementary Table S1.

P19	AAGAAGCGCACCAAGCACTTCATC	RPL32-f	RPL32	FBgn0002626
P20	TCTGTTGTCGATACCCTTGGGCTT	RPL32-r	RPL32	FBgn0002626
P21	TGAGCCACCTCATCCAAATCACCT	daw-f	daw	FBgn0031461
P22	TCGATCACGATGAATGGCCGGTAA	daw-r	daw	FBgn0031461
P23	TGATCATGACATGCGCTGTTGTCG	myo-f	myo	FBgn0026199
P24	ACGTTGGTGCTACAACGAAATGCC	myo-r	myo	FBgn0026199
P95	GGCTTTTGTTTCACGTCCGTGGA	babo-5'	CG8224	FBgn0011300
P96	CTGTTTCAAATATCCTCTTCATAATCTCAT	babo-A-3'	CG8224	FBgn0011300
P97	CTTGGACGGAGATTTGCCGCCC	babo-B-3'	CG8224	FBgn0011300
P98	CTGGTAGGACACCACTGTAATTCTC	babo-C-3'	CG8224	FBgn0011300
P131	CATTTGGGAGCAGTGGTAAGA	Act-beta-f	Act-beta	FBgn0024913
P132	CTTCCACCTCGGCCATTAAA	Act-beta-r	Act-beta	FBgn0024913