Heart-Specific Activin Signaling Promotes Cardiomyopathy and Organismal Aging

through Autophagy Inhibition

Kai Chang[1][§], Ping Kang[1][§], Ying Liu[1][§], Kerui Huang[1], Erika Taylor[2], Rolf Bodmer[2], Karen Ocorr[2],

Hua Bai[1]*

1 Department of Genetics, Development, and Cell Biology, Iowa State University, Ames, IA

50011

2 Development, Aging, and Regeneration Program, Sanford-Burnham-Prebys Medical

Discovery Institute, La Jolla, California 92037

***Corresponding Author:**

Hua Bai

Phone: 515-294-9395

Email: hbai@iastate.edu

§ These authors contributes equally to this work.

22   **Running title:**

23   **Activin promotes cardiac aging**

Abstract

Age-dependent loss of cardiac tissue homeostasis largely impacts heart performance and contributes significantly to cardiovascular diseases later in life. Cellular quality control machinery, such as autophagy/lysosome system, plays a crucial role in maintaining cardiac health and preventing age-induced cardiomyopathy and heart failure. However, how aging alters autophagy/lysosome system to impact cardiac function remain largely unknown. Here using Drosophila model system, we show that cellular autophagic flux and lysosome number decrease in aging heart, which is associated with increased cardiomyopathy and cardiac arrhythmias. Among many known autophagy regulators, activin signaling (a member of TGF-beta superfamily) was identified in our recent study as a negative factor of autophagy and protein homeostasis in flight muscle. In this study, we find that cardiac-specific knockdown of Daw, an activin-like protein in Drosophila, prevents age-dependent increases in cardiac arrhythmias and diastolic dysfunction. Furthermore, cardiac-specific expressed activin type I receptor Babo results in pre-matured cardiac aging phenotypes at young ages. Similar to our previous flight muscle study, Daw silencing strongly promotes early step of autophagy process (i.e. autophagosome formation), and shows less impacts on autophagosome-lysosome fusion. Flies with Daw knockdown also maintain robust autophagic flux in aged fly hearts. Interestingly, reduction in cardiac activin signaling significantly prolongs lifespan and improves the functions of distal tissues (such as age-dependent climbing ability). Thus, our findings highlight the emerging role of activin signaling in autophagic regulation, cardiac aging, as well as systemic control of longevity.

Introduction

Aging is associated with an exponential increase of the incidence of cardiovascular diseases (CVD) (Dai et al., 2012; North and Sinclair, 2012). Resolving the contributing mechanisms of cardiovascular diseases is a pressing goal of basic and translational aging research. Among many plausible mechanisms underlying age-related diseases, the impact of inter-organ communication on tissue aging, including cardiac aging, has recently become an important topic in the field of aging research. Several recently discovered blood-borne factors, such as growth differentiation factor 11 (GDF11), have been linked to systemic aging control and age-associated pathologies (e.g., cardiac hypertrophy) (Castellano et al., 2017; Demontis et al., 2014; Loffredo et al., 2013). However, due to their signal complexity, the precise regulatory role of these circulating factors (especially hormones and cytokines) in tissue aging remains largely unknown. To develop effective strategies targeting humoral factors to treat age-associated diseases (e.g., CVD), it is crucial to decipher the mechanisms underlying systemic aging regulation. These include how systemic factors interact with their receptors, how tissue-specific regulation is achieved, and how they mediate inter-organ communication to maintain organismal homeostasis.

Age-related changes in cardiovascular structure and output have been linked to increased risk of coronary heart disease, sudden cardiac death and stroke in aging population (Lakatta and Levy, 2003). During normal aging, the left ventricular wall of human hearts becomes thickened and the diastolic filling rate of left ventricle gradually decreases with age. On the other hand, the left ventricular systolic performance at rest remains less or shows no change with age (Lakatta and Levy, 2003). Several mechanisms underlying age-associated changes in cardiovascular structure and function are proposed, for example changes in growth factor signaling, decreased cellular quality control, altered calcium handling, elevated extracellular matrix deposition or fibrosis,

increased mitochondria damage, and the production of reactive oxygen species (ROS) (Dai et al.,

2009; North and Sinclair, 2012).

Cellular quality control systems, such as macroautophagy (hereafter autophagy), are

essential to maintain tissue homeostasis during aging (Quarles et al., 2015). Disruption of

autophagy pathways by Atg5 knockout in the mouse heart accelerates cardiac aging, including an

increase in left ventricular hypertrophy and decrease in factional shortening (Taneike et al., 2010).

Although many longevity interventions activating autophagy can greatly preserve cardiac function

during aging (North and Sinclair, 2012), no evidence has indicated that autophagy activation alone

can delay the aging process in animal hearts. Furthermore, how aging negatively impacts and

modulates autophagy is largely unknown (Quarles et al., 2015). Therefore, there is an urgent need

to fully understand the regulation of cellular autophagy during cardiac aging, in order to develop

effective ways to activate autophagy to prevent age-associated tissue damage and cardiac

dysfunction.

We recently identified that activin signaling acts on hearts to regulate autophagy and age-

induced cardiomyopathy in *Drosophila*. Additionally, the regulation of cardiac aging by activin

signaling is ligand-dependent. RNAi against activin-like protein Daw preserved cardiac function

with age, while reduction in GDF11-like protein Myo promotes cardiac aging. We found that Daw

negatively regulates cardiac autophagy, while autophagy inhibition attenuates the positive effects

of Daw RNAi on cardiac aging. Interestingly, despite the positive relationship between Daw and

mTOR (mechanistic target of rapamycin, a major autophagy regulator), we found that activation

of mTOR through TSC1 RNAi did not block the beneficial effects of Daw knockdown during

cardiac aging, suggesting that Daw might regulate cardiac aging through mTOR-independent

pathways. Our findings suggest that Drosophila activin signaling may regulate autophagy and age-

induced cardiomyopathy through novel mTOR-independent mechanisms.

Materials and methods

Fly Stocks, Feeding Protocol, and Chloroquine Treatment

96   UAS-Atg1 RNAi (HB387, BL26731), UAS-Trip attp40 (HB389, BL36304), UAS-Tsc

RNAi (HB359, BL52931), UAS-Tsc1 RNAi (HB361, BL54034), Daw RNAi (HB314,

BL50911), Daw RNAi (HB226, BL34974) were from the Bloomington Stock Center. Daw

RNAi (HB226, BL34974) were backcrossed to ywR in our lab. Heart specific drivers Hand4.2-

Gal4 and TinΔ4-Gal4 were kind gifts from Rolf Bodmer (Sanford Burnham Prebys Medical

Discovery Institute). Unless specialized food is needed, fly strains were maintained on standard

cornmeal molasses agar medium at room temperature (25℃). For aging experiments, fly stains

were transferred to fresh food every 2-3 days. For chloroquine treatments, 100 ul of 20 mM

chloroquine diphosphate salt (Sigma) was added onto the food.

Fly heartbeat analysis

106   A semi-intact Drosophila adult fly heart was prepared according to previously described

protocols (Ocorr, Fink, Cammarato, Bernstein, & Bodmer, 2009; Vogler & Ocorr, 2009) in order

to measure cardiac function parameters. High-speed 3000 frames movies were taken around the

rate of 100 fps using a HAMAMATSU C11440 camera on an Olympus BX51WI microscope

with a 10X water immersion lens. The live images were processed using HCI imaging software

(Hamamatsu Corp). M-modes and quantitative data were generated using SOHA, a MatLab-

based application that was published in K Occor et al., 2007 (Ocorr et al., 2007). The M-mode

provides a snapshot of movement of heart tube wall edges of abdominal A3 segment in Y-axis

over time in the X-axis. Diastolic Interval is the heart relaxation time, and systolic interval is the

heart contraction time. Heart Period is the duration between ends of two consecutive diastolic

intervals. Arrhythmias observed in M-mode can be indicated as an arrhythmia index, which is

obtained by normalizing the standard deviation of all heart periods in each record to the median

heart period for each fly. The above measurements were made in abdominal A3 segment as well

and then analyzed by GraphPad Prism 7 (GraphPad Softwore, Inc). The outliers were identified

using Robust regression and Outlier removal (ROUT) method (Q=1%), and statistical

significances were evaluated by t-test and one-way ANOVA analyses using GraphPad Prism 7.

Immunostaining and imaging

Antibodies for immonustaining included: anti-Atg8a (1:300) (generated in this study) and

anti-rat IgG 594 (Jackson ImmunoResearch). F-actin was visualized by Alexa Fluor 488-

conjugated Phalloidin (Invitrogen). Adult female flies were collected and dissected in PBS.

Hearts were fixed in 4% paraformaldehyde for 15 min at RT. After washing in PBS with 0.1%

Triton X-100 (PBST), the fixed hearted were blocked in 5% normal goat serum diluted in PBST

(5% NGS) for 1 hour at RT. Hearts were then washed with PBST and incubated overnight at 4 ℃

with primary antibodies diluted in 5% NGS. After washing hearts with PBST, the samples were

incubated for 2 hours at RT with the appropriate fluorescence-conjugated secondary antibodies

(Jackson ImmunoResearch) diluted in PBST. Hearts were then washed again with PBST and

mounted in ProLong Gold antifade reagent (Invitrogen). Samples were examined under an

epifluorescence-equipped (Olympus) microscope. The number/area of positive immunostaining

was measured with the "Measure and Count" function provided by Cellsens (Olympus).

Western Blot

Antibodies for western blot included: beta-Actin antibody (1:2000) (Cell Signaling

Technology# 4967S), phospho-4E-BP1 antibody (1:1000) (Cell Signaling Technology# 2855S),

Atg8a antibody (generated from our lab), and HRP conjugated secondary antibodies, anti-Rat-

IgG-HRP (1:5000) (Company), anti-Rabbit-IgG-HRP (1:5000) (Company). KC167 cells were

homogenized in lysis buffer with leupeptin, benzamidine, antipain, PMSF and 2-

Mercaptoethonal. Supernatant was denatured at 95 ℃ for 5 min. About 30ug of denatured protein

was separated on Mini-PROTEAN precast gels (Bio-Rad) and then transfer to PVDF membrane

(Bio-Rad). Following incubation with primary and secondary antibodies, the blots were

visualized with Pierce ECL Western Blotting Substrate (Thermo Fisher Scientific). Band

intensity was quantified with Image Lab software (Bio-Rad).

Statistical analysis

GraphPad Prism 6 (GraphPad Software, La Jolla, CA) was used for statistical analysis.

To compare the mean value of treatment groups versus that of control, either student t-test or

one-way ANOVA was performed using Dunnett's test for multiple comparison. The effects of

mutants during aging was analyzed by two-way ANOVA, including Tukey multiple comparisons

test.

Results

In order to investigate Activin signaling regulation on cardiac aging, heart-specific

drivers were crossed with ywR wild type and daw mutant flies (RNAi). The female flies from the

offspring of above crosses were used to analyze heart performance on their 2-week-old, 4-week-

old, 6-week-old and 8-week-old. On each time point, semi-intact Drosophila adult fly hearts

were prepared according to previously described protocols (Ocorr, Fink, Cammarato, Bernstein,

& Bodmer, 2009; Vogler & Ocorr, 2009) in order to measure cardiac function parameters. The

heart movement was captured in high-speed 3000 frames movies taken at around the rate of 100

fps using a HAMAMATSU C11440 camera on an Olympus BX51WI microscope with a 10X

water immersion lens. The movies were processed using HCI imaging software (Hamamatsu

Corp.) M-modes and quantitative data were generated using SOHA, a MatLab-based application

that was published in K Occor et al., 2007 (Ocorr et al., 2007). The M-mode provides a snapshot

of movement of heart tube wall edges in Y-axis over time in the X-axis. Diastolic Interval is the

heart relaxation time, and systolic interval is the heart contraction time. Heart Period is the

duration between ends of two consecutive diastolic intervals. Arrhythmias observed in M-mode

can be indicated as an arrhythmia index, which is obtained by normalizing the standard deviation

of all heart periods in each record to the median heart period for each fly. In this study, we found

that cardiac-specific Daw knockdown preserves age-dependent increases in cardiac arrhythmia

and diastolic dysfunction (Figure 1).

To investigate Activin signaling on autophagy, Atg8a (homolog of mammals LC3)

antibody generated from our lab was used to detect the autophagosome. In young flies, the

intensity of basal autophagosome levels were very low in wildtype hearts, but much higher in

hearts expressing Daw RNAi. In old flies, even though the autophagosome number in hearts

expressing Daw RNAi was lower than the young hearts, it was still significantly higher than the

autophagosome number in wild type hearts (Figure 2A). In order to verify whether the high level

autophagosome in daw RNAi line is due to elevated upstream autophagosome formation level, or

defect downstream lysosomal activity, the BafA1 was used to inhibit the lysosomal activity. The

accumulation of autophagosome after 1 hour BafA1 incubation was captured to represent the

autophagy flux. Based on my results, the basal autophagy flux was very low and showed no

significant difference between wild type and Daw RNAi hearts of either young or old flies

(Figure 2B). There are two explanations for these results: (1) Activin signaling stimulates cardiac

autophagosome formation but does not affect basal autophagy flux. (2) The basal autophagy flux

was too low to detect any difference between wild type and Daw RNAi flies.

It is well-known that nutrient-sensing pathways, such as mTOR, negatively regulate the

activation of autophagy machinery. We then tested whether Activin/Daw signaling regulates

autophagosome formation through mTOR. We found that reduced expression of Daw decreases

the phosphorylation of 4E-BP, one of the major targets of mTOR signaling, suggesting that Daw

positively regulates mTOR activities (Figure 3A). However, we found that mTOR signaling is

not required for the activation of autophagosome formation by Daw RNAi, indicated by the high

levels of autophagosome in Daw and TSC1 double knockdown flies (Fig. 3B). These results

suggest that Daw regulates autophagosome formation through an unknown mTOR-independent

mechanism. As we observed an unconventional regulation of autophagy by Activin/Daw

signaling, we wonder whether or not Daw might regulate cardiac aging through mTOR. In

contrast to previous reports (Lee et al., 2010), activation of mTOR by Tsc1 RNAi did not affect

cardiac arrhythmicity or accelerated cardiac aging, and mTOR is not required for Daw-mediated

cardiac aging (Fig. 3C). Altogether, Activin/Daw signaling regulates cardiac aging through

autophagy/lysosome system, independent of mTOR.

Atg1 is essential to initiate autophagy machinery in Drosophila, our next step was to test

whether autophagy plays any role in cardiac aging. Autophagy disruption via silencing Atg1 in

heart using two tissue-specific drivers Tin-Delta4-Gal4 and Hand4.2-Gal4 both showed

premature cardiac phenotypes: in 2-week-old female hearts, the Arrhythmia index and diastolic

intervals increased to aged hearts' level (Figure 4). Knocking-down Atg1 in Hand4.2-Gal4 tissue

driver had stronger phenotype than in Tin-Delta4-Gal4 because Hand4.2-Gal4 is more robust

than the other.

To further test whether autophagy plays any role in Daw-regulated cardiac aging, cardiac

tissue-specific drivers Hand4.2-gal4 were crossed with ywR wild type and Daw mutant flies

(RNAi), the adult fly progeny was then fed with lysosomal inhibitor chloroquine (CQ) for 24

hours before analyzing cardiac function by SOHA. CQ treatment increased arrhythmia index of

Daw RNAi flies at advanced age, suggesting disruption of autophagy/lysosome system abolishes

the beneficial effects of Daw knockdown on cardiac arrhythmicity (Figure 5A). Similarly,

autophagy disruption via Atg1 RNAi blocked the beneficial effects of Daw RNAi on age-

dependent cardiac arrhythmicity (Figure 5B).

Discussion

Aging is an extremely complex, multifactorial process accompanied by accumulation of

deleterious changes in cells and tissues that are respond for a wide range of diseases and even

death. During the process, the heart undergoes complex phenotypic changes such as progressive

myocardial remodeling, declined myocardial contractile capacity, increased left ventricular wall

thickness and chamber size, prolonged diastole as well as increased arrhythmia (Lakatta & Levy,

2003; Strait & Lakatta, 2012). All of those biological changes can gradually alter the cardiac

functions and confer vulnerability of the heart to various cardiovascular stresses, thus increase

the chance of developing cardiovascular disease (CVD) dramatically. Even though the death

rates in United States caused by CVD have declined in the United States recent years due to

large amount of research on related field, it is still the leading cause of death, especially in elder

population. There are 30.8% deaths caused by CVD from 2003 to 2013 in United States, 65% of

them occurred after the age of 75 (Mozaffarian et al., 2016). In this regard, it is crucial to

understand the mechanism behind cardiac aging, it can help us not only prevent CVD but also

explore potential treatments for promoting cardiac health.

So far there are large amount of theories for cardiac aging process. It was generally

accepted that age-related CVD results from the accumulation of cholesterol and fatty acids in

tissues, which will further induce the production of inflammatory cytokines and reactive oxygen

species (ROS) (North & Sinclair, 2012). Recent years, a number of longevity genes and their

related pathways have been identified involved in regulating fundamental process of cardiac

aging. For instance, it has been proposed that calories restriction can increase health span and

reduce the incidence of most age-related diseases (i.e. CVDs) by promoting expression of

longevity genes (Ahmet et al., 2011). Even though enriched knowledge of molecular mechanism

of cardiac aging has been revealed, the precise aging process, which cannot explain in a unifying

manner still remains largely unknown. As more novel mechanisms behind cardiac aging being

explored, the unique role of autophagy has been depicted in regulation of biological aging

process.

Autophagy is a highly conserved process that maintains tissue and cellular homeostasis

by degradation and recycling of damaged organelles, protein aggregates and other cytoplasmic

substances. Microautophagy, chaperon-mediated autophagy and macroautophagy are three

identified pathways of autophagy, macroautophagy (simply referred to autophagy hereafter) is

the pathway that will be the focus of our study. The autophagic process initiated with the

isolated membrane, or phagophore that elongates to form a double-membrane structure, the

autophagosome. Then the autophagosome fuses with a lysosome to form an autolysosome to

degrade the enclosed materials along with autophagosomal membrane (Shibutani & Yoshimori,

2014). In Drosophila, the components of autophagy can be divided into several function units,

they are: (1) the serine/threonine protein kinase complex Atg1 (ULK1, ULK2 in mammals),

commonly considered as the initiator of the autophagy machinery; (2) two ubiquitin-like

conjugation systems, Atg12 (covalently conjugated to Atg5), and Atg8 [(LC3 in mammals)

conjugated to the lipid molecule phosphatidylethanolamine (PE)]; and (3) a protein complex

containing the class III phosphatidylinositol 3-kinase Vps34 (PIK3C3 in mammals), that is

required for autophagosome formation (Neufeld, 2012; Wong, Puente, Ganley, & Jiang, 2013).

There are evidence supports that basal autophagy plays essential roles in maintaining normal

heart homeostasis and morphology. Disruption of autophagy by Atg5 knockout in mouse heart

has shown premature phenotype, such as increased left ventricular hypertrophy and decreased

fractional shortening (Taneike et al., 2010). Thus manipulation of autophagy machinery could

potentially help to develop the therapeutic approaches to treat CVDs. However, the most

fundamental questions about autophagy, how cardiac aging regulates autophagy, and what is the

cause of age-associated autophagy alterations are still unsolved. Therefore in my study, I aim to

gain deeper understanding of the fundamental relationship between cardiac aging and autophagy.

Previous literature performed lifespan screening and proposed that the activin signaling

of TGF-b family is involved in regulating muscle aging and autophagy (Bai, Kang, Hernandez,

& Tatar, 2013). Based on my study, the Activin signaling also plays a role in cardiac aging and

cardiac basal autophagosome formation. Besides, the transforming growth factor b (TGF-b)

family signaling is essential in regulating cell growth, differentiation and developmental process

in wide range of biological systems (Massague, 2012). Thus the Activin signaling of TGF-b

family could be a good start point to explore the mechanism of cardiac aging process. Moreover,

the reverse genetics screening of Actin signaling downstream targets can be an alternative

method to identify more candidate genes that are potentially involved in Activin

signaling/cardiac aging mechanism. In Drosophila, fewer signaling components are present

which can simplify mechanistic study of this pathway. As in mammals, TGF-b family signaling

pathway has two branches in Drosophila, they are bone morphogenetic protein (BMP) and

Activin signaling pathway. In both pathways, signaling starts with ligand binding to a receptor

complex composed of type I and type II receptor kinases. Activation of type II receptor kinase by

ligand binding causes phosphorylation of type I receptor kinase, which enables appropriate

phosphorylation of receptor-activated Smad (R-Smad) substrate. R-Smad phosphorylation causes

formation of complex with the Drosophila common-Smad (co-Smad), Medea, which can

translocate to the nucleus and regulate its downstream transcriptional activity. Within the Activin

subfamily, three ligands, activin-b (Actb), Dawdle (Daw), and Myoglianin (Myo) signal through

the type I receptor Baboon (Babo), type II receptor Wit or Punt to activate transcriptional factor

dSmad2, the activated form of dSmad2 then forms a complex with its cofactor Medea and

translocates to nucleus to regulate downstream transcription (Upadhyay, Moss-taylor, Kim,

Ghosh, & Connor, 2017).

Here we uncovered an important role of activin signaling in the regulation of cardiac

aging in Drosophila. Our findings of mTOR-independent autophagy suggests that there are

diverse signaling pathways controlling autophagy processes and each of them might play distinct

roles in tissue homeostasis and aging. This work will contribute to advance our understanding of

the molecular and genetic mechanisms underlying cardiac aging. Activin signaling is one of the

important targets of therapeutic drug development for the treatment of lung cancer and muscle

dystrophy (especially cancer cachexia) (Tsuchida et al., 2009; Zhou et al., 2010). A number of

clinical trials are currently investigating treatments for cancer and muscle wasting using activin

inhibitors (Cohen et al., 2015). Thus, our work is expected to significantly advance the field at

both the basic and applied levels. Due to the signal complexity of activin signaling (for example,

multiple ligands act on a few receptors), the tissue-dependent action of each activin ligand

remains to be carefully examined. Using sophisticated Drosophila genetic tools and tissue-

specific control of gene expression, we anticipate the identification of distinct mechanisms by

which activin signaling regulates tissue homeostasis during aging.

Acknowledgements

We thank Bloomington Drosophila Stock Center and Drosophila Genomics Resource Center for

fly stocks and cDNA clones. We thank Michael O'Connor for the kind advice, activin reagents,

and fly lines.

Author Contributions Statement

H.B., R.B., and K.O. planned research and prepared manuscript. K.C., P.K., Y.L., K.H., E.T.,

H.B. performed research. All authors reviewed the manuscript and approved.

Competing Interests

The authors have declared that no competing interest exists.

References
312
Castellano, J.M., Mosher, K.I., Abbey, R.J., McBride, A.A., James, M.L., Berdnik, D., Shen,
J.C., Zou, B., Xie, X.S., Tingle, M., Hinkson, I.V., Angst, M.S., Wyss-Coray, T., 2017.
Human umbilical cord plasma proteins revitalize hippocampal function in aged mice.
Nature 544, 488-492.
Dai, D.F., Chen, T., Johnson, S.C., Szeto, H., Rabinovitch, P.S., 2012. Cardiac aging: from
molecular mechanisms to significance in human health and disease. Antioxidants &
redox signaling 16, 1492-1526.
Dai, D.F., Santana, L.F., Vermulst, M., Tomazela, D.M., Emond, M.J., MacCoss, M.J.,
Gollahon, K., Martin, G.M., Loeb, L.A., Ladiges, W.C., Rabinovitch, P.S., 2009.
Overexpression of catalase targeted to mitochondria attenuates murine cardiac aging.
Circulation 119, 2789-2797.
Demontis, F., Patel, V.K., Swindell, W.R., Perrimon, N., 2014. Intertissue control of the
nucleolus via a myokine-dependent longevity pathway. Cell reports 7, 1481-1494.

Lakatta, E.G., Levy, D., 2003. Arterial and cardiac aging: major shareholders in cardiovascular
327   disease enterprises: Part II: the aging heart in health: links to heart disease. Circulation
328   107, 346-354.
329   Loffredo, F.S., Steinhauser, M.L., Jay, S.M., Gannon, J., Pancoast, J.R., Yalamanchi, P., Sinha,
330   M., Dall'Osso, C., Khong, D., Shadrach, J.L., Miller, C.M., Singer, B.S., Stewart, A.,
331   Psychogios, N., Gerszten, R.E., Hartigan, A.J., Kim, M.J., Serwold, T., Wagers, A.J.,
332   Lee, R.T., 2013. Growth differentiation factor 11 is a circulating factor that reverses
333   age-related cardiac hypertrophy. Cell 153, 828-839.
334   North, B.J., Sinclair, D.A., 2012. The intersection between aging and cardiovascular disease.
335   Circulation research 110, 1097-1108.
336   Quarles, E.K., Dai, D.F., Tocchi, A., Basisty, N., Gitari, L., Rabinovitch, P.S., 2015. Quality
337   control systems in cardiac aging. Ageing research reviews 23, 101-115.
338   Taneike, M., Yamaguchi, O., Nakai, A., Hikoso, S., Takeda, T., Mizote, I., Oka, T., Tamai, T.,
339   Oyabu, J., Murakawa, T., Nishida, K., Shimizu, T., Hori, M., Komuro, I., Takuji
340   Shirasawa, T.S., Mizushima, N., Otsu, K., 2010. Inhibition of autophagy in the heart
341   induces age-related cardiomyopathy. Autophagy 6, 600-606.
342   Aguilera, M. O., Berón, W., & Colombo, M. I. (2012). The actin cytoskeleton participates in the
343   early events of autophagosome formation upon starvation induced autophagy.
344   Autophagy, 8(11), 1590–1603. https://doi.org/10.4161/auto.21459
345   Bai, H., Kang, P., Hernandez, A. M., & Tatar, M. (2013). Activin Signaling Targeted by
346   Insulin/dFOXO Regulates Aging and Muscle Proteostasis in Drosophila. PLoS
347   Genetics, 9(11). https://doi.org/10.1371/journal.pgen.1003941
348   Ball, R.W., Warren-Paquin, M., Tsurudome, K., Liao, E.H., Elazzouzi, F., Cavanagh, C., An,
349   B.S., Wang, T.T., White, J.H., Haghighi, A.P., 2010. Retrograde BMP signaling
350   controls synaptic growth at the NMJ by regulating trio expression in motor neurons.
351   Neuron 66, 536-549
352   Kast, D.J., Zajac, A.L., Holzbaur, E.L., Ostap, E.M., Dominguez, R., 2015. WHAMM Directs
353   the Arp2/3 Complex to the ER for Autophagosome Biogenesis through an Actin Comet
354   Tail Mechanism. Current biology : CB 25, 1791-1797.
355   Lakatta, E. G., & Levy, D. (2003). Special Review : Clinical Cardiology : New Frontiers Arterial
356   and Cardiac Aging : Major Shareholders in Cardiovascular Disease Enterprises Part II :
357   The Aging Heart in Health : Links to Heart Disease. Circulation, 107(2), 139–146.
358   https://doi.org/10.1161/01.CIR.0000048892.83521.58
359   Lim, H.-Y., Wang, W., Chen, J., Ocorr, K., & Bodmer, R. (2017). ROS Regulate Cardiac
360   Function via a Distinct Paracrine Mechanism. Cell Reports, 7(1), 35–44.
361   https://doi.org/10.1016/j.celrep.2014.02.029
362   Mackay, T. F. C., Richards, S., Stone, E. A., Barbadilla, A., Ayroles, J. F., Zhu, D., … Gibbs, R.
363   A. (2012). The Drosophila melanogaster Genetic Reference Panel. Nature, 482(7384),
364   173–178. Retrieved from http://dx.doi.org/10.1038/nature10811
365   Massague, J. (2012). TGF[beta] signalling in context. Nat Rev Mol Cell Biol, 13(10), 616–630.
366   Retrieved from http://dx.doi.org/10.1038/nrm3434
367   Mozaffarian, D., Benjamin, E. J., Go, A. S., Arnett, D. K., Blaha, M. J., Cushman, M., …
368   Turner, M. B. (2016). Heart disease and stroke statistics-2016 update a report from the
369   American Heart Association. Circulation (Vol. 133).
370   https://doi.org/10.1161/CIR.0000000000000350

Nagy, P., Varga, Á., Kovács, A. L., Takáts, S., & Juhász, G. (2015). How and why to study
autophagy in Drosophila: It's more than just a garbage chute. Methods, 75, 151–161.
https://doi.org/10.1016/j.ymeth.2014.11.016
Neufeld, T. P. (2012). Autophagy and cell growth--the yin and yang of nutrient responses.
Journal of Cell Science, 125(Pt 10), 2359–68. https://doi.org/10.1242/jcs.103333
North, B. J., & Sinclair, D. A. (2012). The intersection between aging and cardiovascular
disease. Circulation Research, 110(8), 1097–1108.
https://doi.org/10.1161/CIRCRESAHA.111.246876
Ocorr, K., Fink, M., Cammarato, A., Bernstein, S., & Bodmer, R. (2009). Semi-automated
Optical Heartbeat Analysis of small hearts. Journal of Visualized Experiments : JoVE,
(31), 3–6. https://doi.org/10.3791/1435
Ocorr, K., Reeves, N. L., Wessells, R. J., Fink, M., Chen, H.-S. V., Akasaka, T., … Bodmer, R.
(2007). KCNQ potassium channel mutations cause cardiac arrhythmias in Drosophila
that mimic the effects of aging. Proceedings of the National Academy of Sciences of
the United States of America, 104(10), 3943–8.
https://doi.org/10.1073/pnas.0609278104
Shibutani, S. T., & Yoshimori, T. (2014). A current perspective of autophagosome biogenesis.
Cell Research, 24(1), 58–68. https://doi.org/10.1038/cr.2013.159
Strait, J., & Lakatta, E. (2012). Aging-associated cardiovascular changes and their relationship to
heart failure. Heart Failure Clinics, 8(1), 143–164.
https://doi.org/10.1016/j.hfc.2011.08.011.Aging-associated
Taneike, M., Yamaguchi, O., Nakai, A., Hikoso, S., Takeda, T., Mizote, I., … Otsu, K. (2010).
Inhibition of autophagy in the heart induces age-related cardiomyopathy. Autophagy,
6(5), 600–606. https://doi.org/10.4161/auto.6.5.11947
Upadhyay, A., Moss-taylor, L., Kim, M., Ghosh, A. C., & Connor, M. B. O. (2017). TGF- b
Family Signaling in Drosophila, (Table 1). https://doi.org/10.1101/cshperspect.a022152
Vogler, G., & Ocorr, K. (2009). Visualizing the beating heart in Drosophila. Journal of
Visualized Experiments : JoVE, (31), 6–8. https://doi.org/10.3791/1425
Wong, P. M., Puente, C., Ganley, I. G., & Jiang, X. (2013). The ULK1 complex sensing nutrient
signals for autophagy activation. Autophagy, 9(2), 124–137.
https://doi.org/10.4161/auto.23323
Woodring, P.J., Hunter, T., Wang, J.Y., 2005. Mitotic phosphorylation rescues Abl from F-actin-
mediated inhibition. The Journal of biological chemistry 280, 10318-10325.
Yogalingam, G., & Pendergast, A. M. (2008). Abl kinases regulate autophagy by promoting the
trafficking and function of lysosomal components. Journal of Biological Chemistry,
283(51), 35941–35953. https://doi.org/10.1074/jbc.M804543200
Yoshida, K., Yamaguchi, T., Natsume, T., Kufe, D., Miki, Y., 2005. JNK phosphorylation of 14-
3-3 proteins regulates nuclear targeting of c-Abl in the apoptotic response to DNA
damage. Nature cell biology 7, 278-285.
410

Figure 1. Arrhythmicity was measured and quantified as arrhythmia indexes across a 6-week and 8-week time courses (Mean ± SEM, 16-38 flies/data points. Among those, larger sample size were used in 6-week and 8-week time points to control increased variance caused by aging). Cardiac performance were measured every 2 weeks. Adult progeny of Dawdle RNAi stains (daw 50911 and daw 34974) crossed with heart tissue drivers (Hand4.2-Gal4 and TinΔ4-Gal4) has a significantly lower arrhythmia index in old flies [6-week-old and 8-week-old, *P < 0.05 (unpaired t-test)].

Figure 2. Knocking down Daw induced basal level autophagosome formation. (2A) The basal autophagosome number in young (2 week old) and old (8 week old) female flies was detected by Atg8a antibodies (Mean ± SEM, 4 flies/data points). The regions of interested (ROI) selected from A1/A2 abdominal segments of adult fly heart were used for *measure and count* analysis provided by CellSens (Olympus Corp.) [**P < 0.01, ***P < 0.001 (student's t-test)]. (2B) Quantification of autophagy flux by autophagosome number fold change after BafA1 treatment in young (2 week old) and old (8 week old) female flies [Mean ± SEM, 4 flies/data points, P < 0.05 (student's t-test)].

Figure 3. Activin/Daw signaling regulates cardiac aging through autophagy/lysosome system, independent of mTOR. (A) Western blots show Daw, not Myo treatment induces phosphorylation of 4E-BP in cultured Kc167 cells. (B) Daw RNAi induced autophagosome formation detected by Atg8a antibodies (C) and age-associated arrhythmicity independent of mTOR/TSC1. [Scale bar: 50 µm. *P < 0.05 (Student's t-test)].

Figure 4. Autophagy disruption via Atg1 RNAi led to premature cardiomyopathy in young flies. High-speed video microscopy and SOHA were used to quantitatively analyze beating D. melanogaster hearts at 2 weeks (Mean ± SEM, 16-20 flies/data points). Adult progeny of Atg1 RNAi stains crossed with heart tissue drivers (Hand4.2-Gal4 and TinΔ4-Gal4) has a significantly higher arrhythmia index. The diastolic interval also elevated significantly in adult progeny crossed with Hand4.2-Gal4 ([*P < 0.05, **P < 0.01, ***P < 0.001 (unpaired t-test)].

Figure 5. Autophagy disruption via CQ feeding in Daw RNAi old flies phenocopied wild type old flies cardiac performance. For chloroquine treatments, 100 ul of 20 mM chloroquine diphosphate salt (Sigma) was added onto the food. High-speed video microscopy and SOHA were used to quantitatively analyze beating D. melanogaster hearts at 6 weeks (Mean ± SEM, 14-20 flies/data points). After 24 hour chloroquine (CQ) treatment, the arrhythmia index for adult progeny of Daw RNAi stains crossed with heart tissue drivers (Hand4.2-Gal4) was brought back to wild type level. [*P < 0.05, (unpaired t-test)].