Activin Signaling Regulates Autophagy and Cardiac Aging through mTORC2

Kai Chang[1][§], Ping Kang[1][§], Ying Liu[1], Kerui Huang[1], Erika Taylor[2], Antonia P. Sagona[3], Ioannis

P. Nezis[3], Rolf Bodmer[2], Karen Ocorr[2], Hua Bai[1]*

1 Department of Genetics, Development, and Cell Biology, Iowa State University, Ames, IA

50011, USA

2 Development, Aging, and Regeneration Program, Sanford-Burnham-Prebys Medical

Discovery Institute, La Jolla, California 92037, USA

3 School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK

*Corresponding Author:

Hua Bai

Phone: 515-294-9395

Email: hbai@iastate.edu

§ These authors contribute equally to this work.

21 **Running title:**

22 **Activin promotes cardiac aging through mTORC2**

Abstract

Age-dependent loss of cardiac tissue homeostasis largely impacts heart performance and

contributes significantly to cardiovascular diseases later in life. Cellular quality control

machinery, such as autophagy/lysosome system, plays a crucial role in maintaining cardiac

health and preventing age-induced cardiomyopathy and heart failure. However, how aging alters

the autophagy/lysosome system to impact cardiac function remains largely unknown. Here using

Drosophila heart as a model system, we show that activin signaling, a member of TGF-beta

superfamily, negatively regulates cardiac autophagy and cardiac health during aging. We found

that cardiac-specific knockdown of *Daw*, an activin-like protein in *Drosophila*, increased cardiac

autophagy and prevented age-related cardiac dysfunction, including arrhythmia and bradycardia

(slow heart rate). Inhibition of autophagy blocked *Daw* knockdown-mediated cardioprotection.

Consistently, cardiac-specific expression of constitutively activated activin type I receptor *Babo*

disrupted cardiac function at young ages. Intriguingly, the key autophagy regulator, mechanistic

target of rapamycin complex 1 (mTORC1), was not involved in activin-mediated autophagy.

Instead, activin signaling genetically interacted with Rictor, the key subunit of mTORC2, to

regulate autophagy and cardiac aging. Knockdown of *Daw* increased the mRNA expression of

Rictor and the phosphorylation of AKT in fly hearts. Finally, cardiac-specific silencing of *Daw*

not only improved cardiac health, but also prolonged lifespan. Thus, our findings highlight an

emerging role of activin signaling and mTORC2 in the regulation of autophagy and cardiac

aging.

Key words: Activin ligand, Dawdle, Atg8a, TSC1, Rictor, mTOR complex 2, Autophagic flux,

Semi-automatic optical heartbeat analysis (SOHA)

Introduction

Aging is associated with an exponential increase of the incidence of cardiovascular
diseases (CVD) [1, 2]. Age-related changes in cardiovascular structure and output have been linked
to increased risk of coronary heart disease, sudden cardiac death and stroke in elderly population
[3]. During normal aging, the left ventricular wall of human hearts becomes thickened and the
diastolic filling rate of left ventricle gradually decreases. On the other hand, the left ventricular
systolic performance at rest remains no change with age [3]. Several mechanisms underlying these
age-associated changes in cardiovascular structure and function have been proposed, for example
changes in growth factor signaling, decreased cellular quality control, altered calcium handling,
elevated extracellular matrix deposition or fibrosis, increased mitochondria damage, and the
production of reactive oxygen species (ROS) [2, 4]. Resolving the contributing mechanisms
underlying age-dependent decline of cardiovascular function is critical for the development of
therapeutic interventions for the treatment of cardiovascular diseases.

Cellular quality control systems, such as macroautophagy (hereinafter referred to as
autophagy), are essential to the maintenance of tissue homeostasis during aging [5, 6]. Autophagy is
a highly conserved process that is responsible for the degradation and recycling of damaged
organelles, protein aggregates and other cytoplasmic substances. The autophagic process initiated
with the isolated membrane, or phagophore that elongates to form a double-membrane structure,
the autophagosome. Then the autophagosome fuses with a lysosome to form an autolysosome to
degrade the cargo within the autolysosome [7]. It is generally accepted that autophagy activity
declines with age [8-10]. Disruption of autophagy pathways often leads to loss of tissue homeostasis
and tissue dysfunction. For example, knocking out autophagy gene Atg5 in the mouse heart
accelerates cardiac aging, including an increase in left ventricular hypertrophy and decrease in
factional shortening [10]. As a crucial autophagy regulator, mechanistic target of rapamycin (mTOR),
especially mTOR complex 1 (mTORC1), plays an important role in the regulation of cardiac tissue
homeostasis [2]. Rapamycin-mediated suppression of mTORC1 induces autophagy and protects
mouse cardiomyocytes from oxidative stress [11]. In *Drosophila*, mTORC1 is known to promote
electrical pacing-induced heart failure [12, 13], and high-fat-diet-induced cardiomyopathy [14]. The
mTOR signaling consists of two complexes, mTORC1 and mTORC2. It is not known whether
mTORC2 also contributes to the regulation of autophagy [15]. Although inhibition of mTOR is one
of the robust longevity interventions [16, 17], it remains unclear whether mTOR pathway plays a

major role in age-dependent decline of autophagy. Notably, a recent study showed that mTORC1

activity does not increase with age in most tissues of C57BL/6J.Nia mice [18]. Thus, the mechanistic

basis for age-dependent alterations in autophagy and the contribution of mTORC1 remains to be

determined.

Autophagy can be activated by various stresses, such as dietary restriction, hypoxia,

oxidative stress, and pathogen infection [19]. The activation of autophagy is tightly controlled by key

cellular signaling pathways, in particular the nutrient signaling pathways like mTORC1 and AMP-

activated protein kinase (AMPK) [20]. These pathways are often involved in control of longevity.

We and other groups recently found that activin signaling, a transforming growth factor beta (TGF-

beta) subfamily, regulates muscle proteostasis and longevity in *Drosophila* [21, 22]. As in mammals,

TGF-beta family in *Drosophila* has two major branches, bone morphogenetic protein (BMP) and

activin signaling pathways [23]. In both pathways, signaling starts with ligand binding to a receptor

complex composed of type I and type II receptor kinases, followed by phosphorylation of receptor-

activated Smad (R-Smad). *Drosophila* activin signaling is known to regulate neuroblast

proliferation, axon guidance, and metabolic homeostasis during development (reviewed in [24]). The

regulation of adult physiology by *Drosophila* activin is not well understood. We previously

showed that knockdown of one of the three activin ligands, *Daw*, prolongs lifespan and increases

autophagosome formation in flies [21]. However, the role of TGF-beta/activin signaling in cardiac

aging remains largely unknown.

It has been shown that TGF-beta factors promote cardiac fibrosis during aging [25]. Mice

with heterozygous mutants for activin receptor-like kinases (ALK4) are protected from pressure

overload-induced fibrosis and dysfunction [26, 27]. Inhibition of ALK5 also reduces myocardial

infarction-induced systolic dysfunction and left ventricular remodeling in rats [28]. However,

conflicting results are reported in other studies showing that activin A protects hearts from

hypoxia/reoxygenation- and ischemia/reperfusion-induced cell death [29]. In addition, it is unknown

whether TGF-beta family proteins, in particular activin, regulate cardiac homeostasis and function

through autophagy. Given that the cardiac development and aging are remarkably conserved

between *Drosophila* and mammals, we investigated the role of activin signaling in the regulation

of cardiac autophagy and cardiac aging using *Drosophila* heart model and cardiac movement

detection techniques [30]. Here we show that activin signaling acts on hearts to regulate autophagy

and age-related cardiomyopathy in *Drosophila*. RNAi-mediated knockdown of the activin-like

protein *Daw* preserved cardiac function with age, while expression of activated activin type I

receptor *Babo* induced cardiomyopathy early in life. Inhibition of autophagy blocked the

cardioprotective effects by *Daw* knockdown. Intriguingly, activin signaling genetically interacted

with mTORC2 subunit Rictor (rapamycin-insensitive companion of mTOR), but not mTORC1 to

control autophagy and cardiac aging. Thus, our study uncovered a novel interaction between

activin and mTORC2 in the regulation of tissue homeostasis, autophagy, and age-related

cardiomyopathy.

Results

Heart-specific knockdown of activin-like ligand *Daw* slows cardiac aging

Our previous study demonstrated that the activin-like protein Daw regulated tissue

homeostasis (especially in flight muscle) and longevity in *Drosophila* [21]. Interestingly, we found

that activin signaling increased in aging fly hearts, as indicated by elevated phosphorylation of

Smad2 in heart nuclei (Fig. 1a). This finding suggest that activin signaling may play a role in

regulating cardiac aging. To investigate this possibility, we knocked down the expression of *Daw*

specifically in the heart with a binary GAL4/UAS system where we crossed *UAS-Daw RNAi*

lines into a cardiac-specific tissue driver (*Hand-gal4*). We analyzed cardiac performance in

young and old flies using a high-speed video imaging system (semi-automatic optical heartbeat

analysis, SOHA) [31]. M-Mode traces from the SOHA analysis indicated that knockdown of *Daw*

preserved cardiac contractility at advanced ages (Fig. 1b). In wild-type flies, normal aging is

associated with increased cardiac arrhythmia, diastolic intervals, heart period (bradycardia, slow

heart rate) (Figs. 1c, 1d, 1e), and decreased cardiac output (Fig. 1f). Systolic intervals normally

remain unchanged (Fig. 1g). Interestingly, cardiac-specific knockdown of *Daw* attenuated age-

dependent increase in arrhythmia, diastolic intervals, and heart period (Figs. 1c, 1d, 1e). *Daw*

knockdown also maintained relatively normal cardiac output at advanced ages (Fig. 1f). Similar

results were observed from three independent Daw RNAi lines (the knockdown efficiency of

Daw RNAi was verified by qRT-PCR in our previous studies [21]).

The *Hand-gal4* line drives gene expression in both cardiomyocytes and pericardial cells

[32]. To test the tissue-specific role of Daw in cardiac aging, we first crossed Daw RNAi into a

cardiomyocyte-specific driver *TinC-gal4* [33]. Similar to the results from *Hand-gal4* (Fig. 1h),

cardiomyocyte-specific knockdown of *Daw* prevented age-related increase in cardiac arrhythmia

(Fig. 1i). We also tested whether *Daw* expression in pericardial cells can regulate cardiac aging

using a pericardial cell-specific driver *Dot-gal4* [34]. Interestingly, knockdown of *Daw* in

pericardial cells also attenuated age-induced arrhythmia (Fig. 1j), but not the increase in diastolic

interval (Fig. 1k). Thus, Daw appears to promote cardiac aging, in particular age-induced

arrhythmia, through both cardiomyocyte-specific regulation and non-autonomous signaling from

pericardial cells.

Cardiomyocyte-specific knockdown of activin receptor *Babo* delays cardiac aging

In *Drosophila*, activin-like ligand Daw signals through the type I receptor Babo, the fly

homolog of mammalian ALK4 (activin receptor type-1B) [23]. To investigate whether Daw acts

through its receptor Babo to modulate cardiac aging, we first asked if Babo plays a similar role in

preventing age-dependent increase in cardiomyopathy. Consistent with the results from *Daw*

knockdown, we found that cardiomyocyte-specific knockdown of *Babo* blocked the age-related

increase in cardiac arrhythmia, diastolic intervals, and heart period (Figs. 2a, 2b, 2c). Two

independent Babo RNAi lines were used and both gave similar results. Systolic intervals

remained unchanged in both controls and *Babo* RNAi flies (Fig. 2d). Knockdown of *Babo* did

not affect age-dependent changes in cardiac output (Data not shown).

In a reciprocal experiment, we used a GeneSwitch heart driver (*Hand-GS-gal4*) to

express a constitutive active form of *Babo* (*Babo-Act, or Babo.Q302D*) specifically in adult

hearts. We found that cardiac expression of activated *Babo* significantly increased cardiac

arrhythmia, diastolic intervals, and heart period already at young ages (Figs. 2e, 2f, 2g). Taken

together, these results suggest that Babo plays an important role in age-related cardiomyopathy,

especially cardiac arrhythmia, diastolic function and heart rate.

Daw negatively regulates autophagy in fly hearts

Autophagy is one of the key mechanisms in maintaining tissue hemostasis, and its

activity normally declines with age [6, 9, 10]. Our previous studies showed that activin signaling

regulates autophagosome formation and Atg8a transcription [21]. It is likely that Daw regulates

cardiac aging by modulating autophagy. To first verify the role of Daw in autophagy-lysosome

activity, we used LysoTracker staining to monitor the lysosomal acidification in adult fat body, a

well-established autophagy marker in flies [35]. Consistent with our previous finding, heterozygous

Daw[11] loss-of-function mutants showed increased LysoTracker staining, suggesting an elevated

basal autophagy/lysosome activities (Fig. 3a). Using a lipidation assay, we furthermore showed

that mutation of *Daw* increased the levels of lipidated Atg8a, the fly homolog of mammalian

LC3/GABARAP family proteins (Fig. 3b). Finally, mosaic analysis revealed that somatic cell

clones in larval fat body expressing RNAi against either Daw or Babo (marked by GFP) showed

increased numbers of autophagosome, indicated by mCherry-Atg8a puncta (Fig. 3c). The data

suggest that activin signaling negatively regulates autophagy and lysosome activities.

Next, we examined whether Daw plays a similar role in modulating autophagy in fly

hearts. Using a commercial antibody for *Drosophila* Atg8a protein (Fig. S1 shows the antibody

validation), we found that the number of autophagosome in the heart was higher in *Daw[11]*

mutants compared to wild-type (Fig. S2a). Because autophagy is a dynamic process, the

increased autophagosome number could result from increased autophagy activities, or blockage

of lysosomal degradation [36]. We next examined the role of Daw in the regulation of autophagic

flux (the turnover of autophagosome). We exposed semi-intact fly hearts to a V-ATPase inhibitor

bafilomycin A1 (Baf A1), which blocks both lysosomal acidification and autophagosome-

lysosome fusion [37]. We found that Baf A1 treatment resulted in a higher increase in the number

of Atg8a-positive autophagosome in *Daw[11]* hearts compared to wild-type hearts (Figs. S2a-

S2c). Thus, the results suggest that Daw negatively regulate autophagic flux in fly hearts.

Aging tissues show reduced autophagy and accumulated cellular damages. We next

examined whether inhibition of activin/Daw signaling in fly hearts can rescue age-dependent

decline of autophagy. Using a mCherry-Atg8a reporter line, we observed a dramatic

accumulation of Atg8a-positive autophagosome in the heart (Fig. 3d, 3e, 3h). The

cardiomyocyte-specific driver *TinC-gal4* was used. The accumulation of autophagosome is

likely due to the blockage of autophagosome turnover, because increases numbers of Atg8a-

positive puncta were only seen in Baf A1-treated young hearts, but not the old hearts (Figs. 3e,

3h, 3j). Consistently, the adaptor protein p62/Ref(2)P, a marker for defective aggrephagy [38], was

accumulated in aging hearts and Atg8a mutants (Fig. S3). Intriguingly, flies with heart-specific

Daw knockdown maintained high levels of cardiac autophagosome turnover (or active

autophagic flux) at both young and old ages, as indicated by the induction of autophagosome

number upon Baf A1 treatment (Figs. 3f, 3h, 3j). Notably, constitutively activated Babo

repressed the autophagic flux at both young and old ages (Figs. 3g, 3h). Activation of Babo also

blocked the accumulation of autophagosome in old fly hearts, which suggests that besides the

regulation of autophagosome turnover, activin signaling also play an important role in

autophagosome initiation and formation (Figs. 3g, 3h). We were able to verify these phenotypes

using another heart-specific driver *Hand-Gal4* (Fig. 3i). Taken together, our findings suggest that

inhibition of activin/Daw signaling promotes autophagosome turnover and maintains healthy

autophagic activity in old hearts.

Inhibition of autophagy, but not activation of mTORC1, blocks *Daw* knockdown-mediated

cardioprotection

Since we observed an induction of autophagy in *Daw* knockdown flies, we next asked

whether autophagy activity is required for Daw-regulated cardiac aging. First, we fed control and

Daw knockdown flies with lysosomal inhibitor chloroquine (CQ). As expected, CQ treatment

increased cardiac arrhythmia in *Daw* knockdown flies at old age (Fig. 4a). To directly test

whether autophagy plays any role in Daw-regulated cardiac aging, we generated double

knockdown flies by combining *UAS-Daw RNAi* and *UAS-Atg1 RNAi* fly lines. Atg1 is a

serine/threonine-protein kinase that is essential for autophagy initiation. Cardiac-specific

knockdown of *Atg1* increased both cardiac arrhythmia and diastolic intervals at young ages (Fig.

4b), similar to the fly heart with activated Babo (Figs. 2e, 2f). Consistent with CQ treatment,

cardiac-specific knockdown of *Atg1* attenuated cardioprotective effects of Daw RNAi, since

simultaneously knockdown of *Atg1* and *Daw* in the heart led to age-dependent increase in

arrhythmia similar to control flies (Fig. 4c). Thus, these results suggest that autophagy and Atg1

are required for Daw-regulated cardiac aging.

mTORC1 is a key negative regulator of autophagy [35]. We then tested whether Daw could

negatively regulate autophagy and cardiac aging through activation of mTORC1 signaling. To

test this possibility, we first generated double knockdown flies by combining *UAS-Daw RNAi*

and *UAS-Tsc1 RNAi*. Tsc1 (Tuberous sclerosis protein 1) is a negative regulator of mTORC1

(Fig. S4a). Surprisingly, activation of mTORC1 via *Tsc1* knockdown had no effects on cardiac

arrhythmia (Fig. 4d), but reduced diastolic intervals at young age (Fig. 4e), which is very

different from the cardiomyopathy resulted from cardiac-specific expression of activated activin

receptor *Babo* (Figs. 2e, 2f). Furthermore, knockdown of *Tsc1* did not rescue the

cardioprotective effects of *Daw* knockdown during aging (Fig. 4f). Simultaneously knockdown

of *Tsc1* and *Daw* did not show age-dependent increase in arrhythmia. Similarly, loss-of-function

mutant *Tsc1*[12] did not rescue the elevated LysoTracker staining in *Daw*[11] (Fig. 4g). These

results suggest that activation of mTORC1 is not required for Daw-regulated autophagy and

cardiac aging.

253   It is known that inhibition of mTORC1 prolongs lifespan [16, 39] and attenuates stress-

induced cardiomyopathies [12, 13, 40, 41]. However, the effects of mTORC1 induction on cardiac

aging are not well understood. Here, we observed unique changes of cardiac function in response

to activation of mTORC1 (via *Tsc1* knockdown), including decreased heart period (Tachycardia,

fast heart rate), and reduced diastolic intervals (Figs. 4d, 4e) These cardiac functional changes

are not the same as age-induced cardiomyopathies, rather they resemble high-fat-diet-induced

cardiac complications where mTORC1 is known to play an important role [14]. Intriguingly, flies

with cardiac-specific knockdown of *Tsc1* showed no age-dependent increases in heart period,

diastolic intervals, and arrhythmia (Figs. S4b, S4c, S4d). These unique alterations in cardiac

function were also observed in other mTORC1 activation mutants, such as 1) Knockdown of

Reptor (Repressed by TOR), a recently identified bZIP transcription factor that is negatively

regulated by mTORC1 [42] (Figs. S4e, S4f, S4g), and 2) Over-expression of *Rheb* (Ras Homolog

Enriched in Brain), a positive regulator of mTORC1 (Figs. S4h, S4i, S4j).

Daw genetically interacts with mTORC2/Rictor to control autophagy and cardiac aging

268   Although Daw did not interact mTORC1 to regulate autophagy and cardiac aging,

cardiac-specific knockdown of *Daw* induced the phosphorylation of AKT, the upstream regulator

of mTORC1 (Figs. 5a, 5b). On the other hand, activated receptor Babo reduced cardiac AKT

phosphorylation (Figs. 5c, 5d). It is known that AKT can be activated and phosphorylated by two

kinases at distinct positions, PDK1 (phosphoinositide-dependent kinase 1) at Thr308 and Rictor

(major component of mTORC2) at Ser473 (Ser505 in flies) [43]. Cardiac-specific over-expression

Rictor can also induce AKT phosphorylation (Figs. 5e, 5f). Interestingly, knockdown of *Daw* up-

regulated the mRNA expression of *Rictor* in fly hearts (Figs. 5g, 5h). These results suggest that

Daw is a negative regulator of mTORC2/Rictor.

277   Unlike mTORC1, the role of mTORC2 in the regulation of autophagy is unknown.

Intriguingly, we found that loss-of-function mutation in *Rictor* repressed the high lysosomal

activities in *Daw*[11] mutants (Figs. 5i, 5j, 5k, 5l). Similarly, mutations in Sin1, another core

component of mTORC2, blocked the elevated lysosomal activity in *Daw* mutants. Furthermore,

mosaic analysis revealed that over-expression of Rictor induced lysosome activity (Fig. 5m).

Interestingly, unlike *Tsc1[12]* mutants, *Rictor[42]* mutants did not repress starvation-induced

lysosome activity (Fig. S5c). Similarly, activated Babo also did not block starvation-induced

lysosome activity (Fig. S5d). These data suggest that activin signaling and mTORC2/Rictor

regulate autophagy differently from mTORC1, and Rictor acts downstream of activin signaling

in the regulation of autophagy.

Given that Rictor positively regulates autophagy, we predict that Rictor is a cardiac

protective factor. Indeed, cardiac-specific over-expression of *Rictor* blocked age-related

increases in diastolic intervals and arrhythmia (Figs. 6a, 6b), while knockdown of *Rictor* induced

arrhythmia, especially at old ages (Fig. 6c). Interestingly, over-expression of *Rictor* attenuated

the elevated diastolic intervals and arrhythmia in fly hearts with constitutively activated *Babo*

(Figs. 6d, 6e), suggesting that fly activin signaling regulates cardiac aging through

mTORC2/Rictor (Fig. 6f).

Cardiac-specific reduction of activin-like protein *Daw* prolongs lifespan

Tissue-specific hormonal signaling in systemic aging control have been previously

reported [8, 44, 45]. However, whether longevity can be achieved by maintaining healthy hearts

remains unclear. Here we tested the effects of cardiac-specific knockdown of activin signaling in

longevity control. Knockdown of *Daw* in the heart using two cardiac tissue drivers (*Hand-gal4*

and *TinC-gal4*) significantly prolongs lifespan (range from 14% to 32% extension of median

lifespan) (Figs. 7a-7e). Compared to *Hand-gal4*, cardiomyocyte-specific driver *TinC-gal4*

produced stronger lifespan extension. Two independent lifespan analyses were performed and

both showed similar lifespan phenotypes (Fig. 7e). In the first lifespan trial, multiple control

lines (dashed lines in Fig. 7a) and two Daw RNAi lines (solid lines in Fig. 7a) were included.

Because Daw is a hormonal factor, it is unclear whether Daw regulates lifespan through cell-

autonomous or non-autonomous mechanisms. However, our previous study showed that Daw

regulates longevity in a tissue-specific manner [21]. Flight muscle-specific knockdown of *Daw*

prolongs lifespan, while fat body-specific knockdown of *Daw* shortens lifespan. Therefore, it is

possible that cardiac-specific Daw signaling regulates longevity through maintenance of cardiac

health and production of unknown systemic factors from the fly heart.

Discussion

TGF-beta signaling plays vital roles in a wide range of human diseases, including cancer and cardiovascular diseases [46]. Originally identified as a reproductive hormone, activin, a TGF-beta subfamily member, has become an emerging target for the treatment of many human diseases [47]. In the present study, we investigated the role of activin in cardiac aging using a *Drosophila* heart model. Similar to our previous study in *Drosophila* flight muscle [21], reduction of activin signaling prevented age-related decline of cardiac function. We think that this is partly due to the activation of autophagy. *Daw* mutant flies exhibited high autophagic flux and elevated lysosome activities. Genetic analysis further showed that autophagy gene Atg1 is required for Daw-regulated cardiac aging. Importantly, we found that activation of mTORC1 did not rescue the autophagy and cardiac aging phenotypes in *Daw* knockdown flies. Instead, activin/Daw signaling interacts with mTORC2 subunit Rictor to regulate autophagy and cardiac function. Thus, our studies uncover an emerging role of activin signaling and mTORC2 in the regulation of autophagy and cardiac tissue homeostasis.

Although previous studies reveal an interesting link between TGF-beta family proteins and animal aging [21, 48-50], the role of activin in cardiac aging is not well understood. It's known that activin A serum levels significantly increase with age [51]. The mRNA expression of *activin A*, as well as serum activin A levels are positively correlated with heart failure [52, 53], and hypertension in the elderly [54]. These findings suggest that activation of activin signaling at old ages might be detrimental to the heart. Studies showed that activin A promotes cardiac fibrosis and myocardial damage after ischemia reperfusion [55]. Consistently, mice with heterozygous mutations of ALK4 are protected from pressure overload-induced fibrosis and dysfunction [26, 27]. Inhibition of ALK5 also reduced myocardial infarction-induced systolic dysfunction and left ventricular remodeling in rats [28]. However, other studies suggest that activin A protects hearts from hypoxia/reoxygenation- and ischemia/reperfusion-induced cell death [29], even though it can promote cardiac apoptosis at higher concentrations [56]. The findings from the present study support the negative regulatory role of activin signaling in cardiac function, in particular age-related cardiac dysfunction (e.g., arrhythmia). This is consistent with the pro-aging role of TGF-beta and activin [21, 50].

Activin signaling has been implicated in the regulation of a wide range of cellular

processes, including cell death and proliferation, inflammation, fibrosis, and metabolic

homeostasis [47, 57-59]. The role of activin signaling in the control of autophagy, a key tissue

maintenance process, was only identified recently [21, 60-63]. In *C. elegans*, reduction of activin-like

protein Daf-7 suppressed beta-amyloid peptide-induced autophagosome accumulation [62, 63].

Another recent study found that activin A blocked oxygen-glucose deprivation-induced

autophagy through the inhibition of JNK and p38 MAPK pathways in neuronal PC12 cells [60].

Our previous study showed that knockdown of *Drosophila* activin-like protein *Daw* in flight

muscle induced autophagosome formation and autophagy gene expression [21]. However, how

activin signaling regulates autophagy and how this regulation contributes to tissue homeostasis

during aging remain largely unknown. Our findings in the present study suggest that the fly

activin-like protein Daw promotes cardiac aging through inhibition of autophagy. It seems that

activin signaling controls both autophagosome formation and turnover. Importantly, fly hearts

with reduced *Daw* expression maintain high levels of autophagosome turnover (autophagic flux)

throughout the life, whereas the autophagosome turnover is largely blocked in aged hearts in

wild-type flies. The active autophagosome turnover (and degradation capacity) at old ages is

likely responsible for the healthy hearts seen in *Daw* knockdown flies.

mTORC1 is a well-known autophagy regulator and it inhibits autophagosome formation

by increasing the phosphorylation of the serine/threonine protein kinase Atg1 (ULK1/ULK2 in

mammals), the initiator of the autophagy machinery [7]. Intriguingly, activation of mTORC1 did

not block the induction of autophagosome formation in *Daw* knockdown flies, suggesting that

there might be unknown mTORC1-independent mechanisms involved in Daw-regulated

autophagy and cardiac aging. mTORC1-independent autophagy regulation has been observed in

several recent studies [64-66]. A genome-wide screen study identified an important role of type III

PI3 kinases in autophagy control under normal nutritional conditions [64]. The screen also revealed

that growth factor signaling pathways, such as MAPK-ERK and AKT/FOXO3, negatively

regulate autophagy by inhibiting the type III PI3 kinase cascade. Interestingly, we noticed that

knockdown of *Daw* led to elevated AKT phosphorylation and the expression of mTORC2

subunit *Rictor*, suggesting mTORC2/AKT signaling may be involved in Daw-regulated

autophagy and cardiac aging. Indeed, we found that mutation of *Rictor* blocked the activation of

autophagy in *Daw* knockdown flies. mTORC2 is one of the two mTOR complexes that is

insensitive to short-term rapamycin treatment, and is involved in glucose homeostasis and actin

cytoskeleton reorganization [67, 68]. The direct regulatory role of mTORC2/Rictor in cellular

autophagy has not been previously established, although a few studies suggest that Rictor may be

involved in autophagy activation under unique circumstances. For example, Rictor is required for

resveratrol-induced autophagy in rat myocardium and H9c2 cardiac myoblast cells [69]. In rat

kidney NRK-52E cells, silencing Rictor blocked cisplatin induced autophagy [70]. Our findings

align with these studies suggesting that Rictor is a positive regulator of autophagy. However,

conflicting findings are reported in other studies showing Rictor negatively regulates

autophagosome formation in mouse skeletal muscle [71], and LC3-II levels in human senescent

endothelial cells [72]. Thus, more studies are needed to better understand the role of

mTORC2/Rictor in the regulation of autophagy and its contribution to tissue homeostasis during

aging.

The positive role of Rictor in autophagy control identified in the present study is

consistent with the cardioprotection effects of Rictor. We found that cardiac-specific over-

expression of *Rictor* slows cardiac aging, similar to *Daw* knockdown. Our findings are consistent

with two recent studies showing that disruption of mammalian Rictor induced cardiac

dysfunction [73, 74]. Cardiac-specific deletion of Rictor in mice induced dilation and decreased

fractional shortening by activating MST1 kinase and hippo pathway [74]. Unlike mTORC1

mutants, Rictor null mutants (in both worms and mice) are short-lived [75, 76], suggesting the

detrimental role of mTORC2/Rictor in regulating tissue aging. The role of *Drosophila* Rictor in

longevity regulation is not yet known, although it has been shown that *Drosophila* mutants of

Rictor and Sin1 decreased the tolerance to heat stress [77] and over-expression of *Rictor* rescued

Pink1 knockdown-induced mitochondrial aggregation in flight muscle [78]. The Rictor-Pink1

interaction suggests that Rictor could be a potential positive regulator for mitophagy, a key

process for the mitochondria quality control during aging.

During normal aging, the heart undergoes complex phenotypic changes such as

progressive myocardial remodeling, reduced myocardial contractile capacity, increased left

ventricular wall thickness and chamber size, prolonged diastole as well as increased arrhythmia

[3]. All of these biological changes can gradually alter cardiac functions and confer vulnerability of

the heart to various cardiovascular stresses, thus increasing the chance of developing

cardiovascular disease dramatically. The present study has uncovered an important role of TGF-

beta/activin signaling pathway in the regulation of autophagy and cardiac aging in *Drosophila*.

We also found an intriguing interaction between activin and mTORC2 in the regulation of

cardiac autophagy, as well as a new role of Rictor in cardiac aging in *Drosophila*. It remains to

be determined how activin regulates Rictor in fly hearts. It is possible that activin signaling

regulates Rictor through both transcriptional activation (e.g., through insulin/FOXO signaling [79])

and post-translational modification.

Additionally, cardiac-specific knockdown of activin-like protein *Daw* extended fly

lifespan, suggesting that longevity can be achieved by maintaining a healthy heart. Our findings

also suggest that Daw itself or other unidentified hormonal factors secreted from cardiomyocytes

play an important role in systemic aging control. Hence, it remains to be determined the

mechanistic basis for activin-mediated control of cardiac aging and longevity, which will

eventually help the development of therapeutic interventions targeting activin for the treatment

of age-related cardiovascular diseases.

Materials and methods

Fly husbandry and stocks

Flies were maintained at 25 ℃, 60% relative humidity and 12-hour light/dark (40%

relative humidity was used in the experiments from Figures 1&2, which were performed in

Bodmer laboratory). Adults were reared on agar-based diet with 0.8% cornmeal, 10% sugar, and

2.5% yeast (unless otherwise noted). Fly stocks used in the present study are: *UAS-Daw RNAi*

(Bloomington *Drosophila* Stock Center (BDSC) #34974, BDSC #50911, and Vienna Drosophila

Resource Center (VDRC) #105309), *UAS-Babo RNAi* (BDSC #25933, BDSC #40866), *UAS-*

Babo-Act (also known as *Babo.Q302D*) [80], *UAS-Tsc1 RNAi* (BDSC #52931, #54034), *UAS-*

Reptor RNAi (BDSC #25983), *UAS-Rheb* (BDSC #9688), *UAS-Atg1 RNAi* (BDSC #26731),

UAS-Rictor RNAi (BDSC #36699), *Daw[11]* [81], *Tsc1[12]* [82], *Rictor[42]* [83] , *Sin1[e03756]* (BDSC

#18188), *UAS-Rictor[77], UAS-HA-Rictor[77], Hand-gal4* (also known as Hand4.2-gal4) [32], and

TinC-gal4 (also known as *TinCΔ4-gal4*) [33], *Dot-gal4* [34], *Hand-GS-gal4* [84], *UAS-mCherry-Atg8a*

(BDSC #37750), *Atg8a[Δ4]* [85].

UAS-Daw RNAi (BDSC #34974), *UAS-Babo RNAi* (BDSC #25933), and *UAS-Myo RNAi*

(BDSC #31114) were backcrossed into *yw[R]* background for 5~7 generations, and *yw[R]* flies were

used as control or wild-type (WT) flies in most of the experiments. For other UAS-RNAi lines

that were not backcrossed to yw^R, following genotypes were used as control: $y^1 sc^* v^1$;

P[VALIUM20-mCherry]attP2 (BDSC #35785), $y^1 v^1$; *P[CaryP]attP2* (BDSC #36303), or $y^1 v^1$;

P[CaryP]attP40 (BDSC # 36304). Female flies were used in all experiments. RU486

(mifepristone, Sigma, St. Louis, MO, USA) were used to activate *Hand-GS-Gal4* at a final

concentration of 200 μM mixed in food.

Mosaic analysis

Two FLPout lines were used: 1). *hs-flp; endogenous P-3x mCherry-Atg8a, UAS-*

GFP/Cyo; Act>CD2>Gal4,UAS-Dcr2/TM6) (kindly provided by Gábor Juhász). 2). *yw, hs-flp,*

UAS-CD8::GFP; Act>y+>Gal4,UAS-GFP.nls;UAS-Dcr2/SM6::TM6B (kindly provided by Jun-

yuan Ji, originally generated by Bruce Edgar). To generate clones, freshly laid eggs (within 4-6

hr) were heated in a 37 ℃ water bath for 45 min. Early L3 larvae (84 hours after egg laying)

were used in the experiments.

Fly heartbeat analysis

To measure cardiac function parameters, semi-intact *Drosophila* adult fly hearts were

prepared according to previously described protocols [31]. High-speed 3000 frames movies were

taken at a rate of 100 frames per second using a Hamamatsu ORCA-Flash4.0 digital CMOS

camera (Hamamatsu Photonics, Hamamatsu, Japan) on an Olympus BX51WI microscope with a

10X water immersion lens (Olympus, Waltham, MA, USA) (Hamamatsu EM-CCD 9300 camera

was used in the experiments from Figures 1&2, which were performed in Bodmer laboratory).

The live images that contain heart tub within abdominal A3 segment were processed using HCI

imaging software (Hamamatsu Photonics, Hamamatsu, Japan). M-modes and cardiac parameters

were generated using SOHA, a MATLAB-based image application as described previously [31, 86].

The M-mode provides a snapshot of movement of heart wall over time. Four cardiac parameters

were analyzed, including heart period, diastolic interval, systolic interval, and arrhythmia index.

Diastolic interval (DI) is the duration for each heart relaxation phase (diastole). Heart period

(HP) is the pause time between the two consecutive diastole (HP is the reciprocal of heart rate).

Systolic interval (SI) was calculated as the HP minus the DI. Arrhythmia index (AI) is the

standard deviation of all HP in each fly normalized to the median HP. Cardiac output was

calculated using following equation: (π r(d)^2 - π r(s)^2) x HR. r(d) is radius of the heart tube at

diastolic phase, while r(d) is radius of the heart tube at systolic phase.

Quantitative RT-PCR

RNA extraction and cDNA synthesis were performed using Cells-to-CT kit (Thermo

Fisher Scientific, Waltham, MA, USA) from ~15 dissected adult hearts. QRT-PCR was

performed with a Quantstudio 3 Real-Time PCR System (Thermo Fisher Scientific, Waltham,

MA, USA). mRNA abundance of each gene was normalized to the expression of ribosomal

protein L32 (*RpL32* or *rp49*) by the method of comparative C_T . Primer sequences are listed in

Supplementary Table S1.

Antibodies and immunostaining

Since commercial antibodies for *Drosophila* Atg8a show non-specific staining in

immunostaining of fly tissues (data not shown), we used a GABARAP (E1J4E) antibody (1:300)

(Cell Signaling Technology (CST) #13733, Danvers, MA, USA) to stain and detect endogenous

Drosophila Atg8a. The epitope used to generate GABARAP antibody has identical amino acid

sequences to *Drosophila* Atg8a protein (personal communication with Cell Signaling

Technology, and see sequence alignment in Fig. S1a). This antibody has been previously used in

several *Drosophila* autophagy studies [87-89], and verified by Kim et al. [90]. We further verified the

specificity of the GABARAP antibody using an Atg8a loss-of-function mutant *Atg8a$^{\Delta 4}$* (Figs.

S1b-S1d). Other antibodies used for immunostaining are list below: Ref(2)P (1:1000) [38],

Phospho-Smad2 antibody (1:500) (CST #3108), Phospho-Akt (Ser473) (1:300) (CST # 4060),

Alexa Fluor 488-conjugated Phalloidin (or Alexa Fluor 594) for F-actin staining (Thermo Fisher

Scientific, Waltham, MA, USA). All fluorescence-conjugated secondary antibodies were from

Jackson ImmunoResearch, West Grove, PA, USA).

For immunostaining, adult female flies were collected and dissected in PBS. Hearts were

fixed in 4% paraformaldehyde for 15 min at room temperature (RT). After washing in PBS with

0.1% Triton X-100 (PBST), the fixed hearted were blocked in 5% normal goat serum (NGS)

diluted in PBST for 1 hour at RT. Hearts were then washed with PBST and incubated overnight

at 4 ℃ with primary antibodies diluted in 5% NGS. After washing with PBST, the samples were

incubated for 2 hours at RT with appropriate fluorescence-conjugated secondary antibodies.

Hearts were mounted in ProLong Gold anti-fade reagent (Thermo Fisher Scientific, Waltham,

MA, USA) before imaged using an epifluorescence-equipped BX51WI microscope (Olympus,

Waltham, MA, USA).

Imaging analysis

For imaging analysis, fluorescence images were first processed using the deconvolution

module in Olympus cellSens Demensions software, and then the number of puncta in a selected

perinuclear region (~707 μm^2) was measured with the "Measure and Count" module in cellSens

software (Olympus, Waltham, MA, USA).

LysoTracker and Magic Red staining

Acidic organelles (including autolysosome) were monitored by staining tissue with 100

nM LysoTracker Red DND-99 (Invitrogen, Grand Island, NY, USA) for 5 minutes at room

temperature. Lysosomal cathepsin B activities were monitored using Magic Red Cathepsin-B

Assay kit (#938, ImmunoChemistry Technologies, Bloomington, MN, USA) following

manufacturer's manual. Nuclei were stained with either DAPI or Hoechst 33342 (1 μg/ml)

(ImmunoChemistry Technologies).

Western blotting for Atg8a lipidation

Antibodies for western blot included: β-actin antibody (1:2000) (CST #4967),

GABARAP (E1J4E) antibody (1:2000) (CST #13733), and HRP-conjugated secondary

antibodies (Jackson ImmunoResearch, West Grove, PA, USA). Flies were first transferred to

centrifuge tube containing 1x laemmli buffer (10 μl buffer for each milligram of fly), and heated

at 100 ℃ for 3 min. The samples were then homogenized and heated again at 100 ℃ for 3 min.

After the centrifuge at 13000x rpm for 5 min, supernatants were collected and loaded onto Mini-

PROTEAN precast gels (Bio-Rad, Hercules, CA, USA). Following incubation with primary and

secondary antibodies, the blots were visualized with Pierce ECL Western Blotting Substrate

(Thermo Fisher Scientific, Waltham, MA, USA).

Bafilomycin A1 and Chloroquine Treatment

For Bafilomycin A1 treatment, semi-intact hearts were incubated with 100 nM of

bafilomycin A1 (Alfa Aesar, Haverhill, MA, USA) in artificial hemolymph (buffer receipt in [31])

for two hours at room temperature prior to the appropriate immunostaining. DMSO was used as

a control.

For chloroquine treatments, 100 μl of 20 mM chloroquine diphosphate salt (Sigma-

Aldrich, St Louis, MO, USA) was added onto the fly food. Flies were fed with chloroquine for at

least 24 hours prior to the cardiac analysis or the western blots.

Demography and survival analysis

Newly enclosed female flies were allowed to mate for two days, then separated from

males and assigned to replicate one-liter demography cages at a density of 100-125 flies per

cage. Three independent cages were set-up per genotype. Food was changed every two days, at

which time dead flies were removed from the cage and counted. Survival analysis was conducted

with JMP statistical software (SAS Institute, Cary, NC, USA), and data from replicate cages

were combined. Mortality distributions were compared by Log-rank test.

Statistical analysis

GraphPad Prism (GraphPad Software, La Jolla, CA) was used for statistical analysis. To

compare the mean value of treatment groups versus that of control, either student t-test or one-

way ANOVA was performed using Dunnett's test for multiple comparison. The effects of

mutants during aging was analyzed by two-way ANOVA, including Tukey multiple comparisons

test. In SOHA analysis, the outliers were identified using Robust regression and Outlier removal

(ROUT) method (Q=1%) prior to the data analysis.

Acknowledgements

We thank Bloomington Drosophila Stock Center, Harvard Medical School, and Drosophila

Genomics Resource Center for fly stocks and cDNA clones. We thank Michael O'Connor for

activin reagents and fly stocks, Duojia Pan, Stocker Hugo, Jongkyeong Chung, Gábor Juhász,

Jun-yuan Ji, Bruce Edgar for fly stocks. Marc Tatar for help with Demography and survival

analysis. This work was supported by NIH/NIA R00 AG048016 and AFAR Research Grants to

HB.

Author Contributions Statement

Conceived and designed the experiments: KC RB KO HB. Performed the experiments: KC PK

YL KH APS IPN HB. Analyzed the data: KC PK YL KH ET APS IPN HB. Wrote the paper: KC

RB KO HB. All authors reviewed and approved the manuscript.

Competing Interests

The authors declare that no competing interest exists.

Figure Legends

Figure 1. Heart-specific knockdown of *Daw* slows cardiac aging.

a. Immunostaining of phospho-Smad2 in fly hearts at young (2 weeks) and old ages (6 weeks).

Arrows indicate cardiomyocyte nuclei and positive pSmad2 staining. Scale bar: 20 μm.

b. Representative M-mode traces (8 seconds) showing age-dependent movement of heart wall in

control and heart-specific *Daw* knockdown flies (*Hand-gal4>UAS-Daw^{RNAi}*).

c-g. Age-related changes in arrhythmia index, diastolic intervals, heart period, cardiac output,

and systolic intervals in control (*Ctrl*) and cardiac-specific *Daw* knockdown flies (*Daw^{RNAi}*).

Flies were cultured at 40% relative humidity. *Hand-gal4* driver was used to knock down gene

expression specifically in cardiac tissues (cardiomyocytes and pericardial cells). Results from

three independent *UAS-Daw^{RNAi}* lines are shown (RNAi #1: BDSC #34974, RNAi #2: VDRC

#105309, RNAi #3: BDSC #50911). Statistical significance is assessed by two-way ANOVA

followed by Tukey multiple comparisons test (* p<0.05, ** p<0.01, *** p<0.001, ns = not

significant). N=7~20. The interaction between genotype and age is statistically significant for

heart period (P = 0.0041) and diastolic interval (P = 0.0243).

h-j. Age-dependent changes in cardiac arrhythmia between control and *Daw* knockdown using

various tissue drivers, *Hand-gal4* (cardiomyocytes and pericardial cells), *TinC-gal4*

(cardiomyocytes), and *Dot-gal4* (pericardial cells). Student t-test (* p<0.05, ** p<0.01, ***

p<0.001). N=25~31.

k. Age-dependent changes in diastolic intervals between control and *Daw* knockdown in

pericardial cells (*Dot-gal4*). Student t-test (* p<0.05, ** p<0.01, *** p<0.001). N=13~26.

Figure 2. Cardiomyocyte-specific knockdown of activin receptor *Babo* delays cardiac aging.

a-d. Age-dependent changes in cardiac arrhythmia, diastolic intervals, heart period, and systolic

intervals in control (*Ctrl*) and cardiomyocyte-specific *Babo* knockdown flies (*Babo^{RNAi}*). Flies

were cultured at 40% relative humidity. *TinC-gal4* driver was used. Results from two

independent *UAS-Babo^{RNAi}* lines are shown (RNAi #1: BDSC #25933, RNAi #2: BDSC

#40866). N=15~30.

e-g. Changes in cardiac arrhythmia, diastolic intervals and heart period in fly hearts expressing

constitutively activated *Babo* (*Babo^{Act}*). GeneSwitch heart driver *Hand-GS-gal4* was used to

induce adult-onset *Babo* activation. RU: RU486 (Mifepristone). Flies were cultured at 40%

relative humidity. N=6~9. One-way ANOVA (*** p<0.001, ** p<0.01, * p<0.05, ns = not

significant).

Figure 3. Daw negatively regulates autophagy in fly hearts.

a. Representative images of LysoTracker staining of young adult fat body between wild-type

(*WT*) and heterozygous *Daw$^{[11]}$* mutants. Scale bar: 20 μm.

b. Western blot analysis on Atg8a lipidation between *WT* and *Daw$^{[11]}$* mutants. GABARAP

antibodies were used to detect Atg8a. The lower band represents lipidated Atg8a. Flies were fed

with chloroquine (CQ, 20 mM) for at least 24 hours prior to western blots.

c. Representative images of mosaic analysis on autophagosome staining in larval fat body. Fat

body clones were generated by crossing *Daw^{RNAi}* and *Babo^{RNAi}* lines into FLPout line carrying

mCherry-Atg8a reporter (*hs-flp; endogenous P-3x mCherry-Atg8a, UAS-GFP/Cyo;*

Act>CD2>Gal4,UAS-Dcr2). RNAi Clones are GFP-positive cells (dashed lines). Scale bar: 20

μm.

d. Representative images of fly hearts expressing autophagosome reporter mCherry-Atg8a at

both young and old ages. Arrows indicate cardiomyocyte nuclei and surrounding

autophagosomes. Abdominal segments A2-A3 are shown. Heart tube is located between two

yellow dashed lines. PC: Pericardial cell.

e-g. Representative images of mCherry-Atg8a reporter surrounding cardiomyocyte nuclei in

control, *Daw^{RNAi}*, and *Babo^{Act}* flies. Semi-intact hearts were incubated with 100 nM of

bafilomycin A1 (Baf A1) for two hours prior to immunostaining. Scale bar: 10 μm.

h. Quantification of age-dependent changes in mCherry-Atg8a-positive puncta in perinuclear

region in control, *Daw^{RNAi}*, and *Babo^{Act}* flies. *TinC-gal4* was used to drive the expression of

Atg8a reporter and gene knockdown. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns

= not significant). N=5~7.

i. Quantification of age-dependent changes in mCherry-Atg8a-positive puncta in perinuclear

region in control and *Daw^{RNAi}*. *Hand -gal4* was used to drive the expression of Atg8a reporter

and *Daw* knockdown. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not

significant). N=5~7.

j. Quantification of Baf A1-induced mCherry-Atg8a-positive puncta between control and

Daw^{RNAi}.

Figure 4. Inhibition of autophagy, but not activation of mTORC1, blocks *Daw* knockdown-

mediated cardioprotection.

a. Cardiac arrhythmia of chloroquine-treated (20 mM, 24 hours) 6-week-old control and *Daw*

knockdown flies (*Hand-gal4* used). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns =

not significant). N=14~38.

b. Cardiac arrhythmia and diastolic intervals in heart-specific knockdown of *Atg1* at young ages

(2-week-old) (*Hand-gal4* used). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not

significant). N=16~19.

c. Age-dependent changes in cardiac arrhythmia in control, *Daw^{RNAi}*, and *Daw^{RNAi}; Atg1^{RNAi}* flies

(*TinC-gal4* used). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant).

N=16~35.

d-e. Cardiac arrhythmia and diastolic intervals of *Tsc1* knockdown flies (*TinC-gal4* used). Two

independent Tsc1 RNAi lines were used (RNAi-1: BDSC #52931, RNAi-2: BDSC #54034).

One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=16~18.

f. Age-dependent changes in cardiac arrhythmia in control, *Daw[RNAi]*, and *Daw[RNAi]; Tsc1[RNAi]* flies

(*TinC-gal4* used). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant).

N=7~30.

g. Representative images of LysoTracker staining in adult fat body of *WT*, *Daw[111]*, *Tsc1[12]* and

double mutant *Daw[111]; Tsc1[12]*. Scale bar: 10 μm.

Figure 5. Daw genetically interacts with mTORC2/Rictor to control autophagy.

a. Representative images of phospho-AKT staining in cardiomyocytes of control and Daw

knockdown flies (*Hand-gal4*). Scale bar: 10 μm.

b. Image quantification of Figure 5a. Student t-test (* $p<0.05$, ** $p<0.01$, *** $p<0.001$). N=5.

c. Representative images of phospho-AKT staining in cardiomyocytes of control and Daw

knockdown flies (*Hand-gal4*). Scale bar: 10 μm.

d. Image quantification of Figure 5c. Student t-test (* $p<0.05$, ** $p<0.01$, *** $p<0.001$). N=5.

e. Representative images of phospho-AKT staining in cardiomyocytes of control and Rictor

over-expression (*Hand-gal4*). Two independent Rictor over-expression used (#1: *UAS-Rictor*,

#2: *UAS-HA-Rictor* [77]). Scale bar: 10 μm.

f. Image quantification of Figure 5e. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns =

not significant). N=5.

g. QRT-PCR analysis of *Rictor* expression in *Daw* knockdown flies (*Hand-gal4*). Student t-test

(* $p<0.05$, ** $p<0.01$, *** $p<0.001$). N=3.

h. QRT-PCR analysis of *Daw* expression in *Daw* knockdown flies (*Hand-gal4*). Student t-test (*

$p<0.05$, ** $p<0.01$, *** $p<0.001$). N=3.

i. Representative images of LysoTracker staining in adult fat body of *WT*, *Daw[111]*, *Rictor[42]* and

double mutant *Rictor[42]; Daw[111]*. Scale bar: 20 μm.

j. Image quantification of Figure 5i. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns =

not significant). N=5.

k. Representative images of Magic Red staining in adult fat body of *WT*, *Daw[111]*, *Rictor[42]* and

double mutant *Rictor[42]; Daw[111]*. Scale bar: 20 μm.

l. Image quantification of Figure 5k. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=5.

m. Representative images of mosaic analysis of LysoTracker staining in larval fat body. Fat body clones were generated by crossing *Rictor* overexpression lines into a FLPout line (*yw, hs-flp, UAS-CD8::GFP; Act>y+>Gal4,UAS-GFP.nls;UAS-Dcr2*). Clones with *Rictor* overexpression are GFP-positive cells (dashed lines). Scale bar: 20 μm.

Figure 6. Heart-specific overexpression of *Rictor* preserved cardiac function during aging.

a-b. Diastolic intervals and arrhythmia in flies with heart-specific overexpression of *Rictor* (*Hand-gal4*). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=19~26.

c. Cardiac arrhythmia in flies with heart-specific knockdown of *Rictor*. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=24.

d-e. Diastolic intervals and arrhythmia in flies with heart-specific expression of *Babo^{Act}*, or both *Babo^{Act}* and *Rictor* (*Hand-gal4*). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant). N=11~27.

f. Proposed model for activin-mTORC2 interaction in the regulation of autophagy and cardiac aging.

Figure 7. Cardiac-specific knockdown of *Daw* prolongs lifespan.

a-b. Survival analysis of cardiomyocyte-specific (*TinC-gal4*) knockdown of *Daw*. Two independent trials and multiple controls (*yw x yw, yw x Daw RNAi, TinC x mCherry RNAi*) were performed. Log-Rank test, $p<0.001$.

c-d. Survival analysis of heart-specific (*Hand-gal4*) knockdown of *Daw*. Two independent trials were performed. Log-Rank test, $p<0.001$.

e. Lifespan table to show sample size and median lifespan of the survival analysis.

Supplemental file

Table S1. Primer list

Figure S1. Validation of GABARAP antibody.

a. Multiple sequence alignment between *Drosophila* Atg8a and human GABARAP using

ClustalW (https://www.genome.jp/tools-bin/clustalw). The red box indicates the predicted

epitope sequences used in producing GABARAP antibody.

b. Representative images of GABARAP immunostaining in fly hearts of *WT* and *Atg8a*$^{\Delta 4}$

mutants. Heart tube is located between two yellow dashed lines. The panels on the right are the

zoomed-in images. Scale bar: 20 μm.

c. Representative images of GABARAP immunostaining in adult fat body of *WT* and *Atg8a*$^{\Delta 4}$

mutants. The panels on the right are the zoomed-in images. Scale bar: 20 μm.

d. Western blots testing GABARAP antibody using *Atg8a*$^{\Delta 4}$ mutants.

Figure S2. Daw inhibits autophagy flux in the heart.

a. Representative images of GABARAP immunostaining in fly hearts of *WT* and *Daw*[11] mutants

treated with or without Baf A1. Heart tube is located between two yellow dashed lines. The

panels on the right are the zoomed-in images. PC: Pericardial cell. Scale bar: 20 μm.

b. Quantification of Figure S2. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not

significant). N=5.

c. Quantification of Baf A1-induced puncta between *WT* and *Daw*[11] mutants.

Figure S3. Aging increases the accumulation of Ref(2)P in the heart.

a-b. Representative images of Ref(2)P in young and old hearts. Scale bar: 10 μm.

c. Representative images of Ref(2)P in the heart of *Atg8a*$^{\Delta 4}$ mutants. Scale bar: 10 μm.

Figure S4. Activation of mTORC1 alters heart period, but not arrhythmia.

a. Schematic diagram for PI3K/AKT/mTOR signaling pathway.

b-d. Heart period, diastolic intervals, and arrhythmia in control (*Ctrl*) and cardiomyocyte-

specific *Tsc1* knockdown flies (*TinC-gal4*). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, *

$p<0.05$, ns = not significant). N=14~18.

e-g. Heart period, diastolic intervals, and arrhythmia in control (*Ctrl*) and cardiomyocyte-specific

Reptor knockdown flies (*TinC-gal4*). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns

= not significant). N=13~28.

h-j. Heart period, diastolic intervals, and arrhythmia in control (*Ctrl*) and flies overexpressing

Rheb (*TinC-gal4*). One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns = not significant).

N=21~31.

Figure S5. Autophagy regulation by mTORC2.

a. Representative images of LysoTracker staining in adult fat body of *WT*, *Daw[11]*, *Sin1[e03756]*

and double mutant *Sin1[e03756]*; *Daw[11]*. Scale bar: 20 μm.

b. Image quantification of Figure S5a. One-way ANOVA (*** $p<0.001$, ** $p<0.01$, * $p<0.05$, ns

= not significant). N=5.

c. Representative images of LysoTracker staining in fed and starved WT, *Tsc1* mutants, and

Rictor mutants. Scale bar: 20 μm.

d. Representative images of mosaic analysis on LysoTracker staining in larval fat body of

Babo[Act] flies upon starvation. Larval fat body clones were generated using a FLPout line (*yw, hs-*

flp, UAS-CD8::GFP; Act>y+>Gal4,UAS-GFP.nls;UAS-Dcr2). Clones with *Babo[Ac]* expression

are GFP-positive cells (dashed lines). Scale bar: 20 μm.

References

1. Dai, D.F., Chen, T., Johnson, S.C., Szeto, H. & Rabinovitch, P.S. Cardiac aging: from
molecular mechanisms to significance in human health and disease. *Antioxidants & redox*
signaling **16**, 1492-1526 (2012).
2. North, B.J. & Sinclair, D.A. The intersection between aging and cardiovascular disease.
Circulation research **110**, 1097-1108 (2012).
3. Lakatta, E.G. & Levy, D. Arterial and cardiac aging: major shareholders in
cardiovascular disease enterprises: Part II: the aging heart in health: links to heart disease.
Circulation **107**, 346-354 (2003).
4. Dai, D.F. *et al.* Overexpression of catalase targeted to mitochondria attenuates murine
cardiac aging. *Circulation* **119**, 2789-2797 (2009).
5. Quarles, E.K. *et al.* Quality control systems in cardiac aging. *Ageing research reviews* **23**,
101-115 (2015).
6. Rubinsztein, D.C., Marino, G. & Kroemer, G. Autophagy and aging. *Cell* **146**, 682-695
(2011).
7. Shibutani, S.T. & Yoshimori, T. A current perspective of autophagosome biogenesis.
Cell research **24**, 58-68 (2014).
8. Demontis, F. & Perrimon, N. FOXO/4E-BP signaling in Drosophila muscles regulates
organism-wide proteostasis during aging. *Cell* **143**, 813-825 (2010).
9. Chang, J.T., Kumsta, C., Hellman, A.B., Adams, L.M. & Hansen, M. Spatiotemporal
regulation of autophagy during Caenorhabditis elegans aging. *eLife* **6** (2017).
10. Taneike, M. *et al.* Inhibition of autophagy in the heart induces age-related
cardiomyopathy. *Autophagy* **6**, 600-606 (2010).
11. Dutta, D., Xu, J., Kim, J.S., Dunn, W.A., Jr. & Leeuwenburgh, C. Upregulated autophagy
protects cardiomyocytes from oxidative stress-induced toxicity. *Autophagy* **9**, 328-344
(2013).
12. Wessells, R. *et al.* d4eBP acts downstream of both dTOR and dFoxo to modulate cardiac
functional aging in Drosophila. *Aging cell* **8**, 542-552 (2009).
13. Luong, N. *et al.* Activated FOXO-mediated insulin resistance is blocked by reduction of
TOR activity. *Cell metabolism* **4**, 133-142 (2006).
14. Birse, R.T. *et al.* High-fat-diet-induced obesity and heart dysfunction are regulated by the
TOR pathway in Drosophila. *Cell metabolism* **12**, 533-544 (2010).
15. Saxton, R.A. & Sabatini, D.M. mTOR Signaling in Growth, Metabolism, and Disease.
Cell **168**, 960-976 (2017).
16. Kaeberlein, M. *et al.* Regulation of yeast replicative life span by TOR and Sch9 in
response to nutrients. *Science* **310**, 1193-1196 (2005).
17. Lee, J.H. *et al.* Sestrin as a feedback inhibitor of TOR that prevents age-related
pathologies. *Science* **327**, 1223-1228 (2010).
18. Baar, E.L., Carbajal, K.A., Ong, I.M. & Lamming, D.W. Sex- and tissue-specific changes
in mTOR signaling with age in C57BL/6J mice. *Aging cell* **15**, 155-166 (2016).
19. Kroemer, G., Marino, G. & Levine, B. Autophagy and the integrated stress response.
Molecular cell **40**, 280-293 (2010).
20. He, C. & Klionsky, D.J. Regulation mechanisms and signaling pathways of autophagy.
Annual review of genetics **43**, 67-93 (2009).

21. Bai, H., Kang, P., Hernandez, A.M. & Tatar, M. Activin signaling targeted by
insulin/dFOXO regulates aging and muscle proteostasis in Drosophila. *PLoS genetics* **9**,
e1003941 (2013).
22. Demontis, F., Patel, V.K., Swindell, W.R. & Perrimon, N. Intertissue control of the
nucleolus via a myokine-dependent longevity pathway. *Cell reports* **7**, 1481-1494 (2014).
23. Upadhyay, A., Moss-Taylor, L., Kim, M.J., Ghosh, A.C. & O'Connor, M.B. TGF-beta
Family Signaling in Drosophila. *Cold Spring Harbor perspectives in biology* (2017).
24. Upadhyay, A., Moss-Taylor, L., Kim, M.J., Ghosh, A.C. & O'Connor, M.B. TGF-beta
Family Signaling in Drosophila. *Cold Spring Harbor perspectives in biology* **9** (2017).
25. Biernacka, A. & Frangogiannis, N.G. Aging and Cardiac Fibrosis. *Aging and disease* **2**,
158-173 (2011).
26. Li, C.Y. *et al.* Partial inhibition of activin receptor-like kinase 4 attenuates pressure
overload-induced cardiac fibrosis and improves cardiac function. *Journal of hypertension*
34, 1766-1777 (2016).
27. Chen, Y.H. *et al.* Haplodeficiency of activin receptor-like kinase 4 alleviates myocardial
infarction-induced cardiac fibrosis and preserves cardiac function. *Journal of molecular*
and cellular cardiology **105**, 1-11 (2017).
28. Tan, S.M., Zhang, Y., Connelly, K.A., Gilbert, R.E. & Kelly, D.J. Targeted inhibition of
activin receptor-like kinase 5 signaling attenuates cardiac dysfunction following
myocardial infarction. *American journal of physiology. Heart and circulatory physiology*
298, H1415-1425 (2010).
29. Oshima, Y. *et al.* Activin A and follistatin-like 3 determine the susceptibility of heart to
ischemic injury. *Circulation* **120**, 1606-1615 (2009).
30. Ocorr, K., Vogler, G. & Bodmer, R. Methods to assess Drosophila heart development,
function and aging. *Methods* **68**, 265-272 (2014).
31. Ocorr, K., Fink, M., Cammarato, A., Bernstein, S. & Bodmer, R. Semi-automated Optical
Heartbeat Analysis of small hearts. *Journal of visualized experiments : JoVE* (2009).
32. Han, Z. & Olson, E.N. Hand is a direct target of Tinman and GATA factors during
Drosophila cardiogenesis and hematopoiesis. *Development* **132**, 3525-3536 (2005).
33. Lo, P.C. & Frasch, M. A role for the COUP-TF-related gene seven-up in the
diversification of cardioblast identities in the dorsal vessel of Drosophila. *Mechanisms of*
development **104**, 49-60 (2001).
34. Zhang, F., Zhao, Y. & Han, Z. An in vivo functional analysis system for renal gene
discovery in Drosophila pericardial nephrocytes. *Journal of the American Society of*
Nephrology : JASN **24**, 191-197 (2013).
35. Scott, R.C., Schuldiner, O. & Neufeld, T.P. Role and regulation of starvation-induced
autophagy in the Drosophila fat body. *Developmental cell* **7**, 167-178 (2004).
36. Klionsky, D.J. *et al.* Guidelines for the use and interpretation of assays for monitoring
autophagy (3rd edition). *Autophagy* **12**, 1-222 (2016).
37. Mauvezin, C., Nagy, P., Juhasz, G. & Neufeld, T.P. Autophagosome-lysosome fusion is
independent of V-ATPase-mediated acidification. *Nature communications* **6**, 7007
(2015).
38. Nezis, I.P. *et al.* Ref(2)P, the Drosophila melanogaster homologue of mammalian p62, is
required for the formation of protein aggregates in adult brain. *The Journal of cell*
biology **180**, 1065-1071 (2008).

39.   Johnson, S.C., Rabinovitch, P.S. & Kaeberlein, M. mTOR is a key modulator of ageing
864   and age-related disease. *Nature* **493**, 338-345 (2013).
865   40.   Shioi, T. *et al.* Rapamycin attenuates load-induced cardiac hypertrophy in mice.
866   *Circulation* **107**, 1664-1670 (2003).
867   41.   Ding, Y. *et al.* Haploinsufficiency of target of rapamycin attenuates cardiomyopathies in
868   adult zebrafish. *Circulation research* **109**, 658-669 (2011).
869   42.   Tiebe, M. *et al.* REPTOR and REPTOR-BP Regulate Organismal Metabolism and
870   Transcription Downstream of TORC1. *Developmental cell* **33**, 272-284 (2015).
871   43.   Sarbassov, D.D., Guertin, D.A., Ali, S.M. & Sabatini, D.M. Phosphorylation and
872   regulation of Akt/PKB by the rictor-mTOR complex. *Science* **307**, 1098-1101 (2005).
873   44.   Hwangbo, D.S., Gershman, B., Tu, M.P., Palmer, M. & Tatar, M. Drosophila dFOXO
874   controls lifespan and regulates insulin signalling in brain and fat body. *Nature* **429**, 562-
875   566 (2004).
876   45.   Zhang, G. *et al.* Hypothalamic programming of systemic ageing involving IKK-beta, NF-
877   kappaB and GnRH. *Nature* **497**, 211-216 (2013).
878   46.   Akhurst, R.J. & Hata, A. Targeting the TGFbeta signalling pathway in disease. *Nature*
879   *reviews. Drug discovery* **11**, 790-811 (2012).
880   47.   Tsuchida, K. *et al.* Activin signaling as an emerging target for therapeutic interventions.
881   *Cell communication and signaling : CCS* **7**, 15 (2009).
882   48.   Luo, S., Kleemann, G.A., Ashraf, J.M., Shaw, W.M. & Murphy, C.T. TGF-beta and
883   insulin signaling regulate reproductive aging via oocyte and germline quality
884   maintenance. *Cell* **143**, 299-312 (2010).
885   49.   Narasimhan, S.D. *et al.* PDP-1 links the TGF-beta and IIS pathways to regulate
886   longevity, development, and metabolism. *PLoS genetics* **7**, e1001377 (2011).
887   50.   Shaw, W.M., Luo, S., Landis, J., Ashraf, J. & Murphy, C.T. The C. elegans TGF-beta
888   Dauer pathway regulates longevity via insulin signaling. *Current biology : CB* **17**, 1635-
889   1645 (2007).
890   51.   Baccarelli, A. *et al.* Activin A serum levels and aging of the pituitary-gonadal axis: a
891   cross-sectional study in middle-aged and elderly healthy subjects. *Experimental*
892   *gerontology* **36**, 1403-1412 (2001).
893   52.   Yndestad, A. *et al.* Elevated levels of activin A in heart failure: potential role in
894   myocardial remodeling. *Circulation* **109**, 1379-1385 (2004).
895   53.   Wei, Q. *et al.* The expression and role of activin A and follistatin in heart failure rats after
896   myocardial infarction. *International journal of cardiology* **168**, 2994-2997 (2013).
897   54.   Tsai, Y.L. *et al.* The Association Between Serum Activin A Levels and Hypertension in
898   the Elderly: A Cross-Sectional Analysis From I-Lan Longitudinal Aging Study.
899   *American journal of hypertension* (2017).
900   55.   Chen, Y. *et al.* Regulation and actions of activin A and follistatin in myocardial
901   ischaemia-reperfusion injury. *Cytokine* **69**, 255-262 (2014).
902   56.   Liu, M., Mao, C., Li, J., Han, F. & Yang, P. Effects of the Activin A-Follistatin System
903   on Myocardial Cell Apoptosis through the Endoplasmic Reticulum Stress Pathway in
904   Heart Failure. *International journal of molecular sciences* **18** (2017).
905   57.   Ghosh, A.C. & O'Connor, M.B. Systemic Activin signaling independently regulates
906   sugar homeostasis, cellular metabolism, and pH balance in Drosophila melanogaster.
907   *Proceedings of the National Academy of Sciences of the United States of America* **111**,
908   5729-5734 (2014).

909   58.   Chng, W.B., Bou Sleiman, M.S., Schupfer, F. & Lemaitre, B. Transforming growth
910   factor beta/activin signaling functions as a sugar-sensing feedback loop to regulate
911   digestive enzyme expression. *Cell reports* **9**, 336-348 (2014).
912   59.   Werner, S. & Alzheimer, C. Roles of activin in tissue repair, fibrosis, and inflammatory
913   disease. *Cytokine & growth factor reviews* **17**, 157-171 (2006).
914   60.   Xue, L.X. *et al.* Activin A/Smads signaling pathway negatively regulates Oxygen
915   Glucose Deprivation-induced autophagy via suppression of JNK and p38 MAPK
916   pathways in neuronal PC12 cells. *Biochemical and biophysical research communications*
917   **480**, 355-361 (2016).
918   61.   Guo, B. *et al.* Genome-wide screen identifies signaling pathways that regulate autophagy
919   during Caenorhabditis elegans development. *EMBO reports* **15**, 705-713 (2014).
920   62.   Florez-McClure, M.L. & Link, C.D. Erratum: "Florez-McClure ML, Hohsfield LA, Fonte
921   G, Bealor MT, Link CD. Decreased insulin-receptor signaling promotes the autophagic
922   degradation of beta-amyloid peptide in C. elegans". Autophagy 2007; 3:569-80.
923   *Autophagy* **5**, 132-133 (2009).
924   63.   Florez-McClure, M.L., Hohsfield, L.A., Fonte, G., Bealor, M.T. & Link, C.D. Decreased
925   insulin-receptor signaling promotes the autophagic degradation of beta-amyloid peptide
926   in C. elegans. *Autophagy* **3**, 569-580 (2007).
927   64.   Lipinski, M.M. *et al.* A genome-wide siRNA screen reveals multiple mTORC1
928   independent signaling pathways regulating autophagy under normal nutritional
929   conditions. *Developmental cell* **18**, 1041-1052 (2010).
930   65.   Li, Y. *et al.* Protein kinase C controls lysosome biogenesis independently of mTORC1.
931   *Nature cell biology* **18**, 1065-1077 (2016).
932   66.   Henson, S.M. *et al.* p38 signaling inhibits mTORC1-independent autophagy in senescent
933   human CD8(+) T cells. *The Journal of clinical investigation* **124**, 4004-4016 (2014).
934   67.   Oh, W.J. & Jacinto, E. mTOR complex 2 signaling and functions. *Cell cycle* **10**, 2305-
935   2316 (2011).
936   68.   Gaubitz, C., Prouteau, M., Kusmider, B. & Loewith, R. TORC2 Structure and Function.
937   *Trends in biochemical sciences* **41**, 532-545 (2016).
938   69.   Gurusamy, N. *et al.* Cardioprotection by resveratrol: a novel mechanism via autophagy
939   involving the mTORC2 pathway. *Cardiovascular research* **86**, 103-112 (2010).
940   70.   Li, J. *et al.* Rictor/mTORC2 protects against cisplatin-induced tubular cell death and
941   acute kidney injury. *Kidney international* **86**, 86-102 (2014).
942   71.   Mammucari, C. *et al.* FoxO3 controls autophagy in skeletal muscle in vivo. *Cell*
943   *metabolism* **6**, 458-471 (2007).
944   72.   Xiong, Y. *et al.* ARG2 impairs endothelial autophagy through regulation of MTOR and
945   PRKAA/AMPK signaling in advanced atherosclerosis. *Autophagy* **10**, 2223-2238 (2014).
946   73.   Shende, P. *et al.* Cardiac mTOR complex 2 preserves ventricular function in pressure-
947   overload hypertrophy. *Cardiovascular research* **109**, 103-114 (2016).
948   74.   Sciarretta, S. *et al.* mTORC2 regulates cardiac response to stress by inhibiting MST1.
949   *Cell reports* **11**, 125-136 (2015).
950   75.   Soukas, A.A., Kane, E.A., Carr, C.E., Melo, J.A. & Ruvkun, G. Rictor/TORC2 regulates
951   fat metabolism, feeding, growth, and life span in Caenorhabditis elegans. *Genes &*
952   *development* **23**, 496-511 (2009).
953   76.   Lamming, D.W. *et al.* Depletion of Rictor, an essential protein component of mTORC2,
954   decreases male lifespan. *Aging cell* **13**, 911-917 (2014).

77.   Jevtov, I. *et al.* TORC2 mediates the heat stress response in Drosophila by promoting the
formation of stress granules. *Journal of cell science* **128**, 2497-2508 (2015).
78.   Wu, Z. *et al.* Tricornered/NDR kinase signaling mediates PINK1-directed mitochondrial
quality control and tissue maintenance. *Genes & development* **27**, 157-162 (2013).
79.   Chen, C.C. *et al.* FoxOs inhibit mTORC1 and activate Akt by inducing the expression of
Sestrin3 and Rictor. *Developmental cell* **18**, 592-604 (2010).
80.   Brummel, T. *et al.* The Drosophila activin receptor baboon signals through dSmad2 and
controls cell proliferation but not patterning during larval development. *Genes &*
development **13**, 98-111 (1999).
81.   Serpe, M. & O'Connor, M.B. The metalloprotease tolloid-related and its TGF-beta-like
substrate Dawdle regulate Drosophila motoneuron axon guidance. *Development* **133**,
4969-4979 (2006).
82.   Gao, X. & Pan, D. TSC1 and TSC2 tumor suppressors antagonize insulin signaling in cell
growth. *Genes & development* **15**, 1383-1392 (2001).
83.   Lee, G. & Chung, J. Discrete functions of rictor and raptor in cell growth regulation in
Drosophila. *Biochemical and biophysical research communications* **357**, 1154-1159
(2007).
84.   Monnier, V. *et al.* dJun and Vri/dNFIL3 are major regulators of cardiac aging in
Drosophila. *PLoS genetics* **8**, e1003081 (2012).
85.   Pircs, K. *et al.* Advantages and limitations of different p62-based assays for estimating
autophagic activity in Drosophila. *PloS one* **7**, e44214 (2012).
86.   Fink, M. *et al.* A new method for detection and quantification of heartbeat parameters in
Drosophila, zebrafish, and embryonic mouse hearts. *BioTechniques* **46**, 101-113 (2009).
87.   Ratliff, E.P., Barekat, A., Lipinski, M.M. & Finley, K.D. Brain trauma and autophagy:
What flies and mice can teach us about conserved responses. *Autophagy* **12**, 2256-2257
(2016).
88.   Ratliff, E.P. *et al.* Assessing Basal and Acute Autophagic Responses in the Adult
Drosophila Nervous System: The Impact of Gender, Genetics and Diet on Endogenous
Pathway Profiles. *PloS one* **11**, e0164239 (2016).
89.   Barekat, A. *et al.* Using Drosophila as an integrated model to study mild repetitive
traumatic brain injury. *Scientific reports* **6**, 25252 (2016).
90.   Kim, M. *et al.* Drosophila Gyf/GRB10 interacting GYF protein is an autophagy regulator
that controls neuron and muscle homeostasis. *Autophagy* **11**, 1358-1372 (2015).
988
989

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Genotype	Sample size	Mean lifespan (days)	Median lifespan (days)	Difference RNAi vs. Control	Prob. (Log-Rank)
Trial #1					
TinC>mCherry^{RNAi}	355	39.1	38	/	/
TinC>Daw^{RNAi-1}	389	44.3	46	21%	<0.001
yw/Daw^{RNAi-1}	301	45.2	44	/	/
Hand>Daw^{RNAi-1}	355	52.3	52	18%	<0.001
Trial #2					
TinC>mCherry^{RNAi}	364	45.3	44	/	/
TinC>Daw^{RNAi-1}	340	58	58	32%	<0.001
Hand>mCherry^{RNAi}	322	55.2	56	/	/
Hand>Daw^{RNAi-1}	315	63.3	64	14%	<0.001

Supplementary file
Supplemental Figure S1

a

```
DmAtg8a    MKFQYKEEHAFEKRRAEGDKIRRKYPDRVPVIVEKAPKARIGDLDKKKYLVPSDLTVGQF
hGABARAP   MKFVYKEEHPFEKRRSEGEKIRKKYPDRVPVIVEKAPKARIGDLDKKKYLVPSDLTVGQF
           ***  ***** :*****:**:***:********************** *************

DmAtg8a    YFLIRKRIHLRPEDALFFFVNNVIPPTSATMGSLYQEHHEEDYFLYIAYSDENVYGMAKI
hGABARAP   YFLIRKRIHLRAEDALFFFVNNVIPPTSATMGQLYQEHHEEDFFLYIAYSDESVYGL---
           ***********.**********************.*******:*********.***:

DmAtg8a    N
hGABARAP   -
```

b

c

d

Supplementary file
Supplemental Figure S2

Supplementary file
Supplemental Figure S3

a **WT (Young)**

b **WT (Old)**

c

Supplementary file
Supplemental Figure S5

Supplementary Table S1.

Primer name	Primer sequence (5'-3')	Symbol	FlyBase ID
RpL32-f	AAGAAGCGCACCAAGCACTTCATC	RpL32	FBgn0002626
RpL32-r	TCTGTTGTCGATACCCTTGGGCTT	RpL32	FBgn0002626
Daw-f	TGAGCCACCTCATCCAAATCACCT	Daw	FBgn0031461
Daw-r	TCGATCACGATGAATGGCCGGTAA	Daw	FBgn0031461
Rictor-f	CCCAACTGGCTAGCGTTTAT	Rictor	FBgn0031006
Rictor-r	GGCCTTTGCTTTCGTTTGAG	Rictor	FBgn0031006