1   **Title: Consequences of NMDA receptor deficiency can be rescued in the adult**
2   **brain**
3

4   **Authors:** Catharine A. Mielnik[1], Mary A. Binko[2], Adam J. Funk[3], Emily M. Johansson[4],
5   Katheron Intson[1], Nirun Sivananthan[1], Yuxiao Chen[1], Rehnuma Islam[2], Marija
6   Milenkovic[1], Wendy Horsfall[1], Ruth A. Ross[1], Shreejoy Tripathy[5], Laurent Groc[3], Ali
7   Salahpour[1], Robert E. McCullumsmith[4], Evelyn K. Lambe[2], and Amy J. Ramsey[1,2]*

8
9   **Affiliations:**
10
11 [1] Department of Pharmacology & Toxicology, University of Toronto, ON, M5S 1A8,
12 Canada
13
14 [2] Department of Physiology, University of Toronto, Toronto, ON, M5S 1A8, Canada
15
16 [3]Department of Neurosciences, University of Toledo, Toledo, OH 43614 USA
17
18 [4]Interdisciplinary Institute for NeuroScience (IINS) CNRS, Université Bordeaux Segalen,
19 33000 Bordeaux, France
20
21 [5]Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health,
22 Department of Psychiatry, University of Toronto, Toronto, ON M5T 1L8, Canada
23
*Correspondance to:
Amy Ramsey
1 King's College Circle
Room 4302 Medical Sciences Building
Toronto, ON M5S 1A8 Canada
(416) 978-2509
a.ramsey@utoronto.ca

31
32
33
34

Abstract: N-methyl-D-aspartate receptors (NMDARs) are required to shape activity-dependent connections in the developing and adult brain. Impaired NMDAR signaling through genetic or environmental insults causes a constellation of neurodevelopmental disorders that manifest as intellectual disability, epilepsy, autism, or schizophrenia. It is not clear whether the developmental impacts of NMDAR dysfunction can be overcome by interventions in adulthood. This question is paramount for neurodevelopmental disorders arising from mutations that occur in the *GRIN* genes, which encode NMDAR subunits, and the broader set of mutations that disrupt NMDAR function. We developed a mouse model where a congenital loss-of-function allele of *Grin1* is restored to wildtype by gene editing with Cre recombinase. Rescue of NMDARs in adult mice yields surprisingly robust improvements in cognitive behaviors, including those that are refractory to treatment with current medications. These results suggest that neurodevelopmental disorders arising from NMDAR deficiency can be effectively treated in adults.

INTRODUCTION

Greater than 1% of children are born with a neurodevelopmental disorder [1], including diagnoses of autism spectrum disorder and pervasive developmental delay [2]. Until recently, most children with global developmental delay were not given a more specific diagnosis that could predict treatment or long-term prognosis. Whole exome sequencing has revolutionized diagnostic assessment and has identified hundreds of genes that can cause intellectual disability and developmental delay through transmitted and *de novo* variants [3].

Through whole-exome sequencing, a new syndrome called *GRIN* encephalopathy has been identified that is caused by mutations in one of the seven *GRIN* genes that encode subunits for NMDA-type glutamate receptors (NMDARs). Deleterious missense and nonsense variants in *GRIN1*, *GRIN2A-D*, and *GRIN3A-B* cause encephalopathies that are often first diagnosed as intellectual disability, global developmental delay, epilepsy, autism, and/or schizophrenia [4]. The variants are often *de novo* heterozygous mutations that act as dominant negatives to reduce NMDAR function, although some variants lead to a gain-of function by altered channel gating properties [4]. Regardless of the nature of the mutation, patients with these deleterious variants have a similar syndrome of intellectual disability, and additional symptoms such as epilepsy, autism, cortical visual impairment, and movement disorders [4].

The identification of pathogenic variants in a *GRIN* gene allows for target-directed pharmacological treatments where approved drugs are available, but gene editing may ultimately be the most effective method to treat neurodevelopmental disorders. The timing of intervention remains a question for the future application of gene editing towards neurodevelopmental disorders. It has been assumed that intervention should occur as early in development as is medically feasible, that waiting risks irremediable damage, and that adults with these conditions are beyond the reach of medical treatment to improve cognitive function. However, these assumptions have not been stringently tested. Currently, there are many adults with these disorders that might also

benefit from gene therapy, and it is unknown whether the developmental consequences

of disease-causing variants can be reversed in adulthood.

The ability to reverse developmental insults is likely to depend on the nature of the

insult. For example, while adult rescue of Rett Syndrome gene *Mecp2* in mice reversed

several phenotypes [5], adult rescue of *Shank3* in mice showed a more selective

improvement to social behaviours [6]. Thus, it is conceivable that developmental insults

to the NMDAR system cannot be overcome with adult intervention, considering the

central role of this receptor. Indeed, NMDARs are required for the proper connectivity of

developing sensory circuits in the thalamus and cortex [7–9], for the establishment of

both inhibitory [10] and excitatory [11] synapses, and for the patterning of neuron

dendritic arborizations [12].

Since there is strong evidence that NMDARs participate in many aspects of

neurodevelopment, we asked whether developmental consequences of NMDAR

deficiency could be reversed in adult mice. Specifically, we asked whether adult

intervention could improve cognitive behaviours mediated by the cortex, since

intellectual disability is a core symptom of patients with *GRIN1* encephalopathy. *GRIN1*

encodes the essential subunit GluN1 that is present in all NMDARs, and null mutations

of *GRIN1* are lethal in humans [13] and in mice [7, 14]. Considering the well-established

role of these receptors in development and synapse refinement, it would be predicted

that NMDAR deficiencies caused by *GRIN1* mutations, in particular, would be refractory

to adult intervention.

Grin1 expression was restored in adulthood and molecular analysis, cellular function

and cognitive behaviours were quantified as outputs to measure the ability to reverse

intellectual disability. Strikingly, we discovered that plasticity at the cellular, synaptic,

and behavioural level was evident in the cortex. Furthermore, this rescue of cognitive

ability was reproduced in a separate adult cohort, and was maintained over a longer

recovery time. This study suggests that plasticity of cognitive circuits extends well into

adulthood, and that there is an inherent ability to upregulate NMDAR activity and to

normalize cognitive outputs.

111
112

RESULTS

Generation of mice with a reversible *Grin1* deficiency

To directly answer whether the developmental consequences of NMDAR deficiency

could be rescued in adults, we generated mice with a reversible hypomorphic mutation

in *Grin1*, the essential subunit of all NMDARs. Our previous studies with a similar

mouse line showed that a 90% knockdown of functional NMDARs is achieved through

the targeted insertion of a *neo* cassette in an intron of the *Grin1* gene [15]. In the new

mouse line, we added *lox*P sites flanking the *neo* cassette to allow for inducible excision

of the mutation, so that Cre recombinase could restore the locus to wildtype in a

conditional manner (Fig. 1A,B).

We intercrossed these mice with Rosa26-CreERT2 mice that ubiquitously express a

tamoxifen-inducible Cre recombinase. We first identified the tamoxifen regimen that

comprehensively induced Cre activity throughout the brain using a Cre reporter line

(Rosa26-dTomato: Supplementary Fig. 1). We then administered tamoxifen to all

genotypes of mice at postnatal day (PD) 70 and measured biochemical and behavioural

endpoints at PD98 (two-week treatment, two-week recovery). Four genotypes of mice

were studied: $Grin1^{+/+}$ (WT), $Grin1^{+/+}$:CreTg (WTCre), $Grin1^{flneo/flneo}$ ($Grin1^{KD}$), and

$Grin1^{flneo/flneo}$:CreTg ($Grin1^{RESCUE}$). We determined that WT and WTCre mice had similar

biochemical and behavioural phenotypes in all of the subsequent studies

(Supplementary Fig. 2), and thus experimental results for WT, $Grin1^{KD}$ and $Grin1^{RESCUE}$

mice were compared. Studies were performed with both male and female mice of equal

number and powered to study the effect of sex.

We determined the extent of molecular recovery of *Grin1* mRNA and GluN1 protein by

fluorescence *in situ* hybridization and immunofluoresence (Fig. 1C), and the regional

levels of NMDAR by [^{3}H]MK-801 radioligand binding (Supplementary Fig. 3). Notable for

our experimental objective, we observed substantial rescue of *Grin1* mRNA and

NMDAR protein complex in the prefrontal cortex (PFC) of $Grin1^{RESCUE}$ mice (Fig. 1C and

Supplemental Fig. 3), affording the ability to test whether cortical functions could

recover from developmental NMDAR deficiency.

We also asked whether recovery of *Grin1* expression was achieved in both

glutamatergic and GABAergic neurons of the cortex (Fig. 2A,B). In WT mice, the levels

of *Grin1* are similar between *Gad1*+ GABAergic and *Vglut1*+ glutamatergic cells. This is

consistent with single cell transcriptomics data reference atlases [16], which indicate

that *Grin1* is normally expressed in both cell types in the adult cortex, with higher levels

observed in *Vglut1*+ cells (Fig. 2C). Interestingly, *Grin1^{KD}* mice have less detectable

Grin1 mRNA in *Gad1*+ cells than *Vglut1*+ cells (Fig 2A,B). Furthermore, while *Grin1*

mRNA was substantially increased in all *Vglut1*+ cortical neurons in the *Grin1^{RESCUE}*

mice, the extent of recovery was less consistent in the Gad1+ neurons of adjacent

slices.

NMDAR currents and synaptic GluN1 protein are restored in mPFC neurons

The extent of functional NMDAR recovery in the cortex was determined through whole

cell electrophysiological recordings from brain slices. Physiological recordings from

layer V pyramidal neurons of medial PFC (mPFC) were performed (Fig. 3A). Bath

applied NMDA elicited an inward current in wildtype cells. This current was greatly

attenuated in *Grin1^{KD}* mice; however, in *Grin1^{RESCUE}* mice NMDA-elicited current was

restored to wildtype levels (Fig. 3B,C). The differences in functional NMDARs occurred

in the presence of largely similar intrinsic membrane properties (Supplementary Table

1). However, capacitance was significantly larger in prefrontal neurons of *Grin1^{RESCUE}*

compared to WT mice (Fig. 3D). Accordingly, we analyzed the current density of the

NMDA-elicited currents (Fig. 3E; effect of genotype, $F_{2,95}$=3.6, p=0.03) and found that

Grin1^{RESCUE} mice also had greatly increased current density compared to *Grin1^{KD}* mice.

As further demonstration of synaptic NMDAR recovery in the PFC, the levels of GluN1

protein were determined by immunoprecipitation with anti-PSD-95 antibody and mass

spectrometry. This procedure isolates the proteins that are part of the PSD-95 post-

synaptic protein complex, and, specifically, the amount of GluN1 peptide IVNIGAVLSTR

was determined relative to the intensity of PSD-95 peptides, to indicate the abundance

of GluN1 protein at PFC synapses. As shown in Fig. 3F, GluN1 peptide throughout the

PFC was reduced in *Grin1^{KD}* mice and restored to wildtype levels in *Grin1^{RESCUE}* mice

(WT: 0.79±0.06, *Grin1^{RESCUE}*: 0.67±0.02, p=0.077, power=1.00). Collectively, these

results indicate that we achieved rescue of NMDAR deficiencies in glutamatergic

neurons of the PFC in order to test the ability of adult intervention to improve cognitive

behaviours.

Cognitive impairments are rescued by adult intervention

We studied several domains of cognition that are used as endophenotypes for autism

related neurodevelopmental disorders: habituation to a novel environment, sensory

processing of acoustic startle, executive function, and social interaction. Although each

of these behaviours relies on more than cortical function for their performance, previous

studies have repeatedly demonstrated the critical role that the PFC plays in these

cognitive tasks. Indeed, cell-selective knockout of NMDARs in cortical neurons is

sufficient to impair habituation, sensory processing of acoustic startle, and social

interaction [17–19].

Habituation to a novel environment requires the cortical and hippocampal processes of

working and spatial memory to reduce exploration activity after a period of time [20, 21].

Habituation was quantified by calculating the habituation index (H.I.), which is the time

required to reach half of the maximal locomotor activity using linear regression. $Grin1^{KD}$

mice showed initial hyperactivity relative to WT in the first 10 minutes of exploration, and

120 minutes later, these mutant mice continued to explore the arena with high levels of

activity (extrapolated H.I.: 149.1±15.6 min; Fig. 4A). In contrast, while $Grin1^{RESCUE}$

rescue mice also showed initial hyperactivity, their habituation to the novel environment

was similar to WT mice (H.I.: WT 53.5±1.2 min, $Grin1^{RESCUE}$ 64.4±3.2 min; Fig. 4A,

p>0.99, power=1.00).

Sensorimotor gating, which is modulated by cortical arousal circuits [22], was measured

with the paradigm of pre-pulse inhibition (PPI) of acoustic startle response (ASR).

Consistent with studies of the original knockdown mutation [23], $Grin1^{KD}$ mice exhibited

deficits in sensorimotor gating at pre-pulse intensities of 4, 8, and 16 dB (Fig. 4B). This

indicated that the pre-cognitive ability to attenuate motor response to a startling sound

was impaired. $Grin1^{RESCUE}$ mice showed a complete restoration of sensory processing

in this test, with PPI levels that were similar to WT littermates (Fig. 4B, p>0.99,

power>0.99).

Executive function was tested in the puzzle box test, which measured the ability of the

mouse to overcome increasingly challenging obstacles and reach a goal box. Mice were

first introduced to the arena with an open doorway leading to the goal, but on

subsequent tests the doorway was blocked, and mice had to use an underpass or dig

through bedding to reach the goal. Thus, the test measured goal-directed behaviour and

cognitive flexibility to respond to different challenges [24]. *Grin1*^{KD} mice routinely failed

the most challenging task of digging through bedding and performed markedly worse

than WT in all trials (Fig. 4C). Impressively, *Grin1*^{RESCUE} mice solved both challenges,

and performed significantly better than *Grin1*^{KD} mice on all trials (Fig. 4C). Indeed, in

three of the seven trials, *Grin1*^{RESCUE} mice performed similar to WT mice. Thus, there

were substantial improvements in executive function as assessed in the puzzle box test.

Lastly, affiliative social behaviour was studied by measuring the amount of time that a

mouse spent investigating a novel C57Bl/6J mouse. The novel mouse was constrained

in one area with a wire cage, and an empty cage was included in the arena to control for

non-social investigation of the cage. As expected, social interaction was significantly

impaired in *Grin1*^{KD} mice relative to WT (Fig. 4D). *Grin1*^{RESCUE} mice displayed social

interaction that was completely restored to WT levels (p>0.99, power=1.00).

Furthermore, *Grin1*^{KD} mutant mice showed excessive interest in exploring the arena and

the empty cage as represented by the activity trace in Fig. 4D, and the time spent per

visit with the novel mouse was substantially reduced compared to WT (Fig. 4D). Not

only was the amount of time spent in social interaction normalized in *Grin1*^{RESCUE} mice,

but the quality of social interaction appeared to improve, as demonstrated by the longer

time spent with each visit to the novel mouse (Fig. 4D).

We observed similar patterns of behavioural deficits in male and female *Grin1*^{KD} mice,

and similar patterns of recovery in *Grin1*^{RESCUE} male and female mice in most tests.

There were notable sex differences in a select behavioural test, the puzzle box test,

where female mice performed better than male mice irrespective of genotype

(Supplementary Fig. 4). It should also be noted that, in the puzzle box test, the female

Grin1^{RESCUE} mice showed a greater improvement than male *Grin1*^{RESCUE} mice.

Cognitive improvements persist with a longer recovery period

Finally, we asked whether these behavioural improvements would persist or would

further improve with a longer recovery period. In a distinct cohort of experimental and

control mice, we induced Cre-mediated rescue of *Grin1* at PD70, as in the original

paradigm, but waited an additional month before testing the animals (six-week recovery

vs. original two-week recovery). In this cohort, there was a similar degree of rescue in

the level of NMDARs (Supplementary Table 2). Behaviourally, the *Grin1*^{RESCUE} mice

showed significant improvement across all measures examined (Fig. 5), in a pattern

consistent with the assessments presented in Fig. 4. *Grin1*^{RESCUE} mice, treated at PD70,

and allowed to recover for six weeks, showed improvement in their habituation index

(HI), sensorimotor gating (PPI), and affiliative social behaviour deficits (AS), that was

similar to WT (HI: $p=0.17$, power=1.00; PPI: (4dB) $p=0.072$, power=1.00, (8dB) $p=1.00$,

power=0.99, (16dB) $p=1.00$, power=0.94; AS: $p>0.99$, power=0.99). This experiment

indicates that adult intervention leads to sustained cognitive improvement that can be

observed as early as two weeks after completion of treatment.

In summary, our data collectively show that the cortical restoration of *Grin1* expression

in adulthood leads to significant and persisting reversals in cognitive impairments.

DISCUSSION

The knockdown of *Grin1* results in viable mutant mice with deficits in cognitive behaviours that parallel the symptoms of *GRIN1* encephalopathy [13]. We did not find evidence of any deleterious effects from the postnatal upregulation of NMDARs. *Grin1*^{RESCUE} mice appeared healthier and were less reactive to handling after Cre induction. We found improvements in every aspect of behaviour that we examined.

Our strategy to achieve temporal rescue of NMDARs took advantage of a tamoxifen-inducible Cre recombinase [25]. The study design allowed us to treat all groups of mice with tamoxifen, reducing the likelihood that the behavioural recovery of *Grin1*^{RESCUE} mice would be obscured by the drug treatment. Vogt et al. (2008) showed that a four-week washout period was sufficient to avoid tamoxifen's effects on cognition [26]. We observed that the biochemical and behavioural measures were remarkably similar with a two- or six-week washout (Fig. 3 and 4, Supplemental Fig. 3, Table 2), suggesting that tamoxifen had little effect on our measures of recovery.

Our results provide striking evidence of the plasticity of the adult brain, particularly in the prefrontal cortex. Within the cortex the highest levels of recovery were observed in *Vglut1+* cells, which normally express the highest levels of *Grin1* [16]. It is possible that very early interventions would provide a more complete recovery in some cell types or brain functions. However, our results suggest that symptoms of intellectual disability, a consistent symptom of *GRIN* encephalopathies [4], can be largely treated with adult intervention. This is particularly surprising since cognitive impairments are refractory to current pharmacological treatment in patients with autism and schizophrenia [27], two conditions associated with impaired NMDAR function [28]. Adult genetic reversal has an even greater clinical impact, as it offers the possibility of stable restoration of normal function even after the brain has completed development [29, 30].

The prevalence of coding variants in *GRIN1,* or in the other *GRIN* genes, is unknown. The first patients to be sequenced had diagnoses of intellectual disability [31] or epilepsy [13]. A recent whole-exome sequencing study reported that 7% of patients with

autism or schizophrenia carry a predicted-deleterious coding mutation in one of six

GRIN genes (25/370 patients with schizophrenia, 15/192 patients with autism) [32].

There have also been numerous genetic and epidemiological studies supporting a

causal role for NMDARs in several neuropsychiatric disorders. Thus, the significance of

our findings is not limited to those patients who have been sequenced to date.

This study highlights the significant potential of therapeutic intervention in adult patients.

It demonstrates that a delay between symptom onset and treatment can be overcome.

The cognitive symptoms of neuropsychiatric and neurodevelopmental conditions

caused by NMDAR hypofunction are amenable to treatment and show persisting

improvement. The mature cortex has sufficient plasticity to recover from insults to this

key developmental system, and adult intervention with the appropriate therapeutic agent

should treat intellectual disabilities.

304
305   **Supplementary information is available at MP's website.**
306

MATERIALS AND METHODS

Animals

Animal housing and experimentation were carried out in accordance with the Canadian

Council in Animal Care (CCAC) guidelines for the care and use of animals and following

protocols approved by the Faculty of Medicine and Pharmacy Animal Care Committee

at the University of Toronto. Mice were group housed with littermates on a 12-h light-

dark cycle and were given *ad libitum* access to water and food (2018 Teklad Global

18% Protein Rodent Diet, Envigo, Madison Wisconsin USA, www.envigo.com).

ROSA26^{CreERT2} mice were obtained from Jackson Laboratory (008463; B6.129-

Gt(ROSA)26Sor$^{tm1(cre/ERT2)Tyj}$/J), and were previously described [25]. *ROSA26^{CreERT2}*

mice were identified by PCR of genomic DNA using the following primers: common

forward 5'-AAAGTCGCTCTGAGTTGTTAT-3', wildtype reverse 5'-

GGAGCGGGAGAAATGGATATG-3', mutant reverse 5'-CCTGATCCTGGCAATTTCG-

3'.

The Cre-reporter mouse line used, *ROSA26^{tdTomato}*, was obtained from Jackson

Laboratory (007914; B6.Cg-*Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J*) [33] and was crossed

with the *ROSA26^{CreERT2}* line. *ROSA26^{tdTomato}* mice were identified by PCR of genomic

DNA using the following primers: wildtype forward 5'-AAGGGAGCTGCAGTGGAGTA-

3', wildtype reverse 5'-CCGAAAATCTGTGGGAAGTC-3', mutant forward 5'-

GGCATTAAAGCAGCGTATCC-3', mutant reverse 5'- CTGTTCCTGTACGGCATGG-3'.

Grin1$^{flneo/flneo}$ mice were generated at the University of Toronto, based on the previously

described *Grin1$^{neo/neo}$* mouse [15]. Identical to the *Grin1$^{neo/neo}$* model, the *Grin1* gene

was modified via homologous recombination with an intervening sequence (neomycin

cassette), and targeted into intron 19, flanked by *loxP* sites (pXena vector; gift of Dr.

Beverly Koller). The linearized targeting construct was electroporated into mouse ES

cells (129/SvlmJ strain) by The Centre for Phenogenomics (TCP, Toronto, ON).

Targeted cell lines were screened by PCR and Southern blot (data not shown) and

chimeras were generated by diploid aggregation. PCR primers for genotyping were:

wildtype forward 5'-TGAGGGGAAGCTCTTCCTGT-3', mutant forward 5'-

GCTTCCTCGTGCTTTACGGTAT-3', common reverse 5'-

AAGCGATTAGACAACTAAGGGT-3'.

Grin1$^{+/flneo}$:CreTg mice were produced by crossing *ROSA26*^{CreERT2} C57Bl/6J congenic

mice to *Grin1*$^{+/flneo}$ C57Bl/6J congenics. The resulting compound heterozygotes were

bred to *Grin1*$^{+/flneo}$ 129/SvlmJ congenics to produce the F1 progeny used for all

experiments as recommended by the Banbury Conference[34].

Tamoxifen Administration

Tamoxifen was administered to all genotypes of mice (WT, WTCre, *Grin1*^{KD},

Grin1^{RESCUE}). Tamoxifen (T5648, Sigma-Aldrich, St. Louis, MO, USA) was administered

via oral gavage (6mg, 20mg/ml dissolved in 100% corn oil at 65°C for 1 hour) on day 1

of treatment, and then mice were given tamoxifen chow (TD.140425, 500mg/kg, Envigo)

ad libitum for 14 days.

Detection of Cre Recombinase Activity

Mice carrying the *ROSA26*^{tdTomato} and *ROSA26*^{CreERT2} alleles were generated and

administered tamoxifen as described above. Three days following completion of

tamoxifen treatment, mice were anaesthetized with 250mg/kg Avertin, which was

composed of [2,2,2-tribromoethanol (Sigma Aldrich) dissolved in 2-methyl-2-butanol

(Sigma Aldrich) 2.5% v/v)]. Whole brains were perfused with 4% paraformaldehyde

(PFA). Frozen sections of 40μm were collected (coronal sections at Bregma 1.78mm,

1.10mm, -1.94mm, -6.00mm). Sections were mounted and visualized using Nikon i80

epifluorescent microscope and Nikon Elements software (NIS-Elements Basic

Research, version 3.1).

Behavioural Testing

Male and female mice of equal numbers were used for behavioural testing. Tests were

administered at PD98 or PD126. All experimental animals were first tested for locomotor

activity on Day 1. Mice were then assigned to one of two groups for subsequent

behavioural tests that spanned three days. The puzzle box test [24] was administered to

mice in Group A over Days 2-4. Mice in Group B were tested in the elevated plus maze

(data not shown) on Day 2, the social affiliative paradigm on Day 3, and pre-pulse

inhibition of acoustic startle on Day 4, as previously described [35–38], and in detail in

Supplementary Information.

Fluorescence In Situ Hybridization

Expression of *Grin1, Vglut1, and Gad1* mRNA in WT, *Grin1*[KD], and *Grin1*[RESCUE] mice

was visualized by RNAscope Multiplex Fluorescent Reagent Kit v2 protocol (ACD Bio;

CA, USA). Fresh frozen mouse brains were used to collect 20μm saggital sections (1.2

mm from midline) onto charged slides (Superfrost Plus, ThermoFisher). Sections were

hybridized to *Grin1* probes (product reference #533691-C1, ACD Bio) and *Vglut1*

probes (#416631-C2), to *Grin1* probes and *Gad1* probes (#400951-C3) or to control

probe mixtures (positive control probes #320881, negative control probes #320871).

Processed slides were stored at $4°C$ before imaging at 20X magnification with an Axio

Scan.Z1 slide scanner (Zeiss, Oberkochen, DEU) or 40X magnification with a

AxioObserverZ1 Inverted Motorized Microscope. Zen Blue software was used to format

images.

Re-analysis of publicly-accessible single-cell transcriptomics data

We obtained single-cell RNAseq data sampled from the adult mouse primary visual

cortex via the Allen Institute for Brain Science's Cell Types database

(http://celltypes.brain-map.org/) [16]. We utilized expression counts for reads mapping

to exons and summarized gene expression per cell using counts per million reads

sequenced (CPM).

GluN1 Immunofluorescent Visualization

Expression of GluN1 protein levels in WT, *Grin1*[KD], and *Grin1*[RESCUE] mice were

visualized by fluorescent immunohistochemistry on fresh frozen sagittal tissue sections

(20 μm thick; Lateral ~1.92mm). Sections were incubated with an in-house rabbit anti-

GluN1 antibody raised against peptide ETEKPRGYQMSTRLK (C) (1:200), and then

with secondary antibody, anti-rabbit Alexa 568 (ThermoFisher, #A11011, 1:500).

[^{3}H]MK-801 Saturation Binding

NMDAR levels in WT, *Grin1^{KD}*, and *Grin1^{RESCUE}* mice were quantified in prefrontal

cortical and hippocampal and tissue. Membranes were prepared and diluted to a

1.6µg/µl working concentration and stored at -80°C. [^{3}H]MK-801 (Perkin Elmer) was

diluted to a working concentration of 120nM (final concentration 40nM) in binding buffer.

Cold MK-801 (Sigma Aldrich) was prepared to a 1200nM working solution (final

concentration 400nM; 10x [^{3}H]MK-801) in binding buffer. Binding assays were

performed with the NMDAR antagonist MK-801 (hot and/or cold), mouse brain

membranes (80µg) and binding buffer (total binding vs. non-specific binding), with a

total assay volume of 150µl. Radioactivity was quantified via liquid scintillation

spectrometry [39].

PSD-95 Immunoprecipitation Mass Spectrometry

As previously described [40], mouse anti-PSD-95 antibody (Millipore, catalogue #

MAB1596) was used to capture PSD-95 protein complexes from flash frozen cortex

samples (3 males and 3 females of each genotype were used). 5µg of PSD-95 antibody

was coupled per 1mg of Dynabeads (Life Technologies; antibody coupling kit protocol

(#14311D). For each sample, 10mg/ml antibody coupled beads were incubated with

mouse brain lysate for 1 hour at room temperature. Captured protein complexes were

eluted and heated to 70°C for 10 minutes then processed for mass spectrometry. All

samples were loaded on a 1.5 mm, 4-12% Bis-Tris Invitrogen NuPage gel

(NP0335BOX) and electrophoresed in 1x MES buffer (NP0002) for 10 minutes at 180v.

Lanes were harvested, cut into ~2mm squares, and subjected to in-gel tryptic digestion

and peptide recovery. Samples were resuspended in 0.1% formic acid. Nano liquid

chromatography coupled electrospray tandem mass spectrometry (nLC-ESI-MS/MS)

analyses were performed on a 5600+ QTOF mass spectrometer (Sciex, Toronto, ON,

Canada) interfaced to an Eksigent (Dublin, CA) nanoLC.ultra nanoflow system.

Peptides were eluted using a variable mobile phase (MP) gradient. The data was

recorded using Analyst-TF (version 1.7) software.

The data independent acquisition (DIA) method was set to go through 1757 cycles for

99 minutes, where each cycle performed one TOF-MS scan type (0.25 second

accumulation time, in a 550.0 to 830.0 m/z window) followed by 56 sequential

overlapping windows of 6 Daltons each. The resulting data were analyzed by Sciex DIA

software to generate peptide intensities.

Electrophysiological Recordings

Mice were euthanized by decapitation under isoflurane anaesthesia. Brains were chilled

in sucrose cutting solution and blocked to obtain the anterior portion and coronal slices

(400 µm) of the medial PFC (mPFC) (1.98 mm-1.34 mm; Paxinos & Franklin Atlas). The

slices were transferred to the stage of an upright microscope for whole cell patch clamp

recordings. Layer V pyramidal neurons in the mPFC were identified by infrared

differential inference contrast microscopy. Experiments were performed in the presence

of CNQX disodium salt (20 µm; Alomone Labs) to block AMPA receptors. NMDA (30

µm; Sigma-Aldrich) was bath applied to assess NMDAR function. Application of APV

(50 µM; Alomone Labs) confirmed the inward currents were mediated by NMDARs. The

peak amplitude of the NMDA currents was measured using Clampfit software

(Molecular Devices). The magnitude of NMDA-elicited inward currents was quantified by

subtracting a 1 second average holding current at the peak from the average holding

current at the baseline.

Quantification and Statistical Analysis

Statistically significant outliers were calculated and excluded, using the Grubb's Test.

Data were analyzed either using a one- or two-way ANOVA (repeated measures) where

indicated, with multiple comparisons and post-hoc Bonferroni's test, as indicated in

figure legends. For electrophysiological recordings, paired t-tests were used to compare

neuronal responses to NMDA before and after APV. Data analysis was not blinded.

Differences in means were considered statistically significant at $p < 0.05$. Significance

levels are as follows; *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, ns – not significant.

All data analyses were performed using the Graphpad Prism 6.0 software and/or IBM

SPSS 23.0 Software.

REFERENCES

REFERENCES

1.   Maulik PK, Mascarenhas MN, Mathers CD, Dua T, Saxena S. Prevalence of
intellectual disability: A meta-analysis of population-based studies. Res Dev
Disabil. 2011;32:419–436.

2.   Fombonne E. Epidemiology of pervasive developmental disorders. Pediatr Res.
2009;65:591–598.

3.   Mohammed S, Scott R, Vogt J, Al-Turki S, Cross G, Smithson S, et al. Large-
scale discovery of novel genetic causes of developmental disorders. Nature.
2014;519:223–228.

4.   XiangWei W, Jiang Y, Yuan H. De novo mutations and rare variants occurring in
NMDA receptors. Curr Opin Physiol. 2018;2:27–35.

5.   Guy J, Gan J, Selfridge J, Cobb S, Bird A. Reversal of neurological defects in a
mouse model of Rett syndrome. Science (80-). 2007;315:1143–1147.

6.   Mei Y, Monteiro P, Zhou Y, Kim J-AA, Gao X, Fu Z, et al. Adult restoration of
Shank3 expression rescues selective autistic-like phenotypes. Nature.
2016;530:481–484.

7.   Li Y, Erzurumlu RS, Chen C, Jhaveri S, Tonegawa S. Whisker-related neuronal
patterns fail to develop in the trigeminal brainstem nuclei of NMDAR1 knockout
mice. Cell. 1994;76:427–437.

8.   Suzuki A, Lo F-S, Zhao S, Itohara S, Hayashi Y, Arakawa H, et al. Thalamic
NMDA Receptor Function Is Necessary for Patterning of the Thalamocortical
Somatosensory Map and for Sensorimotor Behaviors. J Neurosci.
2014;34:12001–12014.

9.   Iwasato T, Datwani A, Wolf AM, Nishiyama H, Taguchi Y, Tonegawa S, et al.
Cortex-restricted disruption of NMDAR1 impairs neuronal patterns in the barrel
cortex. Nature. 2000;406:726–731.

10. Gu X, Zhou L, Lu W. An NMDA Receptor-Dependent Mechanism Underlies
Inhibitory Synapse Development. Cell Rep. 2016;14:471–478.

11. Zhang Z -w., Peterson M, Liu H. Essential role of postsynaptic NMDA receptors in
developmental refinement of excitatory synapses. Proc Natl Acad Sci.
2013;110:1095–1100.

12. Espinosa JS, Wheeler DG, Tsien RW, Luo L. Uncoupling Dendrite Growth and
Patterning: Single-Cell Knockout Analysis of NMDA Receptor 2B. Neuron.
2009;62:205–217.

13.   Lemke JR, Geider K, Helbig KL, Heyne HO, Schütz H, Hentschel J, et al.
498   Delineating the GRIN1 phenotypic spectrum: A distinct genetic NMDA receptor
499   encephalopathy. Neurology. 2016;86:2171–2178.

500   14.   Forrest D, Yuzaki M, Soares HD, Ng L, Luk DC, Sheng M, et al. Targeted
501   disruption of NMDA receptor 1 gene abolishes NMDA response and results in
502   neonatal death. Neuron. 1994;13:325–338.

503   15.   Mohn AR, Gainetdinov RR, Caron MG, Koller BH. Mice with reduced NMDA
504   receptor expression display behaviors related to schizophrenia. Cell.
505   1999;98:427–436.

506   16.   Tasic B, Yao Z, Graybuck LT, Smith KA, Nguyen TN, Bertagnolli D, et al. Shared
507   and distinct transcriptomic cell types across neocortical areas. Nature.
508   2018;563:72–78.

509   17.   Rompala GR, Zsiros V, Zhang S, Kolata SM, Nakazawa K. Contribution of NMDA
510   Receptor Hypofunction in Prefrontal and Cortical Excitatory Neurons to
511   Schizophrenia-Like Phenotypes. PLoS One. 2013;8:e61278.

512   18.   Belforte JE, Zsiros V, Sklar ER, Jiang Z, Yu G, Li Y, et al. Postnatal NMDA
513   receptor ablation in corticolimbic interneurons confers schizophrenia-like
514   phenotypes. Nat Neurosci. 2010;13:76–83.

515   19.   Finlay JM, Dunham GA, Isherwood AM, Newton CJ, Nguyen T V, Reppar PC, et
516   al. Effects of prefrontal cortex and hippocampal NMDA NR1-subunit deletion on
517   complex cognitive and social behaviors. Brain Res. 2015;1600:70–83.

518   20.   Yamaguchi S, Hale LA, D'Esposito M, Knight RT. Rapid prefrontal-hippocampal
519   habituation to novel events. J Neurosci. 2004;24:5356–5363.

520   21.   Ranganath C, Rainer G. Cognitive neuroscience: Neural mechanisms for
521   detecting and remembering novel events. Nat Rev Neurosci. 2003;4:193–202.

522   22.   Li L, Du Y, Li N, Wu X, Wu Y. Top-down modulation of prepulse inhibition of the
523   startle reflex in humans and rats. Neurosci Biobehav Rev. 2009;33:1157–1167.

524   23.   Duncan GE, Moy SS, Perez A, Eddy DM, Zinzow WM, Lieberman JA, et al.
525   Deficits in sensorimotor gating and tests of social behavior in a genetic model of
526   reduced NMDA receptor function. Behav Brain Res. 2004;153:507–519.

527   24.   Ben Abdallah NM-B, Fuss J, Trusel M, Galsworthy MJ, Bobsin K, Colacicco G, et
528   al. The puzzle box as a simple and efficient behavioral test for exploring
529   impairments of general cognition and executive functions in mouse models of
530   schizophrenia. Exp Neurol. 2011;227:42–52.

25. Ventura A, Kirsch DG, McLaughlin ME, Tuveson DA, Grimm J, Lintault L, et al.
Restoration of p53 function leads to tumour regression in vivo. Nature.
2007;445:661–665.

26. Vogt MAA, Chourbaji S, Brandwein C, Dormann C, Sprengel R, Gass P.
Suitability of tamoxifen-induced mutagenesis for behavioral phenotyping. Exp
Neurol. 2008;211:25–33.

27. Millan MJ, Agid Y, Brüne M, Bullmore ET, Carter CS, Clayton NS, et al. Cognitive
dysfunction in psychiatric disorders: Characteristics, causes and the quest for
improved therapy. Nat Rev Drug Discov. 2012;11:141–168.

28. Tarabeux J, Kebir O, Gauthier J, Hamdan FF, Xiong L, Piton A, et al. Rare
mutations in N-methyl-D-aspartate glutamate receptors in autism spectrum
disorders and schizophrenia. Transl Psychiatry. 2011;1:e55.

29. Duyne GD Van. Cre Recombinase. Mob. DNA III, vol. 3, American Society of
Microbiology; 2015. p. 119–138.

30. Speed HE, Kouser M, Xuan Z, Liu S, Duong A, Powell CM. Apparent Genetic
Rescue of Adult Shank3 Exon 21 Insertion Mutation Mice Tempered by
Appropriate Control Experiments. ENeuro. 2019:ENEURO.0317-19.2019.

31. Hamdan FF, Gauthier J, Araki Y, Lin DT, Yoshizawa Y, Higashi K, et al. Excess of
de novo deleterious mutations in genes associated with glutamatergic systems in
nonsyndromic intellectual disability. Am J Hum Genet. 2011;88:306–316.

32. Yu Y, Lin Y, Takasaki Y, Wang C, Kimura H, Xing J, et al. Rare loss of function
mutations in N-methyl-d-aspartate glutamate receptors and their contributions to
schizophrenia susceptibility. Transl Psychiatry. 2018;8:12.

33. Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, et al. A robust
and high-throughput Cre reporting and characterization system for the whole
mouse brain. Nat Neurosci. 2010;13:133–140.

34. Silva AJ, Simpson EM, Takahashi JS, Lipp HP, Nakanishi S, Wehner JM, et al.
Mutant mice and neuroscience: Recommendations concerning genetic
background. Neuron. 1997;19:755–759.

35. Islam R, Trépanier M-O, Milenkovic M, Horsfall W, Salahpour A, Bazinet RP, et al.
Vulnerability to omega-3 deprivation in a mouse model of NMDA receptor. Npj
Schizophr. 2017;3:12.

36. Mielnik CAA, Horsfall W, Ramsey AJJ. Diazepam improves aspects of social
behaviour and neuron activation in NMDA receptor-deficient mice. Genes, Brain
Behav. 2014;13:592–602.

37.   Milenkovic M, Mielnik CAA, Ramsey AJJ. NMDA receptor-deficient mice display
567   sexual dimorphism in the onset and severity of behavioural abnormalities. Genes,
568   Brain Behav. 2014;13:850–862.

569   38.   Moy SS, Nadler JJ, Young NB, Perez A, Holloway LP, Barbaro RP, et al. Mouse
570   behavioral tasks relevant to autism: Phenotypes of 10 inbred strains. Behav Brain
571   Res. 2007;176:4–20.

572   39.   DeBlasi A, O'Reilly K, Motulsky HJ. Calculating receptor number from binding
573   experiments using same compound as radioligand and competitor. Trends
574   Pharmacol Sci. 1989;10:227–229.

575   40.   Sullivan CRCR, Mielnik CACA, O'Donovan SMSM, Funk AJAJ, Bentea E,
576   DePasquale EAEA, et al. Connectivity Analyses of Bioenergetic Changes in
577   Schizophrenia: Identification of Novel Treatments. Mol Neurobiol. 2018;56.

578   41.   Franklin KBJ, Paxinos G. Paxinos and Franklin's The mouse brain in stereotaxic
579   coordinates.

Acknowledgments: The authors would like to acknowledge Beverly Koller for donation
of the pXena targeting construct, and Marc Caron, Bob Lefkowitz, Michael Didriksen,
Jean-Martin Beaulieu, and Stephane Angers for helpful discussion. The authors would
like to acknowledge Chinmaya Sadangi for preliminary work on RNAscope microscopy.
Funding: This work was supported by CIHR funding to AJR (MOP119298), EKL
(MOP89825 and CRC in Developing Cortical Physiology) and AS (MOP206649) and by
NIMH to REM and AJF (MH107916 and L.I.F.E.). The work was also supported by
graduate scholarships from CIHR and OGS to CAM, OMRI to MAB and a postdoctoral
fellowship from Stiftelsen Olle Engkvist Byggmästare to EMJ. **Competing Interests:**
Authors declare no competing interests.

Figure 1. Generation and molecular characterization of the *Grin1* inducible-rescue mouse line. (A) Targeting construct schematic and recombination events at the *Grin1* locus in the generation of the *Grin1*^flneo/flneo^ mouse model (top panel). PGK, PGK promotor; neo, neomycin selection cassette; polyA, polyadenylation sequence. **(B)** Schematic of expected molecular events at the *Grin1* locus. **(C)** RNAScope visualization of *Grin1* mRNA in mouse sagittal sections (20μm) via fluorescent in situ hybridization using *Grin1* probes *(left: whole brain; middle: 20x micrograph of prefrontal cortex)*. At right, sagittal sections (20μm) of prefrontal cortex visualizing GluN1 protein expression via fluorescent immunohistochemistry with a rabbit anti-GluN1 antibody (in-house, 1:200; secondary anti-rabbit Alexa 568) in WT, *Grin1*^KD^, and *Grin1*^RESCUE^ mice.

Figure 2. *Grin1* gene expression levels in adult mouse cortex. *Grin1* mRNA expression in *Vglut*+ **(A)** and *Gad1*+ **(B)** cells in the adult mouse cortex, visualized with RNAScope. *Grin1* (orange), *Vglut1* (green) and *Gad1* (yellow) mRNA was visualized in mouse sagittal sections (20μm) with fluorescent in situ hybridization in WT, *Grin1*^KD, and *Grin1*^RESCUE mice. **(C)** *Grin1* gene expression levels in wild type adult mouse visual cortex glutamatergic (*Vglut1*+) and GABAergic (*Gad1*+) cells based on publicly accessible single cell transcriptomics data provided by the Allen Institute for Brain Science [16]. Data are quantified as counts per million reads sequenced (CPM) and are based on exonic reads only. To aid visualization, only cells up to the 95[th] percentile of *Grin1* expression are shown. Data shown as box and whisker plots, 5-95 percentile, Wilcoxon rank-sum test between GLU and GABA. Solid white arrows indicate cells with *Grin1* expression, red arrows indicate cell without *Grin1* expression.

Figure 3. NMDAR currents and synaptic GluN1 peptide levels are restored in the PFC.
(A) Schematic of mPFC with whole cell patch clamp recording from layer 5[41]. Inset:
electrophysiological signature of *Grin1^RESCUE* layer 5 pyramidal neuron. **(B)** Representative
traces in voltage-clamp (-75 mV) showing prefrontal response to NMDA across the three
genotypes (number of layer 5 pyramidal neurons shown, 5-6 mice per genotype). **(C)**
Quantification of peak amplitude of prefrontal NMDAR-elicited currents. Data shown as mean ±
SEM, ****p<0.0001, ns=not significant, one-way ANOVA, effect of genotype, $F_{2,95}$ =22,
p<0.0001, Bonferroni posthoc. **(D)** Capacitance of prefrontal layer 5 pyramidal neurons. Data
shown as mean ± SEM, *p<0.05, ns =not significant, one-way ANOVA, Bonferroni posthoc. **(E)**
Current density of prefrontal NMDA-elicited currents. Data shown as mean ± SEM, **p<0.01,
****p<0.0001, one-way ANOVA, effect of genotype, $F_{2,95}$ =3.6, p=0.03, Bonferroni posthoc. **(F)**
Synaptic GluN1 peptide (IVNIGAVLSTR) levels in the PFC. Data shown as mean ± SEM,
****p<0.0001, ns=not significant, one-way ANOVA, effect of genotype, PFC: $F_{2,14}$ =64.76,
p<0.0001, Bonferroni posthoc.

Figure 4. Cognitive symptoms show rescue following intervention. (A) Time-course and total distance traveled (cm) in open field (OF) test in WT, *Grin1^KD^*, and *Grin1^RESCUE^* mice. Data shown as mean ± SEM, ****p<0.0001, two-way ANOVA (time-course), one-way ANOVA (total), effect of genotype, $F_{3,164}$=176.34, p<0.0001, Bonferroni posthoc. Habituation index (time to reach half maximal activity) in OF test in WT, *Grin1^KD^*, and *Grin1^RESCUE^* mice. Data shown as mean ± SEM, ****p<0.0001, one-way ANOVA, effect of genotype, $F_{3,167}$=14.13, p<0.0001, Bonferroni posthoc. **(B)** Percent inhibition of startle response (PPI) in WT, *Grin1^KD^*, *Grin1^RESCUE^* mice. Shown on graph, *Grin1^KD^* vs. *Grin1^RESCUE^*: ***p<0.001, ****<0.0001, data shown as mean

± SEM, one-way ANOVA (within pre-pulse decibel), effect of genotype: $F_{3,75}$= (4dB)8.81,
p<0.001, (8dB)14.59, p<0.001, (16db)9.57, p<0.001, Bonferroni posthoc. **(C)** Schematic of
puzzle box test. Time to reach goal zone (seconds; max 300sec) measured in puzzle box
paradigm in WT, *Grin1^{KD}*, *Grin1^{RESCUE}* mice. Shown on graph, #WT vs. *Grin1^{RESCUE}*: ##p<0.01,
####p<0.0001; **Grin1^{KD}* vs. *Grin1^{RESCUE}*: **p<0.01, ***p<0.001, ****p<0.0001. Data shown as
mean ± SEM, two-way ANOVA, effect of genotype: $F_{3,82}$=53.63, p<0.001, Bonferroni posthoc.
N-values: WT – 21, *Grin1^{KD}* – 25, *Grin1^{RESCUE}* - 21 **(D)** Representative traces of modified three-
chamber sociability. Total time spent, number of visits, and time per visit in the social zone
during 10min. test, measured in WT, *Grin1^{KD}*, and *Grin1^{RESCUE}* mice. Data shown as mean ±
SEM, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, ns = not significant, one-way ANOVA,
effect of genotype: (time) $F_{3,75}$=7.14, p<0.0001, (visits) $F_{3,78}$=2.31, p<0.0001, (time/visit)
$F_{3,77}$=12.55, p<0.0001, Bonferroni posthoc.

Figure 5. Replicable and robust cognitive improvements following adult intervention with longer recovery time. (A) Time-course and total distance traveled (inset) in open field (OF) test in WT, *Grin1KD*, and *Grin1RESCUE* mice. Data shown as mean ± SEM, ****p<0.0001,

two-way ANOVA (time-course), one-way ANOVA (total), effect of genotype, $F_{3,168}$=339.43,

p<0.0001, Bonferroni posthoc. Habituation index (time to reach half maximal activity) in OF test

in WT, *Grin1*^{KD}, and *Grin1*^{RESCUE} mice. Data shown as mean ± SEM, ****p<0.0001, one-way

ANOVA, effect of genotype, $F_{3,170}$=26.82, p<0.0001, Bonferroni posthoc. **(B)** Percent inhibition

of startle response (PPI) in WT, *Grin1*^{KD}, *Grin1*^{RESCUE} mice. Shown on graph, *Grin1*^{KD} vs.

Grin1^{RESCUE}: **p<0.01, ***p<0.001, data shown as mean ± SEM, one-way ANOVA, effect of

genotype: $F_{3,80}$= (4dB)20.03, p<0.001, (8dB)11.33, p<0.001, (16db)6.03, p<0.001, Bonferroni

posthoc. **(C)** Time to reach goal zone (seconds; max 300sec) measured in puzzle box

paradigm in WT, *Grin1*^{KD}, *Grin1*^{RESCUE} mice. Shown on graph, #WT vs. *Grin1*^{RESCUE}: #p<0.05;

*Grin1^{KD} vs. *Grin1*^{RESCUE}: ***p<0.001, ****p<0.0001. Data shown as mean ± SEM, two-way

ANOVA, effect of genotype: $F_{3,81}$=56.48, p<0.001, Bonferroni posthoc. N-values: WT – 26,

Grin1^{KD} – 20, *Grin1*^{RESCUE} - 21 **(D)** Total time spent, number of visits, and time per visit in the

social zone during 10min. test, measured in WT, *Grin1*^{KD}, and *Grin1*^{RESCUE} mice. Data shown

as mean ± SEM, **p<0.01, ****p<0.0001, ns = not significant, one-way ANOVA, effect of

genotype: (time) $F_{3,83}$=8.74, p<0.0001, (visits) $F_{3,83}$=96.87, p<0.0001, (time/visit) $F_{2,63}$=11.12,

p<0.0001, Bonferroni posthoc.