The genetic intractability of *Symbiodinium microadriaticum* to standard algal transformation methods

Jit Ern Chen[1], Adrian C. Barbrook[2], Guoxin Cui[1], Christopher J. Howe[2], Manuel Aranda[1]

Author Affiliations

[1]Red Sea Research Center, King Abdullah University of Science and Technology (KAUST), Thuwal,

Saudi Arabia.

[2]Department of Biochemistry, University of Cambridge, Cambridge, United Kingdom

Abstract

Modern transformation and genome editing techniques have shown great success across a broad variety of organisms. However, no study of successfully applied genome editing has been reported in a dinoflagellate despite the first genetic transformation of *Symbiodinium* being published about 20 years ago. Using an array of different available transformation techniques, we attempted to transform *Symbiodinium microadriaticum* (CCMP2467), a dinoflagellate symbiont of reef-building corals, with the view to performing subsequent CRISPR-Cas9 mediated genome editing. Plasmid vectors designed for nuclear transformation containing the chloramphenicol resistance gene under the control of the CaMV p35S promoter as well as several putative endogenous promoters were used to test a variety of transformation techniques including biolistics, electroporation and silicon carbide whiskers. Chloroplast-targeted transformation were attempted using an engineered *Symbiodinium* chloroplast minicircle encoding a modified PsbA protein that confers atrazine resistance. We report that we have been unable to confer chloramphenicol or atrazine resistance to *Symbiodinium microadriaticum* strain CCMP2467.

Introduction

Efforts to understand better the molecular mechanisms which govern the symbiosis between marine Cnidarians and their dinoflagellate symbionts have been hampered by the lack of genetically tractable model organisms. This is especially true for the symbiotic relationship between corals and dinoflagellates from the genus *Symbiodinium*. This interaction forms the bedrock of the coral ecosystem (Muscatine et al., 1981; Muscatine and Porter, 1977) and yet is highly sensitive to relatively small changes in environmental conditions (Brown, 1997). Abnormally high ocean temperatures have been identified as one of the key factors that can precipitate the breakdown of the *Symbiodinium*-coral symbiosis, which can lead to wide-spread, regional and even global coral bleaching events (Hughes et al., 2017). The predicted increase in ocean temperatures due to anthropogenic climate change is expected to accelerate this crisis (Smith et al., 2007).

One of the prevailing theories as to how coral reefs would be able to withstand rising ocean temperatures rests on the assumption that there are certain thermo-tolerant *Symbiodinium* strains which are able to form a more robust relationship with their host (Howells et al., 2012). While the effectiveness of thermo-tolerant *Symbiodinium* strains in maintaining algal-coral symbiosis has been shown to be significant, the genetic basis of this robustness remains unknown. Without genetic tools and the absence of any viable method to carry out traditional genetic studies such as inbreeding and cross-breeding, isolating and confirming the identity of thermo-tolerance genes will be difficult.

Previous studies have described methods for transformation of free-living *Symbiodinium* cells. The first, published in 1997 by Ten Lohuis and Miller describes the transformation of *Symbiodinium* CS-153 using silicon carbide whiskers (ten Lohuis and Miller, 1998). In the ten Lohuis paper it was reported that the Cauliflower Mosaic Virus p35S and *Agrobacterium* nos and p1'2' promoters were able to drive the expression of reporter genes (GUS) and selectable markers (hygromycin and geneticin resistance genes) in *Symbiodinium*. Subsequently, two papers published in 2015 by Ortiz-

Matamoros et al. described transformation using glass beads agitation with and without

Agrobacterium of *Symbiodinium kawagutii*, *Symbiodinium* sp. Mf11.5b.1 and *Symbiodinium*

microadriaticum MAC-CassKB8 (Ortiz-Matamoros et al., 2015a; Ortiz-Matamoros et al., 2015b). In

these publications, the authors used the *nos* promoter to drive the expression of the *ba*r gene which

confers resistance to the herbicide Basta. However, the authors of the paper note that their transiently

transformed cells lost their chlorophyll and were unable to reproduce under herbicide selection.

We used previously published transformation protocols for *Symbiodinium* (ten Lohuis and Miller, 1998)

as well as different standard protocols for alga based on electroporation, biolistics and glass beads

agitation to attempt to transform *S. microadriaticum* CCMP2467. Plasmid constructs utilizing the

Cauliflower Mosaic Virus p35S promoter as well as putative endogenous *Symbiodinium* promoter and

terminator regions identified from the S. *microadriaticum* genome (Aranda et al., 2016) were used to

drive the chloramphenicol resistance gene for nuclear transformation, while artificial *Symbiodinium*

plastid minicircles modified to carry a mutated *psbA* gene that is predicted to confer resistance to the

atrazine herbicide were used to carry out chloroplast-targeted transformations. However, we were

unable to obtain resistant strains under chloramphenicol or atrazine selection using either set of

constructs.

We also carried out an attempted nuclear transformation using a construct carrying the geneticin

resistance gene under the control of the CaMV p35S promoter and NosT terminator with *Symbiodinium*

strain CS-153, the strain that was used in the ten Lohuis and Miller (1998) paper. After 10 weeks, we

did not see any growth on geneticin agar plates or in geneticin liquid cultures.

Results

Antibiotic susceptibility test

Previous studies (Ortiz-Matamoros et al., 2015b; ten Lohuis and Miller, 1998) used notably high

concentrations of hygromycin and geneticin (G418) of around 3 mg/ml in order to select for resistant

Symbiodinium transformants, a process which required about three months as untransformed cells were

able to survive up to eight weeks of antibiotics exposure.

In order to select a more cost-effective selection antibiotic and to confirm that our main transformation

strain CCMP2467 has similar antibiotic tolerance levels to strain CS-153, we used liquid cultures to

test the effectiveness of several antibiotics to determine a more potent and less expensive alternative to

hygromycin and geneticin. We determined that 100 µg/ml of chloramphenicol was just as effective as

2.5 mg/ml of hygromycin in reducing the number of observed *Symbiodinium* CCMP2467 cells in

culture (see Figure 1). In addition, the cost of chloramphenicol is significantly lower than hygromycin

or geneticin weight-for-weight across a range of suppliers (see Supplementary Table 1). Therefore, we

decided to use chloramphenicol instead of hygromycin or geneticin as our main antibiotic to select for

transformants. A similar, independent test was also carried out to determine an effective selection

concentration for atrazine resistance, which was found to be 150 ng/ml (see Supplementary Table 2).

102 *Figure 1*: Growth of *Symbiodinium* CCMP2467 in f/2 liquid medium under antibiotic selection. Results
103 were from three biological replicates, each measured four times using a FlowCAM. The error bars
104 indicate standard error of the mean.
105

Test for chloramphenicol resistance gene function

We tested the functionality of the chloramphenicol resistance gene using *Saccharomyces cerevisiae*

expression vectors in *S. cerevisiae* (see Figure 2). Under galactose induction, yeast cultures were grown

with and without chloramphenicol selection (4 mg/ml) to show that the presence of the chloramphenicol

resistance gene (ChloR) under the control of the *GAL1* promoter was necessary and sufficient to confer

increase chloramphenicol resistance to yeast cells. The minimal growth medium used for this

experiment was supplemented with 3% glycerol and 0.5% galactose in order to force yeast cells to grow

anaerobically while providing a low level of galactose to induce *GAL1* promoter expression. The results

showed that yeast cells transformed with the pYESChloR construct (see Supplementary Data 1), which

contains the ChloR gene under the control of the *GAL1* promoter, became more resistant to

chloramphenicol compared to the control untransformed strain (31019b) and the strain transformed with

the empty vector pYESeGFP (see Supplementary Data 2). We present this as evidence for the ability of

the *ChloR* gene to facilitate chloramphenicol resistance when expressed as a transgene.

Figure 2: *S. cerevisiae* growth under chloramphenicol selection. "Parental" refers to the parental,
untransformed *S. cerevisiae* yeast strain 31019b. "Empty vector" refers to the parental yeast strain
transformed with a pYES2.1 vector carrying an eGFP coding region. "Chlo Resist" refers to the parental
yeast strain transformed with a pYES2.1 vector carrying the chloramphenicol acyltransferase (CAT)
resistance gene (ChloR). Results are from five biological replicates. Asterisks indicate significantly
different means with a p-value of < 0.01 using Student's t-test with unequal variance. T-test was carried
out with a significance value of 0.05. Error bars indicate standard error of the mean.

Promoter region identification

In order to identify suitable promoter regions we chose a set of five expressed genes including those for

Actin, *β-Tubulin A*, *β-Tubulin B*, *Hsp90*, and *PsbJ* from the *S. microadriaticum* genome as confirmed

by transcriptomics data (see Supplementary Table 3) from Chen et al. (2017). To identify the right start

codon for the Actin and PsbJ genes we performed 5' RACE on transcripts to sequence their 5' UTRs.

For the *Actin* gene, the 5' RACE results confirmed the presence of a spliced leader 55 bp upstream of

the first ATG site (see Figure 3). This particular spliced leader sequence is notable for being the most

common spliced leader sequence in the *Symbiodinium kawagutii* transcriptome, being found in 6235

out of 6501 full-length *S. kawagutii* cDNAs containing a complete spliced leader sequence (Zhang et

al., 2013). Using the coding sequence, we did a BLAST search on the *S. microadriaticum* genome

(Aranda et al., 2016) and identified gene model Smic17360 as the best hit sequence for our 5' RACE

result. We then amplified an approximately 2 kb region upstream from the putative start ATG to serve

as the promoter element for the pAct-ChloR-ActT construct. The 3' terminator sequence used for this

construct was the 500 bp region downstream of the stop codon of Smic17360.

GGACACTGACATGGACTGAAGGAGTAGAAA**TCCGTAGCCATTTTGGCTCAAG**TTAG
CCATGTCGGACGAGGTTGTCGCTCTCGTGGTGGACAATGGCAGTGGCATGTGCAAG
GCCGGCTTCTCCGGTGATGACGCTCCGCGAACGGTTTTTCCCTCCCTGATCGGGCG
ACCGAAGCACCGGAGCATTGTGGTGGGGATGGATCAGAGAGACCACTACGTCGGAG
ACGAAGCACAAGCCAAGCGCGGTGTCTTGGCTCTCAAGTACCCTATCGAGCACGGC
ATCGTGACGAACTGGGACGACATGGAGCGGATCTGGCACCACACTTTCTACAA

5' RACE forward primer
5' UTR
Splice leader
Putative Actin coding sequence
Degenerate Actin reverse primer

Figure 3: 5' RACE results of Actin transcript to identify *Actin* 5' UTR sequence.

Within the 2 kb promoter region upstream of Smic17360 (*Actin*), we identified all promoter elements that have been noted as important for proper *trans*-splicing of the leader (see Figure 4A) (Lin et al., 2015). These include a TTTT-box motif 34 bp upstream of a potential transcriptional start site (YYANWYY) and a branch point (YTNAY) 32 bp upstream of the splice acceptor (AG) (see Figure 4B).

Figure 4: Unique promoter architecture in *Symbiodinium*. (A) A schematic view of the proposed promoter (TTTT-box) relative to the putative transcription start site, splice branch point and acceptor site, upstream of the coding region (CDS) in the genomic sequence of *S. kawagutii* genes. For comparison, premature mRNA and mature mRNA are also shown. Figure was adapted from Lin et al.

(2015). (B) Genomic sequence of *S. microadriaticum pAct* promoter 175 bp upstream of the putative
start codon, with promoter elements as described in (A).

We carried out a similar 5' RACE validation experiment to define the *PsbJ* promoter (*pPsbJ*) region

(see Supplementary Figure 1A). In most photosynthetic organisms, the *psbJ* gene is located in the

chloroplast genome as part of the *psbEFLJ* operon. However, in the *S. microadriaticum* genome the

PsbJ gene exists as a three copy tandem repeat in the nucleus (see Supplementary Figure 1B), which

lends support to previous reports that *psbJ* is not encoded on any of the *Symbiodinium* chloroplasts

minicircles (Howe et al., 2008). For these reasons, the 5' region of the *pPsbJ-ChloR* construct was

designed to encode an additional N-terminal peptide which are absent from all other, chloroplast-

encoded orthologues of *PsbJ* and which could be the *Symbiodinium* chloroplast localization peptide

(see Figure 5). Note that because the *Symbiodinium* PsbJ is encoded in the nucleus, we are using the

gene name format for higher plant nuclear genes (capitalized first letter, *PsbJ*) rather than the naming

format for plastid genes (lower case first letter, *psbJ*).

Figure 5: Alignment of putative *S. microadriaticum* PsbJ amino acid sequences against other psbJ
orthologues. The Smic_PsbJ_Long variant is due to an alternate in-frame start codon upstream of the
start codon that would be equivalent to the one found in other psbJ homologues, and this additional N-
teminal polypeptide region could be involved in plastidic sub-cellular compartmentalization signaling.
The NCBI reference sequence ID are as follows: *Phaeodactylum tricornutum* YP_874372.1,
Arabidopsis thaliana NC_000932.1, *Arabidopsis lyrata* NC_034365.1, *Nicotiana tabacum*
NC_001879.2.

For the nuclear constructs based on the β-Tubulin A, β-Tubulin B and Hsp90 promoters, we took the 3

kb region upstream of the predicted start codon as the putative promoter region and a 400 bp 3' region

originating downstream of the endogenous Hsp90 stop codon as a terminator region (see Supplementary

Data 3, 4 and 5). These regions were selected only based on *in silico* analysis, and were not

independently identified via RACE.

Transformation construct design

Plasmids were constructed to carry the chloramphenicol (*ChloR*), geneticin (*GenR*) or the atrazine

(*PsbA^{S262G}*) resistance gene under the control of a series of different gene promoters and terminators.

We use the term *ChloR* here to describe the chloramphenicol acetyltransferase (*CAT*) gene which

confers resistance to chloramphenicol and *GenR* to describe the neomycin phosphotransferase II (*nptII*)

gene that confers resistance to geneticin G148. Similarly, the gene we term *PsbA^{S262G}* is a variant of the

psbA gene that contains an amino acid mutation that is predicted to confer resistance to atrazine.

Table 1: List of constructs. Detailed plasmids maps for all constructs in this table can be found in
Supplementary Data 6 to 13.

Construct name	Plasmid backbone	Promoter origin	Resistance gene	Terminator origin
p35S-ChloR-NosT	pCR4	Cauliflower mosaic virus 35S	Chloramphenicol acetyltransferase	Agrobacterium Nopaline synthase
pAct-ChloR-ActT	pCR2.1	Symbiodinium SmicGene17360	Chloramphenicol acetyltransferase	Symbiodinium SmicGene17360
pBTubA-ChloR	pCR2.1	Symbiodinium SmicGene25527	Chloramphenicol acetyltransferase	(no terminator)
pBTubB-ChloR	pCR2.1	Symbiodinium SmicGene4192	Chloramphenicol acetyltransferase	(no terminator)
pHsp90-ChloR-Hsp90T	pCR2.1	Symbiodinium SmicGene33536	Chloramphenicol acetyltransferase	Symbiodinium SmicGene33536
pPsbJ-ChloR	pCR2.1	Symbiodinium SmicGene31399	Chloramphenicol acetyltransferase	(no terminator)
pPsbA^{S262G}GEM	Chloroplast minicircle	Endogenous PsbA	Mutant PsbA, resistant to atrazine	Endogenous PsbA
pPsbAGEM	Chloroplast minicircle	Endogenous PsbA	none	Endogenous PsbA

The plasmids constructed can be classed into two major groups based on the target site of plasmid

transformation (see Table 1). Chloramphenicol and geneticin constructs, derived from commercial *E.*

coli vector backbones, were designed for nuclear *Symbiodinium* transformation while atrazine

constructs were designed as artificial chloroplast minicircle chromosomes. In addition to the promoter

and terminator regions previously described, the p35S promoter and Nos terminator were also used to

construct the p35S-ChloR-NosT (see Figure 6) and p35S-Neo-2A-eCFP constructs, as it has been

reported previously that these expression elements were sufficient to drive the expression of the

hygromycin resistance gene in *Symbiodinium* (ten Lohuis and Miller, 1998).

213 *Figure 6*: Vector map for plasmid p35S-ChloR-NosT, constructed using the pCR2.1 vector backbone.
214 This vector map was made using PlasMapper (Dong et al., 2004).
215

The artificial chloroplasts minicircle constructs are fusions of a PCR-amplified *psbA* minicircle with

the cloning vector pGEM-T-Easy, creating a simple shuttle vector for use in transformation experiments

(pPsbAGEM). The organization of the shuttle vector is shown in Supplementary Data 10. The *psbA*

minicircle of *S. microadriaticum* was amplified with back-to-back primers. The primers were designed

to anneal to regions of DNA just downstream of the *psbA* coding region, since subsequent inclusion of

the vector sequence here seemed less likely to disrupt promoter or origin of replication sequences. The

shuttle vector pPsbAGEM was then modified by site-directed mutagenesis to create pPsbA^{S262G}GEM

which contains two nucleotide changes that change the serine codon for residue 262 to a glycine codon

and a further synonymous base change in a neighboring codon to act as a marker.

Transformation details (*S. microadriaticum* strain CCMP2467)

A brief summary of the various transformation experiments carried out can be seen in Table 2, and the

details of each transformation can be found in Supplementary Table 4. 45 sets of transformation

experiments were carried out using strain CCMP2467 for a total of 207 transformation samples,

including control samples (i.e. no plasmid control). Various transformation conditions for each

transformation protocol were tested, and all treated cultures were kept under observation for at least 3

months in liquid culture with some of the experiments also selected for on agar plates.

Table 2: Summary list of the number of experiments carried out grouped by transformation method.
Details on each transformation can be found in Supplementary Table 4.
237

Transformation method	Number of experiments	Number of samples
Electroporation	17	69
Biolistics	9	57
Glass beads agitation	9	21
Silicon carbide whiskers agitation	7	31
FuGENE transfection	3	29
Total	45	207

Electroporation was the most used transformation method both in terms of the number of experiments

and the number of samples treated. Both exponential decay and multi-square wave pulses were used,

with field strengths ranging from 20 V to 2.2 kV. For biolistics, both gold and tungsten particles of

various sizes were used with varying rupture disks strengths ranging from 900 to 1550 psi. The least

tested method, FuGENE transfection, was only tested three times, using the recommended

manufacturer's protocol, with the variable factors being the amount of plasmid DNA used and the ratio

of DNA to FuGENE solution (see Supplementary Table 4E).

In both solid and liquid tests, transformations were considered to have failed when cells in the no-

antibiotic/no-herbicide control liquid cultures/agar plates became visibly decoloured due to senescence,

while at the same time no growth could be observed in the selection cultures/plates. For agar plates,

control culture senescence usually took around six weeks and for liquid cultures it was around three

months.

For the selective agar plates, none of the putative transformant cultures tested survived more than four

weeks. For liquid culture selection, any cultures which showed signs of growth after roughly one month

were sub-cultured into fresh f/2 liquid or agar with the appropriate selection antibiotic/herbicide.

Although we did observe that some of these cultures began to grow under selection after several months

in liquid culture, none of these treated cultures were able to grow after sub-culturing in fresh selection

medium (data not shown). We therefore considered these growths as the result of untransformed cells

that were able to outlast the effective period of their respective selection antibiotic/herbicide and not

true stable transformants.

Transformation details (*Symbiodinium* strain CS-153)

We concentrated most of our transformation efforts on the *Symbiodinium* strain CCMP2467 because

the genome of this strain had been sequenced and cultures of this strain are currently being used for

other research projects in our group. However, the published protocol by ten Lohuis and Miller (1998)

used a different *Symbiodinium* strain, known as CS-153, isolated from a different cnidarian and

geographic location from CCMP2467 (see Materials and Methods). In addition, our nuclear

transformation constructs were designed to use chloramphenicol rather than hygromycin or geneticin

as selection markers.

We therefore decided to replicate as closely as feasible the materials and methods used in a previously

published protocol (ten Lohuis and Miller, 1998). We therefore carried out a silicon carbide whiskers

agitation transformation using *Symbiodinium* strain CS-153 with our chloramphenicol and geneticin

constructs (see Table 3). The two geneticin constructs contained the neomycin/geneticin resistance

gene, Transposase Tn5 Aminoglycoside 3' phosphotransferase (GenR), under the control of a

Chlamydomonas reinhardtii chimeric HSP70A/RBCS2 promoter (Supplementary Data 14) and a p35S

promoter (Supplementary Data 15) respectively. These constructs are not identical to the ones that were

used in the original transformation ten Lohuis paper, which unfortunately are no longer available.

Table 3: List of constructs used for CS-153 transformation using silicon carbide whiskers. Plasmid maps
can be found in Supplementary Data 6, 7, 9, 14 and 15.

Construct name	Plasmid backbone	Promoter origin	Resistance gene	Terminator origin
p35S-ChloR-NosT	pCR4	Cauliflower mosaic virus 35S	Chloramphenicol acetyltransferase	Agrobacterium Nopaline synthase
pAct-ChloR-ActT	pCR2.1	Symbiodinium SmicGene17360	Chloramphenicol acetyltransferase	Symbiodinium SmicGene17360
pPsbJ-ChloR	pCR2.1	Symbiodinium SmicGene31399	Chloramphenicol acetyltransferase	(no terminator)
pChlamy3-GenR-GAmCherry	pChlamy_3	Chimeric Chlamydomonas HSP70A/RBCS2	Transposase Tn5 Aminoglycoside 3'-phosphotransferase	(no terminator)
p35S-Neo-2A-eCFP	pChlamy_3	Cauliflower mosaic virus 35S	Transposase Tn5 Aminoglycoside 3'-phosphotransferase	Agrobacterium Nopaline synthase

Using the ten Lohuis and Miller (1998) silicon carbide whiskers agitation protocol, two-week old

Symbiodinium cultures at a cell density of roughly 3×10^5 cells/ml were harvested and transformed with

five different constructs and a "no plasmid" control (see Table 3). Putative CS-153 transformants were

then selected in liquid cultures and on top agar plates with the appropriate antibiotics. In the case of

geneticin selection, we used the same concentration (3 mg/ml) that ten Lohuis and Miller (1998) used.

We did not detect any significant growth in liquid media or visibly growing colonies on selection plates

even after 17 weeks of observation.

Discussion

Our attempts to transform *S. microadriaticum* CCMP2467 stably have not been successful despite the use of a wide variety of constructs and standard transformation methods known to work successfully in the transformation of other microalgae. We have tested both transgenic (p35S) and endogenous promoters (pAct, pBTubA, pBTubB and pHsp90) to drive ChloR expression in the cytoplasm as well as an endogenous promoter of a nuclear-encoded protein targeted to the chloroplasts (pPsbJ). We have also attempted to introduce into *Symbiodinium* chloroplasts artificial minicircles that contain a modified *psbA* gene that is potentially able to confer resistance to atrazine.

In term of transformation methods, we have tried to transform *Symbiodinium* using standard algal transformation techniques such as silicon carbide whisker agitation, biolistics (particle bombardment), electroporation and glass bead agitation. In addition, we used FuGENE transfection media to see if methods more commonly used to transfect animal cells were effective. We did not test *Agrobacterium*-mediated transformation as Ortiz-Matamoros et al. (2015) reported that this method was able to create only transient, rather than stable, transformants that were unable to replicate or maintain normal pigmentation under Basta herbicide selection (Ortiz-Matamoros et al., 2015a).

We also tested if the *Symbiodinium* strain CS-153 was more genetically tractable to the silicon carbide whiskers agitation transformation protocol published by ten Lohuis and Miller (1998). Although both CS-153 and CCMP246 are classified as clade A *Symbiodinium* strains, we do not know the actual genetic distance between the two strains, and they may therefore actually be more akin to separate algal species than strains of the same species. Unfortunately, we did not observe any transformants with CS-153 either, in liquid culture or agar plates. We do acknowledge however that this attempt was only carried out once, and further optimization may be required for successful transformation of CS-153 using silicon carbide whiskers agitation.

Our results illustrate the resistance of *Symbiodinium*, particularly *Symbiodinium microadriaticum*

CCMP2467, to well-established algal transformation methods. Since the first reported publication by

ten Lohuis and Miller (1998), to our knowledge, the only other primary papers concerning

Symbiodinium transformations are those of Ortiz-Matamoros et al. (2015a and 2015b). Ortiz-

Matamoros et al.reported transient expression of GFP in *Symbiodinium* using glass bead

transformations and *Agrobacterium tumefaciens*-assisted glass bead transformations. However, these

methods were associated with loss of photosynthetic pigments and inability for cells to replicate under

Basta herbicide selection. Taken together, these results strongly suggests that this particular algal group

may require novel methods and techniques for successful transformation and editing of the genome.

Materials and Methods

Yeast transformation and Chloramphenicol resistance test

Yeast strain 31019b was kindly provided to us by Prof. Brent N. Kaiser from the University of Sydney.

The S.c. EasyComp™ Transformation Kit (#K505001) (Thermo Fisher Scientific, Waltham,

Massachusetts, US) was used to make transformation competent 31019b cells. The following primers

were used to PCR amplify eGFP and ChloR coding region fragments.

eGFP F1 (5'-TCGACCTGCAGGCGGCCGCAC-3')

eGFP R1 (5'-CTTCTTGTCCATGGTGAAGGGG-3')

ChloR F1 (5'-ATGGAGAAAAAATCACTGGATATACCACCGTTGATATAT-3')

ChloR R1 (5'-TTACGCCCCGCCCTGCCACTCATCGCAGTA-3')

The coding regions were ligated into the yeast expression vector pYES2.1 using a TOPO ligation kit

(#K415001) (Thermo Fisher Scientific, Waltham, Massachusetts, US).

Yeast cells were grown in Yeast Peptone Dextrose medium (#Y1375) (Sigma-Aldrich, St. Louis,

Missouri, US) or Yeast Nitrogenous Base medium (#233520) (BD, Franklin Lakes, New Jersey, US)

with 2% glucose for culture propagation at 30 °C. Experimental cultures were grown in YNB medium

enriched with 2% galactose with various additives such as uracil (final concentration 20 µg/ml) or

chloramphenicol (final concentration 4 mg/ml) depending on the experiment. Yeast growth curve

standards were made using a haemocytometer and measuring the corresponding absorbance value at

OD600 in a 1 cm cuvette using a Nanodrop 200c spectrophotometer (Thermo Fisher Scientific,

Waltham, Massachusetts, US).

Construction of transformation plasmids

Yeast expression plasmids for the chloramphenicol resistance gene function test were synthesized using

the pYES2.1 TOPO TA Yeast Expression Kit. The ChloR gene sequence was amplified from the

pCR2.1-p35S-ChloR-NosT plasmid while the eGFP gene sequence (which was used as a negative

control) was amplified from the pSpCas9n(BB)-2A-GFP (PX461) plasmid from Addgene

(http://www.addgene.org/48140/) (see Table 4).

pCR2.1-based plasmids were synthesized using the pCR2.1 plasmid from the TOPO TA cloning kit as

the vector backbone. The p35S-ChloR-NosT cassette was constructed by amplifying the p35S promoter

and the NosT terminator from the pK7WGF2::hCas9 plasmid from Addgene

(http://www.addgene.org/46965/) and amplifying the chloramphenicol acetyltransferase gene from a

pBC SK+ plasmid. The three fragments were then assembled using assembly PCR to create a p35S-

ChloR-NosT fragment that was subsequently ligated into a pCR2.1 plasmid. Similar methods were used

to construct the various other transformation constructs, although in some cases, Gibson assembly rather

than normal assembly PCR was used. Primers used for these assembly PCRs can be found in Table T7.

PCR for amplification of the *psbA* minicircle was carried out using MasterAmp PCR buffer D

(Epicentre), GoTaq DNA polymerase and primers AIKconstruct2_F and AIKconstruct2_R (see Table

4). PCR cycling conditions were 95 °C 2 minutes 15 seconds, followed by 40 cycles of 95ºC 45 seconds,

57 °C 45 seconds, 72 °C 3 minutes 30 seconds, followed by a final step of 72 °C 10 minutes. Purified

PCR products of the desired size were there ligated using the pGEM-T-Easy Vector System (Promega,

Madison, Wisconsin, US). Correct insertions were selected from the subsequently transformed *E. coli*

clones.

Primers for mutagenesis were designed using the QuikChange Primer Design program. The program

designed a pair of mutagenic primers whose sequences are indicated in Table 4. Mutagenesis reactions

used 125 ng of each primer, 200 µM each NTP, MasterAmp PCR buffer D (Epicentre) and 2.5 U

PfuUltra HF DNA polymerase in a total volume of 50 µl. DNA template concentration was varied and

either 5, 10, 20 or 50 ng of pPsbAGEM DNA was used per reaction. PCR cycling conditions were 95

°C 30 seconds, followed by 16 cycles of 95 °C 30 seconds, 55 °C 1 minute, 72 °C 8 minutes, followed

by a final step of 72 °C 10 minutes. Samples were subsequently treated with 20 U *Dpn*I restriction

enzyme for 1 hour at 37 °C. 5µl of reaction mix was used to transform chemically competent DH5α

cells. Cells were plated out on LB agar plates containing 100 µg/ml ampicillin. Resulting colonies were

picked and cultured. Sequencing of plasmid DNA from these cells revealed the presence of the desired

base changes from the original construct and no other changes.

pChlamy_3-based plasmids conferring geneticin resistance was kindly provided by Dr. Rachel A. Levin

from the School of Biological Earth and Environmental Sciences, University of New South Wales and

Prof. Madeleine van Oppen from the School of BioSciences, University of Melbourne.

Table 4: List of primers used to construct expression cassettes for *Symbiodinium* expression vectors.

Used in plasmid	Name	Function	Sequence	Length
p35S-ChloR-NosT	p35S F1	Amplify p35S	CTAGAGCCAAGCTGATCTCCT	21
p35S-ChloR-NosT	p35S-CAT R1	Amplify p35S	GGTGGTATATCCAGTGATTTTTTTCTCCATACCGGTCGGAGTCCTCTCCAAATGAAATGA	60
p35S-ChloR-NosT	p35S-CAT F2	Amplify ChloR	ATGGAGAAAAAAATCACTGGATAT	24
p35S-ChloR-NosT	CAT-NosT R2	Amplify ChloR	CCAAATGTTTGAACGATCTGCTTGACAAGCCAATTGTTACGCCCCGCCCTGCCACTCATC	60
p35S-ChloR-NosT	CAT-NosT F3	Amplify NosT	GATGAGTGGCAGGGCGGGGCGTAACAATTGGCTTGTCAAGCAGATCGTTCAAACATTTGG	60
p35S-ChloR-NosT	NosT R3	Amplify NosT	TCGATCTAGTAACATAGATG	20
pAct-ChloR-ActT	pActA F1	Amplify pActA	GGAGACCCGGAAACGTGTCG	20
pAct-ChloR-ActT	pActA-CAT R1	Amplify pActA	CCATGTTTAAACACCTTAAGGTCGGGCTAACTCCGAGACACTTCGTGGAA	50
pAct-ChloR-ActT	pActA-CAT F2	Amplify ChloR	TTAGCCCGACCTTAAGGTGTTTAAACATGGAGAAAAAAATCACTGGATAT	50
pAct-ChloR-ActT	CAT-ActT R2	Amplify ChloR	AGACCGTGGGGCGTCGACACGCGTTTACGCCCCGCCCTGCCACTC	45
pAct-ChloR-ActT	CAT-ActT F3	Amplify ActT	CGGGGCGTAAACGCGTGTCGACGCCCCACGGTCTCATTTTGGACTAA	47
pAct-ChloR-ActT	ActT R3	Amplify ActT	ACTACAAGGGCGTTTTTTTTTTTCAGTG	28
pBTubA-ChloR	pBTubA F1	Amplify pBTubA	CTAAATGTCAAGCAACTCAATAT	23
pBTubA-ChloR	pBTubA-CAT R1	Amplify pBTubA	TCCATGTTTAAACCTTAAGGTCGACGGCTGCAAGAAGCGGAAGGTA	46
pBTubA-ChloR	CAT F1	Amplify ChloR	GTCGACCTTAAGGTTTAAACATGGAGAAAAAAATCACTGGATATACCACC	50
pBTubA-ChloR	Chlo R3	Amplify ChloR	TTACGCCCCGCCCTGCCACTCATCGCAGTA	30
pBTubB-ChloR	pBTubB F1	Amplify pBTubB	ACATGGTCATAGACGTTACCTAGCT	25
pBTubB-ChloR	pBTubB-CAT R1	Amplify pBTubB	TCTCCATGTTTAAACCTTAAGGTCGACGGCGGAAGGCTGCAGAAAAAGGA	50
pBTubB-ChloR	CAT F1	Amplify ChloR	GTCGACCTTAAGGTTTAAACATGGAGAAAAAAATCACTGGATATACCACC	50
pBTubB-ChloR	Chlo R3	Amplify ChloR	TTACGCCCCGCCCTGCCACTCATCGCAGTA	30
pHsp90-ChloR-Hsp90T	pHsp90 F1	Amplify pHsp90	GTCGACCTTGCATAGGGGAATCGTACTTCCAA	32
pHsp90-ChloR-Hsp90T	pHsp90-ChloR R1	Amplify pHsp90	CATGTTTAAACCTTAAGGTCGACGGCTCTCTGGACACGCAAGTTTC	46
pHsp90-ChloR-Hsp90T	pHsp90-ChloR F2	Amplify ChloR	GCCGTCGACCTTAAGGTTTAAACATGGAGAAAAAAATCACTGGATATACCAC	52
pHsp90-ChloR-Hsp90T	ChloR-Hsp90T R2	Amplify ChloR	AGCGCTAGCCTTAAGGTTTAAACTTACGCCCCGCCCTGCCA	41
pHsp90-ChloR-Hsp90T	ChloR-Hsp90T F3	Amplify Hsp90T	TAAGTTTAAACCTTAAGGCTAGCGCTGTTTGATGACGGGTGCAGG	45
pHsp90-ChloR-Hsp90T	Hsp90T R3	Amplify Hsp90T	GCTAGCAGGACCCCTGGTCAGTCCCGTT	28
pPsbJ-ChloR	pPsbJ F1	Amplify pPsbJ	GTCGACCTGTCAGCAGCCATTTTGATCACATT	32
pPsbJ-ChloR	pPsbJ-ChloR R1	Amplify pPsbJ	TCCATGTTTAAACCTTAAGGTCGACGCTCACTTGCACCGACCGGCC	46
pPsbJ-ChloR	pPsbJ-ChloR F2	Amplify ChloR	TGAGCGTCGACCTTAAGGTTTAAACATGGAGAAAAAAATCACTGGATATACCA	53
pPsbJ-ChloR	Chlo R3	Amplify ChloR	TTACGCCCCGCCCTGCCACTCATCGCAGTA	30
pYESChloR	Chlo F3	Clone ChloR	ATGGAGAAAAAAATCACTGGATATACCACCGTTGATATAT	40
pYESChloR	Chlo R3	Clone ChloR	TTACGCCCCGCCCTGCCACTCATCGCAGTA	30
pYESeGFP	eGFP clone F1	Clone eGFP	TCGACCTGCAGGCGGCCGCAC	21
pYESeGFP	eGFP R1	Clone eGFP	CTTCTTGTCCATGGTGAAGGGG	22
pPsbAGEM	AIKconstruct2_F	Clone minicircle	GGTCTGGACTTAGTTTATAAGGC	23
pPsbAGEM	AIKconstruct2_R	Clone minicircle	AGACAAGTCTGGTCTGGACG	22
pPsbA^{S262G}GEM	AIKcon2mut1	psbA mutagenesis	GAAGAAGTGTAAACTACGAGAGTTATTGAAGCCAGCATATTGAAAAATTAGTCTGCCAAA	60
pPsbA^{S262G}GEM	AIKcon2mut2	psbA mutagenesis	TTTGGCAGACTAATTTTTCAATATGCTGGCTTCAATAACTCTCGTAGTTTACACTTCTTC	60

Symbiodinium cell culture conditions

Cultures of the dinoflagellate *Symbiodinium microadriaticum* (strain CCMP2467) were obtained from the Bigelow National Center for Marine Algae and Microbiota (NCMA). This strain was originally isolated from a scleractinian coral, *Stylophora pistillata*, in the Gulf of Aqaba (LaJeunesse, 2001; Lajeunesse et al., 2015). Stock cultures were grown in Percival incubators under a 12:12 day:night regiment in f/2 medium in Nunc cell culture-treated TripleFlasks (132913) (ThermoFisher Scientific, Waltham, MA, USA) without shaking. Growth conditions were set at a Photosynthetic Photon Flux Density of 80 µmol photons m^{-2} s^{-1}, growth temperature of 26 °C and growth media salinity of 40 ppt (i.e. the salinity of seawater from the Red Sea). These growth conditions will be referred to henceforth as standard culture conditions.

Cultures of the dinoflagellate *Symbiodinium microadriaticum* (strain CS-153) were obtained from the

Australian National Algae Culture Collection (ANACC) in Hobart, Tasmania. This strain was originally

isolated from a jellyfish, *Cassiopeia xamachana*, from off the coast of Florida, United States of America

(ANACC database). Stock cultures were grown in a New Brunswick Innova 4340 incubator under a

14:10 day:night regiment in f/2 medium in Pyrex conical flasks (Corning Incorporated, Corning, NY,

USA) without shaking. Growth conditions were set at a Photosynthetic Photon Flux Density of 40

µmol photons m^{-2} s^{-1}, growth temperature of 26 °C and growth medium salinity of 34 ppt using

Ultramarine synthetic sea salt (Waterlife Research Industries Ltd, Waterlife Research Industries Ltd,

Middlesex, United Kingdom).

Transformation methods

Silicon carbide whiskers agitation transformation

This protocol was modified from Ten Lohuis and Miller (1998) (ten Lohuis and Miller, 1998).

Approximately 5×10^7 cells were harvested by centrifugation at 3000g for 5 minutes, washed with 5

ml of f/2 medium, repelleted, and then resuspended in 500 µl of f/2 medium. A transformation mixture

containing 40 µl of 50 µg/ml Silar silicon carbide whiskers (997002-5g) (Haydale Technologies

Incorporated, Greer, South Carolina, USA) was sequentially mixed with 20 to 40 µg of plasmid (circular

or linear), 160 µl of PEG8000 (20% wv; filter sterilized) and then f/2 medium was added to a final

volume of 250 µl. For negative DNA controls, plasmids were omitted from the transformation mix.

The 500 µl resuspended cells were then added to the transformation mixture and vortexed over a period

of 2 minutes, pausing for 5 seconds every 10 seconds. 2 ml of f/2 medium with 100 µg/ml of

carbenicillin was added to the transformation mixture. The mixture was then incubated in the dark at

27 °C for 1 to 2 days. The agitated *Symbiodinium* were then grown under selection in either liquid

culture (1 ml of transformant culture in 150 ml of liquid f/2 medium) or on agar plates (150 µl of

transformant culture per agar plate).

 Biolistics

The biolistics protocol was modified from instructions given in the Biorad PDS-1000/He Biolistic

Particle Delivery System manual. Approximately 5×10^7 Symbiodinium cells were harvested by

centrifugation at 3000g for 5 minutes and subsequently washed with 5 ml of filtered, sterilized f/2

medium before being plated onto an f/2 agar plate approximately 1 hour before transformation.

Tungsten or gold microcarriers suspended in 50% glycerol (30 mg/ml) were vortexed for 5 minutes

before a 50 µl aliquot of the mix was placed into a 1.5 ml Eppendorf tube. 5 µg of plasmid DNA was

added to the microcarrier mix (omitted for the negative DNA control) and immediately vortexed for 10

seconds. This was followed by the addition of 50 µl of 2.5 M CaCl₂, vortexing for 10 seconds, and the

addition of 20 µl 0.1 M spermidine. The mixture was then mixed by vortexing for 2 minutes before

being allowed to settle for 1 minute. The microcarriers were then pelleted by pulse centrifugation and

the liquid supernatant was discarded. The pellet was gently washed with 140 µl of 70% ethanol before

the mix was pulse centrifuged and the supernatant was removed. This washing step was repeated with

140 µl of 100% ethanol. The pellet was then re-suspended with 10 or 70 µl of 100% ethanol depending

on the number of macrocarriers being used. 10 µl of the mix was used to dip dry a layer of microcarriers

onto a Bio-Rad biolistics macrocarrier (#1652335) (Bio-Rad, Hercules, California, US) and this step

was repeated seven times when using 7 macrocarriers. The Biorad macrocarrier was then loaded into a

macrocarrier holder and fired at an agar plate of *Symbiodinium* cells according to the manufacturer's

manual for the Biorad PDS-1000/He Biolistic Particle Delivery System. Chamber pressure, rupture disk

pressure and distance of agar plate from the macrocarrier holder were adjusted based on the experiments

carried out as described in the Results section.

After the biolistic bombardment, *Symbiodinium* cells were transferred from the agar plate into 5 ml of

f/2 medium and incubated under standard culture conditions for one day. The transformed

Symbiodinium were then grown under selection in liquid culture (2 ml of transformant culture in 150

ml of liquid f/2 medium) and/or on agar plates (150 µl of transformant culture per agar plate).

Electroporation

The electroporation protocol used in this publication was modified from published protocols used for *Phaeodactylum tricornutum* and *Nannochloropsis* sp. (Kilian et al., 2011; Zhang and Hu, 2014). Approximately 1×10^8 cells were harvested by centrifugation at $3000g$ for 5 minutes. The cells were subsequently washed with 5 ml of f/2 medium and repelleted, before being washed with 1 ml of 375 mM sorbitol, repelleted, and finally resuspended in 100 µl of 375 mM sorbitol. The suspension was then mixed with 2-4 µg of plasmid DNA (or just 5 µl of water for the no plasmid control) and 40 µg of denatured salmon sperm DNA. The mixture was then incubated on ice for 10 minutes before being transferred into a 2 mm electroporation cuvette. Electroporation was performed using a Bio-Rad Gene Pulser Xcell Electroporation system (Biorad, Hercules, California, US). The parameters of the system (exponential decay vs multiple pulse, field strength, capacitance, shunt resistance) were adjusted according the experiments carried out as described in the Results section.

After electroporation, the cells were immediately transferred to a 15 ml Falcon tube containing 10 ml of f/2 medium with 100 µg/ml carbenicillin and left under standard cultures conditions for one day. After this, the cells were then collected by centrifugation at 1500 g for 10 minutes and resuspended in 3 ml of f/2 medium. The electroporated *Symbiodinium* were then grown under selection in liquid culture (1 ml of transformant culture in 150 ml of liquid f/2 medium) and/or on agar plates (150 µl of transformant culture per agar plate).

Glass bead agitation transformation, with cell wall digest step

The glass bead protocol was modified from Kindle (1990) and the cellulase cell wall digest was modified from Levin et al. (2017). Approximately 2×10^6 cells were harvested by centrifugation at $3000g$ for 5 minutes, washed with 5 ml of f/2 medium, repelleted and then resuspended in 1 ml of digestion solution (0.5 M D-sorbitol in filtered autoclaved seawater). Either 0.3 kilounits (KU) of cellulase from *Trichoderma* sp. (#C1794) (Sigma-Aldrich, St. Louis, Missouri, US) or 10 mg of

Snailase (S0100, Beijing Biodee Biotechnology Co., Ltd, Beijing, China) was added to the digest mix.

The mixture was then incubated in the dark at 30 °C for 1 day on a tube rotator.

The digest mixture was then centrifuged at 800g for 5 minutes and the supernatant was discarded. The

pellet was resuspended in 1 ml of protoplast wash solution (0.5 M D-sorbitol, 0.5 M sucrose, 25 mM

CaCl$_2$, and 100 mg/ml kanamycin in filtered autoclaved seawater) before being incubated at 25 °C for

3 hours in the dark on a tube rotator. The cells were centrifuged at 800g for 5 minutes before being

washed with 1 ml of f/2 medium, repelleted, and resuspended in 300 µl of f/2 medium.

100 µl of filter sterilized 20% w/v PEG 6000 (81260, Sigma) was added to the 300 µl of resuspended

cells followed by the addition of 50 µg of salmon sperm carrier DNA (#15632011) (Invitrogen,

Carlsbad, California, US) and 2 µg of plasmid DNA (in various combinations of circular and linearized

plasmids, as described in the results section). 300 µg of autoclave-sterilized 0.5mm glass beads

(#11079105) (BioSpec, Bartlesville, Oklahoma, US) were added to the transformation mixture. The

mixture was vortexed for 30 seconds after which the beads were allowed to settle. The agitated

Symbiodinium were then resuspended in 2 ml of fresh f/2 with 100 µg/ml of carbenicillin. After this,

the transformant cultures were grown under selection in liquid medium (1 ml of transformant culture in

150 ml of liquid f/2 medium) and on agar plates plates (150 µl of transformant culture per agar plate).

FuGENE HD transfection

Approximately 5×10^6 *Symbiodinium* cells were harvested by centrifugation at 3000g for 5 minutes,

washed with 5ml of f/2 medium, repelleted and resuspended in 150 µl of f/2 medium. A 30 µl FuGENE

transfection/DNA mixture was made using f/2 medium with different mixtures of plasmid DNA and

FuGENE HD transfection reagent as according to Supplementary Table 4E. 1 to 30 µl of this mix was

then added to the 150 µl cell suspension and incubated at standard growth conditions for 24 hours. The

cells were then grown under selection in liquid culture (100 µl of transformant culture per 150 ml of

liquid f/2 medium).

f/2 agar plate selection

1.5 % microbiological agar (Fisher Scientific, New Hampshire, US) made up with filtered, autoclaved

seawater enriched with f/2 medium was used to make agar plates for transformation selection. 100

µg/ml chloramphenicol or 200 to 1000 ng/ml atrazine was used as the selection antibiotic/herbicide

depending on the construct being tested. Control plates with no selection antibiotic/herbicide were also

made to confirm that the transformation method was mild enough that some cells survived the treatment.

The samples were inoculated onto the agar plate using 3 ml of top agar made from 0.8 % plant agar

(Duchefa Biochemie, Haarlem, Netherlands) in f/2 medium with no antibiotics. This was done due to

the tendency of *Symbiodinium* cells to clump, which prevented the cells from being easily evenly spread

across agar surfaces. The plates were then grown in a LMS vertical incubator set to 26 °C with a 14:10

hours day:night cycle and 40 µmol photons m^{-2} s^{-1} light intensity for up to four months.

Liquid f/2 selection

536   Transformant cultures were grown in 225 cm^2 EasYFlask Nunc bottles (#159934) (Thermo Scientific,

537   Waltham, Massachusetts, US) filled with 150 ml of filtered, autoclaved seawater enriched with f/2

538   medium. The selection antibiotic/herbicide used was either 100 µg/ml chloramphenicol or 150 ng/ml

539   atrazine and the cultures were kept under standard growth conditions.

Acknowledgements

The authors would like to thank Dr. Rachel A. Levin and Prof. Madeleine van Oppen (School of Biological Earth and Environmental Sciences, The University of New South Wales) for the geneticin constructs and for their valuable feedback during the course of this project.

This publication is based upon work supported by the King Abdullah University of Science and Technology (KAUST) Office of Sponsored Research (OSR) under Awards No. URF/1/1705-01 and OCRF-2014-CRG3-2216.

References

Aranda M, Li Y, Liew YJ, Baumgarten S, Simakov O, Wilson MC, Piel J, Ashoor H, Bougouffa S, Bajic VB, Ryu T, Ravasi T, Bayer T, Micklem G, Kim H, Bhak J, LaJeunesse TC, Voolstra CR (2016) Genomes of coral dinoflagellate symbionts highlight evolutionary adaptations conducive to a symbiotic lifestyle. Sci Rep-Uk **6**: 39734.

Brown BE (1997) Coral bleaching: causes and consequences. Coral Reefs **16**: S129-S138.

Chen JE, Cui G, Wang X, Liew YJ, Aranda M (2017) Recent expansion of heat-activated retrotransposons in the coral symbiont Symbiodinium microadriaticum. The ISME journal.

Dong X, Stothard P, Forsythe IJ, Wishart DS (2004) PlasMapper: a web server for drawing and auto-annotating plasmid maps. Nucleic Acids Res **32**: W660-664.

Howe CJ, Nisbet RE, Barbrook AC (2008) The remarkable chloroplast genome of dinoflagellates. J Exp Bot **59**: 1035-1045.

Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, van Oppen MJH (2012) Coral thermal tolerance shaped by local adaptation of photosymbionts. Nat Clim Change **2**: 116-120.

Hughes TP, Kerry JT, Alvarez-Noriega M, Alvarez-Romero JG, Anderson KD, Baird AH, Babcock RC, Beger M, Bellwood DR, Berkelmans R, Bridge TC, Butler IR, Byrne M, Cantin NE, Comeau S, Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF, Gilmour JP, Harrison HB, Heron SF, Hoey AS, Hobbs JA, Hoogenboom MO, Kennedy EV, Kuo CY, Lough JM, Lowe RJ, Liu G, McCulloch MT, Malcolm HA, McWilliam MJ, Pandolfi JM, Pears RJ, Pratchett MS, Schoepf V, Simpson T, Skirving WJ, Sommer B, Torda G, Wachenfeld DR, Willis BL, Wilson SK (2017) Global warming and recurrent mass bleaching of corals. Nature **543**: 373-377.

587
588 **Kilian O, Benemann CSE, Niyogi KK, Vick B** (2011) High-efficiency homologous
589 recombination in the oil-producing alga Nannochloropsis sp. P Natl Acad Sci USA **108**:
590 21265-21269.

591
592 **Kindle KL** (1990) High-frequency nuclear transformation of Chlamydomonas reinhardtii. P
593 Natl Acad Sci USA **87**: 1228-1232.

594
595 **LaJeunesse TC** (2001) Investigating the biodiversity, ecology, and phylogeny of
596 endosymbiotic dinoflagellates in the genus Symbiodinium using the its region: In search of a
597 "species" level marker. J Phycol **37**: 866-880.

598
599 **Lajeunesse TC, Lee SY, Gil-Agudelo DL, Knowlton N, Jeong HJ** (2015) Symbiodinium
600 necroappetens sp nov (Dinophyceae): an opportunist 'zooxanthella' found in bleached and
601 diseased tissues of Caribbean reef corals. Eur J Phycol **50**: 223-238.

602
603 **Levin RA, Suggett DJ, Nitschke MR, van Oppen MJ, Steinberg PD** (2017) Expanding the
604 Symbiodinium (Dinophyceae, Suessiales) Toolkit Through Protoplast Technology. J
605 Eukaryot Microbiol.

606
607 **Lin SJ, Cheng SF, Song B, Zhong X, Lin X, Li WJ, Li L, Zhang YQ, Zhang H, Ji ZL, Cai**
608 **MC, Zhuang YY, Shi XG, Lin LX, Wang L, Wang ZB, Liu X, Yu S, Zeng P, Hao H, Zou Q,**
609 **Chen CX, Li YJ, Wang Y, Xu CY, Meng SS, Xu X, Wang J, Yang HM, Campbell DA,**
610 **Sturm NR, Dagenais-Bellefeuille S, Morse D** (2015) The Symbiodinium kawagutii genome
611 illuminates dinoflagellate gene expression and coral symbiosis. Science **350**: 691-694.

612
613 **Muscatine L, Mccloskey LR, Marian RE** (1981) Estimating the daily contribution of carbon
614 from zooxanthellae to coral animal respiration. Limnol Oceanogr **26**: 601-611.

615
616 **Muscatine L, Porter JW** (1977) Reef corals - mutualistic symbioses adapted to nutrient-
617 poor environments. Bioscience **27**: 454-460.

618
619 **Ortiz-Matamoros MF, Islas-Flores T, Voigt B, Menzel D, Baluska F, Villanueva MA**
620 (2015a) Heterologous DNA Uptake in Cultured Symbiodinium spp. Aided by Agrobacterium
621 tumefaciens. Plos One **10**: e0132693.

622
623 **Ortiz-Matamoros MF, Villanueva MA, Islas-Flores T** (2015b) Transient transformation of
624 cultured photosynthetic dinoflagellates (Symbiodinium spp.) with plant-targeted vectors.
625 Cienc Mar **41**: 21-32.

626
627 **Smith DM, Cusack S, Colman AW, Folland CK, Harris GR, Murphy JM** (2007) Improved
628 surface temperature prediction for the coming decade from a global climate model. Science
629 **317**: 796-799.

630
631 **ten Lohuis MR, Miller DJ** (1998) Genetic transformation of dinoflagellates (Amphidinium
632 and Symbiodinium): expression of GUS in microalgae using heterologous promoter
633 constructs. Plant Journal **13**: 427-435.

634
635 **Zhang CY, Hu HH** (2014) High-efficiency nuclear transformation of the diatom
636 Phaeodactylum tricornutum by electroporation. Mar Genom **16**: 63-66.

637
638 **Zhang H, Zhuang YY, Gill J, Lin SJ** (2013) Proof that Dinoflagellate Spliced Leader
639 (DinoSL) is a Useful Hook for Fishing Dinoflagellate Transcripts from Mixed Microbial
640 Samples: Symbiodinium kawagutii as a Case Study. Protist **164**: 510-527.

Supporting information

Supplementary Figure 1: Verification of pPsbJ 5' UTR sequence via RACE.

Supplementary Table 1: Price of antibiotics from various major suppliers, taken on 30[th] of April 2017.

Supplementary Table 2: Growth curves of *Symbiodinium microadriaticum* under Atrazine treatment.

Supplementary Table 3: Expression of selected genes from transcriptomics analysis published in Chen et al. (2017).

Supplementary Table 4: Detailed list of transformations carried out, with information on transformation conditions and materials used.

Supplementary Data 1: Vector sequence for plasmids pYESChloR.

Supplementary Data 2: Vector sequence for plasmid pYESeGFP.

Supplementary Data 3: Genomic sequence of putative β-Tubulin A from Smic.scaffold612 published in Aranda et al. (2016).

Supplementary Data 4: Genomic sequence of putative β-Tubulin B from Smic.scaffold51 published in Aranda et al. (2016).

Supplementary Data 5: Genomic sequence of putative Hsp90 from Smic.scaffold975 published in Aranda et al. (2016).

Supplementary Data 6: Vector sequence for plasmid pCR4 p35S-ChloR-NosT.

Supplementary Data 7: Vector sequence for plasmid pCR2.1 pAct-ChloR-ActT.

Supplementary Data 8: Vector sequence for plasmid pCR2.1 pBtubA-ChloR.

Supplementary Data 9: Vector sequence for plasmid pCR2.1 pBtubB-ChloR.

Supplementary Data 10: Vector sequence for plasmid pCR2.1 pHsp90-ChloR-Hsp90T.

Supplementary Data 11: Vector sequence for plasmid pCR2.1 pPsbJ-ChloR.

Supplementary Data 12: Vector sequence for plasmid pPsbA^{S262G}GEM. Atrazine resistance is conferred

by mutation of nucleotide 790 T->G and 791 C->G, causing a 262 Ser->Gly mutation. A silent mutation

for identification purposes was also added to 795 T->C.

Supplementary Data 13: Vector sequence for plasmid pPsbAGEM.

Supplementary Data 14: Vector sequence for plasmid pChlamy3-GenR-GAmCherry.

Supplementary Data 15: Vector sequence for plasmid p35S-GenR-eCFP-NosT.