Undocumented potential for primary productivity in a globally-distributed bacterial photoautotroph

Authors: E.D. Graham[1], J.F. Heidelberg[1,2], B.J. Tully[*1,2]

Author Affiliations:

[1]Department of Biological Sciences, University of Southern California, Los Angeles, CA, USA

[2]Center for Dark Energy Biosphere Investigations, University of Southern California, Los Angeles, CA, USA

*Corresponding Author: Benjamin Tully, tully.bj@gmail.com

Keywords: autotrophy; marine carbon cycle; metagenomics; Alphaproteobacteria; aerobic anoxygenic phototrophs

Abstract: Aerobic anoxygenic phototrophs (AAnPs) are common in the global oceans and are
associated with photoheterotrophic activity. To date, AAnPs have not been identified in the
surface ocean that possess the potential for carbon fixation. Using the *Tara Oceans* metagenomic
dataset, we have reconstructed draft genomes of four bacteria that possess the genomic potential
for anoxygenic phototrophy, carbon fixation via the Calvin-Benson-Bassham cycle, and the
oxidation of sulfite and thiosulfate. Forming a monophyletic clade within the
Alphaproteobacteria and lacking cultured representatives, the organisms compose minor
constituents of local microbial communities (0.1-1.0%), but are globally distributed, present in
multiple samples from the North Pacific, Mediterranean Sea, the East Africa Coastal Province,
and the South Atlantic. These organisms represent a shift in our understanding of microbially-
mediated photoautotrophy in the global oceans and provide a previously undiscovered route of
primary productivity.

24
Significance Statement: In examining the genomic content of organisms collected during the
Tara Oceans expedition, we have identified a novel clade within the *Alphaproteobacteria* that
has the potential for photoautotrophy. Based on genome observations, these organisms have the
potential to couple inorganic sulfur compounds as electron donors to fix carbon into biomass.
They are globally distributed, present in samples from the North Pacific, Mediterranean Sea, East
Africa Coastal Current, and the South Atlantic. This discovery may require re-examination of the
microbial communities in the global ocean to understand and constrain the impacts of this group
of organisms on the global carbon cycle.

Introduction

It has been understood for decades that the basis of the global marine carbon cycle are oxygenic photoautotrophs that perform photoautotrophic processes. Two additional phototrophic processes are common in the ocean and are mediated by proteorhodopsin-containing microorganisms and aerobic anoxygenic phototrophs (AAnPs). Proteorhodopsins and AAnPs have historically been associated with photoheterotrophy[1,2], a process that supplements additional energy to microorganisms beyond what is obtained as part of a heterotrophic metabolic strategy. AAnPs utilize type-II photochemical reaction centers (RCIIs) and bacteriochlorophyll (BChl), are globally distributed[3], and have been identified in phylogenetically diverse groups of microorganisms[4-6]. Though anaerobic microorganisms with RCIIs and BChl are known to fix CO_2[7] and marine AAnPs can incorporate inorganic carbon via anaplerotic reactions[8], marine AAnPs have not been linked to carbon fixation[9]. The identification of marine AAnPs capable of carbon fixation adds to our understanding of microbial photosynthesis in the global oceans and represents a previously undiscovered route of photoautotrophy.

The *Tara Oceans* expedition generated microbial metagenomes during a circumnavigation of the global oceans[10,11]. *Tara Oceans* samples were collected from 63 sites in 10 major ocean provinces, with most sites contributing multiple sampling depths (generally, surface, deep chlorophyll maximum [DCM], and mesopelagic) and multiple size fractions (generally, 'viral', 'girus' [giant virus], 'bacterial', and 'protistan') from each depth (Supplementary Data File 1). We independently assembled each sample and assemblies from all samples within a province were combined and subjected to binning techniques to reconstruct microbial genomes (Fig. 1 and Extended Data Fig. 1). Microbial genomes reconstructed from eight of ten provinces (Mediterranean, Red Sea, Arabian Sea, Indian Monsoon Gyre, East Africa Coastal, South Atlantic, and North Pacific; 36 sites, 134 samples) were annotated using the KEGG Ontology (KO) system[12] and examined for the genes and pathways of interest.

Results and Discussion

From 1,774 metagenome-assembled genomes (MAGs), 53 genomes possessed the genes encoding the core subunits of RCIIs (PufLM). Of those 53, four genomes (MED800, EAC638, SAT68, NP970; Fig. 1) also contained genes for ribulose-1,5-bisphosphate carboxylase (Rubisco; RbsLS; Fig. 2). Rubisco has four major forms, of which three (Types I, II, and III) have been shown to fix CO_2 and two are known to participate in the Calvin-Benson-Bassham (CBB) cycle (Types I and II[13,14]. Phylogenetic placement of the Rubisco large subunits recovered from the genomes revealed them to be of the Type IC/D subgroup[13], suggesting that the identified proteins represent *bona fide* Rubiscos capable of carbon fixation (Fig. 2). Within the Type IC/D subgroup, the Rubisco sequences from the four analyzed genomes formed a distinct cluster with environmental sequences derived from the Global Ocean Survey (GOS) metagenomes[15,16], but lacking sequences from reference organisms.

Similarly, the PufM sequences from the analyzed *Tara* genomes did not cluster with reference sequences, instead grouping with sequences from the GOS metagenomes[15]. Sequences from MED800, SAT68, and NP970 were group together in one cluster, while EAC638 was located in a separate cluster (Fig. 3). The MED800/SAT68/NP970 clade is basal to the previously identified phylogroups E and F, while the EAC638 clade is basal to the *Roseobacter*-related phylogroup G[17]. As MED800, EAC638, SAT68, and NP970 branch in distinct clades on

both the RbsL and PufM trees that consist of entirely environmental sequences, it may be
possible that these clades represent a phylogenetically coherent group of organisms with the
potential for both phototrophy and carbon fixation.

The draft genomes were of high enough quality (66-85% complete; <5.5% duplication;
Table 1) to possess sufficient phylogenetic markers for accurate placement (Extended Data Table
1). The four organisms form a monophyletic clade basal to the Family *Rhodobacteraceae* (Fig.
4). The relationship between the genomes would suggest that NP970, SAT68, and MED800 are
phylogenetically more closely related to each other than either are to EAC638. As is common
with assembled metagenomic sequences, the recovered genomes lack a distinguishable 16S
rRNA gene sequence. However, based on the observed phylogenetic distance in the concatenated
marker tree, we suggest that these organisms represent a new clade within the
Rhodobacteraceae, and possibly a family-level clade previously without a reference sequence
within the *Alphaproteobacteria*. We propose that NP970, SAT68, and MED800 represent three
species within the same genus (tentatively named, '*Candidatus Luxescamonas taraoceani*'), with
EAC638 as a representative of a species in a sister genus (tentatively named, '*Candidatus*
Luxescabacter africus').

In addition to Rubisco, all four genomes contained genes encoding phosphoribulokinase,
an essential gene of the CBB cycle, and 50-89% of the genes necessary to perform complete
carbon fixation (Fig. 5). The RCII genes in MED800, EAC638, SAT68, and NP970 were
accompanied by bacteriochlorophyll biosynthesis and light-harvesting genes (Supplementary
Data File 2). This complement of reaction center, bacteriochlorophyll biosynthesis, and essential
carbon fixation genes support a role for autotrophy within these organisms beyond the identified
marker genes.

All four genomes possessed ATP-binding cassette (ABC) type transporters for
spermidine/putrescine and L- and branched-chain amino acids. These transporters are indicative
of the utilization of organic nitrogen compounds, as spermidine and putrescine are nitrogen rich
organic compounds, while the scavenging of amino acids reduces the overall nitrogen demands
of the cell. Further, the genomes lacked transporters and degradation enzymes for many of the
saccharides common in the marine environment[18] (Fig. 5). However, MED800, SAT68, and
NP970 possessed an ABC-type α-glucoside transporter, an annotated β-glucosidase in SAT68,
and D-xylose and D-ribose ABC-type transporters in EAC638. While autotrophs are generally
considered to not require external sources of organic carbon, saccharide transporters are
commonly observed in classical photoautotrophic organisms[19,20], including the specific example
of α-glucoside transporters in strains of *Synechocystis*[21]. For the four genomes, the minimal
number of carbon transport and degradation genes may suggest that the organisms have a limited
capacity to utilize dissolved organic carbon compounds, but are capable of heterotrophic growth
under certain conditions. As such, the genomic potential of these organisms suggest that NP970,
SAT68, MED800, and EAC638 are likely facultative autotrophs or mixotrophs.

In oxygenic photosynthesis, electrons are donated as a result of the oxidation of water.
Lacking photosystem II, organisms with RCII are incapable of oxidizing water and would
require an alternative electron donor for autotrophic processes. EAC638 and SAT68 contained
the full/partial suite of genes necessary for thiosulfate oxidation, while MED800, NP970, and
SAT68 possessed the genes for oxidizing sulfite. The oxidation of sulfur compounds has
previously been linked to autotrophy in the marine environment[22]. The oxidation of organic
sulfur compounds, like dimethyl sulfide, has been shown to be a source of thiosulfate[23] and

sulfite[24] in the marine environment. Electrons derived from inorganic sulfur sources (thiosulfate
and/or sulfite) could be transferred directly through cytochromes or membrane-bound quinone
dehydrogenases to the electron transport chain. The cycling of electrons through the reaction
centers could generate the proton motive force necessary to generate NADH via reverse electron
flow[25], convert NADH to NADPH via transhydrogenase, and generate ATP for the CBB cycle
(Fig. 5).

The reconstruction of four genomes from the same novel family in four different
provinces (North Pacific, Mediterranean, East Africa coastal current, and South Atlantic)
suggests that the observed genomes represent an *in situ* microbial population from the surface
marine environment. Each of the genomes recruit metagenomic reads from multiple sampling
sites in each province and are present at >0.1% of the microbial relative abundance (range: 0.1-
1.04%; mean: 0.286%) in 20 samples (Fig. 1; Supplemental Information 1). Predominantly, the
genomes were present in samples are located in the surface (n = 5) or DCM (n = 12). These
organisms were collected at depths where light was available for photosynthesis and less
frequently identified at deeper depths (n = 3 mesopelagic samples). When >0.1% relative
abundance, the genomes tend to be more abundant in the 'bacterial'/'girus' size fraction (n = 14),
though were also observed in 'protistan' (n = 4), and 'viral' (n = 2) size fractions (Supplemental
Information 1). The nature of the 'bacterial' size fractions suggests that these organisms are
generally not particle attached and <1.6μm in size. The occurrence in the protistan fraction may
be due to slightly larger cells or attachment to particles, but this data are difficult to interpret as
the 'protistan' and 'bacterial' size fractions can overlap (0.8-1.6μm). As members of the free-
living bacterioplankton, these organisms should be poised to grow in aerobic conditions. All four
genomes possessed the genes encoding for cytochromes involved in aerobic metabolisms (aa_3-
and bc_1-type), and lacked the genes for cytochromes involved in microaerobic metabolisms and
alternative electron acceptors. Further, all four genomes encoded the gene for an oxygen-
dependent ring cyclase (*acsF*), a necessary component in bacteriochlorophyll biosynthesis for
which there is alternative that is oxygen-independent (*bchE*) and used by anaerobic organisms.

With this discovery, the potential photosynthesis in the ocean has expanded beyond
organisms harboring chlorophyll *a* to include *Alphaproteobacteria* with BChl *a*. Though these
organisms have not been cultivated or sequenced before, both PufM and Rubisco in MED800,
EAC638, NP970, and SAT68 are phylogenetically related to environmentally-derived protein
sequences, lending credence to the fact that these organisms may be a persistent element of
oceanic carbon fixation. As such, clades of environmentally sampled genes (*rbsL* and *pufM*) can
now be linked to a previously unrecognized source of marine primary productivity. The
identification of a globally distributed clade of AAnPs in the ocean capable of carbon fixation
continues to expand our understanding of photosynthesis and the marine carbon cycle.

Materials and Methods:

Assembly

All sequences for the reverse and forward reads from each sampled station and depth within the
Tara Oceans dataset were accessed from European Molecular Biology Laboratory (EMBL)[10,11].
Typically, *Tara* sampling sites have multiple metagenomic samples, representing different
sampling depths and size fractions. The common size fractions were used during sampling were:
'bacterial' (0.22-1.6μm) (includes Mediterranean 'girus' samples), 'protistan' (0.8-5.0μm),
'girus' (0.45-0.8μm) and 'viral' (<0.22μm). Surface samples were collected at ~5-m depth, while

deep chlorophyll maximum (DCM) and mesopelagic depths were variable depending the
physiochemical features of the site. Paired-end reads from different filter sizes from each site and
depth (e.g., TARA0007, girus filter fraction, sampled at the DCM) were assembled using
Megahit[26] (v1.0.3; parameters: --preset, meta-sensitive) (Supplementary Data File 1). All of the
Megahit assemblies from each province were pooled in to two tranches based on assembly size,
<2kb and ≥2kb. Longer assemblies (≥2kb) with ≥99% semi-global identity were combined using
CD-HIT-EST[27] (v4.6; -T 90 -M 500000 -c 0.99 -n 10). The reduced set of contiguous DNA
fragments (contigs) ≥2kb was then cross-assembled using Minimus2[28] (AMOS v3.1.0;
parameters: -D OVERLAP=100 MINID=95).

Binning

Contigs from each province were initially clustered into tentative genomic bins using
BinSanity[29]. Due to computational limitations, the South Atlantic, East African Coastal province,
and Mediterranean Sea were initially run with contig size cutoffs of 11.5kbp, 7.5kbp, and 7kbp,
respectively. The BinSanity workflow was run iteratively three times using variable preference
values (v.0.2.5.5; parameters: -p [(1) -10, (2) -5, (3) -3] -m 4000 -v 400 -d 0.95). Between each
of the three main clustering steps, refinement was performed based on sequence composition
(parameters: -p [(1) -25, (2) -10, (3) -3] -m 4000 -v 400 -d 0.95 -kmer 4). After refinement and
before the next pass with BinSanity, bins were evaluated using CheckM[30] (v.1.0.3; parameters:
lineage_wf, default settings) for completion and redundancy. Genomes were considered for
further analysis based on the completeness and contamination metrics. The cutoff values were:
>90% complete with <10% contamination, 80-90% complete with <5% contamination, or 50-
80% complete with <2% contamination. Bins meeting these metrics were reclassified as draft
genomes were removed from subsequent rounds of clustering. After identification of the four
genomes of interest (initially 51-84% complete, <7.0% contamination), binning was performed
with CONCOCT[31] (v.0.4.1; parameters: -c 800 -I 500) on contigs >5kb from each province that
had a produced a genome of interest. To improve completion estimates, overlapping CONCOCT
and BinSanity bins were visualized using Anvi'o[32] (v.2.1.0) and manually refined to improve
genome completion and minimize contamination estimates (Extended Data Fig. 3-6).

Annotation

Putative DNA coding sequences (CDSs) were predicted for each genome using Prodigal[33]
(v.2.6.2; -m -p meta). Putative CDS were submitted for annotation by the KEGG database using
BlastKOALA[12] (taxonomy group, Prokaryotes; database, genus_prokaryotes +
family_eukaryotes; Accessed March 2017) (Supplementary Data File Table 3). Assessment of
pathways and metabolisms of interest were determined using the script KEGG-decoder.py
(www.github.com/bjtully/BioData/tree/master/KEGGDecoder). Genomes of interested were
determined based on the presence of genes assigned as the M subunit of type-II photochemical
reaction center (PufM) and ribulose-1,5-bisphosphate carboxylase (RbsLS). After confirmation
of the genes of interest (see below), additional annotations were performed for the genomes
using the Rapid Annotation using Subsystem Technology (RAST) service (Classic RAST default
parameters - Release70)[34].

Phylogeny

An initial assessment of phylogeny was conducted using pplacer[35] within CheckM. The
Prodigal-derived CDSs were searched for a collection of single-copy marker genes that was
common to all four *Tara* assembled genomes using hidden Markov models collected from the

Pfam database[36] (Accessed March 2017) and HMMER [37] (v3.1b2; parameters: hmmsearch -E
1e-10 --notextw). 17 marker genes were identified that met this criteria[38-40] (Extended Data
Table 1). The 17 markers were identified in 2,889 reference genes from complete and partial
genomes accessed from NCBI Genbank[41] (Supplementary Data File 4). If a genome contained
multiple copies of a single marker gene both were excluded from the final tree. Only genomes
containing ≥10 markers were used for phylogenetic placement. Each marker set was aligned
using MUSCLE[42] (v3.8.31; parameter: -maxiters 8) and trimmed using TrimAL[43] (v.1.2rev59;
parameter: -automated1). Alignments were then manually assessed and concatenated in
Geneious[44]. An approximate maximum likelihood tree was generated using FastTree[45] (v.2.1.10;
parameters: -lg -gamma; Supplementary Data File 5). A simplified version of this phylogenetic
tree was constructed using the same protocol, but with 160 reference genomes for Fig. 2
(Supplementary Data File 6 and 7).

Phylogenetic tree – Rubisco and Type-II Reaction Center

RbsL and PufM sequences representing previously described lineages were collected[13,17]
(Supplementary Data File 8 and 9). Additional reference PufM sequences were collected from
environmentally generated bacterial artificial chromosomes[5] and Integrated Microbial Genomes
(IMG; Accessed Feb 2017)[46]. Protein sequences from IMG were assessed based on genomes
with KEGG Ontology (KO) annotations[47] for the reaction center subunit M (K08929). PufM
sequences from Prodigal predicted CDS (as above) of Global Ocean Survey (GOS) assemblies[16]
were identified using DIAMOND[48] (v.0.8.36.98; parameters: BLASTP, default settings), where
all reference and *Tara* genome sequences were used as a query. Two separate phylogenetic trees
were constructed (RbsL and PufM) using the following methodology. Sequences were aligned
using MUSCLE[42] (parameter: -maxiters 8) and automatically trimmed using TrimAL[43]
(parameter: -automated1) (Supplementary Data File 10 and 11). After manual assessment,
trimmed alignments were used to construct approximately-maximum-likelihood phylogenetic
trees using FastTree[45] (parameters: -lg -gamma) (Supplementary Data File 12 and 13).

Relative abundance of genomes in each sample

Reads from each sample were recruited against all assemblies ≥2kb from the same province
using Bowtie2[49] (parameters: default settings), under the assumptions that contigs <2kb would
include, low abundance bacteria and archaea, bacteria and archaea with high degrees of
repeats/assembly poor regions, fragmented picoeukaryotic genomes, and problematic read
sequences (low quality, sequencing artefacts, etc.). For the four sets of contigs (North Pacific,
Mediterranean, East Africa Coastal province, and South Atlantic), putative CDS were
determined via Prodigal (parameters: see above). In order to estimate the relative abundance of
the four analyzed genomes within the bacteria and archaea portion of the total microbial
community (excluding eukaryotes and viruses), single-copy marker genes were identified using a
collection of previously identified HMMs[50,51] and searched using HMMER[37] (hmmsearch --
notextw --cut_tc). Markers belonging to the four genomes were isolated from the total set of
environmental markers. The number of reads aligned to each marker was determined using
BEDTools[52] (v2.17.0; multicov default parameters). Length-normalized relative abundance
values were determined for each genome as in Equation 1 (Supplementary Data File 1):

251
$$\frac{\sum \text{Reads bp}^{-1} \text{ genome markers}}{\sum \text{Reads bp}^{-1} \text{ metagenome contig markers}} \times 100 \qquad (1)$$

Data availability

Data is available… submission to NCBI is ongoing. [Currently data is available at FigShare, including high resolution copies of figures, contig and protein sequences, and all supplementary data files: **https://figshare.com/s/9f603e9bbef71164e61b**]

References:

1. Béjá, O., Spudich, E. N., Spudich, J. L., Leclerc, M. & DeLong, E. F. Proteorhodopsin phototrophy in the ocean. *Nature* **411,** 786–789 (2001).

2. Harashima, K., Kawazoe, K., Yoshida, I. & Kamata, H. Light-Stimultated Aerobic Growth of Erythrobacter Species Och-114. *Plant and Cell Physiology* **28,** 365–374 (1987).

3. Schwalbach, M. S. & Fuhrman, J. A. Wide-ranging abundances of aerobic anoxygenic phototrophic bacteria in the world ocean revealed by epifluorescence microscopy and quantitative PCR. *Limnology and Oceanography* **50,** 620–628 (2005).

4. Koblížek, M. Ecology of aerobic anoxygenic phototrophs in aquatic environments. *FEMS Microbiol. Rev.* **39,** 854–870 (2015).

5. Béjá, O. *et al.* Unsuspected diversity among marine aerobic anoxygenic phototrophs. *Nature* **415,** 630–633 (2002).

6. Kang, I. *et al.* Genome Sequence of Fulvimarina pelagi HTCC2506T, a Mn(II)-Oxidizing Alphaproteobacterium Possessing an Aerobic Anoxygenic Photosynthetic Gene Cluster and Xanthorhodopsin. *J. Bacteriol.* **192,** 4798–4799 (2010).

7. Frigaard, N.-U. Biotechnology of Anoxygenic Phototrophic Bacteria. *Adv. Biochem. Eng. Biotechnol.* **156,** 139–154 (2016).

8. Hauruseu, D. & Koblížek, M. Influence of Light on Carbon Utilization in Aerobic Anoxygenic Phototrophs. *Appl. Environ. Microbiol.* **78,** 7414–7419 (2012).

9. Moran, M. A. *et al.* Deciphering ocean carbon in a changing world. *Proceedings of the National Academy of Sciences* **113,** 3143–3151 (2016).

10. Pesant, S. *et al.* Open science resources for the discovery and analysis of Tara Oceans data. *Sci. Data* **2,** 150023–16 (2015).

11. Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean microbiome. *Science* **348,** 1261359–1261359 (2015).

12. Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG Tools for Functional Characterization of Genome and Metagenome Sequences. *Journal of Molecular Biology* **428,** 726–731 (2016).

13. Tabita, F. R. *et al.* Function, Structure, and Evolution of the RubisCO-Like Proteins and Their RubisCO Homologs. *Microbiol. Mol. Biol. Rev.* **71,** 576–599 (2007).

14. Badger, M. R. & Bek, E. J. Multiple Rubisco forms in proteobacteria: their functional significance in relation to CO2 acquisition by the CBB cycle. *Journal of Experimental Botany* **59,** 1525–1541 (2007).

15. Rusch, D. B. *et al.* The Sorcerer II Global Ocean Sampling Expedition: Northwest Atlantic through Eastern Tropical Pacific. *Plos Biol* **5,** e77 (2007).

16. Venter, J. C. *et al.* Environmental genome shotgun sequencing of the Sargasso Sea. *Science* **304,** 66–74 (2004).

17. Yutin, N. *et al.* Assessing diversity and biogeography of aerobic anoxygenic phototrophic bacteria in surface waters of the Atlantic and Pacific Oceans using the Global Ocean Sampling expedition metagenomes. *Environ. Microbiol.* **9,** 1464–1475 (2007).

18. Baker, B. J., Lazar, C. S., Teske, A. P. & Dick, G. J. Genomic resolution of linkages in carbon, nitrogen, and sulfur cycling among widespread estuary sediment bacteria. *Microbiome* **3,** 14 (2015).

19. Gómez-Baena, G. *et al.* Glucose Uptake and Its Effect on Gene Expression in Prochlorococcus. *PLoS ONE* **3,** e3416–11 (2008).

20. Michelou, V. K., Cottrell, M. T. & Kirchman, D. L. Light-Stimulated Bacterial Production and Amino Acid Assimilation by Cyanobacteria and Other Microbes in the North Atlantic Ocean. *Appl. Environ. Microbiol.* **73,** 5539–5546 (2007).

21. Kaneko, T. *et al.* Sequence analysis of the genome of the unicellular cyanobacterium Synechocystis sp. strain PCC6803. II. Sequence determination of the entire genome and assignment of potential protein-coding regions. *DNA Res.* **3,** 109–136 (1996).

305   22.   Swan, B. K. *et al.* Potential for Chemolithoautotrophy Among Ubiquitous Bacteria Lineages in the Dark
306   Ocean. *Science* **333,** 1296–1300 (2011).

307   23.   de Zwart, J., Nelisse, P. N. & Kuenen, J. G. Isolation and characterization of Methylophaga sulfidovorans sp
308   nov: An obligately methylotrophic, aerobic, dimethylsulfide oxidizing bacterium from a microbial mat.
309   *FEMS Microbiol. Ecol.* **20,** 261–270 (1996).

310   24.   Kelly, D. P., Baker, S. C., Trickett, J., Davey, M. & Murrell, J. C. Methanesulphonate utilization by a novel
311   methylotrophic bacterium involves an unusual monooxygenase. *Microbiology* **140,** 1419–1426 (1994).

312   25.   Fischer, W. W., Hemp, J. & Johnson, J. E. Evolution of Oxygenic Photosynthesis. *Annu. Rev. Earth Planet.*
313   *Sci.* **44,** 647–683 (2016).

314   26.   Li, D. *et al.* MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced methodologies
315   and community practices. *Methods* **102,** 3–11 (2016).

316   27.   Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-generation sequencing
317   data. *Bioinformatics* **28,** 3150–3152 (2012).

318   28.   Treangen, T. J., Sommer, D. D., Angly, F. E., Koren, S. & Pop, M. Next generation sequence assembly with
319   AMOS. *Curr Protoc Bioinformatics* **Chapter 11,** Unit 11.8 (2011).

320   29.   Graham, E. D., Heidelberg, J. F. & Tully, B. J. BinSanity: unsupervised clustering of environmental
321   microbial assemblies using coverage and affinity propagation. *PeerJ* **5,** e3035–19 (2017).

322   30.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the quality
323   of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* **25,** 1043–1055
324   (2015).

325   31.   Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat Meth* **11,** 1144–1146
326   (2014).

327   32.   Eren, A. M. *et al.* Anviʻo: an advanced analysis and visualization platform for ’omics data. *PeerJ* **3,** e1319
328   (2015).

329   33.   Hyatt, D., LoCascio, P. F., Hauser, L. J. & Uberbacher, E. C. Gene and translation initiation site prediction
330   in metagenomic sequences. *Bioinformatics* **28,** 2223–2230 (2012).

331   34.   Aziz, R. K. *et al.* The RAST Server: Rapid Annotations using Subsystems Technology. *BMC Genomics* **9,**
332   75 (2008).

333   35.   Matsen, F. A., Kodner, R. B. & Armbrust, E. V. pplacer: linear time maximum-likelihood and Bayesian
334   phylogenetic placement of sequences onto a fixed reference tree. *BMC Bioinformatics* **11,** 538 (2010).

335   36.   Bateman, A. *et al.* The Pfam Protein Families Database. *Nucleic Acids Res.* **30,** 276–280 (2002).

336   37.   Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence similarity searching.
337   *Nucleic Acids Res.* **39,** W29–W37 (2011).

338   38.   Wu, D., Jospin, G. & Eisen, J. A. Systematic Identification of Gene Families for Use as ‘Markers’ for
339   Phylogenetic and Phylogeny-Driven Ecological Studies of Bacteria and Archaea and Their Major
340   Subgroups. *PLoS ONE* **8,** e77033–11 (2013).

341   39.   Santos, S. R. & Ochman, H. Identification and phylogenetic sorting of bacterial lineages with universally
342   conserved genes and proteins. *Environ. Microbiol.* **6,** 754–759 (2004).

343   40.   Alexandre, A., Laranjo, M., Young, J. P. W. & Oliveira, S. dnaJ is a useful phylogenetic marker for
344   alphaproteobacteria. *Int. J. Syst. Evol. Microbiol.* **58,** 2839–2849 (2008).

345   41.   Benson, D. A. *et al.* GenBank. *Nucleic Acids Res.* **28,** 15–18 (2000).

346   42.   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids*
347   *Res.* **32,** 1792–1797 (2004).

348   43.   Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated alignment trimming
349   in large-scale phylogenetic analyses. *Bioinformatics* **25,** 1972–1973 (2009).

350   44.   Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop software platform for the
351   organization and analysis of sequence data. *Bioinformatics* **28,** 1647–1649 (2012).

352   45.   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood trees for large
353   alignments. *PLoS ONE* **5,** e9490 (2010).

354   46.   Markowitz, V. M. *et al.* The integrated microbial genomes (IMG) system. *Nucleic Acids Res.* **34,** D344–8
355   (2006).

356   47.   Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M. & Tanabe, M. KEGG as a reference resource for
357   gene and protein annotation. *Nucleic Acids Res.* **44,** D457–D462 (2016).

358   48.   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND. *Nat Meth* **12,**
359   59–60 (2014).

49.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Meth* **9,** 357–359 (2012).

50.   Albertsen, M. *et al.* Genome sequences of rare, uncultured bacteria obtained by differential coverage binning
of multiple metagenomes. *Nat Biotechnol* **31,** 533–538 (2013).
51.   Tully, B. J. & Heidelberg, J. F. Potential Mechanisms for Microbial Energy Acquisition in Oxic Deep-Sea
Sediments. *Appl. Environ. Microbiol.* **82,** 4232–4243 (2016).
52.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26,** 841–842 (2010).
367

Acknowledgments:

We would like to acknowledge and thank Drs. Eric Webb and William Nelson for providing
invaluable comments and critiques in the early stages of this research. We are indebted to the
Tara Oceans consortium for their commitment to open-access data that allows data aficionados
to indulge in the data and attempt to add to the body of science contained within. And we thank
the Center for Dark Energy Biosphere Investigations (C-DEBI) for providing funding to BJT and
JFH (OCE-0939654). This is C-DEBI contribution number ###.

Author contributions: BJT conceived of the research plan, performed analysis, and wrote the
manuscript. EDG performed analysis and wrote the manuscript. JFH provided funding, provided
guidance, and edited the manuscript.

Competing financial interests: The authors declare no conflict of interest.
379

Fig. 1. The approximate locations of *Tara Oceans* sampling sites used to generate metagenomes incorporated in to this study. Each grid represents the three possible sample depths and filter fractions (top row: surface, middle row: DCM, bottom row: mesopelagic). An 'X' denotes that no sample was collected for that depth and size fraction at the site. Circle size represents relative abundance. (A) North Pacific – NP 970, (B) Mediterranean Sea – MED800, (C) South Atlantic – SAT68, and (D) East Africa current – EAC638. Size fraction: orange, 'bacterial' (0.22-1.6μm); purple, 'protistan' (0.8-5.0μm); blue, 'girus'+'viral' (<0.22-0.8μm). The maps in Figure 1 were modified under a CC BY-SA 3.0 license from 'South Atlantic Ocean laea location map' by Tentotwo, 'North America laea location map' by TUBS, 'Location of Mozambique in Africa' by Rei-artur, and 'Blank Map of South Europe and North Africa' by historicair.

Fig. 2. Phylogenetic tree of the ribulose-1,5-bisphosphate carboxylase large subunit (Rubsico, RbsL) with the major forms denoted. Inset (**A**) – A zoomed in view of the Type 1C/D Rubsico subgroup. Purple sequence names denote RbsL proteins from the Global Ocean Survey. Sequences used for this tree can be found in Supplementary Data File 10. Phylogenetic distances and local support values can be found in Supplementary Data File 12. Sequence information, including accession numbers and assignments can be found in Supplementary Data File 8.

Extended Data Fig. 3. Phylogenetic tree of the M subunit of type-II photochemical reaction
center (PufM). Environmental sequences obtained from the Global Ocean Survey (purple, •) and
Béjá *et al.* (2002) (pink, •) are highlighted. Boxes illustrate approximate positions of
phylogroups previously assigned by Yutin *et al.* (2007). Sequences used for this tree can be
found in Supplementary Data File 11. Phylogenetic distances and local support values can be
found in Supplementary Data File 13. Sequence information, including accession numbers and
taxonomies can be found in Supplementary Data File 9.
407

Fig. 4. Approximate maximum likelihood *Alphaproteobacteria* phylogenetic tree of 17
concatenated single-copy marker genes for the *Tara* assembled and 160 reference genomes.
Reference sequences from the *Gammaproteobacteria* used as an outgroup. Reference genome
information, including accession numbers, can be found in Supplementary Data File 4.
Phylogenetic distances and local support values can be found in Supplementary Data File 5.

Fig. 5. Cellular schematic of the four reconstructed genomes. (**A**) The presence of a gene(s) in a genome is represented by a yellow square (MED800), pink star (NP970), green hexagon (SAT68), and/or blue circle (EAC638). Schematic illustrates predicted membrane bound proteins, but does accurately represent cellular localization. (**B**) A detailed view of the proposed flow of electrons from donors to photosynthesis and carbon fixation. Abbreviations: cyt, cytochrome; Q pool, quinone pool; LH, light-harvesting proteins; soeABC, sulfite dehydrogenase (quinone); CBB, Calvin-Benson-Bassham.

424 **Table 1**. Statistics of the four *Tara* assembled genomes.

Genome ID	No. of contigs	Total length (bp)	Max. contig length (bp)	N50	Mean length (bp)	GC (%)	No. of predicted CDS$^\lozenge$	Est. Completeness (%)*	Est. duplication/redundancy (Est. strain heterogeneity) (%)*
MED800	174	2,140,579	55,951	13,831	12,302	36.7	2,184	66.98	2.36 (100.0)
SAT68	192	2,295,506	63,422	13,440	11,956	36.8	2,334	85.02	5.37 (75.0)
EAC638	121	1,868,409	80,916	19,502	15,441	30.7	1,939	68.88	4.75 (75.0)
NP970	120	2,585,639	84,649	32,137	21,547	36.1	2,445	82.04	3.04 (33.33)

425 $\lozenge$, as determined using Prodigal

426 *, as determine using CheckM with the Alphaproteobacteria markers (225 markers in 148 sets)

427 N50 - length of contig for which all contigs longer in length contain half of the total genome

428 CDS - coding DNA sequence