Framework for quality assessment of whole genome, cancer sequences

Justin P. Whalley[1,2], Ivo Buchhalter[3,4], Esther Rheinbay[5,6], Keiran M. Raine[7], Kortine

Kleinheinz[3], Miranda D. Stobbe[1,2], Johannes Werner[3], Sergi Beltran[1,2], Marta Gut[1,2],

Daniel Huebschmann[3,4,8], Barbara Hutter[9], Dimitri Livitz[5,6], Marc Perry[10], Mara

Rosenberg[5,6], Gordon Saksena[5,6], Jean-Rémi Trotta[1,2], Roland Eils[3,4], Jan Korbel[11],

Daniela S. Gerhard[12], Peter Campbell[7], Gad Getz[5,6,13], Matthias Schlesner[3], Ivo G.

Gut*[1,2], PCAWG-Tech, PCAWG-QC & PCAWG Network

[1]CNAG-CRG, Centre for Genomic Regulation (CRG), Barcelona Institute of Science and

Technology (BIST), Barcelona, Spain

[2]Universitat Pompeu Fabra (UPF), Barcelona, Spain

[3]Division of Theoretical Bioinformatics (B080), German Cancer Research Center
(DKFZ), Heidelberg, Germany

[4]Department for Bioinformatics and Functional Genomics, Institute for Pharmacy and
Molecular Biotechnology (IPMB) and BioQuant, Heidelberg University, Heidelberg,
Germany

[5]Massachusetts General Hospital Cancer Center and Department of Pathology, Boston,
USA

[6]Broad Institute of Harvard and MIT, Cambridge, MA, USA

[7]Wellcome Trust Sanger Institute, Hinxton, UK

[8]Department of Pediatric Immunology, Hematology and Oncology, University Hospital
Heidelberg, Heidelberg, Germany

[9]Division of Applied Bioinformatics (G200), German Cancer Research Center (DKFZ),
Heidelberg, Germany

[10]Ontario Institute for Cancer Research, Toronto, Ontario, Canada

[11]Genome Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany

[12]Office of Cancer Genomics, National Cancer Institute, US National Institutes of Health,
Bethesda, MD, USA

[13]Harvard Medical School, Boston, MA, USA

*Corresponding author: ivo.gut@cnag.crg.eu

Abstract

Working with cancer whole genomes sequenced over a period of many years in different

sequencing centres requires a validated framework to compare the quality of these

sequences. The Pan-Cancer Analysis of Whole Genomes (PCAWG) of the International

Cancer Genome Consortium (ICGC), a project a cohort of over 2800 donors provided us

with the challenge of assessing the quality of the genome sequences. A non-redundant set

of five quality control (QC) measurements were assembled and used to establish a star

rating system. These QC measures reflect known differences in sequencing protocol and

provide a guide to downstream analyses of these whole genome sequences. The resulting

QC measures also allowed for exclusion samples of poor quality, providing researchers

within PCAWG, and when the data is released for other researchers, a good idea of the

sequencing quality. For a researcher wishing to apply the QC measures for their data we

provide a Docker Container of the software used to calculate them. We believe that this is

an effective framework of quality measures for whole genome, cancer sequences, which

will be a useful addition to analytical pipelines, as it has to the PCAWG project.

Introduction

Combining whole genome sequencing data from individual projects has many advantages: increased statistical power, the ability to extend hypotheses across several projects and the possibility of asking biological questions covering a wider range of phenomena. However when the genome sequencing data comes from different centres, was sequenced at different times and under different protocols, great care must be taken to ensure that the sequencing data is of comparable quality, to avoid drawing false conclusions. The Pan-Cancer Analysis of Whole Genomes (PCAWG) project provided us with a great opportunity to assemble, test and finalise which quality control measures are important for comparing the quality of whole genome, cancer sequences.

The PCAWG project assembled a cohort of 48 projects encompassed in the International Cancer Genome Consortium (ICGC)[1] and The Cancer Genome Atlas (TCGA)[2] of which we analysed 2959 cancer genomes (normal-tumour genome pairs) from 2830 donors. The size of the dataset and the diversity of the samples, representing many different cancers from varied populations, allow the exploration of many fundamental questions of cancer. Although there were inclusion criteria based on the sequencing platform (Illumina) and minimum sequencing depth, there was a need to ascertain the quality of the sequencing data and how they compared to each other. There were 17 different sequencing centres involved and the sequencing was performed in over a five year time-span (2009-2014) (a time period of during which the sequencing methodology was still evolving rapidly), so to be able to perform analysis across the whole data set, it was necessary that the quality of the sequencing was carefully assessed.

There are advantages in developing a comprehensive set of quality measures. We will be

able to exclude samples of low quality. This will save running downstream analyses,

saving computational and the researchers' time. For researchers in PCAWG studying

driver mutations, we can provide a sanity check. If the driver mutation is only found in

low quality samples, it may not be a good candidate, compared to if it is supported by

high quality samples. As PCAWG will release the data for community to use, our quality

measures will provide a guide to the quality of the whole genome sequences within. For

researchers who wish to assess the quality of their whole genome cancer sequences, we

will also be releasing our methods, in a Docker Container for easy implementation.

To develop a framework needed to determine the quality of samples, we use the methods

employed by the sequencing centres involved in PCAWG as well as results in the

literature. TCGA marker papers (see references[3-5] for examples from 2014-16) all include

quality control (QC) measures such as depth of coverage, batch effects and contamination

levels, as calculated as part of the Firehose analysis infrastructure. Likewise a recent

ICGC paper[6] with samples sequenced from three different centres relied on similar QC

measures computed by the Picard toolkit. Lu et al.[7], carried out meta-analysis of exome

data available from the TCGA for 12 cancer types which is similar, but not identical in

scope, to the data set examined here. Their inclusion criteria were based on coverage

depth and percentage of exome coverage for both the normal and tumour samples. Other

cancer studies have also pointed to the importance of the percentage of the genome

covered[8,9] as well as error rates for each of the paired reads[10] as QC measures. The scale

and diversity of the PCAWG project provides a useful testing ground for QC measures,

both for selection of the measures and the thresholds to use in grading the sequences.

Here we present the results of the work by sequencing centres and research groups

involved in PCAWG to define important quality control measures, and how best to

combine the results from these measures. Based on the PCAWG data we selected

measures covering five important features to assess the quality of cancer genome

sequences: mean coverage, evenness of coverage, somatic mutation calling coverage,

paired reads mapping to different chromosomes and the ratio of difference in edits

between paired reads, an edit being a base in the read which is different to the reference

genome. These measurements we computed for both the normal and tumour samples. To

summarise the five QC measures, we established a star rating system to cover the range

of the highest quality cancer genomes, passing the thresholds set for each measurement,

to those that had many sequencing quality issues.

Results

All our analysis is based on the aligned sequences from the PCAWG core pipeline[11].

Within the aligned sequences we did not use duplicate reads, reads with a mapping

quality of zero and ignored supplementary alignments. The first three quality control

measures; mean coverage, evenness of coverage and somatic mutation calling coverage;

are linked to different aspects of the coverage of the genomic sequence. The other two

measures indicate discrepancies between the paired reads: mapping to different

chromosomes and the ratio of edits between the paired reads compared to the reference

genome. Finally we summarise these five measures into a star rating, for easy comparison

of each of the sample pair's quality.

Mean Coverage When deciding on what depth to sequence cancer genomes to, a trade

off has to be made between the advantages of sequencing deep to the cost of sequencing.

The deeper the cancer genome is sequenced the greater the confidence in calling somatic

events (see Tyler et al.[12] for a comparison of somatic mutation calling at depths up to

300X). A precondition for the inclusion of a patient in the PCAWG study was the

availability of a whole genome sequence of the normal and tumour with 25X coverage or

greater. We found that a number of the submitters calculated coverage differently. For

standardization the mean number of reads covering each position in the genome was

calculated, after low quality and duplicate reads were excluded so to not inflate the

number of reads (see *Supplementary Methods* for exact methods used). As shown in

Supplementary Figure S1, most commonly the normal samples were sequenced to around

30X, while there was a bimodal distribution for the tumour samples with maxima at 38X

and 60X. To provide a meaningful guide to the quality of the genomes in PCAWG, we

therefore set the thresholds for the mean coverage, after aligning, to 25X for normal

samples and 30X for tumour samples. This resulted in 0.4% normal and 2.2% tumour

samples not reaching these minimum criteria (see *Supplementary Figure S1*).

Evenness of Coverage To confidently identify germline variants and somatic mutations

requires an even coverage across the target area[13], in this case the entire genome. Two

different methods, in use by the sequencing centres involved for whole genomes, were

chosen. One measure is to calculate the ratio of the median coverage over the mean

coverage (MoM). An evenly covered sequence should have a ratio of one, with the mean

value the same as the median value, not skewed by very low or high coverage in certain

regions. To decide within what range of values a sample should fall to be regarded as

evenly covered, we used the whiskers of the boxplots in *Figure 1*, 1.5 × I.Q.R

(interquartile range) of the data, which results in the range of 0.99 - 1.06 for a normal

sample and the wider range of 0.92 - 1.09 for the tumour samples (see *Supplementary*

Figure S2). For MoM coverage ratio (and for FWHM described below), there is a greater

range of values for the tumour samples than normal samples, potentially due to

biologically reasons valid for tumours, e.g. large deletions could lead to a more unevenly

covered sample. If the normal sample is unevenly covered, it is more likely due to a

sequencing artefact. Hence, we are more stringent for the normal than the tumour

samples.

The second measure of evenness looks at the variation of the normalised coverage in ten

kilobase (kb) genomic windows, after correction for GC-dependent coverage bias using

the somatic CNV calling algorithm ACEseq[14] (*Figure 2*). The main cloud, which

corresponds to the main copy number state of the sample, is determined (as shown by the

red dots in *Figure 2*). The remaining coverage variation is measured as full width at half

maximum (FWHM) of the main cloud. This measure is insensitive to copy number

aberrations and GC-dependent coverage bias. To determine the thresholds, 1000 WGS

samples from different tumour types were used. We chose the pruning values based on

clustering of these samples and subsequent visual inspection of the "best" samples that

exceeded the threshold to see whether they are valid. Using these results the thresholds

chosen are 0.205 for the normal and the more lenient 0.34 for the tumour, above which

the sample would be regarded as having an uneven coverage (see *Supplementary Figure*

S3).

The two evenness measures tend to identify different samples as having uneven coverage

(see *Figure 3*). Spearman's correlation coefficient for the two measures suggests that

these measures are not correlated for the normal ($\rho = 0.24$) and tumour ($\rho = -0.06$)

samples. FWHM is insensitive to GC bias, as the CNV caller corrects for this while MoM

identifies other evenness outliers.

A sample needs to be in the respective ranges of the MoM and FWHM for the normal and

the tumour to pass the evenness quality measure, of which 6.28% and 5.81% respectively

of the samples were not.

Somatic Mutation Calling Coverage Having both the measure of the depth of coverage

in terms of the mean and evenness of coverage, our next QC measure look at the effect of

these at each base in the cancer genome (so both the normal and the tumour sample). This

measures gives a good summary of how much of the cancer genome is sufficiently

covered to call a somatic mutation event. MuTect[15] with default settings calculates for

each base in the genome, if it has sufficient coverage in both the normal and tumour

sample (least fourteen reads are present in the tumour and eight reads in the matched

normal sample). Based on those requirements, we had to establish the number of bases to

consider the sample sufficiently covered. Ideally the threshold should be high enough to

penalise the less well-sequenced samples, while not unduly penalising tumour samples

that have had large deletions in the genome resulting in fewer bases to sequence. Taking

into account the largest unambiguous mapping for a female donor (so not including the Y

chromosome) would be 2,835,690,481 bases[16], 2.6 gigabases would best suit these two

needs. This results in 5.95% of normal-tumour pairs with fewer bases, than this threshold

(see *Supplementary Figure S4*).

Paired reads mapping to different chromosomes The two reads from a read pair

should represent the ends of a contiguous DNA sequence that depending on the insert

size should be a given distance apart (for PCAWG between 200 and 1,000 bases). Paired

reads mapping to different chromosomes can be due to a rearrangement. However an

excess of reads mapping to different chromosomes points to a technical artefact. So

deciding a threshold based on percentage of paired reads mapping to different

chromosomes, we should not penalise sequences with biological causes of the paired

reads mapping to different chromosomes (such as chromothripsis[17], or more generally,

interchromosomal rearrangements). We set the threshold to 3%, which even samples with

confirmed high levels of rearrangements and chromothripsis do not exceed (which in our

experience, do not have more than 1% of paired reads mapping to different

chromosomes). Of the normal sequences 14.5% exceed the threshold, as do 13.0%

tumour sequences (see *Supplementary Figure S5*). Interestingly there are more normal

samples failing this measure, which cannot be explained by biological processes. A

possible explanation may be that for lower quality samples in preparing libraries with

PCR amplification, this amplification step causes an increase in two fragments of DNA

from different parts of the genome being fused together, as has previously been noted[18].

Consequently, this translates to an increase in percentage of paired reads mapping to

different chromosomes.

Ratio of difference in edits between paired reads Damage in sequencing runs has been

linked to a global imbalance in edits (where the base in read is different compared to the

reference) between read 1 and read 2 in paired end sequencing[19]. Therefore the ratio of

the sum of edits between paired reads for a well-sequenced sample should be close to

one. We adjudged samples with a two-fold ratio of edits between the paired reads, or

greater, as having something gone wrong in the sequencing cycle resulting in lower data

quality. Based on this threshold 4.66% and 4.49% normal and tumour samples failed

respectively.

Summary The five quality measures were selected to provide minimal redundancy in

flagging quality issues in normal/tumour paired genome sequences - that each measure

reflects a facet of sequencing quality that other measure does not. The best way to

summarise these comparisons between the different measures is with a Venn diagram

(*Figure 4*). There is some overlap between certain measures, for example 75 sample pairs

are penalised by both having a high percentage paired reads mapping to different

chromosomes and uneven coverage. However a much higher number of samples

penalised by one of these measures and not the other. Having defined these five, non-

redundant QC measures our next step was to summarise them, to give an overall score for

quality for the other researchers in PCAWG to use.

Star rating system

We used the five quality measures to construct a star rating for each cancer genome

(normal/tumour whole genome sequence). For each QC measure a star is awarded if both

the normal and tumour sample pass the threshold. Half a star is awarded if only the

normal passes the threshold for the respective QC measures. For somatic mutation calling

coverage, a whole star is awarded for passing, none otherwise. The reasons for the extra

weighting of the normal sample for the other four measures are that there is no biological

reason for low quality in the normal sequence and a well-sequenced normal sample is

important for calling somatic mutations.

Summing the stars earned for each of the five QC measure results in 66.4% of the

normal/tumour sample pairs of the PCAWG being rated as 5 stars. Looking specifically

at the different projects (see *Figure 5*), a more nuanced picture is available. The quality

does not seem to be biased by tissue type (see *Supplementary Figure S7*) based on

detailed molecular subtypes of the tumours in PCAWG[20], the difference seems to be

more at the project level. Unfortunately, there is only limited project metadata on when

and which protocol was used to sequence the samples. Detailed metadata was available

for 95 donors of the CLLE-ES project (concerning Chronic Lymphocytic Leukaemia), so

it could be used as an example. Changes in protocol had an effect on the quality of the

sequencing over the four years in which CLLE-ES samples were sequenced. For the

CLLE-ES project, most notable was the change to a no PCR proband in 2012, which

resulted in improvements to the measures of paired reads mapping to different

chromosomes and evenness of coverage. This in turn resulted in a measurable change in

somatic mutation calling coverage and improvement in star ratings (*Supplementary*

Figure S8). We found similar results for a subset of 348 samples sequenced at the Broad

Institute (see *supplementary Figure S9*), which had metadata recorded in CGHub[21] about

the time and instruments used to sequence. We hypothesise that this will be true for other

projects as well.

Having calculated the star rating for the sequences, it was interesting to see how our QC

measures relate to the calling of somatic single nucleotide variants (SNVs)[11], somatic

insertion and deletions (indels)[11] and somatic structural variants (SVs)[22] in PCAWG. An

advantage of using these PCAWG datasets is that four callers were used for each.

Looking at the proportion of calls, which all four callers supported, gives us a good idea

how the quality of sequencing influences the identification of unambiguous somatic

mutations. While the proportion of calls supporting the four callers varies greatly by

sample, we find that the samples with four stars or greater tended to have higher

proportions than samples with less than four stars for SNVs, indels and SVs (with p-

values of $\sim 10^{-5}$, $\sim 10^{-5}$, $\sim 10^{-18}$ respectively, using the Mann-Whitney-U test, also see

Figure 6).

Taking this analysis further we used linear regression models, to further analyse the

relation between the proportion of calls supported by four callers and the actual QC

measures (see *Supplementary Tables S1-S3*). The results show that, significantly, an

increasing percentage of paired reads mapping to different chromosomes in tumour

samples, has a negative effect on the proportion of calls supported by four callers for

SNVs, indels and SVs. More specifically, for SNVs an increasing mean coverage in

tumours has a significant positive effect on the proportion of calls supported by four

callers. While in indels there is a significant effect negative effect on the proportion of

calls supported by four calls by increasing unevenness (as measured by FWHM) in

tumours. As in indels, the unevenness effect is also true in SVs as well as significant

negative effects by increasing percentage of paired reads mapping to different

chromosomes in normal samples and ratio of difference in edits between paired reads in

tumour samples.

The results from this analysis suggest the quality of sequencing as measured by star

rating does have a measurable effect the downstream analyses. However the QC

measures which make up the star rating effect the different downstream analyses in

different ways. As our QC measures reflect different aspects of sequencing quality, they

also have varying levels of importance in using these sequences in the downstream

analyses of calling SNVs, indels and SVs.

Discussion

The established star rating system allows grading the normal and tumour sample

sequences by quality in absence of information on how sequencing was carried out, what

protocols were used and what problems may have occurred during the sequencing

process. The system is not designed to be all encompassing, instead using a small amount

of computational resources and time (compared to the actual aligning of the sequences),

we get a good snapshot of the quality of the normal-tumour sample pair sequences on

which to call somatic mutations. Likewise having graded the cancer genomes with our

five-star system, we do not intend researchers to necessarily exclude the lower ranked

cancer genomes, just to be wary of any conclusions based solely on the lower scoring

genomes.

With our star rating system, we sent several samples to the exclusion list due to their poor

performance in one of the QC measures. Due to the timing, this did not prevent the

downstream analyses being performed. Though anecdotally it would have saved 55 days

computational runtime for our one star sample. For all samples that remained, the QC star

rating was embedded in the header of the variant call format files for use of the

researchers within PCAWG, and when the data is released, to all researchers.

For those projects in PCAWG, which we had metadata, we found that sequencing quality

has definitely improved over the time period 2009-2014 in which the samples sequenced.

Our results for the CLLE-ES project suggest that in part a protocol change to PCR-free

methods improved sequencing, as in line with best practices from a recent benchmarking

exercise.[12]

Another advantage of our quality control is the link to the downstream analyses. In

aggregate, the higher the quality of the sequences, had a higher proportion of the

consensus somatic SNVs, indels, SVs called. These results suggest overall that higher

quality sequence will identify the true positive somatic mutations with higher probability.

Our data would suggest that when pre-amplification of DNA will be needed for WGS,

e.g. DNA isolated from formalin fixed, paraffin embedded tissue, the star rating system

will be helpful when the variants and mutations are interpreted.

We believe that our method can be adapted for similar projects that look to use whole

genome sequences from a variety of sources. The thresholds we used based on our

experience and applied to this dataset of 2959 cancer genomes can also be used as guide

to quality of sequences. It is worth noting that they represent a trade-off of being severe

enough to penalise poor quality while not discriminating against samples with valid

biological causes. We also would recommend using our methods to ascertain the quality

before downstream analyses by other groups. To enable others to use our approach, there

is a Docker Container, which can be accessed at https://github.com/eilslabs/PanCanQC.

We provide a framework for quality assessment, which opens the door to do large-scale

meta-analysis in a more robust framework.

Acknowledgements

The authors would like to thank Jennifer Jennings and her colleagues at the Ontario

Institute for Cancer Research (OICR) for their help in the administration of this working

group.

JPW, MDS, SB, MG, JT and IGG are supported by the Ministerio de Economía, Industria

y Competitividad and European Regional Development Fund (MINECO/FEDER

BIO2015-71792-P), the Instituto de Salud Carlos III (ISCIII) and the Generalitat de

Catalunya. In addition we have received funding from ELIXIR-EXCELERATE (EC

H2020 #676559) and RD-Connect (EC FP7/2007-2013 #305444).

The work done by IB, KK, JW, DH, BH, RE and MS was supported by the BMBF-

funded Heidelberg Center for Human Bioinformatics (HD-HuB) within the German

Network for Bioinformatics Infrastructure (de.NBI) (#031A537A, #031A537C) and the

BMBF-funded German ICGC-projects (ICGC-PedBrain: 109252 (German Cancer Aid),

01KU1201A,B; ICGC-MMML: 01KU1002B and ICGC-DE-MINING: 01KU1505E).

ER, DL, MR, GS and GG would like to acknowledge G.G. MGH startup package and

Broad funds.

KMR and PC are members of the Cancer Genome Project supported by a Wellcome

Trust grant (098051).

Author contributions
JPW, IB, ER, KMR, KK, MDS and JW wrote the manuscript, helped develop and apply

the methods and analysed the results.

SB, MG, DH, BH, DL, MP, MR, GS and JT contributed to the development of the

methods.

RE, JK, DSG, PC, GG, MS and IGG provided project supervision; through feedback and

the reviewing of the work done, as well as editing of the manuscript.

IB and JW constructed the Docker Container with code contributions from KK and

KMR.

PCAWG-Tech and PCAWG-Network provided the data, metadata and the framework for

this research.

344
Competing financial interests
The authors declare no competing financial interests

References

1.   International Cancer Genome Consortium et al. International network of cancer
genome projects. Nature 464, 993–8 (2010).

2.   Cancer Genome Atlas Research Network et al. The cancer genome atlas pan-
cancer analysis project. Nat Genet 45, 1113–20 (2013).

3.   Ceccarelli, M. et al. Molecular profiling reveals biologically discrete subsets and
pathways of progression in diffuse glioma. Cell 164, 550–63 (2016).

4.   Cancer Genome Atlas Network. Comprehensive genomic characterization of head
and neck squamous cell carcinomas. Nature 517, 576–82 (2015).

5.   Cancer Genome Atlas Research Network. Comprehensive molecular
characterization of urothelial bladder carcinoma. Nature 507, 315–22 (2014).

6.   Biankin, A. V. et al. Pancreatic cancer genomes reveal aberrations in axon
guidance pathway genes. Nature 491, 399–405 (2012).

7.   Lu, C. et al. Patterns and functional implications of rare germline variants across
12 cancer types. Nat Commun 6, 10086 (2015).

8.   Liu, J. et al. Genome and transcriptome sequencing of lung cancers reveal diverse
mutational and splicing events. Genome Res 22, 2315–27 (2012).

9.   Ramkissoon, L. A. et al. Genomic analysis of diffuse pediatric low-grade gliomas
identifies recurrent oncogenic truncating rearrangements in the transcription
factor mybl1. Proc Natl Acad Sci U S A 110, 8188–93 (2013).

10.   Berger, M. F. et al. The genomic complexity of primary human prostate cancer.
Nature 470, 214–20 (2011).

11.   Simpson, J. et al. Detecting Somatic Mutations in 2,834 Cancer Whole Genomes.
In preparation.

12.   Alioto, T. S. et al. A comprehensive assessment of somatic mutation detection in
cancer using whole-genome sequencing. Nat Commun 6, 10001 (2015).

13.   Mokry, M et al. Accurate SNP and mutation detection by targeted custom
microarray-based genomic enrichment of short-fragment sequencing libraries.
Nucleic Acids Res (2010).

14.   Kleinheinz et al. Copy-number variants from ACESeq. In preparation.

15. Cibulskis, K. Et al. Sensitive detection of somatic point mutations in impure and
heterogeneous cancer samples. Nat Biotechnol 31, 213–9 (2013).

16. Zook, J. M. et al. Integrating human sequence data sets provides a resource of
benchmark snp and indel genotype calls. Nat Biotechnol 32, 246–51 (2014).

17. Korbel, J. O. & Campbell, P. J. Criteria for inference of chromothripsis in cancer
genomes. Cell 152, 1226–36 (2013).

18. Oyola, S. O. *et al.* Optimizing illumina next-generation sequencing library
preparation for extremely at-biased genomes. BMC Genomics 13, 1 (2012).

19. Chen, L., Liu, P., Evans, T. C. & Ettwiller, L. M. DNA damage is a pervasive
cause of sequencing errors, directly confounding variant identification. Science
355, 752– 756 (2017).

20. Hoadley, K. et al. Supervised and unsupervised molecular classification of diverse
tumour types from whole genome sequencing data. In preparation.

21. Wilks, C. et al. The cancer genomics hub (CGHub): overcoming cancer through
the power of torrential data. Database (2014).

22. PCAWG-6. Patterns of structural variations, signatures, genomic correlations,
retrotransposons, mobile elements. In preparation.

Figures

Figure 1: Distribution of the median coverage over mean coverage ratios for normal and
tumour samples. The horizontal dashed bar at 1 represents the value of an evenly covered
sample. As shown in the plot the tumour samples have a greater spread of values than the
normal, we hypothesize this is to be expected as tumours are more likely to have deletions
and structural rearrangements, which will lead to less evenly covered sequence. The
whiskers on each of the boxplots (0.99-1.06 for the normal and 0.92-1.09 for the tumour)
were taken as thresholds for this measure.

a)

b)

Figure 2: GC content versus the normalised coverage for evenly covered sample (a) and
unevenly covered sample (b). The main cloud, corresponding to the main copy number
state of the samples, is indicated in red. Other clouds (yellow in the (a)) represent
different copy number states of copy- number aberrant regions. FWHM is calculated on
the main copy number state.

a)

b)

Figure 3: Density scatter plot comparing the two evenness of coverage measures for
normal (a) and tumour (b). The number of points overlapping is reflected by the colour at
that point as shown by the legend. The dashed lines reflect the thresholds for the evenness
measures. These graphs show while there both methods pick out certain samples as
unevenly covered, they also show individually samples which do not have even coverage.

Figure 4: Venn diagram showing for which QC measure sample pairs were penalised for.
The outside numbers show that each QC measures penalises a fair number of sample
pairs uniquely. Looking at the overlaps between QC measures, while some measures are
closer to each other than others, they all maintain a large degree of independence.

Figure 5: Distribution of the star ratings for the PCAWG genomes, grouped by tissue
type (as labelled along the x-axis), and then project. The project name and number of
samples in the project are labelled at the top of the bar. The colour of the bar reflects
what percentage of samples in the project have that star rating (corresponding to the
legend). The bar on the far left shows the results for all samples. The plot demonstrates
the varying quality of different projects - differences we believe come from when the
genome was sequenced and the sequencing protocol used.

434
Figure 6a: Samples with four stars or greater tend to have a higher the proportion of somatic single nucleotide variants (SNV) calls supported by four callers than samples with fewer than four stars. This is significant using the Mann-Whitney U test, with p-value ~ 10^{-5}.

Figure 6b: Samples with four stars or greater tend to have a higher the proportion of somatic insertion and deletion (indel) calls supported by four callers than samples with fewer than four stars. This is significant using the Mann-Whitney U test, with p-value ~ 10^{-5}.

445   *Figure 6c: Samples with four stars or greater tend to have a higher the proportion of*
446   *somatic structural variant (SV) calls supported by four callers than samples with fewer*
447   *than four stars. This is significant using the Mann-Whitney U test, with p-value ~ 10^{-8}.*