Title: Drosophila Ca$_V$2 channels harboring human migraine mutations cause synapse hyperexcitability that can be suppressed by inhibition of a Ca^{2+} store release pathway

Short Title: Drosophila models of Ca$_V$2 migraine mutations

Authors and Affiliations:

Douglas J. Brusich [1,4], Ashlyn M. Spring [1,2,5], Thomas D. James [1,3], Catherine J. Yeates[1,3], Timothy H. Helms [1,6], C. Andrew Frank [1,2,3]

[1] – Department of Anatomy and Cell Biology, University of Iowa Carver College of Medicine, Iowa City, IA 52242

[2] – Interdisciplinary Graduate Program in Genetics, University of Iowa, Iowa City, IA 52242

[3] – Interdisciplinary Graduate Program in Neuroscience, University of Iowa, Iowa City, IA 52242

[4] – Present Address: Human Biology, University of Wisconsin-Green Bay, Green Bay, WI 54311

[5] – Present Address: Department of Biology, University of North Carolina, Chapel Hill, NC 27599

[6] – Present Address: Department of Veterinary Biosciences, Ohio State University, Columbus, OH 43210

Address for Correspondence: C. Andrew Frank. Department of Anatomy and Cell Biology, University of Iowa Carver College of Medicine, 51 Newton Road, 1-661A. Iowa City, IA 52242. USA.

andy-frank@uiowa.edu

ABSTRACT

Gain-of-function mutations in the human $Ca_V2.1$ gene *CACNA1A* cause familial hemiplegic migraine type 1 (FHM1). To characterize cellular problems potentially triggered by $Ca_V2.1$ gains of function, we engineered mutations encoding FHM1 amino-acid substitutions S218L (SL) and R192Q (RQ) into transgenes of *Drosophila melanogaster* Ca_V2/*cacophony*. We expressed the transgenes pan-neuronally. Phenotypes were mild for RQ-expressing animals. By contrast, single mutant SL- and double mutant (DM) SL, RQ-expressing animals showed overt phenotypes, including sharply decreased viability. By electrophysiology, SL- and DM-expressing neuromuscular junctions (NMJs) exhibited enhanced evoked discharges, supernumerary epileptiform discharges, and an increase in the amplitudes and frequencies of spontaneous events. Some spontaneous events were gigantic (10-40 mV), multi-quantal events. Gigantic spontaneous events were eliminated by application of TTX – or by lowered or chelated Ca^{2+} – suggesting that gigantic events were elicited by spontaneous nerve firing. A follow-up genetic approach revealed that some neuronal hyperexcitability phenotypes were reversed after knockdown or mutation of Drosophila homologs of phospholipase Cβ (PLCβ), IP_3 receptor, or ryanodine receptor (RyR) – all factors known to mediate Ca^{2+} release from intracellular stores. Pharmacological inhibitors of intracellular Ca^{2+} store release produced similar effects. Interestingly, however, the decreased viability phenotype was not reversed by genetic impairment of intracellular Ca^{2+} release factors. On a cellular level, our data suggest inhibition of signaling that triggers intracellular Ca^{2+} release could counteract hyperexcitability induced by gains of $Ca_V2.1$ function.

AUTHOR SUMMARY

Prior research has demonstrated that gain-of-function mutations in a gene important for neurotransmission (*CACNA1A*) are known to cause migraine in humans. We attempted to mimic some of those gain-of-function mutations in a simple genetic model organism and to examine neurotransmission by electrophysiology. Our findings yield potential clues as to how particular migraine-causing mutations may impact neurophysiology on a cellular level. We used the fruit fly *Drosophila melanogaster* and its

model synapse, the neuromuscular junction (NMJ) to perform our studies. We document three main advances: 1) characterization of fruit fly models harboring gain-of-function calcium channel alterations known to cause human familial hemiplegic migraine type 1 (FHM1); 2) characterization of hyperactive neurotransmission caused by one of these alterations; and 3) an ability to quell hyperactive neurotransmission by impairing intracellular Ca^{2+} store release, through both genetic and pharmacological means. Our work contributes to a broader understanding of how pathological mutations could impact cellular physiology. More generally, the utilization of genetic model organisms promises to uncover potential ways to reverse those impacts.

INTRODUCTION

Episodic neurological disorders like migraine, epilepsy, and ataxia can result from underlying ion channel dysfunctions (1-3). For many such disorders, little is known about how aberrant channel functions affect neuronal signaling paradigms. Cell-based and model organism-based examinations of disease-causing mutations could offer insights into disease-relevant biological processes. One Mendelian form of migraine – familial hemiplegic migraine type 1 (FHM1) – results from gain-of-function missense mutations in human *CACNA1A*, which encodes the $\alpha 1$ subunit of $Ca_V2.1$ (P/Q)-type calcium channels (4). Two FHM1-causing amino-acid substitutions alter highly conserved $Ca_V2.1$ $\alpha 1$ amino-acid residues, R192 and S218 (4, 5). The R192Q amino-acid substitution (RQ) causes "pure" FHM1, while the S218L substitution (SL) causes a severe combination of FHM1, seizures, and susceptibility to edema following head injury (4, 5). These two FHM1-causing amino-acid substitutions have been studied intensely (6), most notably in knock-in mouse models of FHM1 (7-9).

FHM1 knock-in mice display gain-of-function $Ca_V2.1$ phenotypes at neurons and synapses. Model synapses studied include the diaphragm neuromuscular junction (NMJ) (10, 11), the calyx of Held (12-14), the trigeminal sensory neuron pathway (15-17), and cortical neurons (18, 19). At the mouse NMJ, both RQ and SL increase the frequency of spontaneous excitatory potentials (10, 11). These increases in quantal frequency are dependent on mutation dose and are more pronounced in SL versus

RQ. SL also elicits broadening of evoked end-plate potentials at the mouse NMJ (11). At the calyx of

Held, both substitutions result in enhanced excitatory postsynaptic currents (EPSCs) (12-14), and it has

been reported that SL causes an increase in the resting intracellular neuronal calcium, which could be

responsible for some potentiation of synapse function (12).

It was recently reported that 2,5′-di(tertbutyl)-1,4,-benzohydroquinone (BHQ) reverses aspects of

SL-induced gating dysfunction and short-term plasticity (20). As part of that study, we found that BHQ

also restores short-term synaptic plasticity to NMJs in fruit fly larvae expressing a transgene that

encodes an S161L amino-acid substitution in Drosophila Ca_V2/Cacophony – the functional equivalent of

human $Ca_V2.1$ S218L (20). Independent follow-up work in the mouse S218L model demonstrated that

BHQ application also blunts cortical spreading depression (CSD) susceptibility (21). Given these

collective results, a further examination of fruit fly synapses could be valuable for uncovering relevant

molecular and electrophysiological consequences of $Ca_V2.1$ gains of function.

For the present study, we characterized the fruit fly as a way to model neuronal effects of FHM1-

causing mutations. We neuronally expressed *cacophony* transgenes harboring the *Drosophila*

melanogaster equivalents of RQ or SL – or both RQ and SL concurrently (denoted as "DM" for "double

mutant"). On the organismal level, neuronal expression of SL or DM transgenes drastically impaired

overall health. On the synapse level, SL and DM transgenes markedly enhanced aspects of evoked and

spontaneous neurotransmission, consistent with prior studies in mice. Through a combination of

genetics, RNA interference, pharmacology, and electrophysiology, we uncovered evidence that

impairment of a conserved intracellular signaling pathway that triggers store Ca^{2+} release reverses

hyperexcitability phenotypes in the context of gain-of-function Drosophila Ca_V2.

RESULTS

Transgenic Drosophila Ca_V2 "FHM1" channels cause coarse larval phenotypes and fly lethality

We undertook a reverse-translational approach in *Drosophila melanogaster* to study the impact

that FHM1-inducing $Ca_V2.1$ amino-acid substitutions may exert on the level of individual synapses.

Drosophila *cacophony* encodes the $\alpha1$ subunit of fruit fly Ca_V2-type channels. We cloned two amino-acid

substitutions that cause human FHM1 (S218L and R192Q) into the analogous codons of a functional

Drosophila *UAS-cacophony (cac)-eGFP* transgene (22). Single mutant transgenes were termed "SL"

(*UAS-cac-eGFP^{S161L}*) (20) or "RQ" (*UAS-cac-eGFP^{R135Q}*) (Fig. 1A). We also generated a double mutant

transgene containing both mutations on the same cDNA clone, termed "DM" (*UAS-cac-eGFP$^{R135Q, S161L}$*).

"WT" signifies previously characterized wild-type *UAS-cac-eGFP^{WT}* transgenes (22).

We expressed WT, DM, SL, and RQ *UAS-cac-eGFP* transgenes in post-mitotic Drosophila

neurons using the *elaV(C155)-Gal4* driver and the *Gal4/UAS* expression system (23, 24). Initially, we

examined transgenic animals qualitatively for visible phenotypes. Neuronal expression of either SL or

DM caused larvae to move in a jerky, uncoordinated manner. At the early third instar stage, SL- and DM-

expressing animals developed protruding, anterior spiracles prematurely – well before the normal time

point of wandering third instar stage and pupation (Fig 1B).

Our initial observations indicated that SL- and DM-expressing animals were not present in

expected Mendelian proportions. For each transgene (WT, DM, SL, and RQ), we set up test crosses

(*elaV(C155)-Gal4* females x *Balancer Chromosome/UAS-cac-eGFP* males) and counted the number of

transgenic *UAS-cac-GFP*-expressing adult progeny and the number of sibling flies carrying a balancer

chromosome. We also set up *Gal4* and balancer chromosome control crosses lacking any *UAS-cac-*

eGFP transgenes (Table 1). Compared to animals expressing the WT transgene, viability was

dramatically diminished for animals expressing the SL and DM transgenes (Fig. 1C, Table 1). It was also

diminished for SL- and DM-expressing animals compared to genetically matched control siblings carrying

the *elaV(C155)-Gal4* driver and a balancer chromosome (Table 1). WT- and RQ-expressing animals did

not show significant defects in viability or statistical differences from the Gal4 control cross (Fig. 1C,

Table 1). As expected, there was some depressed viability in animals carrying a balancer chromosome

alone (Table 1).

Sex or dose of the SL and DM transgenes could influence viability. In Drosophila, X-linked

dosage compensation equalizes the expression of X-linked genes by doubling X-linked gene

transcription in males (25-27). The X-linked neuronal enhancer trap Gal4 line *elaV(C155)-Gal4* should be

expressed at higher levels in hemizygous *elaV(C155)-Gal4/Y* males than in heterozygous *elaV(C155)-*

Gal4/+ females. Thus, effects of driving *UAS* transgenes could be stronger in males. Counting male vs.

female progeny of SL- and DM-expressing flies revealed that while viability was starkly diminished for

both sexes, it was also significantly lower in SL- and DM-expressing males than in SL- and DM-

expressing females (Fig. 1C, Table 1).

Table 1. Test Crosses and Survival of Adult Progeny.

elaV(C155)-Gal4 x *Balancer /&*	*GAL4/+; &/+* progeny (female)	*GAL4/+; Balancer/+* progeny (female)	Normalized Viability Index for & (female)	*GAL4/Y; &/+* progeny (male)	*GAL4/Y; Balancer/+* progeny (male)	Normalized Viability Index for & (male)
& = +	52 (30.6%)	37 (21.8%)	85.1	53 (31.2%)	28 (16.5%)	114.6
& = *UAS-cac-eGFP^{WT}*	190 (30.3%)	115 (18.3%)	100.0	205 (32.6%)	118 (18.8%)	105.2
& = *UAS-cac-eGFP^{DM}*	**132 (12.7%)**	**499 (48.0%)**	**16.0 *****	**38 (3.7%)**	**371 (35.7%)**	**6.2 *** ###**
& = *UAS-cac-eGFP^{SL}*	**38 (12.3%)**	**136 (44.2%)**	**16.9 *****	**17 (5.5%)**	**117 (38.0%)**	**8.8 *** #**
& = *UAS-cac-eGFP^{RQ}*	157 (29.0%)	128 (23.6%)	74.2	147 (27.1%)	110 (20.3%)	80.9

Crosses were performed utilizing *elaV(C155)-Gal4* virgin females x *w/Y; Balancer/&*
(UAS-cac-eGFP or +) males. Male and female progeny were counted separately. As
expected for fruit fly balancer chromosomes, there was some lethality associated with
inheriting a balancer chromosome. Separately, there was profound lethality associated
with inheriting either the *UAS-cac-eGFP^{DM}* or *UAS-cac-eGFP^{SL}* transgenes driven by pan-
neuronal *elaV(C155)-Gal4*. *&* - denotes the chromosome utilized opposite the balancer in
males in the test cross. Balancers used depended on the cross and transgene being
balanced (*CyO-GFP* was used for *& = +* and for *& = UAS-cac-eGFP^{DM}* and TM6b was
used for the others). Normalized Viability Index scores were calculated from the non-
Balancer/Balancer progeny ratio of a single sex; this value was then normalized against
the female progeny ratio for the cross where *& = UAS-cac-eGFP^{WT}* – i.e., 190/115 = a
normalized baseline value of 100.0. For statistical analyses, raw progeny counts were
used. *** $p < 0.001$ by Fisher's exact test, compared to sex-matched progeny counts,
utilizing the *UAS-cac-eGFP^{WT}* as the control. ### $p < 0.001$, # $p = 0.05$, comparing DM- and
SL- expressing males vs. DM- and SL-expressing females respectively.

We also assessed adult fly longevity, comparing WT and DM transgenic flies (Figs. 1D, E). For

females, transgenic WT (mean survival: 63 days, *n* = 23) and driver control *elaV(C155)-Gal4/+* animals

(66 days, *n* = 16) did not differ with respect to survival. Transgenic DM females (22.5 days, *n* = 28) had

severely stunted longevity (Fig. 1D). The results for males were consistent: longevity of transgenic WT

males (median survival: 56 days, *n* = 38) and driver control *elaV(C155)-Gal4/Y* animals (54.5 days, *n* =

22) did not differ statistically. By contrast, the survival of transgenic DM males (14.5 days, *n* = 10) was

markedly diminished (Fig. 1E).

Cac-GFP localizes normally and levels are comparable across transgenic constructs

We investigated why SL- and DM-expressing animals were showing overt phenotypes. We

considered the possibility that excessive quantities of α1 protein generated via the *GAL4/UAS*

expression system could reduce viability. Opposing this idea, prior studies showed that neuronal

overexpression of the WT *UAS-cac-eGFP* transgene renders no discernible structural, behavioral, or

electrophysiological abnormalities. Additionally, overexpressed Cac-GFP protein efficiently localizes to

active zone structures at synapses like the larval neuromuscular junction (NMJ) (20, 22, 28-31).

Using wandering third instar larvae and *elaV(C155)-Gal4* driver, we first checked Cac-GFP

localization of several transgenic lines: WT (published line, *UAS-cac-eGFP[786c]*) (22), DM (*UAS-cac-*

eGFP[DM(2M)]) (this study), SL (*UAS-cac-eGFP[S/L(3-2M)]*) (20), and   RQ (*UAS-cac-eGFP[R/Q(1M)]*) (this study).

We used an anti-GFP antibody to detect Cac-GFP and co-stained with a monoclonal antibody against

the presynaptic ELKS/CAST active zone protein Bruchpilot (Brp) (32). In all cases, Cac-GFP localized as

expected in the larval central nervous system (Figs. 2A-D, red channel). It also predominantly localized

to presynaptic active zone sites at neuromuscular junction (NMJ), as expected (Figs. 2E-H), consistent

with the reports for the original WT constructs (22, 28).

We checked Cac-GFP levels for the different transgenic constructs by Western Blot. We drove

the transgenes neuronally using *elaV(C155)-Gal4* and collected adult heads for analysis, blotting for

Cac-GFP (239 kDa) with anti-GFP and anti-Actin as a loading control. Compared to *elaV(C155)-Gal4* line

controls, each *UAS-cac-eGFP* transgenic line showed an additional, faint band that migrated at a size

consistent with Cac-GFP (Fig. 2I). This band was expressed at comparable levels between the WT, DM,

SL, and RQ lines (Fig. 2J), with no appreciable difference in control levels of actin between the lines (Fig.

2K).

DM-expressing NMJs show small changes in bouton number and glutamate receptor density

Even in the absence of localization or expression-level differences, transgenic mutant Cac-GFP

expression could affect synapse growth or development. Previously, we found no significant

abnormalities in NMJ synaptic growth for SL-expressing flies (20). We extended our analysis to the DM

transgene line by co-staining third instar larval NMJs with antibodies against the Drosophila PSD-95

homolog, Discs Large (Dlg) and the GluRIIA glutamate receptor subunit (Figs. 3A-D). We observed a

very small decrease in the number of Dlg-positive synaptic boutons at DM-expressing NMJs compared to

control WT-expressing NMJs. This decrease was statistically significant only for segment A2, muscle 6/7

(Fig. 3E). We found no significant change in the number of glutamate receptor clusters per NMJ

comparing WT-expressing synapses and DM-expressing synapses (Fig. 3F).

At DM-expressing NMJs, the percentage of the synaptic area covered by the GluRIIA clusters –

normalized to total Dlg area – was slightly but significantly increased (Fig. 3G). We also observed a very

small (~ 7%), but statistically significant increase in the average intensity of GluRIIA-containing glutamate

receptor clusters, normalized to Dlg intensity (Fig. 3H) – while average Dlg intensity itself was unchanged

between WT- and DM-expressing NMJs. In principle, an expansion of the synaptic area capable of

receiving neurotransmitter and a slightly denser packing of GluRIIA-containing receptors could yield

gains in synaptic transmission, especially in quantal size (33). The magnitude of any such change based

on this postsynaptic staining profile (< 10% increase in receptor density) would likely be small, but was

uncertain based on these measures alone. We needed to conduct finer analyses by electrophysiology,

both to document possible changes in synaptic function and also to test for potential presynaptic

contributions when mutant *cac* transgenes were expressed.

DM-, SL- and RQ-expressing NMJs display hyperexcitable evoked synaptic discharges

Coarse phenotypes from neuronally expressed DM and SL transgenes (Fig. 1) suggested

abnormal neuronal or synapse function. Neuronal expression of gain-of-function *UAS-cac-GFP*

transgenes could result in enhanced evoked NMJ neurotransmission in Drosophila, similar to the knock-

in mouse FHM1 models. Expression of both SL and DM significantly increased EPSP amplitudes across

a range of low extracellular $[Ca^{2+}]$ (0.2–0.5 mM) (Fig. 4A, data for 0.4 mM $[Ca^{2+}]_e$ are shown) (20).

Expression of RQ numerically increased average NMJ EPSP amplitudes, but this increase was not

statistically significant (Fig. 4A). Neither estimated quantal content (QC) (Fig. 4B) nor calcium

cooperativity of release for mutant lines were significantly different than WT across this range of 0.2–0.5

mM $[Ca^{2+}]$ (Fig. 4C) (20) (but see more detailed quantal analyses later).

We noted that the EPSP waveforms of RQ, SL, and DM animals were sometimes abnormal (Figs.

4D, E). In addition to increases in EPSP amplitude (Fig. 4D), we observed two distinct EPSP waveform

phenotypes: 1) 'extra discharges' ("ED"), in which supernumerary spiking events occurred during the

decay phase of the EPSP waveform (Fig. 4E, left); and 2) 'shoulders,' in which there was an extended

discharge during the decay phase of the EPSP (Fig. 4E, right), causing a discontinuity in the decay.

These phenotypes were somewhat reminiscent of a broadening of the end-plate potential previously

reported at the NMJs of SL knock-in mice (11). The SL-expressing NMJs produced only the extra

discharge type of abnormal waveform, whereas the RQ-expressing NMJs produced only the shoulder

form (Figs. 4E, F). Consistent with both mutations being present in DM, those NMJs exhibited both types

of abnormal waveform (Fig. 4F). We were also able to generate "RQ only" animals – null *cac^HC129* mutant

larvae rescued to viability by neuronal expression of the RQ transgene. In the case of "RQ only", the

waveform dysfunction closely matched that shown by the RQ-expressing NMJs (Fig. 4F) – i.e. a shoulder

waveform phenotype was present. We were unable to generate "SL only" or "DM only" animals, possibly

due to deleterious gains of function from the SL mutation.

We assessed the severity of the extra discharge phenotype by counting the number of extra

discharge events per 30 evoked pulses (30 recording sweeps at 1 Hz per NMJ). Quantification confirmed

that SL- and DM-expressing NMJs were highly dysfunctional, suggesting neuronal hyperexcitability (Fig.

4G). A previous study in Drosophila demonstrated that higher levels of magnesium in the recording

saline can mask hyperexcitability of neurons (34). Therefore, we conducted additional WT and DM recordings in saline with lowered [$MgCl_2$] (6 mM vs. 10 mM for normal saline, see Materials and Methods). DM-expressing NMJs displayed extreme dysfunction in low $MgCl_2$, both in terms of the percentage of NMJs that produced any supernumerary discharges (100%, Fig. 4H) and the number of extra discharges counted per 30 presynaptic pulses (Fig. 4I). By contrast, WT-expressing NMJs showed almost no such dysfunction (Figs. 4H, I).

Finally, we conducted an additional series of recordings in normal saline, this time with the larval CNS left intact to check if the hyperexcitability might reflect an *in vivo* state for Drosophila larvae. With this experimental maneuver, it was possible to discern "native circuit" differences between WT- and DM-expressing animals. The "CNS intact" condition resulted in some epileptiform spontaneous activity. Compared to WT, the DM-expressing NMJs displayed a high degree of spontaneous epileptiform activity, marked by rapid, continuous large pulses (Fig. 4J; see several measures and explanation in legend).

In conclusion, SL- and DM-expressing NMJs displayed evoked gain-of-function phenotypes consistent with prior mammalian FHM1 mutant analyses. By contrast, RQ-expressing NMJs only displayed a mild gain-of-function shoulder phenotype.

SL- and DM-expressing NMJs show enhanced spontaneous miniature EPSPs with respect to both amplitude and frequency

Mammalian models of FHM1 also show dysfunctional spontaneous neurotransmission (10, 11). We extended our electrophysiological analyses at the Drosophila NMJ to quantal neurotransmission. Immediately, we observed a striking phenotype: for SL- and DM-expressing NMJs, there was an enhancement in both amplitude and frequency of spontaneous miniature EPSPs (mEPSPs) (Figs. 5A-E, Table 2). By contrast, neither an increase in spontaneous mEPSP amplitude nor mEPSP frequency were observed for RQ- or WT-expressing NMJs compared to non-transgenic w^{1118} controls (Figs. 5B-E, Table 2).

Since the mutations examined are in a voltage-gated calcium channel, it was important to document electrophysiological behavior at various calcium concentrations. At both 0.5 mM and 0.4 mM

extracellular $[Ca^{2+}]$, analyses of thousands of individual spontaneous events revealed that increases in spontaneous amplitudes were due to an overall increase in the size distribution of the events at SL- and DM-expressing NMJs (Figs. 5D-G). Additionally, at both 0.5 mM and 0.4 mM extracellular $[Ca^{2+}]$, we noted that the spontaneous events at SL- and DM-expressing NMJs included a minority of gigantic spontaneous events (10-40 mV) that were never seen in w^{1118} or WT-expressing controls or in RQ-expressing NMJs (Figs. 5A, D-G). Notably, these gigantic events were seen in the complete absence of presynaptic nerve stimulation in nerves that had already been severed from the central nervous system.

Table 2. Raw electrophysiological data of selected spontaneous (mEPSP) events

Line	Saline	n	Average mEPSP (mV)	mEPSP Freq (Hz)	Median mEPSP (mV)	Maximum mEPSP (mV)	Resting Membrane V (mV)
w^{1118}	0.5 mM $[Ca^{2+}]$	13	0.86 ± 0.07	4.6 ± 0.4	0.69	11.53	-62.9 ± 0.9
GAL4 > WT		17	0.77 ± 0.05	2.8 ± 0.2	0.67	7.24	-67.8 ± 0.9
GAL4 > DM		25	**1.75 ± 0.22*****	**5.8 ± 0.7****	**1.04$^{\#\#\#}$**	36.91	-68.2 ± 1.2
GAL4 > SL		12	1.32 ± 0.16	**6.7 ± 1.0****	**0.76$^{\#\#\#}$**	44.42	-65.4 ± 0.9
GAL4 > RQ		13	0.77 ± 0.05	3.4 ± 0.4	0.61	3.88	-64.7 ± 1.0
w^{1118}	0.4 mM $[Ca^{2+}]$ (and BHQ controls) + 5 µM BHQ	15	0.70 ± 0.03	3.7 ± 0.2	0.61	3.37	-61.4 ± 0.4
GAL4 > WT		25	0.79 ± 0.05	3.1 ± 0.2	0.66	5.41	-64.2 ± 0.9
GAL4 > DM		17	**1.48 ± 0.13*****	**6.0 ± 0.7****	**1.10$^{\#\#\#}$**	41.17	-62.2 ± 0.6
GAL4 > SL		14	**1.66 ± 0.19*****	**6.6 ± 1.1*****	**1.18$^{\#\#\#}$**	57.90	-65.1 ± 1.6
GAL4 > RQ		12	0.79 ± 0.06	4.3 ± 0.4	0.69	3.38	-66.5 ± 1.7
GAL4 > WT		14	0.79 ± 0.06	2.5 ± 0.8	0.67	4.44	-63.4 ± 0.9
GAL4 > DM		10	1.69 ± 0.42	4.2 ± 0.6	**0.88$^{\#\#\#}$**	31.23	-61.2 ± 1.0
GAL4 > SL		17	1.37 ± 0.32	**3.8 ± 0.5***	**0.91$^{\#\#\#}$**	36.86	-64.7 ± 1.2
w^{1118}	0.2 mM $[Ca^{2+}]$	9	0.64 ± 0.03	3.9 ± 0.3	0.56	2.73	-61.2 ± 0.9
GAL4 > WT		12	0.70 ± 0.06	2.5 ± 0.3	0.59	2.88	-67.1 ± 1.3
GAL4 > DM		19	**1.34 ± 0.08*****	**5.2 ± 0.6***	**1.11$^{\#\#\#}$**	19.95	-61.8 ± 0.9
GAL4 > SL		14	0.94 ± 0.08	**7.0 ± 1.1*****	**0.77$^{\#\#\#}$**	5.93	-65.6 ± 1.6
GAL4 > RQ		8	0.77 ± 0.04	2.0 ± 0.4	0.66	2.88	-58.7 ± 0.5
GAL4 > WT	0 mM $[Ca^{2+}]$	9	0.73 ± 0.05	2.7 ± 0.3	0.61	3.31	-60.1 ± 1.4
GAL4 > DM		10	**1.15 ± 0.09***	8.4 ± 1.6	**0.98$^{\#\#\#}$**	6.59	-58.0 ± 1.6
GAL4 > SL		11	**1.17 ± 0.11****	**12.6 ± 2.3****	**0.93$^{\#\#\#}$**	5.42	-58.7 ± 0.6
GAL4 > WT	0.5 mM $[Ca^{2+}]$ (BAPTA and TTX controls)	18	0.73 ± 0.02	4.2 ± 0.4	0.65	3.04	-66.5 ± 1.2
GAL4 > DM		16	**1.29 ± 0.14*****	**7.4 ± 0.7*****	**0.92$^{\#\#\#}$**	41.56	-64.4 ± 0.8

GAL4 > WT	+10 µM BAPTA-AM	9	0.62 ± 0.06	**1.6 ± 0.2****	**0.51[###]**	3.67	-60.4 ± 1.9
GAL4 > DM		8	1.16 ± 0.10	**2.0 ± 0.2*****	0.97	5.50	-59.5 ± 1.0
GAL4 > WT	+3 µM TTX	7	0.66 ± 0.05	3.8 ± 0.4	**0.54[###]**	2.82	-62.1 ± 1.1
GAL4 > DM		18	1.09 ± 0.07	6.9 ± 0.7	**0.85[###]**	9.82	-62.5 ± 0.7

Average mEPSP amplitudes ± SEM and mEPSP frequencies ± SEM for selected conditions. Also given: median mEPSP amplitudes and maximum mEPSP amplitudes achieved for spontaneous events analyzed (~100 per NMJ). w^{1118} is a non-transgenic wild-type control. WT, DM, SL, and RQ are shorthand for the indicated *UAS-cac-eGFP* transgene being driven in male progeny presynaptically by the *elaV(C155)-Gal4* driver. These data illustrate differential effects when lowering extracellular [Ca^{2+}], chelating Ca^{2+} with BAPTA-AM, or inactivating Na_V channels with TTX. Electrophysiological data were analyzed in two ways as average per NMJ and as cumulative distributions. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. control by one-way ANOVA with Tukey's post-hoc (control is *GAL4 > WT* for most, except in the cases of BHQ, BAPTA-AM, and TTX, in which case the control is the same genotype without treatment). [###] $p < 0.001$ vs. control, examining cumulative distributions by Kruskal-Wallis test with Dunn's post-hoc for multiple comparisons.

It was uncertain if enhanced spontaneous excitability was due to real-time expression of gains of function in Ca_V2 channel gating kinetics, long-term developmental alterations at the synapse – or if both factors could contribute. We considered altered Ca_V2 kinetics. It was previously demonstrated that the SL mutation causes complex biophysical alterations to $Ca_V2.1$ gating function, both by enhancing voltage-dependent activation (8, 9, 35) and by inhibiting calcium-dependent facilitation (36). Follow-up work showed that the drug 2,5′-di(tertbutyl)-1,4,-benzohydroquinone (BHQ) opposes those effects, reversing SL-induced gains of function (20). As part of the same study, we showed that BHQ restores a form of short-term synaptic plasticity at SL-expressing Drosophila NMJs (20). We extended those prior analyses of BHQ effects on Ca_V2 gating, this time by examining the distribution of spontaneous events. We found that acute application of 5 µM BHQ was partially effective at reversing the increased size distribution of events for SL- and DM-expressing NMJs, without changing the distribution of WT events (Figs. 5F, G; Table 2). Notably, 5 µM BHQ did not abolish gigantic events (Figs. 5F, G). Interestingly, a higher concentration of 10 µM BHQ did abolish gigantic events for SL- and DM-expressing NMJs, but it also significantly decreased the size distribution of WT mEPSPs, which could indicate off-target

postsynaptic effects (Fig. 5F). Our BHQ application data are consistent with the idea that that

spontaneous neurotransmission gain-of-function phenotypes are driven in part through gating changes at

Ca_V2 channels.

To test if long-term developmental alterations at the synapse could also play a role, we

engineered stage-specific *UAS-cac* transgene expression. We utilized the temperature-sensitive

Gal80^{TS}/TARGET system to temporally control expression of the DM transgene (37). To conduct this

experiment, we generated *elaV(C155)-Gal4 >> UAS-cac-eGFP^{DM}* animals with a ubiquitous *Gal80^{TS}*

transgene (37). Gal80^{TS} protein halts Gal4-induced gene expression at permissive temperatures (25℃)

but not at restrictive temperatures (29℃). For our experiment, animals raised at 25℃ throughout life had

no discernible spontaneous neurotransmission hyperexcitability (Fig. 5H). By contrast, animals started at

25℃ and shifted to 29℃ for the final 24 or 48 hours before third instar NMJ recording showed

progressively more spontaneous hyperexcitability (Figs. 5H, I). This experiment indicates that

developmentally regulated expression of gain-of-function Ca_V2 channel subunits also underlies some of

the spontaneous neurotransmission gain-of-function phenotypes.

Gigantic spontaneous events require extracellular calcium and sodium channel activity

Prior work proposed that mammalian neuronal dysfunction downstream of FHM1 mutations may

be calcium-dependent (12). We tested whether the observed effects on quantal size in our model could

be calcium-dependent. First, we reduced the extracellular $[Ca^{2+}]$ in the recording saline to 0.2 mM.

Consistent with classic characterizations of Drosophila NMJ properties (38), low calcium did little to

change the distribution of mEPSP size, the median mEPSP size, or the 25^{th}-75^{th} percentiles of mEPSP

size – all of which remained normal for WT and elevated for SL- and DM-expressing NMJs (Figs. 6A, B,

Table 2). However, lowering extracellular $[Ca^{2+}]$ almost completely abrogated gigantic (10-40 mV)

spontaneous events at SL- and DM-expressing NMJs – and it completely eliminated the very largest

ones (Figs. 6A, B). This suggested that these gigantic events somehow relied on a sufficient driving force

of presynaptic calcium influx – and potentially on spontaneous presynaptic nerve firing.

We extended these analyses by altering the recording saline in three additional ways: 1) zero extracellular calcium; 2) adding the membrane-permeable calcium chelator, 1,2-Bis (2-aminophenoxy) ethane-N,N,N',N'-tetra acetic acid tetrakis (acetoxymethyl ester) (BAPTA-AM, 10 µM); or 3) adding tetrodotoxin (TTX, 3 µM) to block voltage-gated sodium channels. We compared WT-expressing and DM-expressing NMJs (and SL-expressing NMJs in the case of zero calcium). All three manipulations produced a similar effect on mEPSP size for the gain-of-function mutants: an elimination of gigantic spontaneous events, but a persistence of overall elevated mEPSP size (Figs. 6C-F, Table 2). By contrast, these manipulations had little to no effect on the distribution of mEPSP amplitudes at WT-expressing NMJs (Figs. 6C-D, Table 2).

Finally, we recorded spontaneous events in more *in vivo*-like condition, using an intact CNS, without severing the motor nerve. In order to do this, we revisited the intact CNS condition (Fig. 4J) – this time adding TTX to the recording saline (0.5 mM $[Ca^{2+}]$). This left the full network anatomy intact, while quieting any spontaneous epileptiform activity. Under these conditions, the spontaneous event amplitude profile of DM-expressing NMJs was still larger than that of WT-expressing NMJs – and as expected, there were no gigantic events (Figs. 6G, H). Interestingly, however, the difference between WT-expressing NMJs and DM-expressing NMJs was muted (Figs. 6G, H; compare to Figs. 5D, E). These data suggest that in living animals, network effects could potentially influence the spontaneous gain-of-function activity.

Large spontaneous events are due to multi-vesicular release

The presence of gigantic spontaneous mEPSPs that were sensitive to low calcium, calcium chelation, and TTX treatment suggested the possibility of spontaneous multi-vesicular release at SL- and DM-expressing NMJs. If this were true, traditional analysis of spontaneous mEPSPs would result in an overestimation of average quantal size (Fig. 5B) and underestimation of average QC (Fig. 4B) for SL- and DM-expressing NMJs.

We utilized the method of failures to better resolve questions about quantal size and QC. At very low concentrations of extracellular calcium, synapses like the NMJ are essentially limited to a one-or-

none evoked response in which stimulation of the presynaptic nerve either leads to the release of a

single vesicle or fails to release any vesicles (39). By conducting failure analyses, it is possible to

measure the distribution of quantal events and also to estimate QC in a way that eliminates confounds of

higher concentrations of calcium. First, we conducted failure analysis recordings at 0.14 mM $[Ca^{2+}]_e$ for

WT-, DM-, and SL-expressing NMJs (Figs. 7A-C). For this condition, the evoked events for SL- and DM-

expressing NMJs were far larger on average than those observed WT-expressing NMJs (Fig. 7C –

EPSP). This was due to a large proportion of events of > 2 mV for the SL- and DM-expressing conditions

(compare Figs. 7A, B). Furthermore, even in this low level of extracellular Ca^{2+}, many of the DM and SL

events represented multi-vesicular release rather than the release of a single large vesicle. We

calculated values of QC of > 2 for both mutant conditions at 0.14 mM $[Ca^{2+}]_e$ (QC = m = ln[(# trials)/(#

failures)] (40)) (Fig. 7C).

To test if lower calcium could generate a leftward shift in event size, we applied a more restrictive

condition of 0.1 mM $[Ca^{2+}]_e$ to DM-expressing NMJs. At 0.1 mM $[Ca^{2+}]_e$ the proportion of failures was very

high for DM-expressing NMJs, with events over 4 mV all but absent, and events greater than 1.5 mV also

less prevalent (Fig. 7D). The first peak in the distribution of events, which is reflective of single vesicle

size (40), was centered near 0.7 mV (Fig. 7D), a value consistent with single-vesicle responses of normal

size for the Drosophila NMJ (38). Together, these data suggested that the observed large events at SL-

and DM-expressing NMJs – regardless of whether spontaneous or failure analysis-evoked – were likely

due to multi-vesicular release (see Fig. 7E, spontaneous and failure analyses distributions side-by-side).

If larger spontaneous events are multi-vesicular (or at the very least include a proportion of multi-

vesicular events), this property should also be reflected in slowed spontaneous event rise time kinetics.

We analyzed the rise time kinetics of several thousand spontaneous events for w^{1118}, WT-, SL-, DM-, and

RQ-expressing NMJs. Average rise times were slowed only for DM-expressing NMJs (Fig. 7F). However,

the rise times for larger events were markedly slower for *all* genotypes, not just DM (Fig. 7G). For SL-

and DM-expressing NMJs there was a much larger proportion of such events. Collectively, our data

suggest that large events (> 2 mV) include several that are multi-vesicular.

PLCβ loss genetically suppresses spontaneous excitability

For SL- and DM-expressing NMJs, we hypothesized that specific cellular cues could dictate the various electrophysiological phenotypes we documented: multi-vesicular quantal events, gigantic TTX-sensitive spontaneous events, and enhanced NMJ excitability. We inquired as to what the molecular nature of those cues might be. Our experiments indicated that intracellular calcium or intracellular calcium signaling processes might be important (Fig. 6). Additionally, recent data from the mouse calyx of Held demonstrated that S218L knock-in synapses have enhanced resting intracellular calcium (12). We hypothesized that altered intracellular calcium signaling or handling could impact myriad intracellular signals and investigated which signaling pathways might be relevant. This line of inquiry spurred a genetic approach examining regulators of intracellular calcium to test if inhibition of any of these factors may influence gain-of-function Ca_V2 phenotypes at the synapse (Fig. 8A). We sought to identify suppressors capable of reversing gains of Ca_V2 function caused by the SL and DM transgenes.

Prior studies of Drosophila NMJ homeostatic synaptic plasticity, which involves the potentiation of Ca_V2 function, suggested some possible candidate molecules (41, 42). Additionally, we previously showed that the Drosophila PLCβ homolog *phospholipase-C at 21C* (*Plc21C*) is necessary for this same neuronal homeostatic potentiation mechanism (43). Plc21C is one of two Drosophila Phospholipase-Cβ (PLCβ) family members, and is expressed in the nervous system (44). Canonically, PLCβ proteins cleave phosphatidylinositol 4,5-bisphosphate (PIP_2) to generate soluble inositol triphosphate (IP_3), as well as membrane-bound diacylglycerol (DAG). These signaling factors influence synaptic transmission in a variety of ways, including direct modulation of Ca_V2 (45), and they have been shown to act at several synapses, including the NMJ (46-51).

We targeted *Plc21C* gene expression in neurons with a previously verified *UAS-Plc21C(RNAi)* construct, *Plc21C^{GD11359}* (43, 52). Compared to the NMJs of w^{1118} and WT controls, those in which only *Plc21C* had been knocked down presynaptically exhibited no discernable baseline changes in mEPSP size (Fig. 8B, Table 3) – or as previously documented, EPSP size, or QC (43). By contrast, in DM-expressing NMJs such *Plc21C* knockdown alleviated aspects of NMJ hyperexcitability. Specifically, there

402 was a leftward shift in the distribution of spontaneous events (Fig. 8B, Table 3). Interestingly, there was

403 not a significant reversal of the enhanced mEPSP frequency phenotype (Table 3).

405 **Table 3. Raw electrophysiological data of spontaneous (mEPSP) events – impairment of**
406 **intracellular Ca^{2+} release pathway.**

Line	Experiment (all 0.5 mM Ca^{2+})	n	Average mEPSP (mV)	mEPSP Freq (Hz)	Median mEPSP (mV)	Maximum mEPSP (mV)	Resting Membrane V (mV)
w^{1118} females	Plc21C RNAi baseline	27	0.81 ± 0.05	4.3 ± 0.3	0.64	7.16	-62.2 ± 0.6
Plc21C(RNAi) males		12	0.80 ± 0.06	**2.0 ± 0.2****	0.61	4.50	-66.5 ± 0.9
Plc21C(RNAi) females		6	0.83 ± 0.02	3.8 ± 0.9	0.59	7.68	-61.6 ± 0.3
GAL4 > DM females	suppression of DM	19	1.28 ± 0.08	5.8 ± 0.5	1.01	7.62	-67.0 ± 1.2
GAL4 > DM + Plc21C(RNAi) females		13	**0.78 ± 0.04*****	5.2 ± 0.6	**0.68**###	3.89	-64.9 ± 1.6
GAL4; $itpr^{ug3}$/+	suppression of DM	9	0.74 ± 0.04	3.7 ± 0.2	0.64	3.34	-61.3 ± 0.6
GAL4 > DM; $itpr^{ug3}$/+		14	**1.05 ± 0.06***	6.1 ± 0.5	**0.86**###	16.21	-65.8 ± 0.7
GAL4; RyR^{E4340K}/+	suppression of DM	13	0.83 ± 0.03	3.3 ± 0.3	0.70	3.38	-63.0 ± 0.5
GAL4 > DM; RyR^{E4340K}/+		17	**0.91 ± 0.04*****	5.1 ± 0.8	**0.76**###	5.13	-62.5 ± 0.9
GAL4 > WT	XestC and LiCl controls	19	0.80 ± 0.02	2.3 ± 0.3	0.71	2.59	-69.6 ± 1.1
GAL4 > DM		30	1.63 ± 0.13	6.6 ± 0.7	1.08	51.03	-65.8 ± 0.7
GAL4 > WT	+ 5 µM XestC	7	0.89 ± 0.03	3.9 ± 0.9	0.78	3.80	-69.8 ± 2.2
GAL4 > DM		14	**1.14 ± 0.11***	6.4 ± 1.1	**0.81**###	30.76	-67.2 ± 1.3
GAL4 > WT	+ 10 mM LiCl	11	0.81 ± 0.03	3.2 ± 0.4	0.73	4.09	-66.6 ± 1.3
GAL4 > DM		12	**1.2 ± 0.06***	4.1 ± 0.4	**0.98**###	6.34	-68.8 ± 1.7

Average mEPSP amplitudes ± SEM and mEPSP frequencies ± SEM for selected
experimental conditions. Also given are the median mEPSP amplitude and the maximum
mEPSP amplitudes achieved for all spontaneous events analyzed per genotype (~100 per

NMJ). w^{1118} is a non-transgenic wild-type control. WT and DM are shorthand for the
indicated *UAS-cac-eGFP* transgene being driven in progeny presynaptically by the
elaV(C155)-Gal4 driver. This table illustrates differential effects when impairing an
intracellular calcium release signaling pathway through mutation of the *Plc21C*, *itpr*, and
RyR genes, or through pharmacological application of Xestospongin C or LiCl.
Electrophysiological data were analyzed in two ways as average per NMJ and as
cumulative distributions. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. control by one-way
ANOVA with Tukey's post-hoc (for all cases, the appropriate control is the same genotype
without treatment; some control data are on Table 2). # $p < 0.05$, ## $p < 0.01$, ### $p < 0.001$
vs. control to examine cumulative distributions by Kruskal-Wallis test with Dunn's post-
hoc.

IP$_3$R and RyR point mutations strongly suppress hyperexcitability

We hypothesized that Plc21C could exert effects on spontaneous neurotransmission via one of

several components of its canonical signaling pathway (e.g. PIP$_2$, DAG, or IP$_3$). Notably, IP$_3$ acts through

the IP$_3$ receptor (IP$_3$R), an intracellular calcium channel located on the endoplasmic reticulum (ER). At

other model synapses, release of Ca^{2+} from the intracellular stores can promote the release of

neurotransmitter-laden vesicles and contribute to the amplitudes of spontaneous events (53-56).

Moreover, IP$_3$R has been proposed to play a role in spontaneous vesicle release through calcium-

induced calcium release (CICR) (57), and increased ER Ca^{2+} release was recently shown to potentiate

synaptic transmission at the Drosophila NMJ (58).

We examined the Drosophila IP$_3$R gene (*itpr*). Homozygous *itpr* loss-of-function mutations are

lethal, so we tested a heterozygous loss-of-function condition. Since IP$_3$R clusters consist of multiple

units, we hypothesized that we might be able to partially disrupt them through a loss-of-function point

mutation, *itpr^{ug3}*, a mutant possessing a missense mutation in the IP$_3$R ligand-binding domain (59).

itpr^{ug3}/+ phenocopied *Plc21C* knockdown at DM-expressing NMJs: the mEPSP amplitude was partially

reduced toward WT levels, and the number of giant, spontaneous events was diminished (Figs. 8C, D,

Table 3). Importantly, on its own *itpr^{ug3}/+* did not significantly affect the baseline amplitude or distribution

of mEPSPs (Fig. 8C, Table 3). Finally, as with the RNAi experiment, the increased mEPSP frequency

phenotype was not suppressed (Table 3).

We performed analogous experiments with a Drosophila ryanodine receptor gene (*RyR*)

mutation. Tetrameric RyR channels have been reported to contribute to CICR downstream of IP$_3$Rs (57).

Additionally, gigantic spontaneous miniature potentials at other model synapses are mediated by RyR

and rapid expulsion of calcium from presynaptic stores (60-64). We found that the heterozygous *RyR*

point mutant RyR^{E4340K}/+ (65) almost completely suppressed the increased average mEPSP amplitude in

the DM-expressing background (Figs. 8C, E, Table 3). Additionally, the gigantic spontaneous events

were abrogated (Figs. 8C, E, Table 3). Control recordings showed that RyR^{E4340K}/+ did not affect the

baseline amplitude or distribution of mEPSPs (Figs. 8C, E). As with *Plc21C* and *itpr*, impairment of *RyR*

function did not significantly suppress the enhanced mEPSP frequency phenotype of DM-expressing

NMJs (Table 3).

Because the RyR^{E4340K}/+ background provided such a strong suppression of spontaneous

mEPSP hyperexcitability at DM-expressing NMJs, we checked if it could also suppress hyperexcitability

in the context of evoked excitation. As shown before, when incubated in low extracellular magnesium,

100% of the DM-expressing NMJs showed a hyperexcitability dysfunction, with high expressivity of extra,

epileptiform discharges (Figs. 3H,I; 8F). In a heterozygous RyR^{E4340K}/+ genetic background, this

hyperexcitability phenotype was partially suppressed, in terms of both the penetrance of NMJs with extra

evoked discharges and the expressivity of the extra discharge dysfunction at individual NMJs (Fig. 8F).

On its own, the RyR^{E4340K}/+ condition shows almost no baseline hyperexcitability phenotype (Fig. 8F).

Spontaneous mEPSP hyperexcitability can be suppressed pharmacologically

Our data for genetic manipulations affecting Plc21C, IP$_3$R, and RyR show that it is possible to

attenuate DM-induced gain-of-function mEPSP amplitude and excitability phenotypes by genetically

impairing factors known to promote intracellular Ca^{2+} release. We wondered if pharmacological

manipulations could also be effective. We turned to two agents to test this idea: lithium (10mM LiCl in

larval food) and Xestospongin C (5 μM in recording saline). Chronic exposure to LiCl inhibits inositol

monophosphate phosphatase, eventually resulting in a disruption of the recycling process that generates

PIP$_2$ (66, 67). Xestospongin C has been previously characterized as a membrane-permeable inhibitor of

IP$_3$ receptors (68, 69). Either chronically feeding larvae LiCl or applying Xestospongin C to the recording

bath caused a significant leftward shift in the overall size distribution of spontaneous amplitudes (Figs.

9A-C), reminiscent of the effects observed for *Plc21C*, *itpr*, and *RyR* losses of function. The acute

Xestospongin C application seemed to exert a stronger suppression effect in this regard, while the

chronic LiCl application exerted a stronger suppression of the gigantic spontaneous events (Figs. 9A-C,

Table 3). Notably, neither pharmacological manipulation diminished baseline spontaneous

neurotransmission in WT-expressing control NMJs, nor did either manipulation significantly suppress the

elevated mEPSP frequency phenotype for DM-expressing NMJs (Figs 9A-C, Table 3).

Mutations targeting intracellular calcium release signaling do not restore viability

Since genetic mutations that target intracellular calcium release ameliorate hyperexcitability

phenotypes, we reasoned that the same (or similar) mutations might ameliorate the lethality phenotypes

associated with expressing the DM transgene. We conducted lethality test crosses and progeny counts

in a similar manner as before (Table 1). This time, however, we crossed females bearing the *UAS-cac-*

eGFP^{DM} transgene to males carrying both the *elaV(C155)-Gal4* driver and a collection of loss-of-function

genetic manipulations on *Drosophila melanogaster* Chromosome II for the *Plc21C*, *RyR*, or *Gq* genes. In

addition to *Plc21C* and *RyR*, we chose *Gq* because canonical PLCβ signaling is downstream of Gαq

function. Our prior work showed that *Plc21C* and *Gq* play a role in the maintenance of homeostatic

plasticity at the NMJ (43). The hypothesis to test was that female progeny carrying the driver, the DM

transgene, and the intracellular calcium release manipulation could have improved viability versus to

female progeny carrying only the driver and the DM transgene. Male progeny siblings would not carry the

driver – and would therefore not express the DM transgene – and could be used to control for

parameters affecting lethality, independent of the DM transgene.

As expected, female progeny carrying the driver, the DM transgene, and no balancer

chromosome had reduced viability compared to their male sibling counterparts (Table 4; see "& = +"

data). However, introducing heterozygous loss-of-function manipulations affecting *Plc21C*, *RyR*, and *Gq*

did not ameliorate this phenotype. Instead, those manipulations almost always further reduced viability,

often strongly (Table 4). The effect was particularly strong for all *Plc21C* and *Gq* loss-of-function

conditions examined (Table 4). For *RyR*, the effect was strong only for the *RyR^{16}* deletion allele (Table

4). Heterozygous *RyR* point mutant manipulations did not further enhance lethality in a statistically

significant way, but they did not ameliorate the lethality phenotype either (Table 4). No manipulation

examined resulted in significantly higher male lethality, compared to control males (Table 4).

Table 4. Loss-of-function mutations in an intracellular Ca^{2+} store release pathway enhance adult
lethality phenotypes

w/w; CyO-GFP/UAS-cac-eGFP^{DM} x *C155/Y; CyO-GFP/&*	Non-Curly females	Curly females	Normalized Viability Index for & (female)	Non-Curly males	Curly males	Normalized Viability Index for & (male)
& = +	100 (14.1%)	218 (30.7%)	100.0	152 (21.4%)	240 (33.8%)	138.1 [#]
Plc21C(RNAi)	**25 (4.9%)**	**183 (36.0%)**	**29.8 ******	106 (20.9%)	194 (38.2%)	119.1 [####]
Plc21C$^{Df2L(BSC4)}$	**5 (4.9%)**	**44 (42.7%)**	**24.8 ****	18 (17.4%)	36 (35.0%)	109.0 [##]
Plc21C$^{Df(p60A)}$	**14 (5.4%)**	**86 (33.0%)**	**35.5 *****	55 (21.1%)	106 (40.6%)	113.1 [###]
Plc21C^{MI01911}	**31 (6.6%)**	**162 (34.4%)**	**41.7 *****	102 (21.7%)	176 (37.4%)	126.3 [####]
RyR^{E4340K}	19 (8.1%)	49 (20.8%)	84.5	55 (23.3%)	113 (47.9%)	106.1
RyR^{R4305C}	17 (6.3%)	57 (21.2%)	65.0	76 (28.3%)	119 (44.2%)	139.2 [#]
RyR^{k0943}	16 (9.9%)	52 (32.1%)	67.1	40 (24.7%)	54 (33.3%)	161.5 [#]
RyR^{16}	**16 (5.0%)**	**94 (29.3%)**	**37.1 *****	65 (20.2%)	146 (45.5%)	97.1 [##]
RyR^{Q3878X}	15 (7.2%)	47 (22.7%)	69.6	59 (28.5%)	86 (41.5%)	149.6 [#]

RyR^{Y4452X}	18 (7.8%)	47 (20.3%)	83.5	51 (22.1%)	115 (49.8%)	96.7
Gq^{28}	**14 (4.4%)**	**94 (29.6%)**	**32.5 *****	70 (22.0%)	140 (44.0%)	109.0 ####
Gq^{221c}	**13 (3.5%)**	**121 (32.4%)**	**23.4 ******	101 (27.0%)	139 (37.2%)	158.4 ####

Viability enhancement/suppression test crosses were performed utilizing *w/w; CyO-GFP/UAS-cac-eGFP*^{DM} virgin females x *elaV(C155-Gal4)/Y; CyO-GFP/(& = mutant or UAS-RNAi or +)* males. Balancer or non-Balancer (Curly and non-Curly) female and male progeny were counted. Raw progeny counts and relative proportions are shown. Changes in the proportion of non-Curly female progeny acquired could indicate a suppression or enhancement effect on viability by *&*. A normalized viability index number was set = (proportion of non-Curly females for *&* tested)/(proportion of non-Curly females where *& = +*). In all cases for losses of function of *Plc21C*, *RyR*, and *Gq* gene function, the normalized viability index decreased numerically. Fisher's exact tests were performed for each cross to test for differences in female to male Cy:non-Cy ratios. This type of analysis controlled for any lethality caused by the *&* genetic manipulation itself. For female progeny ratios, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ vs. *& = +* by Fisher's exact test between crosses. Within a cross, # $p < 0.05$, ## $p < 0.01$, ### $p < 0.001$, #### $p < 0.0001$ for male vs. female viable progeny ratios. Most crosses meet a significant threshold by this latter criterion because the *DM* transgene causes lethality itself, and it is only expressed in females.

DISCUSSION

We generated fruit flies designed to mimic the effects of FHM1-inducing Ca$_V$2.1 channel mutants, R192Q and S218L. Flies expressing the SL and DM transgenes for Drosophila Ca$_V$2/Cacophony displayed overt phenotypes, including reduced viability (Fig. 1). They also displayed synaptic phenotypes, including enhanced evoked excitability (Fig. 4), stark increases in quantal size and frequency (Fig. 5), giant, spontaneous, sodium channel-dependent events (Figs. 5, 6), and enhanced probability of release at very low calcium (Fig. 7). All of these neurotransmission phenotypes occurred without major alterations in active zone localization or overall synaptic architecture (Figs. 2, 3). RQ-expressing NMJs had only a mild phenotype: EPSP discharges with shoulder-like waveforms (Fig. 4). Genetic knockdown of Drosophila PLCβ or mutations affecting the receptors that gate intracellular

calcium stores (IP$_3$ receptor and Ryanodine receptor) partially alleviated some of the electrophysiological

phenotypes (Fig. 8), as did pharmacological manipulations targeting the same processes (Fig. 9). These

results suggest that intracellular Ca^{2+} signaling through IP$_3$ receptors and Ryanodine receptors could

influence physiological dysfunction in a gain-of-function Ca$_V$2 background (Fig. 9D). Additionally, given

the ability of TTX to block gigantic spontaneous events – and given our ability to quiet that phenotype

through genetic and pharmacological means – impairment of the IP$_3$ Receptor/Ryanodine Receptor

pathway may limit spontaneous neuronal firing by as-yet undetermined mechanisms (Fig. 9D).

Similarities between fly mutations and FHM1-causing human mutations

Evoked Neurotransmission. Our discovery that SL- and DM-expressing Drosophila NMJs

displayed increased evoked excitation, especially at low [Ca^{2+}]$_e$ (Figs. 4, 7) (20), was consistent with

findings from diaphragm NMJs in SL knock-in mice (11). In that context, the end-plate potential (EPP)

amplitudes were significantly increased at low levels of calcium (0.2 mM), but did not differ from those at

wild-type NMJs at physiological calcium (2 mM) (11). Interestingly, at the SL knock-in calyx of Held,

excitatory postsynaptic currents (EPSCs) were increased, but this effect was most pronounced at high

levels of [Ca^{2+}]$_e$ (12). The epileptiform EPSP discharges caused by expression of the SL-containing

transgenic constructs in flies (Fig. 4) were reminiscent of the EPP broadening at SL knock-in NMJs (11).

Finally, the severity of the dysfunction in the Drosophila NMJ waveform in the context of decreased

extracellular magnesium (6 mM) (Figs. 4, 8) was consistent with a marked increase in calcium current in

response to long action potential waveforms in calyces of Held expressing the RQ or SL mutant protein

(12, 14).

Enhanced Quantal Frequency. The enhanced mEPSP frequency at SL- and DM-expressing

Drosophila NMJs (Fig. 5, Table 1) was reminiscent of observations in prior FHM1 studies. In the RQ and

SL knock-in mice, the NMJs exhibited significant increases in the frequency of mEPPs (9-11).

Theoretically, this spontaneous activity could ultimately lead to a buildup of intracellular calcium or a

change in intracellular calcium dynamics. In support of this view, at the calyx of Held in SL knock-in mice

the frequency of spontaneous mEPSCs was enhanced and resting [Ca^{2+}]$_i$ was elevated (12). In that

case, the increase in quantal frequency was partially reversed by adding the cell-permeable calcium

chelator EGTA-AM (12).

Evidence from several model synapses suggests that Ca_V2 channels can play a prominent role in

spontaneous release. In granule cells of the hippocampus, stochastic activity of $Ca_V2.2$-type channels

potentiates spontaneous miniature events, and the application of either BAPTA-AM or EGTA-AM is

sufficient to inhibit them (70). Other studies demonstrated that P/Q-, N-, and R-type calcium channels

also promote spontaneous release (71). Notably, the differences in the spontaneous miniature

phenotype between mice harboring the SL and RQ knock-in substitution, or fruit flies expressing

mimicking mutations, suggest that the differences in cellular outcomes occur downstream of the Ca_V2

channel. This highlights a need for genetic approaches to uncover pathways that might contribute to the

divergent phenotypes, as well those that are shared, to better understand the biology downstream of

gain-of-function calcium channel activity.

Differences between fly mutations and FHM1-causing human mutations

Quantal amplitudes. FHM1 mutations have been shown to enhance spontaneous miniature

quantal release frequency in other systems (8-11), but there has been no report of increases in

spontaneous miniature quantal size due to these mutations. In theory, an increase in the amplitude of

mEPSP events at the Drosophila NMJ could be explained by an alteration to the expression and

localization of postsynaptic proteins. Yet immunostaining of postsynaptic markers showed only a slight

increase in postsynaptic glutamate receptor clustering (Figs. 3G, H). Instead, a combination of quantal

analyses (Figs. 5-7) pointed to alterations to the nature of spontaneous, presynaptic vesicle release –

namely, that a certain percentage of quantal events in SL- and DM-expressing NMJs were multi-

vesicular.

Why do SL- and DM-expressing NMJs in Drosophila show spontaneous multi-vesicular release

while other models do not? The synaptic preparation examined is likely to be critical. Evidence from other

systems has demonstrated that calcium-channel activity can have a profound effect on quantal size. For

example, work at the *C. elegans* NMJ has demonstrated that calcium from intracellular and extracellular

sources combines to dictate quantal size and frequency (54). Additionally, spontaneous miniature events

with large amplitudes ("maximinis") have been documented at fast inhibitory synapses of the cerebellum

(53, 72). Similar to the NMJ activity documented in our study, these maximinis rely on the ability of

ryanodine-sensitive stores to support spontaneous calcium transients large enough to cause multi-

vesicular release. It is possible that the architecture of a giant synapse like the Drosophila NMJ – which

contains hundreds of active zones clustered into individual boutons and has a low level of spontaneous,

multi-vesicular release (73) – makes it exquisitely sensitive to small changes in intracellular calcium

handling.

Evoked waveforms. As is the case for the Drosophila NMJ EPSPs (Fig. 4), the diaphragm NMJ of

FHM1 knock-in mice displayed EPP broadening (11). However, the extra, epileptiform discharges we find

at the Drosophila NMJ do not seem to be documented for the mammalian NMJ. An instructive parallel

may be drawn between our data and cultures of Drosophila giant neurons, in which manipulation of the

voltage-gated potassium current generated altered waveforms, including extra and extended discharges

(74, 75). This raises the possibility that some aspects of the FHM1 phenotypes may be caused by the

perturbation of other voltage-activated currents, and by synapse excitability more generally (76). This

possibility is also consistent with the fact that mutations in the Na^+/K^+ ATPase gene cause pure FHM

(77). Given the effectiveness of the Drosophila system for uncovering complex relationships amongst ion

channel activities, in particular potassium currents, the fly may be a good model for studying the cellular

bases of disorders such as FHM1 (78-82).

Implications for gain-of-function calcium channel and migraine signaling pathways

In this study, we have shown that the expression of gain-of-function transgenes bearing

mutations mimicking those associated with migraine in humans causes overt spontaneous and/or evoked

phenotypes. Further, we have shown that genetic or pharmacological impairment of an intracellular

calcium release signaling pathway suppresses some gain-of-function Ca_V2 electrophysiological

phenotypes (Figs. 8, 9). Moreover, impairment of RyR function partially suppresses evoked

hyperexcitability (Fig. 8F). A simple explanation – but certainly not the only explanation – is that $Ca_V2.1$

gains of function potentiate the baseline activities of the IP$_3$R and RyR channels (83-86). Further work

would be required to test this possibility rigorously.

One strength of Drosophila is the power of genetic manipulation. The finding that components

involved in a signaling pathway controlling intracellular calcium release can be targeted to counteract

electrophysiological problems caused by Ca$_V$2.1 gain-of-function mutations is novel and has not been

directly suggested by other studies of FHM1 animal models. However, in hindsight, the implication of

PLCβ activity and intracellular calcium in hyperexcitability does fit with results of prior studies. One recent

RNA profiling analysis of the cerebellum of SL knock-in mice revealed an overrepresentation of several

signaling pathway components, including PLCβ (87). Moreover, PLCβ and the release of calcium from

intracellular stores have been implicated in signaling by calcitonin gene related peptide (CGRP) (88-90),

whose levels are correlated with migraine (91-93). Further investigation will be needed to establish

whether there is a causative link between the action of intracellular calcium stores either in inducing

migraine or in precipitating neurological events that precede some forms of migraine, like aura and

cortical spreading depression.

Finally, we must note that impairing intracellular calcium release signaling pathways did not

reverse all phenotypes associated with SL- and DM-expressing NMJs. For instance, in the case of

increased mEPSP frequency, there is no significant reversal (Table 3). Moreover, in the case of DM-

induced lethality, there is often a significant enhancement, rather than a suppression (Table 4). These

results point to the fact that gain-of-function Ca$_V$2 substitutions may cause multiple, separable, pleiotropic

effects. One possibility worth future consideration is that the neuronal hyperexcitability phenotypes are

somehow protective for fruit fly viability or health – or are a reflection of a protective process that gets

blunted when PLCβ and RyR are diminished.

MATERIALS AND METHODS

Gain-of-function *cacophony* constructs: To generate *UAS-cac-eGFP^{SL}* transgenes, we used PCR to

alter the serine 161 codon to leucine in the pUAST-based *UAS-cac-eGFP* DNA construct (20, 22). This

substitution corresponds to S218L in mammalian CACNA1A. To generate *UAS-cac-eGFP^{RQ}* transgenes,

we used PCR to change the arginine 135 codon to glutamine. This substitution corresponds to R192Q in

mammalian CACNA1A. For the *UAS-cac-eGFP*^{DM} transgene, both mutations were incorporated into the

same *UAS-cac-eGFP* construct using PCR to link the overlapping RQ and SL fragments. Transgenic

lines were generated by injection of *UAS-cac-eGFP* constructs into a *w*^{1118} background (The Best Gene,

Chino Hills, CA) and mapped and backcrossed.

Drosophila Stocks, Genetics, and Husbandry: Animals used for viability counts and electrophysiology

were generated by driving neuronal expression of *UAS-cac-eGFP* transgenes with *elaV(C155)-GAL4*

(23). Multiple *UAS-cac-eGFP* transgenic lines were initially examined to control for possible differences

caused by independent *UAS* genomic insertions: WT: *UAS-cac-eGFP*^{786c} (22), *UAS-cac-eGFP*^{422a} (22);

SL: *UAS-cac-eGFP*$^{S/L(3-2M)}$, *UAS-cac-eGFP*$^{S/L(3-6M)}$, *UAS-cac-eGFP*$^{S/L(3-8M)}$; RQ: *UAS-cac-eGFP*$^{R/Q(1M)}$,

UAS-cac-eGFP$^{R/Q(2-4M)}$; DM: *UAS-cac-eGFP*$^{DM(1M)}$, *UAS-cac-eGFP*$^{DM(2M)}$.

w^{1118} (94) was used as a non-transgenic wild-type control. Other Drosophila mutant alleles used

were *Df2L(BSC4)* (K. Cook to flybase.org), *Plc21C*^{p60A} (95), *Plc21C*^{MI01911} (96), *itpr*^{ug3} (97), *RyR*^{E4340K} (65),

RyR^{R4305C} (65), *RyR*^{k09q3} (98), *RyR*^{16} (99), *RyR*^{Q3878X} (65), *RyR*^{Y4452X} (65), *Gq*^{28} (100), and Gq^{221c} (101).

Mutant Drosophila stocks were obtained either from the Bloomington Drosophila Stock Center (BDSC,

Bloomington, Indiana) or directly from the labs that generated them. The *UAS-Plc21C(RNAi)*

transformant lines 26557 and 26558 (*Plc21C*^{GD11359}) (102) were obtained from the Vienna Drosophila

Resource Center (VDRC, Vienna, Austria). A *Gal80*^{TS} expression line (37) was employed for a temporal

Gal4 expression experiment. Flies were raised at 25°C (or 29°C for one temperature shift experiment) in

humidity- and light-controlled Percival incubators (Geneva Scientific, Fontana, WI), in glass vials on a

standard Drosophila food containing water, agar, molasses, yellow cornmeal, and yeast.

Electrophysiology and Analysis: Wandering third-instar larvae were selected for analysis. Larvae were

dissected in a modified HL3 saline with the following components (and concentrations): NaCl (70 mM),

KCl (5 mM), MgCl$_2$ (10 mM or 6 mM as noted), NaHCO$_3$ (10 mM), sucrose (115 mM = 3.9%), trehalose

(4.2 mM = 0.16%), HEPES (5.0 mM = 0.12%), and CaCl$_2$ (0.5 mM, unless otherwise noted). The central

nervous system was removed, except for specific instances noted (Figures 4 and 6). Pharmacological

agents tetrodotoxin (TTX, Tocris/R&D Systems), BAPTA-AM (Sigma), Xestospongin C (Tocris/R&D), or

lithium chloride (LiCl, Sigma) were added as noted for some experiments. For the experiment using TTX

(select agent toxin), all appropriate federal regulations and protocols established for the Select Agent

Program established by the Centers for Disease Control and Prevention (CDC) and the US Department

of Agriculture (USDA) were followed.

Electrophysiological data were collected using Axopatch 200B or Axoclamp 900A amplifiers

(Molecular Devices, Sunnyvale, CA). Sharp electrode (> 10 MΩ) recordings were taken from muscle 6 of

abdominal segments 2 and 3, as described previously (29, 30, 103). Prior to muscle V_m measurements,

the Axoclamp 900A was bridge balanced. For the Axopatch 200B, the amplifier was placed in bridge

mode (using I-CLAMP FAST for sharp electrode recordings). Before recording from each muscle,

electrode resistance was measured and properly compensated by applying a step input and adjusting

series resistance. Muscles with a V_m more hyperpolarized than -60 mV and an input resistance of greater

than 5 MΩ were deemed suitable for recording (29). Data were digitized using a Digidata 1440A data

acquisition system (Molecular Devices), and recorded using the pCLAMP 10 acquisition software

(Molecular Devices). Spontaneous activity was recorded, followed by evoked activity. For presynaptic

nerve stimulation, a Master-8 pulse stimulator (A.M.P. Instruments, Jerusalem, Israel) and an ISO-Flex

isolation unit (A.M.P. Instruments) were utilized to deliver suprathreshold stimuli (1 ms unless otherwise

indicated) to the appropriate segmental nerve. For each NMJ, the average amplitude of spontaneous

miniature excitatory postsynaptic potential EPSPs (mEPSPs) was quantified by measuring approximately

100-200 individual spontaneous release events per NMJ. The average per-NMJ mEPSP amplitudes

were then averaged for each genotype. Evoked EPSP amplitude was calculated for each NMJ as the

average of 30 events (1 Hz). Quantal content (QC) was determined in two different ways. At very low

extracellular [Ca^{2+}], QC was calculated by the method of failures, as m = ln[(# trials)/(# failures)], as

described elsewhere (40). At higher extracellular [Ca^{2+}], QC was calculated by dividing EPSP/mEPSP,

as described in the text. For analyses conducted across different calcium concentrations, QC was

corrected for non-linear summation (104). For histograms displaying mEPSP amplitude frequencies, the

same number of spontaneous events was analyzed for each NMJ (per genotype or experimental

condition). This ensured that no individual NMJs were overrepresented or underrepresented in the

aggregate analyses.

Immunostaining and Image Analysis: Third instar larvae were filleted in HL3 saline. Dissected animals

were fixed for 3 minutes in Bouin's fixative (Ricca Chemical Company, Arlington, TX), washed using

standard procedures, and incubated in primary antibodies overnight at 4°C. This was followed by

additional washes and a two-hour incubation in secondary antibody at room temperature. Staining was

performed using the following primary antibodies: mouse anti-GluRIIA (8B4D2) at 1:250 (bouton/cluster

counting) or 1:500 (intensity analyses) (Developmental Studies Hybridoma Bank (DSHB), University of

Iowa); rabbit anti-Dlg 1:30,000 (105, 106), mouse anti-Brp (nc82) 1:250 (32) (deposited to DSHB by

Buchner, E.), rabbit anti-GFP 1:250 (Torrey Pines Biolabs Inc. TP401). The following fluorophore-

conjugated antibodies were also used (Jackson ImmunoResearch Laboratories): goat anti-mouse-488

1:1000 (DyLight); and goat anti-rabbit-549 1:2000 (DyLight). Larval preparations were mounted in

Vectashield (Vector Laboratories) and imaged at room temperature using Zen software on a Zeiss 700

LSM mounted on an Axio Observer.Z1. An EC Plan-Neofluar 40X Oil DIC Objective (aperture 1.30) or an

EC Plan-Apochromat 63x Oil DIC Objective (aperture 1.40) (Carl Zeiss Microscopy) was used.

For analysis of fluorescence intensity and area, experimental and control larval preparations were

stained in the same container, mounted on the same slide, imaged using identical acquisition settings,

and analyzed using the same procedure and thresholds. Bouton and glutamate receptor cluster numbers

were quantified semi-automatically using the 'Spots' function in Imaris x64 v7.6.0 (Bitplane, Zurich

Switzerland). Any errors in automated counting were corrected by hand to arrive at the final value.

GluRIIA and Dlg levels were assessed using ImageJ 1.48s/Java 1.6.0_24 (64-bit) with Fiji plugins. Z-

stack images were compressed using the maximum projection function; ROIs were hand drawn to

exclude non-synaptic structures; a minimum threshold was set for each channel to eliminate background

fluorescence; and the Measure function was used to assess fluorescence intensity and area.

Western Blotting: 10 adult fly heads/sample were prepared in sample buffer using standard methods.

SDS-PAGE was performed using the Novex NuPAGE SDS-PAGE system with 4%-12% Bis-Tris gels run

at 125 V for 10 minutes and 150 V for 2.5 hours. Transfer to PVDF membrane (Bio-Rad, Hercules, CA) was performed using a Trans-Blot-SDSemi-Dry Transfer Cell (Bio-Rad, Hercules, CA). Blocking was performed in 5% BSA for GFP blots or 5% milk for actin blots in 1X PBS with 0.1% Tween 20. Primary antibodies were obtained from the DSHB, mouse anti-actin (JLA20) 1:1000, or from Torrey Pines Biolabs, rabbit anti-GFP 1:2000. Horseradish peroxidase-conjugated goat anti-mouse secondary antibody (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) was used at 1:5000 for actin blots. Horseradish peroxidase-conjugated goat anti-rabbit secondary antibody (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) was used at 1:5000 for GFP blots. All antibodies were diluted in blocking buffer. Blots were developed with Super-Signal West Pico Chemiluminescent Substrate (Thermo Scientific, Waltham, MA) and imagedwith Amersham Hyperfilm ECL film (GE Healthcare Limited, Buckinghamshire, UK). Band intensity was quantified using ImageJ.

Statistical Analyses and Data Plots: Most electrophysiological comparisons were made across multiple data sets. As appropriate, statistical significance was either assessed by one-way ANOVA with Tukey's post-hoc analysis for multiple comparisons (assumes Gaussian distribution), or a non-parametric Kruskal-Wallis ANOVA with Dunn's post-hoc analysis for multiple comparisons (does not assume Gaussian distribution). Other statistical tests utilized included Fisher's exact tests for viability counts and for counts of gigantic mEPSP events; Log-rank tests for survivability curves; linear regression analyses for calcium cooperativity; and Student's T-Tests for direct comparisons between one control group and one experimental group. p values of * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and **** $p < 0.0001$ were considered significant. The values reported or plotted on regular bar graphs are mean ± SEM. The values reported and plotted on box-and-whisker graphs are: box (25^{th} – 75^{th} percentiles), whiskers (1^{st} – 99^{th} percentiles), line (median), + symbol (average), and individual raw data points plotted outside the 1^{st} and 99^{th} percentiles. Statistical analyses were performed in GraphPad Prism (GraphPad Software).

746 **ACKNOWLEDGEMENTS:** We thank Martin Müller, Chun-Fang Wu and members of the Frank lab for

747 helpful comments on earlier versions of this manuscript, and the laboratories of Tina Tootle and Fang Lin

748 for helpful discussions. We thank the Bloomington Drosophila Stock Center and the Vienna Drosophila

749 Resource Center for several fly stocks detailed in the Materials and Methods section.

REFERENCES

1.   Kullmann DM. Neurological channelopathies. Annu Rev Neurosci. 2010;33:151-72.

2.   Russell JF, Fu YH, Ptacek LJ. Episodic neurologic disorders: syndromes, genes, and mechanisms. Annu Rev Neurosci. 2013;36:25-50.

3.   Ryan DP, Ptacek LJ. Episodic neurological channelopathies. Neuron. 2010;68(2):282-92.

4.   Ophoff RA, Terwindt GM, Vergouwe MN, van Eijk R, Oefner PJ, Hoffman SM, et al. Familial hemiplegic migraine and episodic ataxia type-2 are caused by mutations in the Ca2+ channel gene CACNL1A4. Cell. 1996;87(3):543-52.

5.   Kors EE, Terwindt GM, Vermeulen FL, Fitzsimons RB, Jardine PE, Heywood P, et al. Delayed cerebral edema and fatal coma after minor head trauma: role of the CACNA1A calcium channel subunit gene and relationship with familial hemiplegic migraine. Ann Neurol. 2001;49(6):753-60.

6.   Pietrobon D. Calcium channels and migraine. Biochimica et biophysica acta. 2013;1828(7):1655-65.

7.   Eikermann-Haerter K, Dilekoz E, Kudo C, Savitz SI, Waeber C, Baum MJ, et al. Genetic and hormonal factors modulate spreading depression and transient hemiparesis in mouse models of familial hemiplegic migraine type 1. The Journal of clinical investigation. 2009;119(1):99-109.

8.   van den Maagdenberg AM, Pietrobon D, Pizzorusso T, Kaja S, Broos LA, Cesetti T, et al. A Cacna1a knockin migraine mouse model with increased susceptibility to cortical spreading depression. Neuron. 2004;41(5):701-10.

9.   van den Maagdenberg AM, Pizzorusso T, Kaja S, Terpolilli N, Shapovalova M, Hoebeek FE, et al. High cortical spreading depression susceptibility and migraine-associated symptoms in Ca(v)2.1 S218L mice. Ann Neurol. 2010;67(1):85-98.

10.   Kaja S, van de Ven RC, Broos LA, Veldman H, van Dijk JG, Verschuuren JJ, et al. Gene dosage-dependent transmitter release changes at neuromuscular synapses of CACNA1A R192Q knockin mice are non-progressive and do not lead to morphological changes or muscle weakness. Neuroscience. 2005;135(1):81-95.

11.   Kaja S, Van de Ven RC, Broos LA, Frants RR, Ferrari MD, Van den Maagdenberg AM, et al. Severe and progressive neurotransmitter release aberrations in familial hemiplegic migraine type 1 Cacna1a S218L knock-in mice. Journal of neurophysiology. 2010;104(3):1445-55.

12.   Di Guilmi MN, Wang T, Inchauspe CG, Forsythe ID, Ferrari MD, van den Maagdenberg AM, et al. Synaptic gain-of-function effects of mutant Cav2.1 channels in a mouse model of familial hemiplegic migraine are due to increased basal [Ca2+]i. J Neurosci. 2014;34(21):7047-58.

13.   Inchauspe CG, Urbano FJ, Di Guilmi MN, Ferrari MD, van den Maagdenberg AM, Forsythe ID, et al. Presynaptic CaV2.1 calcium channels carrying familial hemiplegic migraine mutation R192Q allow faster recovery from synaptic depression in mouse calyx of Held. Journal of neurophysiology. 2012;108(11):2967-76.

14.   Inchauspe CG, Urbano FJ, Di Guilmi MN, Forsythe ID, Ferrari MD, van den Maagdenberg AM, et al. Gain of function in FHM-1 Cav2.1 knock-in mice is related to the shape of the action potential. Journal of neurophysiology. 2010;104(1):291-9.

15.   Hullugundi SK, Ansuini A, Ferrari MD, van den Maagdenberg AM, Nistri A. A hyperexcitability
phenotype in mouse trigeminal sensory neurons expressing the R192Q Cacna1a missense mutation of
familial hemiplegic migraine type-1. Neuroscience. 2014;266:244-54.

16.   Park J, Moon H, Akerman S, Holland PR, Lasalandra MP, Andreou AP, et al. Differential
trigeminovascular nociceptive responses in the thalamus in the familial hemiplegic migraine 1 knock-in
mouse: a Fos protein study. Neurobiology of disease. 2014;64:1-7.

17.   Pietrobon D, Moskowitz MA. Pathophysiology of migraine. Annu Rev Physiol. 2013;75:365-91.

18.   Vecchia D, Tottene A, van den Maagdenberg AM, Pietrobon D. Abnormal cortical synaptic
transmission in CaV2.1 knockin mice with the S218L missense mutation which causes a severe familial
hemiplegic migraine syndrome in humans. Frontiers in cellular neuroscience. 2015;9:8.

19.   Vecchia D, Tottene A, van den Maagdenberg AM, Pietrobon D. Mechanism underlying unaltered
cortical inhibitory synaptic transmission in contrast with enhanced excitatory transmission in CaV2.1
knockin migraine mice. Neurobiology of disease. 2014;69:225-34.

20.   Inagaki A, Frank CA, Usachev YM, Benveniste M, Lee A. Pharmacological Correction of Gating
Defects in the Voltage-Gated Cav2.1 Ca(2+) Channel due to a Familial Hemiplegic Migraine Mutation.
Neuron. 2014;81(1):91-102.

21.   Eikermann-Haerter K, Arbel-Ornath M, Yalcin N, Yu ES, Kuchibhotla KV, Yuzawa I, et al.
Abnormal synaptic Ca(2+) homeostasis and morphology in cortical neurons of familial hemiplegic
migraine type 1 mutant mice. Ann Neurol. 2015;78(2):193-210.

22.   Kawasaki F, Zou B, Xu X, Ordway RW. Active zone localization of presynaptic calcium channels
encoded by the cacophony locus of Drosophila. J Neurosci. 2004;24(1):282-5.

23.   Lin DM, Goodman CS. Ectopic and increased expression of Fasciclin II alters motoneuron growth
cone guidance. Neuron. 1994;13(3):507-23.

24.   Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates and
generating dominant phenotypes. Development. 1993;118(2):401-15.

25.   Breen TR, Lucchesi JC. Analysis of the dosage compensation of a specific transcript in
Drosophila melanogaster. Genetics. 1986;112(3):483-91.

26.   Cline TW, Meyer BJ. Vive la difference: males vs females in flies vs worms. Annual review of
genetics. 1996;30:637-702.

27.   Lucchesi JC, Kuroda MI. Dosage compensation in Drosophila. Cold Spring Harbor perspectives
in biology. 2015;7(5).

28.   Fouquet W, Owald D, Wichmann C, Mertel S, Depner H, Dyba M, et al. Maturation of active zone
assembly by Drosophila Bruchpilot. J Cell Biol. 2009;186(1):129-45.

29.   Frank CA, Kennedy MJ, Goold CP, Marek KW, Davis GW. Mechanisms underlying the rapid
induction and sustained expression of synaptic homeostasis. Neuron. 2006;52(4):663-77.

30.   Frank CA, Pielage J, Davis GW. A presynaptic homeostatic signaling system composed of the
Eph receptor, ephexin, Cdc42, and CaV2.1 calcium channels. Neuron. 2009;61(4):556-69.

31.   Gaviño MA, Ford KJ, Archila S, Davis GW. Homeostatic synaptic depression is achieved through
a regulated decrease in presynaptic calcium channel abundance. eLife. 2015;4.

32.   Wagh DA, Rasse TM, Asan E, Hofbauer A, Schwenkert I, Durrbeck H, et al. Bruchpilot, a protein
with homology to ELKS/CAST, is required for structural integrity and function of synaptic active zones in
Drosophila. Neuron. 2006;49(6):833-44.

33.   DiAntonio A, Petersen SA, Heckmann M, Goodman CS. Glutamate receptor expression regulates
quantal size and quantal content at the Drosophila neuromuscular junction. J Neurosci. 1999;19(8):3023-
32.

34.   Feng Y, Ueda A, Wu CF. A modified minimal hemolymph-like solution, HL3.1, for physiological
recordings at the neuromuscular junctions of normal and mutant Drosophila larvae. Journal of
neurogenetics. 2004;18(2):377-402.

35.   Tottene A, Conti R, Fabbro A, Vecchia D, Shapovalova M, Santello M, et al. Enhanced excitatory
transmission at cortical synapses as the basis for facilitated spreading depression in Ca(v)2.1 knockin
migraine mice. Neuron. 2009;61(5):762-73.

36.   Adams PJ, Rungta RL, Garcia E, van den Maagdenberg AM, MacVicar BA, Snutch TP.
Contribution of calcium-dependent facilitation to synaptic plasticity revealed by migraine mutations in the
P/Q-type calcium channel. Proc Natl Acad Sci U S A. 2010;107(43):18694-9.

37.   McGuire SE, Le PT, Osborn AJ, Matsumoto K, Davis RL. Spatiotemporal rescue of memory
dysfunction in Drosophila. Science. 2003;302(5651):1765-8.

38.   Jan LY, Jan YN. Properties of the larval neuromuscular junction in Drosophila melanogaster. J
Physiol. 1976;262(1):189-214.

39.   Martin AR. Quantal Nature of Synaptic Transmission. Physiol Rev. 1966;46:51-66.

40.   Del Castillo J, Katz B. Quantal components of the end-plate potential. J Physiol. 1954;124(3):560-
73.

41.   Frank CA. Homeostatic plasticity at the Drosophila neuromuscular junction. Neuropharmacology.
2014;78:63-74.

42.   Frank CA. How voltage-gated calcium channels gate forms of homeostatic synaptic plasticity.
Frontiers in cellular neuroscience. 2014;8:40.

43.   Brusich DJ, Spring AM, Frank CA. A single-cross, RNA interference-based genetic tool for
examining the long-term maintenance of homeostatic plasticity. Frontiers in cellular neuroscience.
2015;9:107.

44.   Shortridge RD, Yoon J, Lending CR, Bloomquist BT, Perdew MH, Pak WL. A Drosophila
phospholipase C gene that is expressed in the central nervous system. J Biol Chem.
1991;266(19):12474-80.

45.   Tedford HW, Zamponi GW. Direct G protein modulation of Cav2 calcium channels. Pharmacol
Rev. 2006;58(4):837-62.

46.   Cremona O, De Camilli P. Phosphoinositides in membrane traffic at the synapse. Journal of cell
science. 2001;114(Pt 6):1041-52.

47.   Goni FM, Alonso A. Structure and functional properties of diacylglycerols in membranes.
Progress in lipid research. 1999;38(1):1-48.

48.   Huang FD, Woodruff E, Mohrmann R, Broadie K. Rolling blackout is required for synaptic vesicle
exocytosis. J Neurosci. 2006;26(9):2369-79.

49.   Peters C, Bayer MJ, Buhler S, Andersen JS, Mann M, Mayer A. Trans-complex formation by
proteolipid channels in the terminal phase of membrane fusion. Nature. 2001;409(6820):581-8.

50.   Rohrbough J, Broadie K. Lipid regulation of the synaptic vesicle cycle. Nat Rev Neurosci.
2005;6(2):139-50.

51.   Wu L, Bauer CS, Zhen XG, Xie C, Yang J. Dual regulation of voltage-gated calcium channels by
PtdIns(4,5)P2. Nature. 2002;419(6910):947-52.

52.   Dahdal D, Reeves DC, Ruben M, Akabas MH, Blau J. Drosophila pacemaker neurons require g
protein signaling and GABAergic inputs to generate twenty-four hour behavioral rhythms. Neuron.
2010;68(5):964-77.

53.   Llano I, Gonzalez J, Caputo C, Lai FA, Blayney LM, Tan YP, et al. Presynaptic calcium stores
underlie large-amplitude miniature IPSCs and spontaneous calcium transients. Nat Neurosci.
2000;3(12):1256-65.

54.   Liu Q, Chen B, Yankova M, Morest DK, Maryon E, Hand AR, et al. Presynaptic ryanodine
receptors are required for normal quantal size at the Caenorhabditis elegans neuromuscular junction. J
Neurosci. 2005;25(29):6745-54.

55.   Collin T, Marty A, Llano I. Presynaptic calcium stores and synaptic transmission. Curr Opin
Neurobiol. 2005;15(3):275-81.

56.   Emptage NJ, Reid CA, Fine A. Calcium stores in hippocampal synaptic boutons mediate short-
term plasticity, store-operated Ca2+ entry, and spontaneous transmitter release. Neuron.
2001;29(1):197-208.

57.   Simkus CR, Stricker C. The contribution of intracellular calcium stores to mEPSCs recorded in
layer II neurones of rat barrel cortex. J Physiol. 2002;545(Pt 2):521-35.

58.   Wong CO, Chen K, Lin YQ, Chao Y, Duraine L, Lu Z, et al. A TRPV channel in Drosophila motor
neurons regulates presynaptic resting Ca2+ levels, synapse growth, and synaptic transmission. Neuron.
2014;84(4):764-77.

59.   Joshi R, Venkatesh K, Srinivas R, Nair S, Hasan G. Genetic dissection of itpr gene function
reveals a vital requirement in aminergic cells of Drosophila larvae. Genetics. 2004;166(1):225-36.

60.   Conti R, Tan YP, Llano I. Action potential-evoked and ryanodine-sensitive spontaneous Ca2+
transients at the presynaptic terminal of a developing CNS inhibitory synapse. J Neurosci.
2004;24(31):6946-57.

61.   Dunn TW, Syed NI. Ryanodine receptor-transmitter release site coupling increases quantal size
in a synapse-specific manner. The European journal of neuroscience. 2006;24(6):1591-605.

62.   Gordon GR, Bains JS. Noradrenaline triggers multivesicular release at glutamatergic synapses in
the hypothalamus. J Neurosci. 2005;25(49):11385-95.

63.   Sharma G, Vijayaraghavan S. Modulation of presynaptic store calcium induces release of
glutamate and postsynaptic firing. Neuron. 2003;38(6):929-39.

64.   Dunn TW, McCamphill PK, Syed NI. Activity-induced large amplitude postsynaptic mPSPs at
soma-soma synapses between Lymnaea neurons. Synapse. 2009;63(2):117-25.

65.   Dockendorff TC, Robertson SE, Faulkner DL, Jongens TA. Genetic characterization of the 44D-
45B region of the Drosophila melanogaster genome based on an F2 lethal screen. Molecular & general
genetics : MGG. 2000;263(1):137-43.

66.   Hallcher LM, Sherman WR. The effects of lithium ion and other agents on the activity of myo-
inositol-1-phosphatase from bovine brain. J Biol Chem. 1980;255(22):10896-901.

67.   Lenox RH, Wang L. Molecular basis of lithium action: integration of lithium-responsive signaling
and gene expression networks. Mol Psychiatry. 2003;8(2):135-44.

68.   Gafni J, Munsch JA, Lam TH, Catlin MC, Costa LG, Molinski TF, et al. Xestospongins: potent
membrane permeable blockers of the inositol 1,4,5-trisphosphate receptor. Neuron. 1997;19(3):723-33.

69.   Wilcox RA, Primrose WU, Nahorski SR, Challiss RA. New developments in the molecular
pharmacology of the myo-inositol 1,4,5-trisphosphate receptor. Trends Pharmacol Sci. 1998;19(11):467-
75.

70.   Goswami SP, Bucurenciu I, Jonas P. Miniature IPSCs in hippocampal granule cells are triggered
by voltage-gated Ca2+ channels via microdomain coupling. J Neurosci. 2012;32(41):14294-304.

71.   Ermolyuk YS, Alder FG, Surges R, Pavlov IY, Timofeeva Y, Kullmann DM, et al. Differential
triggering of spontaneous glutamate release by P/Q-, N- and R-type Ca2+ channels. Nat Neurosci.
2013;16(12):1754-63.

72.   Xu-Friedman MA, Regehr WG. Maximinis. Nat Neurosci. 2000;3(12):1229-30.

73.   Melom JE, Akbergenova Y, Gavornik JP, Littleton JT. Spontaneous and evoked release are
independently regulated at individual active zones. J Neurosci. 2013;33(44):17253-63.

74.   Yao WD, Wu CF. Auxiliary Hyperkinetic beta subunit of K+ channels: regulation of firing
properties and K+ currents in Drosophila neurons. J Neurophysiol. 1999;81(5):2472-84.

75.   Berke BA, Lee J, Peng IF, Wu CF. Sub-cellular Ca2+ dynamics affected by voltage- and Ca2+-
gated K+ channels: Regulation of the soma-growth cone disparity and the quiescent state in Drosophila
neurons. Neuroscience. 2006;142(3):629-44.

76.   Gao Z, Todorov B, Barrett CF, van Dorp S, Ferrari MD, van den Maagdenberg AM, et al.
Cerebellar ataxia by enhanced Ca(V)2.1 currents is alleviated by Ca2+-dependent K+-channel activators
in Cacna1a(S218L) mutant mice. J Neurosci. 2012;32(44):15533-46.

77.   Vanmolkot KR, Kors EE, Turk U, Turkdogan D, Keyser A, Broos LA, et al. Two de novo mutations
in the Na,K-ATPase gene ATP1A2 associated with pure familial hemiplegic migraine. European journal
of human genetics : EJHG. 2006;14(5):555-60.

78.   Bergquist S, Dickman DK, Davis GW. A hierarchy of cell intrinsic and target-derived homeostatic
signaling. Neuron. 2010;66(2):220-34.

79.   Lee J, Ueda A, Wu CF. Pre- and post-synaptic mechanisms of synaptic strength homeostasis
revealed by slowpoke and shaker K+ channel mutations in Drosophila. Neuroscience. 2008;154(4):1283-
96.

80.   Lee J, Ueda A, Wu CF. Distinct roles of Drosophila cacophony and Dmca1D Ca(2+) channels in
synaptic homeostasis: genetic interactions with slowpoke Ca(2+) -activated BK channels in presynaptic
excitability and postsynaptic response. Developmental neurobiology. 2014;74(1):1-15.

81.   Peng IF, Wu CF. Drosophila cacophony channels: a major mediator of neuronal Ca2+ currents
and a trigger for K+ channel homeostatic regulation. J Neurosci. 2007;27(5):1072-81.

82.   Parrish JZ, Kim CC, Tang L, Bergquist S, Wang T, Derisi JL, et al. Kruppel mediates the selective
rebalancing of ion channel expression. Neuron. 2014;82(3):537-44.

83.   Li P, Chen SR. Molecular basis of Ca(2)+ activation of the mouse cardiac Ca(2)+ release channel
(ryanodine receptor). The Journal of general physiology. 2001;118(1):33-44.

84.   Rahman T. Dynamic clustering of IP3 receptors by IP3. Biochemical Society transactions.
2012;40(2):325-30.

85.   Zahradnik I, Gyorke S, Zahradnikova A. Calcium activation of ryanodine receptor channels--
reconciling RyR gating models with tetrameric channel structure. The Journal of general physiology.
2005;126(5):515-27.

86.   Laver DR. Ca2+ stores regulate ryanodine receptor Ca2+ release channels via luminal and
cytosolic Ca2+ sites. Clinical and experimental pharmacology & physiology. 2007;34(9):889-96.

87.   de Vries B, Eising E, Broos LA, Koelewijn SC, Todorov B, Frants RR, et al. RNA expression
profiling in brains of familial hemiplegic migraine type 1 knock-in mice. Cephalalgia : an international
journal of headache. 2014;34(3):174-82.

88.   Drissi H, Lasmoles F, Le Mellay V, Marie PJ, Lieberherr M. Activation of phospholipase C-beta1
via Galphaq/11 during calcium mobilization by calcitonin gene-related peptide. J Biol Chem.
1998;273(32):20168-74.

89.   Morara S, Wang LP, Filippov V, Dickerson IM, Grohovaz F, Provini L, et al. Calcitonin gene-
related peptide (CGRP) triggers Ca2+ responses in cultured astrocytes and in Bergmann glial cells from
cerebellar slices. Eur J Neurosci. 2008;28(11):2213-20.

90.   Drissi H, Lieberherr M, Hott M, Marie PJ, Lasmoles F. Calcitonin gene-related peptide (CGRP)
increases intracellular free Ca2+ concentrations but not cyclic AMP formation in CGRP receptor-positive
osteosarcoma cells (OHS-4). Cytokine. 1999;11(3):200-7.

91.   Goadsby PJ, Edvinsson L, Ekman R. Release of vasoactive peptides in the extracerebral
circulation of humans and the cat during activation of the trigeminovascular system. Ann Neurol.
1988;23(2):193-6.

92.   Goadsby PJ, Edvinsson L, Ekman R. Vasoactive peptide release in the extracerebral circulation
of humans during migraine headache. Ann Neurol. 1990;28(2):183-7.

93.   Kaiser EA, Russo AF. CGRP and migraine: could PACAP play a role too? Neuropeptides.
2013;47(6):451-61.

94.   Hazelrigg T, Levis R, Rubin GM. Transformation of white locus DNA in drosophila: dosage
compensation, zeste interaction, and position effects. Cell. 1984;36(2):469-81.

95.   Weinkove D, Neufeld TP, Twardzik T, Waterfield MD, Leevers SJ. Regulation of imaginal disc cell
size, cell number and organ size by Drosophila class I(A) phosphoinositide 3-kinase and its adaptor. Curr
Biol. 1999;9(18):1019-29.

96.   Venken KJ, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, et al. MiMIC: a highly
versatile transposon insertion resource for engineering Drosophila melanogaster genes. Nature methods.
2011;8(9):737-43.

97.   Deshpande M, Venkatesh K, Rodrigues V, Hasan G. The inositol 1,4,5-trisphosphate receptor is
required for maintenance of olfactory adaptation in Drosophila antennae. J Neurobiol. 2000;43(3):282-8.

98.   Spradling AC, Stern D, Beaton A, Rhem EJ, Laverty T, Mozden N, et al. The Berkeley Drosophila
Genome Project gene disruption project: Single P-element insertions mutating 25% of vital Drosophila
genes. Genetics. 1999;153(1):135-77.

99.   Sullivan KM, Scott K, Zuker CS, Rubin GM. The ryanodine receptor is essential for larval
development in Drosophila melanogaster. Proc Natl Acad Sci U S A. 2000;97(11):5942-7.

100.   Yao CA, Carlson JR. Role of G-proteins in odor-sensing and CO2-sensing neurons in Drosophila.
J Neurosci. 2010;30(13):4562-72.

101.   Banerjee S, Joshi R, Venkiteswaran G, Agrawal N, Srikanth S, Alam F, et al. Compensation of
inositol 1,4,5-trisphosphate receptor function by altering sarco-endoplasmic reticulum calcium ATPase
activity in the Drosophila flight circuit. J Neurosci. 2006;26(32):8278-88.

102.   Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, et al. A genome-wide transgenic
RNAi library for conditional gene inactivation in Drosophila. Nature. 2007;448(7150):151-6.

103.   Davis GW, DiAntonio A, Petersen SA, Goodman CS. Postsynaptic PKA controls quantal size and
reveals a retrograde signal that regulates presynaptic transmitter release in Drosophila. Neuron.
1998;20(2):305-15.

104.   Martin AR. A further study of the statistical composition on the end-plate potential. J Physiol.
1955;130(1):114-22.

105.   Budnik V, Koh YH, Guan B, Hartmann B, Hough C, Woods D, et al. Regulation of synapse
structure and function by the Drosophila tumor suppressor gene dlg. Neuron. 1996;17(4):627-40.

106.   Pielage J, Cheng L, Fetter RD, Carlton PM, Sedat JW, Davis GW. A presynaptic giant ankyrin
stabilizes the NMJ through regulation of presynaptic microtubules and transsynaptic cell adhesion.
Neuron. 2008;58(2):195-209.

1010

FIGURE LEGENDS

Figure 1: SL- and DM-expressing flies exhibit coarse phenotypes. (A) Schematic of Ca$_V$2-type calcium channel α1a subunit, with substitutions to Drosophila Cacophony (Cac) residues indicated (mammalian residues in parentheses) and a CLUSTAL-Omega alignment of Cac, human CACNA1A, and mouse CACNA1A amino acids spanning the relevant region ([*] - fully conserved; [:] - strongly similar; [.] - weakly similar). **(B, C)** Visible phenotypes for larvae resulting from crosses of *elaV(C155)-Gal4* females x *Balancer/UAS-cac-eGFP*$^{MUT \ or \ WT}$ males. **(B)** Premature spiracle protrusion in larvae expressing the *UAS-cac-eGFP*^{SL} (shown) or *UAS-cac-eGFP*^{DM} (not shown) transgenic lines. The spiracle phenotype did not occur in larvae expressing *UAS-cac-eGFP*^{RQ} or *UAS-cac-eGFP*^{WT}. **(C)** Same crosses as in (B) showing diminished *UAS-cac-eGFP* mutant viability. "*UAS-cac* Viability Index" = # *UAS-cac-eGFP* transgenic adult progeny/# Balancer Chromosome siblings, normalized to 100% for WT female progeny counts (Table 1 for raw counts; for all comparisons, $n \geq 115$ Balancer sibling progeny were counted). *** $p < 0.001$ by Fisher's Exact test compared to WT sex-specific control. # $p = 0.05$, ### $p < 0.001$ by Fisher's Exact test between sexes for the SL or DM genotypes. **(D, E)** For both females (D) and males (E), there was starkly diminished longevity for adult flies expressing the DM transgene. **** $p < 0.0001$ by Log-rank test.

Figure 2: Localization and expression levels of Cac-GFP transgenes are normal. (A-D) Images of larval central nervous systems from animals expressing Cac-GFP protein (WT or mutant). Anti-GFP (red), and anti-Bruchpilot (Brp - green) staining are shown. Scale bar 100 μm. **(E-H)** Wild-type and mutant Cac-GFP successfully localized to NMJ active zones, as indicated by co-staining with anti-Brp (green) and anti-GFP (red). Scale bar 5 μm. **(I)** Western blots of fruit fly head lysates (10 heads/lane, single sex per Western), from flies expressing either *elaV-Gal4* alone or the indicated *UAS-cac-eGFP* transgene driven by *elaV-Gal4*. Blots were probed with anti-GFP (top) and anti-Actin (bottom) antibodies. The band corresponding to Cac-GFP is indicated. Other bands are non-specific. **(J)** Compared to WT,

1037 there was no statistically significant change in Cac-GFP expression for any of the transgenic lines utilized

1038 in this study (band normalized to actin; $p > 0.65$, one-way ANOVA with Dunnett's multiple comparisons

1039 vs. WT; GAL4 alone control excluded from analysis). **(K)** Actin levels were also steady across all

1040 transgenic lines ($p > 0.71$, one-way ANOVA with Dunnett's multiple comparisons vs. WT).

1041

Figure 3: Hallmarks of NMJ development are generally normal when Cac-GFP transgenes are

expressed

(A-D) NMJ images of the synapses on Muscle 6/7 of WT- and DM-expressing third-instar larvae,

immunostained with anti-Discs Large (Dlg) and anti-GluRIIA antibodies. Scale bars, 25 μm. **(E)** For DM-

expressing NMJs, average synaptic bouton numbers were normal, except for a slight undergrowth

detected for synapse A2 muscle 6/7 (* $p < 0.05$, Student's T-test vs. WT, $n \geq 8$ NMJs for all genotypes

and segments). **(F)** The number of glutamate receptor clusters per synapse at DM-expressing NMJs was

not statistically significantly different than WT-expressing NMJs ($p > 0.1$, Student's T-test, $n \geq 8$ NMJs for

all genotypes and segments). **(G)** For DM-expressing NMJs, there was a small increase in GluRIIA-

containing receptor area coverage. **(H)** For DM-expressing NMJs there was a small increase in anti-

GluRIIA staining intensity normalized for anti-Dlg intensity (* $p < 0.05$ by Student's T-test vs. WT for both

measures, $n \geq 15$ NMJs for each genotype).

1054

Figure 4: SL- and DM-expressing NMJs display hyperexcitability in evoked neurotransmission.

(A) Average EPSP amplitudes at 0.4 mM $[Ca^{2+}]_e$ for non-transgenic control (w^{1118}) or Cac-GFP-

expressing lines (** $p < 0.01$ by one-way ANOVA with Tukey's post-hoc vs. w^{1118}; or # $p < 0.05$ and ### p

< 0.001 vs. WT; $n \geq 12$ for all genotypes). **(B)** Average quantal content (QC, estimated as

EPSP/mEPSP) at 0.4 mM $[Ca^{2+}]_e$ ($p > 0.15$ by one-way ANOVA with Tukey's post-hoc for all genotypes,

compared to both w^{1118} and WT controls). **(C)** Log-log plots of extracellular calcium concentration vs. QC

corrected for non-linear summation (NLS QC). There are no statistically significant differences in calcium

cooperativity between genotypes (p = 0.16, linear regression analysis). **(D, E)** Example

electrophysiological traces of **(D)** normal and **(E)** abnormal EPSP waveforms. **(F)** Effect of genotype on

EPSP waveforms in response to 30 presynaptic pulses. "RQ only" signifies larvae with a null

endogenous *cac* mutation rescued to viability by the RQ-expressing transgene. **(G)** Effect of genotype on

number of extra discharges observed per 30 presynaptic pulses (* $p < 0.05$ and *** $p < 0.001$ vs. WT by

one-way Kruskal-Wallis ANOVA with Dunn's post-hoc). **(H)** Penetrance and **(I)** severity of DM-associated

extra discharge waveform dysfunction in low extracellular Mg^{2+} (6 mM). **(J)** NMJ recordings of 2 min

spontaneous neurotransmission with an intact CNS. Measurements assessed: epileptiform activity > 2

sec in duration at any point in the recording; epileptiform activity with postsynaptic events > 4 mV;

epileptiform activity with postsynaptic events > 10 mV; any observed postsynaptic event (epileptiform or

not) > 10 mV; any recording that was entirely epileptiform throughout (n = 9 for WT, n = 10 for DM; * $p <$

0.05, ** $p < 0.01$ by Fisher's Exact Test). All genotypes abbreviated (WT, DM, SL, RQ) are *elaV(C155)-*

Gal4/Y; UAS-cac-eGFP$^{(X)}$/+ or *w^{1118}* for non-transgenic wild type. Data bars represent the average value

and error bars the SEM.

1076

Figure 5: SL- and DM-expressing NMJs have enhanced mEPSPs. (A) Electrophysiological traces of

spontaneous activity at WT- and DM-expressing NMJs. Example traces with two different scales show

variable severity of spontaneous neurotransmission phenotypes, in terms of frequency severity (left) or

amplitude severity (right). **(B)** Effects of genotype on average spontaneous mEPSP amplitude (** $p <$

0.01 vs. *w^{1118}* by one-way ANOVA with Tukey's post-hoc; ### $p < 0.001$ vs. WT by one-way ANOVA with

Tukey's post-hoc. **(C)** Effects of genotype on spontaneous mEPSP frequency. ## $p < 0.01$ vs. WT by

one-way ANOVA with Tukey's post-hoc; $n \geq 12$ NMJs, all genotypes. **(D)** Box and whisker plots of

mEPSP amplitude range at 0.5 mM extracellular [Ca^{2+}]. Box denotes 25^{th}-75^{th} percentile; line denotes

median; + sign denotes average; whiskers range from 1^{st}-99^{th} percentile; individual data points outside

the 1^{st} and 99^{th} percentiles are plotted; (*** $p < 0.001$ by Kruskal-Wallis ANOVA with Dunn's post-hoc vs.

either *w^{1118}* or WT; $n > 1400$ mEPSPs for each genotype). **(E)** Cumulative probability histogram of the

data in (D) showing a marked rightward shift in mEPSP amplitudes for SL- and DM-expressing NMJs. **(F)**

Box and whisker plot (as in (D)) of mEPSP amplitude at 0.4 mM extracellular $[Ca^{2+}]$ with and without the

$Ca_V2.1$ channel modifier BHQ (*** $p < 0.001$ by Kruskal-Wallis ANOVA with Dunn's post-hoc vs. identical

genotype +/- BHQ; $n > 985$ mEPSPs for each genotype). **(G)** Cumulative probability histogram of a

subset of data in (F). 5 μM BHQ causes a partial leftward shift in the distribution of events for DM- and

SL-expressing NMJs while not affecting WT-expressing NMJs. **(H)** Box and whisker plot (as in (D)) of

mEPSP amplitude range when expressing the DM transgene for acute periods of developmental time

(*** $p < 0.001$ by Kruskal-Wallis ANOVA with Dunn's post-hoc vs. DM 0 hr., ### $p < 0.001$ by vs. DM 24

hr.; $n > 1095$ mEPSPs for each genotype). **(I)** Cumulative probability histogram of the data in (H)

showing a rightward shift in mEPSP amplitudes for longer periods of DM expression.

1098

Figure 6: Gigantic spontaneous events vanish in response to diminished Ca^{2+}, buffered Ca^{2+}, or

blocked Na_V. (A) Box and whisker plot of mEPSP amplitudes at 0.2 mM extracellular Ca^{2+}. Plot as in

Fig. 5 (*** $p < 0.001$ by Kruskal-Wallis ANOVA with Dunn's post-hoc vs. either w^{1118} or WT; $n > 780$

mEPSPs for each genotype). **(B)** Cumulative probability histogram of the data in (A) showing a rightward

shift in mEPSP amplitudes for SL- and DM-expressing NMJs – but less so than for 0.5 mM Ca^{2+}, with

smaller and fewer gigantic events (compare to Figure 5). **(C)** Box and whisker plots demonstrating

elimination of gigantic spontaneous events by various manipulations. (*** $p < 0.001$ by Fisher's exact test

examining the incidence of gigantic mEPSPs > 10 mV vs. DM or SL alone, as appropriate). **(D-F)**

Cumulative probability histograms of mEPSP size separately showing the effects of zero extracellular

Ca^{2+} (D); application of BAPTA-AM in 0.5 mM Ca^{2+} (E); application of TTX in 0.5 mM Ca^{2+} (F). In each

case, the rightward shift in mEPSP size distribution persists due to DM expression. However, the gigantic

spontaneous events are eliminated (see frequency shift at arrowheads). **(G)** Box and whisker plot of

spontaneous event amplitudes at 0.5 mM extracellular Ca^{2+} + TTX, with an intact central nervous

system. (*** $p < 0.001$ by Mann-Whitney U Test of WT vs. DM; $n = 900$ mEPSPs for each genotype). **(H)**

Cumulative probability histogram of the data in (G).

Figure 7: Failure analysis: SL- and DM-expressing NMJs show elevated release probability at very low extracellular calcium. (A-B) Frequencies of evoked amplitudes at very low extracellular Ca^{2+} (0.14 mM) for **(A)** WT-expressing NMJs and **(B)** DM-expressing NMJs. For the DM-expressing NMJs, there is a clear rightward shift in the size distribution of DM-expressing events, as well as a marked decrease in the frequency of failures (categorized as 0 mV events). **(C)** For WT-, DM-, and SL-expressing NMJs, the average EPSP size for successfully evoked events, as well as estimated QC by failure analyses (0.14 mM Ca^{2+}) (* $p < 0.05$; ** $p < 0.01$ by one-way ANOVA with Tukey's post-hoc compared to WT). **(D)** Further lowering extracellular Ca^{2+} (0.1 mM) for DM reveals a leftward shift in size distribution and an increase in failure percentage compared to (B). **(E)** Box and whisker data are presented as in Figs. 5 and 6 – this time showing the size distributions of spontaneous mEPSP events (WT, DM, SL), as well as failure analysis (FA) evoked events for the same genotypes (failures excluded). **(F, G)** Box and whisker plots for mEPSP rise times (0.5 mM Ca^{2+}, see Fig. 5D) show a significant increase only for DM-expressing NMJs (F), as well as a dramatic slowdown for events > 2 mV in size, regardless of genotype (G).

Figure 8: Inhibition of an intracellular Ca^{2+} release pathway dampens gain-of-function phenotypes associated with FHM1-mimicking mutations. (A) Schematic of an RNA interference (RNAi)-based approach to identify suppressors of gain-of-function electrophysiological phenotypes. **(B)** Knockdown of *Plc21C* gene function reverses the increase in spontaneous mEPSP amplitude elicited by DM expression. **(C)** Box and whisker plots (as before) and **(D, E)** cumulative probability histograms (as before) demonstrate that heterozygous, loss-of-function point mutations in genes encoding the IP_3 receptor (*Itpr*^{ug3}/+) and the Ryanodine receptor (*RyR*^{E4340K}/+) also significantly diminish the gain-of-function spontaneous mEPSP phenotypes in DM-expressing NMJs. *** $p < 0.001$ by Kruskal-Wallis ANOVA with Dunn's multiple comparisons test vs. DM alone. **(F)** The *RyR*^{E4340K}/+ condition also diminishes evoked EPSP hyperexcitability phenotypes in a DM-expressing background (# of extra discharges [ED] per muscle – see also Figure 4).

Figure 9: Pharmacological inhibition of intracellular Ca^{2+} release dampens gain-of-function

phenotypes associated with FHM1-mimicking mutations. (A-C) Data displayed and analyzed as

before. Box and whisker plots (A) and cumulative probability histograms (B, C) demonstrate that acute

application of either LiCl (to block PIP$_2$ recycling) or Xestospongin C (to block IP$_3$ receptors) both

suppress the gain-of-function spontaneous mEPSP phenotypes in DM-expressing NMJs. *** $p < 0.001$

by Kruskal-Wallis ANOVA with Dunn's multiple comparisons test vs. DM alone. **(D)** Cartoon model

depicting neuronal components implicated in this study of regulating neurophysiology downstream of

migraine-mimicking amino-acid substitutions. Red – Ca$_V$2 channels; gray – IP$_3$ receptors; blue –

Ryanodine receptors; yellow – Na$_V$ channels.

A

R135Q (R192Q)

S161L (S218L)

EF Hand

IQ Motif

	RQ		SL	
D.m. Cac	TGAMTIFAEANIDVDLR	MLRSFRVLRPLKLVSRIPSLQVVLGLLVLF	178
H.s. CACNA1A	TGIL---ATVGTEFDLR	TLRAVRVLRPLKLVSGIPSLQVVLKS	IMKAMIPLLQIGLLLFF	235
M.m. CACNA1A	TGIL---ATVGTEFDLR	TLRAVRVLRPLKLVSGIPSLQVVLKS	IMKAMIPLLQIGLLLFF	237

** *   :   *   . :.:.*** **:*:**********   *************:*** .********.:*

B

WT

SL

C

UAS-cac Viability Index (% WT Females)

Females | Males

WT | DM | SL | RQ

p = 0.08   0.15

*** ###   # *** *** *** ***

D

Females ♀

—— *elaV-Gal4/+*
—— *elaV-Gal4/+; UAS-cac^{WT}/+*
—— *elaV-Gal4/+; UAS-cac^{DM}/+*

% Survival

Days

****

E

Males ♂

—— *elaV-Gal4/Y*
—— *elaV-Gal4/Y; UAS-cac^{WT}/+*
—— *elaV-Gal4/Y; UAS-cac^{DM}/+*

% Survival

Days

****

E   Brp   WT   Cac-GFP

F   DM

G   SL

H   RQ

I

260 kDa
Cac-GFP
140 kDa

Actin

J

% WT Cac-GFP/Actin

K

% WT Actin