Genome-wide analysis of the association of transposable elements with gene regulation suggests that *Alu* elements have the largest overall regulatory impact

Lu Zeng[1, 2&], Stephen M. Pederson[2&], Danfeng Cao[1], Zhipeng Qu[2], Zhiqiang Hu[1], David L. Adelson[2*], and Chaochun Wei[1*]

[1]School of Life Sciences and Biotechnology

Shanghai Jiao Tong University

Shanghai

P. R. China

[2]School of Biological Sciences

The University of Adelaide

Adelaide, SA

Australia

RUNNING TITLE: Repetitive elements and gene regulation

[&]These authors contributed equally to this work

Author for Corresponding:

[1]Chaochun Wei, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai, China, [1]Tel: (86) (21) 34204348, Fax: (86)(21) 34204348, Email: ccwei@sjtu.edu.cn

[2]David L. Adelson, School of Biological Sciences, The University of Adelaide, Adelaide, SA, Australia, [2]Tel: +61 8 83137555, Fax: +61 8 83133262, Email: david.adelson@adelaide.edu.au

ABSTRACT

Nearly half of the human genome is made up of transposable elements (TEs) and there is evidence that TEs are involved in gene regulation. Here, we have integrated publicly available genomic, epigenetic and transcriptomic data to investigate this in a genome-wide manner. A bootstrapping statistical method was applied to minimize the confounder effects from different repeat types. Our results show that although most TE classes are primarily associated with reduced gene expression, *Alu* elements are associated with up regulated gene expression. Furthermore, *Alu* elements had the highest probability of any TE class of contributing to regulatory regions of any type defined by chromatin state. This suggests a general model where clade specific SINEs may contribute more to gene regulation than ancient/ancestral TEs. Finally, non-coding regions were found to have a high probability of TE content within regulatory sequences, most notably in repressors. Our exhaustive analysis has extended and updated our understanding of TEs in terms of their global impact on gene regulation, and suggests that the most recently derived types of TEs, *i.e.* clade or species specific SINES, have the greatest overall impact on gene regulation.

Keywords: Transposable elements/Epigenetic/Gene expression/Gene ontology/Regulatory elements.

INTRODUCTION

Repetitive elements are similar or identical DNA sequences present in multiple copies throughout the genome. The majority of the repetitive sequences in the human genome are derived from transposable elements (TEs) (Lander, et al. 2001; Smit 1999) that can move within the genome, potentially giving rise to mutations or altering genome size and structure. Typical eukaryotic genomes contain millions of copies of transposable elements (TEs) and other repetitive sequences. TEs fall into two major classes: those moving/replicating via a copy and paste mechanism and an RNA intermediate (retrotransposons) and those moving via direct cut and paste of their DNA sequences (DNA transposons). Retrotransposons can be subdivided into two groups: Those with long terminal repeats (LTRs), and those without LTRs (non-LTRs). Human LTR elements are related to endogenous retroviruses (HERVs), which along with similar elements account for nearly 8% of the human genome (Cordaux and Batzer 2009). Non-LTR retrotransposons include two sub-types: autonomous long interspersed elements (LINEs) and non-autonomous short interspersed elements (SINEs), which are dependent on autonomous elements for their replication; both LINEs and SINEs are widespread in eukaryotic genomes. LINE-1 (long interspersed element 1) and *Alu* elements are two TEs that belong to non-LTR retrotransposons, which account for approximately one-quarter of the human genome (Lander, et al. 2001).

A number of existing studies have shown that TEs can influence host genes by providing novel promoters, splice sites or post-transcriptional modification to re-wire different developmental regulatory and transcriptional networks (Cowley and Oakey 2013; Kunarso, et al. 2010; Lynch, et al. 2011). TEs tend to regulate gene expression through several mechanisms (Britten 1996; Cowley and Oakey 2013; Pereira, et al. 2009; van de Lagemaat, et al. 2003). For example, in rodents, the expression levels of protein coding genes containing repetitive elements are significantly associated with the number of repetitive elements in those genes (Pereira, et al. 2009). L1 elements show a stronger negative correlation with expression levels than the gene length (Jjingo, et al. 2011), and the presence of L1 sequences within genes can lower transcriptional activity (Han, et al. 2004). Moreover,

TEs have been shown to influence gene expression through non-coding RNAs, resulting in the

reduction or silencing of gene expression (Rebollo, et al. 2012). For example, the expression of long

intergenic non-coding RNAs (lincRNAs) was strongly correlated with HERVH transcriptional

regulatory signals (Kelley and Rinn 2012). Past studies have found that TEs have contributed to

nearly half of the active regulatory elements of the human genome (Jacques, et al. 2013), by altering

gene promoters and creating alternative promoters and enhancers to regulate gene activity (Conley, et

al. 2008; Franchini, et al. 2011; Medstrand, et al. 2001). According to previous research, 60% of TEs

in both human and mouse were located in intronic regions and all TE families in human and mouse

can exonize, supporting the view that TEs may create new genes and exons by promoting the

formation of novel or alternative transcripts (Piriyapongsa, et al. 2007; Sela, et al. 2007). The

association between repetitive elements and RNAs has also been investigated. For example, *Alu*

elements in lncRNAs can lead to *STAU1* mediated mRNA decay by duplexing with complementary

Alu elements in the 3'UTRs of mRNAs (Hadjiargyrou and Delihas 2013), and the insertion of TEs

may also drive the evolution of lincRNAs and alter their biological functions (Kelley and Rinn 2012).

In this study, TEs in the human genome were analyzed using genome-wide datasets associated with

gene regulation. These datasets enabled an assessment of the association of TEs with Regulatory

Elements (RE) based on chromatin states, as marked by histone modification within six human cell

lines, lincRNA, Gene Ontology (GO) enrichment, as well as overall transcriptome profiles.

Although, a number of studies have analyzed the association between TEs and gene regulation in the

human genome, they did not consider whether gene intervals contained multiple repeat types or not.

This can cause a confounding factor when attempting to determine the regulatory impact of a specific

repeat type. In our study, we eliminated confounding effects by focusing on gene intervals that only

contained one type of repeat and used a weighted bootstrap approach to minimize any influence of co-

occurring elements. Our method presents a novel way to investigate the association of specific TE

type with gene expression.

RESULTS

The Distribution Of Repetitive Elements In The Human Genome

Initially, we compared the distribution of repetitive elements in the human genome. We found many

repetitive elements overlapped with gene models from the human RefSeq gene datasets, and their

distributions with respect to components of gene models are shown in Figure 1. Fewer repetitive

elements were found in coding exons and non-coding exons such as the 5' and 3' UTRs. This

indicates that repeat accumulation in these regions is selected against in order to preserve regulatory

and coding functions (Figure 1).

Transposable Elements And Gene Regulation By Chromatin States.

It is known that TE-derived sequences can contribute to transcription factor binding sites, promoters

and enhancers, and insulators/silencers (de Souza, et al. 2013; Lynch, et al. 2011). To look for

enrichment of TEs within regulatory elements, we looked at the proportions of nucleotides with a TE

in six defined regulatory elements (Ernst, et al. 2011) as they appear in different components of the

gene model. This represents the probability of a given nucleotide within a Regulatory Element (RE)

being from a transposable element (TE), i.e. p(TE|RE) (see Methods for details) across the set of

genic regions (Figure 2A). One major observation from Figure 2A is that TE have higher p(TE|RE) in

insulators indicating that they are present at higher levels in insulators relative to most other

regulatory regions. While Active Promoters have the lowest repeat level in regulatory regions. The

proportion of TEs in RE is lower than the genome wide average based on the p(TE|RE) values with

exceptions being most Weak Enhancers in non-coding regions. Unsurprisingly, CDS exons have the

lowest repeat content. The overall under-representation of TE in RE is likely indicative of negative

selection for TE in regulatory regions.

Pairwise comparisons of p(TE|RE) between different gene model compartments are shown in Figure

2B in order to identify significant differences (see methods for description of statistical methods used)

in repeat content between regulatory regions/compartments. These reveal that for all regulatory

elements, TEs are more sparsely distributed across regulatory elements within CDS exons than

regulatory elements in other genic regions. Likewise, regulatory elements in the 3'UTR were more

sparsely populated with TEs in comparison to those in other regions, with the sole exception being

Active Promoters in the 5'UTR. In contrast, Polycomb Repressed Regions and Insulators in intergenic

regions were enriched for TEs in comparison to these elements in other gene model compartments.

Different Classes Of Transposable Elements And Their Associations With Chromatin State.

In order to systematically characterize the potential role of different repeat classes within the

regulatory elements mentioned above, the distribution of regulatory elements within specific TEs

classes were investigated as below, using the probability of a nucleotide also belonging to a TE

element p(TE|RE) (See Methods for details). Six TE classes (*Alu*, L1, L2, LTR, MIR and DNA

transposons) were examined (Figure 3A). The most significant observation from Figure 3A was that

Alu had the highest p(TE|RE) across all regulatory regions, especially in Weak Promoter, Strong

Enhancer and Weak Enhancer, while recent inserted L1 elements were least likely to be within

regulatory elements compared to the other five repeat classes. Although L1 elements are the most

abundant mobile elements in the human genome, accounting for 17 % of the human genome, they had

lower p (TE|RE) compared to *Alu* elements, which may indicate a negative selection for L1 element

insertions in gene regulatory regions. Interestingly, MIR and L2, ancient repeats that have been

inactive since prior to the mammalian radiation, are present at very low levels in the human genome

and they have a relatively high p(TE|RE) in Strong Enhancer regions.

In order to identify significant differences within specific repeat types between regulatory regions,

pair-wise comparisons of p(TE|RE) were implemented and shown in Figure 3B (see methods for

description of statistical methods used). Among six defined regulatory regions, *Alu* elements

contributed more to Active Promoter, Weak Promoter, Strong Enhancer and Weak Enhancer regions

compared to the other five repeat types. Ancestral L2 and MIR repeats were found to be retained

more than recent L1 across all RE. Another indication of potential exaptation of ancestral repeats was

the observation that recent inserted L1 had the lowest probability of being in a RE compared to other

repeat types (Figure 3B). However, LTR elements and DNA transposons were also more likely to be

retained than recent L1s, suggesting some degree of exaptation for those repeats as well.

TEs Are Abundantly Present In Long Intergenic Non-Coding RNAs In Regulatory Elements

Because ncRNAs are known to regulate gene expression, we decided to examine the relationships

between lincRNAs and TEs in REs (Figure 4). The observation that >30% of nucleotides from many

of the regulatory elements were derived from TEs was quite striking. Specifically, lincRNA exonic

regions contained the highest RE density for Polycomb Repressed Regions (Figure 4A), with nearly a

third of these nucleotides being derived from TEs, suggesting that the presence of transposable

elements in lincRNA exons may be associated with regulation of lincRNA expression and

subsequently linked to gene regulation. Unsurprisingly, we found that TEs made up a lower

proportion of nucleotides in CDS-exons across all regulatory elements, when compared to CDS-

introns, lincRNA exons and lincRNA introns (Figure 4B).

Identification of biases due to multiple TE types and transcript length on gene expression

We first summarized the overall distribution of transposable elements within various compartments of

gene models by finding genes containing TEs in either single or multiple compartments (Figure S1,

Table 1), and genes containing one or more type of TE (Figure S2, Table 2). Because gene/transcript

length can also cause bias in the estimation of gene expression we examined the relationship between

gene length and which compartments of a gene contained a TE (Figure S3), as well as the relationship

between gene length and the presence of a specific type of TE (Figure S4), using a Wilcoxon Test

(Tables S2 & S3). We found that only genes with TEs in the 3'UTR or in multiple genic regions

showed a bias towards longer length, whilst for TEs exclusively within the proximal promoter or

5'UTR there was a bias towards shorter genes (Figure S3; Table S2). When assessing the relationship

between gene length and the presence of a specific TE class, the length of genes with *Alu*, L2 or MIR

elements alone were very similar to genes with no TE, whilst L1 and LTR elements showed a bias

towards shorter genes, and the presence of multiple elements biased towards longer genes (Figure S4;

Table S3).

Effects On The Probability Of A Gene Being Detected As Expressed Due To The Presence Of A TE Across The Different Gene Model Compartments.

As chromatin states are not always indicative of changes in transcriptional activity, we further investigated any effects on human gene expression due to the presence of specific TE classes within each of the four regulatory regions, i.e., Proximal Promoter, 5'UTR, CDS and 3'UTR. However, as TEs are far less frequent in CDS regulatory regions (Figure S1), the subsequent analysis focused on the other three regions. Six human tissue transcriptome datasets (adipose, brain, kidney, liver, skeletal muscle and testes tissue) were selected from the Illumina BodyMap2 dataset for this analysis, and global patterns of gene expression were investigated based on the presence or absence of each TE within these three genic regions respectively.

In order to exclude confounder effects caused by multiple repeat types on the expression of single genes, the weighted bootstrap method was applied to both the probability of a gene being detected as expressed (Figure 5A), and to the overall expression levels for those genes detected as expressed (Figure 5B). This revealed that *Alu* elements are commonly associated with a higher probability of expression when located in either the 5'UTR or the 3'UTR across the majority of tissues. In contrast to the presence of an *Alu*, the presence of L1 elements in the Proximal Promoter showed a negative impact on the probability of a gene being detected as expressed in 3 out of the 6 tissues, with the remaining tissues being directionally consistent and quite likely to be Type II errors (Supplementary Figure S6A).

Effects On The Levels Of Gene Expression Due To The Presence Of A TE In Each Gene Model Compartment.

Again using the weighted bootstrap approach to minimize any confounding effects of co-occurring elements, the presence of an *Alu* in the 5'UTR was found to be associated with increased expression levels in five of the six tissues investigated (Figure 5B). Similarly, *Alu* elements in the Proximal Promoter were associated with increased expression in two of the tissues (skeletal muscle and testis).

Alu elements in 3'UTR were associated with elevated expression levels only in the Kidney sample.

The presence of L1, L2 and MIR elements showed varying degrees of reduced gene expression across

the tissues when located in the 3'UTR only. It was also noted that whilst strongly controlling the

family-wise Type-I error rate (FWER), the adjusted confidence intervals would result in an increase

in the Type-II error rate where true differences cannot be detected. As such, the point at which the

confidence intervals would include zero was found and taken as a proxy for the p-value. Confidence

intervals based on these p values to an FDR of 0.05 are shown in Supplementary Figure S6 with the p

values given in Supplementary Table S5. It is clear from this additional approach that the role of TEs

such as L1 elements in Proximal Promoters and 3'UTRs, LTR elements in 5'UTRs and many of the

elements in the 3'UTR may have been considerably understated as a result of our more conservative

approach.

Analysis Of Genes With Exapted Or Exonized TEs

In order to determine if exapted or exonized TEs might influence gene function, genic regions

(proximal promoter, 5'UTR and 3'UTR) that overlapped with TEs were analyzed to assess the

association of TEs with different gene functions (annotated using Gene Ontology).

The three fundamental GO categories are: cellular component, molecular function and biological

process. Enrichment information for each GO category is listed in Supplementary Table S6. First, we

looked at the association of gene function and repeats in the biological process category (Figure 6,

Table S6a). Figure 6 represents that 5'UTR regions containing repeats showed a strong enrichment in

genes with regulatory processes, mainly involving metabolic processes, while promoter regions

containing repeats were rarely associated with biological process. A similar pattern was found with

respect to cellular component (Figure S7, Table S6b) and molecular function (Figure S8, Table S6b),

repeats in 5'UTR were predominantly enriched in intracellular/cytoplasmic structures and associated

with binding activities respectively.

The same method was applied to investigate the association of repeats with function in protein coding

regions (CDS). We found that genes with protein coding exons containing Alus were enriched for the

GO   term   "intracellular   non-membrane-bounded   organelle".   Interestingly,   these

exonization/exaptation events were found associated with splice variants incorporating *Alu* sequences

(Table S7 & Figure S9, S10).

Moreover, according to our analysis of TEs and alternative splicing data, we found that 2.98% of

alternatively spliced transcripts contained TEs within protein coding exons (Table 3). *Alu* and MIR

were the two most common repeats to be involved in alternative splicing/exonization, but LINEs and

LTR also contributed to a significant proportion of exonization. This suggests a role for exonization

of TEs in the evolution of increasing coding and regulatory versatility of the transcriptome.

DISCUSSION

In this work, we analyzed the distribution of various classes of transposable elements and their

association with regulatory elements (active chromatin) for the first time. We have also analyzed the

association of specific repeat types with gene expression in the human genome using a new method

that minimizes confounders. Based on the analysis of the TE distributions in genic regions and

corresponding gene expression patterns, the presence of specific TEs was found to be associated with

changes in gene expression. Further, gene function as defined by GO term analysis differed

depending on the TE insertion site within the gene. Finally, we looked at TEs present in ncRNAs,

specifically lincRNAs, and found that repetitive elements were present at higher levels in lincRNAs

than in coding exons.

TEs And Gene Regulation

It is known that epigenetic modification of TEs can regulate transcription by altering active

chromatin, and this chromatin modification has evolved from epigenetic 'defense' mechanisms to

silence TE activity (Slotkin and Martienssen 2007). Here we present a comprehensive analysis of TE

contribution to each gene compartment associated with different types of active chromatin. A role for

TEs in silencing is consistent with our finding that they are enriched in Poly-comb repressed regions

and Insulators (Figure 2). However, the enrichment of TEs in Weak Enhancer and Strong Enhancer

regions (Figure 2) is suggestive of exaptation of TEs as active regulators of gene transcription. We

also found that regulatory regions overlapping with genes had a greater proportion of sequence

originating from TEs in the 5' and 3'UTRs compared to coding exons (Fig 2). This is not surprising

considering the potential adverse effect of TE insertion in a protein coding sequence (Belancio, et al.

2006; Belancio, et al. 2008).

TEs have been implicated in altering gene expression through a number of mechanisms, for example,

TE can be exonized through alternative splicing and they can alter transcription factor binding

because they contain transcription factor binding sites (Cordaux and Batzer 2009; Polak and Domany

2006). However, it is difficult to analyze the impact of specific TEs on gene expression, due to the co-

occurrence of other TE types in or near individual genes. To overcome this problem, we used a

bootstrapping approach for gene expression analysis, effectively allowing us to focus on genes

containing single TE types. Figures 5B and S6B summarize our findings with regard to gene

expression, and indicate that *Alu* elements in proximal promoter regions, 5'UTR and 3'UTR are

commonly associated with increased gene expression. Moreover, *Alu* elements were found to have a

high probability of contributing to regulatory elements, especially with respect to Weak Promoter,

Strong Enhancer and Weak Promoter regions (Figure 3). These results are consistent with previous

reports that motifs in TEs such as Alus can be exapted as transcription factor binding sites and are

involved in gene regulation (Kim, et al. 2004; Levanon, et al. 2004; Lin, et al. 2009). However, in

contrast to our finding that *Alu* elements were associated with increased gene expression, the

exonization of *Alu* sequences from alternative splicing seems to cause only decreased expression of

the alternatively spliced transcript (Lin, et al. 2009). Furthermore, the persistence of *Alu* insertions

would most likely be selected against, which would suggest that an *Alu* element can rarely take on a

regulatory role when it is inserted into a gene (Deininger 2011).

Although L1 elements are the most abundant repeats in the human genome, they were associated with

decreased gene expression (Fig 5), and were less prevalent in regulatory elements or active chromatin,

when compared to other repeat classes (Fig 3). Considering the size of L1 elements (~6kb), it is easy

to understand that the insertion of L1 elements, even if truncated, are likely to cause disruption of host

gene function (Kazazian, et al. 1988; Meischl, et al. 2000; Perepelitsa-Belancio and Deininger 2003).

Therefore, newly inserted L1 elements are probably under negative selection in regulatory regions

and are less likely be exapted as a RE.

L2 and MIR are two ancient TE families conserved in mammals, and are inactive or fossil TE

elements (Deininger and Batzer 2002). According to previous studies MIR elements have been

exapted as functional enhancers in the human genome, and are enriched in transcription factor binding

sites (Jjingo, et al. 2014). However, our results showed that MIR and L2 elements were associated

with reduced gene expression when located in 3'UTRs (Figure 5), but not with increased gene

expression. This indicates that their exaptation as enhancers has the opposite effect of their exaptation

within gene models. Figure 5 also showed that LTRs were associated with repression of gene

expression, which is in contrast to previous work that implicated LTRs as alternative promoters

(Cohen, et al. 2009). ERVL LTRs have been reported to provide molecular mechanisms for stochastic

rewiring of gene expression and evolution in the mammalian germline (Franke, et al. 2017). Our

analysis suggests that this rewiring does not affect the majority of expressed genes except for LTR

inserted into 3'UTR, which are associated with decreased gene expression.

The Association Between TEs And LincRNA

LincRNAs are known to regulate gene expression through epigenetic mechanisms (Engreitz, et al.

2013) and can be important regulators of the human innate immune response (NE, et al. 2014). One

possible factor driving lincRNA evolution and biological function is the insertion of TE elements, TE

insertions into lincRNA were confirmed significantly reduced expression in many tissues and cell

lines than lincRNAs that are devoid with TEs (Kelley and Rinn 2012). This transcriptional repression

can be supported with our results, as we found lincRNA containing TEs were clearly enriched in

Insulators and Polycomb repressed regions (Figure4). However, we also found that lincRNA

containing TEs were highly enriched in Weak Enhancer regions (Figure 4), suggesting that TEs

present in lincRNA may have an 'active' role in gene expression.

The Exaptation of TEs

In addition to potentially altering gene expression by insertion into regulatory elements, TEs may also

be associated with specific functional characteristics of expressed protein coding genes. When we

examined the functional annotation of repeat containing genes, we found that some functions were

over-represented (Table S5). Perhaps the most interesting of these associations was that genes with

Alu insertions were found to contribute to coding exons through alternative splice variants. One

explanation of this observation is that *Alu*-induced alternative transcripts may result in nonsense-

mediated decay of alternative transcripts (Lewis, et al. 2003). Two examples of alternatively spliced

genes of this type with implications for human disease are *DISC1* and *NOS3* (Table S7 and Figure S9

& S10). *DISC1* alternative transcripts are known to contribute to increased risk of schizophrenia

(Callicott, et al. 2005; Rapoport, et al. 2005) and *NOS3* transcript variants are associated with

cardiovascular disease phenotypes (Hingorani, et al. 1999; Pacanowski, et al. 2009). Based on

previous research, nearly 4% of protein-coding sequences include transposable elements, and one-

third of them are *Alu* insertions (Nekrutenko and Li 2001). Therefore, *Alu* exonization in protein-

coding genes may play an important role in modifying gene expression.

Implications For Our Perception Of Genome Evolution

While there are many publications implicating TEs in the regulation of individual genes, our work

clarifies some previous uncertainties and resolves some contradictions, confirming that this role of

TEs is significant across the genome. In general, most TEs would appear to be strongly associated

with repression of gene expression, either through the 5'UTR or perhaps as components of lincRNA

exons. However, the presence of Alus in 3'UTR and proximal promoter regions may act to increase

gene expression. These results are supported by previous studies (Jjingo, et al. 2011) and provide a

new understanding of how repeats are associated with epigenetic regulation of gene expression.

Finally, while exapted TEs may contribute to the generation of transcripts that undergo nonsense

mediated decay as part of gene regulation, we speculate that they may also provide an opportunity for

alternative splicing and novel exaptation. TEs therefore are important agents of change with respect to

the evolution of gene expression networks.

MATERIAL AND METHODS

Theoretical Framework And Methods

We constructed pipelines to analyze the distribution of repetitive elements in different parts of the

human genome. Repetitive elements overlapping with protein coding regions, non-coding regions and

regulatory elements were identified. GO term over-representation and expression analyses were

carried out for repetitive elements overlapping with protein-coding regions. The pipelines and related

materials are described below.

Tools Used To Develop Pipelines For Repetitive Element Analysis

The identification and classification of TEs from the human genome was conducted by developing a

pipeline with Perl, R (R Core Team 2014), and BEDTools (Quinlan and Hall 2010). Perl was used to

extract information from different datasets. R was used to build graphs to illustrate the repeat

distribution in different genic regions, the identification of repetitive elements with respect to

functional elements, GO term over-representation analysis and expression analysis of TEs. BED

format file intersection was used to extract the overlapping regions between different datasets, with a

lower limit of 25-bps. The UCSC Genome Browser (Dreszer, et al. 2012; Kent, et al. 2002) was used

to download genome sequence data and genome annotations including RefSeq genes. RSEM

(Trapnell, et al. 2012) was adopted to assemble RNA-Seq reads into transcripts and estimate their

abundance (measured as transcripts per million (TPM)). Plots were generated using ggplot2 in R

(Wickham 2009).

Datasets

Genome And Annotations. NCBI's Human genome and its annotation datasets (RefSeq hg19)

(Pruitt, et al. 2007) were downloaded from the UCSC Genome Browser (Dreszer, et al. 2012; Ernst, et

al. 2011). A total of 37,697 human RefSeq transcripts were merged into 18,777 genes by taking the

longest transcript(s) that represented each distinct gene locus. Repetitive elements were downloaded

from the RepeatMasker (http://www.repeatmasker.org) track of the UCSC Genome Browser. All

repetitive sequence intervals were also de-duplicated to deal with potential overlapping repeat

annotations. Overall, there were 5,298,130 human repetitive elements, which represented

approximately 1.467Gb in the human genome.

Regulatory Element Datasets From Six Human Cell Lines. The regulatory element datasets from

six human cell lines were downloaded from the UCSC Genome Browser. Each cell line dataset

contained the annotation of six regulatory elements: 1) Active Promoters, 2) Weak Promoters, 3)

Strong Enhancers, 4) Weak Enhancers, 5) Insulators and 6) Polycomb Repressed Regions. These

regulatory element annotations were derived from different chromatin states that have been marked

by histone methylation, acetylation and histone variants H2AZ, PolIII, and CTCF (Ernst, et al. 2011).

Gene Expression Datasets From Six Human Tissues. Human RNA-seq data from the Illumina

bodyMap2 transcriptome (See Table S4 for detail) (http://www.ebi.ac.uk/ena/data/view/ERP000546)

dataset was used to measure the association between TEs and the expression levels of genes

containing TEs in six tissues.

The Distribution Of Repetitive Elements In The Human Genome. To assess how human TEs were

distributed in genes, we compared different genic regions containing TEs. Based on Repbase (Jurka,

et al. 2005; Kohany, et al. 2006) annotations identified by RepeatMasker

(http://www.repeatmasker.org), repeat elements in human were divided into two categories: human-

specific repeats, and repeats shared with different species. Human-specific repeats were those

annotated with "Homo sapiens" or "primates" as their origin (See Table S8 for the list of human-

specific repeat classes), whilst those remaining were categorized as shared repeats. Intergenic regions

as well as the exons and introns within 5'UTR, CDS and 3'UTR regions from RefSeq genes (Meyer,

et al. 2013) were then compared with these different categories of repeats. Next, we generated the

summarised distributions of repetitive elements overlapping these regions by calculating the

proportions of bases belonging to repetitive elements within each of the combined sets of regions, i.e.:

$$proportion = \frac{total\ length\ of\ repeats \in defined\ regions\ (bps)}{total\ length\ of\ defined\ regions\ (bps)} \qquad (1)$$

The   code   repository   for   the   above   can   be   found   at

https://github.com/UofABioinformaticsHub/RepeatElements.

The Occurrence Of Transposable Elements Within Regulatory Regions. We further explored the

association between any TE and the regulatory elements defined above, by calculating the proportion

of nucleotides within each of the five sets of genic regions (5'UTR, CDS-exon, CDS-intron, 3'UTR

and Intergenic) that were part of a regulatory element for each of the six human cell lines. The

proportion of nucleotides that were TEs within each regulatory element were also calculated for each

genic region. All proportions were subsequently transformed using the logit function for model fitting

across tissues (Table S1) using the model

$$y_{ijk} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + \varepsilon_{ijk} \qquad (2)$$

where y_{ijk} is the logit transformed proportions representing $p(\text{TE}|\text{RE})$ across each genic region i,

each regulatory element j and tissue k, such that μ is the overall mean, α_i is the effect due to each

genic region, β_j is the effect due to each regulatory element with $(\alpha\beta)_{ij}$ representing any changes

not accounted for in the first two terms. Tests for normality and homoscedasticity were performed

using the Shapiro-Wilk test and Levene's test respectively. Where violations of homoscedasticity

were found robust standard errors were obtained using the sandwich estimator (Zeileis 2004).

Confidence Intervals for pairwise comparisons were obtained as implemented in the R package

multcomp (Westfall 2008) in order to control the Type I error at α=0.05 across the entire set of

comparisons.

A specific TE analysis was then performed using six of the major human TE classes based on the

Repbase classification system: *Alu*, L1, L2, MIR, LTR and DNA transposons. L1 elements were

further resolved into either ancestral (L1M, L1PB and L1PA subfamilies) or recent/clade-specific

(L1HS subfamily), based on their Repbase annotations. The proportions of nucleotides within each

TE type that were also regulatory elements were calculated giving tissue-specific estimates of

$p(\text{TE}|\text{RE})$. Proportions were again transformed using the logit function, and the same analysis as

above was performed.

Quality Control And Preprocessing Of The Gene Expression Data In Different Human Tissues.

RNA-seq reads of six human tissues were first assessed using FastQC software

(www.bioinformatics.babraham.ac.uk), to provide an overview of whether the raw RNA-Seq data

contained any problems or biases before further analysis. Reads with poor-quality bases were

trimmed (based on the results of FastQC with MINLEN set to 26) for subsequent data analysis. Table

4 showed the numbers of reads in raw RNA-Seq datasets and the statistics after the QC process by

using Trimmomatic-0.32 (Bolger, et al. 2014). Then, we built transcript reference sequences using

rsem-prepare-reference (Li and Dewey 2011) from Hg19 human RefSeq genes. The references were

then input to rsem-calculate-expression (Li and Dewey 2011) using default parameters for all 6

tissues to obtain TPM based expression values.

Proximal promoter regions were defined as 1,000bp upstream of the gene transcription start sites

based on the longest transcripts for each gene. *Alu*, MIR, L1, L2 and LTR repeat regions were then

identified within the proximal promoters, 5'UTR, CDS and 3'UTR regions.

The Weighted Bootstrap Procedure For Assessing The Effects Of A TE In Each Genic Region.

Many genes contain multiple transposable elements, with only a minority of genes containing a single

TE (Figure S2). In order to assess any effects on transcription due to the presence of a single TE, a

weighted bootstrap approach was devised. For a given TE within each genic region within each

individual tissue, the frequencies of co-occurring TEs and combinations of TEs were noted. Uniform

sampling probabilities were then used for the set of genes containing a specific TE in a specific region,

whilst sampling weights were assigned to genes lacking the specific TE based on TE composition,

such that the TE content of the sampled set of reference genes matched that of the test set of genes,

based on the defined categories. Gene length was divided into 10 bins and these were included as an

additional category when defining sampling weights. This ensured that two gene sets were obtained

for each bootstrap iteration, which were matched in length and TE composition with the sole

difference being the presence of the specific TE within each specific genic region (Figure S5). The

mean difference in expression level, as measured by log(TPM), and the difference in the proportions

of genes detected as expressed were then used as the variables of interest in the bootstrap procedure.

The bootstrap was performed on sets of 1000 genes for 10,000 iterations using the proximal promoter

as defined above, along with 5'UTR and 3'UTRs. When comparing expression levels, genes with

zero read counts were omitted prior to bootstrapping. In order to compensate for multiple testing

considerations, confidence intervals were obtained across the $m = 90$ tests at the level $1 - \alpha/m$,

which is equivalent to the Bonferroni correction, giving confidence intervals, which controlled the

FWER at the level $\alpha = 0.05$. Approximate two-sided p-values were also calculated by finding the

point at which each confidence interval crossed zero, and additional significance was determined by

estimating the FDR on these sets of p-values using the Benjamini-Hochberg method.

Long Intergenic Non-Coding Rnas And Tes. Annotations for 8,196 previously described putative

human lincRNAs were downloaded (Cabili, et al. 2011) and the distribution of TEs within regulatory

elements in lincRNA exons and introns was obtained using the same methods as above. The

previously described regression models were then used to analyse this dataset.

Association Of Functional Elements With Human Repetitive Elements. To demonstrate the

potential functional significance of repetitive elements, the Database for Annotation, Visualization

and Integrated Discovery (DAVID) (Dennis, et al. 2003) was used to perform the GO classification.

We first extracted Gene-IDs from overlapping regions between different gene categories (1000bp

proximal promoter, 5'UTR, 3'UTR, and the combination of these 3 regions) and TEs. These gene-

lists were then submitted to the DAVID Functional Classification Tool. We chose the third level of

GO terms to describe the over-represented functional terms for the three datasets and visualized the

functional over-representation of overlapped genes using the R package *heatmap.2*. The p-value was

applied in the GO analysis as the standard index to determine the degree of enrichment. The threshold

for over-represented GO terms was set to an FDR (Benjamini-Hochberg method) less than 0.05.

Protein-coding genes with Alus were also visualised with the UCSC genome browser

(http://genome.ucsc.edu/) to compare their mRNA with various gene datasets and annotations.

Association Of Alternative Splicing And Protein Coding Regions Containing TEs. In order to

assess the relationship between transposable elements and exonization, an alternative splicing

annotation dataset (SIB Alt-splicing) was downloaded from the UCSC Genome Browser

(http://genome.ucsc.edu/). These data were generated from RefSeq genes, Genbank RNAs and ESTs

that aligned to the human genome. A total of 46,973 alternatively spliced transcripts were intersected

with gene models containing transposable elements.

ACKNOWLEDGEMENT

The authors wish to thank Dan Kortschak, Atma Ivancevic, Joy Raison, Reuben Buckley and Sim Lin
Lim for valuable discussions and critical reading of drafts. This work was supported by grants from
the National Natural Science Foundation of China (61272250 and 61472246) and the National Basic
Research Program of China (2013CB956103). The funders had no role in study design, data
collection and analysis, decision to publish, or preparation of the manuscript.
470

AUTHOR CONTRIBUTION

DLA and CCW conceived, designed and managed the study. LZ collected the datasets, implemented the

analysis pipeline, and analyzed the data. SMP analyzed the data. ZPQ, DFC, ZQH prepared datasets. LZ,

DLA and CCW wrote and revised the manuscript. All authors reviewed and approved the final manuscript.

CONFLICT OF INTEREST

The author(s) declare that they have no competing interests.

REFERENCES

Belancio VP, Hedges DJ, Deininger P 2006. LINE-1 RNA splicing and influences on
mammalian gene expression. Nucleic Acids Res 34: 1512-1521. doi:
10.1093/nar/gkl027
Belancio VP, Roy-Engel AM, Deininger P 2008. The impact of multiple splice sites in
human L1 elements. Gene 411: 38-45. doi: 10.1016/j.gene.2007.12.022
Bolger AM, Lohse M, Usadel B 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30: 2114-2120. doi: 10.1093/bioinformatics/btu170
Britten RJ 1996. DNA sequence insertion and evolutionary variation in gene regulation.
Proc Natl Acad Sci U S A 93: 9374-9377.
Cabili MN, et al. 2011. Integrative annotation of human large intergenic noncoding RNAs
reveals global properties and specific subclasses. Genes Dev 25: 1915-1927. doi:
10.1101/gad.17446611
Callicott JH, et al. 2005. Variation in DISC1 affects hippocampal structure and function
and increases risk for schizophrenia. Proc Natl Acad Sci U S A 102: 8627-8632. doi:
10.1073/pnas.0500515102
Cohen CJ, Lock WM, Mager DL 2009. Endogenous retroviral LTRs as promoters for
human genes: a critical assessment. Gene 448: 105-114. doi:
10.1016/j.gene.2009.06.020
Conley AB, Piriyapongsa J, Jordan IK 2008. Retroviral promoters in the human genome.
Bioinformatics 24: 1563-1567. doi: 10.1093/bioinformatics/btn243
Cordaux R, Batzer MA 2009. The impact of retrotransposons on human genome
evolution. Nat Rev Genet 10: 691-703. doi: 10.1038/nrg2640
Cowley M, Oakey RJ 2013. Transposable elements re-wire and fine-tune the
transcriptome. PLoS Genet 9: e1003234. doi: 10.1371/journal.pgen.1003234
de Souza FS, Franchini LF, Rubinstein M 2013. Exaptation of transposable elements into
novel cis-regulatory elements: is the evidence always strong? Mol Biol Evol 30: 1239-
1251. doi: 10.1093/molbev/mst045
Deininger P 2011. Alu elements: know the SINEs. Genome Biol 12: 236. doi:
10.1186/gb-2011-12-12-236
Deininger PL, Batzer MA 2002. Mammalian retroelements. Genome Res 12: 1455-1465.
doi: 10.1101/gr.282402
Dennis G, Jr., et al. 2003. DAVID: Database for Annotation, Visualization, and Integrated
Discovery. Genome Biol 4: P3.
Dreszer TR, et al. 2012. The UCSC Genome Browser database: extensions and updates
2011. Nucleic Acids Res 40: D918-923. doi: 10.1093/nar/gkr1055
Engreitz JM, et al. 2013. The Xist lncRNA exploits three-dimensional genome
architecture to spread across the X chromosome. Science 341: 1237973. doi:
10.1126/science.1237973
Ernst J, et al. 2011. Mapping and analysis of chromatin state dynamics in nine human
cell types. Nature 473: 43-49. doi: 10.1038/nature09906
Franchini LF, et al. 2011. Convergent evolution of two mammalian neuronal enhancers
by sequential exaptation of unrelated retroposons. Proc Natl Acad Sci U S A 108: 15270-
15275. doi: 10.1073/pnas.1104997108
Franke V, et al. 2017. Long terminal repeats power evolution of genes and gene
expression programs in mammalian oocytes and zygotes. Genome Res. doi:
10.1101/gr.216150.116
Gong C, Maquat LE 2011. lncRNAs transactivate STAU1-mediated mRNA decay by
duplexing with 3' UTRs via Alu elements. Nature 470: 284-288. doi:
10.1038/nature09701

Hadjiargyrou M, Delihas N 2013. The Intertwining of Transposable Elements and Non-
Coding RNAs. Int J Mol Sci 14: 13307-13328. doi: 10.3390/ijms140713307
Han JS, Szak ST, Boeke JD 2004. Transcriptional disruption by the L1 retrotransposon
and implications for mammalian transcriptomes. Nature 429: 268-274. doi:
10.1038/nature02536
Hingorani AD, et al. 1999. A common variant of the endothelial nitric oxide synthase
(Glu298-->Asp) is a major risk factor for coronary artery disease in the UK. Circulation
100: 1515-1520.
Jacques PE, Jeyakani J, Bourque G 2013. The majority of primate-specific regulatory
sequences are derived from transposable elements. PLoS Genet 9: e1003504. doi:
10.1371/journal.pgen.1003504
Jjingo D, et al. 2014. Mammalian-wide interspersed repeat (MIR)-derived enhancers and
the regulation of human gene expression. Mob DNA 5: 14. doi: 10.1186/1759-8753-5-
14
Jjingo D, Huda A, Gundapuneni M, Marino-Ramirez L, Jordan IK 2011. Effect of the
transposable element environment of human genes on gene length and expression.
Genome Biol Evol 3: 259-271. doi: 10.1093/gbe/evr015
Jurka J, et al. 2005. Repbase Update, a database of eukaryotic repetitive elements.
Cytogenet Genome Res 110: 462-467. doi: 10.1159/000084979
Kazazian HH, Jr., et al. 1988. Haemophilia A resulting from de novo insertion of L1
sequences represents a novel mechanism for mutation in man. Nature 332: 164-166.
doi: 10.1038/332164a0
Kelley D, Rinn J 2012. Transposable elements reveal a stem cell-specific class of long
noncoding RNAs. Genome Biol 13: R107. doi: 10.1186/gb-2012-13-11-r107
Kent WJ, et al. 2002. The human genome browser at UCSC. Genome Res 12: 996-1006.
doi: 10.1101/gr.229102. Article published online before print in May 2002
Kim DD, et al. 2004. Widespread RNA editing of embedded alu elements in the human
transcriptome. Genome Res 14: 1719-1725. doi: 10.1101/gr.2855504
Kohany O, Gentles AJ, Hankus L, Jurka J 2006. Annotation, submission and screening of
repetitive elements in Repbase: RepbaseSubmitter and Censor. BMC Bioinformatics 7:
474. doi: 10.1186/1471-2105-7-474
Kunarso G, et al. 2010. Transposable elements have rewired the core regulatory
network of human embryonic stem cells. Nat Genet 42: 631-634. doi: 10.1038/ng.600
Lander ES, et al. 2001. Initial sequencing and analysis of the human genome. Nature
409: 860-921. doi: 10.1038/35057062
Lavie L, Maldener E, Brouha B, Meese EU, Mayer J 2004. The human L1 promoter:
variable transcription initiation sites and a major impact of upstream flanking sequence
on promoter activity. Genome Res 14: 2253-2260. doi: 10.1101/gr.2745804
Lee JY, Ji Z, Tian B 2008. Phylogenetic analysis of mRNA polyadenylation sites reveals a
role of transposable elements in evolution of the 3'-end of genes. Nucleic Acids Res 36:
5581-5590. doi: 10.1093/nar/gkn540
Levanon EY, et al. 2004. Systematic identification of abundant A-to-I editing sites in the
human transcriptome. Nat Biotechnol 22: 1001-1005. doi: 10.1038/nbt996
Lewis BP, Green RE, Brenner SE 2003. Evidence for the widespread coupling of
alternative splicing and nonsense-mediated mRNA decay in humans. Proc Natl Acad Sci
U S A 100: 189-192. doi: 10.1073/pnas.0136770100
Li B, Dewey CN 2011. RSEM: accurate transcript quantification from RNA-Seq data with
or without a reference genome. BMC Bioinformatics 12: 323. doi: 10.1186/1471-2105-
12-323

Lin L, et al. 2009. Large-scale analysis of exonized mammalian-wide interspersed
repeats in primate genomes. Hum Mol Genet 18: 2204-2214. doi: 10.1093/hmg/ddp152
Lynch VJ, Leclerc RD, May G, Wagner GP 2011. Transposon-mediated rewiring of gene
regulatory networks contributed to the evolution of pregnancy in mammals. Nat Genet
43: 1154-1159. doi: 10.1038/ng.917
McCue AD, Slotkin RK 2012. Transposable element small RNAs as regulators of gene
expression. Trends Genet 28: 616-623. doi: 10.1016/j.tig.2012.09.001
Medstrand P, Landry JR, Mager DL 2001. Long terminal repeats are used as alternative
promoters for the endothelin B receptor and apolipoprotein C-I genes in humans. J Biol
Chem 276: 1896-1903. doi: 10.1074/jbc.M006557200
Meischl C, Boer M, Ahlin A, Roos D 2000. A new exon created by intronic insertion of a
rearranged LINE-1 element as the cause of chronic granulomatous disease. Eur J Hum
Genet 8: 697-703. doi: 10.1038/sj.ejhg.5200523
Meyer LR, et al. 2013. The UCSC Genome Browser database: extensions and updates
2013. Nucleic Acids Res 41: D64-69. doi: 10.1093/nar/gks1048
NE II, et al. 2014. Long non-coding RNAs and enhancer RNAs regulate the
lipopolysaccharide-induced inflammatory response in human monocytes. Nat Commun
5: 3979. doi: 10.1038/ncomms4979
Nekrutenko A, Li WH 2001. Transposable elements are found in a large number of
human protein-coding genes. Trends Genet 17: 619-621.
Pacanowski MA, Zineh I, Cooper-Dehoff RM, Pepine CJ, Johnson JA 2009. Genetic and
pharmacogenetic associations between NOS3 polymorphisms, blood pressure, and
cardiovascular events in hypertension. Am J Hypertens 22: 748-753. doi:
10.1038/ajh.2009.81
Pereira V, Enard D, Eyre-Walker A 2009. The effect of transposable element insertions
on gene expression evolution in rodents. PLoS One 4: e4321. doi:
10.1371/journal.pone.0004321
Perepelitsa-Belancio V, Deininger P 2003. RNA truncation by premature
polyadenylation attenuates human mobile element activity. Nat Genet 35: 363-366. doi:
10.1038/ng1269
Piriyapongsa J, Polavarapu N, Borodovsky M, McDonald J 2007. Exonization of the LTR
transposable elements in human genome. BMC Genomics 8: 291. doi: 10.1186/1471-
2164-8-291
Polak P, Domany E 2006. Alu elements contain many binding sites for transcription
factors and may play a role in regulation of developmental processes. BMC Genomics 7:
133. doi: 10.1186/1471-2164-7-133
Pruitt KD, Tatusova T, Maglott DR 2007. NCBI reference sequences (RefSeq): a curated
non-redundant sequence database of genomes, transcripts and proteins. Nucleic Acids
Res 35: D61-65. doi: 10.1093/nar/gkl842
Quinlan AR, Hall IM 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26: 841-842. doi: 10.1093/bioinformatics/btq033
R : A language and environment for statistical computing [Internet]. Vienna, Austria: R
Foundation for Statitical Computing; 2014. Available from: http://www.R-project.org
Rapoport JL, Addington AM, Frangou S, Psych MR 2005. The neurodevelopmental model
of schizophrenia: update 2005. Mol Psychiatry 10: 434-449. doi:
10.1038/sj.mp.4001642
Rebollo R, Romanish MT, Mager DL 2012. Transposable elements: an abundant and
natural source of regulatory sequences for host genes. Annu Rev Genet 46: 21-42. doi:
10.1146/annurev-genet-110711-155621

Sela N, et al. 2007. Comparative analysis of transposed element insertion within human
and mouse genomes reveals Alu's unique role in shaping the human transcriptome.
Genome Biol 8: R127. doi: 10.1186/gb-2007-8-6-r127
Slotkin RK, Martienssen R 2007. Transposable elements and the epigenetic regulation of
the genome. Nat Rev Genet 8: 272-285. doi: 10.1038/nrg2072
Smit AF 1999. Interspersed repeats and other mementos of transposable elements in
mammalian genomes. Curr Opin Genet Dev 9: 657-663.
Trapnell C, et al. 2012. Differential gene and transcript expression analysis of RNA-seq
experiments with TopHat and Cufflinks. Nat Protoc 7: 562-578. doi:
10.1038/nprot.2012.016
van de Lagemaat LN, Landry JR, Mager DL, Medstrand P 2003. Transposable elements in
mammals promote regulatory variation and diversification of genes with specialized
functions. Trends Genet 19: 530-536. doi: 10.1016/j.tig.2003.08.004
Westfall THaFBaP 2008. Simultaneous Inference in General Parametric Models.
Biometrical Journal 50: 346--363.
Wickham H. 2009. ggplot2: elegant graphics for data analysis: Springer New York.
Zeileis A 2004. Econometric Computing with HC and HAC Covariance Matrix Estimators.
Journal of Statistical Software 11(10): 1-17.
643

FIGURE LEGENDS

Fig. 1. Distribution of repetitive elements overlapping with different human gene regions.

Gene regions are shown on the x-axis and the y-axis shows the percentages of the genomic regions

containing repetitive elements. Human-specific repeats were those annotated with "Homo sapiens" or

"primates" as their origin (see Table S8 for the list of human-specific repeat classes). The remaining

repetitive regions were categorized as shared repeats.

Fig.2. Analysis of the co-occurrence of Transposable Elements and Regulatory Elements across

multiple genomic regions A) Probability estimates are those of an individual base within each region

being part of a Regulatory Element, i.e. p(RE), or a Transposable Element within a Regulatory

Element, i.e. p(TE|RE). Error bars indicate ±1 Std. Error as calculated on the logit-transformed

values. B) Pairwise comparisons of TE content in RE based on $1 - \alpha$ Confidence Intervals for the

difference between logit-transformed probabilities p(TE|RE), adjusted for multiple comparisons at the

level $\alpha = 0.05/6$ within each RE (60). Intervals highlighted in red show significant pairwise

differences (confidence intervals do not cross the 0 difference value).

Fig.3. Analysis of the occurrence of Regulatory Elements within specific classes of Transposable

Elements. A) Probability estimates are for an individual base within each type of element belonging

to each of the regulatory elements, i.e. p(RE|TE). Error bars indicate ±1 Std. Error as calculated on the

logit-transformed values. B) Pairwise comparisons of TE content in RE based on $1 - \alpha$ Confidence

Intervals for the difference between logit-transformed probabilities p(RE|TE), adjusted for multiple

comparisons at the level $\alpha = 0.05/6$ within each RE (60). Intervals highlighted in red show

significant pairwise differences (confidence intervals do not cross the 0 difference value).

Fig.4. Analysis of the co-occurrence of Transposable Elements and Regulatory Elements across

non-coding regions A) Probability estimates for an individual base within each type of non-coding

region being part of a Regulatory Element p(RE) or a Transposable Element within each Regulatory

Element p(TE|RE). Error bars indicate ±1 Std. Error as calculated on the logit-transformed values. B)

Pairwise comparisons of TE content in RE based on $1 - \alpha$ Confidence Intervals for the difference

between logit-transformed probabilities p(TE|RE), adjusted for multiple comparisons at the level

$\alpha = 0.05/6$ within each RE (60). Intervals highlighted in red show significant pairwise differences

(confidence intervals do not cross the 0 difference value).

Fig.5. Effects on gene expression of the presence of specific TEs in each gene compartment. A)

Confidence Intervals for the difference in the probability of a gene being detected as expressed due to

the presence of each TE in each genic region. B) Confidence Intervals for the difference in mean

log2(TPM) counts. For both A) and B), Confidence Intervals were obtained using the weighted

bootstrap and are $1 - \alpha/m$ intervals, where $\alpha = 0.05$ and $m = 90$ as the total number of intervals

presented. Dots represent the median value from the bootstrap procedure, whilst the vertical line

indicates zero. Intervals which do not cross zero are coloured red, and indicate a rejection of the null

hypothesis, $H_0: \Delta\theta = 0$, where θ represents the parameter of interest.

Fig.6. Enrichment of GO terms of genes containing TEs in promoter, 5'UTR and 3'UTR

regions.

Enrichment of GO terms of genes containing TEs in "Biological Process". Genes containing different

types of repetitive elements in the proximal promoter regions are labeled as "Promoter with Repeats",

and Genes containing repetitive elements in UTR regions are labeled as "5/3UTR with Repeats".

Genes named "Combined Repeats" are the combined data from 3 regions we mentioned above. The

darker the color, the greater the GO term enrichment as determined by FDR.

Biological Process

GO:0006935 chemotaxis, taxis, locomotory behavior
GO:0006909 rapid-early process
GO:0007586 blood coagulation
GO:0009611 response to wounding
GO:0051851 neuromuscular synaptic process
GO:0009653 neuromuscular biosynthetic process
GO:0007155 cell adhesion
GO:0009888 regulation of biosynthetic process
GO:0032535 regulation of metabolic process
GO:0051252 regulation of cellular metabolic process
GO:0048519 cellular nitrogen compound biosynthetic process
GO:0016109 cellular aromatic compound metabolic process
GO:0006820 anion transport
GO:0015671 regulation of heteroatom process
GO:0006952 regulation of body fluid levels
GO:0009058 regulation of nitrogen compound metabolic process
GO:0009059 regulation of cellular biosynthetic process
GO:0010556 regulation of primary metabolic process