1

Detailed analyses of stall force generation in *Mycoplasma mobile* gliding

Masaki Mizutani[a], Isil Tulum[a,b], Yoshiaki Kinosita[c], Takayuki Nishizaka[c] and Makoto Miyata[a,b]

[a] Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

[b] The OCU Advanced Research Institute for Natural Science and Technology (OCARINA), Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

[c] Department of Physics, Faculty of Science, Gakushuin University, 1-5-1 Mejiro, Toshima-ku, Tokyo 171-8588, Japan

Corresponding author: (Makoto Miyata, Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan, +81-6-6605-3157, miyata@sci.osaka-cu.ac.jp)

Keywords: gliding motility, optical tweezers, force measurement, gear ratio, energy conversion

ABSTRACT

Mycoplasma mobile is a bacterium that uses a unique mechanism to glide on solid surfaces at a velocity of up to 4.5 µm/s. Its gliding machinery comprises hundreds of units that generate the force for gliding based on the energy derived from ATP; the units catch and pull on sialylated oligosaccharides fixed to solid surfaces. In the present study, we measured the stall force of wild-type and mutant strains of *M. mobile* carrying a bead manipulated using optical tweezers. The strains that had been enhanced for binding exhibited weaker stall forces than the wild-type strain, indicating that stall force is related to force generation rather than to binding. The stall force of the wild-type strain decreased linearly from 113 to 19 pN following the addition of 0–0.5 mM free sialyllactose (a sialylated oligosaccharide), with a decrease in the number of working units. Following the addition of 0.5 mM sialyllactose, the cells carrying a bead loaded using optical tweezers exhibited stepwise movements with force increments. The force increments ranged from 1 to 2 pN. Considering the 70-nm step size, this small unit force may be explained by the large gear ratio involved in the *M. mobile* gliding machinery.

SIGNIFICANCE

Mycoplasma is a genus of bacteria that parasitizes animals. Dozens of *Mycoplasma* species glide over the tissues of their hosts during infection. The gliding machinery of *Mycoplasma mobile*, the fastest species, includes intracellular motors and hundreds of legs on the cell surface. In the present study, we precisely measured force generation using a highly focused laser beam arrangement (referred to as optical tweezers) under various conditions. The measurements obtained in this study suggest that the rapid gliding exhibited by *M. mobile* arises from the large gear ratio of its gliding machinery.

INTRODUCTION

Members of the bacterial class *Mollicutes*, which includes the genus *Mycoplasma*, are parasitic and occasionally commensal bacteria that are characterized by small cells and genomes, and by the absence of a peptidoglycan layer (1, 2). Dozens of parasitic *Mycoplasma* species, such as the fish pathogen *Mycoplasma mobile* (3-5) and the human pathogen *Mycoplasma pneumoniae* (6-8), have protrusions, and exhibit gliding

motility in the direction of the protrusions on solid surfaces, which enables

mycoplasmas to parasitize other organisms.   Interestingly, *Mycoplasma* gliding does

not involve flagella or pili, and is completely unrelated to other bacterial motility

systems, or the conventional motor proteins that are common in eukaryotic motility.

M. mobile, which can be isolated from the gills of freshwater fish, is a fast-gliding

Mycoplasma (9-13).   It glides smoothly and continuously on glass at an average speed

of 2.0–4.5 µm/s, or 3–7 times the length of the cell per second.   A working model,

called the centipede or power-stroke model, has been proposed to explain the gliding

mechanism of *M. mobile*: the cells repeatedly catch, pull, drag, and release the sialylated

oligosaccharides (SOs) on solid surfaces (3-5, 14).   The gliding machinery comprises

internal and surface structures (15-19).   The internal structure includes the α- and β-

subunit paralogs of F-type ATPase/synthase, and generates the force for gliding, based

on the energy derived from ATP (18-22).   The force is probably transmitted across the

cell membrane to the surface structure, which is composed of at least three huge

proteins, Gli123, Gli521, and Gli349 (15-17, 23).   Gli521, the crank protein, transmits

the interior force of the cell to Gli349, with actual structural changes (17, 24).   Gli349,

the leg protein, extends after thermal fluctuation, and catches the SOs, which are the

predominant structures on animal cell surfaces (Fig 1*A*) (9, 25-27).   The cells always

glide in the direction of the machinery, which may be a result of the directed binding of

the cells on solid surfaces (3, 28).   In theory, hundreds of gliding units on the cell

surface should act cooperatively to enable smooth gliding (Fig. 1*B*) (15, 27, 29).   To

examine this working model in detail, it is necessary to investigate the behavior of the

individual units.   Recently, the discrete movements involved in gliding motility, which

are possibly attributable to single leg movements, have been observed by controlling the

working leg number following the addition of free SOs (22, 27).   In the current study,

we focused on the pulling force exerted on the solid surface, because the generation of

force has not been properly investigated, with the exception of the final stall force of the

wild-type strain (11).   Therefore, in the present study, we quantitatively measured the

pulling forces of several strains of whole cells under various conditions using optical

tweezers (11, 28, 30-32), and characterized the force generated by the gliding

machinery.

RESULTS

Stall force measurement.　The propelling force in *M. mobile* cells has been measured using optical tweezers (11).　A bead bound to a cell was trapped by a highly focused laser beam, and the force was calculated by measuring the distance between the center of the bead and the trap; the force acting on the bead increased linearly with the displacement from the trap center (11, 28, 32).　In the present study, *M. mobile* cells, which had been biotinylated and suspended in phosphate-buffered saline with glucose (PBS/G), were inserted into a tunnel chamber.　Polystyrene beads coated with avidin (1-µm in diameter) were subsequently added to the tunnel.　A bead trapped using optical tweezers was attached to the back end of the gliding cell by exploiting the avidin–biotin interaction (28, 33, 34).　The cell pulled the bead from the trap center (Fig. 2*A* and Movie S1). Starting from 0 s, the pulling force increased and reached a plateau at approximately 40 s.　The maximum value of the average of 25 data points was used as the stall force (Fig. 2*B*).　The stall force of the wild-type strain was 113 ± 32 pN ($n = 50$).

To determine the proteins involved in force generation or force transmission, we measured the stall force in the wild-type strain of *M. mobile* to compare it with that in six previously isolated strains with known gliding speeds and/or binding activities (12, 21, 35).　The *gli521* (P476R) mutant of protein Gli521 has a single amino acid substitution (proline for arginine at the 476th position) (21), but mutations in other strains have not been described.　The pulling forces of the mutants increased from 0 s and stalled at 20–50 s.　The stall force of the m14 strain—110 ± 29 pN ($n = 29$)—was not significantly different from that of the wild-type strain ($P = 0.1 > 0.05$, according to the Student's *t*-test).　The stall forces of the *gli521* (P476R), m6, m27, m29, and m34 mutants were significantly reduced to 64–81% of that of the wild-type strain ($P = 6 \times 10^{-3}$, 4×10^{-4}, 9×10^{-5}, 8×10^{-9}, and 4×10^{-4}, respectively; $P < 0.01$, according to the Student's *t*-test) (Fig. 2*B Inset*).　The *gli521* (P476R) mutant, which was characterized by enhanced binding, exhibited a smaller force than the wild-type strain, suggesting that the stability of binding is not the only determinant of stall force.

Genome sequencing of various strains.　To determine the proteins and mutations associated with the decreases in stall force, we sequenced the genomes of the strains

using a MiSeq sequencer.   The genome of the *gli521* (P476R) mutant has been

sequenced for a 30,469-bp DNA region encoding four open reading frames (ORFs):

gli123, *gli349*, *gli521*, and *gli42*.   However, the other regions remain unknown (21).

The genome sequencing result for the *gli521* (P476R) mutant was consistent with the

previous report of the mutation in *gli521* (21), and showed additional mutations in other

regions (Table S1).   One of the additional mutations causes an amino acid substitution

in MvspB, a surface protein (23, 36-39).   However, the reduced stall force should be

caused by the mutation in Gli521, because MvspB accounts for only 1.2% of the mass

of all the surface proteins, and the antibody against an abundant and closely related

protein, MvspI, did not influence the gliding motility (23, 39).   The genomes of the

m6, m14, m27, m29, and m34 strains have not been sequenced yet, and we identified

various mutations in the present study.   We suggest that the decrease in the stall force

in the m27 strain in the present study was caused by the mutation in *gli521*.   All the

strains had the same single amino acid substitution (serine to isoleucine) as the 354th

residue in MMOB1700, a homolog of ABC transporter permease based on a Basic

Local Alignment Search Tool (BLAST) search.   This mutation may have been derived

from a substitution that occurred on the clone used for the reported genome sequencing,

because it was derived from the same origin (ATCC 43663) as the present study (36).

Interestingly, MMOB1700 exhibited 5 other mutations in the 10 strains analyzed,

suggesting a special mechanism that causes a high rate of mutation in this gene.   Next,

we sequenced the genomes of nonbinding strains m12, m13, and m23, which have

mutations in *gli123*, *gli349*, and *gli349*, respectively (12, 15, 16, 35).   Genome

sequencing revealed that the identified mutations were consistent with a previous report

(35), although additional mutations were identified in other regions.

Binding and gliding of various strains.   To systematically clarify the relationship

between the features and mutation in the genome, we examined the binding activities

and the gliding speeds for the wild-type, *gli521* (P476R), m6, m14, m27, m29, and m34

mutants, which can glide.   The cell suspensions were adjusted to the same optical

density and inserted into a tunnel chamber.   After 15 min, we videoed the cells to

count the numbers on the glass for the bound-cell ratio, and to determine their gliding

speed, as previously reported (Fig. 3*A*) (29, 35, 40).   The binding activities and the

gliding speeds were averaged for 20 independent images and 100 cells, respectively.

The binding activity and gliding speed of the *gli521* (P476R) mutant were consistent

with those reported in the previous study (35).   Other strains have not been analyzed

by the method used here.   The characteristics of the binding activities allowed

classification of the strains into three groups (Fig. 3*B* left): (i) m6 had 44% of the

activity of the wild-type strain; (ii) m14, m27, and m29 had 80%, 92%, and 73% of the

activity, respectively; and (iii) the *gli521* (P476R) and m34 mutants had 159% and

145% of the activity of the wild-type strain, respectively.   We compared these data to

the binding activities previously estimated from hemadsorption, the adsorption of

erythrocytes onto the surface of colonies (12).   The hemadsorption values of m6, m14,

m27, m29, and m34 were 24%, 93%, 113%, 96%, and 122%, respectively; i.e., except

for m27, they were consistent with the results of the analyses in the present study.   The

gliding speed of the wild-type strain was 3.7 ± 0.2 µm/s and the relative gliding speeds

of the mutants ranged from 80% to 103% of the wild-type strain, showing that the

gliding speeds differed less than the binding activities (Fig. 3*B* right).   Of the strains

enhanced for binding, the *gli521* (P476R) and m34 mutants had reduced stall forces,

suggesting that stall force is not determined simply by binding activity.

Interestingly, we noticed that the proportions of the nongliding bound cells were much

higher in the m6, m14, m27, and m29 strains than in the wild-type strain.   The

proportion of nongliding cells in the cells on glass was 6% for the wild-type strain, as

previously reported (40), but those for the m6, m27, and m29 strains were higher (18%,

19%, and 16%, respectively), and that of the m14 strain was much higher (62%) (Fig.

S1).   This observation can be explained by assuming that gliding can be achieved by

the concerted action of many molecules and interactions.

Effect of SOs on stall force.   An *M. mobile* cell glides as a result of the integrated

movement of its many legs.   However, if the number of working legs is reduced, it is

possible to detect the pulling force of single unit.   Previous studies have shown that the

number of working legs can be reduced by adding free SOs (9, 22, 25, 27, 40, 41).   We

therefore added various concentrations of free sialyllactose (SL)—an SO—to gliding *M.*

mobile cells, and measured the stall force at 200 and 500 frames/s with 0–0.25, 0.33,

and 0.5 mM free SL.   The cells reached a plateau after 30–40 s, and the time taken to

reach the stall increased with the concentration of free SL.   The stall force decreased

from 113 to 19 pN following the addition of up to 0.50 mM free SL (Fig. 4*A* and *B*).

These results suggest that the stall force is the sum of the pulling force of many units.

To detect the pulling force attributable to smaller numbers of units, we reduced the laser

power of the optical tweezers to achieve a higher trace resolution.   In this experiment,

we applied 0.25 mM SL with a reduction of the trap stiffness of the optical tweezers

from 0.5–0.7 to 0.1 pN/nm, whereby the trapped cells were able to escape from the trap

center, and then measured the force under these conditions.   The force increments with

repeated small peaks were detected, and the peaks were measured using a peak-finding

algorithm, as summarized in Fig. 5*A*.   The distribution of these increments was

determined by fitting the sum of four Gaussian curves whose peaks were positioned at

1.1, 2.0, 3.2, and 4.4 pN (Fig. 5*B*).   The peaks were positioned at twice, three, and four

times the value of the first peak, suggesting that these peaks reflect single, double,

triple, and quadruple the minimum force increment (42).   In the same way, the

individual increments were analyzed for the *gli521* (P476R) mutant, and were also

determined by fitting the sum of four Gaussian curves whose peaks were positioned at

0.9, 1.8, 2.6, and 3.5 pN (Fig. 5*B*).

Stepwise force increments.   Because the force increments were detected for limited

leg numbers, it was possible to derive the force increments from the force generated by

a single leg or a minimum force generation unit.   Next, we added 0.50 mM SL to limit

the number of working legs, and reduced the trap stiffness to 0.06–0.07 pN/nm to detect

the minimum force increments more precisely.   Very small ratios of cells remained on

the glass surface under these conditions, and we attached a bead to the cells using the

optical tweezers.   Displacements were detected in 51 of 63 cells.   Eight cells glided

by creeping displacements with occasional discontinuous increments, which were

mostly stepwise (Fig. 6*A*) and sometimes exhibited small peaks, as shown in Figure 5.

Three cells exhibited force increments with more than six continuous steps (Fig. 6*B*).

The average value of the force increments from 46 steps or peaks in 11 cell trajectories

was 1.45 ± 0.44 pN (Fig. 6*C*).   These results suggest that the minimum force increment

was 1–2 pN, which should be the minimum unit force for gliding.

DISCUSSION

Mutations influencing gliding.   In the present study, we sequenced the whole

genomes of *M. mobile* mutants characterized by binding ability, gliding motility, and

colony spreading.   Based on these results, we propose the proteins responsible for or

related to these characteristics.   The m6 strain, which exhibited reduced binding, a

lower gliding speed, and a weaker stall force, has mutations in FtsH, MvspI, and SecY

(37-39, 43, 44).   The substituted amino acid in FtsH is not conserved in other

mycoplasmas except for *Mycoplasma pulmonis*, a closely related species.   MvspI is

probably not involved in gliding, as described above (23).   SecY is generally essential

for protein secretion in bacteria (44).   The substitution in SecY probably affects the

secretion of gliding proteins, resulting in a reduction in the binding activity of cells,

because the amino acid substituted in the mutant is conserved in many mycoplasmas,

including *Mycoplasma hominis*, *Mycoplasma bovis*, and *M. pulmonis*.   The m14 strain

was characterized by reduced binding and gliding, and its glucokinase, which

phosphorylates glucose to glucose-6-phosphate at the first step of glycolysis, has an

amino acid substitution (45).   Interestingly, the m14 colony was less well dispersed

than the wild-type colony, suggesting that glucokinase is related to gliding including

chemotaxis, because colony shape is largely influenced by motility in many bacteria

(12).   The m27 strain, which was characterized by a small proportion of gliding cells,

has a substitution at the 1461st amino acid of a total 4727 amino acids in the coded

Gli521 protein, suggesting that the structure around this position is indispensable for

gliding.   The m34 strain was characterized by enhanced binding, and has a substitution

in the conserved amino acid of the β subunit of F-type ATPase (46).   The membrane

potential may be related to binding activity, because F-type ATPase is responsible for

the membrane potential in *M. mobile*.

Cell behavior in stall.   Each cell was stalled using optical tweezers focused on the

bead bound to the back end of the cell.   What events can be expected in the stall?

The stall force decreased with the addition of free SL (Fig. 4).   This observation

suggests that the legs repeatedly catch, pull, and release SOs, even in the stalled state

(Fig. 7), because the force under free SOs should increase to the stall force in the

absence of SOs if the legs do not detach in stall.   In our gliding model, it was

suggested that the legs detach as a result of the tension caused by continuous cell

displacement in gliding (14, 28, 41).   The putative detachment in the stall may suggest

that directed detachment occurs with much shorter displacement than expected from a

95-nm long leg structure, because the detachment also occurs in the stall.   This

assumption can explain the observation that the *gli521* (P476R) and m34 mutants

exhibited a smaller stall force than the wild-type strain, although they had a higher ratio

of cells bound to the glass (Fig. 2 and 3).   The higher ratio of bound cells was probably

due to the reduced force required to detach the post-stroke legs.

Unit number of gliding machinery.   The minimum force increment did not change

significantly with the distance from the laser.   Because the force increments occurred

additionally to the previous ones, the gliding unit should have generated the same force

constantly over a rather long distance ranging from 0 to 200 nm (Fig. 6).

In the present study, the stall force and the minimum force increments of a cell were

approximately 113 and 1.5 pN, respectively (Fig. 2*B* and 6*C*).   Previously, the number

of legs in *M. mobile* was estimated to be 450 (16).   The number of working units was

calculated from the stall force and the minimum force increments of a cell; 113 over 1.5

pN is calculated to be approximately 75-fold, suggesting that 75 minimum units can

work simultaneously.   Assuming that the minimum unit of force corresponds to a

single molecule of Gli349, one-sixth of the Gli349 molecules simultaneously suggested

to participate in force generation in the stalled state (Fig. 1*B*).   The friction occurring at

the interface between an *M. mobile* cell and water flow in the interlamellar of a carp gill

is calculated to be 34 pN for the maximum, based on Stokes' law (11, 47).   This

number is three-fold smaller than the stall force of a cell, suggesting that a cell can glide

against water flow using the force generated by the simultaneous strokes of many legs.

***M. mobile* gliding is characterized by large steps and a small force.**   We compared

the step size and the force of *M. mobile* gliding with those of conventional motor

proteins such as myosin, dynein, and kinesin, which perform stepwise movements along

the rail proteins driven by the energy derived from ATP.   The step sizes and the forces

of myosin-II, cytoplasmic dynein, kinesin, and myosin-V have been reported as 5.3, 8,

8, and 36 nm, and 3–5, 7–8, 8, and 2–3 pN, respectively (48-53). The long step and

small force of myosin-V are caused by the lever effect of 26-nm arms (54). The step

size of *M. mobile* is 70 nm; much larger than that of conventional motor proteins (22).

The minimum unit force calculated here (1–2 pN) suggests the gear effect in the gliding

machinery. Gli521, which is the force transmitter, forms a triskelion with 100-nm

arms (24), and Gli349, which is the leg, is shaped like an eighth note in musical notation

with a 50-nm flexible string (26, 55), suggesting that these proteins cause the lever

effects.

Energy conversion efficiency of *M. mobile* gliding. The direct energy source for *M.*

mobile gliding is ATP; based on experiments that permeabilize cells, a "gliding ghost"

can be reactivated by ATP (21). A gliding ghost exhibits stepwise movement with a

dwell time that is dependent on the ATP concentration used, suggesting that the step is

coupled to ATP hydrolysis (22). Based on a minimum unit force of 1–2 pN and a

spring constant of 0.06–0.07 pN/nm, the work done per step, W_{step}, is calculated to be 8–

33 pN nm from the equation $W_{step} = 1/2 \times$ spring constant $\times$ displacement^2. Assuming

that one ATP molecule is consumed per step, the energy conversion efficiency of *M.*

mobile gliding can be calculated to be approximately 10–40%, because generally

approximately 80 pN nm free energy is available from the hydrolysis of one ATP

molecule.

F-type ATPase attains 100% energy conversion efficiency (56). It has been suggested

that the gliding machinery of *M. mobile* is driven by the α- and β-subunit paralogs of F-

type ATPase (18). The force transmission from this motor to the solid surfaces through

several large components including Gli521 and Gli349 may be related to the putative

energy loss.

MATERIALS AND METHODS

Strains and cultivation. The *M. mobile* strain 163K (ATCC 43663) was used as the

wild-type, and its 9 mutants were grown in Aluotto medium at 25°C, as previously

described (12, 35, 40, 57).

Surface modifications of *M. mobile* cells and polystyrene beads. The cultured cells

were washed with PBS/G consisting of 75 mM sodium phosphate (pH 7.3), 68 mM

NaCl, and 20 mM glucose, suspended in 1.0 mM Sulfo-NHS-LC-LC-biotin (Thermo

Scientific, Waltham, MA) in PBS/G, and kept for 15 min at room temperature (RT), as

previously described (28, 29, 33, 34, 40).   Polystyrene beads (1.0 μm in diameter)

(Polysciences, Warrington, PA) were coated with avidin (Sigma-Aldrich, St. Louis,

MO), as previously described (28).

Force measurements.   The avidin-coated beads were attached to biotinylated cells in

two different ways, according to the concentrations of free SL used in the experiments.

In force measurements under 0–0.13 mM SL conditions, the biotinylated cells were

inserted into a tunnel chamber that had been precoated with 10% horse serum (20, 27,

29).   Avidin-coated beads were sonicated, inserted into the tunnel chamber with various

concentrations of free SL in PBS/G, and bound to the cells (32).   In force

measurements under 0.25–0.50 mM SL conditions, avidin-coated beads were sonicated

and mixed with biotinylated cells in a microtube, and kept for 10–30 min at RT.   Then,

the mixture was inserted into a tunnel chamber and kept for 15 min at RT.   The

chamber was washed with PBS/G and the PBS/G was replaced by various

concentrations of free SL in PBS/G.   Both ends of the tunnel were sealed with nail

polish.   The bead movements were recorded at 200 or 500 frames/s and analyzed by

displacement of up to 250 nm from the trap center (the linear range of the laser trap)

using ImageJ 1.43u (http://rsb.info.nih.gov/ij/) and IGOR Pro 6.33J (WaveMetrics,

Portland, OR) software packages (22, 28, 32, 58).

Genome sequencing of the various strains.   All the strains were plated and isolated

as previously described (19).   The genomic DNAs were isolated using a QIAGEN

DNeasy Blood & Tissue kit (QIAGEN, Hilden, Germany).   The isolated genomic

DNA was sequenced using MiSeq (Illumina Inc., San Diego, CA) and mapped by CLC

Genomics Workbench 8 (QIAGEN, Hilden, Germany).

Characterization of binding and gliding of the various strains.   All the strains were

cultured to reach an optical density at 600 nm of 0.08.   They were then suspended and

inserted into a tunnel chamber (27, 29).   Cell behavior was recorded and analyzed as

previously reported (27, 35).

ACKNOWLEDGMENTS

This work was supported by a Grant-in-Aid for Scientific Research in the innovative

area "Harmonized Supramolecular Motility Machinery and Its Diversity" (MEXT

KAKENHI; Grant Number 24117002), and by Grants-in-Aid for Scientific Research

(B) and (A) (MEXT KAKENHI; Grant Numbers 24390107 and 17H01544) to MM.

REFERENCES

1. Razin S, Yogev D, & Naot Y (1998) Molecular biology and pathogenicity of
mycoplasmas. *Microbiol Mol Biol Rev* 62(4):1094-1156.

2. Razin S & Hayflick L (2010) Highlights of mycoplasma research--an historical
perspective. *Biologicals* 38(2):183-190.

3. Miyata M (2008) Centipede and inchworm models to explain *Mycoplasma* gliding.
Trends Microbiol 16(1):6-12.

4. Miyata M (2010) Unique centipede mechanism of *Mycoplasma* gliding. *Annu Rev*
Microbiol 64:519-537.

5. Miyata M & Hamaguchi T (2016) Prospects for the gliding mechanism of
Mycoplasma mobile. *Curr Opin Microbiol* 29:15-21.

6. Nakane D, Kenri T, Matsuo L, & Miyata M (2015) Systematic structural analyses
of attachment organelle in *Mycoplasma pneumoniae*. *PLoS Pathog*
11(12):e1005299.

7. Miyata M & Hamaguchi T (2016) Integrated information and prospects for gliding
mechanism of the pathogenic bacterium *Mycoplasma pneumoniae*. *Front Microbiol*
7(960).

8. Kawamoto A, *et al.* (2016) Periodicity in attachment organelle revealed by electron
cryotomography suggests conformational changes in gliding mechanism of
Mycoplasma pneumoniae. *MBio* 7(2):e00243-00216.

9. Morio H, Kasai T, & Miyata M (2015) Gliding direction of *Mycoplasma mobile*. *J*
Bacteriol 198(2):283-290.

10. Lee W, *et al.* (2015) Three-dimensional superlocalization imaging of gliding
Mycoplasma mobile by extraordinary light transmission through arrayed nanoholes.
ACS Nano 9(11):10896-10908.

11. Miyata M, Ryu WS, & Berg HC (2002) Force and velocity of *Mycoplasma mobile*
gliding. *J Bacteriol* 184(7):1827-1831.

12. Miyata M, *et al.* (2000) Gliding mutants of *Mycoplasma mobile*: relationships
between motility and cell morphology, cell adhesion and microcolony formation.
Microbiology 146 (Pt 6):1311-1320.

13. Rosengarten R & Kirchhoff H (1987) Gliding motility of *Mycoplasma* sp. nov.
strain 163K. *J Bacteriol* 169(5):1891-1898.

14. Chen J, Neu J, Miyata M, & Oster G (2009) Motor-substrate interactions in

mycoplasma motility explains non-Arrhenius temperature dependence. *Biophys J*

97(11):2930-2938.

15. Uenoyama A, Kusumoto A, & Miyata M (2004) Identification of a 349-kilodalton

protein (Gli349) responsible for cytadherence and glass binding during gliding of

Mycoplasma mobile. *J Bacteriol* 186(5):1537-1545.

16. Uenoyama A & Miyata M (2005) Identification of a 123-kilodalton protein

(Gli123) involved in machinery for gliding motility of *Mycoplasma mobile*. *J*

Bacteriol 187(16):5578-5584.

17. Seto S, Uenoyama A, & Miyata M (2005) Identification of a 521-kilodalton protein

(Gli521) involved in force generation or force transmission for *Mycoplasma mobile*

gliding. *J Bacteriol* 187(3):3502-3510.

18. Nakane D & Miyata M (2007) Cytoskeletal "Jellyfish" structure of *Mycoplasma*

mobile. *Proc. Natl. Acad. Sci. USA* 104(49):19518-19523.

19. Tulum I, Yabe M, Uenoyama A, & Miyata M (2014) Localization of P42 and F_1-

ATPase alpha-subunit homolog of the gliding machinery in *Mycoplasma mobile*

revealed by newly developed gene manipulation and fluorescent protein tagging. *J*

Bacteriol 196(10):1815-1824.

20. Jaffe JD, Miyata M, & Berg HC (2004) Energetics of gliding motility in

Mycoplasma mobile. *J Bacteriol* 186(13):4254-4261.

21. Uenoyama A & Miyata M (2005) Gliding ghosts of *Mycoplasma mobile*. *Proc Natl*

Acad Sci USA 102(36):12754-12758.

22. Kinosita Y*, et al.* (2014) Unitary step of gliding machinery in *Mycoplasma mobile*.

Proc. Natl. Acad. Sci. USA 111(23):8601-8606.

23. Kusumoto A, Seto S, Jaffe JD, & Miyata M (2004) Cell surface differentiation of

Mycoplasma mobile visualized by surface protein localization. *Microbiology* 150(Pt

12):4001-4008.

24. Nonaka T, Adan-Kubo J, & Miyata M (2010) Triskelion structure of the Gli521

protein, involved in the gliding mechanism of *Mycoplasma mobile*. *J Bacteriol*

192(3):636-642.

25. Nagai R & Miyata M (2006) Gliding motility of *Mycoplasma mobile* can occur by

repeated binding to *N*-acetylneuraminyllactose (sialyllactose) fixed on solid

surfaces. *J Bacteriol* 188(18):6469-6475.

26. Adan-Kubo J, Uenoyama A, Arata T, & Miyata M (2006) Morphology of isolated
Gli349, a leg protein responsible for *Mycoplasma mobile* gliding via glass binding,
revealed by rotary shadowing electron microscopy. *J Bacteriol* 188(8):2821-2828.

27. Kasai T*, et al.* (2013) Role of binding in *Mycoplasma mobile* and *Mycoplasma*
pneumoniae gliding analyzed through inhibition by synthesized sialylated
compounds. *J Bacteriol* 195(3):429-435.

28. Tanaka A, Nakane D, Mizutani M, Nishizaka T, & Miyata M (2016) Directed
binding of gliding bacterium, *Mycoplasma mobile*, shown by detachment force and
bond lifetime. *MBio* 7(3):e00455-00416.

29. Nakane D & Miyata M (2012) *Mycoplasma mobile* cells elongated by detergent and
their pivoting movements in gliding. *J Bacteriol* 194(1):122-130.

30. Ashkin A, Dziedzic JM, Bjorkholm JE, & Chu S (1986) Observation of a single-
beam gradient force optical trap for dielectric particles. *Opt Lett* 11(5):288.

31. Nishizaka T, Miyata H, Yoshikawa H, Ishiwata S, & Kinosita K, Jr. (1995)
Unbinding force of a single motor molecule of muscle measured using optical
tweezers. *Nature* 377(6546):251-254.

32. Mizutani M & Miyata M (2017) Force measurement on *Mycoplasma mobile*
gliding using optical tweezers. *Bio-protocol* 7(3):e2127

33. Hiratsuka Y, Miyata M, & Uyeda TQ (2005) Living microtransporter by uni-
directional gliding of *Mycoplasma* along microtracks. *Biochem Biophys Res*
Commun 331(1):318-324.

34. Hiratsuka Y, Miyata M, Tada T, & Uyeda TQ (2006) A microrotary motor powered
by bacteria. *Proc Natl Acad Sci U S A* 103(37):13618-13623.

35. Uenoyama A, Seto S, Nakane D, & Miyata M (2009) Regions on Gli349 and
Gli521 protein molecules directly involved in movements of *Mycoplasma mobile*
gliding machinery, suggested by use of inhibitory antibodies and mutants. *J*
Bacteriol 191(6):1982-1985.

36. Jaffe JD*, et al.* (2004) The complete genome and proteome of *Mycoplasma mobile*.
Genome Res 14(8):1447-1461.

37. Adan-Kubo J, Yoshii SH, Kono H, & Miyata M (2012) Molecular structure of
isolated MvspI, a variable surface protein of the fish pathogen *Mycoplasma mobile*.

J Bacteriol 194(12):3050-3057.

38. Wu HN, Kawaguchi C, Nakane D, & Miyata M (2012) "Mycoplasmal antigen
modulation," a novel surface variation suggested for a lipoprotein specifically
localized on *Mycoplasma mobile*. *Curr Microbiol* 64(5):433-440.

39. Wu HN & Miyata M (2012) Whole surface image of *Mycoplasma mobile*,
suggested by protein identification and immunofluorescence microscopy. *J*
Bacteriol 194(21):5848-5855.

40. Kasai T, Hamaguchi T, & Miyata M (2015) Gliding motility of *Mycoplasma mobile*
on uniform oligosaccharides. *J Bacteriol* 197(18):2952-2957.

41. Kasai T & Miyata M (2013) Analyzing inhibitory effects of reagents on
Mycoplasma gliding and adhesion. *Bio-protocol* 3(14):e829.

42. Leidel C, Longoria RA, Gutierrez FM, & Shubeita GT (2012) Measuring molecular
motor forces in vivo: implications for tug-of-war models of bidirectional transport.
Biophys J 103(3):492-500.

43. Ito K & Akiyama Y (2005) Cellular functions, mechanism of action, and regulation
of FtsH protease. *Annu Rev Microbiol* 59:211-231.

44. Mori H & Ito K (2006) Different modes of SecY-SecA interactions revealed by site-
directed in vivo photo-cross-linking. *Proc Natl Acad Sci U S A* 103(44):16159-
16164.

45. Gaurivaud P*, et al.* (2016) *Mycoplasma agalactiae* secretion of beta-(1-->6)-glucan,
a rare polysaccharide in prokaryotes, is governed by high-frequency phase
variation. *Appl Environ Microbiol* 82(11):3370-3383.

46. Beven L*, et al.* (2012) Specific evolution of F1-like ATPases in mycoplasmas. *PLoS*
One 7(6):e38793.

47. Lauder GV (1984) Pressure and water flow patterns in the respiratory tract of the
bass (*Micropterus salmoides*). *Journal of Experimental Biology* 113:151-164.

48. Schnitzer MJ & Block SM (1997) Kinesin hydrolyses one ATP per 8-nm step.
Nature 388(6640):386-390.

49. Kojima H, Muto E, Higuchi H, & Yanagida T (1997) Mechanics of single kinesin
molecules measured by optical trapping nanometry. *Biophys J* 73(4):2012-2022.

50. Takagi Y, Homsher EE, Goldman YE, & Shuman H (2006) Force generation in
single conventional actomyosin complexes under high dynamic load. *Biophys J*

90(4):1295-1307.

51. Fujita K, Iwaki M, Iwane AH, Marcucci L, & Yanagida T (2012) Switching of

myosin-V motion between the lever-arm swing and brownian search-and-catch. *Nat*

Commun 3(956).

52. Park PJ & Lee KJ (2013) A modified active Brownian dynamics model using

asymmetric energy conversion and its application to the molecular motor system. *J*

Biol Phys 39(3):439-452.

53. Gennerich A, Carter AP, Reck-Peterson SL, & Vale RD (2007) Force-induced

bidirectional stepping of cytoplasmic dynein. *Cell* 131(5):952-965.

54. Vilfan A (2005) Elastic lever-arm model for myosin V. *Biophys J* 88(6):3792-3805.

55. Metsugi S*, et al.* (2005) Sequence analysis of the gliding protein Gli349 in

Mycoplasma mobile. *Biophysics (Nagoya-shi)* 1:33-43.

56. Kinosita K, Jr., Yasuda R, Noji H, & Adachi K (2000) A rotary molecular motor

that can work at near 100% efficiency. *Philos Trans R Soc Lond B Biol Sci*

355(1396):473-489.

57. Aluotto BB, Wittler RG, Williams CO, & Faber JE (1970) Standardized

bacteriologic techniques for the characterization of *Mycoplasma* species. *Int. J.*

Syst. Bacteriol. 20:35-58.

58. Kinosita Y, Uchida N, Nakane D, & Nishizaka T (2016) Direct observation of

rotation and steps of the archaellum in the swimming halophilic archaeon

Halobacterium salinarum. *Nat Microbiol* 1(11):16148.

 FIGURE LEGENDS

Fig. 1. Schematic illustration of the gliding machinery. (*A*) Magnified illustration of

a unit. The single unit consists of an internal structure (upper blue) and three huge

proteins: Gli123 (purple), Gli349 (red), and Gli521 (green) on the cell surface. The

force generated by the internal structure based on ATP hydrolysis is transmitted through

Gli521 and pulls Gli349 repeatedly. Gli349 catches and pulls SOs on the solid surface;

the unit force was estimated to be approximately 1.5 pN in the present study (see Fig.

6). (*B*) Approximately 75 legs (red) projecting from the cell can work simultaneously.

The cell glides in the direction of the yellow arrow. Unbound legs are not illustrated.

Fig. 2. Stall force of the various strains. (*A*) Left: Illustration and micrograph of

measurement using optical tweezers. The polystyrene bead (circle) (1 µm in diameter)

bound to the cell is trapped by a focused laser beam (yellow hourglass) and glides in the

direction of the white arrow. Black and pink crosses indicate the focal point of the

laser and the bead center, respectively. Right: Optical micrographs of the trapped cell.

The cell with the bead (large black ring with white center) was trapped at 0 s and stalled

at 80 s. (*B*) Representative traces. The line colors correspond to the bars in the inset.

Inset: Averages were normalized to the wild-type (WT) value, and are presented with

standard deviations (SDs) ($n = 50, 42, 26, 31$, and 35). **, $P < 0.01$ (the difference

from the WT was supported by the Student's *t*-test). The red broken line indicates the

stall force for the WT.

Fig. 3. Binding and gliding properties of the various strains. (*A*) The cell trajectories

are presented as a stack for 5 s, changing color from red to blue. (*B*) Averages of

binding activity (right) and gliding speed (left) were normalized to the WT value and

are presented with SDs.

Fig. 4. Effect of SL on stall force. (*A*) Representative traces at various

concentrations of SL. (*B*) Concentration dependency on SL. Averages were plotted

with SDs ($n = 50, 13, 23, 16, 16$, and 11 for 0, 0.05, 0.13, 0.25, 0.33, and 0.50 mM,

respectively).

Fig. 5.   Detection of force increments under low load and 0.25 mM SL.   (*A*)
Representative traces of force transition in the WT and *gli521* (P476R) mutants are
shown in the upper and lower panels, respectively.   Green and cyan triangles in each
panel indicate small peak positions taken by a peak-finding algorithm adapted to IGOR
Pro 6.33J.   (*B*) Distributions of peak values detected by the peak-finding algorithm
were fitted to the sum of four Gaussian curves.   The first, second, third, and fourth tops
of the Gaussian curves were 1.1, 2.0, 3.2, and 4.4 pN in the WT, and 1.0, 1.8, 2.6, and
3.5 pN in the *gli521* (P476R) mutant, respectively (n = 976 and 1067).

Fig. 6.   Detection of force increments under low load and 0.50 mM SL.   (*A*) Two
representative time courses of force generation.   The trajectories in the dashed
rectangular areas are magnified as insets and marked as green and blue lines for
increments and dwell times.   (*B*) Representative time course of continuous stepwise
trajectory.   The histogram of PDF analysis for indicated steps is shown in the right
inset.   (*C*) The histogram of force increments for 46 steps or peaks.   The averaged
value is indicated by a black triangle.

Fig. 7.   Leg behavior during stall.   During pre-stall, the legs colored red repeatedly
catch, pull, drag, and detach the SOs on the glass surface, thereby enabling cell
propulsion.   During stall, the force-transmitting legs are in equilibrium between
attached and detached states.   The yellow and black arrows indicate the gliding
direction and the pulling force generated by the leg strokes, respectively.

Fig. S1.   Distributions of gliding speed of the various strains over 10 s.   Experimental
data were fitted to the sum of two Gaussian curves colored blue.

Movie S1.   Stall force of gliding cell carrying a bead measured using optical tweezers.

Movie S2.   Stepwise movement of gliding cell carrying a bead detected with a weak
trap.

Fig. 1. (Single column)

Fig. 2. (Single column)

Fig. 3. (Double column)

A WT | gli521 (P476R) | m6 | m14 | m27

m29 | m34

10 μm

B
Binding activity (%) — WT, gli521 (P476R), m6, m14, m27, m29, m34

Gliding speed (%) — WT, gli521 (P476R), m6, m14, m27, m29, m34

Fig. 4. (Single column)

Fig. 5. (Single column)

Fig. 6. (Double column)

Fig. 7. (Single column)

Fig. S1.

TABLE S1 Whole genome sequence of various strains

Strain	Nucleotide change[a]	Amino acid change[b]	Previously reported	Mutation type	ORF	Annotation[c]
WT	C232735A	S354I		missense	MMOB1700	Hypothetical
gli521 (P476R)	C149908G	P476R	P476R	missense	MMOB1040	Gli521
	C232735A	S354I		missense	MMOB1700	Hypothetical
	399903T399904	M8−36*		frame shift	MMOB3220	MvspB
m6	C232735A	S354I		missense	MMOB1700	Hypothetical
	G308962A	G350R		missense	MMOB2540	SecY preprotein translocase subunit
	C423006-	E225−230*		frame shift	MMOB3340	MvspI
	G676032C	P191R		missense	MMOB5400	FtsH
m12	C136968T	Q523*	Q523*	nonsense	MMOB1020	Gli123
	T214208-	F672−695*		frame shift	MMOB1570	Oligopeptide ABC transporter binding protein
	C232735A	S354I		missense	MMOB1700	Hypothetical
	A232903T	I298N		missense	MMOB1700	Hypothetical
	C424498A	H195N		missense	MMOB3350	PolC truncated DNA polymerase III
m13	C142646T	Q1257*	Q1257*	nonsense	MMOB1030	Gli349
	C232735A	S354I		missense	MMOB1700	Hypothetical
m14	C232735A	S354I		missense	MMOB1700	Hypothetical
	233458T233459	I113−142*		frame shift	MMOB1700	Hypothetical
	C272657A	P156Q		missense	MMOB2030	Glucokinase
m23	C147186T	S2770L	S2770L	missense	MMOB1030	Gli349
	T214208-	F672−695*		frame shift	MMOB1570	Oligopeptide ABC transporter binding protein
	C232735A	S354I		missense	MMOB1700	Hypothetical
	A233098C	I233S		missense	MMOB1700	Hypothetical
	A345995T	N18Y		missense	MMOB2830	HsdR Type I restriction enzyme r protein
m27	A73556G	L101L		silent	MMOB0550	Frr ribosome recycling factor
	C152863T	P1461L		missense	MMOB1040	Gli521
	C232735A	S354I		missense	MMOB1700	Hypothetical
m29	C232735A	S354I		missense	MMOB1700	Hypothetical
	A587414T	K29N		missense	MMOB4660	TruB tRNA pseudouridine synthase B
	C654504T	W82*		nonsense	MMOB5190	Hypothetical
	A754250-	I126*		frame shift	MMOB6120	Hypothetical
m34	C232735A	S354I		missense	MMOB1700	Hypothetical
	A232753G	I348T		missense	MMOB1700	Hypothetical
	T365326A	D304V		missense	MMOB2960	AtpD F_oF_1 ATPase beta subunit

[a] Numbering is as in the sequence under accession number NC_006908.
[b] Asterisks indicate stop codons.
[c] Annotations are based on Molligen 3.0.