# CA1 and CA3 differentially support spontaneous retrieval of episodic contexts within human hippocampal subfields

Halle R. Dimsdale-Zucker, Maureen Ritchey, Arne D. Ekstrom, Andrew P. Yonelinas, Charan Ranganath

## Abstract

The hippocampus plays a critical role in supporting spatial and episodic memory. Mechanistic models predict that the hippocampal subfields have computational specializations that support memory in different ways. However, there is little empirical evidence to suggest substantial differences between the subfields, particularly in humans. To clarify how hippocampal subfields support human spatial and episodic memory, we used multivariate analyses of high-resolution functional magnetic resonance imaging (fMRI) data from a novel virtual reality paradigm. Multi-voxel pattern similarity analyses revealed that CA1 represented items that shared an episodic context as more similar than those from different episodic contexts. CA23DG showed the opposite pattern, leading to a subfield-by-condition interaction. The complementary characteristics of these subfields explain how we can parse our experiences into cohesive episodes while retaining the specific details that support vivid recollection.

## Main text

Considerable evidence suggests that the hippocampus is essential for episodic memory and plays a particular role in binding information about items and the context in which they were encountered (Davachi, 2006; Eichenbaum, Yonelinas, & Ranganath, 2007; Knierim, Lee, & Hargreaves, 2006). Most mechanistic models suggest that the hippocampal subfields play complementary roles in the representation of episodic context (Kesner & Rolls, 2015). Although these models would lead to the expectation of

large differences between coding in CA1 and CA3, between-subfield differences at the level of single-units in rodents and in overall activity in human fMRI studies have been relatively modest. Indeed, both CA1 and CA3 have been implicated in representations of temporal (Kraus, Robinson, White, Eichenbaum, & Hasselmo, 2013; Salz et al., 2016) and spatial (Kyle, Smuda, Hassan, & Ekstrom, 2015; Lee, Jerman, & Kesner, 2005) contextual information. Leutgeb and Leutgeb (2007) argued that the different computations supported by CA3 and CA1 should be most apparent when one analyzes the population-level activity patterns elicited by different contexts—whereas CA3 should differentiate between specific experiences in the same context, CA1 should globally differentiate between different contexts.

Here, we used high-resolution fMRI and multivariate analysis methods to test how different hippocampal subfields contribute to representations of spatial and episodic context. We designed a virtual reality environment consisting of two houses (spatial contexts; Figure 1). After becoming familiarized with the spatial layouts of each house, participants viewed a series of 20 videos (episodic contexts) depicting first-person navigation through each house while they encountered a series of objects. Each object was studied only once; it was uniquely placed in a single house that was shown in a single video. We scanned participants while they performed an item recognition test that required them to differentiate between studied and novel objects. Although the items were displayed without any contextual information, based on cognitive models of recognition memory and models of human hippocampal function, we predicted that recollection-based item recognition should trigger reactivation of information about the context in which that item was encountered (Davachi, 2006; Eichenbaum et al., 2007).

Accordingly, we tested whether multi-voxel patterns elicited during item recognition carried information about the associated spatial (house) or episodic (same house and video) context.

Figure 1. Experimental Approach. During encoding, participants studied objects uniquely located within one of two spatial locations (Spatial Contexts) across a series of 20 videos (Episodic Contexts). At object recognition (scanned), participants made a memory judgment (remember/familiar/new) to old and new objects presented without any contextual information. We used representational similarity analyses to compare the voxel patterns for each object to other objects that had been studied in the same (or different) house or video.

Recognition data collected in the scanner indicated high rates of recollection for studied items (mean "Remember" hit rate = 0.68 [SD = 0.17]; Supplemental Table 1). These subjective ratings were corroborated by high hit rates on the post-scan context

memory test (mean context memory hit rate = 0.71 [SD = 0.11]; Supplemental Results). Consistent with previous studies, univariate fMRI analyses revealed that hippocampus was more active for correctly recollected items than familiar or missed items (Supplemental Figure 2A). Having established hippocampal recruitment for item recollection, we proceeded to investigate whether hippocampal activity patterns carried information related to spontaneous retrieval of spatial and episodic contexts for these recollected trials.

As shown in Figure 1, we estimated single-trial multi-voxel patterns within regions of interest (ROIs) corresponding to CA1 and a combined CA2/CA3/dentate gyrus (CA23DG) subregion within the body of hippocampus. Specifically, we computed voxel pattern similarity (PS) between trial pairs for recollected items that shared the *same episodic context* (i.e., same-video/same-house), shared the *same spatial context* but different episodic contexts (different-video/same-house), or were associated with *different episodic and spatial contexts* (different-video/different-house). To maximize the likelihood of identifying trials that were associated with successful context retrieval, we restricted analyses to trials that were associated with correct recollection-based item recognition *and* correct identification of spatial context (house) in the post-scan context memory test.

To test whether regions carried information about an item's encoding context, we fitted a mixed model with a random effect of subject (Gordon, Rissman, Kiani, & Wagner, 2014) testing for effects of ROI (CA1, CA23DG), Context Similarity (same episodic, same spatial, different context), and Hemisphere (left, right), as well as their interactions on PS values. There was a significant ROI x Context Similarity x Hemisphere interaction

($\chi^2(2)$ = 13.30, $p$ = 0.001). Follow up analyses revealed that this was driven by a reliable

interaction between ROI and Context Similarity in left ($\chi^2(2)$ = 15.64, $p < .001$) but not

right ($\chi^2(2)$ = 1.65, $p$ = 0.44) hemisphere. This finding indicates that left CA1 and

CA23DG were differentially sensitive to Context Similarity. To further break down this

interaction, we conducted separate analyses restricted to left hemisphere in our ROIs to

assess representation of Context Similarity.

To investigate whether regions carried information about an item's spatial

encoding context, we compared PS values for items that had been studied in the same

house or different house. In the same house condition, we eliminated trial pairs that had

been studied within the same video to ensure that any observed effects could uniquely

be attributed to Spatial Context Similarity and not Episodic Context Similarity. Based on

traditional models (Kesner & Rolls, 2015; Marr, 1971), we expected to see greater PS

for same-house as compared to different-house pairs in CA23DG but not in CA1. As

can be seen in Figure 2, neither CA23DG ($\chi^2(1)$ = 0.08, $p$ = 0.78) nor CA1 ($\chi^2(1)$ = 0.03,

$p$ = 0.86) systematically differed in their representation based on an item's spatial

context. These results indicate that neither CA1 nor CA23DG were differentially

sensitive to Spatial Context Similarity alone.

Figure 2. Neural pattern similarity for spatial and episodic contexts. Pattern similarity was higher in left CA1 for items studied in the same episode (Same Video Same House) than for items in different episodes (Different Video Same House). Left CA23DG showed a reversal of this pattern such that pattern similarity was higher for items studied between episodes versus within the same episode. Neither CA1 nor CA23DG patterns were sensitive to spatial context similarity.

To investigate whether activity patterns in hippocampal subfields carried information about episodic context, we compared PS values for items that were studied in the same video (which necessarily meant that that the items had also been studied in the same spatial context) against items that were studied in different videos that depicted the same spatial context. Some models (Kesner & Rolls, 2015; Leutgeb & Leutgeb, 2007) predict that CA1 should treat items from the same episodic context as more similar to one another than CA23DG. Indeed, in CA1, activity patterns were more

similar across pairs of items from the same episodic context than across pairs from

different episodes ($\chi^2$(1) = 6.50, $p$ = 0.01). Intriguingly, CA23DG showed the reverse

pattern; that is, PS was significantly *lower* for items in the same episode than for items

in different episodes ($\chi^2$(1) = 10, $p$ = 0.002). We observed a significant interaction,

($\chi^2$(1) = 15.45, $p$ < .001; Figure 2), indicating that the PS profiles of CA1 and CA23DG

were qualitatively different for Episodic Context Similarity. Control analyses that

matched the number of trials across conditions revealed consistent results, ruling out

the possibility that this effect can be explained by differing numbers of trial pairs in each

condition.

Our results reveal striking differences in retrieval of contextual information across

the hippocampal subfields and provide a rare statistical dissociation between CA1 and

CA23DG. In CA1, PS was *higher* when recollecting items encountered in the same

episodic context (same video) than for items associated with different episodic contexts

(different videos); in CA23DG, PS was *lower* when recollecting items encountered in the

same episodic context than when recollecting items from different episodic contexts.

These results are consistent with the idea that CA1 represents global contextual

regularities across items ("pattern completion"), whereas CA23DG exaggerates

differences between items that have competing associations within the same episodic

context ("pattern separation"). Together, CA1 and CA23DG can play complementary

roles in supporting episodic memory by allowing one to remember specific items, as

well as their relationships, within a shared context.

Differences between the subfields are a prominent component of models of

subfield function (Kesner & Rolls, 2015; Leutgeb & Leutgeb, 2007), which, in turn, are

based on anatomical differences in the inputs, connections, and firing properties of the subfields. Studies in rodents have reported different effects of lesions (Farovik, Dupont, & Eichenbaum, 2010; Ji & Maren, 2008) and in population-level coding (Gusev, Cui, Alkon, & Gubin, 2005; Lee, Rao, & Knierim, 2004; Leutgeb & Leutgeb, 2007) between CA3 and CA1 for representations of both spatial and non-spatial contextual information. However, we are not aware of any prior findings in humans that have reported dissociations between the roles of CA1 and CA23DG during spontaneous retrieval of contextual information.

One previous human fMRI study reported a statistical dissociation between CA1 and CA23DG as participants monitored the spatial layouts of cities that varied in similarity (Stokes, Kyle, & Ekstrom, 2015). Stokes et al. found that CA23DG represented cities with the same layout as more similar than cities with different layouts whereas CA1 was not sensitive to the change in layouts (Stokes et al., 2015). One critical difference between their study and ours is that Stokes et al. (2015) directly assessed memory for spatial layouts whereas we assessed incidental retrieval of contextual information during item recognition. Our results challenge traditional notions that CA3 is particularly sensitive to changes in spatial contexts (Leutgeb & Leutgeb, 2007; Stokes et al., 2015) in that we did not see a difference in pattern similarity for items that were studied within the same spatial context or between spatial contexts.

The present results converge with other findings (Kyle, Smuda, et al., 2015) in showing that, when spatial information alone is insufficient to resolve context, spatial information does not drive representations within CA3. Our findings also accord with

Leutgeb and Leutgeb's (2007) proposal that, through rate remapping, CA3 can differentiate between different sensory cues that are encountered in the same place.

Our findings dovetail with extant evidence that CA1 plays a critical role in representing time (Kraus et al., 2013), in representing sequences of ordered information (Allen, Salz, McKenzie, & Fortin, 2016; Farovik et al., 2010), and in using this temporal information to define episodes (Wang & Diana, 2016) under demands that mimic a realistic, real-world episodic context. Additionally, our findings support the idea that CA1 is critical for distinguishing similar temporal contexts (Kesner & Rolls, 2015), given that we saw greater pattern similarity in CA1 for items within the same episode as compared with those that occurred between episodes.

The fact that CA23DG showed lower neural similarity for items encountered during the same episode as compared to items encountered during different episodes seems at odds with theories proposing a critical role for CA3 in episodic memory retrieval (Kesner & Rolls, 2015). Examined more closely, however, the results align with recent findings indicating that the hippocampus differentiates between related information in an episode (Chanales, Oza, Favila, & Kuhl, 2017; Favila, Chanales, & Kuhl, 2016; LaRocque et al., 2013; Schlichting, Mumford, & Preston, 2015). For instance, building on the theory of Marr (1971), it has been argued that dentate gyrus and CA3 work together to distinguish related information (i.e., "pattern separation"). Our finding that CA23DG is more likely to individuate items within an episode—resulting in lower neural similarity—is consistent with the idea that CA3 pushes apart representations of similar items that were encountered in the same episode (see also

(Kim, Norman, & Turk-Browne, 2017; Kyle, Stokes, Lieberman, Hassan, & Ekstrom, 2015; Schapiro, Kustner, & Turk-Browne, 2012) for related findings).

Another possibility is that the coding scheme of CA3 may be flexible based on the elements that have priority in the task (Guzowski, Knierim, & Moser, 2004). Successful performance on our task requires one to differentiate between representations of items encountered in the same video. However, if we could construct a task in which there were lower demands to orthogonalize item-specific features, we might expect CA3 to show a representational scheme more consistent with pattern completion. That is, we would expect CA3 to show increased similarity for items in the same context relative to items encountered in different contexts. Emerging evidence suggests that CA3 can represent commonalities between mental states (Aly & Turk-Browne, 2016), but it remains to be seen whether, under task conditions emphasizing similarities among items, what coding scheme CA3 would adopt.

# References

Allen, T. A., Salz, D. M., McKenzie, S., & Fortin, N. J. (2016). Nonspatial Sequence Coding in CA1

Neurons. *The Journal of Neuroscience*, *36*(5), 1547–1563.

https://doi.org/10.1523/JNEUROSCI.2874-15.2016

Aly, M., & Turk-Browne, N. B. (2016). Attention promotes episodic encoding by stabilizing

hippocampal representations. *Proceedings of the National Academy of Sciences*,

201518931. https://doi.org/10.1073/pnas.1518931113

Chanales, A. J., Oza, A., Favila, S. E., & Kuhl, B. A. (2017). Overlap among spatial memories

triggers divergence of hippocampal representations. *bioRxiv*, 99226.

https://doi.org/10.1101/099226

Davachi, L. (2006). Item, context and relational episodic encoding in humans. *Current Opinion in*

*Neurobiology*, *16*(6), 693–700. https://doi.org/10.1016/j.conb.2006.10.012

Eichenbaum, H., Yonelinas, A. P., & Ranganath, C. (2007). The medial temporal lobe and

recognition memory. *Annual Review of Neuroscience*, *30*, 123–152.

https://doi.org/10.1146/annurev.neuro.30.051606.094328

Farovik, A., Dupont, L. M., & Eichenbaum, H. (2010). Distinct roles for dorsal CA3 and CA1 in

memory for sequential nonspatial events. *Learning & Memory (Cold Spring Harbor, N.Y.)*,

*17*(1), 12–17. https://doi.org/10.1101/lm.1616209

Favila, S. E., Chanales, A. J. H., & Kuhl, B. A. (2016). Experience-dependent hippocampal pattern

differentiation prevents interference during subsequent learning. *Nature*

*Communications*, *7*, 11066. https://doi.org/10.1038/ncomms11066

Gordon, A. M., Rissman, J., Kiani, R., & Wagner, A. D. (2014). Cortical Reinstatement Mediates

the Relationship Between Content-Specific Encoding Activity and Subsequent

Recollection Decisions. *Cerebral Cortex*, *24*(12), 3350–3364.

https://doi.org/10.1093/cercor/bht194

Gusev, P. A., Cui, C., Alkon, D. L., & Gubin, A. N. (2005). Topography of Arc/Arg3.1 mRNA

expression in the dorsal and ventral hippocampus induced by recent and remote spatial

memory recall: dissociation of CA3 and CA1 activation. *The Journal of Neuroscience: The

Official Journal of the Society for Neuroscience*, *25*(41), 9384–9397.

https://doi.org/10.1523/JNEUROSCI.0832-05.2005

Guzowski, J. F., Knierim, J. J., & Moser, E. I. (2004). Ensemble Dynamics of Hippocampal Regions

CA3 and CA1. *Neuron*, *44*(4), 581–584. https://doi.org/10.1016/j.neuron.2004.11.003

Ji, J., & Maren, S. (2008). Differential roles for hippocampal areas CA1 and CA3 in the contextual

encoding and retrieval of extinguished fear. *Learning & Memory (Cold Spring Harbor,

N.Y.)*, *15*(4), 244–251. https://doi.org/10.1101/lm.794808

Kesner, R. P., & Rolls, E. T. (2015). A computational theory of hippocampal function, and tests of

the theory: new developments. *Neuroscience and Biobehavioral Reviews*, *48*, 92–147.

https://doi.org/10.1016/j.neubiorev.2014.11.009

Kim, G., Norman, K. A., & Turk-Browne, N. B. (2017). Neural differentiation of incorrectly

predicted memories. *Journal of Neuroscience*, 3272–16.

https://doi.org/10.1523/JNEUROSCI.3272-16.2017

Knierim, J. J., Lee, I., & Hargreaves, E. L. (2006). Hippocampal place cells: Parallel input streams,

    subregional processing, and implications for episodic memory. *Hippocampus*, *16*(9),

    755–764. https://doi.org/10.1002/hipo.20203

Kraus, B. J., Robinson, R. J., White, J. A., Eichenbaum, H., & Hasselmo, M. E. (2013).

    Hippocampal "time cells": time versus path integration. *Neuron*, *78*(6), 1090–1101.

    https://doi.org/10.1016/j.neuron.2013.04.015

Kyle, C. T., Smuda, D. N., Hassan, A. S., & Ekstrom, A. D. (2015). Roles of human hippocampal

    subfields in retrieval of spatial and temporal context. *Behavioural Brain Research*, *278C*,

    549–558. https://doi.org/10.1016/j.bbr.2014.10.034

Kyle, C. T., Stokes, J. D., Lieberman, J. S., Hassan, A. S., & Ekstrom, A. D. (2015). Successful

    retrieval of competing spatial environments in humans involves hippocampal pattern

    separation mechanisms. *eLife*, e10499. https://doi.org/10.7554/eLife.10499

LaRocque, K. F., Smith, M. E., Carr, V. A., Witthoft, N., Grill-Spector, K., & Wagner, A. D. (2013).

    Global similarity and pattern separation in the human medial temporal lobe predict

    subsequent memory. *The Journal of Neuroscience: The Official Journal of the Society for*

    *Neuroscience*, *33*(13), 5466–5474. https://doi.org/10.1523/JNEUROSCI.4293-12.2013

Lee, I., Jerman, T. S., & Kesner, R. P. (2005). Disruption of delayed memory for a sequence of

    spatial locations following CA1- or CA3-lesions of the dorsal hippocampus. *Neurobiology*

    *of Learning and Memory*, *84*(2), 138–147. https://doi.org/10.1016/j.nlm.2005.06.002

Lee, I., Rao, G., & Knierim, J. J. (2004). A double dissociation between hippocampal subfields:

    differential time course of CA3 and CA1 place cells for processing changed environments.

    *Neuron*, *42*(5), 803–815. https://doi.org/10.1016/j.neuron.2004.05.010

Leutgeb, S., & Leutgeb, J. K. (2007). Pattern separation, pattern completion, and new neuronal

    codes within a continuous CA3 map. *Learning & Memory (Cold Spring Harbor, N.Y.)*,

    *14*(11), 745–757. https://doi.org/10.1101/lm.703907

Marr, D. (1971). Simple memory: a theory for archicortex. *Philosophical Transactions of the*

    *Royal Society of London. Series B, Biological Sciences*, *262*(841), 23–81.

Salz, D. M., Tiganj, Z., Khasnabish, S., Kohley, A., Sheehan, D., Howard, M. W., & Eichenbaum, H.

    (2016). Time Cells in Hippocampal Area CA3. *The Journal of Neuroscience*, *36*(28), 7476–

    7484. https://doi.org/10.1523/JNEUROSCI.0087-16.2016

Schapiro, A. C., Kustner, L. V., & Turk-Browne, N. B. (2012). Shaping of Object Representations

    in the Human Medial Temporal Lobe Based on Temporal Regularities. *Current Biology*,

    *22*(17), 1622–1627. https://doi.org/10.1016/j.cub.2012.06.056

Schlichting, M. L., Mumford, J. A., & Preston, A. R. (2015). Learning-related representational

    changes reveal dissociable integration and separation signatures in the hippocampus

    and prefrontal cortex. *Nature Communications*, *6*, 8151.

    https://doi.org/10.1038/ncomms9151

Stokes, J., Kyle, C., & Ekstrom, A. D. (2015). Complementary roles of human hippocampal

    subfields in differentiation and integration of spatial context. *Journal of Cognitive*

    *Neuroscience*, *27*(3), 546–559. https://doi.org/10.1162/jocn_a_00736

Wang, F., & Diana, R. A. (2016). Temporal context processing within hippocampal subfields.

    *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2016.03.048