Deep transcriptome annotation suggests that small and large proteins encoded in

the same genes often cooperate

Sondos Samandi[1,7] † , Annie V. Roy[1,7] † , Vivian Delcourt[1,7,8], Jean-François Lucier[2],

Jules Gagnon[2], Maxime C. Beaudoin[1,7], Benoît Vanderperre[1], Marc-André Breton[1], Julie

Motard[1,7], Jean-François Jacques[1,7], Mylène Brunelle[1,7], Isabelle Gagnon-Arsenault[6,7],

Isabelle Fournier[8], Aida Ouangraoua[3], Darel J. Hunting[4], Alan A. Cohen[5], Christian R.

Landry[6,7], Michelle S. Scott[1], Xavier Roucou[1,7]∗

[1]Department of Biochemistry, [2]Department of Biology and Center for Computational

Science, [3]Department of Computer Science, [4]Department of Nuclear Medicine &

Radiobiology, [5]Department of Family Medicine, Université de Sherbrooke, Québec,

Canada; [6]Département de biologie and IBIS, Université Laval, Québec, Canada;

[7]PROTEO, Québec Network for Research on Protein Function, Structure, and

Engineering, Québec, Canada; [8]Université Lille, INSERM U1192, Laboratoire

Protéomique, Réponse Inflammatoire & Spectrométrie de Masse (PRISM) F-59000 Lille,

France

†These authors contributed equally to this work

∗Correspondance to Xavier Roucou: Département de biochimie (Z8-2001), Faculté de

Médecine et des Sciences de la Santé, Université de Sherbrooke, 3201 Jean Mignault,

Sherbrooke, Québec J1E 4K8, Canada, Tel. (819) 821-8000x72240; Fax. (819) 820 6831;

E-Mail: <u>xavier.roucou@usherbrooke.ca</u>

Abstract

Recent studies in eukaryotes have demonstrated the translation of alternative open

reading frames (altORFs) in addition to annotated protein coding sequences (CDSs). We

show that a large number of small proteins could in fact be coded by altORFs. The

putative alternative proteins translated from altORFs have orthologs in many species and

evolutionary patterns indicate that altORFs are particularly constrained in CDSs that

evolve slowly. Thousands of predicted alternative proteins are detected in proteomic

datasets by reanalysis using a database containing predicted alternative proteins. Protein

domains and co-conservation analyses suggest a potential functional relationship between

small and large proteins encoded in the same genes. This is illustrated with specific

examples, including altMiD51, a 70 amino acid mitochondrial fission-promoting protein

encoded in *MiD51/Mief1/SMCR7L*, a gene encoding an annotated protein promoting

mitochondrial fission. Our results suggest that many coding genes code for more than one

protein that are often functionally related.

Introduction

Current protein databases are cornerstones of modern biology but are based on a number of assumptions. In particular, a mature mRNA is predicted to contain a single CDS; yet, ribosomes can select more than one translation initiation site (TIS)[1-3] on any single mRNA. Also, minimum size limits are imposed on the length of CDSs, resulting in many RNAs being mistakenly classified as non-coding (ncRNAs)[4-11]. As a result of these assumptions, the size and complexity of most eukaryotic proteomes have probably been greatly underestimated[12-15]. In particular, few small proteins (defined as of 100 amino acids or less) are annotated in current databases. The absence of annotation of small proteins is a major bottleneck in the study of their function and their roles in health and disease. This is further supported by classical and recent examples of small proteins of functional importance, for instance many critical regulatory molecules such as the F0 subunit of the F0F1-ATPsynthase[16], the sarcoplasmic reticulum calcium ATPase regulator phospholamban[17], and the key regulator of iron homeostasis hepcidin[18]. This limitation also impedes our understanding of the process of origin of new genes, which are thought to contribute to evolutionary innovations. Because these genes generally code for small proteins [19-22], they are difficult to unambiguously detect by proteomics and in fact are impossible to detect if they are not included in proteomics databases.

Functional annotation of ORFs encoding small proteins is particularly challenging since an unknown fraction of small ORFs may occur by chance in the transcriptome, generating a significant level of noise [13]. However, given that many small proteins have

important functions and are ultimately one of the most important sources of functional

novelty, it is time to address the challenge of their functional annotations[13].

We systematically reanalyzed several eukaryotic transcriptomes to annotate previously

unannotated ORFs which we term alternative ORFs (altORFs), and we annotated the

corresponding hidden proteome. Here, altORFs are defined as potential protein-coding

ORFs in ncRNAs, in UTRs or in different reading frames from annotated CDSs in

mRNAs (Figure 1a). For clarity, predicted proteins translated from altORFs are termed

alternative proteins and proteins translated from annotated CDSs are termed reference

proteins.

Our goal was to provide functional annotations of alternative proteins by (1) analyzing

relative patterns of evolutionary conservation between alternative and reference proteins

and their corresponding coding sequences; (2) estimating the prevalence of alternative

proteins both by bioinformatics analysis and by detection in large experimental datasets;

(3) detecting functional signatures in alternative proteins; and (4) predicting and testing

functional cooperation between alternative and reference proteins.

Results

Prediction of altORFs and alternative proteins. We predicted a total of 539,134

altORFs compared to 68,264 annotated CDSs in the human transcriptome (Figure 1b,

Table 1). Because identical ORFs can be present in different RNA isoforms transcribed

from the same genomic locus, the number of unique altORFs and CDSs becomes 183,191

and 54,498, respectively. AltORFs were also predicted in other organisms for comparison

(Table 1). By convention, only reference proteins are annotated in current protein

databases. As expected, altORFs are on average small, with a size ranging from 30 to

1480 codons. Accordingly, the median size of predicted human alternative proteins is 45

amino acids compared to 460 for reference proteins (Figure 1c), and 92.96 % of

alternative proteins have less than 100 amino acids. Thus, the bulk of the translation

products of altORFs would be small proteins. The majority of altORFs either overlap

annotated CDSs in a different reading frame (35.98%) or are located in 3'UTRs (40.09%)

(Figure 1d). 9.83% of altORFs are located in repeat sequences (Figure 1-figure

supplement 1a), compared to 2.45% of CDSs. To assess whether observed altORFs could

be attributable solely to random occurrence, due for instance to the base composition of

the transcriptome, we estimated the expected number of altORFs generated in 100

shuffled human transcriptomes. Overall, we observed 62,307 more altORFs than would

be expected from random occurrence alone (Figure 1e; $p < 0.0001$). This analysis suggests

that a large number are expected by chance alone but that at the same time, a large

absolute number could potentially be maintained and be functional. The density of

altORFs observed in the CDSs, 3'UTRs and ncRNAs (Figure 1f) was markedly higher

than in the shuffled transcriptomes, suggesting that these are maintained at frequencies

higher than expected by chance, again potentially due to their coding function. In

contrast, the density of altORFs observed in 5'UTRs was much lower than in the shuffled

transcriptomes, supporting recent claims that negative selection eliminates AUGs (and

thus the potential for the evolution of altORFs) in these regions[23,24].

Although the majority of human annotated CDSs do not have a TIS with a Kozak motif

(Figure 1g)[25], there is a correlation between a Kozak motif and translation efficiency[26].

We find that 27,539 (15% of 183,191) human altORFs encoding predicted alternative

proteins have a Kozak motif (A/GNNAUGG), as compared to 19,745 (36% of 54,498)

for annotated CDSs encoding reference proteins (Figure 1g). The number of altORFs

with Kozak motifs is significantly higher in the human transcriptome compared to

shuffled transcriptomes (Figure 1-figure supplement 2), again supporting their potential

role as protein coding.

Conservation analyses. Next, we compared evolutionary conservation patterns of

altORFs and CDSs. A large number of human alternative proteins have homologs in other

species. In mammals, the number of homologous alternative proteins is higher than the

number of homologous reference proteins (Figure 2a), and 9 are even conserved from

human to yeast (Figure 2b), supporting a potential functional role. As phylogenetic

distance from human increases, the number and percentage of genes encoding

homologous alternative proteins decreases more rapidly than the percentage of genes

encoding reference proteins (Figures 2a and c). This observation indicates either that

altORFs evolve more rapidly than CDSs or that distant homologies are less likely to be

detected given the smaller sizes of alternative proteins. Another possibility is that they

evolve following the patterns of evolution of genes that evolve *de novo*, with a rapid birth

and death rate, which accelerates their turnover over time[20].

Since the same gene may contain a conserved CDS and one or several conserved

altORFs, we analyzed the co-conservation of orthologous altORF-CDS pairs. Our results

show a very large fraction of co-conserved alternative-reference protein pairs in several species

(Figure 3). Detailed results for all species are presented in Table 2.

If altORFs play a functional role, they would be expected to be under purifying selection.

The first and second positions of a codon experience stronger purifying selection than the

third because of redundancy in the genetic code[27]. In the case of CDS regions

overlapping altORFs with a shifted reading frame, the third codon positions of the CDSs

are either the first or the second in the altORFs, and should thus also undergo purifying

selection. We analyzed conservation of third codon positions of CDSs for 100 vertebrate

species for 1,088 altORFs completely nested within and co-conserved across vertebrates

(human to zebrafish) with their 889 CDSs from 867 genes (Figure 4). We observed that

in regions of the CDS overlapping altORFs, third codon positions were evolving at

significantly more extreme speeds (slow or quick) than third codon positions of random

control sequences from the entire CDS (Figure 4), reaching up to 67-fold for conservation

at $p<0.0001$ and 124-fold for accelerated evolution at $p<0.0001$. We repeated this

analysis with the 53,862 altORFs completely nested within the 20,814 CDSs from 14,677

genes, independently of their co-co-conservation. We observed a similar trend, with a 22-

fold for conservation at $p<0.0001$, and a 24-fold for accelerated evolution at $p<0.0001$

(Figure 4-figure supplement 1).This is illustrated with three altORFs located within the

CDS of *NTNG1*, *RET* and *VTI1A* genes (Figure 5). These three genes encode a protein

promoting neurite outgrowth, the proto-oncogene tyrosine-protein kinase receptor Ret

and a protein mediating vesicle transport to the cell surface, respectively. Two of these

alternative proteins have been detected by ribosome profiling (*RET*, IP_182668.1) or

mass spectrometry (*VTI1A*, IP_188229.1) (see below, Supplementary files 1 and 2).

Evidence of expression of alternative proteins. We provide two lines of evidence

indicating that thousands of altORFs are translated into proteins. First, we re-analyzed

detected TISs in publicly available ribosome profiling data[28,29], and found 26,531 TISs

mapping to annotated CDSs and 12,616 mapping to altORFs in these studies (Figure 6a;

Supplementary file 1). Only a small fraction of TISs detected by ribosomal profiling

mapped to altORFs[3'] even if those are more abundant than altORF[5'] relative to shuffled

transcriptomes, likely reflecting a recently-resolved technical issue which prevented TIS

detection in 3'UTRs [30]. New methods to analyze ribosome profiling data are being

developed and will likely uncover more translated altORFs [9]. In agreement with the

presence of functional altORFs[3'], cap-independent translational sequences were recently

discovered in human 3'UTRs[31]. Second, we re-analyzed proteomic data using our

composite database containing alternative proteins in addition to annotated reference

proteins (Figure 6b; Supplementary file 2). We selected four studies representing

different experimental paradigms and proteomic applications: large-scale [32] and targeted

[33] protein/protein interactions, post-translational modifications [34], and a combination of

bottom-up, shotgun and interactome proteomics [35] . In the first dataset, we detected 3,957

predicted alternative proteins in the interactome of reference proteins[32], providing a

framework to uncover the function of these proteins. In a second proteomic dataset

containing about 10,000 reference human proteins[35], a total of 549 predicted alternative

proteins were detected. Using a phosphoproteomic large data set[34], we detected 384

alternative proteins. The biological function of these proteins is supported by the

observation that some alternative proteins are specifically phosphorylated in cells

stimulated by the epidermal growth factor, and others are specifically phosphorylated

during mitosis (Figure 7; Supplementary file 3). We provide examples of spectra

validation (Figure 7-figure supplement 1). A fourth proteomic dataset contained 77

alternative proteins in the epidermal growth factor receptor interactome[33] (Figure 6b). A

total of 4,872 different alternative proteins were detected in these proteomic data. The

majority of these proteins are coded by altORF^{CDS}, but there are also significant

contributions of altORF$^{3'}$, altORF^{nc} and altORF$^{5'}$ (Figure 6c). Overall, by mining the

proteomic and ribosomal profiling data, we detected the translation of a total of 17,371

unique alternative proteins. 467 of these alternative proteins were detected by both MS

and ribosome profiling (Figure 8), providing a high-confidence collection of small

alternative proteins for further studies.

Functional annotations of alternative proteins. An important goal of this study is to

associate potential functions to alternative proteins, which we can do through

annotations. Because the sequence similarities and the presence of particular signatures

(families, domains, motifs, sites) are a good indicator of a protein's function, we analyzed

the sequence of the predicted alternative proteins in several organisms with InterProScan,

an analysis and classification tool for characterizing unknown protein sequences by

predicting the presence of combined protein signatures from most main domain

databases[36] (Figure 9; Figure 9-figure supplement 1). We found 41,511 (23%) human

alternative proteins with at least one InterPro signature (Figure 9b). Of these, 37,739 (or

20.6%) are classified as small proteins. Interestingly, the reference proteome has a

smaller proportion (840 or 1.68%) of small proteins with at least one InterPro signature,

supporting a biological activity for alternative proteins.

Similar to reference proteins, signatures linked to membrane proteins are abundant in the

alternative proteome and represent more than 15,000 proteins (Figures 9c-e; Figure 9-

figure supplement 1). With respect to the targeting of proteins to the secretory pathway or

to cellular membranes, the main difference between the alternative and the reference

proteomes lies in the very low number of proteins with both signal peptides and

transmembrane domains. Most of the alternative proteins with a signal peptide do not

have a transmembrane segment and are predicted to be secreted (Figures 9c, d),

supporting the presence of large numbers of alternative proteins in plasma[37]. The majority

of predicted alternative proteins with transmembrane domains have a single membrane

spanning domain but some display up to 27 transmembrane regions, which is still within

the range of reference proteins that show a maximum of 33 (Figure 9e).

We extended the functional annotation using the Gene Ontology. A total of 585

alternative proteins were assigned 419 different InterPro entries, and 343 of them were

tentatively assigned 192 gene ontology terms (Figure 10). 15.5% (91/585) of alternative

proteins with an InterPro entry were detected by MS or/and ribosome profiling, compared

to 13.7% (22,055/161,110) for alternative proteins without an InterPro entry (p-value =

1.13e-05, Fisher's exact test and chi-square test). Thus, predicted alternative proteins with

InterPro entries are more likely to be detected, supporting their functional role. The most

abundant class of predicted alternative proteins with at least one InterPro entry are C2H2

zinc finger proteins with 110 alternative proteins containing 187 C2H2-type/integrase

DNA-binding domains, 91 C2H2 domains and 23 C2H2-like domains (Figure 11a).

Eighteen of these (17.8%) were detected in public proteomic and ribosome profiling

datasets, a percentage that is similar to reference zinc finger proteins (20.1%) (Figure 6,

Table 3). Alternative proteins have between 1 and 23 zinc finger domains (Figure 11b).

Zinc fingers mediate protein-DNA, protein-RNA and protein-protein interactions[38]. The

linker sequence separating adjacent finger motifs matches or resembles the consensus

TGEK sequence in nearly half the annotated zinc finger proteins[39]. This linker confers

high affinity DNA binding and switches from a flexible to a rigid conformation to

stabilize DNA binding. The consensus TGEK linker is present 46 times in 31 alternative

zinc finger proteins (Supplementary file 4). These analyses show that a number of

alternative proteins can be classified into families and will help deciphering their

functions.

Evidence of functional relationships between reference and alternative proteins

coded by the same genes. Since one gene may code for both a reference and one or

several alternative proteins, we asked whether paired (encoded in the same gene)

alternative and reference proteins have functional relationships. The functional

associations discussed here are potential functional interactions that do not necessarily

imply physical interactions; however, there are a few known examples of functional

linkage between different proteins encoded in the same gene (Table 4). If there is a

functional relationship, one would expect orthologous alternative- reference protein pairs

to be co-conserved more often than expected by chance[40]. Our results show a large

fraction of co-conserved alternative-reference protein pairs in several species (Figure 3).

Detailed results for all species are presented in Table 2.

Another mechanism that could show functional relationships between alternative and

reference proteins encoded in the same gene would be that they share protein domains.

We compared the functional annotations of the 585 alternative proteins with an InterPro

entry with the reference proteins expressed from the same genes. Strikingly, 89 of 110

altORFs coding for zinc finger proteins (Figure 11) are present in transcripts in which the

CDS also codes for a zinc finger protein. Overall, 138 alternative/reference protein pairs

share at least one InterPro entry and many pairs share more than one entry (Figure 12a).

The number of shared entries was much higher than expected by chance (Figure 12b,

$p<0.0001$). The correspondence between InterPro domains of alternative proteins and

their corresponding reference proteins coded by the same genes also indicates that even

when entries are not identical, the InterPro terms are functionally related (Figure 12c;

Figure 12-figure supplement 1), overall supporting a potential functional linkage between

reference and predicted alternative proteins. Domain sharing remains significant

($p<0.001$) even when the most frequent domains, zinc fingers, are not considered (Figure

12-figure supplement 2).

Recently, the interactome of 118 human zinc finger proteins was determined by affinity

purification followed by mass spectrometry [41]. This study provides a unique opportunity

to test if, in addition to possessing zinc finger domains, thus being functionally connected

some pairs of reference and alternative proteins coded by the same gene also interact. We

re-analyzed the MS data using our alternative protein sequence database to detect

alternative proteins in this interactome (Supplementary file 5). Five alternative proteins

(IP_168460.1, IP_168527.1, IP_270697.1, IP_273983.1, IP_279784.1) were identified

within the interactome of their reference zinc finger proteins. This number was higher

than expected by chance ($p<10^{-6}$) based on 1 million binomial simulations of randomized

interactomes. This result strongly supports the hypothesis of functional relationships

between alternative and reference proteins coded by the same genes, and indicates that

there are examples of physical interactions.

Finally, we integrated the co-conservation and expression analyses to produce a high-

confidence list of alternative proteins predicted to have a functional relationship with

their reference proteins and found 2,715 alternative proteins in mammals (*H. sapiens* to

B. taurus), and 44 in vertebrates (H. *sapiens* to *D. rerio*) (Supplementary file 6). In order

to further test for functional relationship between alternative/reference protein pairs in

this list, we focused on alternative proteins detected with at least two peptide spectrum

matches or with high TIS reads. From this subset, we selected altMiD51 (IP_294711.1)

among the top 2% of alternative proteins detected with the highest number of unique

peptides in proteomics studies, and altDDIT3 (IP_211724.1) among the top 2% of

altORFs with the most cumulative reads in translation initiation ribosome profiling

studies.

AltMiD51 is a 70 amino acid alternative protein conserved in vertebrates[42] and co-

conserved with its reference protein MiD51 from humans to zebrafish (Supplementary

file 6). Its coding sequence is present in exon 2 of the *MiD51/MIEF1/SMCR7L* gene. This

exon forms part of the 5'UTR for the canonical mRNA and is annotated as non-coding in

current gene databases (Figure 13a). Yet, altMiD51 is robustly detected by MS in several

cell lines (Supplementary file 2: HEK293, HeLa Kyoto, HeLa S3, THP1 cells and gut

tissue), and we validated some spectra using synthetic peptides (Figure 13-figure

supplement 1), and it is also detected by ribosome profiling (Supplementary file 1)[37,42,43].

We confirmed co-expression of altMiD51 and MiD51 from the same transcript (Figure

13b). Importantly, the tripeptide LYR motif predicted with InterProScan and located in

the N-terminal domain of altMiD51 (Figure 13a) is a signature of mitochondrial proteins

localized in the mitochondrial matrix[44]. Since *MiD51/MIEF1/SMCR7L* encodes the

mitochondrial protein MiD51, which promotes mitochondrial fission by recruiting

cytosolic Drp1, a member of the dynamin family of large GTPases, to mitochondria[45], we

tested for a possible functional connection between these two proteins expressed from the

same mRNA. We first confirmed that MiD51 induces mitochondrial fission (Figure 13-

figure supplement 2). Remarkably, we found that altMiD51 also localizes at the

mitochondria (Figure 13c; Figure 13-figure supplement 3) and that its overexpression

results in mitochondrial fission (Figure 13d). This activity is unlikely to be through

perturbation of oxidative phosphorylation since the overexpression of altMiD51 did not

change oxygen consumption nor ATP and reactive oxygen species production (Figure 13-

figure supplement 4). The decrease in spare respiratory capacity in altMiD51-expressing

cells (Figure 13-figure supplement 4a) likely resulted from mitochondrial fission[46]. The

LYR domain is essential for altMiD51-induced mitochondrial fission since a mutant of

the LYR domain, altMiD51(LYR→AAA) was unable to convert the mitochondrial

morphology from tubular to fragmented (Figure 13d). Drp1(K38A), a dominant negative

mutant of Drp1 [47], largely prevented the ability of altMiD51 to induce mitochondrial

fragmentation (Figure 13d; Figure 13-figure supplement 5a). In a control experiment, co-

expression of wild-type Drp1 and altMiD51 proteins resulted in mitochondrial

fragmentation (Figure 13-figure supplement 5b). Expression of the different constructs

used in these experiments was verified by western blot (Figure 13-figure supplement 6).

Drp1 knockdown interfered with altMiD51-induced mitochondrial fragmentation (Figure

14), confirming the proposition that Drp1 mediates altMiD51-induced mitochondrial

fragmentation. It remains possible that altMiD51 promotes mitochondrial fission

independently of Drp1 and is able to reverse the hyperfusion induced by Drp1

inactivation. However, Drp1 is the key player mediating mitochondrial fission and most

likely mediates altMiD51-induced mitochondrial fragmentation, as indicated by our

results.

AltDDIT3 is a 31 amino acid alternative protein conserved in vertebrates and co-

conserved with its reference protein DDIT3 from human to bovine (Supplementary file

6). Its coding sequence overlaps the end of exon 1 and the beginning of exon 2 of the

DDIT3/CHOP/GADD153 gene. These exons form part of the 5'UTR for the canonical

mRNA (Figure 15a). To determine the cellular localization of altDDIT3 and its possible

relationship with DDIT3, confocal microscopy analyses were performed on HeLa cells

co-transfected with altDDIT3^{GFP} and DDIT3^{mCherry}. Expression of these constructs was

verified by western blot (Figure 15-figure supplement 1). Interestingly, both proteins

were mainly localized in the nucleus and partially localized in the cytoplasm (Figure

15b). This distribution for DDIT3 confirms previous studies [48,49]. Both proteins seemed to

co-localize in these two compartments (Pearson correlation coefficient 0.92, Figure 15c).

We further confirmed the statistical significance of this colocalization by applying

Costes' automatic threshold and Costes' randomization colocalization analysis and

Manders Correlation Coefficient (Figure 15d; Figure 15-figure supplement 2) [50]. Finally,

in lysates from cells co-expressing altDDIT3^{GFP} and DDIT3^{mCherry}, DDIT3^{mCherry} was

immunoprecipitated with GFP-trap agarose, confirming an interaction between the small

altDDTI3 and the large DDIT3 proteins encoded in the same gene (Figure 15e).

Discussion

We have provided the first functional annotation of altORFs with a minimum size of 30

codons in different genomes. The comprehensive annotation of *H sapiens* altORFs is

freely available to download at https://www.roucoulab.com/p/downloads (Homo sapiens

functional annotation of alternative proteins based on RefSeq GRCh38 (hg38)

predictions). In light of the increasing evidence from approaches such as ribosome

profiling and MS-based proteomics that the one mRNA-one canonical CDS assumption

is strongly challenged, our findings provide the first clear functional insight into a new

layer of regulation in genome function. While many observed altORFs may be

evolutionary accidents with no functional role, several independent lines of evidence

support translation and a functional role for thousands of alternative proteins: (1)

overrepresentation of altORFs relative to shuffled sequences; (2) overrepresentation of

altORF Kozak sequences; (3) active altORF translation detected via ribosomal profiling;

(4) detection of thousands of alternative proteins in multiple existing proteomic datasets;

(5) correlated altORF-CDS conservation, but with overrepresentation of highly conserved

and fast-evolving altORFs; (6) overrepresentation of identical InterPro signatures

between alternative and reference proteins encoded in the same mRNAs; (7) several

thousand co-conserved paired alternative-reference proteins encoded in the same gene;

and (8) presence of clear, striking examples in altMiD51, altDDIT3 and 5 alternative

proteins interacting with their reference zinc finger proteins. While 5 of these 8 lines of

evidence support an unspecified functional altORF role, 4 of them (5, 6, 7 and 8)

independently support a specific functional/evolutionary interpretation of their role: that

alternative proteins and reference proteins have paired functions. Note that this

hypothesis does not require binding, just functional cooperation such as activity on a

shared pathway.

The presence of different coding sequences in the same gene provides a coordinated

transcriptional regulation. Consequently, the transcription of different coding sequences

can be turned on or off together, similar to prokaryotic operons. Thus, the observation

that alternative-reference protein pairs encoded in the same genes have functional

relationships is not completely unexpected. This observation is also in agreement with

increasing evidence that small proteins regulate the function of larger proteins[51]. We

speculate that clustering of a CDS and one or more altORFs in the same genes, and thus

in the same transcription unit allows cells to adapt more quickly with optimized energy

expenditure to environmental changes.

Upstream ORFs here labeled altORFs[5'] are important translational regulators of canonical

CDSs in vertebrates[52]. Interestingly, the altORF[5'] encoding altDDIT3 was characterized

as an inhibitory upstream ORF [53,54], but evidence of endogenous expression of the

corresponding small protein was not sought. The detection of altMiD51 and altDDIT3

suggests that a fraction of altORFs[5'] may have dual functions as translation regulators and

functional proteins.

Our results raise the question of the evolutionary origins of these altORFs. A first

possible mechanism involves the polymorphism of initiation and stop codons during

evolution [55,56]. For instance, the generation of an early stop codon in the 5'end of a CDS

could be followed by the evolution of another translation initiation site downstream,

creating a new independent ORF in the 3'UTR of the canonical gene. This mechanism of

altORF origin, reminiscent of gene fission, would at the same time produce a new altORF

that shares protein domains with the annotated CDS, as we observed for a substantial

fraction (24%) of the 585 alternative proteins with an InterPro entry. A second

mechanism would be de novo origin of ORFs, which would follow the well-established

models of gene evolution *de novo*[20,57,58] in which new ORFs are transcribed and

translated and have new functions or await the evolution of new functions by mutations.

The numerous altORFs with no detectable protein domains may have originated this way

from previously non-coding regions or in regions that completely overlap with CDS in

other reading frames.

Detection is an important challenge in the study of small proteins. A TIS detected by

ribosome profiling does not necessarily imply that the protein is expressed as a stable

molecule, and proteomic analyses more readily detect large proteins that generate several

peptides after enzymatic digestion. In addition, evolutionarily novel genes tend to be

poorly expressed, again reducing the probability of detection [20]. Here, we used a

combination of five search engines, thus increasing the confidence and sensitivity of hits

compared to single-search-engine processing[59,60]. This strategy led to the detection of

several thousand alternative proteins. However, ribosome profiling and MS have technical caveats and the comprehensive contribution of small proteins to the proteome will require more efforts, including the development of new tools such as specific antibodies.

Only a relatively small percentage of alternative proteins (22.6%) are functionally annotated with Interpro signatures, compared to reference proteins (96.9%). An obvious explanation is the small size of alternative proteins with a median size of 45 amino acids, which may not be able to accommodate large domains. It has been proposed that small proteins may be precursors of new proteins but require an elongation of their coding sequence before they display a useful cellular activity [19,51]. According to this hypothesis, it is possible that protein domains appear only after elongation of the coding sequence. Alternatively, InterPro domains were identified by investigating the reference proteome, and alternative proteins may have new domains and motifs that remain to be characterized. Finally, an unknown fraction of predicted altORFs may not be translated or may code for non-functional peptides.

In conclusion, our deep annotation of the transcriptome reveals that a large number of small eukaryotic proteins, which may even represent the majority, are still officially unannotated. Our results also suggest that many small and large proteins coded by the same mRNA may cooperate by regulating each other's function or by functioning in the same pathway, confirming the few examples in the literature of unrelated proteins encoded in the same genes and functionally cooperating[61–65]. To determine whether or not

this functional cooperation is a general feature of small/large protein pairs encoded in the

same gene will require much more experimental evidence, but our results strongly

support this hypothesis.

Materials and methods

Generation of alternative open reading frames (altORFs) and alternative protein

databases. Throughout this manuscript, annotated protein coding sequences and proteins

in current databases are labelled annotated coding sequences or CDSs and reference

proteins, respectively. For simplicity reasons, predicted alternative protein coding

sequences are labelled alternative open reading frames or altORFs.

To generate MySQL databases containing the sequences of all predicted alternative

proteins translated from reference annotation of different organisms, a computational

pipeline of Perl scripts was developed as previously described with some modifications[37].

Genome annotations for *H. sapiens* (release hg38, Assembly: GCF_000001405.26), *P.*

troglodytes (Pan_troglodytes-2.1.4, Assembly: GCF_000001515.6), *M. musculus*

(GRCm38.p2, Assembly: GCF_000001635.22), *D. melanogaster* (release 6, Assembly:

GCA_000705575.1), *C. elegans* (WBcel235, Assembly: GCF_000002985.6) and *S.*

cerevisiae (Sc_YJM993_v1, Assembly: GCA_000662435.1) were downloaded from the

NCBI website (http://www.ncbi.nlm.nih.gov/genome). For *B. taurus* (release UMD

3.1.86), *X. tropicalis* (release JGI_4.2) and *D. rerio* (GRCz10.84), genome annotations

were downloaded from Ensembl (http://www.ensembl.org/info/data/ftp/). Each annotated

transcript was translated *in silico* with Transeq[66]. All ORFs starting with an AUG and

ending with a stop codon different from the CDS, with a minimum length of 30 codons

(including the stop codon) and identified in a distinct reading frame compared to the

annotated CDS when overlapping the CDS, were defined as altORFs.

An additional quality control step was performed to remove initially predicted altORFs

with a high level of identity with reference proteins. Such altORFs typically start in a

different coding frame than the reference protein but through alternative splicing, end

with the same amino acid sequence as their associated reference protein. Using BLAST,

altORFs overlapping CDSs chromosomal coordinates and showing more than 80%

identity and overlap with an annotated CDS were rejected.

AltORF localization was assigned according to the position of the predicted translation

initiation site (TIS): altORFs$^{5'}$, altORFs^{CDS} and altORFs$^{3'}$ are altORFs with TISs located

in 5'UTRs, CDSs and 3'UTRs, respectively. Non-coding RNAs (ncRNAs) have no

annotated CDS and all ORFs located within ncRNAs are labelled altORFs^{nc}.

The presence of the simplified Kozak sequence (A/GNNATGG) known to be favorable

for efficient translation initiation was also assessed for each predicted altORF[67].

Identification of TISs. The global aggregates of initiating ribosome profiles data were

obtained from the initiating ribosome tracks in the GWIPS-viz genome browser[28] with

ribosome profiling data collected from five large scale studies[2,9,68–70]. Sites were mapped

to hg38 using a chain file from the UCSC genome browser

(http://hgdownload.soe.ucsc.edu/goldenPath/hg19/liftOver/hg19ToHg38.over.chain.gz)

and CrossMap v0.1.6 (RRID:SCR_001173). Similar to the methods used in these studies,

an altORF is considered as having an active TIS if it is associated with at least 10 reads at

one of the 7 nucleotide positions of the sequence NNNAUGN (AUG is the predicted

altORF TIS). An additional recent study was also included in our analysis[29]. In this study,

a threshold of 5 reads was used. Raw sequencing data for ribosome protected fragments

in harringtonine treated cells was aligned to the human genome (GRCh38) using bowtie2

$(2.2.8)$[71]. Similar to the method used in this work, altORFs with at least 5 reads

overlapping one position in the kozak region were considered as having an

experimentally validated TIS.

Generation of shuffled transcriptomes. Each annotated transcript was shuffled using

the Fisher-Yates shuffle algorithm. In CDS regions, all codons were shuffled except the

initiation and stop codons. For mRNAs, we shuffled the 5'UTRs, CDSs and 3'UTRs

independently to control for base composition. Non-coding regions were shuffled at the

nucleotide level. The resulting shuffled transcriptome has the following features

compared to hg38: same number of transcripts, same transcripts lengths, same nucleotide

composition, and same amino-acid composition for the proteins translated from the

CDSs. Shuffling was repeated 100 times and the results are presented with average values

and standard deviations. The total number of altORFs is 539,134 for hg38, and an

average of 489,073 for shuffled hg38. AltORFs and kozak motifs in the 100 shuffled

transcriptomes were detected as described above for hg38.

Identification of paralogs/orthologs in alternative proteomes. Both alternative and

reference proteomes were investigated. Pairwise ortholog and paralog relationships

between the human proteomes and the proteomes from other species, were calculated

using an InParanoid-like approach[72], as described below (RRID:SCR_006801). The

following BLAST (RRID:SCR_001010) procedure was used. Comparisons using our

datasets of altORFs/CDS protein sequences in multiple FASTA formats from

Saccharomyces cerevisiae, Caenorhabditis elegans, Drosophila melanogaster, Danio

rerio, Xenopus tropicalis Bos taurus, Mus musculus, Pan troglodytes, Homo sapiens were

performed between each pair of species (*Homo sapiens* against the other species),

involving four whole proteome runs per species pair: pairwise comparisons (organism A

vs organism B, organism B vs organism A), plus two self-self runs (organism A vs

organism A, organism B vs organism B). BLAST homology inference was accepted when

the length of the aligned region between the query and the match sequence equalled or

exceeded 50% of the length of the sequence, and when the bitscore reached a minimum

of 40[73]. Orthologs were detected by finding the mutually best scoring pairwise hits

(reciprocal best hits) between datasets A-B and B-A. The self-self runs were used to

identify paralogy relationships as described[72].

Co-conservation analyses. For each orthologous alternative protein pair A-B between

two species, we evaluated the presence and the orthology of their corresponding

reference proteins A'-B' in the same species. In addition, the corresponding altORFs and

CDSs had to be present in the same gene.

In order to develop a null model to assess co-conservation of alternative proteins and

their reference pairs, we needed to establish a probability that any given orthologous

alternative protein would by chance occur encoded on the same transcript as its paired,

orthologous reference protein. Although altORFs might in theory shift among CDSs (and

indeed, a few examples have been observed), transposition events are expected to be

relatively rare; we thus used the probability that the orthologous alternative protein is

paired with any orthologous CDS for our null model. Because this probability is by

definition higher than the probability that the altORF occurs on the paired CDS, it is a

conservative estimate of co-conservation. We took two approaches to estimating this

percentage, and then used whichever was higher for each species pair, yielding an even

more conservative estimate. First, we assessed the percentage of orthologous reference

proteins under the null supposition that each orthologous alternative protein had an equal

probability of being paired with any reference protein, orthologous or not. Second, we

assessed the percentage of non-orthologous alternative proteins that were paired with

orthologous reference proteins. This would account for factors such as longer CDSs

having a higher probability of being orthologous and having a larger number of paired

altORFs. For example, between *Homo sapiens* and *Mus musculus*, we found that 22,304

of 54,498 reference proteins (40.9%) were orthologs. Of the 157,261 non-orthologous

alternative proteins, 106,987 (68%) were paired with an orthologous reference protein.

Because 68% is greater than 40.9%, we used 68% as the probability for use in our null

model. Subsequently, our model strongly indicates co-conservation (Fig. 3 and Table 2;

$p<10^{-6}$ based on 1 million binomial simulations; highest observed random percentage

=69%, much lower than the observed 96% co-conservation).

Analysis of third codon position (wobble) conservation. Basewise conservation scores

for the alignment of 100 vertebrate genomes including *H. sapiens* were obtained from

UCSC genome browser

(http://hgdownload.soe.ucsc.edu/goldenPath/hg38/phyloP100way/)

(RRID:SCR_012479). Conservation PhyloP scores relative to each nucleotide position

within codons were extracted using a custom Perl script and the Bio-BigFile module

version 1.07 (see code file). The PhyloP conservation score for the wobble nucleotide of

each codon within the CDS was extracted. For the 53,862 altORFs completely nested

inside 20,814 CDSs, the average PhyloP score for wobble nucleotides within the altORF

region was compared to the average score for the complete CDS. To generate controls,

random regions in CDSs with a similar length distribution as altORFs were selected and

PhyloP scores for wobble nucleotides were extracted. We compared the differences

between altORF and CDS PhyloP scores (altORF PhyloP – CDS PhyloP) to those

generated based on random regions. We identified expected quantiles of the differences

("DQ" column in the table), and compared these to the observed differences. Because

there was greater conservation of wobble nucleotide PhyloP scores within altORFs

regions located farther from the center of their respective genes ($r = 0.08$, $p < 0.0001$),

observed differences were adjusted using an 8-knot cubic basis spline of percent distance

from center. These observed differences were also adjusted for site-specific signals as

detected in the controls.

Human alternative protein classification and in silico functional annotation.

Repeat and transposable element annotation

RepeatMasker, a popular software to scan DNA sequences for identifying and classifying

repetitive elements (RRID:SCR_012954), was used to investigate the extent of altORFs

derived from transposable elements[74]. Version 3-3-0 was run with default settings.

Alternative protein analysis using InterProScan (RRID:SCR_005829)

InterProScan combines 15 different databases, most of which use Hidden Markov models

for signature identification[75]. Interpro merges the redundant predictions into a single

entry and provides a common annotation. A recent local version of InterProScan 5.14-

53.0 was run using default parameters to scan for known protein domains in alternative

proteins. Gene ontology (GO) and pathway annotations were also reported if available

with -goterm and -pa options. Only protein signatures with an E-value $\leq 10^{-3}$ were

considered.

We classified the reported InterPro hits as belonging to one or several of three clusters;

(1) alternative proteins with InterPro entries; (2) alternative proteins with signal peptides

(SP) and/or transmembrane domains (TM) predicted by at least two of the three SignalP,

PHOBIUS, TMHMM tools and (3) alternative proteins with other signatures.

The GO terms assigned to alternative proteins with InterPro entries were grouped and

categorised into 13 classes within the three ontologies (cellular component, biological

process, molecular function) using the CateGOrizer tool[76] (RRID:SCR_005737).

Each unique alternative protein with InterPro entries and its corresponding reference

protein (encoded in the same transcript) were retrieved from our InterProscan output.

Alternative and reference proteins without any InterPro entries were ignored. The overlap

in InterPro entries between alternative and reference proteins was estimated as follows.

We went through the list of alternative/reference protein pairs and counted the overlap in

the number of entries between the alternative and reference proteins as

100*intersection/union. All reference proteins and the corresponding alternative proteins

were combined together in each comparison so that all domains of all isoforms for a

given reference protein were considered in each comparison. The random distribution of

the number of alternative/reference protein pairs that share at least one InterPro entry was

computed by shuffling the alternative/reference protein pairs and calculating how many

share at least one InterPro entry. This procedure was repeated 1,000 times. Finally, we

compared the number and identity of shared InterPro entries in a two dimensional matrix

to illustrate which Interpro entries are shared. In many instances, including for zinc-finger

coding genes, InterPro entries in alternative/reference protein pairs tend to be related

when they are not identical.

Mass Spectrometry identification. Wrapper Perl scripts were developed for the use of

SearchGUI v2.0.11[77] (RRID:SCR_012054) and PeptideShaker v1.1.0[59]

(RRID:SCR_002520) on the *Université de Sherbrooke's* 39,168 core high-performance

Mammouth Parallèle 2 computing cluster

(http://www.calculquebec.ca/en/resources/compute-servers/mammouth-parallele-ii).

SearchGUI was configured to run the following proteomics identification search engines:

X!Tandem[78], MS-GF+[79], MyriMatch[80], Comet[81], and OMSSA[82]. SearchGUI parameters

were set as follow: maximum precursor charge, 5; maximum number of PTM per peptide,

5; X!Tandem minimal fragment m/z, 140; removal of initiator methionine for Comet, 1. A

full list of parameters used for SearchGUI and PeptideShaker is available in

Supplementary file 2, sheet 1. For PXD000953 dataset[35], precursor and fragment

tolerance were set 0.006 Da and 0.1 Da respectively, with carbamidomethylation of C as

a fixed modification and Nter-Acetylation and methionine oxidation as variable

modifications. For PXD000788[33] and PXD000612[34] datasets, precursor and fragment

tolerance were set to 4.5 ppm and 0.1 Da respectively with carbamidomethylation of

cysteine as a fixed modification and Nter-Acetylation, methionine oxidation and

phosphorylation of serine, threonine and tyrosine as variable modifications. For

PXD002815 dataset[32], precursor and fragment tolerance were set to 4.5 ppm and 0.1 Da

respectively with carbamidomethylation of cysteine as a fixed modification and Nter-

Acetylation and methionine oxidation as variable modifications.Datasets were searched

using a target-decoy approach against a composite database composed of a target

database [Uniprot canonical and isoform reference proteome (16 January 2015) for a total

of 89,861 sequences + custom alternative proteome resulting from the in silico translation

of all human altORFs (available to download at

https://www.roucoulab.com/p/downloads)], and their reverse protein sequences from the

target database used as decoys. In order to separate alternative and reference proteins for

FDR analyses, PeptideShaker output files were extracted with target and decoy hits.

PSMs matching reference target or decoy proteins were separated from those matching

alternative targets or decoys as previously described [83,84]. PSMs that matched both

reference and alternative proteins were automatically moved to the reference database

group. PSMs were then ranked according to their PeptideShaker score and filtered at 1%

FDR separately. Validated PSMs were selected to group proteins using proteoQC R tool

[85], and proteins were separately filtered again using a 1% FDR cut-off.

Only alternative proteins identified with at least one unique and specific peptide were

considered valid[59]. Any peptide matching both a canonical (annotated in Uniprot) and an

alternative protein was attributed to the canonical protein. For non-unique peptides, i.e.

peptides matching more than one alternative protein, the different accession IDs are

indicated in the MS files. For subsequent analyses (e.g. conservation, protein

signature…), only one protein is numbered in the total count of alternative proteins; we

arbitrarily selected the alternative protein with the lowest accession ID.

Peptides matching proteins in a protein sequence database for common contaminants

were rejected[86].

For spectral validation (Figure 6-figure supplement1, 2, 3, and 4), synthetic peptides were

purchased from the peptide synthesis service at the *Université de Sherbrooke*. Peptides

were solubilized in 10% acetonitrile, 1% formic acid and directly injected into a Q-

Exactive mass spectrometer (Thermo Scientific) via an electro spray ionization source

(Thermo Scientific). Spectra were acquired using Xcalibur 2.2 (RRID:SCR_014593) at

70000 resolution with an AGC target of 3e6 and HCD collision energy of 25. Peaks were

assigned manually by comparing monoisotopic m/z theoretical fragments and

experimental (PeptideShaker) spectra.

In order to test if the interaction between alternative zinc-finger/reference zinc-finger

protein pairs (encoded in the same gene) may have occurred by chance only, all

interactions between alternative proteins and reference proteins were randomized with an

in-house randomisation script. The number of interactions with reference proteins for

each altProt was kept identical as the number of observed interactions. The results

indicate that interactions between alternative zinc-finger/reference zinc-finger protein

pairs did not occur by chance ($p<10^{-6}$) based on 1 million binomial simulations; highest

observed random interactions between alternative zinc-finger proteins and their reference

proteins = 3 (39 times out of 1 million simulations), compared to detected interactions=5.

Code availability. Computer codes are available upon request with no restrictions.

Data availability. Alternative protein sequence databases for different species can be

accessed at https://www.roucoulab.com/p/downloads with no restrictions.

Cloning and antibodies. Human Flag-tagged altMiD51(WT) and

altMiD51(LYR→AAA), and HA-tagged DrP1(K38A) were cloned into pcDNA3.1

(Invitrogen) using a Gibson assembly kit (New England Biolabs, E26115). The cDNA

corresponding to human MiD51/MIEF1/SMCR7L transcript variant 1 (NM_019008) was

also cloned into pcDNA3.1 by Gibson assembly. In this construct, altMiD51 and MiD51

were tagged with Flag and HA tags, respectively. MiD51^{GFP} and altMiD51^{GFP} were also

cloned into pcDNA3.1 by Gibson assembly. For MiD51^{GFP}, a LAP tag[32] was inserted

between MiD51 and GFP. gBlocks were purchased from IDT. Human altDDIT3^{mCherry}

was cloned into pcDNA3.1 by Gibson assembly using coding sequence from transcript

variant 1 (NM_004083.5) and mCherry coding sequence from pLenti-myc-GLUT4-

mCherry (Addgene plasmid # 64049). Human DDIT3^{GFP} was also cloned into pcDNA3.1

by Gibson assembly using CCDS8943 sequence.

For immunofluorescence, primary antibodies were diluted as follow: anti-Flag (Sigma,

F1804) 1/1000, anti-TOM20 (Abcam, ab186734) 1/500. For western blots, primary

antibodies were diluted as follow: anti-Flag (Sigma, F1804) 1/1000, anti-HA (BioLegend,

901515) 1/500, anti-actin (Sigma, A5441) 1/10000, anti-Drp1 (BD Transduction

Laboratories, 611112) 1/500, anti-GFP (Santa Cruz Biotechnology, sc-9996) 1/10000,

anti-mCherry (Abcam, ab125096) 1/2000.

Cell culture, immunofluorescence, knockdown and western blots. HeLa cells (ATCC

CRM-CCL-2, authenticated by STR profiling, RRID:CVCL_0030) cultures tested

negative for mycoplasma contamination (ATCC 30-1012K), transfections,

immunofluorescence, confocal analyses and western blots were carried out as previously

described[87]. Mitochondrial morphology was analyzed as previously described[88]. A

minimum of 100 cells were counted (n=3 or 300 cells for each experimental condition).

Three independent experiments were performed.

For Drp1 knockdown, 25,000 HeLa cells in 24-well plates were transfected with 25 nM

Drp1 SMARTpool: siGENOME siRNA (Dharmacon, M-012092-01-0005) or ON-

TARGET plus Non-targeting pool siRNAs (Dharmacon, D-001810-10-05) with

DharmaFECT 1 transfection reagent (Dharmacon, T-2001-02) according to the

manufacturer's protocol. After 24h, cells were transfected with pcDNA3.1 or altMiD51,

incubated for 24h, and processed for immunofluorescence or western blot. Colocalization

analyses were performed using the JACoP plugin (Just Another Co-localization Plugin) [50]

implemented in Image J software.

Immunoprecipitations. Immunoprecipitations experiments were conducted using GFP-

Trap (ChromoTek) protocol with minor modifications. Briefly, cells were lysed with co-

ip lysis buffer (0.5 % NP40, Tris-HCl 50 mM pH 7.5, NaCl 150 mM and two EDTA-free

Roche protease inhibitors per 50 mL of buffer). After 5 mins of lysis on ice, lysate was

sonicated twice at 11 % amplitude for 5 s with 3 minutes of cooling between sonication

cycles. Lysate was centrifuged, supernatant was isolated and protein content was assessed

using BCA assay (Pierce). GFP-Trap beads were conditioned with lysis buffer. 40 μL of

beads were added to 2 mg of proteins at a final concentration of 1 mg/mL. After

overnight immunoprecipitation, beads were centrifuged at 5000 rpm for 5 minutes and

supernatant was discarded. Beads were then washed three times with wash buffer (0.5 %

NP40, Tris-HCl 50 mM pH 7.5, NaCl 200 mM and two EDTA-free Roche protease

inhibitors per 50 mL of buffer) and supernatants were discarded. Immunoprecipitated

proteins were eluted from beads by adding 40 µL of Laemmli buffer and boiling at 95 °C

for 15 minutes. Eluate was split in halfs which were loaded onto 10 % SDS-PAGE gels to

allow western blotting of GFP and mCherry tagged proteins. 40 µg of initial lysates were

loaded into gels as inputs.

Mitochondrial localization, parameters and ROS production. Trypan blue quenching

experiment was performed as previously described[89].

A flux analyzer (XF96 Extracellular Flux Analyzer; Seahorse Bioscience, Agilent

technologies) was used to determine the mitochondrial function in HeLa cells

overexpressing altMiD51^{Flag}. Cells were plated in a XF96 plate (Seahorse Biosciences) at

1×10^4 cells per well in Dulbecco's modified Eagle's medium supplemented with 10%

FBS with antibiotics. After 24 hours, cells were transfected for 24 hours with an empty

vector (pcDNA3.1) or with the same vector expressing altMiD51^{Flag} with GeneCellin

tranfection reagent according to the manufacturer's instructions. Cells were equilibrated

in XF assay media supplemented with 25 mM glucose and 1 mM pyruvate and were

incubated at 37 °C in a CO_2-free incubator for 1 h. Baseline oxygen consumption rates

(OCRs) of the cells were recorded with a mix/wait/measure times of 3/0/3 min

respectively. Following these measurements, oligomycin (1 µM), FCCP (0.5 µM), and

antimycin A/rotenone (1 μM) were sequentially injected, with oxygen consumption rate

measurements recorded after each injection. Data were normalized to total protein in each

well. For normalization, cells were lysed in the 96-well XF plates using 15 μl/well of

RIPA lysis buffer (1% Triton X-100, 1% NaDeoxycholate, 0.1% SDS, 1 mM EDTA, 50

mM Tris-HCl pH 7.5). Protein concentration was measured using the BCA protein assay

reagent (Pierce, Waltham, MA, USA).

Reactive oxygen species (ROS) levels were measured using Cellular ROS/Superoxide

Detection Assay Kit (Abcam #139476). HeLa cells were seeded onto 96-well black/clear

bottom plates at a density of 6,000 cells per well with 4 replicates for each condition.

After 24 hours, cells were transfected for 24 hours with an empty vector (pcDNA3.1) or

with the same vector expressing altMiD51^{Flag} with GeneCellin according to the

manufacturer's instruction. Cells were untreated or incubated with the ROS inhibitor (N-

acetyl-L-cysteine) at 10 mM for 1 hour. Following this, the cells were washed twice with

the wash solution and then labeled for 1 hour with the Oxidative Stress Detection

Reagent (green) diluted 1:1000 in the wash solution with or without the positive control

ROS Inducer Pyocyanin at 100 μM. Fluorescence was monitored in real time. ROS

accumulation rate was measured between 1 to 3 hours following induction. After the

assay, total cellular protein content was measured using BCA protein assay reagent

(Pierce, Waltham, MA, USA) after lysis with RIPA buffer. Data were normalised for

initial fluorescence and protein concentration.

ATP synthesis was measured as previously described[90] in cells transfected for 24 hours

with an empty vector (pcDNA3.1) or with the same vector expressing altMiD51^{Flag}.

Acknowledgements

This research was supported by CIHR grants MOP-137056 and MOP-136962 to X.R;

MOP-299432 and MOP-324265 to C.L; a *Université de Sherbrooke* institutional research

grant made possible through a generous donation by Merck Sharp & Dohme to X.R; a

FRQNT team grant 2015-PR-181807 to C.L. and X.R; Canada Research Chairs in

Functional Proteomics and Discovery of New Proteins to X.R, in Evolutionary Cell and

Systems Biology to C.L and in Computational and Biological Complexity to A.O; A.A.C

is supported by a CIHR New Investigator Salary Award; M.S.S is a recipient of a *Fonds*

de Recherche du Québec – Santé Research Scholar Junior 2 Career Award; V.D is

supported in part by fellowships from *Région Nord-Pas de Calais* and PROTEO; A.A.C,

D.J.H, M.S.S and X.R are members of the *Fonds de Recherche du Québec Santé-*

supported *Centre de Recherche du Centre Hospitalier Universitaire de Sherbrooke*. We

thank the staff from the Centre for Computational Science at the *Université de*

Sherbrooke, Compute Canada and Compute Québec for access to the Mammouth

supercomputer.

References

1. Ingolia, N. T., Lareau, L. F. & Weissman, J. S. Ribosome profiling of mouse embryonic stem cells reveals the complexity and dynamics of mammalian proteomes. *Cell* **147,** 789–802 (2011).

2. Lee, S. S. *et al.* Global mapping of translation initiation sites in mammalian cells at single-nucleotide resolution. *Proc. Natl. Acad. Sci. U. S. A.* **109,** E2424-2432 (2012).

3. Mouilleron, H., Delcourt, V. & Roucou, X. Death of a dogma: eukaryotic mRNAs can code for more than one protein. *Nucleic Acids Res.* **44,** 14–23 (2015).

4. Pauli, A. *et al.* Toddler: An Embryonic Signal That Promotes Cell Movement via Apelin Receptors. *Science* **343,** 1248636–1248636 (2014).

5. Anderson, D. M. *et al.* A Micropeptide Encoded by a Putative Long Noncoding RNA Regulates Muscle Performance. *Cell* **160,** 595–606 (2015).

6. Zanet, J. *et al.* Pri sORF peptides induce selective proteasome-mediated protein processing. *Science* **349,** 1356–1358 (2015).

7. Nelson, B. R. *et al.* A peptide encoded by a transcript annotated as long noncoding RNA enhances SERCA activity in muscle. *Science (80-.).* **351,** 271–275 (2016).

8. Bazzini, A. A. *et al.* Identification of small ORFs in vertebrates using ribosome footprinting and evolutionary conservation. *EMBO J.* **33,** 981–993 (2014).

9. Ji, Z., Song, R., Regev, A. & Struhl, K. Many lncRNAs, 5'UTRs, and pseudogenes are translated and some are likely to express functional proteins. *Elife* **4,** e08890 (2015).

10. Prabakaran, S. *et al.* Quantitative profiling of peptides from RNAs classified as

noncoding. *Nat. Commun.* **5,** 5429 (2014).

11.   Slavoff, S. a *et al.* Peptidomic discovery of short open reading frame-encoded

peptides in human cells. *Nat. Chem. Biol.* **9,** 59–64 (2013).

12.   Andrews, S. J. & Rothnagel, J. A. Emerging evidence for functional peptides

encoded by short open reading frames. *Nat. Rev. Genet.* **15,** 193–204 (2014).

13.   Landry, C. R., Zhong, X., Nielly-Thibault, L. & Roucou, X. Found in translation:

Functions and evolution of a recently discovered alternative proteome. *Curr. Opin.*

Struct. Biol. **32,** 74–80 (2015).

14.   Fields, A. P. *et al.* A Regression-Based Analysis of Ribosome-Profiling Data

Reveals a Conserved Complexity to Mammalian Translation. *Mol. Cell* **60,** 816–

827 (2015).

15.   Saghatelian, A. & Couso, J. P. Discovery and characterization of smORF-encoded

bioactive polypeptides. *Nat. Chem. Biol.* **11,** 909–16 (2015).

16.   Stock, D., Leslie, A. G. & Walker, J. E. Molecular architecture of the rotary motor

in ATP synthase. *Science* **286,** 1700–1705 (1999).

17.   Schmitt, J. P. *et al.* Dilated cardiomyopathy and heart failure caused by a mutation

in phospholamban. *Science* **299,** 1410–1413 (2003).

18.   Nemeth, E. *et al.* Hepcidin regulates cellular iron efflux by binding to ferroportin

and inducing its internalization. *Science* **306,** 2090–2093 (2004).

19.   Carvunis, A.-R. *et al.* Proto-genes and de novo gene birth. *Nature* 3–7 (2012).

doi:10.1038/nature11184

20.   Schlötterer, C. Genes from scratch--the evolutionary fate of de novo genes. *Trends*

Genet. **31,** 215–9 (2015).

21.   McLysaght, A. & Hurst, L. D. Open questions in the study of de novo genes: what,

818   how and why. *Nat. Rev. Genet.* **17,** 567–578 (2016).

819   22.   Sabath, N., Wagner, A. & Karlin, D. Evolution of viral proteins originated de novo

820   by overprinting. *Mol. Biol. Evol.* **29,** 3767–80 (2012).

821   23.   Iacono, M., Mignone, F. & Pesole, G. uAUG and uORFs in human and rodent

822   5'untranslated mRNAs. *Gene* **349,** 97–105 (2005).

823   24.   Neafsey, D. E. & Galagan, J. E. Dual modes of natural selection on upstream open

824   reading frames. *Mol. Biol. Evol.* **24,** 1744–51 (2007).

825   25.   Smith, E. *et al.* Leaky ribosomal scanning in mammalian genomes: significance of

826   histone H4 alternative translation in vivo. *Nucleic Acids Res.* **33,** 1298–1308

827   (2005).

828   26.   Pop, C. *et al.* Causal signals between codon bias, mRNA structure, and the

829   efficiency of translation and elongation. *Mol. Syst. Biol.* **10,** 770 (2014).

830   27.   Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of

831   nonneutral substitution rates on mammalian phylogenies. *Genome Res.* **20,** 110–

832   121 (2010).

833   28.   Michel, A. M. *et al.* GWIPS-viz: development of a ribo-seq genome browser.

834   *Nucleic Acids Res.* **42,** D859-864 (2014).

835   29.   Raj, A. *et al.* Thousands of novel translated open reading frames in humans

836   inferred by ribosome footprint profiling. *Elife* **5,** 1–24 (2016).

837   30.   Miettinen, T. P. & Björklund, M. Modified ribosome profiling reveals high

838   abundance of ribosome protected mRNA fragments derived from 3' untranslated

839   regions. *Nucleic Acids Res.* **43,** 1019–1034 (2015).

31. Weingarten-Gabbay, S. *et al.* Systematic discovery of cap-independent translation

sequences in human and viral genomes. *Science (80-.).* **351,** 1–24 (2016).

32. Hein, M. Y. *et al.* A Human Interactome in Three Quantitative Dimensions

Organized by Stoichiometries and Abundances. *Cell* **163,** 712–723 (2015).

33. Tong, J., Taylor, P. & Moran, M. F. Proteomic analysis of the epidermal growth

factor receptor (EGFR) interactome and post-translational modifications associated

with receptor endocytosis in response to EGF and stress. *Mol. Cell. Proteomics* **13,**

1644–1658 (2014).

34. Sharma, K. *et al.* Ultradeep Human Phosphoproteome Reveals a Distinct

Regulatory Nature of Tyr and Ser/Thr-Based Signaling. *Cell Rep.* **8,** 1583–1594

(2014).

35. Rosenberger, G. *et al.* A repository of assays to quantify 10,000 human proteins by

SWATH-MS. *Sci. data* **1,** 140031 (2014).

36. Mitchell, A. *et al.* The InterPro protein families database: the classification

resource after 15 years. *Nucleic Acids Res.* **43,** D213-221 (2014).

37. Vanderperre, B. *et al.* Direct detection of alternative open reading frames

translation products in human significantly expands the proteome. *PLoS One* **8,**

e70698 (2013).

38. Wolfe, S. A., Nekludova, L. & Pabo, C. O. DNA recognition by Cys2His2 zinc

finger proteins. *Annu. Rev. Biophys. Biomol. Struct.* **29,** 183–212 (2000).

39. Laity, J. H., Lee, B. M. & Wright, P. E. Zinc finger proteins: new insights into

structural and functional diversity. *Curr. Opin. Struct. Biol.* **11,** 39–46 (2001).

40. Karimpour-Fard, A., Detweiler, C. S., Erickson, K. D., Hunter, L. & Gill, R. T.

Cross-species cluster co-conservation: a new method for generating protein

interaction networks. *Genome Biol.* **8,** R185 (2007).

41.   Schmitges, F. W. *et al.* Multiparameter functional diversity of human C2H2 zinc

finger proteins. *Genome Res.* **26,** 1742–1752 (2016).

42.   Andreev, D. E. *et al.* Translation of 5' leaders is pervasive in genes resistant to

eIF2 repression. *Elife* **4,** e03971 (2015).

43.   Kim, M.-S. *et al.* A draft map of the human proteome. *Nature* **509,** 575–581

(2014).

44.   Angerer, H. Eukaryotic LYR Proteins Interact with Mitochondrial Protein

Complexes. *Biology (Basel).* **4,** 133–150 (2015).

45.   Losón, O. C., Song, Z., Chen, H. & Chan, D. C. Fis1, Mff, MiD49, and MiD51

mediate Drp1 recruitment in mitochondrial fission. *Mol. Biol. Cell* **24,** 659–667

(2013).

46.   Motori, E. *et al.* Inflammation-Induced Alteration of Astrocyte Mitochondrial

Dynamics Requires Autophagy for Mitochondrial Network Maintenance. *Cell*

Metab. **18,** 844–859 (2013).

47.   Smirnova, E., Shurland, D. L., Ryazantsev, S. N. & van der Bliek, A. M. A human

dynamin-related protein controls the distribution of mitochondria. *J. Cell Biol.*

143, 351–358 (1998).

48.   Cui, K., Coutts, M., Stahl, J. & Sytkowski, A. J. Novel interaction between the

transcription factor CHOP (GADD153) and the ribosomal protein FTE/S3a

modulates erythropoiesis. *J. Biol. Chem.* **275,** 7591–6 (2000).

49.   Chiribau, C.-B., Gaccioli, F., Huang, C. C., Yuan, C. L. & Hatzoglou, M.

Molecular symbiosis of CHOP and C/EBP beta isoform LIP contributes to

endoplasmic reticulum stress-induced apoptosis. *Mol. Cell. Biol.* **30,** 3722–31

(2010).

50.   Bolte, S. & Cordelières, F. P. A guided tour into subcellular colocalization analysis

in light microscopy. *J. Microsc.* **224,** 213–32 (2006).

51.   Couso, J.-P. & Patraquim, P. Classification and function of small open reading

frames. *Nat. Rev. Mol. Cell Biol.* (2017). doi:10.1038/nrm.2017.58

52.   Johnstone, T. G., Bazzini, A. A. & Giraldez, A. J. Upstream ORFs are prevalent

translational repressors in vertebrates. *EMBO J.* (2016).

doi:10.15252/embj.201592759

53.   Jousse, C. *et al.* Inhibition of CHOP translation by a peptide encoded by an open

reading frame localized in the chop 5'UTR. *Nucleic Acids Res.* **29,** 4341–51

(2001).

54.   Young, S. K., Palam, L. R., Wu, C., Sachs, M. S. & Wek, R. C. Ribosome

Elongation Stall Directs Gene-specific Translation in the Integrated Stress

Response. *J. Biol. Chem.* **291,** 6546–6558 (2016).

55.   Lee, Y. C. G. & Reinhardt, J. A. Widespread Polymorphism in the Positions of

Stop Codons in Drosophila melanogaster. *Genome Biol. Evol.* **4,** 533–549 (2012).

56.   Andreatta, M. E. *et al.* The Recent De Novo Origin of Protein C-Termini. *Genome*

Biol. Evol. **7,** 1686–701 (2015).

57.   Knowles, D. G. & McLysaght, A. Recent de novo origin of human protein-coding

genes. *Genome Res.* **19,** 1752–9 (2009).

58.   Neme, R. & Tautz, D. Phylogenetic patterns of emergence of new genes support a

model of frequent de novo evolution. *BMC Genomics* **14,** 117 (2013).

59. Vaudel, M. *et al.* PeptideShaker enables reanalysis of MS-derived proteomics data

sets. *Nat. Biotechnol.* **33,** 22–24 (2015).

60. Shteynberg, D. *et al.* iProphet: multi-level integrative analysis of shotgun

proteomic data improves peptide and protein identification rates and error

estimates. *Mol. Cell. Proteomics* **10,** M111.007690 (2011).

61. Quelle, D. E., Zindy, F., Ashmun, R. A. & Sherr, C. J. Alternative reading frames

of the INK4a tumor suppressor gene encode two unrelated proteins capable of

inducing cell cycle arrest. *Cell* **83,** 993–1000 (1995).

62. Abramowitz, J., Grenet, D., Birnbaumer, M., Torres, H. N. & Birnbaumer, L.

XLalphas, the extra-long form of the alpha-subunit of the Gs G protein, is

significantly longer than suspected, and so is its companion Alex. *Proc. Natl.*

Acad. Sci. U. S. A. **101,** 8366–8371 (2004).

63. Bergeron, D. *et al.* An out-of-frame overlapping reading frame in the ataxin-1

coding sequence encodes a novel ataxin-1 interacting protein. *J. Biol. Chem.* **288,**

21824–35 (2013).

64. Lee, C. -f. C., Lai, H.-L. H.-L., Lee, Y.-C., Chien, C.-L. C.-L. & Chern, Y. The

A2A Adenosine Receptor Is a Dual Coding Gene: A NOVEL MECHANISM OF

GENE USAGE AND SIGNAL TRANSDUCTION. *J. Biol. Chem.* **289,** 1257–

1270 (2014).

65. Yosten, G. L. C. *et al.* A 5′-Upstream short open reading frame encoded peptide

regulates angiotensin type 1a receptor production and signaling via the beta-

arrestin pathway. *J. Physiol.* **6,** n/a-n/a (2015).

66.   Rice, P., Longden, I. & Bleasby, A. EMBOSS: the European Molecular Biology

Open Software Suite. *Trends Genet.* **16,** 276–277 (2000).

67.   Kozak, M. Pushing the limits of the scanning mechanism for initiation of

translation. *Gene* **299,** 1–34 (2002).

68.   Fritsch, C. *et al.* Genome-wide search for novel human uORFs and N-terminal

protein extensions using ribosomal footprinting. *Genome Res.* **22,** 2208–2218

(2012).

69.   Stern-Ginossar, N. *et al.* Decoding human cytomegalovirus. *Science* **338,** 1088–93

(2012).

70.   Gao, X. *et al.* Quantitative profiling of initiating ribosomes in vivo. *Nat. Methods*

12, 147–53 (2015).

71.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*

Methods **9,** 357–9 (2012).

72.   Sonnhammer, E. L. L. & Östlund, G. InParanoid 8: orthology analysis between

273 proteomes, mostly eukaryotic. *Nucleic Acids Res.* **43,** D234-239 (2015).

73.   Remm, M., Storm, C. E. & Sonnhammer, E. L. Automatic clustering of orthologs

and in-paralogs from pairwise species comparisons. *J. Mol. Biol.* **314,** 1041–1052

(2001).

74.   Tarailo-Graovac, M. & Chen, N. Using RepeatMasker to identify repetitive

elements in genomic sequences. *Curr. Protoc. Bioinformatics* **Chapter 4,** Unit

4.10 (2009).

75.   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.

Bioinformatics **30,** 1236–1240 (2014).

76.   Na, D., Son, H. & Gsponer, J. Categorizer: a tool to categorize genes into user-

956   defined biological groups based on semantic similarity. *BMC Genomics* **15,** 1091

957   (2014).

958   77.   Vaudel, M., Barsnes, H., Berven, F. S., Sickmann, A. & Martens, L. SearchGUI:

959   An open-source graphical user interface for simultaneous OMSSA and X!Tandem

960   searches. *Proteomics* **11,** 996–999 (2011).

961   78.   Craig, R. & Beavis, R. C. TANDEM: matching proteins with tandem mass spectra.

962   *Bioinformatics* **20,** 1466–1467 (2004).

963   79.   Kim, S. & Pevzner, P. A. MS-GF+ makes progress towards a universal database

964   search tool for proteomics. *Nat. Commun.* **5,** 5277 (2014).

965   80.   Tabb, D. L., Fernando, C. G. & Chambers, M. C. MyriMatch: highly accurate

966   tandem mass spectral peptide identification by multivariate hypergeometric

967   analysis. *J. Proteome Res.* **6,** 654–661 (2007).

968   81.   Eng, J. K., Jahan, T. A. & Hoopmann, M. R. Comet: an open-source MS/MS

969   sequence database search tool. *Proteomics* **13,** 22–24 (2013).

970   82.   Geer, L. Y. *et al.* Open mass spectrometry search algorithm. *J. Proteome Res.* **3,**

971   958–64 (2004).

972   83.   Menschaert, G. & Fenyö, D. Proteogenomics from a bioinformatics angle: A

973   growing field. *Mass Spectrom. Rev.* **36,** 584–599 (2015).

974   84.   Woo, S. *et al.* Proteogenomic strategies for identification of aberrant cancer

975   peptides using large-scale next-generation sequencing data. *Proteomics* **14,** 2719–

976   30 (2014).

85.   Gatto, L., Breckels, L. M., Naake, T. & Gibb, S. Visualization of proteomics data

using R and Bioconductor. *Proteomics* **15,** 1375–1389 (2015).

86.   Perkins, D. N., Pappin, D. J. C., Creasy, D. M. & Cottrell, J. S. Probability-based

protein identification by searching sequence databases using mass spectrometry

data. *Electrophoresis* **20,** 3551–3567 (1999).

87.   Vanderperre, B. *et al.* An overlapping reading frame in the PRNP gene encodes a

novel polypeptide distinct from the prion protein. *FASEB J.* **25,** 2373–86 (2011).

88.   Palmer, C. S. *et al.* MiD49 and MiD51, new components of the mitochondrial

fission machinery. *EMBO Rep.* **12,** 565–573 (2011).

89.   Vanderperre, B. *et al.* MPC1-like: a Placental Mammal-Specific Mitochondrial

Pyruvate Carrier Subunit Expressed in Post-Meiotic Male Germ Cells. *J. Biol.*

Chem. (2016). doi:10.1074/jbc.M116.733840

90.   Vives-Bauza, C., Yang, L. & Manfredi, G. Assay of Mitochondrial ATP Synthesis

in Animal Cells and Tissues. *Methods Cell Biol* **80,** 155–171 (2007).

FIGURE LEGENDS

Figure 1. Annotation of human altORFs.

(**a**) AltORF nomenclature. AltORFs partially overlapping the CDS must be in a different reading frame. (**b**) Pipeline for the identification of altORFs. (**c**) Size distribution of alternative (empty bars, vertical and horizontal axes) and reference (grey bars, secondary horizontal and vertical axes) proteins. Arrows indicate the median size. The median alternative protein length is 45 amino acids (AA) compared to 460 for the reference proteins. (**d**) Distribution of altORFs in the human hg38 transcriptome. (**e, f**) Number of total altORFs (**e**) or number of altORFs/10kbs (**f**) in hg38 compared to shuffled hg38. Means and standard deviations for 100 replicates obtained by sequence shuffling are shown. Statistical significance was determined by using one sample t-test with two-tailed *p*-values. **** *p*<0.0001. (**g**) Percentage of altORFs with an optimal Kozak motif. The total number of altORFs with an optimal Kozak motif is also indicated at the top.

Figure 1-figure supplement 1. 10% of altORFs are present in different classes of repeats.

While more than half of the human genome is composed of repeated sequences, only 9.83% or 18,003 altORFs are located inside these repeats (**a**), compared to 2,45% or 1,677 CDSs (**b**). AltORFs and CDSs are detected in non-LTR retrotransposons (LINEs, SINEs, SINE-VNTR-Alus), LTR repeats, DNA repeats, satellites and other repeats. Proportions were determined using RepeatMasker (version 3.3.0).

Figure 1-figure supplement 2. The proportion of altORFs with a translation

initiation site (TIS) with a Kozak motif in hg38 is significantly different from 100

shuffled hg38 transcriptomes.

Percentage of altORFs with a TIS within an optimal Kozak sequence in hg38 (dark blue)

compared to 100 shuffled hg38 (light blue). Mean and standard deviations for sequence

shuffling are displayed, and significant difference was defined by using one sample t test.

**** *P*<0.0001. Note that shuffling all transcripts in the hg38 transcriptome generates a

total of 489,073 altORFs on average, compared to 539,134 altORFs in hg38. Most

transcripts result from alternative splicing and there are 183,191 unique altORFs in the

hg38 transcriptome, while the 489,073 altORFs in shuffled transcriptomes are all unique.

Figure 1g shows the percentage of unique altORFs with a kozak motif (15%), while the

current Fig. shows the percentage of altORFs with a kozak motif relative to the total

number of altORFs (14%).

1028

Figure 2. Conservation of alternative and reference proteins across different species.

(**a**) Number of orthologous and paralogous alternative and reference proteins between *H.*

sapiens and other species (pairwise study). (**b**) Phylogenetic tree: conservation of

alternative (blue) and reference (red) proteins across various eukaryotic species. (**c**)

Number and fraction of genes encoding homologous reference proteins or at least 1

homologous alternative protein between *H. sapiens* and other species (pairwise study).

1035

1036 **Figure 3. Number of orthologous and co-conserved alternative and reference**

1037 **proteins between *H. sapiens* and other species (pairwise).** For the co-conservation

1038   analyses, the percentage of observed (Obs.), expected (Exp.) and corresponding *p*-values

1039   relative to the total number of reference-alternative protein pairs are indicated on the right

1040   (see Table 2 for details).

1041

Figure 4. AltORFs completely nested within CDSs show more extreme PhyloP values

(more conserved or faster evolving) than their CDSs. Differences between altORF and

CDS PhyloP scores (altORF PhyloP – CDS PhyloP, *y*-axis) are plotted against PhyloPs

for their respective CDSs (*x*-axis). We restricted the analysis to altORF-CDS pairs that

were co-conserved from humans to zebrafish. The plot contains 889 CDSs containing at

least one fully nested altORF, paired with one of its altORFs selected at random (to avoid

problems with statistical non-independence). PhyloPs for both altORFs and CDSs are

based on 3rd codons in the CDS reading frame, calculated across 100 vertebrate species.

We compared these differences to those generated based on five random regions in CDSs

with a similar length as altORFs. Expected quantiles of the differences ("DQ" columns)

were identified and compared to the observed differences. We show the absolute numbers

("n") and observed-to-expected ratios ("O/E") for each quantile. There are clearly

substantial over-representations of extreme values (red signaling conservation DQ 0.95,

and blue signaling accelerated evolution DQ 0.05) with 317 of 889 altORFs (35.7%). A

random distribution would have implied a total of 10% (or 89) of altORFs in the extreme

values. This suggests that 25.7% (35.7%-10%) of these 889 altORFs undergo specific

selection different from random regions in their CDSs with a similar length distribution.

This percentage is very similar to the 26.2% obtained from an analysis of altORFs

without restriction based on co-conservation in vertebrates (see Figure 4-figure

supplement 1), a total which would imply that there are about 4,458 altORFs fully nested

in CDSs undergoing conserved or accelerated evolution relative to their CDSs.

1063

Figure 4-figure supplement 1. AltORFs completely nested within CDSs show more

extreme PhyloP values (more conserved or faster evolving) than their CDSs.

Differences between altORF and CDS PhyloP scores (altORF PhyloP – CDS PhyloP, y-

axis) are plotted against PhyloPs for their respective CDSs (x-axis). The plot contains all

20,814 CDSs containing at least one fully nested altORF, paired with one of its altORFs

selected at random (to avoid problems with statistical non-independence). PhyloPs for

both altORFs and CDSs are based on third codon positions in the CDS reading frame,

calculated across 100 vertebrate species. We compared these differences to those

generated based on five random regions in CDSs with a similar length as altORFs.

Expected quantiles of the differences ("DQ" columns) were identified and compared to

the observed differences. We show the absolute numbers ("n") and observed-to-expected

ratios ("O/E") for each quantile. There are clearly substantial over-representations of

extreme values (red signalling conservation DQ≥0.95, and blue signalling accelerated

evolution DQ≤0.05) with 6,428 of 19,705 altORFs (36.2%). A random distribution would

have implied a total of 10% (or 1,970) of altORFs in the extreme values. This suggests

that 26.2% (36.2%-10%) of altORFs (or 4,458) undergo specific selection different from

random regions in their CDSs with a similar length distribution.

1081

Figure 5. First, second, and third codon nucleotide PhyloP scores for 100 vertebrate

species for the CDSs of the *NTNG1*, *RET* and *VTI1A* genes. Chromosomal coordinates

for the different CDSs and altORFs are indicated on the right. The regions highlighted in

red indicate the presence of an altORF characterized by a region with elevated PhyloP

scores for wobble nucleotides. The region of the altORF is indicated by a black bar above

each graph.

1088

Figure 6. Expression of human altORFs.

(**a**) Percentage of CDSs and altORFs with detected TISs by ribosomal profiling and

footprinting of human cells[23]. The total number of CDSs and altORFs with a detected TIS

is indicated at the top. (**b**) Alternative and reference proteins detected in three large

proteomic datasets: human interactome[32], 10,000 human proteins[35], human

phosphoproteome[34], EGFR interactome[33]. Numbers are indicates above each column. (**c**)

Percentage of altORFs encoding alternative proteins detected by MS-based proteomics.

The total number of altORFs is indicated at the top. Localization "Unknown" indicates

that the detected peptides can match more than one alternative protein. Localization ">1"

indicates that the altORF can have more than one localization in different RNA isoforms.

1099

Figure 6-figure supplement 1. Spectra validation for altSLC35A4[5']

Example of validation for altSLC35A4[5'] specific peptide

RVEDEVNSGVGQDGSLLSSPFLK. (**a**) Experimental MS/MS spectra (PeptideShaker

graphic interface output). (**b**) MS/MS spectra of the synthetic peptide.

Matching peaks are shown with blue masks. A diagram of the transcript with its accession

number and the localization of the altORF and the CDS is shown at the top.

1106

Figure 6-figure supplement 2. Spectra validation for altRELT[5']

Example of validation for altRELT[5'] specific peptide VALELLK. (**a**) Experimental

MS/MS spectra (PeptideShaker graphic interface output). (**b**) MS/MS spectra of the

synthetic peptide.

Matching peaks are shown with blue masks. A diagram of the transcript with its accession

number and the localization of the altORF and the CDS is shown at the top.

1113

Figure 6-figure supplement 3. Spectra validation for altLINC01420[nc]

Example of validation for altLINC01420[nc] specific peptide

WDYPEGTPNGGSTTLPSAPPPASAGLK. (**a**) Experimental MS/MS spectra

(PeptideShaker graphic interface output). (**b**) MS/MS spectra of the synthetic peptide.

Matching peaks are shown with blue masks. A diagram of the transcript with its accession

number and the localization of the altORF is shown at the top.

1120

Figure 6-figure supplement 4. Spectra validation for altSRRM2[CDS]

Example of validation for altSRRM2[CDS] specific peptide EVILDPDLPSGVGPGLHR.

(**a**) Experimental MS/MS spectra (PeptideShaker graphic interface output). (**b**) MS/MS

spectra of the synthetic peptide.

Matching peaks are shown with blue masks. A diagram of the transcript with its accession

number and the localization of the altORF and the CDS is shown at the top.

1127

1128 **Figure 7. The alternative phosphoproteome in mitosis and EGF-treated cells.**

1129 Heatmap showing relative levels of spectral counts for phosphorylated peptides following

1130   the indicated treatment[34]. For each condition, heatmap colors show the percentage of

1131   spectral count on total MS/MS phosphopeptide spectra. Blue bars on the right represent

1132   the number of MS/MS spectra; only proteins with spectral counts above 10 are shown.

1133

Figure 7-figure supplement 1: Example of a phosphorylated peptide in mitosis -

alternative protein AltLINC01420^{nc} (LOC550643, IP_305449.1).

(**a**) AltLINC01420^{nc} amino acid sequence with detected peptides underlined and

phosphorylated peptide in bold (73,9% sequence coverage). (**b**) MS/MS spectrum for the

phosphorylated peptide (PeptideShaker graphic interface output). The phosphorylation

site is the tyrosine residue, position 2. (**c**) MS/MS spectrum for the non-phosphorylated

peptide. The mass difference between the precursor ions between both spectra

corresponds to that of a phosphorylation, confirming the specific phosphorylation of this

residue in mitosis.

1143

1144   **Figure 8. Number of alternative proteins detected by ribosome profiling and mass**

1145   **spectrometry.**

1146   The expression of 467 alternative proteins was detected by both ribosome profiling

1147   (translation initiation sites, TIS) and mass spectrometry (MS).

1148

Figure 9. Human alternative proteome sequence analysis and classification using

InterProScan.

(**a**) InterPro annotation pipeline. (**b**) Alternative and reference proteins with InterPro

signatures. (**c**) Number of alternative and reference proteins with transmembrane domains

(TM), signal peptides (S) and both TM and SP. (**d**) Number of all alternative and

reference proteins predicted to be intracellular, membrane, secreted and membrane-

spanning and secreted. [1]Proteins with at least one InterPro signature; [2]Proteins with no

predicted signal peptide or transmembrane features. (**e**) Number of predicted TM regions

for alternative and reference proteins.

1158

Figure 9-figure supplement 1: Alternative proteome sequence analysis and

classification in *P. troglodytes*, *M. musculus*, *B. Taurus*, *D. melanogaster* and *S.*

***cerevisiae*.**

For each organism, the number of InterPro signatures (top graphs) and proteins with

transmembrane (TM), signal peptide (SP), or TM+SP features (bottom pie charts) is

indicated for alternative and reference proteins.

1165

Figure 10. Gene ontology (GO) annotations for human alternative proteins.

GO terms assigned to InterPro entries are grouped into 13 categories for each of the three

ontologies. (**a**) 34 GO terms were categorized into cellular component for 107 alternative

proteins. (**b**) 64 GO terms were categorized into biological process for 128 alternative

proteins. (**c**) 94 GO terms were categorized into molecular function for 302 alternative

proteins. The majority of alternative proteins with GO terms are predicted to be

intracellular, to function in nucleic acid-binding, catalytic activity and protein binding

and to be involved in biosynthesis and nucleic acid metabolism processes.

1174

1175 **Figure 11. Main InterPro entries in human alternative proteins.** (**a**) The top 10

1176 InterPro families in the human alternative proteome. (**b**) A total of 110 alternative

1177 proteins have between 1 and 23 zinc finger domains.

1178

Figure 12. Reference and alternative proteins share functional domains.

(**a**) Distribution of the number of shared InterPro entries between alternative and

reference proteins coded by the same transcripts. 138 pairs of alternative and reference

proteins share between 1 and 4 protein domains (InterPro entries). Only

alternative/reference protein pairs that have at least one domain are considered (n = 298).

(**b**) The number of reference/alternative protein pairs that share domains (n = 138) is

higher than expected by chance alone. The distribution of expected pairs sharing domains

and the observed number are shown. (**c**) Matrix of co-occurrence of domains related to

zinc fingers. The entries correspond to the number of times entries co-occur in reference

and alternative proteins. The full matrix is available in figure 12-figure supplement 1.

1189

1190 **Figure 12-figure supplement 1. Matrix of co-occurrence of InterPro entries between**

1191 **alternative/reference protein pairs coded by the same transcript.**

1192 Pixels show the number of times entries co-occur in reference and alternative proteins.

1193 Blue pixels indicate that these domains are not shared, white pixels indicate that they are

1194 shared once, and red that they are shared twice or more.

1195

Figure 12-figure supplement 2. Reference and alternative proteins share functional

domains.

The number of reference/alternative protein pairs that share domains (n = 49) is higher

than expected by chance alone ($p<0.001$). The distribution of expected pairs sharing

domains and the observed number are shown. This is the same analysis as the one

presented in figure 12b, with the zinc finger domains taken out.

1202

Figure 13. AltMiD51$^{5'}$ expression induces mitochondrial fission.

(**a**) AltMiD51$^{5'}$ coding sequence is located in exon 2 or the *MiD51/MIEF1/SMCR7L* gene

and in the 5'UTR of the canonical mRNA (RefSeq NM_019008). +2 and +1 indicate

reading frames. AltMiD51 amino acid sequence is shown with the LYR tripeptide shown

in bold. Underlined peptides were detected by MS. (**b**) Human HeLa cells transfected

with empty vector (mock), a cDNA corresponding to the canonical MiD51 transcript with

a Flag tag in frame with altMiD51 and an HA tag in frame with MiD51, altMiD51^{Flag}

cDNA or MiD51^{HA} cDNA were lysed and analyzed by western blot with antibodies

against Flag, HA or actin, as indicated. (**c**) Confocal microscopy of mock-transfected

cells, cells transfected with altMiD51^{WT}, altMiD51$^{LYR\rightarrow AAA}$ or Drp1^{K38A} immunostained

with anti-TOM20 (red channel) and anti-Flag (green channel) monoclonal antibodies. In

each image, boxed areas are shown at higher magnification in the bottom right corner. %

of cells with the most frequent morphology is indicated: mock (tubular), altMiD51^{WT}

(fragmented), altMiD51(LYR$\rightarrow$AAA) (tubular), Drp1(K38A) (elongated). Scale bar, 10

mm. (**d**) Bar graphs show mitochondrial morphologies in HeLa cells. Means of three

independent experiments per condition are shown (100 cells for each independent

experiment). ***$p<0.0005$ (Fisher's exact test) for the three morphologies between

altMiD51(WT) and the other experimental conditions.

1221

Figure 13-figure supplement 1. Spectra validation for altMiD51.

Example of validation for altMiD51 specific peptides YTDRDFYFASIR and

GLVFLNGK. (**a,c**) Experimental MS/MS spectra (PeptideShaker graphic interface

output). (**b,d**) MS/MS spectra of the synthetic peptides.

Matching peaks are shown with blue masks. A diagram of the transcript with its accession

number and the localization of the altORF and the CDS is shown at the top.

1228

Figure 13-figure supplement 2. MiD51 expression results in mitochondrial fission.

(**a**) Confocal microscopy of HeLa cells transfected with MiD51^{GFP} immunostained with

anti-TOM20 (red channel) monoclonal antibodies. In each image, boxed areas are shown

at higher magnification in the bottom right corner. The localization of MiD51 in fission

sites is shown in merged higher magnification inset. Scale bar, 10 mm. (**b**) Human HeLa

cells transfected with empty vector (mock) or MiD51^{GFP} were lysed and analyzed by

western blot to confirm MiD51^{GFP} expression.

1236

Figure 13-figure supplement 3: AltMiD51 is localized in the mitochondrial matrix.

Trypan blue quenching experiment performed on HeLa cells stably expressing the

indicated constructs. The fluorescence remaining after quenching by trypan blue is shown

relative to Matrix-Venus (Mx-Venus) indicated by the dashed line. (**** p <0,0001, one-

way ANOVA). The absence of quenching of the fluorescence compared to IMS-Venus

indicates the matricial localization of altMiD51. n≥3 cells were quantified per

experiment, and results are from 6 independent experiments. Data are mean ±SEM.

1244

Figure 13-figure supplement 4. Mitochondrial function parameters.

(**a**) Oxygen consumption rates (OCR) in HeLa cells transfected with empty vector (mock) or altMiD51[Flag]. Mitochondrial function parameters were assessed in basal conditions (basal), in the presence of oligomycin to inhibit the ATP synthase (oxygen consumption that is ATP-linked), FCCP to uncouple the mitochondrial inner membrane and allow for maximum electron flux through the respiratory chain (maximal OCR), and antimycin A/rotenone to inhibit complex III (non-mitochondrial). The balance of the basal OCR comprises oxygen consumption due to proton leak and nonmitochondrial sources. The mitochondrial reserve capacity (maximal OCR- basal OCR) is an indicator of rapid adaptation to stress and metabolic changes. Mean values of replicates are plotted with error bars corresponding to the 95% confidence intervals. Statistical significance was estimated using a two-way ANOVA with Tukey's post-hoc test (**p = 0,004). (**b**) ROS production in mock and altMiD51-expressing cells. Cells were untreated, treated with a ROS inducer or a ROS inhibitor. Results represent the mean value out of three independent experiments, with error bars corresponding to the standard error of the mean (s.e.m.). Statistical significance was estimated using unpaired T-test. (**c**) ATP synthesis rate in mock and altMiD51-expressing cells. No significant differences in ATP production were observed between mock and altMiD51 transfected cells. Results represent the mean of three independent experiments (8 technical replicates each). Error bars represent the standard error of the mean. At the end of the experiments, cells were collected and proteins analyzed by western blot with antibodies against the Flag tag (altMiD51) or actin, as indicated, to verify the expression of altMiD51. A representative western blot is shown on the right. Molecular weight markers are shown

1268   on the left (kDa).

1269

Figure 13-figure supplement 5. Representative confocal images of cells co-expressing

altMiD51^{GFP} and Drp1(K38A)^{HA}.

(**a**) Confocal microscopy of HeLa cells co-transfected with altMiD51^{GFP} and

Drp1(K38A)^{HA} immunostained with anti-TOM20 (blue channel) and anti-HA (red

channel) monoclonal antibodies. In each image, boxed areas are shown at higher

magnification in the bottom right corner. % of cells with the indicated morphology is

indicated on the TOM20 panels. (**b**) Confocal microscopy of HeLa cells co-transfected

with altMiD51^{GFP} and Drp1(wt)^{HA} immunostained with anti-TOM20 (blue channel) and

anti-HA (red channel) monoclonal antibodies. In each image, boxed areas are shown at

higher magnification in the bottom right corner. Scale bar, 10 mm.

1280

Figure 13-figure supplement 6. Protein immunoblot showing the expression of

different constructs in HeLa cells.

HeLa cells were transfected with empty vector (pcDNA3.1), altMiD51(WT)^{Flag},

altMID51(LYR→AAA)^{Flag}, Drp1(K38A)^{HA}, or Drp1(K38A)^{HA} and altMiD51(WT)^{Flag}, as

indicated. Proteins were extracted and analyzed by western blot with antibodies against

the Flag tag (altMiD51), the HA tag (Drp1K38A) or actin, as indicated. Molecular weight

markers are shown on the left (kDa). Representative experiment of three independent

biological replicates.

1289

Figure 14. AltMiD51-induced mitochondrial fragmentation is dependent on Drp1.

(**a**) Bar graphs show mitochondrial morphologies in HeLa cells treated with non-target or

Drp1 siRNAs. Cells were mock-transfected (pcDNA3.1) or transfected with

altMiD51^{Flag}. Means of three independent experiments per condition are shown (100 cells

for each independent experiment). ***p<0.0005 (Fisher's exact test) for the three

morphologies between altMiD51 and the other experimental conditions. (**b**) HeLa cells

treated with non-target or Drp1 siRNA were transfected with empty vector (pcDNA3.1)

or altMiD51^{Flag}, as indicated. Proteins were extracted and analyzed by western blot with

antibodies against the Flag tag (altMiD51), Drp1 or actin, as indicated. (**c**) Confocal

microscopy of Drp1 knockdown cells transfected with altMiD51^{GFP} immunostained with

anti-TOM20 (blue channel) and anti-Drp1 (red channel) monoclonal antibodies. In each

image, boxed areas are shown at higher magnification in the bottom right corner. % of

cells with the indicated morphology is indicated on the TOM20 panels. Scale bar, 10 mm.

(**d**) Control Drp1 immunostaining in HeLa cells treated with a non-target siRNA. For (**c**)

and (**d**), laser parameters for Drp1 and TOM20 immunostaining were identical.

1305

Figure 15. AltDDIT3$^{5'}$ co-localizes and interacts with DDIT3.

(**a**) AltDDIT3$^{5'}$ coding sequence is located in exons 1 and 2 or the

DDIT3/CHOP/GADD153 gene and in the 5'UTR of the canonical mRNA (RefSeq

NM_004083.5). +2 and +1 indicate reading frames. AltDDIT3 amino acid sequence is

also shown. (**b**) Confocal microscopy analyses of HeLa cells co-transfected with

altDDIT3^{GFP} (green channel) and DDIT3^{mCherry} (red channel). Scale bar, 10 µm. (**c, d**)

Colocalization analysis of the images shown in (**b**) performed using the JACoP plugin

(Just Another Co-localization Plugin) implemented in Image J software (two independent

biological replicates). (**c**) Scatterplot representing 50 % of green and red pixel intensities

showing that altDDIT3^{GFP} and DDIT3^{mCherry} signal highly correlate (with Pearson

correlation coefficient of 0.92 (*p*-value < 0.0001)). (**d**) Binary version of the image shown

in (**b**) after Costes' automatic threshold. White pixels represent colocalization events (*p*-

value < 0.001, based on 1000 rounds of Costes' randomization colocalization analysis).

The associated Manders Correlation Coefficient, M_1 and M_2, are shown in the right upper

corner. M_1 is the proportion of altDDIT3^{GFP} signal overlapping DDIT3^{mCherry} signal and

M_2 is the proportion of DDIT3^{mCherry} signal overlapping altDDIT3^{GFP}. (**e**) Representative

immunoblot of co-immunoprecipitation with GFP-Trap agarose beads performed on

HeLa lysates co-expressing DDIT3^{mcherry} and altDDIT3^{GFP} or DDIT3^{mcherry} with

pcDNA3.1^{GFP} empty vector (two independent experiments).

1325

Figure 15-figure supplement 1. Protein immunoblot showing the expression of

different constructs in HeLa cells.

HeLa cells were co-transfected with GFP and mCherry, or altDDIT3^{GFP} and

DDIT3^{mCherry}, as indicated. Proteins were extracted and analyzed by western blot with

antibodies, as indicated. Molecular weight markers are shown on the left (kDa).

AltDDIT3 has a predicted molecular weight of 4.28 kDa and thus migrates at its expected

molecular weight when tagged with GFP (~32 kDa). Representative experiment of two

independent biological replicates.

1334

Figure 15–figure supplement 2. Colocalization of altDDIT3 with DDIT3.

Scatter plots of Pearson's Correlation Coefficient and Manders' Correlation Coefficient

after Costes' automatic threshold (p-value < 0.001, based on 1000 rounds of Costes'

randomization colocalization analysis). M1 is the proportion of altDDIT3^{GFP} signal

overlapping DDIT3^{mCherry} signal and M2 is the proportion of DDIT3^{mCherry} signal

overlapping altDDIT3^{GFP}. Error bars represent the mean +/- SD of three independent

experiments (28 cells).

SUPPLEMENTARY FILES

1343

1344 **Supplementary file 1: 12,616 alternative proteins and 26,531 reference proteins with**

1345 **translation initiation sites detected by ribosome profiling after re-analysis of large**

1346 **scale studies. Sheet 1: general information.** Sheet 2: list of alternative proteins; sheet 3:

1347 pie chart of corresponding altORFs localization. Sheet 4: Sheet 2: list of reference

1348 proteins

1349

1350 **Supplementary file 2: 4,872 alternative proteins detected by mass spectrometry**

1351 **(MS) after re-analysis of large proteomic studies.** Sheet 1: MS identification

1352 parameters; sheet 2: raw MS output; sheet 3: list of detected alternative proteins; sheet 4:

1353 pie chart of corresponding altORFs localization.

1354

1355 **Supplementary file 3: list of phosphopeptides.**

1356

1357 **Supplementary file 4: linker sequences separating adjacent zinc finger motifs.**

1358

1359 **Supplementary file 5: 383 alternative proteins detected by mass spectrometry in the**

1360 **interactome of 118 zinc finger proteins.** Sheet 1: MS identification parameters; sheet 2:

1361 raw MS output; sheet 3: list of detected alternative proteins.

1362

1363 **Supplementary file 6: high-confidence list of predicted functional and co-operating**

1364 **alternative proteins based on co-conservation and expression analyses.** Sheet 1: high

1365   confidence list in mammals; sheet 2: high confidence list in in vertebrates.

1 **Table 1: AltORFs and alternative protein annotations in different organisms**

Genomes	Features					
	Transcripts		Current annotations		Annotations of alternative protein coding sequences	
	mRNAs	Others[1]	CDSs	Proteins	altORFs	Alternative proteins
H. sapiens GRCh38 RefSeq GCF_000001405.26	67,765	11,755	68,264	54,498	539,134	183,191
P. troglodytes 2.1.4 RefSeq GCF_000001515.6	55,034	7,527	55,243	41,774	416,515	161,663
M. musculus GRCm38p2, RefSeq GCF_000001635.22	73,450	18,886	73,551	53,573	642,203	215,472
B. Taurus UMD3.1.86	22,089	838	22,089	21,915	79,906	73,603
X. tropicalis Ensembl JGI_4.2	28,462	4,644	28,462	22,614	141,894	69,917
D rerio Ensembl ZV10.84	44,198	8,196	44,198	41,460	214,628	150,510
D. melanogaster RefSeq GCA_000705575.1	30,255	3,474	30,715	20,995	174,771	71,705
C. elegans WBcel235, RefSeq GCF_000002985.6	28,653	25,256	26,458	25,750	131,830	45,603
S. cerevisiae YJM993_v1, RefSeq GCA_000662435.1	5,471	1,463	5,463	5,423	12,401	9,492

3 [1]Other transcripts include miRNAs, rRNAs, ncRNAs, snRNAs, snoRNAs, tRNAs. [2]Annotated retained-intron and processed transcripts were

4 classified as mRNAs.

Table 2: orthology and co-conservation assessment of alternative-reference protein pairs between H. sapiens and other species

	A	B	C	D	E	F	G	H	I	J
						Observed	Mean expected		Max expected	
	Orthologous altProts (of 183,191 total)	Orthologous refProts	Co-conserved altProt-refProt pairs	Non-orthologous altProts	Non-orthologous altProts paired with orthologous refProts	Co-conservation (C/A)	% orthologous refProts (B/54,498)	% non-orthologous altProts paired with an orthologous refProt (E/D)	Max % of 1 million binomial simulations, p=max(G, H), n=A	Inferred p-value
P. troglodytes	113,687	25,755	100,839	69,504	30,772	88.69	47.20	44.27	50.39	<1e-06
M. musculus	25,930	22,304	24,862	157,261	106,987	95.88	40.09	68.031	69.39	<1e-06
B. taurus	25,868	16,887	24,426	157,323	99,369	94.42	30.98	63.16	64.67	<1e-06
X. tropicalis	2,470	12,458	1,974	180,721	95,499	79.91	22.85	52.84	57.81	<1e-06
D. rerio	2,023	12,791	1,203	181,168	94,426	59.46	23.47	52.12	57.29	<1e-06
D. melanogaster	115	4,881	51	183,076	34,352	44.34	8.95	18.76	38.26	<1e-06
C. elegans	34	3,954	8	183,157	26,839	23.52	7.25	14.65	50.00	0.02
S. cerevisiae	6	1,854	2	183,185	10,935	33.33	3.40	5.96	83.33	0.04

In order to compare the observed co-conservation to expected co-conservation, we used the more conservative of two expected values: either the percentage of all refProts (called here reference proteins) that were defined as orthologous (column G), or the percentage of non-orthologous altProts (called here alternative proteins) that were paired with an orthologous refProt. Both of these methods are themselves conservative, as they do not account for the conservation of the pairing. The larger of these values for each species was then used to generate 1 million random binomial distributions with n=# of orthologous altProts; the maximum of these percentages is reported in column I.

1 **Table 3: alternative zinc finger proteins detected by mass spectrometry (MS) and ribosome**
2 **profiling (RP)**

Alternative protein accession	Detection method[1]	Gene	Amino acid sequence	AltORF localization
IP_238718.1	MS	RP11	MLVEVACSSCRSLLHKGAGASEDGAALEPAHTGGKENGATT	nc
IP_278905.1	RP	ZNF761	MSVARPLVGSHILYAIIDFILERNLISVMSVARTLVRSHPLYAT IDFILERNLTSVMSVARPLVRSQTLHAIVDFILEKNKCNECGE VFNQQAHLAGHHRIHTGEKP	CDS
IP_278745.1	MS and RP	ZNF816	MSVARPSVRNHPFNAIIYFTLERNLTNVKNVTMFTFADHTLK DIGRFILERDHTNVRFVTRFSGVIHTLQNIREFILERNHTSVINV AGVSVGSHPFNTIIHFTLERNLTHVMNVARFLVEEKTLHVIID FMLERNLTNVKNVTKFSVADHTLKDIGEFILGKNHTNVRFVT RLSGVIHALQTIREFILERNLTSVINVRRFLIKKESLHNIREFILE RNLTSVMNVARFLIKKQALQNIREFILQRNLTSVMSVAKPLL DSQHLFTIKQSMGVGKLYKCNDCHKVFSNATTIANHYRIHIE ERSTSVINVANFSDVIHNL	CDS
IP_138289.1	MS	ZSCAN3 1	MNIGGATLERNPINVRSVGKPSVPAMASLDTEESTQGKNHM NAKCVGRLSSSAHALFSIRGYTLERSAISVVSVAKPSFRMQGF SSISESTLVRNPISAVSAVNSLVSGHFLRNIRKSTLERDHKGDE FGKAFSHHCNLIRHFRIHTVPAELD	CDS
IP_278564.1	MS	ZNF808	MIVTKSSVTLQQLQIIGESMMKRNLLSVINVACFSDIVHTLQFI GNLILERNLTNVMIEARSSVKLHPMQNRRIHTGEKPHKCDDC GKAFTSHSHLVGHQRIHTGQKSCKCHQCGKVFSPRSLLAEHE KIHF	3'UTR
IP_275012.1	MS	ZNF780 A	MKPCECTECGKTFSCSSNIVQHVKIHTGEKRYNVRNMGKHLL WMISCLNIRKFRIVRNFVTIRSVDKPSLCTKNLLNTRELILMRN LVNIKECVKNFHHGLGFAQLLSIHTSEKSLSVRNVGRFIATLN TLEFGEDNSCEKVFE	3'UTR
IP_270595.1[2]	MS	ZNF440	MHSVERPYKCKICGRGFYSAKSFQIHEKSYTGEKPYECKQCG KAFVSFTSFRYHERTHTGENPYECKQFGKAFRSVKNLRFHKR THTGEKPCECKKCRKAFHNFSSLQIHERMHRGEKLCECKHCG KAFISAKIL	CDS
IP_270643.1[2]	MS	ZNF763	MKKLTLERNPINACHVVKPSIFPVPFSIMKGLTLERNPMSVSV GKPSDVPHTFEGMVGLTGEKPYECKECGKAFRSASHLQIHER TQTHIRIHSGERPYKCKTCGKGFYSPTSFQRHEKTHTAEKPYE CKQCGKAFSSSSSFWYHERTHTGEKPYECKQCGKAFRSASIQ MHAGTHPEEKPYECKQCGKAFRSAPHLRIHGRTHTGEKPYEC KECGKAFRSAKNLRIIHERTQTHVRMHSVERPYKCKICGKGFY SAKSFQIPEKSYTGEKPYECKQCGKAFISFTSFR	3'UTR
IP_270597.1[3]	MS	ZNF440	MKNLTLERNPMSVSNVGKPLFPSLPFDIMKGLTLERTPMSVS NLGKPSDLSKIFDFIKGHTLERNPVNVRNVEKHSIISLLCKYM KGCTEERSSVNVSIVGKHSYLPRSFEYMQEHTMERNPMNVK NAEKHSACLLPFIDMKRLTLEGNTMNASNVAKLSLLPVLFNI MKEHTREKPYQCKQCAKAFISSTSFQYHERTHMGEKPYECM PSGKAFISSSSLQYHERTHTGEKPYEYKQCGKAFRSASHLQM HGRTHTGEKPYECKQYGKAFRPDKIL	3'UTR
IP_270609.1[3]	MS	ZNF439	MNVSNVAKAFTSSSSFQYHERTHTGEKPYQCKQCGKAVRSA SRLQMHGSTHTWQKLYECKQYGKAFRSARIL	3'UTR
IP_270663.1[3]	MS	ZNF844	MHGRTHTQEKPYECKQCGKAFIFSTSFRYHERTHTGEKPYEC KQCGKAFRSATQLQMHRKIHTGEKPYECKQCGKAYRSVSQL LVHERTHTVEQPYEYKQYGKAFRFAKNLQIQTMNVNN	CDS
IP_270665.1[3]	MS	ZNF844	MHRKIHTGEKPYECKQCGKAYRSVSQLLVHERTHTVEQPYE YKQYGKAFRFAKNLQIQTMNVNN	CDS
IP_270668.1[3]	MS	ZNF844	MSSTAFQYHEKTHTREKHYECKQCGKAFISSGSLRYHERTHT GEKPYECKQCGKAFRSATQLQMHRKIHTGEKPYECKQCGKA YRSVSQLLVHERTHTVEQPYEYKQYGKAFRFAKNLQIQTMN	3'UTR

			VNN	
IP_138139.1	MS	ZNF322	MLSPSRCKRIHTGEQLFKCLQCQLCCRQYEHLIGPQKTHPGEKPQQV	3'UTR
IP_204754.1	RP	ZFP91-CNTF	MPGETEEPRPPEQQDQEGGEAAKAAPEEPQQRPPEAVAAAPAGTTSSRVLRGGRDRGRAAAAAAAAAVSRRRKAEYPRRRRSSPSARPPDVPGQQPQAAKSPSPVQGKKSPRLLCIEKVTTDKDPKEEKEEEDDSALPQEVSIAASRPSRGWRSSRTSVSRHRDTENTRSSRSKTGSLQLICKSEPNTDQLDYDVGEEHQSPGGISSEEEEEEEEEMLISEEEIPFKDDPRDETYKPHLERETPKPRRKSGKVKEEKEKKEIKVEVEVEVKEEENEIREDEEPPRKRGRRRKDDKSPRLPKRRKKPPIQYVRCEMEGCGTVLAHPRYLQHHIKYQHLLKKKYVCPHPSCGRLFRLQKQLLRHAKHHTDQRDYICEYCARAFKSSHNLAVHRMIHTGEKPLQCEICGFTCRQKASLNWHMKKHDADSFYQFSCNICGKKFEKKDSVVAHKAKSHPEVLIAEALAANAGALITSTDILGTNPESLTQPSDGQGLPLLPEPLGNSTSGECLLLEAEGMSKSYCSGTERSIHR	nc
IP_098649.1	RP	INO80B-WBP1	MSKLWRRGSTSGAMEAPEPGEALELSLAGAHGHGVHKKKHKKHKKKHKKKHHQEEDAGPTQPSPAKPQLKLKIKLGGQVLGTKSVPTFTVIPEGPRSPSPLMVVDNEEEPMEGVPLEQYRAWLDEDSNLSPSPLRDLSGGLGGQEEEEEQRWLDALEKGELDDNGDLKKEINERLLTARQRALLQKARSQPSPMLPLPVAEGCPPPALTEEMLLKREERARKRRLQAARRAEEHKNQTIERLTKTAATSGRGGRGGARGERRGGRAAAPAPMVRYCSGAQGSTLSFPPGVPAPTAVSQRPSPSGPPPRCSVPGCPHPRRYACSRTGQALCSLQCYRINLQMRLGGPEGPGSPLLATFESCAQE	nc
IP_115174.1	RP	ZNF721	MYIGEFILERNPTHVENVAKPLDSLQIFMRIRKFILERNPTRVETVAKPLDSLQIFMHIRKFILEIKPYKCKECGKAFKSYYSILKHKRTHTRGMSYEGDECRGL	CDS
IP_275016.1	RP	ZNF780A	MNVRSVGKALIVVHTLFSIRKFIPMRNLLYVGNVRWPLDIIANLLNILEFILVTSHLNVKTVGRPSIVAQALFNIRVFTLVRSPMNVRSVGRLLDFTYNFPNIRKLTQVKNHLNVRNVGNSFVVVQILINIEVFILERNPLNVRNVGKPFDFICTLFDIRNCILVRNPLNVRSVGKPFDFICNLFDIRNCILVRNPLNVRNVERFLVFPPSLIAIRTFTQVRRHLECKECGKSFNRVSNHVQHQSIRAGVKPCECKGCGKGFICGSNVIQHQKIHSSEKLFVCKEWRTTFRYHYHLFNITKFTLVKNPLNVKNVERPSVF	CDS or 3'UTR
IP_278870.1	RP	ZNF845	MNVARFLIEKQNLHVIIEFILERNIRNMKNVTKFTVVNQVLKDRRIHTGEKAYKCKSL	CDS
IP_278888.1	RP	ZNF765	MSVARPSAGRHPLHTIIDFILDRNLTNVKIVMKLSVSNQTLKDIGEFILERNYTCNECGKTFNQELTLTCHRRLHSGEKPYKYEELDKAYNFKSNLEIHQKIRTEENLTSVMSVARP	CDS
IP_278918.1	RP	ZNF813	MNVARVLIGKHTLHVIIDFILERNLTSVMNVARFLIEKHTLHIIIDFILEINLTSVMNVARFLIKKHTLHVTIDFILERNLTSVMNVARFLIKKQTLHVIIDFILERNLTSLMSVAKLLIEKQSLHIIIQFILERNKCNECGKTFCHNSVLVIHKNSYWRETSVMNVAKFLINKHTFHVIIDFIVERNLRNVKHVTKFTVANRASKDRRIHTGEKAYKGEEYHRVFSHKSNLERHKINHTAEKP	CDS
IP_280349.1	RP	ZNF587	MNAVNVGNHFFPALRFMFIKEFILDKSLISAVNVENPFLNVPVSLNTGEFTLEKGLMNAPNVEKHFSEALPSFIIRVHTGERPYECSEYGKSFAEASRLVKHRRVHTGERPYECCQCGKHQNVCCPRS	CDS
IP_280385.1	RP	ZNF417	MNAMNVGNHFFPALRFMFIKEFILDKSLISAVNVENPLLNVPVSLNTGEFTLEKGLMNVPNVEKHFSEALPSFIIRVHTGERPYECSEYGKSFAETSRLIKHRRVHTGERPYECCQSGKHQNVCSPWS	CDS

5 [2] These two proteins were not detected with unique peptides but with shared peptides. One protein only was counted in subsequent
6 analyses.

7 [3] These five proteins were not detected with unique peptides but with shared peptides. One protein only was counted in subsequent
8 analyses.

1 **Table 4: Examples of proteins encoded in the same gene and functionally interacting**

Gene	Polypeptides[1]	Reference	altORF localization	altORF size aa	Conservation	Summary of functional relationship with the annotated protein
CDKN2A, INK4	Cyclin-dependent kinase inhibitor 2A or p16-INK4 (P42771), and p19ARF (Q8N726)	(61)	5'UTR	169	Unknown	the unitary inheritance of p16INK4a and p19ARF may underlie their dual requirement in cell cycle control.
GNAS, XLalphas	Guanine nucleotide-binding protein G(s) subunit alpha isoforms XL□s (Q5JWF2) and Alex (P84996)	(62)	5'UTR	+700	Human, mouse, rat	Both subunits transduce receptor signals into stimulation of adenylyl cyclase.
ATXN1	Ataxin-1 (P54253) and altAtaxin-1	(63)	CDS	185	Unknown	Direct interaction
Adora2A	A2A adenosine receptor (P30543) and uORF5	(64)	5'UTR	134	Human, chimpanzee, rat, mouse	A2AR stimulation increases the level of the uORF5 protein via post-transcriptional regulation.

AGTR1	Angiotensin type 1a receptor (P25095) and PEP7	(65)	5'UTR	7	Highly conserved across mammalian species	Inhibits non-G protein-coupled signalling of angiotensin II, without altering the classical G protein-coupled pathway activated by the ligand.

2   [1]The UniProtKB accession is indicated when available.

Figure 1

Figure 1. Annotation of human altORFs.
(**a**) AltORF nomenclature. AltORFs partially overlapping the CDS must be in a different reading
frame. (**b**) Pipeline for the identification of altORFs. (**c**) Size distribution of alternative (empty
bars, vertical and horizontal axes) and reference (grey bars, secondary horizontal and vertical
axes) proteins. Arrows indicate the median size. The median alternative protein length is 45
amino acids (AA) compared to 460 for the reference proteins. (**d**) Distribution of altORFs in the
human hg38 transcriptome. (**e, f**) Number of total altORFs (**e**) or number of altORFs/10kbs (**f**) in
hg38 compared to shuffled hg38. Means and standard deviations for 100 replicates obtained by
sequence shuffling are shown. Statistical significance was determined by using one sample t-test
with two-tailed *p*-values. **** *p*<0.0001. (**g**) Percentage of altORFs with an optimal Kozak motif.
The total number of altORFs with an optimal Kozak motif is also indicated at the top.

a

b

Figure 1-figure supplement 1: 10% of altORFs are present in different classes of repeats.

While more than half of the human genome is composed of repeated sequences, only 9.83% or 18,003 altORFs are located inside these repeats (**a**), compared to 2,45% or 1,677 CDSs (**b**). AltORFs and CDSs are detected in non-LTR retrotransposons (LINEs, SINEs, SINE-VNTR-Alus), LTR repeats, DNA repeats, satellites and other repeats. Proportions were determined using RepeatMasker (version 3.3.0).

Figure 1-figure supplement 2: The proportion of altORFs with a translation initiation site (TIS) with a Kozak motif in hg38 is significantly different from 100 shuffled hg38 transcriptomes.
Percentage of altORFs with a TIS within an optimal Kozak sequence in hg38 (dark blue) compared to 100 shuffled hg38 (light blue). Mean and standard deviations for sequence shuffling are displayed, and significant difference was defined by using one sample t test. **** *P*<0.0001. Note that shuffling all transcripts in the hg38 transcriptome generates a total of 489,073 altORFs on average, compared to 539,134 altORFs in hg38. Most transcripts result from alternative splicing and there are 183,191 unique altORFs in the hg38 transcriptome, while the 489,073 altORFs in shuffled transcriptomes are all unique. Figure 1g shows the percentage of unique altORFs with a kozak motif (15%), while the current Fig. shows the percentage of altORFs with a kozak motif relative to the total number of altORFs (14%).

Figure 2

a

b

c

Figure 2. Conservation of alternative and reference proteins across different species.
(**a**) Number of orthologous and paralogous alternative and reference proteins between *H. sapiens* and other species (pairwise study). (**b**) Phylogenetic tree: conservation of alternative (blue) and reference (red) proteins across various eukaryotic species. (**c**) Number and fraction of genes encoding homologous reference proteins or at least 1 homologous alternative protein between *H. sapiens* and other species (pairwise study).

Figure 3

Figure 3. Number of orthologous and co-conserved alternative and reference proteins between *H. sapiens* and other species (pairwise). For the co-conservation analyses, the percentage of observed (Obs.), expected (Exp.) and corresponding *p*-values relative to the total number of reference-alternative protein pairs are indicated on the right (see Table 2 for details).

Figure 4

Figure 4. AltORFs completely nested within CDSs show more extreme PhyloP values (more conserved or faster evolving) than their CDSs. Differences between altORF and CDS PhyloP scores (altORF PhyloP – CDS PhyloP, *y*-axis) are plotted against PhyloPs for their respective CDSs (*x*-axis). We restricted the analysis to altORF-CDS pairs that were co-conserved from humans to zebrafish. The plot contains 889 CDSs containing at least one fully nested altORF, paired with one of its altORFs selected at random (to avoid problems with statistical non-independence). PhyloPs for both altORFs and CDSs are based on 3rd codons in the CDS reading frame, calculated across 100 vertebrate species. We compared these differences to those generated based on five random regions in CDSs with a similar length as altORFs. Expected quantiles of the differences ("DQ" columns) were identified and compared to the observed differences. We show the absolute numbers ("n") and observed-to-expected ratios ("O/E") for each quantile. There are clearly substantial over-representations of extreme values (red signaling conservation DQ 0.95, and blue signaling accelerated evolution DQ 0.05) with 317 of 889 altORFs (35.7%). A random distribution would have implied a total of 10% (or 89) of altORFs in the extreme values. This suggests that 25.7% (35.7%-10%) of these 889 altORFs undergo specific selection different from random regions in their CDSs with a similar length distribution. This percentage is very similar to the 26.2% obtained from an analysis of altORFs without restriction based on co-conservation in vertebrates (see Figure 4-figure supplement 1), a total which would imply that there are about 4,458 altORFs fully nested in CDSs undergoing conserved or accelerated evolution relative to their CDSs.

Figure 4-figure supplement 1. AltORFs completely nested within CDSs show more extreme PhyloP values (more conserved or faster evolving) than their CDSs. Differences between altORF and CDS PhyloP scores (altORF PhyloP – CDS PhyloP, *y*-axis) are plotted against PhyloPs for their respective CDSs (*x*-axis). The plot contains all 20,814 CDSs containing at least one fully nested altORF, paired with one of its altORFs selected at random (to avoid problems with statistical non-independence). PhyloPs for both altORFs and CDSs are based on third codon positions in the CDS reading frame, calculated across 100 vertebrate species. We compared these differences to those generated based on five random regions in CDSs with a similar length as altORFs. Expected quantiles of the differences ("DQ" columns) were identified and compared to the observed differences. We show the absolute numbers ("n") and observed-to-expected ratios ("O/E") for each quantile. There are clearly substantial over-representations of extreme values (red signalling conservation DQ≥0.95, and blue signalling accelerated evolution DQ≤0.05) with 6,428 of 19,705 altORFs (36.2%). A random distribution would have implied a total of 10% (or 1,970) of altORFs in the extreme values. This suggests that 26.2% (36.2%-10%) of altORFs (or 4,458) undergo specific selection different from random regions in their CDSs with a similar length distribution.

Figure 5

Figure 5. First, second, and third codon nucleotide PhyloP scores for 100 vertebrate species for the CDSs of the *NTNG1*, *RET* and *VTI1A* genes. Chromosomal coordinates for the different CDSs and altORFs are indicated on the right. The regions highlighted in red indicate the presence of an altORF characterized by a region with elevated PhyloP scores for wobble nucleotides. The region of the altORF is indicated by a black bar above each graph.

Figure 6

Figure 6. Expression of human altORFs.
(**a**) Percentage of CDSs and altORFs with detected TISs by ribosomal profiling and footprinting of human cells[23]. The total number of CDSs and altORFs with a detected TIS is indicated at the top. (**b**) Alternative and reference proteins detected in three large proteomic datasets: human interactome[32], 10,000 human proteins[35], human phosphoproteome[34], EGFR interactome[33]. Numbers are indicates above each column. (**c**) Percentage of altORFs encoding alternative proteins detected by MS-based proteomics. The total number of altORFs is indicated at the top. Localization "Unknown" indicates that the detected peptides can match more than one alternative protein. Localization ">1" indicates that the altORF can have more than one localization in different RNA isoforms.

Figure 7

Figure 7. The alternative phosphoproteome in mitosis and EGF-treated cells. Heatmap showing relative levels of spectral counts for phosphorylated peptides following the indicated treatment[34]. For each condition, heatmap colors show the percentage of spectral count on total MS/MS phosphopeptide spectra. Blue bars on the right represent the number of MS/MS spectra; only proteins with spectral counts above 10 are shown.

AltLINC01420^{nc}

MGDQPCASGRSTLPPGNAREAKPPKKRCLLAPR**WDYPEGTPNGGSTTLPSAPPPASAGLK**SHPPPPEK

b

m/z = 917.09
[M+3H]$^{3+}$ = 2751.27 Da

c

m/z = 890.43.09
[M+3H]$^{3+}$ = 2671.29 Da

Figure 7-figure supplement 1: Example of a phosphorylated peptide in mitosis - alternative protein AltLINC01420[nc] (LOC550643, IP_305449.1).

(**a**) AltLINC01420[nc] amino acid sequence with detected peptides underlined and phosphorylated peptide in bold (73,9% sequence coverage). (**b**) MS/MS spectrum for the phosphorylated peptide (PeptideShaker graphic interface output). The phosphorylation site is the tyrosine residue, position 2. (**c**) MS/MS spectrum for the non-phosphorylated peptide. The mass difference between the precursor ions between both spectra corresponds to that of a phosphorylation, confirming the specific phosphorylation of this residue in mitosis.

Figure 8

Figure 8. Number of alternative proteins detected by ribosome profiling and mass spectrometry.
The expression of 467 alternative proteins was detected by both ribosome profiling (translation initiation sites, TIS) and mass spectrometry (MS).

Figure 9

Figure 9. Human alternative proteome sequence analysis and classification using InterProScan.
(**a**) InterPro annotation pipeline. (**b**) Alternative and reference proteins with InterPro signatures. (**c**) Number of alternative and reference proteins with transmembrane domains (TM), signal peptides (S) and both TM and SP. (**d**) Number of all alternative and reference proteins predicted to be intracellular, membrane, secreted and membrane-spanning and secreted. [1]Proteins with at least one InterPro signature; [2]Proteins with no predicted signal peptide or transmembrane features. (**e**) Number of predicted TM regions for alternative and reference proteins.

P. troglodytes

SP+TM: 75 SP: 4,872

Alternative proteins

TM: 11,931 16,878

SP+TM: 2,539 SP: 3,598

Reference proteins

TM: 8,090 14,227

M. musculus

SP+TM: 121 SP: 6,494

Alternative proteins

TM: 13,094 19,709

SP+TM: 2,485 SP: 6,228

Reference proteins

TM: 12,005 20,718

B. taurus

SP+TM: 29 SP: 2,267

Alternative proteins

TM: 5,013 7,309

SP+TM: 1,230 SP: 3,180

Reference proteins

TM: 5,746 10,156

D. melanogaster

SP+TM: 75 SP: 4,872

Alternative proteins

TM: 11,931 16,878

SP+TM: 2,539 SP: 3,598

Reference proteins

TM: 8,090 14,227

S. cerevisiae

SP+TM: 3 SP: 225

Alternative proteins

TM: 819 1,047

SP+TM: 91 SP: 251

Reference proteins

TM: 966 1,308

**Figure 9-figure supplement 1: Alternative proteome sequence analysis and classification in
P. troglodytes, *M. musculus*, *B. Taurus*, *D. melanogaster* and *S. cerevisiae*.**
For each organism, the number of InterPro signatures (top graphs) and proteins with
transmembrane (TM), signal peptide (SP), or TM+SP features (bottom pie charts) is indicated
for alternative and reference proteins.

Figure 10

Figure 10. Gene ontology (GO) annotations for human alternative proteins.
GO terms assigned to InterPro entries are grouped into 13 categories for each of the three
ontologies. (**a**) 34 GO terms were categorized into cellular component for 107 alternative proteins.
(**b**) 64 GO terms were categorized into biological process for 128 alternative proteins. (**c**) 94 GO
terms were categorized into molecular function for 302 alternative proteins. The majority of
alternative proteins with GO terms are predicted to be intracellular, to function in nucleic acid-
binding, catalytic activity and protein binding and to be involved in biosynthesis and nucleic acid
metabolism processes.

Figure 11

Figure 11. Main InterPro entries in human alternative proteins. (**a**) The top 10 InterPro families in the human alternative proteome. (**b**) A total of 110 alternative proteins have between 1 and 23 zinc finger domains.

Figure 12

Figure 12. Reference and alternative proteins share functional domains.
(**a**) Distribution of the number of shared InterPro entries between alternative and reference
proteins coded by the same transcripts. 138 pairs of alternative and reference proteins share
between 1 and 4 protein domains (InterPro entries). Only alternative/reference protein pairs that
have at least one domain are considered (n = 298). (**b**) The number of reference/alternative
protein pairs that share domains (n = 138) is higher than expected by chance alone. The
distribution of expected pairs sharing domains and the observed number are shown. (**c**) Matrix of
co-occurrence of domains related to zinc fingers. The entries correspond to the number of times
entries co-occur in reference and alternative proteins. The full matrix is available in figure 12-
figure supplement 1.

Figure 12-figure supplement 1: Matrix of co-occurrence of InterPro entries between alternative/reference protein pairs coded by the same transcript.
Pixels show the number of times entries co-occur in reference and alternative proteins. Blue pixels indicate that these domains are not shared, white pixels indicate that they are shared once, and red that they are shared twice or more.

Figure 12-figure supplement 2: Reference and alternative proteins share functional domains.
The number of reference/alternative protein pairs that share domains (n = 49) is higher than expected by chance alone (p<0.001). The distribution of expected pairs sharing domains and the observed number are shown. This is the same analysis as the one presented in figure 12b, with the zinc finger domains taken out.

Figure 13

Figure 13. AltMiD51⁵' expression induces mitochondrial fission.
(**a**) AltMiD51⁵' coding sequence is located in exon 2 or the *MiD51/MIEF1/SMCR7L* gene and in the 5'UTR of the canonical mRNA (RefSeq NM_019008). +2 and +1 indicate reading frames. AltMiD51 amino acid sequence is shown with the LYR tripeptide shown in bold. Underlined peptides were detected by MS. (**b**) Human HeLa cells transfected with empty vector (mock), a cDNA corresponding to the canonical MiD51 transcript with a Flag tag in frame with altMiD51 and an HA tag in frame with MiD51, altMiD51^Flag cDNA or MiD51^HA cDNA were lysed and analyzed by western blot with antibodies against Flag, HA or actin, as indicated. (**c**) Confocal microscopy of mock-transfected cells, cells transfected with altMiD51^WT, altMiD51^LYR→AAA or Drp1^K38A immunostained with anti-TOM20 (red channel) and anti-Flag (green channel) monoclonal antibodies. In each image, boxed areas are shown at higher magnification in the bottom right corner. % of cells with the most frequent morphology is indicated: mock (tubular), altMiD51^WT (fragmented), altMiD51(LYR→AAA) (tubular), Drp1(K38A) (elongated). Scale bar, 10 μm. (**d**) Bar graphs show mitochondrial morphologies in HeLa cells. Means of three independent experiments per condition are shown (100 cells for each independent experiment). ***$p<0.0005$ (Fisher's exact test) for the three morphologies between altMiD51(WT) and the other experimental conditions.

AltMiD51⁵′

188   397 499   1890

AltMiD51   MiD51

NM_019008.4

a. Experimental peptide

Intensity

NH2-Y T D R D F Y F A S I R-COOH

b. Synthetic peptide

Relative abundance

y₄⁺

y₅⁺

y₁⁺

y₃⁺

b₅⁺

b₆⁺

y₆⁺

b₇⁺

y₂⁺

y₇⁺

y₈⁺

YTDRDFYFASIR

m/z

Figure 13-figure supplement 1: Spectra validation for altMiD51.
Example of validation for altMiD51 specific peptides YTDRDFYFASIR and
GLVFLNGK. (**a,c**) Experimental MS/MS spectra (PeptideShaker graphic interface
output). (**b,d**) MS/MS spectra of the synthetic peptides.
Matching peaks are shown with blue masks. A diagram of the transcript with its accession
number and the localization of the altORF and the CDS is shown at the top.

Figure 13-figure supplement 2: MiD51 expression results in mitochondrial fission.
(**a**) Confocal microscopy of HeLa cells transfected with MiD51^{GFP} immunostained with anti-TOM20 (red channel) monoclonal antibodies. In each image, boxed areas are shown at higher magnification in the bottom right corner. The localization of MiD51 in fission sites is shown in merged higher magnification inset. Scale bar, 10 μm. (**b**) Human HeLa cells transfected with empty vector (mock) or MiD51^{GFP} were lysed and analyzed by western blot to confirm MiD51^{GFP} expression.

Figure 13-figure supplement 3: AltMiD51 is localized in the mitochondrial matrix.
Trypan blue quenching experiment performed on HeLa cells stably expressing the indicated
constructs. The fluorescence remaining after quenching by trypan blue is shown relative to
Matrix-Venus (Mx-Venus) indicated by the dashed line. (**** p <0,0001, one-way ANOVA).
The absence of quenching of the fluorescence compared to IMS-Venus indicates the matricial
localization of altMiD51. n≥3 cells were quantified per experiment, and results are from 6
independent experiments. Data are mean ±SEM.

Figure 13-figure supplement 4: Mitochondrial function parameters.
(**a**) Oxygen consumption rates (OCR) in HeLa cells transfected with empty vector (mock) or altMiD51[Flag]. Mitochondrial function parameters were assessed in basal conditions (basal), in the presence of oligomycin to inhibit the ATP synthase (oxygen consumption that is ATP-linked), FCCP to uncouple the mitochondrial inner membrane and allow for maximum electron flux through the respiratory chain (maximal OCR), and antimycin A/rotenone to inhibit complex III (non-mitochondrial). The balance of the basal OCR comprises oxygen consumption due to proton leak and nonmitochondrial sources. The mitochondrial reserve capacity (maximal OCR- basal OCR) is an indicator of rapid adaptation to stress and metabolic changes. Mean values of replicates are plotted with error bars corresponding to the 95% confidence intervals. Statistical significance was estimated using a two-way ANOVA with Tukey's post-hoc test (**$p = 0,004$). (**b**) ROS production in mock and altMiD51-expressing cells. Cells were untreated, treated with a ROS inducer or a ROS inhibitor. Results represent the mean value out of three independent experiments, with error bars corresponding to the standard error of the mean (s.e.m.). Statistical significance was estimated using unpaired T-test. (**c**) ATP synthesis rate in mock and altMiD51-expressing cells. No significant differences in ATP production were observed between mock and altMiD51 transfected cells.
Results represent the mean of three independent experiments (8 technical replicates each). Error bars represent the standard error of the mean. At the end of the experiments, cells were collected and proteins analyzed by western blot with antibodies against the Flag tag (altMiD51) or actin, as indicated, to verify the expression of altMiD51. A representative western blot is shown on the right. Molecular weight markers are shown on the left (kDa).

Figure 13-figure supplement 5: Representative confocal images of cells co-expressing altMiD51^{GFP} and Drp1(K38A)^{HA}.

(**a**) Confocal microscopy of HeLa cells co-transfected with altMiD51^{GFP} and Drp1(K38A)^{HA} immunostained with anti-TOM20 (blue channel) and anti-HA (red channel) monoclonal antibodies. In each image, boxed areas are shown at higher magnification in the bottom right corner. % of cells with the indicated morphology is indicated on the TOM20 panels. (**b**) Confocal microscopy of HeLa cells co-transfected with altMiD51^{GFP} and Drp1(wt)^{HA} immunostained with anti-TOM20 (blue channel) and anti-HA (red channel) monoclonal antibodies. In each image, boxed areas are shown at higher magnification in the bottom right corner. Scale bar, 10 μm.

Figure 13-figure supplement 6: Protein immunoblot showing the expression of different constructs in HeLa cells.

HeLa cells were transfected with empty vector (pcDNA3.1), altMiD51(WT)[Flag], altMID51(LYR→AAA)[Flag], Drp1(K38A)[HA], or Drp1(K38A)[HA] and altMiD51(WT)[Flag], as indicated. Proteins were extracted and analyzed by western blot with antibodies against the Flag tag (altMiD51), the HA tag (Drp1K38A) or actin, as indicated. Molecular weight markers are shown on the left (kDa). Representative experiment of three independent biological replicates.

Figure 14

Figure 14. AltMiD51-induced mitochondrial fragmentation is dependent on Drp1.
(**a**) Bar graphs show mitochondrial morphologies in HeLa cells treated with non-target or Drp1 siRNAs. Cells were mock-transfected (pcDNA3.1) or transfected with altMiD51^{Flag}. Means of three independent experiments per condition are shown (100 cells for each independent experiment). ***p<0.0005 (Fisher's exact test) for the three morphologies between altMiD51 and the other experimental conditions. (**b**) HeLa cells treated with non-target or Drp1 siRNA were transfected with empty vector (pcDNA3.1) or altMiD51^{Flag}, as indicated. Proteins were extracted and analyzed by western blot with antibodies against the Flag tag (altMiD51), Drp1 or actin, as indicated. (**c**) Confocal microscopy of Drp1 knockdown cells transfected with altMiD51^{GFP} immunostained with anti-TOM20 (blue channel) and anti-Drp1 (red channel) monoclonal antibodies. In each image, boxed areas are shown at higher magnification in the bottom right corner. % of cells with the indicated morphology is indicated on the TOM20 panels. Scale bar, 10 μm. (**d**) Control Drp1 immunostaining in HeLa cells treated with a non-target siRNA. For (**c**) and (**d**), laser parameters for Drp1 and TOM20 immunostaining were identical.

Figure 15

Figure 15. AltDDIT3⁵' co-localizes and interacts with DDIT3.
(**a**) AltDDIT3$^{5'}$ coding sequence is located in exons 1 and 2 or the *DDIT3/CHOP/GADD153* gene and in the 5'UTR of the canonical mRNA (RefSeq NM_004083.5). +2 and +1 indicate reading frames. AltDDIT3 amino acid sequence is also shown. (**b**) Confocal microscopy analyses of HeLa cells co-transfected with altDDIT3^{GFP} (green channel) and DDIT3^{mCherry} (red channel). Scale bar, 10 μm. (**c, d**) Colocalization analysis of the images shown in (**b**) performed using the JACoP plugin (Just Another Co-localization Plugin) implemented in Image J software (two independent biological replicates). (**c**) Scatterplot representing 50 % of green and red pixel intensities showing that altDDIT3^{GFP} and DDIT3^{mCherry} signal highly correlate (with Pearson correlation coefficient of 0.92 (*p*-value < 0.0001)). (**d**) Binary version of the image shown in (**b**) after Costes' automatic threshold. White pixels represent colocalization events (*p*-value < 0.001, based on 1000 rounds of Costes' randomization colocalization analysis). The associated Manders Correlation Coefficient, M_1 and M_2, are shown in the right upper corner. M_1 is the proportion of altDDIT3^{GFP} signal overlapping DDIT3^{mCherry} signal and M_2 is the proportion of DDIT3^{mCherry} signal overlapping altDDIT3^{GFP}. (**e**) Representative immunoblot of co-immunoprecipitation with GFP-Trap agarose beads performed on HeLa lysates co-expressing DDIT3^{mcherry} and altDDIT3^{GFP} or DDIT3^{mcherry} with pcDNA3.1^{GFP} empty vector (two independent experiments).

Figure 15-figure supplement 1: Protein immunoblot showing the expression of different constructs in HeLa cells.
HeLa cells were co-transfected with GFP and mCherry, or altDDIT3^{GFP} and DDIT3^{mCherry}, as indicated. Proteins were extracted and analyzed by western blot with antibodies, as indicated. Molecular weight markers are shown on the left (kDa). AltDDIT3 has a predicted molecular weight of 4.28 kDa and thus migrates at its expected molecular weight when tagged with GFP (~32 kDa). Representative experiment of two independent biological replicates.

Figure 15–figure supplement 2 : Colocalization of altDDIT3 with DDIT3.
Scatter plots of Pearson's Correlation Coefficient and Manders' Correlation Coefficient after Costes' automatic threshold (p-value < 0.001, based on 1000 rounds of Costes' randomization colocalization analysis). M1 is the proportion of altDDIT3[GFP] signal overlapping DDIT3[mCherry] signal and M2 is the proportion of DDIT3[mCherry] signal overlapping altDDIT3[GFP]. Error bars represent the mean +/- SD of three independent experiments (28 cells).