**Reviving a Lost Species: The Case of the Floreana Galápagos Giant Tortoise *Chelonoidis elephantopus***

Joshua M. Miller[1†]*, Maud C. Quinzin[1†], Nikos Poulakakis[2,3], James P. Gibbs[4], Luciano B. Beheregaray[5], Ryan C. Garrick[6], Michael A. Russello[7], Claudio Ciofi[8], Danielle L. Edwards[9], Elizabeth A. Hunter[10], Washington Tapia[11,12], Danny Rueda[12], Jorge Carrión[12], Andrés A. Valdivieso[1], Adalgisa Caccone[1]*

1. Department of Ecology and Evolutionary Biology, Yale University, 21 Sachem St. New Haven, Connecticut, 06520, United States of America
2. Department of Biology, School of Sciences and Engineering, University of Crete, Vasilika Vouton, Gr-71300, Heraklio, Crete, Greece
3. Natural History Museum of Crete, School of Sciences and Engineering, University of Crete, Knossos Av., GR-71409, Heraklio, Crete, Greece
4. College of Environmental Science & Forestry, State University of New York, Syracuse, New York, 13210, United States of America
5. Molecular Ecology Lab, School of Biological Sciences, Flinders University, GPO Box 2100, Adelaide, SA, 5001, Australia
6. Department of Biology, University of Mississippi, Oxford, Mississippi, 38677, United States of America
7. Department of Biology, University of British Columbia, Okanagan Campus, Kelowna, BC V1V 1V7, Canada
8. Department Biology, University of Florence, 50019 Sesto Fiorentino (FI), Italy
9. Life and Environmental Sciences, University of California, Merced, 5200 N Lake Rd, Merced, California, 95343, United States of America
10. Department of Natural Resources and Environmental Science, University of Nevada – Reno, Max Fleischmann Agricultural Building, Reno NV 89557 USA
11. Galapagos Conservancy, Fairfax, Virginia, 22030, United States of America
12. Galápagos National Park Directorate, Puerto Ayora, Galápagos, Ecuador

*Corresponding authors: joshua.miller@yale.edu, adalgisa.caccone@yale.edu

†These authors contributed equally to this work

**Abstract**

Species are being lost at an unprecedented rate due to human-driven environmental changes. The cases in which species declared extinct can be revived are extremely rare. However, here we report that a remote volcano in the Galápagos Islands hosts many giant tortoises with extremely high ancestry from a species previously declared as extinct: *Chelonoidis elephantopus* or the Floreana tortoise. Of 150 individuals with distinctive morphology sampled from the volcano, genetic analyses revealed that 65 had *C. elephantopus* ancestry Thirty-two were translocated from the volcano's slopes to a captive breeding center. A genetically informed captive breeding program now being initiated will, over the next decades, return *C. elephantopu*s tortoises to Floreana Island to serve as engineers of the island's ecosystems. Ironically, it was the haphazard translocations by mariners killing tortoises for food centuries ago that created the unique opportunity to revive this "lost" species today.

**Introduction**

Human activities have generated an extreme and rapid loss of biodiversity[1]. Many actions have been undertaken to prevent species extinctions, including creating laws to protect endangered species and critical habitats[2], translocations of individuals among populations or into new habitats[3,4], and *ex situ* management including captive breeding[5]. For many species, such actions come too late to facilitate recovery[6–8]. Although 'de-extinction' using laboratory techniques is currently being debated[9], such methods are viable only for some taxa and will generate significant anticipated, and unanticipated, risks[10,11]. Generally, extinction is final and cases where lost species can be revived will be extremely rare.

Despite their insularity, even remote oceanic islands are not exempt from rapid anthropogenic changes. For example, the ecosystems of the Galápagos Islands, located ~900 kilometers off the Pacific coast of Ecuador, have been degraded by human activities since the archipelago was discovered in 1535. The resulting loss of biodiversity, including many endemic species, has been chiefly due to the introduction of non-native species[12,13]. In response, concerted efforts have been taken to restore ecosystems on the islands, including removal of introduced pests[14,15] and population restoration via captive breeding and repatriation of threatened native species (e.g. [16,17]).

Galápagos giant tortoises (*Chelonoidis* spp.) are flagship species for ongoing restoration
efforts[18] in this archipelago and play an important functional role as mega-herbivores in the
islands' ecosystems[19]. Galápagos giant tortoises can be classified into 15 species based on
genetic data[20]. Generally, there is a single species per island with two exceptions: Isabela Island
that has a different endemic species associated with each of its five volcanoes, and Santa Cruz
Island that contains two species, one recently described[20] (Figure 1a). The 15 species exhibit two
general carapace shapes: "domed," a rounded cupola-like form (Figure 1c), and a "saddle-
backed" form, with a high anterior opening creating the shape of a saddle (Figure 1d). Five of the
15 species have the saddle-backed morphology: *C. elephantopus* (Floreana), *C. hoodensis*
(Española), *C. abingdoni* (Pinta), *C. ephippium* (Pinzón), and *C. chathamensis* (San Cristóbal;
Figure 1a).

Over the past three centuries, all giant tortoise populations experienced a ~90% decline[21],
having been killed mostly for food and oil by whalers, sealers, buccaneers, and early
colonists[22,23]. Four species have been declared Extinct[24], including two of the five saddle-backed
species: *C. elephantopus* from Floreana Island and, most recently, *C. abingdoni,* from Pinta
Island. The latter species was represented by a single individual, Lonesome George, until his
death in 2012.

Surprisingly, recent research found living, wild tortoises with genetic ancestry from two
of the extinct saddle-backed species, *C. elephantopus* and *C. abingdoni* (hereafter referred to as
the Floreana and Pinta tortoises, respectively) outside their native range[25–28]. These individuals,
likely the descendants of tortoises translocated among islands by mariners[22,23,29], were discovered
on the remote Volcano Wolf on Isabela Island (Figure 1a,b). Among >1600 tortoises sampled
from Volcano Wolf in 2008 during an exploratory expedition, 105 individuals were admixed
between the locally endemic species, *C. becki*, and the Floreana ($n = 86$)[5] or Pinta ($n = 17$)[4]
species. The majority of these genetically admixed individuals were found on the Volcano
Wolf's western slopes, facing Banks Bay (also known as Puerto Bravo, PBR: Figure 1b), with a
smaller number located on the volcano's northwestern slopes near Piedras Blancas (PBL)[25,26].
No purebred individuals of either of the two non-native species were found in 2008, but genetic
simulations and the young age of some mixed ancestry individuals indicated that purebred
Floreana and Pinta tortoises might still be present on Volcano Wolf[25,26].

Here we build on this previous work. In November 2015 we mounted a 10-day-long

search involving ~70 field personnel combined with helicopter- and ship-support. Focusing our

search, we explored zones of Volcano Wolf most likely to contain individuals with ancestry from

these two "lost" species (Figure 1b) and restricted genetic sampling to those individuals with

saddle-backed morphology among the thousands of locally endemic domed *C. becki*. We then

assigned ancestry to these tortoises using reference databases containing both extant and extinct

species that have previously been used to assign ancestry to tortoises in the wild and captivity[25–

31]. Based on these assignments, we determined suitability of the tortoises for a genetically

informed captive breeding program aimed at reintroducing these key ecosystem engineers to

their native islands.

**Results & Discussion**

In total, we encountered 144 individuals with saddle-backed morphology. Of those, 112

were released after taking blood samples, and 32 with pronounced saddle-backed morphology

were transported to the Galápagos National Park Service's captive tortoise breeding facility on

Santa Cruz Island[32–34]. We assigned ancestry to all 144 of these individuals along with six

saddle-backed tortoises known to have Floreana ancestry[35] already residing at the breeding

center using information from ~700-bp of mitochondrial DNA sequence and diploid genotypes

from 12 nuclear microsatellite loci. These loci have previously been shown to accurately assign

individuals to tortoise species[25–31].

Thirty-five of the 150 individuals analyzed had a mitochondrial DNA haplotype

diagnostic of the Floreana species (Supplementary Figure S1). The remaining individuals either

had haplotypes diagnostic of the Critically Endangered Española Island species ($n$ = 70),

haplotypes shared between the Santiago Island and Volcano Wolf species ($n$ = 44), or a

haplotype shared between the species from San Cristóbal and Santa Cruz Islands ($n$ = 1). The

proportions of haplotypes associated with the Española and Floreana Island species reported here

(46% and 23%, respectively) are substantially higher than previously detected among individuals

from Volcano Wolf[25,26] (5% and 2%). This is likely due to our targeting of saddle-backed

individuals in 2015 versus sampling broadly during previous surveys[25–28].

Bayesian clustering analyses of microsatellite genotypic data using the method

implemented in STRUCTURE[37] revealed that 127 of the 150 tortoises sampled have ancestry

assignments (Q-values) to the extinct Floreana species (average Q-value ± SD = 0.87 ± 0.21;

range 0.16–0.99; Figure 2). Twenty-three individuals did not show evidence of Floreana

ancestry, being assigned to the two genetically distinct populations (PBL and PBR) of *C. becki*,

the endemic Volcano Wolf species. Of those individuals with Floreana Q-values 30 had Floreana

mitochondrial haplotypes (Figure 2).

We conducted additional assignment tests using genotypes generated by simulated

matings within and among four possible parental lineages (PBL, PBR, Española Island, and

Floreana Island). These identified two tortoises with strong assignment to the Floreana species,

being classified as either a purebred Floreana tortoise or a backcross between a $F_1$ × purebred

Floreana tortoise (Figure 3, Supplementary Table S2). An additional 63 tortoises were assigned

to categories with Floreana ancestry (Table 1, Q-values between 0.40–0.77). Admixture between

the saddle-backed species from Floreana and Española Islands (*n* = 43) was more common than

between Floreana and either of the two endemic domed *C. becki* populations (PBL and PBR; *n* =

22), indicative of positive assortative mating between the two saddle-backed species.

Neither mitochondrial DNA nor microsatellite data identified individuals with ancestry

from the Pinta Island species. However, a previous genetic estimate suggested that only 60–70

tortoises with Pinta ancestry are present on Volcano Wolf[25], whereas capture-mark-recapture

methods employed during our expedition estimated that a total of ~5,000–6,000 tortoises

occurred in the area searched. Moreover, despite our substantial search effort, we explored only

~26% of the total tortoise-occupied range on the volcano (J. Gibbs unpublished data). Therefore,

it is possible that individuals with Pinta ancestry still live on Volcano Wolf, but went undetected

in 2015.

Of the 38 individuals currently housed in the captive breeding center, 23 tortoises (9

males and 14 females) were found to have nuclear ancestry from Floreana across multiple

assignment methods (Table 1, Q-values between 0.44–0.77). In addition, 12 of these 23

individuals have mitochondrial haplotypes from Floreana (Supplementary Table S2). Together,

these 23 individuals now form the core of a genetically-informed captive breeding program

aimed at repatriation of tortoises to Floreana Island. For the purposes of the breeding program,

all 23 individuals will be included, regardless of mitochondrial lineage, in order to capture and

maintain as much nuclear genetic diversity from Floreana as possible. The program is modeled

after another one developed for the Española species initiated with only 15 founders. Over 50

years, the Española tortoise program generated >2000 repatriates with nearly 1,000 surviving

tortoises now reproducing independently on their native island[17,38,39].

A key attribute for success of such breeding programs is that the founding individuals are

unrelated, as high relatedness can lead to inbreeding depression. Accordingly, we examined

relatedness among the 23 individuals with Floreana ancestry in the breeding center. The analysis

showed that most individuals were unrelated (average Queller and Goodnight's[40] relatedness = -

0.04, range -0.58–0.63; Supplementary Figure 3). Although the 23 tortoises in captivity represent

a promising founding population, 44 tortoises found during the 2015 expedition, but left on

Volcano Wolf, are also good candidates for the Floreana breeding and repatriation program

(Table 1). These individuals, once re-located on Volcano Wolf, could be incorporated into the

breeding program to further expand the genetic diversity of the founder population.

The Floreana tortoise breeding program will be designed to maximize founder

contributions and *C. elephantopus* genome representation in the resulting progeny, while

promoting *in situ* population growth and minimizing costs to the Galápagos National Park

Directorate. Depending on the goals and priorities of Park decision-makers, complete genome

recovery may not be reached before releases of offspring begin. However, the high proportion of

Floreana ancestry and low relatedness evident in the current breeding individuals indicate that

the 23 founding individuals and their resulting progeny will provide a good starting point for

restoring the species.

Our discovery raises the possibility that the extinct Floreana species could be revived. In

this case, tortoises with Floreana ancestry are living 'genomic archives' that retain the

evolutionary legacy of the extinct species, removing the need for the cloning methods that have

been proposed to bring back extinct species[41]. The Floreana tortoise breeding program is

anticipated to generate thousands of offspring over the next few decades. When repatriated to

Floreana Island, these tortoises can once again play their critical role as ecosystem engineers[19].

In addition, giant tortoises are a major tourist attraction in Galápagos[42]; tortoise restoration on

Floreana Island should create new economic opportunities for the island's few human residents.

Ironically, the opportunity to revive this "lost species" today was created by the same early

visitors to the archipelago whose activities imperiled most giant tortoise species and drove some

into outright extinction.

## Materials and Methods

### Sampling and lab methods

All samples were collected under CITES permit 15US209142/9, Galápagos Park Permit

PC-75-16, and in accordance with Yale Institutional Animal Care and Use Committee (IACUC)

permit number 2016-10825. Samples were collected over a 10-day expedition in November 2015

by a team of 36 researchers plus Galápagos National Park rangers. Groups of 3–4 searchers were

assigned distinct areas on Volcano Wolf, collectively totaling ~36 km$^2$. The search area was

chosen based on a previous survey of tortoises on Volcano Wolf, which indicated that

individuals with Pinta and Floreana ancestry were found to be in their highest densities on the

western slopes of Volcano Wolf[25]. For all tortoises encountered, sex, age, and GPS coordinates

were recorded; for saddle-backed individuals, photographs were taken, blood samples were

collected for DNA analysis, and each individual was injected with a passive integrated

transponder (PIT) tag under the skin for identification in the future. In cases where blood was

taken, ~2 ml of blood was collected from the brachial vein of one of the front legs of the tortoise

and preserved in a lysis buffer containing 0.1 M Tris buffer, 0.1 M EDTA, 0.2 m NaCl, and 1%

SDS, pH 8.0. All tortoises were uniquely marked with paint when first encountered. Starting on

the sixth day, we re-searched areas and recorded whether any individual had been previously

encountered for the purposes of a capture-mark-recapture estimate of population size. In total,

1,333 tortoises were encountered, of which 144 had saddle-backed morphology. The 32 tortoises

removed from Volcano Wolf were initially carried in nets by helicopter from the flanks of the

volcano to the expedition ship anchored in Puerto Bravo harbor, and then transported to the

Galápagos National Park captive tortoise breeding facility on Santa Cruz Island.

DNA was extracted from 150 blood samples using Qiagen blood and tissue extraction

kits. These samples included the 144 saddle-backed individuals mentioned above along with six

individuals already housed at the Galápagos National Park Breeding Center on Santa Cruz Island

that were previously identified to have Floreana ancestry[35]. All samples were sequenced at

~700bp of the mitochondrial DNA control and were genotyped at 12 dinucleotide

microsatellites, using previously developed protocols (detailed procedures in Supplementary

Methods).

### Ancestry assignment

For ancestry assignment based on mitochondrial DNA, the new sequences were aligned to a reference database of 123 previously observed haplotypes, representing all extant and extinct species[29]. Ancestry was assigned by determining shared haplotypes using the program TCS version 1.21[43]. Evolutionary relationships were viewed with the program Network version 5.0 (fluxus-engineering.com; Supplementary Figure 1).

For ancestry assignment based on microsatellite genotypic data, we took a two-step process using STRUCTURE version 2.3.4[37,44]. First, we ran STRUCTURE with a reference dataset of 277 samples including all extant and extinct species to confirm the number of expected genetic clusters ($K$) present in the archipelago (for full parameters see Supplementary Methods). In this case, the optimal $K$ was 12, which corresponds to previously described results[25,26] (Supplementary Figure 2). Second, with archipelago-wide $K$ established, we reduced the reference dataset to the seven clusters previously found to be on Volcano Wolf[29]: Española, San Cristóbal, Central Isabela (La Cazuela, Volcano Alcedo, and Volcano Darwin), Floreana, Pinta, and the two genetically distinct endemic Volcano Wolf populations, PBL, and PBR. To this reduced dataset (155 reference samples in total), the new samples were added and we set $K$ to 7, leaving all other parameters unchanged.

To further quantify the genetic ancestry of individuals, assignment tests were undertaken using three additional methods. Prior to implementing these methods, we expanded the reference database to include simulated genotypes. These genotypes were simulated using HYBRIDLAB version 1.0[45], and corresponded to individuals arising from crosses within and among the genetically distinct *C. becki* populations on Volcano Wolf (PBL and PBR), and the two saddle-backed species from Española and Floreana (see Supplementary Methods). These four lineages were chosen due to their high prevalence in the mitochondrial DNA-based assignments and their contribution to microsatellite genotypes in admixed individuals. Simulated individuals represented explicit ancestry categories: pure parental populations, first-generation ($F_1$) crosses, second-generation ($F_2$) crosses, and backcrosses (i.e., matings between $F_1$'s and their respective purebred lineages).

With these new genotypes added into the reference set, we first used GeneClass version 2.0[46] to calculate the probability that an individual's genotype assigns to a population. Second, we used NEWHYBRIDS version 1.1[47] to compute the posterior probability of various hybrid classes for each individual. In this case, only four pairwise combinations of parental populations

were considered to focus on identifying potential hybrids involving Floreana ancestry: 1)

Española × Floreana, 2) PBL × Floreana, 3) Floreana × PBR, and 4) Española × PBR. Third, we

assigned ancestry using a multivariate approach, discriminant analysis of principal components

(DAPC), as implemented in the R package *adegenet* version 2.0.1[48,49]. In this case, the simulated

genotypes from crosses among Floreana, Española, PBL and PBR were initially used as a

training dataset to define the principal components and discriminant functions. The empirical

genotypes for putative hybrids were then transformed with principal components analysis (PCA)

based on the centering and scaling of the training set, and positioned onto the discriminant

functions. The NEWHYBRIDS and DAPC analyses were carried out only on individuals with

ancestry from Floreana based on the results from the GeneClass analyses (Supplementary Table

S2).

**Relatedness Analysis**

For the 23 individuals in the breeding center with Floreana ancestry, we calculated

pairwise relatedness using the estimator of Queller and Goodnight[40]. This was the highest ranked

of eight tested relatedness estimators tested by the program irelr[50] which considers a composite

score that incorporates estimates of bias, variance, skewness, and kurtosis. The empirical

distribution of relatedness values was compared to the distributions of 10,000 simulated pairs of

individuals for each of four relatedness categories (unrelated, half sibs, full sibs, and parent–

offspring). The pairwise relatedness distribution of both the empirical and simulated data were

calculated using irelr[50].

**<u>References</u>**

1.     Dirzo, R. *et al.* Defaunation in the Anthropocene. *Science (80-. )*. **345,** 401 LP-406 (2014).

2.     Novacek, M. J. & Cleland, E. E. The current biodiversity extinction event: Scenarios for

mitigation and recovery. *Proc. Natl. Acad. Sci. U. S. A.* **98,** 5466–5470 (2001).

3.     Corlett, R. T. Restoration, Reintroduction, and Rewilding in a Changing World. *Trends

Ecol. Evol.* **31,** 453–462 (2016).

4.     Weeks, A. R. *et al.* Assessing the benefits and risks of translocations in changing

environments: a genetic perspective. *Evol. Appl.* **4,** 709–725 (2011).

5.     Canessa, S. *et al.* Planning for ex-situ conservation in the face of uncertainty. *Conserv.

Biol.* **30,** 599–609 (2016).

6.     Hoffmann, M. *et al.* The Impact of Conservation on the Status of the World's Vertebrates.

*Science (80-. ).* **330,** 1503–1509 (2010).

7.     Doremus, H. & Pagel, J. E. Why Listing May Be Forever: Perspectives on Delisting under

the U.S. Endangered Species Act. *Conserv. Biol.* **15,** 1258–1268 (2001).

8.     Taylor, M. F. J., Suckling, K. F. & Rachlinski, J. J. The Effectiveness of the Endangered

Species Act: A Quantitative Analysis. *Bioscience* **55,** 360–367 (2005).

9.     Sherkow, J. S. & Greely, H. T. What If Extinction Is Not Forever? *Science (80-. ).* **340,**

32–33 (2013).

10.    Cohen, S. The Ethics of De-Extinction. *Nanoethics* **8,** 165–178 (2014).

11.    Jones, K. E. From dinosaurs to dodos: who could and should we de-extinct? *Front.*

*Biogeogr.* **6,** (2014).

12.    Phillips, R. B., Wiedenfeld, D. A. & Snell, H. L. Current status of alien vertebrates in the

Galápagos Islands: invasion history, distribution, and potential impacts. *Biol. Invasions*

**14,** 461–480 (2012).

13.    Guézou, A. *et al.* An Extensive Alien Plant Inventory from the Inhabited Areas of

Galapagos. *PLoS One* **5,** e10276 (2010).

14.    Harper, G. A. & Carrion, V. Introduced rodents in the Galapagos: colonisation, removal

and the future. *Isl. Invasives Erad. Manag.* 63–66 (2011).

15.    Campbell, K., Donlan, C. J., Cruz, F. & Carrion, V. Eradication of feral goats Capra

hircus from Pinta Island, Galápagos, Ecuador. *Oryx* **38,** 328–333 (2004).

16.    Fabiani, A. *et al.* Conservation of Galápagos land iguanas: genetic monitoring and

predictions of a long-term program on the island of Santa Cruz. *Anim. Conserv.* **14,** 419–

429 (2011).

17.    Milinkovitch, M. C. *et al.* Genetic analysis of a successful repatriation programme: giant

Galapagos tortoises. *Proc. R. Soc. B-Biological Sci.* **271,** 341–345 (2004).

18.    Benitez-Capistros, F., Hugé, J., Dahdouh-Guebas, F. & Koedam, N. Exploring

conservation discourses in the Galapagos Islands: A case study of the Galapagos giant

tortoises. *Ambio* **45,** 706–724 (2016).

19.    Gibbs, J. P., Sterling, E. J. & Zabala, F. J. Giant Tortoises as Ecological Engineers: A

Long-term Quasi-experiment in the Galápagos Islands. *Biotropica* **42,** 208–214 (2010).

20.    Poulakakis, N. *et al.* Description of a New Galapagos Giant Tortoise Species

308           (Chelonoidis; Testudines: Testudinidae) from Cerro Fatal on Santa Cruz Island. *PLoS One*
309           **10,** e0138779 (2015).

310   21.   Márquez, C. *et al.* Estado actual de las poblaciones de tortugas terrestres gigantes
311           (Geochelone spp., Chelonia: Testudinae) en las islas Galápagos. *Ecol. Apl.* **3,** (2004).

312   22.   Porter, D. *Journal of a cruise made to the Pacific Ocean*. (Wiley & Halsted, 1822).

313   23.   Townsend, C. H. The Galápagos tortoises in relation to the whaling industry: A study of
314           old logbooks. *Zoologica* **4,** 55–135 (1925).

315   24.   The IUCN Red List of Threatened Species. Version 2016-2. (2016). Available at:
316           www.iucnredlist.org. (Accessed: 10th November 2016)

317   25.   Edwards, D. L. *et al.* The genetic legacy of Lonesome George survives: Giant tortoises
318           with Pinta Island ancestry identified in Galapagos. *Biol. Conserv.* **157,** 225–228 (2013).

319   26.   Garrick, R. C. *et al.* Genetic rediscovery of an 'extinct' Galapagos giant tortoise species.
320           *Curr. Biol.* **22,** R10–R11 (2012).

321   27.   Poulakakis, N. *et al.* Historical DNA analysis reveals living descendants of an extinct
322           species of Galapagos tortoise. *Proc. Natl. Acad. Sci. U. S. A.* **105,** 15464–15469 (2008).

323   28.   Russello, M. A. *et al.* Lonesome George is not alone among Galápagos tortoises. *Curr.*
324           *Biol.* **17,** R317–R318 (2007).

325   29.   Poulakakis, N., Russello, M., Geist, D. & Caccone, A. Unravelling the peculiarities of
326           island life: vicariance, dispersal and the diversification of the extinct and extant giant
327           Galapagos tortoises. *Mol. Ecol.* **21,** 160–173 (2012).

328   30.   Benavides, E. *et al.* Lineage Identification and Genealogical Relationships Among
329           Captive Galapagos Tortoises. *Zoo Biol.* **31,** 107–120 (2012).

330   31.   Garrick, R. C. *et al.* Lineage fusion in Galapagos giant tortoises. *Mol. Ecol.* **23,** 5276–
331           5290 (2014).

332   32.   The Galapagos Conservancy. The Wolf Expedition: Resurrecting the extinct tortoise
333           species from Floreana and Pinta. (2015). Available at:
334           http://www.galapagos.org/newsroom/wolf-tortoise-search-outcome/.

335   33.   Nicholls, H. Lonesome George 'look-alike' captured for tortoise-breeding effort. *Nature*
336           (2015). doi:10.1038/nature.2015.18955

337   34.   Blakeslee, S. Scientists Hope to Bring a Galápagos Tortoise Species Back to Life. *The*
338           *New York Times* (2015).

339    35.    Russello, M. A. *et al.* DNA from the Past Informs Ex Situ Conservation for the Future: An
340            "Extinct'' Species of Galapagos Tortoise Identified in Captivity. *PLoS One* **5,** (2010).

341    36.    McMahon, B. J., Teeling, E. C. & Höglund, J. How and why should we implement
342            genomics into conservation? *Evol. Appl.* **7,** 999–1007 (2014).

343    37.    Falush, D., Stephens, M. & Pritchard, J. K. Inference of population structure using
344            multilocus genotype data: Linked loci and correlated allele frequencies. *Genetics* **164,**
345            1567–1587 (2003).

346    38.    Milinkovitch, M. *et al.* Giant Galapagos tortoises; molecular genetic analyses identify a
347            trans-island hybrid in a repatriation program of an endangered taxon. *BMC Ecol.* **7,**
348            doi:10.1186/1472-6785-7-2 (2007).

349    39.    Gibbs, J. P., Hunter, E. A., Shoemaker, K. T., Tapia, W. H. & Cayot, L. J. Demographic
350            Outcomes and Ecosystem Implications of Giant Tortoise Reintroduction to Española
351            Island, Galapagos. *PLoS One* **9,** e110742 (2014).

352    40.    Queller, D. C. & Goodnight, K. F. Estimating relatedness using genetic-markers.
353            *Evolution (N. Y)*. **43,** 258–275 (1989).

354    41.    Shapiro, B. Mammoth 2.0: will genome engineering resurrect extinct species? *Genome*
355            *Biol.* **16,** 228 (2015).

356    42.    Medina, C., Aravena, C. & Vásquez, F. *Economic valuation of the giant tortoise*
357            *conservation in Galapagos Islands Carlos*. (2012).

358    43.    Clement, M., Posada, D. & Crandall, K. A. TCS: a computer program to estimate gene
359            genealogies. *Mol. Ecol.* **9,** 1657–1659 (2000).

360    44.    Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using
361            multilocus genotype data. *Genetics* **155,** 945–959 (2000).

362    45.    Nielsen, E. E., Bach, L. A. & Kotlicki, P. HYBRIDLAB (version 1.0): a program for
363            generating simulated hybrids from population samples. *Mol. Ecol. Notes* **6,** 971–973
364            (2006).

365    46.    Piry, S. *et al.* GENECLASS2: A Software for Genetic Assignment and First-Generation
366            Migrant Detection. *J. Hered.* **95,** 536–539 (2004).

367    47.    Anderson, E. C. & Thompson, E. A. A model-based method for identifying species
368            hybrids using multilocus genetic data. *Genetics* **160,** 1217–1229 (2002).

48.    Jombart, T. adegenet: a R package for the multivariate analysis of genetic markers.

*Bioinformatics* **24,** 1403–1405 (2008).

49.   Jombart, T. & Ahmed, I. adegenet 1.3-1: new tools for the analysis of genome-wide SNP

data. *Bioinformatics* **27,** 3070–3071 (2011).

50.   Gonçalves da Silva, A. & Russello, M. A. iRel: software for implementing pairwise

relatedness estimators and evaluating their performance. *Conserv. Genet. Resour.* **3,** 69–

71 (2011).

51.   Team, R. C. R: A language and environment for statistical computing, reference index

version 3.2.1. (2014).

52.   Hijmas, R. J. raster: Geographic Data Analysis and Modeling version 2.5-8. (2016).

**Acknowledgements**

This work was enabled by substantial logistical and personnel support of the Galápagos National

Park Directorate through its dedication to science-based conservation of giant tortoises via the

"Giant Tortoise Restoration Initiative" undertaken in conjunction with the Galapagos

Conservancy. Expedition expenses and laboratory analyses were also supported by grants from

the Galapagos Conservancy, the Mohamed bin Zayed Species Conservation Fund, National

Geographic Society, and the Oak Foundation to AC, the US National Science Foundation (DEB

1258062) to JG, the Belgian American Educational Foundation to MQ, and the Yale Institute for

Biospheric Studies. C. Gottsegen helped with genetic data collection, A. Antoniou provided

assistance with DAPC analyses. L. Boitani, L. Cayot, C. Parker, and A. Rhodin helped procure

funds and provided encouragement. J. Flanagan provided critical veterinary support.

**Author Contributions**

JMM, NP, LB, CC, EAH, WT, DR, JC, JPG and AC assisted with field logistics and sample

collection. JMM, MCQ, and AV conducted laboratory analyses. JMM, MCQ, and NP conducted

data analyses. LBB, RCG, MAR, and DLE provided technical guidance and conceptual advice

for data analyses. JMM drafted the manuscript with assistance from MCQ, LBB, MAR, JPG, and

AC. JMM, MCQ, NP, LBB, RCG, MAR, CC, DLE, EAH, WT, JPG, and AC discussed the

results and implications and commented on the manuscript at all stages.

**Additional Information**

Microsatellite genotypes and mitochondrial sequences are available on a server hosted by the

College of Environmental Science & Forestry, State University of New York, Syracuse

(http://www.esf.edu/efb/gibbs/Miller_et_al_data_archive.zip). The authors have no financial

conflicts of interest. Correspondence and requests for materials should be addressed to either

joshua.miller@yale.edu or adalgisa.caccone@yale.edu.

**Figures**

Figure 1: Distribution of tortoises among Galápagos Islands and representative photos of tortoise
carapace morphology. A) Map of the distribution of tortoises among Galápagos Islands
along with cartoons indicating carapace morphology for each. Light grey shading
indicates domed morphology, unshaded indicates saddle-backed. Extinct species are noted
with †. B) Larger view of Volcano Wolf on northern Isabela Island. The circle indicates
the approximate field location of the current study. Examples of Galápagos giant tortoises
with domed (C) saddle-backed (D) morphology. Maps were created with R (version
3.2.4[51]) using the raster package (version 2.5-8[52]).

Figure 2: STRUCTURE[37,44] plot for 155 reference samples along with the 150 newly collected individuals. Each individual is represented as a vertical bar, with colors denoting the different genetic clusters, as indicted below. The proportion of color in a bar is equal to the ancestry (Q-value) to a given cluster. Reference samples are those under the black horizontal bar, the newly collected samples are those under the grey horizontal bar with shades and labels corresponding to mitochondrial lineage (the final sample on the far right has a haplotype shared by the Islands of San Cristóbal and Santa Cruz).

Figure 3: Scatterplot of the first two principal components of DAPC showing assignment of unknown individuals to ancestry categories. The rows show pairwise combinations of parental populations. Top: Española × Floreana; middle: Floreana × PBL; bottom: Floreana × PBR. Parental populations are positioned at the ends of each plot with color coded first generation ($F_1$), second generation ($F_2$), and each backcross (BC) individuals between them. Ellipses encompass ~67% of the cloud of points for each group. These genotypes were used as a training set to define the discriminant functions and optimal number of PCs to retain. The Volcano Wolf tortoises with mixed ancestry are identified by black squares where placement is based on the previously defined discriminant functions.

Table 1: Number of individuals assigned to ancestry categories by DAPC (A) and NEWHYBRIDS (B)

A)

| Location | BC to ESP | BC to FLO | BC to PBL | BC to PBR | FLO/ESP $F_1$ or $F_2$ | FLO/PBL $F_1$ or $F_2$ | FLO/PBR $F_1$ or $F_2$ | Total |
|---|---|---|---|---|---|---|---|---|
| PNG | 9 | 2 | | | 8 | 1 | 3 | 23 |
| VW | 13 | 5 | 1 | 3 | 12 | 2 | 8 | 44 |
| Total | 22 | 7 | 1 | 3 | 20 | 3 | 11 | 67 |

B)

| Location | BC to ESP | $F_2$ with PBL or PBR | FLO/ESP $F_2$ | PBL or PBR BC to FLO | Total |
|---|---|---|---|---|---|
| PNG | 10 | 3 | 4 | 2 | 32 |
| VW | 6 | 13 | 17 | 4 | 27 |
| Total | 16 | 16 | 21 | 6 | 59 |

The current location of individuals is listed, as Galápagos National Park Breeding Center (PNG) or Volcano Wolf (VW). Abbreviations are: BC = backcross, $F_1$ = first generation mating, $F_2$ = second generation mating ($F_1$ x $F_1$), ESP = Española, and FLO = Floreana. Note that the total number of samples assigned differs between the two methods as some individuals did not exceed the posterior probability threshold for classification in NEWHYBRIDS (see Supplementary Methods)