Identification of Genetically Important Individuals of the Rediscovered Floreana Galápagos Giant Tortoise (*Chelonoidis elephantopus*) Provide Founders for Species Restoration Program

Joshua M. Miller[1][†]*, Maud C. Quinzin[1][†], Nikos Poulakakis[2,3], James P. Gibbs[4], Luciano B. Beheregaray[5], Ryan C. Garrick[6], Michael A. Russello[7], Claudio Ciofi[8], Danielle L. Edwards[9], Elizabeth A. Hunter[10], Washington Tapia[11,12], Danny Rueda[12], Jorge Carrión[12], Andrés A. Valdivieso[1], Adalgisa Caccone[1]*

1. Department of Ecology and Evolutionary Biology, Yale University, 21 Sachem St. New Haven, Connecticut, 06520, United States of America
2. Department of Biology, School of Sciences and Engineering, University of Crete, Vasilika Vouton, Gr-71300, Heraklio, Crete, Greece
3. Natural History Museum of Crete, School of Sciences and Engineering, University of Crete, Knossos Av., GR-71409, Heraklio, Crete, Greece
4. College of Environmental Science & Forestry, State University of New York, Syracuse, New York, 13210, United States of America
5. Molecular Ecology Lab, School of Biological Sciences, Flinders University, GPO Box 2100, Adelaide, SA, 5001, Australia
6. Department of Biology, University of Mississippi, Oxford, Mississippi, 38677, United States of America
7. Department of Biology, University of British Columbia, Okanagan Campus, Kelowna, BC V1V 1V7, Canada
8. Department Biology, University of Florence, 50019 Sesto Fiorentino (FI), Italy
9. Life and Environmental Sciences, University of California, Merced, 5200 N Lake Rd, Merced, California, 95343, United States of America
10. Department of Natural Resources and Environmental Science, University of Nevada – Reno, Max Fleischmann Agricultural Building, Reno NV 89557 USA
11. Galapagos Conservancy, Fairfax, Virginia, 22030, United States of America
12. Galápagos National Park Directorate, Puerto Ayora, Galápagos, Ecuador

*Corresponding authors: joshua.miller@yale.edu, adalgisa.caccone@yale.edu
†These authors contributed equally to this work

Abstract

Species are being lost at an unprecedented rate due to human-driven environmental changes. The

cases in which species declared extinct can be revived are rare. However, here we report that a

remote volcano in the Galápagos Islands hosts many giant tortoises with high ancestry from a

species previously declared as extinct: *Chelonoidis elephantopus* or the Floreana tortoise. Of 150

individuals with distinctive morphology sampled from the volcano, genetic analyses revealed

that 65 had *C. elephantopus* ancestry and thirty-two were translocated from the volcano's slopes

to a captive breeding center. A genetically informed captive breeding program now being

initiated will, over the next decades, return *C. elephantopu*s tortoises to Floreana Island to serve

as engineers of the island's ecosystems. Ironically, it was the haphazard translocations by

mariners killing tortoises for food centuries ago that created the unique opportunity to revive this

"lost" species today.

Introduction

Human activities have generated an extreme and rapid loss of biodiversity[1]. Many actions

have been undertaken to prevent species extinctions, including creating laws to protect

endangered species and critical habitats[2], translocations of individuals among populations or into

new habitats[3,4], and *ex situ* management including captive breeding[5]. For many species, such

actions come too late to facilitate recovery[6–8]. Although 'de-extinction' using laboratory

techniques is currently being debated[9], such methods are viable only for some taxa and will

generate significant anticipated, and unanticipated, risks[10,11]. Generally, extinction is final and

cases where lost species can be revived will be extremely rare.

Despite their insularity, even remote oceanic islands are not exempt from rapid

anthropogenic changes. For example, the ecosystems of the Galápagos Islands, located ~900

kilometers off the Pacific coast of Ecuador, have been degraded by human activities since the

archipelago was discovered in 1535. The resulting loss of biodiversity, including many endemic

species, has been chiefly due to the introduction of non-native species[12,13]. In response, concerted

efforts have been taken to restore ecosystems on the islands, including removal of introduced

pests[14,15] and population restoration via captive breeding and repatriation of threatened native

species (e.g. [16,17]).

Galápagos giant tortoises (*Chelonoidis* spp.) are flagship species for ongoing restoration
efforts[18] in this archipelago and play an important functional role as mega-herbivores in the
islands' ecosystems[19]. Galápagos giant tortoises can be classified into 15 species based on
genetic data[20]. Generally, there is a single species per island with two exceptions: Isabela Island
that has a different endemic species associated with each of its five volcanoes, and Santa Cruz
Island that contains two species, one recently described[20] (Figure 1a). The 15 species exhibit two
general carapace shapes: "domed," a rounded cupola-like form (Figure 1c), and a "saddle-
backed" form, with a high anterior opening creating the shape of a saddle (Figure 1d). Five of the
15 species have the saddle-backed morphology: *C. elephantopus* (Floreana), *C. hoodensis*
(Española), *C. abingdoni* (Pinta), *C. ephippium* (Pinzón), and *C. chathamensis* (San Cristóbal;
Figure 1a).

Over the past three centuries, all giant tortoise populations experienced a ~90% decline[21],
having been killed mostly for food and oil by whalers, sealers, buccaneers, and early
colonists[22,23]. Four species have been declared Extinct[24], including two of the five saddle-backed
species: *C. elephantopus* from Floreana Island and, most recently, *C. abingdoni,* from Pinta
Island. The latter species was represented by a single individual, Lonesome George, until his
death in 2012.

Surprisingly, recent research found living, wild tortoises with genetic ancestry from two
of the extinct saddle-backed species, *C. elephantopus* and *C. abingdoni* (hereafter referred to as
the Floreana and Pinta tortoises, respectively) outside their native range[25–28]. These individuals,
likely the descendants of tortoises translocated among islands by mariners[22,23,29], were discovered
on the remote Volcano Wolf on Isabela Island (Figure 1a,b). Among >1600 tortoises sampled
from Volcano Wolf in 2008 during an exploratory expedition, 105 individuals were admixed
between the locally endemic species, *C. becki*, and the Floreana ($n = 86$)[5] or Pinta ($n = 17$)[4]
species. The majority of these genetically admixed individuals were found on Volcano Wolf's
western slopes, facing Banks Bay (also known as Puerto Bravo, PBR: Figure 1b), with a smaller
number located on the volcano's northwestern slopes near Piedras Blancas (PBL)[25,26]. No
purebred individuals of either of the two non-native species were found in 2008, but genetic
simulations and the young age of some mixed ancestry individuals indicated that purebred
Floreana and Pinta tortoises might still be present on Volcano Wolf[25,26].

Here we build on this previous work. In November 2015 we mounted a 10-day-long
search involving ~70 field personnel combined with helicopter- and ship-support. Focusing our
search, we explored zones of Volcano Wolf most likely to contain individuals with ancestry from
these two "lost" species (Figure 1b) and restricted genetic sampling to those individuals with
saddle-backed morphology among the thousands of locally endemic domed *C. becki*. We then
assigned ancestry to these tortoises using reference databases containing both extant and extinct
species that have previously been used to assign ancestry to tortoises in the wild and captivity[25–
31]. Based on these assignments, we determined suitability of the tortoises for a genetically
informed captive breeding program aimed at reintroducing these key ecosystem engineers to
their native island.
102
Results & Discussion
In total, we encountered 144 individuals with saddle-backed morphology. Of those, 112
were released after taking blood samples, and 32 with pronounced saddle-backed morphology
were transported to the Galápagos National Park Service's captive tortoise breeding facility on
Santa Cruz Island[32–34]. We assigned ancestry to all 144 of these individuals along with six
saddle-backed tortoises known to have Floreana ancestry[35] already residing at the breeding
center using information from ~700-bp of mitochondrial DNA sequence and diploid genotypes
from 12 nuclear microsatellite loci. These loci have previously been shown to accurately assign
individuals to tortoise species[25–31].
Thirty-five of the 150 individuals analyzed had a mitochondrial DNA haplotype
diagnostic of the Floreana species (Supplementary Figure S1). The remaining individuals either
had haplotypes diagnostic of the Critically Endangered Española Island species ($n = 70$),
haplotypes shared between the Santiago Island and Volcano Wolf species ($n = 44$), or a
haplotype shared between the species from San Cristóbal and Santa Cruz Islands ($n = 1$). The
proportions of haplotypes associated with the Española and Floreana Island species reported here
(46% and 23%, respectively) are substantially higher than previously detected among individuals
from Volcano Wolf[25,26] (5% and 2%). This is likely due to our targeting of saddle-backed
individuals in 2015 versus sampling broadly during previous surveys[25–28].
Bayesian clustering analyses of microsatellite genotypic data using the method
implemented in STRUCTURE[36] revealed that 127 of the 150 tortoises sampled have ancestry

assignments (Q-values) to the extinct Floreana species (average Q-value ± SD = 0.87 ± 0.21;

range 0.16–0.99; Figure 2; Supplementary Figures S4-S6). Twenty-three individuals did not

show evidence of Floreana ancestry, being assigned to the two genetically distinct populations

(PBL and PBR) of *C. becki*, the endemic Volcano Wolf species. Of those individuals with

Floreana Q-values, 30 had Floreana mitochondrial haplotypes (Figure 2).

We conducted additional assignment tests using genotypes generated by simulated

matings within and among four possible parental lineages (PBL, PBR, Española Island, and

Floreana Island). Use of simulated individuals in assignment tests has been suggested to improve

accuracy and efficiency when distinguishing hybrid individuals[37]. Q-values from STRUCTURE

for the simulated hybrids were in the range expected for each "class" of hybrid (Supplementary

Figure S4), indicating we have the ability to identify individuals with Floreana ancestry across

various levels of admixture.

Analyses with the program GeneClass2 version 2.0[38] identified two tortoises with strong

assignment to the Floreana species, being classified as either purebreds or backcrosses between a

F_1 × purebred Floreana tortoise (Supplementary Table S2). This program also identified

additional 63 tortoises that were assigned to categories with Floreana ancestry. Further analysis

of these individuals found a large number of F_1 hybrids and backcrosses both when re-running

STRUCTURE (Q-values between 0.40–0.77; Supplementary Table S2), as well as when we used

NEWHYBRIDS version 1.1[39] (Table 1) and discriminant analysis of principal components

(DAPC ; Figure 3, Table 1). Admixture between the saddle-backed species from Floreana and

Española Islands (*n* = 43) was more common than between Floreana and either of the two

endemic domed *C. becki* populations (PBL and PBR; *n* = 22), indicative of positive assortative

mating between the two saddle-backed species.

A critical factor when planning a captive breeding program is having accurate ancestry

assignments. To examine the precision and accuracy of our ancestry assignments we conducted

additional analyses in STRUCTURE. First, jackknifing the loci to test robustness of Q-values to

reductions in the number of markers, and using the ANCESTDIST option in STRUCTURE,

which collects information about the distribution of Q-values for each individual in our dataset.

Jackknifing showed that individual Q-values were robust to reductions in the number of markers:

average variation in individual Q-values was 0.027 (SD=0.033, range <0.001-0.149;

Supplementary Figure S5). Second, use of the ANCESTDIST option highlighted that, as

expected, for hybrid individuals there is uncertainty around the specific estimate of Q-values

(range 0.002-0.484; Supplementary Figure S6). Inclusion of the simulated individual in our

STRUCTURE run led to a general decrease in the magnitude of Q-values observed

(Supplementary Table S2 and Supplementary Figure S7). Taken together, these analyses suggest

that Q-values should not be taken as direct measure of proportional ancestry, but that our

markers are powerful in detecting the presence of ancestry from the extinct Floreana species.

Neither mitochondrial DNA nor microsatellite data identified individuals with ancestry

from the Pinta Island species. However, a previous genetic estimate suggested that only 60–70

tortoises with Pinta ancestry are present on Volcano Wolf[25], whereas capture-mark-recapture

methods employed during our expedition estimated that a total of ~5,000–6,000 tortoises

occurred in the area searched. Moreover, despite our substantial search effort, we explored only

~26% of the total tortoise-occupied range on the volcano (J. Gibbs unpublished data). Therefore,

it is possible that individuals with Pinta ancestry still live on Volcano Wolf, but went undetected

in 2015.

Of the 38 individuals currently housed in the captive breeding center, 23 tortoises (9

males and 14 females) were found to have nuclear ancestry from Floreana across multiple

assignment methods (Table 1) with Q-values from STRUCTURE ranging between 0.44–0.77,

when simulated individuals were also included in the analyses. In addition, 12 of these 23

individuals have mitochondrial haplotypes from the Floreana species (Supplementary Table S2).

Together, these 23 individuals now form the core of a genetically-informed captive breeding

program aimed at repatriation of tortoises to Floreana Island. For the purposes of the breeding

program, all 23 genetically important individuals will be included, regardless of mitochondrial

lineage, in order to capture and maintain as much nuclear genetic diversity from the Floreana

species as possible. The program is modeled after another one developed for the Española

species initiated with only 15 founders. Over 50 years, the Española tortoise program generated

>2000 repatriates with nearly 1,000 surviving tortoises now reproducing independently on their

native island[17,40,41].

A key attribute for success of such breeding programs is that the founding individuals are

unrelated, as high relatedness can lead to inbreeding depression. Accordingly, we examined

relatedness among the 23 individuals with Floreana ancestry in the breeding center. The analysis

showed that most individuals were unrelated (average Queller and Goodnight's[42] relatedness = -

0.04, range -0.58–0.63; Supplementary Figure 3). Although the 23 tortoises in captivity represent

a promising founding population, 44 tortoises identified during the 2015 expedition, but left on

Volcano Wolf, are also good candidates for the Floreana breeding and repatriation program

(Table 1). These individuals, once re-located on Volcano Wolf, could be incorporated into the

breeding program to further expand the genetic diversity of the founder population.

The Floreana tortoise breeding program will be designed to maximize founder

contributions and *C. elephantopus* genome representation in the resulting progeny, while

promoting *in situ* population growth and minimizing costs to the Galápagos National Park

Directorate. Depending on the goals and priorities of Park decision-makers, complete genome

recovery may not be reached before releases of offspring begin. However, the high proportion of

Floreana ancestry and low relatedness evident in the current breeding individuals indicate that

the 23 founding individuals and their resulting progeny will provide a good starting point for

restoring the species.

Our discovery raises the possibility that the extinct Floreana species could be revived. In

this case, tortoises with Floreana ancestry are living 'genomic archives' that retain the

evolutionary legacy of the extinct species, removing the need for the cloning methods that have

been proposed to bring back extinct species[43]. The Floreana tortoise breeding program is

anticipated to generate thousands of offspring over the next few decades. When repatriated to

Floreana Island, these tortoises can once again play their critical role as ecosystem engineers[19].

In addition, giant tortoises are a major tourist attraction in Galápagos[44]; tortoise restoration on

Floreana Island should create new economic opportunities for the island's few human residents.

Ironically, the opportunity to revive this "lost species" today was created by the same early

visitors to the archipelago whose activities imperiled most giant tortoise species and drove some

into outright extinction.

Materials and Methods

Sampling and lab methods

All samples were collected under CITES permit 15US209142/9, Galápagos Park Permit

PC-75-16, and in accordance with Yale Institutional Animal Care and Use Committee (IACUC)

permit number 2016-10825. Samples were collected over a 10-day expedition in November 2015

by a team of 36 researchers plus Galápagos National Park rangers. Groups of 3–4 searchers were

assigned distinct areas on Volcano Wolf, collectively totaling ~36 km^2. The search area was
chosen based on a previous survey of tortoises on Volcano Wolf, which indicated that
individuals with Pinta and Floreana ancestry were found to be in their highest densities on the
western slopes of Volcano Wolf[25]. For all tortoises encountered, sex, age, and GPS coordinates
were recorded; for saddle-backed individuals, photographs were taken, blood samples were
collected for DNA analysis, and each individual was injected with a passive integrated
transponder (PIT) tag under the skin for identification in the future. In cases where blood was
taken, ~2 ml of blood was collected from the brachial vein of one of the front legs of the tortoise
and preserved in a lysis buffer containing 0.1 M Tris buffer, 0.1 M EDTA, 0.2 m NaCl, and 1%
SDS, pH 8.0. All tortoises were uniquely marked with paint when first encountered. Starting on
the sixth day, we re-searched areas and recorded whether any individual had been previously
encountered for the purposes of a capture-mark-recapture estimate of population size. In total,
1,333 tortoises were encountered, of which 144 had saddle-backed morphology. The 32 tortoises
removed from Volcano Wolf were initially carried in nets by helicopter from the flanks of the
volcano to the expedition ship anchored in Puerto Bravo harbor, and then transported to the
Galápagos National Park captive tortoise breeding facility on Santa Cruz Island.

DNA was extracted from 150 blood samples using Qiagen blood and tissue extraction
kits. These samples included the 144 saddle-backed individuals mentioned above along with six
individuals already housed at the Galápagos National Park Breeding Center on Santa Cruz Island
that were previously identified to have Floreana ancestry[35]. All samples were sequenced at
~700bp of the mitochondrial DNA control and were genotyped at 12 dinucleotide
microsatellites, using previously developed protocols (detailed procedures in Supplementary
Methods).

Ancestry assignment

For ancestry assignment based on mitochondrial DNA, the new sequences were aligned
to a reference database of 123 previously observed haplotypes, representing all extant and extinct
species[29]. Ancestry was assigned by determining shared haplotypes using the program TCS
version 1.21[45]. Evolutionary relationships were viewed with the program Network version 5.0
(fluxus-engineering.com; Supplementary Figure 1).

For ancestry assignment based on microsatellite genotypic data, we took a two-step
process using STRUCTURE version 2.3.4[36,46]. First, we ran STRUCTURE with a reference

dataset of 277 samples including all extant and extinct species to confirm the number of expected

genetic clusters (K) present in the archipelago (for full parameters see Supplementary Methods).

In this case, the optimal K was 12, which corresponds to previously described results[25,26]

(Supplementary Figure 2). Second, with archipelago-wide K established, we reduced the

reference dataset to the seven clusters previously found to be on Volcano Wolf[29]: Española, San

Cristóbal, Central Isabela (La Cazuela, Volcano Alcedo, and Volcano Darwin), Floreana, Pinta,

and the two genetically distinct endemic Volcano Wolf populations, PBL, and PBR. To this

reduced dataset (155 reference samples in total), the new samples were added and we set K to 7,

leaving all other parameters unchanged.

To further quantify the genetic ancestry of individuals, assignment tests were undertaken

using three additional methods. Prior to implementing these methods, we expanded the reference

database to include simulated genotypes. These genotypes were simulated using HYBRIDLAB

version 1.0[47], and corresponded to individuals arising from crosses within and among the

genetically distinct *C. becki* populations on Volcano Wolf (PBL and PBR), and the two saddle-

backed species from Española and Floreana (see Supplementary Methods). These four lineages

were chosen due to their high prevalence in the mitochondrial DNA-based assignments and their

contribution to microsatellite genotypes in admixed individuals. Simulated individuals

represented explicit ancestry categories: pure parental populations, first-generation (F_1) crosses,

second-generation (F_2) crosses, and backcrosses (i.e., matings between F_1's and their respective

purebred lineages).

With these new genotypes added into the reference set, we first used GeneClass version

2.0[38] to calculate the probability that an individual's genotype assigns to a population. Second,

we used NEWHYBRIDS version 1.1[39] to compute the posterior probability of various hybrid

classes for each individual. In this case, only four pairwise combinations of parental populations

were considered to focus on identifying potential hybrids involving Floreana ancestry: 1)

Española × Floreana, 2) PBL × Floreana, 3) Floreana × PBR, and 4) Española × PBR. Third, we

assigned ancestry using a multivariate approach, DAPC, as implemented in the R package

adegenet version 2.0.1[48,49]. In this case, the simulated genotypes from crossess among Floreana,

Española, PBL and PBR were initially used as a training dataset to define the principal

components and discriminant functions. The empirical genotypes for putative hybrids were then

transformed with principal components analysis (PCA) based on the centering and scaling of the

training set, and positioned onto the discriminant functions. The NEWHYBRIDS and DAPC

analyses were carried out only on individuals with ancestry from Floreana based on the results

from the GeneClass analyses (Supplementary Table S2). Finally, we re-ran STRUCTURE using

the simulated crosses as the reference populations.

Precision and accuracy of ancestry estimates were tested with two additional sets of

STRUCTURE analyses. First, we jackknifed our loci, sequentially removing one locus from the

dataset and rerunning STRUCTURE. Second, we used the ANCESTDIST option within

STRUCTURE. See Supplementary Materials for full methods.

Relatedness Analysis

For the 23 individuals in the breeding center with Floreana ancestry, we calculated

pairwise relatedness using the estimator of Queller and Goodnight[42]. This was the highest ranked

of eight tested relatedness estimators tested by the program irelr[50] which considers a composite

score that incorporates estimates of bias, variance, skewness, and kurtosis. The empirical

distribution of relatedness values was compared to the distributions of 10,000 simulated pairs of

individuals for each of four relatedness categories (unrelated, half sibs, full sibs, and parent–

offspring). The pairwise relatedness distribution of both the empirical and simulated data were

calculated using irelr[50].

<u>References</u>

1.　Dirzo, R. *et al.* Defaunation in the Anthropocene. *Science (80-.).* **345,** 401 LP-406 (2014).

2.　Novacek, M. J. & Cleland, E. E. The current biodiversity extinction event: Scenarios for

mitigation and recovery. *Proc. Natl. Acad. Sci. U. S. A.* **98,** 5466–5470 (2001).

3.　Corlett, R. T. Restoration, Reintroduction, and Rewilding in a Changing World. *Trends*

Ecol. Evol. **31,** 453–462 (2016).

4.　Weeks, A. R. *et al.* Assessing the benefits and risks of translocations in changing

environments: a genetic perspective. *Evol. Appl.* **4,** 709–725 (2011).

5.　Canessa, S. *et al.* Planning for ex-situ conservation in the face of uncertainty. *Conserv.*

Biol. **30,** 599–609 (2016).

6.　Hoffmann, M. *et al.* The Impact of Conservation on the Status of the World's Vertebrates.

Science (80-.). **330,** 1503–1509 (2010).

7.　Doremus, H. & Pagel, J. E. Why Listing May Be Forever: Perspectives on Delisting under

the U.S. Endangered Species Act. *Conserv. Biol.* **15,** 1258–1268 (2001).

8.   Taylor, M. F. J., Suckling, K. F. & Rachlinski, J. J. The Effectiveness of the Endangered
Species Act: A Quantitative Analysis. *Bioscience* **55,** 360–367 (2005).

9.   Sherkow, J. S. & Greely, H. T. What If Extinction Is Not Forever? *Science (80-.).* **340,**
32–33 (2013).

10.   Cohen, S. The Ethics of De-Extinction. *Nanoethics* **8,** 165–178 (2014).

11.   Jones, K. E. From dinosaurs to dodos: who could and should we de-extinct? *Front.*
Biogeogr. **6,** (2014).

12.   Phillips, R. B., Wiedenfeld, D. A. & Snell, H. L. Current status of alien vertebrates in the
Galápagos Islands: invasion history, distribution, and potential impacts. *Biol. Invasions*
14, 461–480 (2012).

13.   Guézou, A. *et al.* An Extensive Alien Plant Inventory from the Inhabited Areas of
Galapagos. *PLoS One* **5,** e10276 (2010).

14.   Harper, G. A. & Carrion, V. Introduced rodents in the Galapagos: colonisation, removal
and the future. *Isl. Invasives Erad. Manag.* 63–66 (2011).

15.   Campbell, K., Donlan, C. J., Cruz, F. & Carrion, V. Eradication of feral goats Capra
hircus from Pinta Island, Galápagos, Ecuador. *Oryx* **38,** 328–333 (2004).

16.   Fabiani, A. *et al.* Conservation of Galápagos land iguanas: genetic monitoring and
predictions of a long-term program on the island of Santa Cruz. *Anim. Conserv.* **14,** 419–
429 (2011).

17.   Milinkovitch, M. C. *et al.* Genetic analysis of a successful repatriation programme: giant
Galapagos tortoises. *Proc. R. Soc. B-Biological Sci.* **271,** 341–345 (2004).

18.   Benitez-Capistros, F., Hugé, J., Dahdouh-Guebas, F. & Koedam, N. Exploring
conservation discourses in the Galapagos Islands: A case study of the Galapagos giant
tortoises. *Ambio* **45,** 706–724 (2016).

19.   Gibbs, J. P., Sterling, E. J. & Zabala, F. J. Giant Tortoises as Ecological Engineers: A
Long-term Quasi-experiment in the Galápagos Islands. *Biotropica* **42,** 208–214 (2010).

20.   Poulakakis, N. *et al.* Description of a New Galapagos Giant Tortoise Species
(Chelonoidis; Testudines: Testudinidae) from Cerro Fatal on Santa Cruz Island. *PLoS One*
10, e0138779 (2015).

21.   Márquez, C. *et al.* Estado actual de las poblaciones de tortugas terrestres gigantes
(Geochelone spp., Chelonia: Testudinae) en las islas Galápagos. *Ecol. Apl.* **3,** (2004).

22.   Porter, D. *Journal of a cruise made to the Pacific Ocean*. (Wiley & Halsted, 1822).

23.   Townsend, C. H. The Galápagos tortoises in relation to the whaling industry: A study of
342   old logbooks. *Zoologica* **4,** 55–135 (1925).

343   24.   IUCN. The IUCN Red List of Threatened Species. Version 2016-2. (2016). Available at:
344   www.iucnredlist.org. (Accessed: 10th November 2016)

345   25.   Edwards, D. L. *et al.* The genetic legacy of Lonesome George survives: Giant tortoises
346   with Pinta Island ancestry identified in Galapagos. *Biol. Conserv.* **157,** 225–228 (2013).

347   26.   Garrick, R. C. *et al.* Genetic rediscovery of an 'extinct' Galapagos giant tortoise species.
348   *Curr. Biol.* **22,** R10–R11 (2012).

349   27.   Poulakakis, N. *et al.* Historical DNA analysis reveals living descendants of an extinct
350   species of Galapagos tortoise. *Proc. Natl. Acad. Sci. U. S. A.* **105,** 15464–15469 (2008).

351   28.   Russello, M. A. *et al.* Lonesome George is not alone among Galápagos tortoises. *Curr.*
352   *Biol.* **17,** R317–R318 (2007).

353   29.   Poulakakis, N., Russello, M., Geist, D. & Caccone, A. Unravelling the peculiarities of
354   island life: vicariance, dispersal and the diversification of the extinct and extant giant
355   Galapagos tortoises. *Mol. Ecol.* **21,** 160–173 (2012).

356   30.   Benavides, E. *et al.* Lineage Identification and Genealogical Relationships Among
357   Captive Galapagos Tortoises. *Zoo Biol.* **31,** 107–120 (2012).

358   31.   Garrick, R. C. *et al.* Lineage fusion in Galapagos giant tortoises. *Mol. Ecol.* **23,** 5276–
359   5290 (2014).

360   32.   The Galapagos Conservancy. The Wolf Expedition: Resurrecting the extinct tortoise
361   species from Floreana and Pinta. (2015). Available at:
362   http://www.galapagos.org/newsroom/wolf-tortoise-search-outcome/.

363   33.   Nicholls, H. Lonesome George 'look-alike' captured for tortoise-breeding effort. *Nature*
364   (2015). doi:10.1038/nature.2015.18955

365   34.   Blakeslee, S. Scientists Hope to Bring a Galápagos Tortoise Species Back to Life. *The*
366   *New York Times* (2015).

367   35.   Russello, M. A. *et al.* DNA from the Past Informs Ex Situ Conservation for the Future: An
368   "Extinct'' Species of Galapagos Tortoise Identified in Captivity. *PLoS One* **5,** (2010).

369   36.   Falush, D., Stephens, M. & Pritchard, J. K. Inference of population structure using
370   multilocus genotype data: Linked loci and correlated allele frequencies. *Genetics* **164,**

1567–1587 (2003).

37.   Vähä, J.-P. & Primmer, C. R. Efficiency of model-based Bayesian methods for detecting

hybrid individuals under different hybridization scenarios and with different numbers of

loci. *Mol. Ecol.* **15,** 63–72 (2006).

38.   Piry, S. *et al.* GENECLASS2: A Software for Genetic Assignment and First-Generation

Migrant Detection. *J. Hered.* **95,** 536–539 (2004).

39.   Anderson, E. C. & Thompson, E. A. A model-based method for identifying species

hybrids using multilocus genetic data. *Genetics* **160,** 1217–1229 (2002).

40.   Milinkovitch, M. *et al.* Giant Galapagos tortoises; molecular genetic analyses identify a

trans-island hybrid in a repatriation program of an endangered taxon. *BMC Ecol.* **7,**

doi:10.1186/1472-6785-7-2 (2007).

41.   Gibbs, J. P., Hunter, E. A., Shoemaker, K. T., Tapia, W. H. & Cayot, L. J. Demographic

Outcomes and Ecosystem Implications of Giant Tortoise Reintroduction to Española

Island, Galapagos. *PLoS One* **9,** e110742 (2014).

42.   Queller, D. C. & Goodnight, K. F. Estimating relatedness using genetic-markers.

Evolution (N. Y). **43,** 258–275 (1989).

43.   Shapiro, B. Mammoth 2.0: will genome engineering resurrect extinct species? *Genome*

Biol. **16,** 228 (2015).

44.   Medina, C., Aravena, C. & Vásquez, F. *Economic valuation of the giant tortoise*

conservation in Galapagos Islands Carlos. (2012).

45.   Clement, M., Posada, D. & Crandall, K. A. TCS: a computer program to estimate gene

genealogies. *Mol. Ecol.* **9,** 1657–1659 (2000).

46.   Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using

multilocus genotype data. *Genetics* **155,** 945–959 (2000).

47.   Nielsen, E. E., Bach, L. A. & Kotlicki, P. HYBRIDLAB (version 1.0): a program for

generating simulated hybrids from population samples. *Mol. Ecol. Notes* **6,** 971–973

(2006).

48.   Jombart, T. adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics **24,** 1403–1405 (2008).

49.   Jombart, T. & Ahmed, I. adegenet 1.3-1: new tools for the analysis of genome-wide SNP

data. *Bioinformatics* **27,** 3070–3071 (2011).

50. Gonçalves da Silva, A. & Russello, M. A. iRel: software for implementing pairwise

relatedness estimators and evaluating their performance. *Conserv. Genet. Resour.* **3,** 69–

71 (2011).

51. R Core Team. R: A language and environment for statistical computing, reference index

version 3.2.2. (2015).

52. Hijmas, R. J. raster: Geographic Data Analysis and Modeling version 2.5-8. (2016).

Acknowledgements

This work was enabled by substantial logistical and personnel support of the Galápagos National

Park Directorate through its dedication to science-based conservation of giant tortoises via the

"Giant Tortoise Restoration Initiative" undertaken in conjunction with the Galapagos

Conservancy. Expedition expenses and laboratory analyses were also supported by grants from

the Galapagos Conservancy, the Mohamed bin Zayed Species Conservation Fund, National

Geographic Society, and the Oak Foundation to AC, the US National Science Foundation (DEB

1258062) to JG, the Belgian American Educational Foundation to MQ, and the Yale Institute for

Biospheric Studies. C. Gottsegen helped with genetic data collection, A. Antoniou provided

assistance with DAPC analyses. L. Boitani, L. Cayot, C. Parker, and A. Rhodin helped procure

funds and provided encouragement. J. Flanagan provided critical veterinary support. We would

also like to thank C. Cullingham for insightful comments on the manuscript.

Author Contributions

JMM, NP, LB, CC, EAH, WT, DR, JC, JPG and AC assisted with field logistics and sample

collection. JMM, MCQ, and AV conducted laboratory analyses. JMM, MCQ, and NP conducted

data analyses. LBB, RCG, MAR, and DLE provided technical guidance and conceptual advice

for data analyses. JMM drafted the manuscript with assistance from MCQ, LBB, MAR, JPG, and

AC. JMM, MCQ, NP, LBB, RCG, MAR, CC, DLE, EAH, WT, JPG, and AC discussed the

results and implications and commented on the manuscript at all stages.

Additional Information

Microsatellite genotypes and mitochondrial sequences are available on a server hosted by the

College of Environmental Science & Forestry, State University of New York, Syracuse

(http://www.esf.edu/efb/gibbs/Miller_et_al_data_archive.zip). The authors have no financial

conflicts of interest. Correspondence and requests for materials should be addressed to either

joshua.miller@yale.edu or adalgisa.caccone@yale.edu.

Figures

Figure 1: Distribution of tortoises among Galápagos Islands and representative photos of tortoise

carapace morphology. A) Map of the distribution of tortoises among Galápagos Islands

along with cartoons indicating carapace morphology for each. Light grey shading

indicates domed morphology, unshaded indicates saddle-backed. Extinct species are noted

with †. B) Larger view of Volcano Wolf on northern Isabela Island. The circle indicates

the approximate field location of the current study. Examples of Galápagos giant tortoises

with domed (C) saddle-backed (D) morphology. Maps were created with R (version

3.2.4[51]) using the raster package (version 2.5-8[52]).

Figure 2: STRUCTURE[36,46] plot for 155 reference samples along with the 150 newly collected
individuals. Each individual is represented as a vertical bar, with colors denoting the
different genetic clusters, as indicted below. The proportion of color in a bar is equal to
the ancestry (Q-value) to a given cluster. Reference samples are those under the black
horizontal bar, the newly collected samples are those under the grey horizontal bar with
shades and labels corresponding to mitochondrial lineage (the final sample on the far right
has a haplotype shared by the Islands of San Cristóbal and Santa Cruz).

Figure 3: Scatterplot of the first two principal components of DAPC showing assignment of unknown individuals to ancestry categories. The rows show pairwise combinations of parental populations. Top: Española × Floreana; middle: Floreana × PBL; bottom: Floreana × PBR. Parental populations are positioned at the ends of each plot with color coded first generation (F_1), second generation (F_2), and each backcross (BC) individuals between them. Ellipses encompass ~67% of the cloud of points for each group. These genotypes were used as a training set to define the discriminant functions and optimal number of PCs to retain. The Volcano Wolf tortoises with mixed ancestry are identified by black squares where placement is based on the previously defined discriminant functions.

Table 1: Number of individuals assigned to ancestry categories by DAPC (A) and NEWHYBRIDS (B)

A)

Location		BC to ESP	BC to FLO	BC to PBL	BC to PBR	FLO/ESP F_1 or F_2	FLO/PBL F_1 or F_2	FLO/PBR F_1 or F_2	Total
	PNG	9	2	0	0	8	1	3	23
	VW	13	5	1	3	12	2	8	44
Total		22	7	1	3	20	3	11	67

B)

Location		BC to ESP	BC to FLO	BC to PBL	BC to PBR	FLO/ESP F_1 or F_2	FLO/PBL F_1 or F_2	FLO/PBR F_1 or F_2	Total
	PNG	0	12	0	0	4	1	2	19
	VW	0	7	0	0	17	4	8	36
Total		0	19	0	0	21	5	10	55

The current location of individuals is listed, as Galápagos National Park Breeding Center (PNG) or Volcano Wolf (VW). Abbreviations are: BC = backcross, F_1 = first generation mating, F_2 = second generation mating (F_1 x F_1), ESP = Española, and FLO = Floreana. Note that the total number of samples assigned differs between the two methods as some individuals did not exceed the posterior probability threshold for classification in NEWHYBRIDS (see Supplementary Methods).