**Recipient-biased competition for a cross-fed nutrient is required**

**for coexistence of microbial mutualists**

Alexandra L. McCully, Breah LaSarre, James B. McKinlay[#]

Department of Biology, Indiana University, Bloomington, IN

Running title: Biased competition within a mutualism

[#]Corresponding author. 1001 E 3[rd] Street, Jordan Hall, Bloomington, IN 47405

Phone: 812-855-0359

Email: jmckinla@indiana.edu

**Conflict of interest.**

The authors declare no conflict of interest.

**Abstract.** Many mutualistic microbial relationships are based on nutrient cross-feeding. Traditionally,

cross-feeding is viewed as being unidirectional from the producer to the recipient. This is likely true when

a producer's metabolic waste, such as fermentation products, provides carbon for a recipient. However, in

some cases the cross-fed nutrient holds value for both the producer and the recipient. In such cases, there

is potential for nutrient reacquisition by producer cells in a population, leading to competition against

recipients. Here we investigate the consequences of inter-partner competition for cross-fed nutrients on

mutualism dynamics using an anaerobic coculture pairing fermentative *Escherichia coli* and phototrophic

*Rhodopseudomonas palustris*. In this coculture, *E. coli* excretes waste organic acids that provide carbon

for *R. palustris*. In return*, R. palustris* cross-feeds *E. coli* ammonium ($NH_4^+$), a valuable nitrogen

compound that both species prefer. To explore the potential for inter-partner competition, we first used a

kinetic model to simulate cocultures with varied affinities for $NH_4^+$ in each species. The model predicted

that inter-partner competition for cross-fed $NH_4^+$ could profoundly impact population dynamics. We then

experimentally tested the predictions by culturing mutants lacking $NH_4^+$ transporters in both $NH_4^+$

competition assays and cooperative cocultures. Both theoretical and experimental results indicated that

the recipient must have a competitive advantage in acquiring valuable cross-fed $NH_4^+$ to avoid collapse of

the mutualism. Thus, the very metabolites that form the basis for cooperative cross-feeding can also be

subject to competition between mutualistic partners.

**Significance.** Mutualistic relationships, particularly those based on nutrient cross-feeding, promote

stability of diverse ecosystems and drive global biogeochemical cycles. Cross-fed nutrients within these

systems can be either waste products valued only by one partner or nutrients that both partners value.

Here, we explore how inter-partner competition for a communally-valuable cross-fed nutrient impacts

mutualism dynamics. We discovered that mutualism stability necessitates that the recipient have a

competitive advantage against the producer in obtaining the cross-fed nutrient. We propose that the

requirement for recipient-biased competition is a general rule for mutualistic coexistence based on the

transfer of communally valuable resources, microbial or otherwise.

## Introduction

Mutualistic cross-feeding of resources between microbes can have broad impacts ranging from influencing host health (1, 2) to driving global biogeochemical cycles (3–6). Cross-fed metabolites are often regarded as nutrients due to the value they provide to a dependent partner, the recipient. However, for the partner producing the nutrient, the producer, a cross-fed nutrient's value can vary. On one extreme, the cross-fed metabolite is valued by the recipient but not the producer, as is the case for fermentative waste products (7–10). In other cases, a cross-fed metabolite holds value for both the recipient and the producer, as is the case for vitamin $B_{12}$ (6, 11, 12) and ammonium ($NH_4^+$) (13, 14). Such communally-valuable cross-fed nutrients are subject to partial privatization (15), wherein the producer has mechanisms to retain a portion of the nutrient pool for itself. While most mutualism cross-feeding studies only consider unidirectional metabolite transfer from producer to recipient, we wondered whether these mechanisms for partial privatization could lead to competition between partner populations for communally-valuable cross-fed nutrients. It seems likely that such competition could influence mutualism stability, as is known to be the case for competition for exogenous limiting resources (8, 16–19). To the best of our knowledge inter-partner competition for cross-fed nutrients and its impact on mutualism dynamics have never been investigated.

One example of cross-feeding that could involve competition between mutualistic partners is $NH_4^+$ excretion by $N_2$-fixing bacteria (Fig. 1A), called $N_2$-fixers (13, 14). During $N_2$ fixation, the enzyme nitrogenase converts $N_2$ gas into two $NH_3$ (20). In an aqueous environment, $NH_3$ is in equilibrium with $NH_4^+$. At neutral pH, $NH_4^+$ is the predominant form but small amounts of $NH_3$ can potentially leave the cell by diffusion across the membrane (21) (Fig. 1B). This inherent 'leakiness' for $NH_3$ likely fosters $NH_4^+$ cross-feeding, as extracellular $NH_3$ is available to neighboring microbes. Importantly, these neighbors can include clonal $N_2$-fixers, as $NH_3/NH_4^+$ is a preferred nitrogen source for most microbes. At concentrations above 20 μM, $NH_3$ can be acquired by passive diffusion; below 20 μM, $NH_4^+$ is bound and transported as $NH_3$ by AmtB transporters (Fig. 1B) (22). AmtB-like transporters are conserved throughout all domains of life (23). There is growing evidence that AmtB is used by $N_2$-fixers to recapture $NH_3$ lost

by passive diffusion, as $\Delta$AmtB mutants accumulate $NH_4^+$ in culture supernatants whereas wild-type

strains do not (24–26). Thus, during $NH_4^+$ cross-feeding, AmtB likely facilitates both $NH_4^+$ acquisition by

the mutualistic partner and recapture of $NH_4^+$ by the $N_2$-fixer.

Assessing the effects of inter-partner competition for a cross-fed nutrient would require a level of

experimental control not possible in most natural settings. However, synthetic microbial communities, or

cocultures, are well-suited to address such questions (27–29). We previously developed a bacterial

coculture that features cross-feeding of waste products (organic acids) from *Escherichia coli*, and a

communally-valuable nutrient ($NH_4^+$) from *Rhodopseudomonas palustris* Nx (Fig. 1A) (26). Here, using

both a kinetic model and genetic manipulation to alter the affinity of each species in the coculture for

$NH_4^+$, we demonstrate that inter-partner competition for cross-fed $NH_4^+$ plays a direct role in maintaining

coexistence. Specifically, insufficient competition by *E. coli* for $NH_4^+$ resulted in a collapse of the

mutualism. Mutualism collapse could be delayed or potentially avoided through higher net $NH_4^+$

excretion by *R. palustris* or increased *E. coli* population size. Our results suggest that, as a general rule,

competition for a cross-fed nutrient in an obligate mutualism must be biased in favor of the recipient to

avoid mutualism collapse and the potential extinction of both species.

**Results**

**Competition for cross-fed $NH_4^+$ is predicted to shape mutualism population dynamics.** Within

our coculture (Fig. 1A), *E. coli* (*Ec*) ferments sugars into waste organic acids, providing essential carbon

and electrons to *R. palustris* (*Rp*) Nx. *R. palustris* Nx is genetically engineered to excrete low micromolar

amounts of $NH_4^+$, providing essential nitrogen for *E. coli* (26). $NH_4^+$ excretion by *R. palustris* Nx is due to

mutation of NifA, the master transcriptional regulator of nitrogenase, which results in constitutive

nitrogenase activity even in the presence of normally inhibitory $NH_4^+$ (30). In contrast to organic acids,

which are only useful to *R. palustris*, $NH_4^+$ produced by *R. palustris* Nx is essential for the growth of both

species; *R. palustris* uses some $NH_4^+$ for its own biosynthesis and excretes the rest, which serves as the

nitrogen source for *E. coli*. However, *R. palustris* Nx can also take up $NH_4^+$ (30). Thus, we hypothesized

that competition for cross-fed $NH_4^+$ between the *R. palustris* Nx producer population and the *E. coli*

recipient population could influence mutualism dynamics.

We first explored whether competition for cross-fed $NH_4^+$ could affect the mutualism using

SyFFoN, a mathematical model describing our coculture (26, 31). SyFFoN simulates population and

metabolic dynamics in batch cocultures using Monod equations with experimentally-determined

parameter values. As previous versions described $NH_4^+$ uptake kinetics only for *E. coli* (26, 31), we

amended SyFFoN to include both an *R. palustris* $NH_4^+$ uptake affinity ($K_m$) and higher *R. palustris*

maximum growth rate ($\mu_{MAX}$) when $NH_4^+$ is used (SI Appendix Table S1). We then simulated batch

cocultures wherein the relative affinity for $NH_4^+$ varied between the two species (Fig. 2). The model

predicted that coexistence is maintained when the *R. palustris* affinity for $NH_4^+$ is low relative to that of

*E. coli* (*Rp:Ec* < 1); sufficient $N_2$ is converted to $NH_4^+$ to support *R. palustris* growth and enough $NH_4^+$ is

cross-fed to support *E. coli* growth. In contrast, when the *R. palustris* affinity for $NH_4^+$ is high relative to

that of *E. coli* (*Rp:Ec* > 1), *E. coli* growth is no longer supported because *E. coli* cannot compete for

excreted $NH_4^+$. However, high *R. palustris* cell densities were still predicted (Fig. 2) due to persistent,

low-level organic acid cross-feeding stemming from *E. coli* maintenance metabolism, which can support

*R. palustris* growth even when *E. coli* is not growing (31).

**Genetic disruption of AmtB $NH_4^+$ transporters affects relative affinities for $NH_4^+$.** Bacterial cells

generally acquire $NH_4^+$ through two mechanisms: passive diffusion of $NH_3$, or active uptake by AmtB

transporters (Fig. 1B). We hypothesized that deleting *amtB* genes in either species would result in a lower

affinity for $NH_4^+$ in that species and thus could be used to test how relative $NH_4^+$ affinity impacts

coculture dynamics. We generated ΔAmtB mutants of both *E. coli* and *R. palustris* and first characterized

the effect of the mutations in monoculture. Deletion of *amtB* in *E. coli* had no effect on growth or

fermentation profiles when $NH_4Cl$ was in excess (SI Appendix Fig. S1), consistent with previous

observations where ΔAmtB growth defects were only apparent at $NH_4^+$ concentrations below 20 μM (22).

In *R. palustris* ΔAmtB monocultures with $N_2$ as the nitrogen source, growth trends were equivalent to

those of the parent strain; however, *R. palustris* ΔAmtB excreted more $NH_4^+$ than the parent strain and

about a third of that excreted by *R. palustris* Nx (SI Appendix Fig. S1C and D). In line with our

hypothesis, $NH_4^+$ excretion by *R. palustris* ΔAmtB could be due to a decreased ability to reacquire $NH_4^+$

lost by diffusion, resulting in increased net $NH_4^+$ excretion. Alternatively, we considered that $NH_4^+$

excretion by *R. palustris* ΔAmtB could be due to improper nitrogenase regulation. In several other $N_2$-

fixers, proper nitrogenase regulation requires AmtB, for example to induce post-translational nitrogenase

inhibition (switch-off) in response to $NH_4^+$ (25, 32). We tested whether *R. palustris* ΔAmtB exhibits

$NH_4^+$-induced switch-off by adding $NH_4Cl$ to exponentially growing cultures and measuring $H_2$

production, an obligate product of the nitrogenase reaction (33), as a proxy for nitrogenase activity. Upon

adding $NH_4Cl$, $H_2$ production stopped in *R. palustris* ΔAmtB cultures. In contrast, $H_2$ production only

slowed slightly in *R. palustris* Nx cultures (SI Appendix Fig. S2), consistent with previous observations

for NifA* strains (34, 35). Additionally, like the parent strain, *R. palustris* ΔAmtB did not produce $H_2$

when grown with $NH_4^+$, unlike *R. palustris* Nx (SI Appendix Fig. S3). These observations demonstrate

that *R. palustris* ΔAmtB is competent for $NH_4^+$-induced nitrogenase repression, and thus $NH_4^+$ excretion

by *R. palustris* ΔAmtB is likely due to a poor ability to reacquire $NH_4^+$ lost by diffusion.

To test our hypothesis that deleting *amtB* would lower cellular affinity for $NH_4^+$, we directly

competed all possible *E. coli* and *R. palustris* strain combinations in competition assays where ample

carbon was available for each species but the $NH_4^+$ concentration was kept low; specifically, a small

amount of $NH_4^+$ was added every hour to bring the final $NH_4^+$ concentration to 5 μM (Fig. 3). In this

competition assay, the species that is more competitive for $NH_4^+$ should reach a higher cell density than

the other species. In all cases, WT *E. coli* was more competitive for $NH_4^+$ than *R. palustris*. However,

each *R. palustris* strain was able to outcompete *E. coli* ΔAmtB (Fig. 3), even though the *R. palustris*

maximum growth rate is 4.6-times slower than that of *E. coli* (SI Appendix Fig. S1). Even *R. palustris*

strains lacking AmtB outcompeted *E. coli* ΔAmtB (Fig. 3), indicating that *R. palustris* has a higher

affinity for $NH_4^+$ than *E. coli* independent of AmtB. These data confirmed that deletion of *amtB* was an

effective means by which to lower the relative affinity for $NH_4^+$ in each mutualistic partner.

**Altering relative NH$_4^+$ affinities affects mutualistic partner frequencies.** We then examined how

relative affinities for NH$_4^+$ influenced mutualism dynamics by comparing the growth trends of cocultures

containing either WT *E. coli* or *E. coli* ΔAmtB, paired with either *R. palustris* ΔAmtB, *R. palustris* Nx, or

*R. palustris* NxΔAmtB, the latter of which we previously determined to exhibit 3-fold higher NH$_4^+$-

excretion levels than the Nx strain in monoculture (26). For each *R. palustris* partner, cocultures with *E.*

*coli* ΔAmtB grew slower than cocultures with WT *E. coli* (Fig. 4A,B). *E. coli* ΔAmtB also constituted a

lower percentage of the population and achieved lower cell densities compared to WT *E. coli* when paired

with the same *R. palustris* strain (Fig. 4C). These lower frequencies were consistent with the competitive

disadvantage of *E. coli* ΔAmtB for excreted NH$_4^+$ (Fig. 3).

For *R. palustris* strains lacking AmtB, the effects on population trends varied. Consistent with our

previous work, *R. palustris* NxΔAmtB supported higher WT *E. coli* percentages and cell densities (Fig.

4C) (26). With high NH$_4^+$ excretion levels from *R. palustris* NxΔAmtB, faster *E. coli* growth leads to

rapid organic acid accumulation, which acidifies the environment, inhibits *R. palustris* growth, and leaves

organic acids unconsumed (Fig. 4D) (26). Surprisingly, although *R. palustris* ΔAmtB excreted less NH$_4^+$

than *R. palustris* Nx in monoculture, *R. palustris* ΔAmtB supported a higher WT *E. coli* population in

coculture and consumable organic acids accumulated (Fig. 4C, D). These trends resemble those from

cocultures with *R. palustris* NxΔAmtB (Fig. 4C, D), which has a high level of NH$_4^+$ excretion (SI

Appendix Fig. S1D). Unlike Nx strains, which have constitutive nitrogenase activity due to a mutation in

the transcriptional activator NifA (30), *R. palustris* ΔAmtB has WT NifA. Thus, *R. palustris* ΔAmtB can

likely still regulate nitrogenase expression, and thereby its activity, in response to nitrogen starvation. We

hypothesized that in coculture with WT *E. coli*, *R. palustris* ΔAmtB might experience heightened

nitrogen starvation, as NH$_4^+$ consumption by WT *E. coli* would limit NH$_4^+$ reacquisition by *R. palustris*

ΔAmtB (in an *R. palustris* ΔAmtB monoculture any lost NH$_4^+$ would remain available to *R. palustris*).

We therefore tested whether coculture conditions stimulated higher nitrogenase activity using an

acetylene reduction assay. In agreement with our hypothesis, *R. palustris* ΔAmtB had increased

nitrogenase activity in coculture conditions compared to monocultures, whereas *R. palustris* Nx, which

exhibits constitutive nitrogenase activity, showed similar levels in both conditions (SI Appendix Fig. S4).

Thus, the relatively high WT *E. coli* population in coculture with *R. palustris* ΔAmtB is likely due to both

the competitive advantage for acquiring $NH_4^+$ over *R. palustris* ΔAmtB (Fig. 3) and higher $NH_4^+$ cross-

feeding levels due to increased nitrogenase activity.

**_E. coli_ must have a competitive advantage for $NH_4^+$ acquisition to avoid mutualism collapse.** We

were surprised to observe that cocultures of *R. palustris* ΔAmtB paired with *E. coli* ΔAmtB showed little

growth when started from a single colony of each species (Fig. 4A), a method that we routinely use to

initiate cocultures (26, 31). We reasoned that the higher *R. palustris* ΔAmtB affinity for $NH_4^+$ relative to

*E. coli* ΔAmtB (Fig. 3) likely led to community collapse as predicted by SyFFoN (Fig. 2). Even though

SyFFoN had predicted *R. palustris* growth when outcompeting *E. coli* for $NH_4^+$ (Fig. 2), SyFFoN likely

underestimates the time required to achieve these densities, if they would be achieved at all, as SyFFoN

does not take into account cell death, which is known to occur when *E. coli* growth is prevented (31).

Consistent with the hypothesis that poor coculture growth was due to a competitive disadvantage of *E.*

*coli* ΔAmtB for $NH_4^+$, SyFFoN simulations indicated that starting with a larger *E. coli* ΔAmtB population

would increase the probability that any given *E. coli* cell would acquire $NH_4^+$ versus *R. palustris* and

thereby overcome the competitive disadvantage of *E. coli* ΔAmtB for $NH_4^+$ (SI Appendix Fig. S5).

Indeed, we observed greater growth of both species when cocultures were inoculated with equal or higher

relative densities of *E. coli* ΔAmtB versus *R. palustris* ΔAmtB (SI Appendix Fig. S5).

The explanation that mutualism collapse was due to a competitive advantage of *R. palustris* ΔAmtB

over *E. coli* ΔAmtB for $NH_4^+$ called into question why cocultures pairing *E. coli* ΔAmtB with either *R.*

*palustris* Nx or *R. palustris* NxΔAmtB did not collapse as well (Fig. 4), given that in all of these pairings

*E. coli* ΔAmtB is at competitive disadvantage (Fig. 3). We hypothesized that a relatively high $NH_4^+$

excretion level by these latter *R. palustris* strains (SI Appendix Fig. S1D) could compensate for a low *E.*

*coli* $NH_4^+$ affinity. To explore this hypothesis we simulated cocultures with the *R. palustris* affinity for

$NH_4^+$ set high relative to that of *E. coli* (*Rp:Ec* = 1000) and varied the *R. palustris* $NH_4^+$ excretion level

(Fig. 5). Indeed, increasing *R. palustris* $NH_4^+$ excretion was predicted to overcome a low *E. coli* affinity

for $NH_4^+$ and support growth of both species (Fig. 5). The only exception was at the highest levels of

$NH_4^+$ excretion, where *R. palustris* growth was predicted to be inhibited due to rapid *E. coli* growth and

subsequent accumulation of organic acids that acidify the environment (Fig. 5) (26). These simulations

suggested that *R. palustris* Nx and NxΔAmtB supported coculture growth with *E. coli* ΔAmtB due to

higher $NH_4^+$ excretion levels (SI Appendix Fig. S1D), whereas a combination of low $NH_4^+$ excretion by

*R. palustris* ΔAmtB (SI Appendix Fig. S1D) and a low affinity for $NH_4^+$ by *E. coli* ΔAmtB led to collapse

of the mutualism in this pairing.

So far, we had only considered the effect of severe discrepancies in $NH_4^+$ affinities between the two

species (e.g., 1000-fold difference in $K_m$ values in our simulations) as a mechanism leading to coculture

collapse within the time period of a single culturing. However, we wondered if a subtle discrepancy in

$NH_4^+$ affinities could lead to coculture collapse if given more time. We therefore simulated serial transfers

of cocultures with partners having different relative $NH_4^+$ affinities (Fig. 6A, B). At equivalent $NH_4^+$

affinities (Fig. 6A), both species were predicted to be maintained over serial transfers. However, when the

relative affinities approached a threshold (relative *Rp*:*Ec* = 2.75), cell densities of both species were

predicted to decrease over serial transfers (Fig. 6B). This decline in coculture growth is due to *E. coli*

being slowly but progressively outcompeted for $NH_4^+$ by *R. palustris*. As the difference between the *R.*

*palustris* and *E. coli* populations expands, *R. palustris* cells have a greater chance of acquiring $NH_4^+$ than

the smaller *E. coli* population, further starving *E. coli* and simultaneously cutting off *R. palustris* from its

supply of organic acids from *E. coli*.

The above prediction prompted us to investigate if cocultures pairing *R. palustris* Nx with *E. coli*

ΔAmtB were stable through serial transfers. We focused on cocultures with *R. palustris* Nx rather than *R.*

*palustris* NxΔAmtB because *R. palustris* Nx has AmtB and would therefore be most likely to outcompete

*E. coli* ΔAmtB. Strikingly, after eight serial transfers of cocultures pairing *R. palustris* Nx with *E. coli*

ΔAmtB we observed coculture collapse (Fig. 6C). This observation is in stark contrast to cocultures of *R.*

*palustris* Nx paired with WT *E. coli*, which we have serially transferred for over 100 times with no

extinction events (unpublished data). These results indicate that the recipient population must have a

competitive advantage for a cross-fed nutrient versus the producer population to avoid mutualism

collapse.

**Discussion**

Here we demonstrate that mutualistic partners can compete for a cross-fed nutrient upon which the

mutualistic interaction is based, in this case $NH_4^+$. This competition can impact partner frequencies and

mutualism stability. Efficient nutrient reacquisition by the producer can render nutrient excretion levels

insufficient for cooperative growth, starving the recipient and leading to tragedy of the commons (36).

Conversely, recipient-biased competition for a cross-fed nutrient drives cooperative directionality in

nutrient exchange and thereby promotes mutualism stability. One implication of these results is that inter-

partner competition can influence the level of resource privatization. Within microbial interdependencies,

partial privatization has primarily been thought to depend on mechanisms used by the producer to retain a

portion of a communally-valuable resource (15). Our data indicate that for excreted resources having a

transient availability to both mutualists, recipient acquisition mechanisms can also influence the level of

producer privatization, as the competition impacts how much of a cross-fed resource will be shared versus

re-acquired. In effect, recipient-biased competition avoids tragedy of the commons by enforcing partial

privatization of a communally-valuable resource. The importance of the recipient having the upper hand

in inter-partner competition likely applies to other synthetic cocultures and natural microbial mutualisms

that are based on the cross-feeding of communally-valuable nutrients, including amino acids (37, 38) and

vitamin $B_{12}$ (6, 11). The same rule could also apply to inter-kingdom and non-microbial examples of

cross-feeding (e.g., plants and pollinators, nutrient transfer between plants and bacteria or fungi (39)) and

cooperative feeding (e.g., honeyguide bird and human harvesting of bee hives (40), cooperative hunting

between grouper fish and moray eels (41)). In such cases, increased privatization of a cross-fed or shared

resource, for example through producer-biased competition, could threaten the mutualism upon which

both species depend (15, 39, 42).

In our system, AmtB transporters were crucial determinants of inter-partner competition for $NH_4^+$.

We were intrigued to find that when both species lacked AmtB, *R. palustris* out-competed *E. coli* for

$NH_4^+$ (Fig. 5), enough so to collapse the mutualism within a single culturing (Fig. 3). Whether by

maximizing $NH_4^+$ retention or re-acquisition, *R. palustris*, and perhaps other $N_2$-fixers, might have

additional mechanisms aside from AmtB to minimize loss of $NH_4^+$ as $NH_3$. These mechanisms could

include a relatively low internal pH to favor $NH_4^+$ over $NH_3$, negatively-charged surface features, or

relatively high affinities by $NH_4^+$-assimilating enzymes such as glutamine synthetase. There are several

reasons why it would be beneficial for $N_2$-fixers to minimize $NH_4^+$ loss. First, $N_2$ fixation is expensive,

both in terms of the enzymes involved (43) and the reaction itself, costing 16 ATP to convert one $N_2$ into

two $NH_3$ (33). Passive loss of $NH_3$ would only add to this cost, as more $N_2$ would have to be fixed to

compensate. Second, loss of $NH_4^+$ could benefit nearby microbes competing against an $N_2$-fixer for

separate limiting nutrients (14, 44). The possibility that $N_2$-fixers could have a superior ability to retain or

acquire $NH_4^+$ independently of AmtB is not farfetched. Bacteria are known to exhibit differential abilities

to compete for nutrients. For example, iron acquisition commonly involves iron-binding siderophores, but

siderophores can be chemically distinct and thereby differ in their affinity for iron (45). Strategies to

utilize siderophores as a shared resource are also numerous, leading to different cooperative or

competitive outcomes in microbial communities (45, 46). One must consider that additional mechanisms

for acquiring $NH_4^+$ beyond AmtB might likewise exist. As our results have raised the potential for inter-

partner competition for cross-fed resources themselves, understanding the physiological mechanisms that

confer competitive advantages for nutrient acquisition between species will undoubtedly aid in describing

the interplay between competition and cooperation within mutualisms.

**Materials and Methods**

**Strains and growth conditions.** Strains, plasmids, and primers are listed in SI Appendix Table S2. All *R.*

*palustris* strains contained *ΔuppE* and *ΔhupS* mutations to facilitate accurate colony forming unit (CFU)

measurements by preventing cell aggregation (47) and to prevent $H_2$ uptake, respectively. *E. coli* was

cultivated on Luria-Burtani (LB) agar and *R. palustris* on defined mineral (PM) (48) agar with 10 mM

succinate. $(NH_4)_2SO_4$ was omitted from PM agar for determining *R. palustris* CFUs. Monocultures and

cocultures were grown in 10-mL of defined M9-derived coculture medium (MDC) (26) in 27-mL

anaerobic test tubes. To make the medium anaerobic, MDC was bubbled with $N_2$, then tubes were sealed

with rubber stoppers and aluminum crimps, and then autoclaved. After autoclaving, MDC was

supplemented with cation solution (1 % v/v; 100 mM $MgSO_4$ and 10 mM $CaCl_2$) and glucose (25 mM),

unless indicated otherwise. *E. coli* monocultures were also supplemented with 15mM $NH_4Cl$. All cultures

were grown at 30°C laying horizontally under a 60 W incandescent bulb with shaking at 150 rpm. Starter

cocultures were inoculated with 200 μL MDC containing a suspension of a single colony of each species.

Test cocultures were inoculated using a 1% inoculum from starter cocultures. Serial transfers were also

inoculated with a 1% inoculum. Kanamycin and gentamycin were added to a final concentration of 100

μg/ml for *R. palustris* and 15 μg/ml for *E. coli* where appropriate.

**Generation of *R. palustris* mutants.** *R. palustris* mutants were derived from wild-type CGA009 (49).

Generation of strains CGA4004, CGA4005, and CGA4021 was described previously (26). For generation

of strain CGA4026 (*R. palustris* ΔAmtB) the WT *nifA* gene was amplified using primers JBM1 and

JBM2, digested with XbaI and BamHI, and ligated into plasmid pJQ200SK to make pJQnifA16. This

suicide vector was then introduced into CGA4021 by conjugation, and sequential selection and screening

was performed as described (50) to replace *nifA\** with WT *nifA*. Reintroduction of the WT *nifA* gene was

confirmed by PCR and sequencing.

**Generation of the *E. coli* ΔAmtB mutant.** P1 transduction (51) was used to introduce Δ*amtB::Km* from

the Keio collection strain JW0441-1 (52) into MG1655. The Δ*amtB::Km* genotype of kanamycin-resistant

colonies was confirmed by PCR and sequencing.

**Analytical procedures.** Cell density was assayed by optical density at 660 nm ($OD_{660}$) using a Genesys

20 visible spectrophotometer (Thermo-Fisher, Waltham, MA, USA). Growth curve readings were taken

in culture tubes without sampling (i.e., Tube $OD_{660}$). Specific growth rates were determined using

readings between 0.1-1.0 $OD_{660}$ where there is linear correlation between cell density and $OD_{660}$. Final

$OD_{660}$ measurements were taken in cuvettes and samples were diluted into the linear range as necessary.

$H_2$ was quantified using a Shimadzu (Kyoto, Japan) gas chromatograph (GC) with a thermal conductivity

detector as described (53). Glucose, organic acids, formate and ethanol were quantified using a Shimadzu

high-performance liquid chromatograph (HPLC) as described (54). $NH_4^+$ was quantified using an

indophenol colorimetric assay as described (26).

**Nitrogenase activity.** Nitrogenase activity was measured using an acetylene reduction assay (43). Cells

from 10-mL cultures were harvested and resuspended in 10-mL fresh MDC medium in 27-mL sealed

tubes pre-flushed with argon gas. Suspensions were incubated in light for 1 h at 30°C to recover. Then,

250 µl of 100% acetylene gas was injected into the headspace to initiate the assay, and ethylene

production was measured over time by gas chromatography as described (43). Ethylene levels were

normalized to total *R. palustris* CFUs in the 10-ml volume.

**$NH_4^+$ competition assay.** Fed-batch cultures were performed in custom anaerobic 75-ml serum vials with

side sampling ports. Each vial contained a stir bar and 30-mL of MDC, and was sealed at both ends with

rubber stoppers and aluminum crimps. Each vial was supplemented with 25 mM glucose, 1 % v/v cation

solution and 20 mM sodium acetate. Starter monocultures of each species were grown to equivalent

CFUs/mL in MDC tubes containing limiting nutrients (3 mM sodium acetate for *R. palustris* and 1.5 mM

$NH_4Cl$ for *E. coli*), and 1 mL of each species was inoculated into the serum vials. These competition

cocultures were incubated at 30°C under a 60 W incandescent bulb with stirring at 200 rpm (Thermo

Scientific) for 96 h. Each serum vial was constantly flushed with Ar to maintain anaerobic conditions.

$NH_4Cl$ was fed from a 500 µM $NH_4Cl$ stock using a peristaltic pump (Watson-Marlow) on an automatic

timer (Intermatic DT620) at a rate of 0.33 mL/min once an hour for a final concentration of ~ 5 µM upon

each addition. Samples were taken at 0 and 96 h for quantification of CFUs.

**Mathematical modeling.** A Monod model describing bi-directional cross-feeding in batch cultures,

called SyFFoN_v3 (Syntrophy between Fermenter and Fixer of Nitrogen), was modified from our

previous model (31) to allow for competition between *E. coli* and *R. palustris* for $NH_4^+$ as follows: (i) an

equation for *R. palustris* growth rate on $NH_4^+$ was added to boost the *R. palustris* growth rate when

acquiring $NH_4^+$ and (ii) the ability for *R. palustris* to consume $NH_4^+$ was added along with a $K_m$ of *R.*

*palustris* for $NH_4^+$ ($K_{AR}$). Equations and default parameter values are in the SI Appendix and Table S1.

SyFFoN_v3 runs in R studio and is available for download at: https://github.com/McKinlab/Coculture-

Mutualism.

**Acknowledgments**

We thank Richard Phillips (Indiana University) for providing equipment for the $NH_4^+$ competition assay.

We also thank Jay Lennon (Indiana University) for helpful discussions on the manuscript. This work was

supported in part by the U.S. Department of Energy, Office of Science, Office of Biological and

Environmental Research, under Award Number DE-SC0008131, by the U.S. Army Research Office,

grant W911NF-14-1-0411, and by the Indiana University College of Arts and Sciences.

**References**

1.  Flint HJ, Duncan SH, Scott KP, Louis P. (2007) Interactions and competition within the microbial

community of the human colon: Links between diet and health. *Environ Microbiol.* **9**(5):1101–1111.

2.  Hammer ND, et al. (2014) Inter- and intraspecies metabolite exchange promotes virulence of antibiotic-

resistant *Staphylococcus aureus*. *Cell Host Microbe.* **16**(4):531–537.

3.  McInerney MJ, Sieber JR, Gunsalus RP. (2010) Syntrophy in anaerobic global carbon cycles. *Curr Opin*

*Biotechnol.* **20**(6):623–632.

4.  Durham BP, et al. (2015) Cryptic carbon and sulfur cycling between surface ocean plankton. *Proc Natl*

*Acad Sci U S A*. **112**(2):453–457.

5.  Reeburgh WS. (2007) Oceanic Methane Biogeochemistry. *Chem Rev.* **107**(2):486–513.

6.  Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. (2005) Algae acquire vitamin $B_{12}$

through a symbiotic relationship with bacteria. *Nature.* **438**:90–93.

7.  McInerney MJ, et al. (2008) Physiology, ecology, phylogeny, and genomics of microorganisms capable of

syntrophic metabolism. *Ann N Y Acad Sci.* **1125**:58–72.

8.  Hillesland KL, Stahl DA. (2010) Rapid evolution of stability and productivity at the origin of a microbial

mutualism. *Proc Natl Acad Sci U S A.* **107**(5):2124–2129.

9.  Schink B. (1997) Energetics of syntrophic cooperation in methanogenic degradation. *Microbiol Mol Biol*

*Rev.* **61**(2):262–280.

10. Stams AJM. (1994) Metabolic interactions between anaerobic bacteria in methanogenic environments.

*Antonie van Leeuwenhoek, Int J Gen Mol Microbiol.* **66**:271–294.

11. Grant MA, Kazamia E, Cicuta P, Smith AG. (2014) Direct exchange of vitamin $B_{12}$ is demonstrated by

modeling the growth dynamics of algal-bacterial cocultures. *ISME J* **8**(7):1–10.

12. Seth EC, Taga ME. (2014) Nutrient cross-feeding in the microbial world. *Front Microbiol.* **5**:1-6.

13. Behrens S, et al. (2008) Linking microbial phylogeny to metabolic activity at the single-cell level by using

enhanced element labeling-catalyzed reporter deposition fluorescence in situ hybridization (EL-FISH) and

NanoSIMS. *Appl Environ Microbiol.* **74**(10):3143–3150.

14. Adam B et al. (2016) $N_2$-fixation, ammonium release and N-transfer to the microbial and classical food

web within a plankton community. *ISME J.* **10**:450-459.

15. Estrela S, Morris JJ, Kerr B (2016) Private benefits and metabolic conflicts shape the emergence of

microbial interdependencies. *Environ Microbiol.* **18**(5):1415–1427.

16. Meyer JS, Tsuchiya HM. (1975) Dynamics of mixed populations having complementary metabolism.

*Biotechnol Bioeng.* **17**:1065–1081.

17. Kim HJ, Boedicker JQ, Choi JW, Ismagilov RF. (2008) Defined spatial structure stabilizes a synthetic

multispecies bacterial community. *Proc Natl Acad Sci U S A.* **105**(47):18188–18193.

18. Miura Y, Tanaka H, Okazaki M. (1980) Stability analysis of commensal and mutual relations with

competitive assimilation in continuous mixed culture. *Biotechnol Bioeng.* **22**(5):929–946.

19. Estrela S, Trisos CH, Brown SP. (2012) From metabolism to ecology: cross-feeding interactions shape the

balance between polymicrobial conflict and mutualism. *Am Nat.* **180**(5):566–576.

20. Bulen WA, LeComte JR. (1966) The nitrogenase system from *Azotobacter*: two-enzyme requirement for

$N_2$ reduction, ATP-dependent $H_2$ evolution, and ATP hydrolysis. *Proc Natl Acad Sci U S A.* **56**(3):979–

986.

21. Walter A, Gutknecht J. (1986) Permeability of small nonelectrolytes through lipid bilayer membranes. *J*

*Membr Biol.* **90**(3):207–217.

22. Kim M, et al. (2012) Need-based activation of ammonium uptake in *Escherichia coli. Mol Syst Biol*

**8**(616):1–10.

23. Peng J, Huang CH. (2006) Rh proteins vs Amt proteins: an organismal and phylogenetic perspective on

$CO_2$ and $NH_3$ gas channels. *Transfus Clin Biol.* **13**(1–2):85–94.

24. Barney BM, Eberhart LJ, Ohlert JM, Knutson CM, Plunkett MH. (2015) Gene deletions resulting in

increased nitrogen release by *Azotobacter vinelandii*: Application of a novel nitrogen biosensor. *Appl*

*Environ Microbiol.* **81**(13):4316–4328.

25. Zhang T, et al. (2012) Involvement of the ammonium transporter AmtB in nitrogenase regulation and

ammonium excretion in *Pseudomonas stutzeri* A1501. *Res Microbiol.* **163**(5):332–339.

26. LaSarre B, McCully AL, Lennon JT, McKinlay JB. (2017) Microbial mutualism dynamics governed by

dose-dependent toxicity of cross-fed nutrients. *ISME J.* **11**:337–348.

27. Widder S, et al. (2016) Challenges in microbial ecology: building predictive understanding of community

function and dynamics. *ISME J.* **10**:2557–2568.

28. Momeni B, Chen CC, Hillesland KL, Waite A, Shou W. (2011) Using artificial systems to explore the

ecology and evolution of symbioses. *Cell Mol Life Sci.* **68**(8):1353–1368.

29. Lindemann SR, et al. (2016) Engineering microbial consortia for controllable outputs. *ISME J.* **10**:2077–

2084.

30. McKinlay JB, Harwood CS. (2010) Carbon dioxide fixation as a central redox cofactor recycling

mechanism in bacteria. *Proc Natl Acad Sci U S A.* **107**(26):11669–11675.

31. McCully AL, LaSarre B, McKinlay JB. (2017) Growth-independent cross-feeding modifies boundaries for

coexistence in a bacterial mutualism. *biorXiv*. **doi:** https://doi.org/10.1101/083386

32. Yakunin AF, Hallenbeck PC. (2002) AmtB is necessary for $NH_4^+$-induced nitrogenase switch-off and

ADP-ribosylation in *Rhodobacter capsulatus. J Bacteriol.* **184**(15):4081–4088.

33. Hoffman BM, Lukoyanov D, Dean DR, Seefeldt LC. (2013) Nitrogenase: A draft mechanism. *Acc Chem*

*Res.* **46**(2):587–595.

34. Rey FE, Heiniger EK, Harwood CS. (2007) Redirection of metabolism for biological hydrogen

production. *Appl Environ Microbiol.* **73**(5):1665–1671.

35. Heiniger EK, Oda Y, Samanta SK, Harwood CS. (2012) How posttranslational modification of

nitrogenase is circumvented in *Rhodopseudomonas palustris* strains that produce hydrogen gas

constitutively. *Appl Environ Microbiol.* **78**(4):1023–1032.

36. Rankin DJ, Bargum K, Kokko H. (2007) The tragedy of the commons in evolutionary biology. *Trends*

*Ecol Evol.* **22**(12):643–651.

37. Pande S, et al. (2014) Fitness and stability of obligate cross-feeding interactions that emerge upon gene

loss in bacteria. *ISME J.* **8**(5):953–62.

38. Harcombe W. (2010) Novel cooperation experimentally evolved between species. *Evolution.* **64**(7):2166–

2172.

39. Bronstein JL. (2001) The Exploitation of Mutualisms. *Ecol Lett.* **4**:277–287.

40. Isack HA, Reyer H-U. (1989) Honeyguides and honey gatherers: Interspecific communication in a

symbiotic relationship. *Science.* **243**(4896):1343–1346.

41. Bshary R, Hohner A, Ait-el-Djoudi K, Fricke H. (2006) Interspecific communicative and coordinated

hunting between groupers and giant moray eels in the red sea. *PLoS Biol* **4**(12):2393–2398.

42. Sachs JL, Mueller UG, Wilcox TP, Bull JJ. (2004) The evolution of cooperation. *Q Rev Biol.* **51**(2):211–

244.

43. Oda Y, et al. (2005) Functional genomic analysis of three nitrogenase isozymes in the photosynthetic

bacterium *Rhodopseudomonas palustris*. *Am Soc Microbiol.* 187(22):7784–7794.

44. Morris JJ, Lenski RE, Zinser ER. (2012) The black queen hypothesis: Evolution of dependencies through

adaptive gene loss. *mBio.* **3**(2):1–7.

45. Joshi F, Archana G, Desai A. (2006) Siderophore cross-utilization amongst rhizospheric bacteria and the

role of their differential affinities for $Fe^{3+}$ on growth stimulation under iron-limited conditions. *Curr*

*Microbiol.* **53**(2):141–147.

46. Niehus R, Picot A, Oliveira NM, Mitri S, Foster KR. (2017) The evolution of siderophore production as a

competitive trait. *Evolution*. doi:10.1111/evo.13230.

47. Fritts RK, Lasarre B, Stoner AM, Posto AL, Mckinlay JB (2017) A *Rhizobiales*-specific unipolar

polysaccharide adhesin contributes to *Rhodopseudomonas palustris* biofilm formation across diverse

photoheterotrophic conditions. *Appl Environ Microbiol.* **83**(4):1–14.

48. Kim M-K, Harwood CS (1991) Regulation of benzoate-CoA ligase in *Rhodopseudomonas palustris*.

*FEMS Microbiol Lett.* **83**:199–203.

49. Larimer FW, et al. (2004) Complete genome sequence of the metabolically versatile photosynthetic

bacterium *Rhodopseudomonas palustris*. *Nat Biotechnol.* **22**(1):55–61.

50. Rey FE, Oda Y, Harwood CS. (2006) Regulation of uptake hydrogenase and effects of hydrogen

utilization on gene expression in *Rhodopseudomonas palustris*. *J Bacteriol.* **188**(17):6143–6152.

51. Thomason LC, Costantino N, Court DL. (2007) *E. coli* Genome manipulation by P1 transduction. *Curr*

*Protoc Mol Biol.* doi 10.1002/0471142727.mb0117s79

52. Baba T, et al. (2006) Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the

Keio collection. *Mol Syst Biol.* **2**:2006.0008.

53. Huang JJ, Heiniger EK, McKinlay JB, Harwood CS. (2010) Production of hydrogen gas from light and the

inorganic electron donor thiosulfate by *Rhodopseudomonas palustris*. *Appl Environ Microbiol.*

**76**(23):7717–7722.

54. McKinlay JB, Zeikus JG, Vieille C. (2005) Insights into *Actinobacillus succinogenes* fermentative

metabolism in a chemically defined growth medium. *Appl Env Microbiol.* **71**(11):6651–6656.

    **Figure Legends**

**Fig. 1. Mechanisms of NH$_4^+$ transfer within an obligate bacterial mutualism based on cross-feeding**

**of essential nutrients.** (**A**) *Escherichia coli* (*Ec*) anaerobically ferments glucose into organic acids,

supplying *Rhodopseudomonas palustris* Nx (*Rp* Nx) with essential carbon. *R. palustris* Nx fixes N$_2$ gas

and excretes NH$_4^+$, supplying *E. coli* with essential nitrogen. For, formate; EtOH, ethanol. (**B**) NH$_4^+$ can

be passively lost from cells as NH$_3$. Both species encode high-affinity NH$_4^+$ transporters, AmtB, that

facilitate NH$_4^+$ uptake. NH$_4^+$ is the predominant form at neutral pH, as indicated by an enlarged arrow

head on double-sided arrows.

**Fig. 2. Simulations suggest that *E. coli* must have a competitive advantage for NH$_4^+$ acquisition**

**relative to *R. palustris* to support mutualistic growth**. Final cell densities (solid lines) of *R. palustris*

(*Rp*, purple) and *E. coli* (*Ec*, orange) after 300 h in simulated batch cultures for a range of relative NH$_4^+$

affinities. Initial cell densities are indicated by dotted lines. Relative NH$_4^+$ affinity values represent the

relative *E. coli* K$_m$ for NH$_4^+$ (K$_A$) divided by that of *R. palustris* (K$_{AR}$).

**Fig. 3. AmtB is important for competitive NH$_4^+$ acquisition.** Competitive indices for *E. coli* after 96 h
in NH$_4^+$-limited competition assay cocultures. Cocultures were inoculated with *E. coli* and *R. palustris* at
equivalent cell densities with excess carbon available for both species (25 mM glucose for *E. coli* and 20
mM sodium acetate for *R. palustris*). NH$_4^+$ was added to cocultures to a final concentration of 0.5 μM
every hour for 96 h. The dotted line indicates a competitive index value of 1, where both species are
equally competitive for NH$_4^+$. Filled triangles, WT *E. coli*; open triangles *E. coli* ΔAmtB. Error bars
indicate SD, n=3. Different letters indicate statistical differences between *E. coli* competitive index
values, p < 0.05, determined by one-way ANOVA with Tukey's multiple comparisons post test.

**Fig. 4. NH$_4^+$ transporters influence population and metabolic trends of both partners in coculture.**
Growth curves (**A**), growth rates (**B**), final cell densities after one culturing (**C**), and fermentation product
yields (**D**) from cocultures of all combinations of mutants lacking AmtB in each species. Final cell

densities and fermentation product yields were taken after one week, within 24 h into stationary phase.

ND, not determined. Error bars indicate SD, n=4. Different letters indicate statistical differences, p <

0.05, determined by one-way ANOVA with Tukey's multiple comparisons post test.

**Fig. 5. Higher *R. palustris* NH$_4^+$ excretion levels are predicted to compensate for a low *E. coli* NH$_4^+$**

**affinity.** 300 h batch cultures were simulated with a relative *R. palustris* : *E. coli* (*Ec* : *Rp*) K$_m$ value for

NH$_4^+$ of 0.001 over different *R. palustris* NH$_4^+$ excretion levels (R$_A$). Final cell densities, solid lines;

initial cell densities, dotted lines.

**Fig. 6. A low *E. coli* NH$_4^+$ affinity results in coculture collapse over serial transfers when paired**

**with *R. palustris* Nx. (A,B)** 300 h batch cultures were simulated and serial transferred used a 1%

inoculum based on the cell density at 300 h for the previous culture. Relative NH$_4^+$ affinity values

represent the relative *E. coli* K$_m$ for NH$_4^+$ (K$_A$) divided by that of *R. palustris* (K$_{AR}$). **(C)** Final cell

densities of *R. palustris* Nx and *E. coli* ΔAmtB of cocultures grown for one week, less than 24 h into

stationary phase. A 1% inoculum was used for each subsequent serial transfer. Error bars indicate SD,

n=4. Final *E. coli* cell percentages +/- SD for each transfer are shown.

**Recipient-biased competition for a cross-fed nutrient is required**

**for coexistence of microbial mutualists**

Alexandra L. McCully, Breah LaSarre, James B. McKinlay

Department of Biology, Indiana University, Bloomington

**SI Appendix.**

**SyFFoN_v3 description.**

Equations 1 – 4 were used to describe *E. coli* and *R. palustris* growth rates:
8
9    Eq. 1: *E. coli* growth rate; $\mathbf{\mu_{Ec}} = \mu_{EcMAX} \cdot [G/(K_G+G)] \cdot [A/(K_A+A)] \cdot [b_{Ec}/(b_{Ec}+10^{(f+C)})]$
10
11   Eq. 2: *R. palustris* growth rate ($N_2$); $\mathbf{\mu_{Rpn}} = \mu_{RpMAX} \cdot [C/(K_C+C)] \cdot [N/(K_N+N)] \cdot [b_{Rp}/(b_{Rp}+10^{(f+C)})]$
12
13   Eq. 3: *R. palustris* growth rate ($NH_4^+$); $\mathbf{\mu_{Rpa}} =$
14                    $\mu_{RpMAX2} \cdot [C/(K_C+C)] \cdot [A/(K_{AR}+A)] \cdot [b_{Rp}/(b_{Rp}+10^{(f+C)})]$
15
16   Eq 4: Total *R. palustris* growth rate; $\mathbf{\mu_{Rp}} = \mu_{Rpn} + \mu_{Rpa}$
17
18   Equations 5-14 were used to describe temporal changes in cell densities and extracellular compounds.
19   Numerical constants in product excretion equations are used to account for molar stoichiometric
20   conversions. Numerical constants used in sigmoidal functions are based on those values that resulted in
21   simulations resembling empirical trends. All R and r parameters are expressed in terms of glucose
22   consumed except for $R_A$, which is the amount of $NH_4^+$ produced per *R. palustris* cell (Table S1).
23
24   Eq. 5: Glucose; $d\mathbf{G}/d\mathbf{t} = -\mu_{Ec} \cdot Ec/Y_G - \mu_{Ec} \cdot Ec \cdot (R_c+R_f+R_e+R_{CO2}) -$
25                    $Ec \cdot (G/(K_G+G)) \cdot (10/(10+1.09^{(1000 \cdot \mu Ec)})) \cdot (b_{Ec}/(b_{Ec}+10^{(f+C)})) \cdot ((100/(100+6^C)) \cdot$
26                    $(r_C+r_f+r_e+r_{CO2}) + r_{C\_mono} + r_{f\_mono} + r_{e\_mono} + r_{CO2\_mono})$
27
28   Eq. 6: $N_2$; $d\mathbf{N}/d\mathbf{t} = -\mu_{Rp} \cdot Rp \cdot 0.5 \cdot Ra \cdot (1-(40/(40+1.29^N))) - \mu_{Rp} \cdot Rp/Y_N$
29
30   Eq. 7: Consumable organic acids; $d\mathbf{C}/d\mathbf{t} = Ec \cdot \mu_{Ec} \cdot R_c \cdot 2 + Ec \cdot 2 \cdot (G/(K_G+G))$
31          $\cdot (10/(10+1.09^{(1000 \cdot \mu Ec)})) \cdot (b_{Ec}/(b_{Ec}+10^{(f+C)})) \cdot (r_C \cdot (100/(100+6^C))+r_{C\_mono}) - (\mu_{Rp} \cdot Rp/Y_C)$
32          $- 0.25 \cdot Rp \cdot \mu_{Rp} \cdot Rh_{Rp} - 0.25 \cdot Rp \cdot r_{Hp} \cdot (C/(K_C+C)) \cdot (40/(40+1.29^N)) \cdot (b_{Rp}/(b_{Rp}+10^{(f+C)}))$
33
34   Eq. 8: Formate; $d\mathbf{f}/d\mathbf{t} = (Ec \cdot \mu_{Ec} \cdot R_f \cdot 6) + Ec \cdot 6 \cdot (G/(K_G+G)) \cdot (10/(10+1.09^{(1000 \cdot \mu Ec)}))$
35          $\cdot (b_{Ec}/(b_{Ec}+10^{(f+C)})) \cdot (r_f \cdot (100/(100+6^C)) + r_{f\_mono}$
36
37   Eq. 9: $NH_4^+$; $d\mathbf{A}/d\mathbf{t} = Rp \cdot \mu_{Rp} \cdot R_A \cdot (1-(40/(40+1.29^N))) - \mu_{Ec} \cdot Ec/Y_A - (\mu_{Rp} \cdot Rp/Y_{AR}) \cdot (A/(K_{AR}+A))$
38
39   Eq. 10: *E. coli*; $d\mathbf{Ec}/d\mathbf{t} = \mu_{Ec} \cdot Ec$
40
41   Eq. 11: *R. palustris*; $d\mathbf{Rp}/d\mathbf{t} = \mu_{Rp} \cdot Rp$
42
43   Eq. 12: Ethanol; $d\mathbf{e}/d\mathbf{t} = Ec \cdot 3 \cdot (\mu_{Ec} \cdot R_e + (G/(K_G+G)) \cdot (10/(10+1.09^{(1000 \cdot \mu Ec)}))$
44                    $\cdot (b_{Ec}/(b_{Ec}+10^{(f+C)})) \cdot (r_e \cdot (100/(100+6^C)) + r_{e\_mono}))$
45

Eq. 13: $CO_2$; $dCO2/dt = Ec \cdot 6 \cdot (\mu_{Ec} \cdot R_{CO2} + (G/(K_G+G)) \cdot (10/(10+1.09^{(1000 \cdot \mu Ec)}))$

$\cdot (b_{Ec}/(b_{Ec}+10^{(f+C)})) \cdot (r_{co2} \cdot (100/(100+6^C)) + r_{co2\_mono}))$

$+ Rp \cdot 0.5 \cdot (\mu_{Rp} \cdot Rh_{Rp} + r_{Hp} \cdot (C/(K_C+C)) \cdot (40/(40+1.29^N)) \cdot (b_{Rp}/(b_{Rp}+10^{(f+C)})))$

Eq. 14: $H_2$; $dH/dt = Rp \cdot (\mu_{Rp} \cdot R_{HRp} + r_{Hp} \cdot (C/(K_C+C)) \cdot (40/(40+1.29^N)) \cdot$

$(b_{Rp}/(b_{Rp}+10^{(f+C)}))) + Ec \cdot (\mu_{Ec} \cdot R_{HEc} + (G/(K_G+G)) \cdot (10/(10+1.09^{(1000 \cdot \mu Ec)})) \cdot$

$(b_{Ec}/(b_{Ec}+10^{(f+C)})) \cdot (r_H \cdot (100/(100+6^C)) + r_{H\_mono}))$

Where,

$\mu$ is the specific growth rate of the indicated species ($h^{-1}$).

$\mu_{MAX}$ is the maximum specific growth rate of the indicated species ($h^{-1}$).

G, A, C, N, f, e, H and CO2 are the concentrations (mM) of glucose, $NH_4^+$, consumable organic acids, $N_2$, formate, ethanol, $H_2$, and $CO_2$, respectively. All gasses are assumed to be fully dissolved. Consumable organic acids are those that *R. palustris* can consume, namely, lactate (3 carbons), acetate (2 carbons), and succinate (4 carbons). All consumable organic acids were simulated to have three carbons for convenience. Only net accumulation of formate, ethanol, $CO_2$ and $H_2$ are described in accordance with observed trends.

K is the half saturation constant for the indicated substrate (mM).

Ec and Rp are the cell densities (cells/ml) of *E. coli* and *R. palustris*, respectively.

b is the ability of a species to resist the inhibiting effects of acid (mM).

Y is the *E. coli* or *R. palustris* cell yield from the indicated substrate (cells / μmol glucose). Y values were determined in MDC with the indicated substrate as the limiting nutrient.

R is the fraction of glucose converted into the indicated compound per *E. coli* cell during growth (μmol of glucose / *E. coli* cell), except for $R_A$. Values were adjusted to accurately simulate product yields measured in cocultures and in MDC with and without added $NH_4Cl$.

$R_A$ is the ratio of $NH_4^+$ produced per *R. palustris* cell during growth (μmol / *R. palustris* cell). The default value was based on that which accurately simulated empirical trends.

r is the growth-independent rate of glucose converted into the indicated compound (μmol / cell / h). Default values are based on those which accurately simulated empirical trends in coculture.

$r_{mono}$ is the growth-independent rate of glucose converted into the indicated compound by *E. coli* when consumable organic acids accumulate. Default values are based on linear regression of products accumulated over time in nitrogen-free cell suspensions of *E. coli* (4).

**Table S1. Default parameter values used in the model unless stated otherwise**

| Parameter | Value | Description (Units); Source |
|---|---|---|
| $\mu_{EcMAX}$ | 0.2800 | *E. coli* max growth rate ($h^{-1}$); Monoculture |
| $\mu_{RpMAX}$ | 0.0772 | *R. palustris* max growth rate ($h^{-1}$); Monoculture |
| $\mu_{RpMAX2}$ | 0.0152 | Boost on *R. palustris* growth rate in presence of $NH_4^+$ ($h^{-1}$); Monoculture[a] |
| G | 25 | Glucose (mM) |
| A | 0.00005 | $NH_4^+$ (mM); from initial $(NH_4)_6Mo_7O_{24} \cdot 4H_2O$ concentration |
| C | 0 | Consumable organic acids (those that *R. palustris* was observed to consume: lactate, acetate, and succinate; mM) |
| N | 70 | $N_2$ (assumed to be fully dissolved; mM) |
| f | 0 | Formate (mM) |
| e | 0 | Ethanol (mM) |
| CO2 | 0 | Carbon dioxide (mM) |
| $K_G$ | 0.02 | *E. coli* affinity (Michaelis-Menten constant ($K_m$)) for glucose (mM); (1) |
| $K_C$ | 0.01 | *R. palustris* affinity ($K_m$) for consumable organic acids (mM); Assumed |
| $K_A$ | 0.01 | *E. coli* affinity for $NH_4^+$ (mM); (2) |
| $K_{AR}$ | 0.01 | *R. palustris* affinity for $NH_4^+$ (mM); Assumed[b] |
| $K_N$ | 6 | *R. palustris* affinity ($K_m$) for $N_2$ (mM) |
| Ec | $0.4 \times 10^7$ | *E. coli* cell density (cells / ml) |
| Rp | $3.6 \times 10^7$ | *R. palustris* cell density (cells / ml) |
| $b_{Ec}$ | $10^{43}$ | Resistance of *E. coli* to low pH (mM) |
| $b_{Rp}$ | $10^{32}$ | Resistance of *R. palustris* to low pH (mM) |
| $Y_G$ | $8 \times 10^7$ | Glucose-limited *E. coli* growth yield (cells / μmol glucose); Glucose-limited *E. coli* culture |
| $Y_A$ | $1 \times 10^9$ | $NH_4^+$-limited *E. coli* growth yield (cells / μmol $NH_4^+$); $NH_4^+$-limited *E. coli* culture |
| $Y_C$ | $2.5 \times 10^8$ | Organic acid-limited *R. palustris* growth yield (cells / μmol organic acid); Acetate-limited *R. palustris* culture |
| $Y_N$ | $5 \times 10^8$ | $N_2$-limited *R. palustris* growth yield cells / μmol $N_2$; $N_2$-limited *R. palustris* culture |
| $R_C$ | $1.9 \times 10^{-8}$ | Fraction of glucose converted to organic acids (μmol glucose / cell) |
| $R_f$ | $8 \times 10^{-9}$ | Fraction of glucose converted to formate (μmol glucose / cell) |
| $R_e$ | $4.5 \times 10^{-9}$ | Fraction of glucose converted to ethanol (μmol glucose / cell) |
| $R_{CO2}$ | $5 \times 10^{-10}$ | Fraction of glucose converted to $CO_2$ (μmol glucose / cell) |
| $R_{HRp}$ | $2 \times 10^{-9}$ | *R. palustris* $H_2$ production (μmol $H_2$ / *R. palustris* cell) |
| $R_{HEc}$ | $5 \times 10^{-9}$ | *E. coli* $H_2$ production (μmol $H_2$ / *E. coli* cell) |
| $R_A$ | $0.15 \times 10^{-9}$ | *R. palustris* $NH_4^+$ production (μmol $NH_4^+$ / cell) |
| $r_C$ | $300 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to consumable organic acids (μmol glucose / cell / h) (3) |
| $r_f$ | $47 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to formate (μmol glucose / cell / h) (3) |
| $r_e$ | $15 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to ethanol (μmol glucose / cell / h) (3) |
| $r_{CO2}$ | $2 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to $CO_2$ (μmol glucose / cell / h) (3) |

| | | |
|---|---|---|
| $r_H$ | $2 \times 10^{-11}$ | *E. coli* specific growth-independent rate of $H_2$ production (μmol $H_2$ / cell / h) (3) |
| $r_{C\_mono}$ | $1.2 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to consumable organic acids when consumable organic acids accumulate (μmol glucose / cell / h); (4) |
| $r_{f\_mono}$ | $0.83 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to formate when consumable organic acids accumulate (μmol glucose / cell / h); (4) |
| $r_{e\_mono}$ | $0.5 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to ethanol when consumable organic acids accumulate (μmol glucose / cell / h); (4) |
| $r_{co2\_mono}$ | $1.3 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to $CO_2$ when consumable organic acids accumulate (μmol glucose / cell / h); (4) |
| $r_{H\_mono}$ | $0.83 \times 10^{-11}$ | *E. coli* specific growth-independent rate of glucose conversion to $H_2$ when consumable organic acids accumulate (μmol glucose / cell / h); (4) |
| $r_{Hp}$ | $27 \times 10^{-11}$ | *R. palustris* specific growth-independent rate of $H_2$ production (μmol $H_2$ / cell / h) |

88    [a] Increased growth rate in presence of $NH_4^+$ versus $N_2$ based on the difference in experimentally determined
89    growth rates in *R. palustris* monocultures grown with either $NH_4^+$ or $N_2$ as a nitrogen source.
90    [b] $K_{AR}$ was assumed to be equivalent to the published *E. coli* $K_m$ (2) for $NH_4^+$ ($K_A$).

91   **Table S2. Strains, plasmids, and primers used in this study**

| Strain or plasmid | Description or Sequence (5'-3'); Designation | Source or Purpose |
|---|---|---|
| ***R. palustris* strains** | | |
| CGA009 | Wild-type strain; spontaneous Cm$^R$ derivative of CGA001 | (5) |
| CGA4004 | CGA009 Δ*hupS* Δ*rpa2750*; Parent | (4) |
| CGA4005 | CGA4004 *nifA**; Nx | (4) |
| CGA4021 | CGA4005 Δ*amtB1* Δ*amtB2*; NxΔAmtB | (4) |
| CGA4026 | CGA4004 Δ*amtB1* Δ*amtB2;* ΔAmtB | This study |
| ***E. coli* strains** | | |
| MG1655 | Wild-type K12 strain, WT | (6) |
| K-12 JW0441-1 | Keio collection Δ*amtB::Km* | (7) |
| MG1655ΔAmtB | MG1655 Δ*amtB::Km*; ΔAmtB | This study |
| **Plasmids** | | |
| pJQnifA16 | Gm$^R$; WT *nifA* gene flanked by XbaI/BamHI cloned into pJQ200SK | This study |
| **Primers** | | |
| ALM6f | TTCGTCGCTGAATTGCAACG | *amtB* upstream flanking region (*E. coli*) |
| ALM6r | TCAGGAAGGGGTGATGCGTA | *amtB* downstream flanking region (*E. coli*) |
| JBM1 | CGTCTAGACCGGCGCATCGC | *nifA16* upstream primer; XbaI |
| JBM6 | GGGGATCCTGGTTCGCAGAGG | *nifA16* downstream primer; BamHI |

  **SI Appendix Figures.**

**Fig. S1.** *E. coli* ΔAmtB and *R. palustris* ΔAmtB **monoculture growth and metabolic trends. (A,B)**

Growth rates (**A**) and fermentation product yields (**B**) from WT *E. coli* (filled) or without (open) *E. coli*

ΔAmtB monocultures grown in MDC with 25 mM glucose and 15 mM $NH_4Cl$. Fermentation profiles were

generated from stationary monocultures. Error bars indicate SD, n=3. (**C,D**) Growth curves (**C**) and relative

$NH_4^+$ excretion (**D**) of *R. palustris* monocultures grown in MDC with 3 mM sodium acetate and a 100% $N_2$

headspace. Error bars indicate SD, n=4.

**Fig. S2. *R. palustris* ΔAmtB responds to NH$_4^+$-induced shutoff of nitrogenase.** The effect of either

NH$_4$Cl or NaCl on growth (**A**,**C**) and H$_2$ production (**B**,**D**) in *R. palustris* Nx or *R. palustris* ΔAmtB

monocultures. *R. palustris* monocultures were grown in MDC with 20 mM sodium acetate and a 100% N$_2$

headspace until mid-exponential phase and then supplemented with either 15 mM NH$_4$Cl or 15 mM NaCl

at the time indicated by the arrow.

**Fig. S3. Unlike *R. palustris* Nx, *R. palustris* ΔAmtB does not produce H$_2$ when grown with NH$_4^+$.** *R.*

*palustris* monocultures were grown in MDC with 20 mM sodium acetate and a 100% N$_2$ headspace with

(grey) or without (white) 15mM NH$_4$Cl. Samples for determining H$_2$ yields were taken one week after

inoculation, within 24 hours into stationary phase. Error bars indicate SD, n=3.

**Fig. S4. *R. palustris* ΔAmtB nitrogenase activity increases in coculture.** Normalized nitrogenase activity of *R. palustris* in monoculture (Mono) or coculture (Co) measured by an acetylene reduction assay. Ethylene levels were divided by total *R. palustris* CFUs in the test tube and then normalized to the *R. palustris* Nx monoculture value. Error bars indicate SD, n=4. *, statistical difference between monoculture and coculture conditions, $p < 0.05$, determined using multiple two-tailed t-tests; ns, no significant difference.

**Fig. S5. Higher initial cell densities of *E. coli* ΔAmtB can partially compensate for a low *E. coli* $NH_4^+$ affinity.** Simulations (**A**) and empirical data (**B**,**C**) showing the effect of initial *E. coli* (*Ec*) cell density on population and coculture growth trends when *E. coli* has a lower affinity for $NH_4^+$ compared to *R. palustris* (*Rp*). (**A**) 300 h batch cultures were simulated with a relative *R. palustris* : *E. coli* (*Rp* : *Ec*) $K_m$ value for $NH_4^+$ of 0.001. (**B, C**) Change in cell densities after one week of growth (**B**) and growth curves (**C**) of cocultures inoculated at different species ratios. (**A-C**) A ratio value of 1 represents $2.7 \times 10^6$ CFUs/mL, which was experimentally measured from the starting inoculum for both species before diluting to achieve the indicated ratios. Error bars indicate SD, n=3.

**SI Appendix References.**

1. Buhr A, Daniels GA, Erni B. (1992) The glucose transporter of *Escherichia coli*: Mutants with impaired

translocation activity that retain phosphorylation activity. *J Biol Chem.* **267**(6):3847–3851.

2. Khademi S, et al. (2004) Mechanism of ammonia transport by Amt/MEP/Rh: structure of AmtB at 1.35 A.

*Science.* **305**(5690):1587–1594.

3. McCully AL, LaSarre B, McKinlay JB. (2017) Growth-independent cross-feeding modifies boundaries for

coexistence in a bacterial mutualism. *biorXiv*. **doi:** https://doi.org/10.1101/083386

4. LaSarre B, McCully AL, Lennon JT, McKinlay JB. (2017) Microbial mutualism dynamics governed by

dose-dependent toxicity of cross-fed nutrients. *ISME J.* **11**:337–348.

5. Larimer FW, et al. (2004) Complete genome sequence of the metabolically versatile photosynthetic bacterium

*Rhodopseudomonas palustris. Nat Biotechnol.* **22**(1):55–61.

6. Hayashi K, et al. (2006) Highly accurate genome sequences of *Escherichia coli* K-12 strains MG1655 and

W3110. *Mol Syst Biol.* **2**(1):2006.0007.

7. Baba T, et al. (2006) Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the

Keio collection. *Mol Syst Biol.* **2**:2006.0008.