Genome-enabled insights into the ecophysiology of the comammox bacterium *Candidatus* Nitrospira nitrosa

Pamela Y. Camejo[1], Jorge Santo Domingo[2], Katherine D. McMahon[1,3], Daniel R. Noguera[1,]*

[1]Department of Civil and Environmental Engineering, University of Wisconsin - Madison, Madison, WI, USA

[2]Environmental Protection Agency, Cincinnati, OH, USA

[3]Department of Bacteriology, University of Wisconsin - Madison, Madison, WI, USA

Email addresses:

camejo.pamela@gmail.com

Santodomingo.Jorge@epa.gov

trina.mcmahon@wisc.edu

noguera@engr.wisc.edu

* Corresponding author: Daniel R. Noguera, 1415 Engineering Drive, Madison, WI 53706.

Email: noguera@engr.wisc.edu; Tel: 608-263-7783; Fax: 608-262-5199

ABSTRACT

The recently discovered comammox bacteria have the potential to completely oxidize ammonia to nitrate. These microorganisms are part of the *Nitrospira* genus and are present in a variety of environments, including Biological Nutrient Removal (BNR) systems. However, the physiological traits within and between comammox- and nitrite oxidizing bacteria (NOB)-like *Nitrospira* species have not been analyzed in these ecosystems. In this study, we identified *Nitrospira* strains dominating the nitrifying community of a sequencing batch reactor (SBR) performing BNR under micro-aerobic conditions. We recovered metagenomes-derived draft genomes from two *Nitrospira* strains: (1) *Nitrospira* sp. UW-LDO-01, a comammox-like organism classified as *Candidatus* Nitrospira nitrosa, and (2) *Nitrospira* sp. UW-LDO-02, a nitrite oxidizing strain belonging to the *Nitrospira defluvii* species. A comparative genomic analysis of these strains with other *Nitrospira*-like genomes identified genomic differences in *Ca.* Nitrospira nitrosa mainly attributed to each strains' niche adaptation. Traits associated with energy metabolism also differentiate comammox from NOB-like genomes. We also identified several transcriptionally regulated adaptive traits, including stress tolerance, biofilm formation and micro-aerobic metabolism, which might explain survival of *Nitrospira* under multiple environmental conditions. Overall, our analysis expanded our understanding of the genetic functional features of *Ca.* Nitrospira nitrosa, and identified genomic traits that further illuminate the phylogenetic diversity and metabolic plasticity of the *Nitrospira* genus.

INTRODUCTION

Nitrification is a microbiological process that plays an important role in the nitrogen (N) cycle. This process has been conventionally known as a two-step reaction. The first step, oxidation of ammonia to nitrite, is performed by ammonia-oxidizing bacteria (AOB) or archaea (AOA) and the second step, oxidation of nitrite to nitrate, is carried out by nitrite-oxidizing bacteria (NOB). Recently, the discovery of a new player with the potential to completely oxidize ammonia to nitrate, as in the case of complete ammonia oxidizing (comammox) organisms,[1, 2] has dramatically changed our understanding of microbial mediated N transformations in engineered and natural systems.

Comammox bacteria have been classified within the genus *Nitrospira*. Members of this genus were conventionally regarded as NOB and were thought to rely only on nitrite for growth. However, the genomes of the four comammox-like *Nitrospira* identified to date (*Ca.* Nitrospira nitrosa, *Ca.* Nitrospira nitrificans, *Ca.* Nitrospira inopinata and *Nitrospira* sp. Ga0074138[1-3]), encode the genes necessary for ammonia and nitrite oxidation, suggesting that *Nitrospira* are much more metabolically versatile organisms. Furthermore, comammox-like *Nitrospira* have been identified in a variety of habitats, including groundwater wells, drinking water biofilters, wastewater treatment plants (WWTPs) and other soil and aquatic environments.[4] These findings have prompted questions regarding the ecological significance and lifestyle of these organisms in each of these ecosystems.

Nutrient removal in WWTPs relies on nitrifying organisms to remove N from the wastewater. *Nitrospira*-like bacteria appear to be the dominant nitrite-oxidizers[5-7] in most WWTPs and laboratory scale reactors. The abundance of comammox in WWTPs has been briefly surveyed,

and preliminary results show this functional group is present in these systems.[4] However, genetic

and functional adaptations of comammox to this environment have not been addressed.

In this study, the community performing N removal in a Biological Nutrient Removal (BNR)

lab-scale reactor was analyzed to explore the genomic basis for comammox ecophysiology. A

sequencing batch reactor (SBR) was operated under cyclic anaerobic and micro-aerobic conditions

using two different operational conditions. During the first stage (nitrite addition during a micro-

aerobic phase), two *Nitrospira*-like strains were enriched in the reactor. Draft genome sequences

of these two strains were assembled from metagenomic data; one of them was identified as a

commamox organism, the other as an NOB. Here, we used the draft genomes of these strains, as

well as genomes from both NOB- and comammox-related bacteria, to perform a comparative

genome analysis of the genus *Nitrospira*.

MATERIAL AND METHODS

Operation of Lab-Scale Sequencing Batch Reactor

A laboratory-scale SBR was originally inoculated with activated sludge obtained from the

Nine Springs WWTP in Madison, WI, which uses a modified University of Cape Town (UCT)

process designed to achieve biological P removal[8] and operates with high aeration rates.[9] Synthetic

wastewater containing acetate as the sole carbon source was used for the feed, as described

elsewhere.[10] The hydraulic retention time (HRT) and solids retention time (SRT) were 24 h and

80 days, respectively. The pH in the system was controlled to be between 7.0 and 7.5.

The 2-liter reactor was operated under alternating anaerobic and low oxygen cycles. During

Stage 1 of operation the cycles consisted of 2 h anaerobic, 5 h micro-aerobic, 50 min settling and

10 min decanting. At the beginning of the micro-aerobic phase, sodium nitrite was added to reach

an in-reactor concentration of 10 mg $N\text{-}NO_2^-/L$ and stimulate the use of nitrite as an electron donor.

In addition, an on/off control system was used to limit the amount of oxygen pumped to the reactor

(0.02 L/min) and maintain low dissolved oxygen (DO) concentrations in the mixed liquor, as

described elsewhere.[10] After 100 days of operation, the nitrite supplement was eliminated and the

reactor cycle was changed to: 1.5 h anaerobic, 5.5 h micro-aerobic, 50 min settling and 10 min

decanting (Stage 2).

Sample Collection and Analytical Tests

To monitor reactor performance, mixed liquor and effluent samples were collected, filtered

through a membrane filter (0.45 μm; Whatman, Maidstone, UK) and analyzed for acetate, PO_4^{3-}–

P, NH_4^+-N, NO_3^--N, and NO_2^--N. The concentrations of PO_4^{3-}–P were determined according to

Standard Methods.[11] Total ammonia ($NH_3 + NH_4^+$) concentrations were analyzed using the

salicylate method (Method 10031, Hach Company, Loveland, CO). Acetate, nitrite and nitrate

were measured using high-pressure liquid chromatography as previously described.[10]

Biomass samples from the reactors were collected weekly and stored at -80°C until DNA

extraction was performed. DNA was extracted using UltraClean® Soil DNA Isolation Kit (MoBIO

Laboratories, Carlsbad, CA). Extracted DNA was quantified using a NanoDrop spectrophotometer

(Thermo Fisher Scientific, Waltham, MA) and stored at -80°C.

Metagenome Sequencing, Assembly and Binning

Samples from day 100 (Stage 1) and days 317, 522 and 674 (Stage 2) were selected for

metagenomic analysis. Illumina TruSeq DNA PCR free libraries were prepared for DNA extracts

according to the manufacturer's protocol and paired-end sequenced on either the Illumina HiSeq

2000 platform (v4 chemistry, 2×150 bp; 522-day sample), or the Illumina MiSeq platform (v3

chemistry, 2 × 250 bp; other samples). Unmerged reads were quality-trimmed and filtered with

Sickle (https://github.com/ucdavis-bioinformatics/sickle.git), using a minimum phred score of 20

and a minimum length of 50 bp. The metagenomic reads from the 100-day sample (Stage 1) were

assembled using IDBA-UD.[12] Individual genome bins were extracted from the metagenome

assembly with the R package 'mmgenome'[13] using the differential coverage principle.[14] The bins

were initially extracted by plotting the genome coverage of contigs in metagenomes from day 100

and 317. During the bin extraction, GC content and taxonomy of contigs were also taken into

consideration.

After binning, SSPACE was used to filter small scaffolds (length < 1,000 bp), extend

scaffolds, and to fill gaps.[15] Genome completeness and contamination was estimated using

CHECKM 0.7.1.[16] Table S1 displays quality metrics of the draft genomes after each of the steps

previously described. Two putative *Nitrospira*-like bins were identified and annotated using

MetaPathways v 2.0.[17] To further reduce contamination in these assembled bins, scaffolds with

Open Reading Frames (ORFs) having less than 85% nucleotide identity and 0% protein identity

to other *Nitrospira* genomes were removed from the bins.

Average Nucleotide Identity (ANI)

Pair-wise ANI values of *Nitrospira*-like genomes were obtained using the ANIm method[18]

and implemented in the Python script 'calculate_ani.py' available at

https://github.com/ctSkennerton/scriptShed/blob/master/calculate_ani.py.

Phylogenetic Analyses

The phylogeny of the draft genomes was assessed by constructing a phylogenetic tree using

a concatenated alignment of marker genes. First, PhyloSift v 1.0.1[19] was used to extract a set of 38

marker genes from each genome. Then, the extracted marker protein sequences were concatenated

into a continuous alignment to construct a maximum-likelihood (ML) tree, using RAxML v

7.2.8.[20] RAxML generated 100 rapid bootstrap replicates followed by a search for the best-scoring

ML tree.

For phylogenetic analyses of ammonia monooxygenase subunit A (*amoA*), hydroxylamine

reductase (*hao*) and nitrite oxido-reductase subunit A (*nxrA*) genes, full nucleotide datasets were

downloaded from the NCBI GenBank database.[21] Alignment was performed on sequences

retrieved from the NCBI and the draft genomes using the 'AlignSeqs' command in the DECIPHER

"R" package.[22] Phylogenetic trees were calculated using neighbor-joining criterion with 1,000

bootstrap tests for every node, using the MEGA6 software package.[23] Trees were visualized with

the assistance of TreeGraph.[24]

Population Structure by Metagenomic Analysis

To estimate the abundance of currently known ammonia oxidizers, comammox, and nitrite

oxidizers in the reactor over time, paired-end DNA reads from the metagenomic datasets (days

100, 317, 522 and 674) were competitively mapped to the published genome sequence of 14 AOB

(*Nitrosomonas*, *Nitrosospira* and *Nitrosococcus* genera), 6 AOA (*Nitrososphaera*,

Nitrosoarchaeum and *Nitrosopumilus*), 5 NOB (*Nitrospira* and *Nitrobacter* lineage), 5 anaerobic

ammonia-oxidizing (anammox) bacteria (*Ca.* Brocadia fulgida, *Ca.* Brocadia caroliniensis, *Ca.*

Kuenenia stuttgartiensis, *Ca.* Brocadia sinica and *Ca.* Jettenia caeni), 4 comammox bacteria (*Ca.*

Nitrospira nitrosa, *Ca.* Nitrospira nitrificans, *Ca.* Nitrospira inopinata and *Nitrospira* sp.

Ga0074138) and the two *Nitrospira*-like draft genomes retrieved from the reactor, using the

software package BBMap version 35.85 (https://sourceforge.net/projects/bbmap). A list of the

genomes included in this analysis and the number of reads mapping to each sequence is found in

Table S2. For each organism, the number of unambiguous reads (best hit) mapping to the genomic

sequence with a minimum alignment identity of 90% was quantified and normalized by the number

of reads in each metagenome, paired-end reads average length and genome size (Table S2).

Orthologous Genes Clusters

To assess the degree of homology in the proteomes of the two *Nitrospira*-like genomes,

orthologous genes clusters (OCs) were determined using OrthoMCL.[25] OrthoMCL was run with a

BLAST E-value cut-off of 1e-5, and an inflation parameter of 1.5.

Protein products of each ortholog set were compared against the Eggnog database[26] using

the eggNOG-mapper tool (https://github.com/jhcepas/eggnog-mapper) (*e*-value $<10e^{-2}$) to

determine the COG functional and super-functional category to which they belong.

Accession Numbers

Raw reads and draft genome sequences have been submitted to NCBI and are accessible

under the BioProject identifier PRJNA322674.

RESULTS AND DISCUSSION

Nutrient Removal in Lab-Scale Reactor

Results from a typical cycle of the lab-scale SBR at steady-state operation during Stage 1 are

shown in Fig. 1. During the anaerobic phase, acetate added at the beginning of the anaerobic phase

was completely consumed within an hour (Fig. 1B). P release to the mixed liquor during this

condition was not observed (Fig. 1B), indicating the absence of polyphosphate accumulating

organisms (PAO) in the reactor. Denitrification was incomplete, with only ~ 60% of the nitrate

removed in the anaerobic phase (Fig. 1A), even though the reactor received acetate in this phase.

This suggests that efficient acetate uptake was likely performed by glycogen accumulating

organisms (GAO), without affecting P concentrations.[27] In addition, nitrite production during this

phase (~ 10%) is an indicator of partial denitrification (Fig. 1A).

Complete nitrification occurred in the subsequent micro-aerobic stage, where 91% ± 4% of

ammonia and 91% ± 8% of the added nitrite were removed (Fig. 1A), with nitrate accumulation

accounting for 50% ± 2% of the oxidized nitrogen. During the period of active nitrification, the DO

remained below 0.05 mg/L (Fig. 1C), as the oxygen supplied balanced the oxygen uptake rate. The

oxygen uptake rate decreased after nitrification ceased, and correspondingly, DO increased. To

maintain a low-DO environment, aeration was stopped when DO exceeded the set point (0.2 mg

O_2/L) and resumed when DO decreased below the set point. This operation effectively maintained

DO below 0.4 mg/L (Fig. 1C).

After 100 days of reactor operation under these conditions, the operational parameters were

changed to promote simultaneous nitrification, denitrification and P removal (Stage 2). Unlike

Stage 1, no nitrite was added to the reactor during micro-aerobic conditions. During this second

stage, acetate added at the beginning of the anaerobic phase was used by PAOs for P-cycling, and

nitrite and nitrate produced by ammonia oxidization were used as electron acceptors by PAOs

during micro-aerobiosis, achieving simultaneous removal of N and P. Results of this stage were

described elsewhere.[10]

***Nitrospira*-like Genome Binning**

Using a combination of bidimensional coverage and tetranucleotide frequency, two

Nitrospira-like draft genomes were assembled from a sample collected at the end of Stage 1. The

two draft genomes (*Nitrospira* sp. UW-LDO-1 and UW-LDO-2) had 3.9 and 3.5 Mbp in total with

average GC content of 54.9% and 59.2%, respectively (Table S1). The reconstructed genomes

were assessed to be nearly complete (completeness $\geqq 90\%$) with low contamination ($\leqq 5\%$),

according to the presence of 43 single-copy reference genes (Table S1).

Since the composite genomes did not encode complete 16S rRNA genes, the average

nucleotide sequence identity (ANI) between the draft genomes assembled herein and formerly

published *Nitrospira*-like genomes, was used to determine whether UW-LDO-1 and UW-LDO-2

represented distinct species, as this method has been shown to correlate well with previously

defined 16S rRNA gene species boundaries.[28] The calculated ANI and fraction of alignment for

the *Nitrospira* genomes (Fig. 2) showed that UW-LDO-01 is a representative of the *Ca. Nitrospira*

nitrosa species (ANI >94%, Fraction Aligned = 74.9%), while UW-LDO-02 had the closest

nucleotide identity to *Nitrospira defluvii* (ANI = 92.4%, Fraction Aligned = 72.4%). None of the

other ANI values were greater than 88%, indicating that the two genomes were different from each

other, and supports their classification as *Nitrospira nitrosa* UW-LDO-01 and *Nitrospira defluvii*

UW-LDO-02, respectively.

Phylogenetic Analysis

A genome tree constructed from a concatenated protein alignment of 38 universally

distributed single-copy marker genes[29] confirms the affiliation of *Nitrospira* sp. UW-LDO-01 and

UW-LDO-02 with *Ca.* Nitrospira nitrosa and *Nitrospira defluvii*, respectively (Fig. 3). Consistent

with this phylogeny, UW-LDO-01 encoded the *amoCAB* operon, responsible for ammonia

oxidation, while UW-LDO-02 did not.

The *amoA* and *hao* genes are functional genes involved in redox nitrogen transformations

and are also considered phylogenetic markers to study the diversity of ammonia oxidizing

microorganisms (AOM).[30-33] The phylogenetic tree topologies based on these genes (Fig. S1)

further confirm the classification of UW-LDO-01 as related to *Ca.* Nitrospira nitrosa, although

two paralogs of the *amoA* gene are present in the genome of *Ca. Nitrospira nitrosa*, while only

one *amoA* gene was found in UW-LDO-1.

In addition, the *Nitrospira* sp. UW-LDO-1 and UW-LDO-2 genomes encoded the key

enzyme for nitrite oxidation, *nxr*, which can also be used as a phylogenetic biomarker. UW-LDO-

1 encoded two paralogs of the periplasmic NXR enzyme while *Nitrospira* UW-LDO-2 only

encoded one copy. The affiliation of UW-LDO-1 with *Ca.* Nitrospira nitrosa and UW-LDO-2 with

N. defluvii was supported by phylogeny based on the *nxrA* gene sequence (Fig. S2), consistent

with the phylogenetic analysis of *amoA* and *hao*.

Nitrifying Prokaryotes in Lab-Scale Reactor

The metagenomic analysis of the Stage 1 sample, which corresponds to the operational

conditions when nitrite and ammonia were both present under micro-aerobic conditions, did not

result in the assembly of any other genome of nitrifying microorganisms. Thus, in order to assess

the relative abundance of other known nitrifying prokaryotes present in the reactor, we mapped

metagenomic reads to published genomes of comammox, anammox, AOB, AOA and NOB,

including *Nitrospira* sp. UW-LDO-01 and UW-LDO-02 (Fig. 3). After a competitive mapping of

short-reads from metagenomic samples to each genome (>90% identity), the number of mapping

reads was normalized to both metagenome size and reference genome size, and used as a proxy of

genome abundance.

The metagenomic data show little evidence of AOA and anammox bacteria during the two

stages (0.06% and 0.25% of mapping reads, respectively). AOB were detected in the system, albeit

representing a small fraction of the community (0.17% and 0.05% of total number of reads during

Stage 1 and 2, respectively). Noticeably, *Nitrospira*-like sequences (including comammox- and

NOB-like genomes) recruited the greatest number of metagenomic reads (14.0% of total number

of reads) in the Stage 1 sample (Fig. 4A). Within this genus, *Nitrospira sp.* UW-LDO-01 retrieved

32.3% of the reads competitively mapping to the *Nitrospira*-like genomes (Fig. 4B). The published

genome of *Ca.* Nitrospira nitrosa retrieved 2.3% of the reads, while less than 4% mapped to other

comammox genomes. Therefore, with only a small fraction of reads mapping to other ammonia

oxidizers, we propose that *Nitrospira* sp. UW-LDO-01 was the main comammox in the reactor,

and the main contributor to ammonia oxidation during Stage 1.

Nitrospira sp. UW-LDO-02 appeared to be the most abundant NOB in the reactor during

Stage 1, retrieving 42.4% of the *Nitrospira*-like reads (Fig. 4B), although a large fraction of reads

competitively mapping to *N. defluvii* may indicate the presence of other nitrite oxidizing strains in

the reactor. Therefore, the nitrite oxidation activity in the reactor was carried out by comammox

and NOB.

The metagenomic analysis of Stage 2 samples reveals an overall decrease in the relative

abundance of nitrifying organisms in the reactor after transitioning to this operational

configuration (Fig. 4A). During this stage, metagenomic reads mapping to NOB and comammox

genomes (including *Nitrospira* sp. UW-LDO-01 and UW-LDO-02) decreased to less than 1% of

the total number of reads. This was in part due to the removal of nitrite addition during Stage 2.

However, the decrease in comammox did not correspond to an increase in the number of reads

mapping to other known AOM (Fig. 4A), suggesting the presence of still unrecognized AOM in

reactors operated with low DO conditions, as previously reported.[34]

Differential Gene Content Among *Ca*. Nitrospira nitrosa Genomes

Since *Nitrospira* sp. UW-LDO-01 is the second comammox genome representative of *Ca.*

Nitrospira nitrosa, and the first comammox genome recovered from a nutrient removal bioreactor,

a comparative analysis of its genetic content was carried out. First, a comparison of gene content

among *Ca.* Nitrospira nitrosa (CZQA00000000) and *Nitrospira* sp. UW-LDO-01 was conducted

by BlastP comparison of the translated coding DNA sequence (CDS) set, clustering of ortholog

proteins, and annotation of representatives of each ortholog cluster (OC) and genome-unique

CDSs.

Overall, sequencing and annotation of the UW-LDO-01 genome revealed a genomic

inventory highly similar to the genome of *Ca.* Nitrospira nitrosa.[2] The two genomes shared 81.2%

of the OCs (3,202 OCs), with UW-LDO-01 and *Ca.* Nitrospira nitrosa having 740 and 857 unique

OCs, respectively (Fig. 5A).

OCs belonging to the shared and genome-specific fractions of the two genomes were

classified according to their predicted functional role (Fig. 5B-C). 2,906 and 3,009 OCs, in the

UW-LDO-01 and *Ca*. Nitrospira nitrosa genomes, respectively, were categorized into functional

Clusters of Orthologous Groups (COG) categories. The majority of OCs belonged to the group of

"Unknown Function (S)" (23% of shared OCs, and 19% and 23% of the genome-specific fraction

in UW-LDO-01 and *Ca.* Nitrospira nitrosa, respectively), indicating a large set of metabolic

features not yet elucidated (Fig. 5B-C).

Besides the genes of unknown function, OCs within this shared fraction were mostly

represented by the "Cell wall/membrane/envelope biogenesis (M)" (7%), "Signal transduction

mechanisms (T)" (6%) and "Energy production and conversion (C)" (6%) families, indicating

general conservation of energy metabolism and regulatory mechanisms.

In both genomes, the functional group "Replication, recombination and repair (L)" was over-

represented within the genome-specific fraction. This functional category includes transposases,

integrases and other mobile genetic elements, and their extensive representation in the genome-

specific fraction indicates that horizontal gene transfer has likely played a significant role in the

diversification of *N. nitrosa* strains. The greatest difference among the genome-specific fractions

of UW-LDO-01 and *Ca*. Nitrospira nitrosa was the proportion of OCs represented by the "Cell

wall/membrane/envelope biogenesis (M)" and "Intracellular trafficking, secretion, and vesicular

transport (U)" functional families (Fig. 5C). Glycosyltransferases[35-38], Type IV[39-41] secretion

system proteins and other enzymes involved in polysaccharide formation (main component of the

biofilm matrix) were enriched within these COG categories in the genome-specific fraction of *Ca*.

Nitrospira nitrosa. Differences in biofilm formation capabilities between these strains may relate

to specific niche adaptation: *Ca*. Nitrospira nitrosa was enriched in a biofilm, whereas UW-LDO-

01 was found in a planktonic habitat in wastewater. Analogous findings have been observed in

other genera, where differences among biofilm formation capabilities within the same genus were

linked to the genome content of different strains.[42-44] Similar to the results presented here, these

genetic differences included the presence of type IV secretion systems and enzymes involved in

protein glycosylation.

The comparative genomic analysis also indicated a higher proportion of gene clusters

associated with "Lipid transport and metabolism (I)" in *Nitrospira* sp. UW-LDO-01 (Fig. 5C).

Genes related to ß-oxidation of long-chain fatty acids to acetyl-CoA were present in the genome

of UW-LDO-01, but absent in *Ca*. Nitrospira nitrosa. These genes include a long-chain fatty acid-

CoA ligase, acyl-CoA dehydrogenase, enoyl-CoA hydratase, 3-hydroxyacyl-CoA dehydrogenase

and acetyl-CoA acetyltransferase. Presence of these lipid-related metabolic genes in other

Nitrospira strains was confirmed, although the complete pathway is lacking in *Nitrospira defluvii*,

Ca. Nitrospira nitrificans and *Nitrospira* sp. Ga0074138. This feature may represent a competitive

advantage of some *Nitrospira* strains in habitats rich in long-chain fatty acids, such as WWTPs.[45]

Metabolic Features in *Nitrospira* Genomes

To explore the diverse metabolic capabilities and provide insights into the common and unique metabolic features encoded in the genome of NOB- and comammox-like strains, we compared the gene inventory of 9 complete and draft genomes classified as *Nitrospira*. The analysis was focused on traits associated to energy production, which are summarized in Table S3.

In agreement with previous analyses, only comammox-like genomes encoded ammonia monooxygenase (*amoCAB*) and hydroxylamine dehydrogenase (*haoAB-cycAB*) gene clusters, responsible for ammonia oxidation to nitrite (Table S3), reflecting the capability of this novel *Nitrospira* sub-lineage to perform full-nitrification from ammonia to nitrate.

Analysis of nitrite reducing genes revealed that all *Nitrospira* strains encoded a copper-containing dissimilatory nitrite reductases (*nirK*), which catalyzes the nitrite reduction to nitric oxide, a key step in the denitrification process. Despite the widespread presence of this enzyme across the *Nitrospira* genus, former studies have documented no activity of this protein in NOB-like[46] or comammox-like strains[1], where N loss caused by formation of gaseous compounds was not observed. Since it has been predicted that the NXR complex of *Nitrospira* can reduce nitrate to nitrite,[46] these microorganisms appear genetically capable of converting nitrate (product of nitrification) to nitric oxide. Additional experiments are still needed to obtain more insights into this *Nitrospira* trait. Other denitrification genes, such as nitrate reductase (*nar*), nitric oxide reductase (*nor*) or nitrous oxide reductase (*nos*), were not found in the *Nitrospira* strains analyzed here.

All comammox-like genomes, including UW-LDO-01, encoded the machinery to hydrolyze urea: the *ureABCDFG* urease operon and the *urtABCDE* urea transport system, suggesting that this *Nitrospira* sub-division possesses a high-affinity uptake system for urea and, thus, is adapted

to habitats where urea is present at low levels. A gene cluster involved in urea metabolism was also found in *N. moscoviensis* (Table S3), although the urea-binding proteins *urtBCDE* were lacking in the genome. Ureolytic activity of both *N. moscoviensis*, *Ca.* Nitrospira nitrosa and *Ca.* Nitrospira nitrificans was formerly tested by incubation of these strains with urea-containing media, where urea hydrolysis to ammonium was observed in both cases.[2, 46] Former studies have also shown the presence of genes for urea utilization in *Nitrospira lenta,*[47] a novel *Nitrospira* species enriched under low temperatures, suggesting that the ureolytic activity might be associated with this lineage. Since several ammonia oxidizers also possess the capability of hydrolyzing urea,[48-50] the presence of this trait in *Nitrospira* might evidence horizontal gene transfer among these functional groups.

A contrasting difference among NOB-like and comammox-like genomes was the capability to convert cyanate into ammonia. Only NOB-like genomes encoded a cyanase hydratase enzyme; former studies have experimentally confirmed cyanate degradation in *N. moscoviensis*.[51] Cyanate is produced intracellularly from urea and carbamoyl phosphate decomposition,[52, 53] and in the environment from the chemical/physicochemical decomposition of urea or cyanide.[54, 55] The presence of a cyanase enzyme benefits nitrite oxidizers because it allows them to detoxify cyanate, and the formed ammonium is then available for assimilation and might also serve as a source of energy for ammonia oxidizers in a process described as "reciprocal feeding".[46, 51] Further experiments analyzing the effect of cyanate in the growth on comammox-like bacteria are needed to understand how cyanate degradation would confer them a biological advantage, besides generation of ammonia.

The analysis also revealed the presence of the gene inventory for the uptake and oxidation of formate, an exclusive feature of NOB. Growth on formate as electron donor has been confirmed

in *N. moscoviensis* (under both micro-oxic and anoxic incubations),[46] *Nitrospira japonica*[56] and

uncultured *Nitrospira* in activated sludge.[57] Although formate oxidation should be advantageous

for organisms thriving in hypoxic or anoxic habitats, which also includes comammox-like bacteria,

this feature is absent in the genome of these microorganisms.

The genome of *N. moscoviensis* encodes a group 2a [Ni-Fe] hydrogenase (*hupS* and *hupL*)

and accessory proteins involved in the maturation and transcriptional regulation of hydrogenases

(*hypFCDEAB* and *hoxA*). Furthermore, experiments showed that *N. moscoviensis* was capable of

growing by aerobic respiration of H_2.[58] Although the comammox-like genomes lack the subunits

of the [Ni-Fe] hydrogenase (Hup), the five genomes analyzed here encoded a group 3 [Ni-Fe]

sulfur-reducing hydrogenase gene set (*hydBGDA and hybD*) positioned at the same locus where

Hup is located in *N. moscoviensis*. This hydrogenase complex is an heterotetramer with both

hydrogenase activity and sulfur reductase activity, which might play a role in hydrogen cycling

during fermentative growth.[59] Its beta and gamma subunits, that form the sulfur reducing

component, catalyze the cytoplasmic production of hydrogen sulfide in the presence of elemental

sulfur. The presence of this complex in the genomes indicates the potential of these

microorganisms for oxidizing H_2 using sulfur as electron acceptor, a trait that has not been

analyzed in comammox before, but that could confer this sub-group an advantage when growing

under anaerobic conditions.

Furthermore, the presence of a hyf-like operon (*hyfBCEFGI*), which encodes a putative

group 4 hydrogenase complex, was detected in every NOB-like genome, as well as *Ca.* Nitrospira

nitrosa, *Nitrospira* sp. Ga0074138 and UW-LDO-01. In *Escherichia coli*, this hydrogenase

complex forms part of a second formate hydrogenlyase pathway (oxidation of formate to CO_2 and

reduction of $2H^+$ to H_2 under fermentative conditions).[60] This is likely the case for the

hydrogenase-4 present in the genome of NOB-like strains, which co-occur with genes encoding

formate dehydrogenase. In comammox, however, the role of this distinct hydrogenase is not as

clear. In *Ca.* Nitrospira nitrosa, this complex is found immediately adjacent to a carbon monoxide

dehydrogenase (CODH), an enzyme that catalyzes the interconversion of CO and CO_2[61], a

genomic feature that would allow this strain to obtain energy from carbon monoxide.[62] Conversely,

the genomes of *Nitrospira* sp. Ga0074138 and UW-LDO-01 lack the CODH at this position, which

in the case of UW-LDO-01, was confirmed by alignment of the metagenomic reads to this gene.

No other neighboring gene of the hydrogenase-4 complex could be associated with this enzyme in

these two strains, therefore, the biological role of these genes is still unclear.

Altogether, these results reveal specific traits characterizing the NOB and comammox

functional groups: while comammox-like *Nitrospira* has the genomic potential of ammonia and

nitrite oxidation and potentially, sulfur reduction, NOB-like strains are distinguished by their

cyanate degradation and formate oxidation capabilities; and both urea hydrolysis and H_2

respiration are common traits shared by multiple *Nitrospira* strains.

The Role of Transcriptional Regulation in *Nitrospira*

Transcriptional regulation of gene expression is the most commonly used strategy to control

many of the biological processes in an organism, including progression through the cell cycle,

metabolic and physiological balance, and responses to environmental stress. This regulation is

generally orchestrated by several transcriptional factors (TFs) that directly coordinate the activity

of genes by binding to their promoters. Each *Nitrospira*-like genome codes for at least 100

transcriptional regulators, which account for ~3% of the estimated total number of genes, in

agreement with TFs in other microorganisms.[63-65] A comparative genomic analysis of full and draft

Nitrospira genomes was used to investigate the repertoire of TFs potentially involved in the

survival of these microorganisms under diverse environmental conditions (Table 1).

Among the TFs analyzed, the formate hydrogenlyase transcriptional activator (fhlA)[60, 66] was

the only one shared across all the *Nitrospira* genomes, although only NOB-like genomes encode

genes of its known regulon, the formate hydrogenase complex. The presence of this transcriptional

activator in comammox microorganisms, which appear to be genetically incapable of formate

oxidation (Table 1), might represent an ancestral trait shared by *Nitrospira* and lost during

diversification. This theory would also support the presence of the group 4 hydrogenase

(associated with the formate-hydrogenlyase complex in *E. coli*) in both NOB- and comammox-

like groups.

A common feature among some NOB and comammox bacteria is the presence of the

transcriptional regulators NhaR[67, 68] and OxyR[69] (Table 1). The first one is associated with stress

response to alkaline, acidic, saline and osmotic conditions. OxyR regulates hydrogen peroxide-

inducible genes, such as alkyl hydroperoxide reductase (*ahpCF*) and glutaredoxin (*grxA*), encoded

in all the *Nitrospira* genomes. Presence of these genes would confer *Nitrospira* an improved fitness

advantage over other nitrifying bacteria. For instance, NhaR is lacking in *Nitrosomonas* and

Nitrobacter and OxyR is not present in *Nitrosomonas* and *Nitrosospira* (based on genome search).

Furthermore, the role of NhaR during regulation of *pga* expression,[68] allows the biofilm formation

process to be considered as a flexible and dynamic developmental process driven by external

conditions, representing another means by which NhaR could promote survival of *Nitrospira*.

Likewise, the presence of the chemotaxis regulator CheZ in *Nitrospira* suggests chemotaxis as

another important mechanism by which these microorganisms efficiently and rapidly respond to

changes in the chemical composition of their environment.

To date, the role of the Fnr-type regulatory protein in *Nitrospira* has not been determined.

In other microorganisms, Fnr is part of the signaling involved in the adaptation to micro-oxic

environments,[70-74] where it acts as oxygen sensor and regulator of genes involved in anaerobic and

micro-aerobic metabolism. In *Nitrospi*ra, we predict that this TF would regulate similar genes,

such as the *frd* operon (fumarate reductase), *sdh* operon (succinate dehydrogenase), *ndh* (NADH

dehydrogenase) and *ccb3* complex (cytochrome c oxidase). At least one copy of Fnr in the

genomes of UW-LDO-01, *N. moscoviensis*, *N. defluvii* and N. sp. OLB-3 was located upstream of

a copper-containing nitrite reductase gene (nirK), suggesting a possible mechanism that controls

expression of this denitrification enzyme. The presence of multiple paralog copies of Fnr in several

Nitrospira genomes may indicate a rigorous regulation of metabolism when these microorganisms

are exposed to low levels of oxygen, an important factor affecting *Nitrospira* community

compositions in nitrifying systems.[75]

Overall, this study sheds light about differences in the physiological role of NOB and

comammox-like *Nitrospira*. Specifically, the comparative genomic results evidence traits

associated with energy metabolism as characteristic to each of these functional groups.

Furthermore, the analysis of TFs in *Nitrospira* reveals the alternative use of organic compounds,

response to environmental stress, chemotaxis and anaerobic metabolism as some of the key

mechanisms for the adaptive metabolism of the genus to multiple and adverse conditions. Further

studies in the field should include experiments that combine omics-analysis (transcriptomics,

metabolomics, and proteomics) with chemical data to confirm the ecological role and functionality

of each of these functional groups and their interactions with other microorganisms.

ACKNOWLEDGMENTS

This work was partially supported by funding from the National Science Foundation (CBET-

1435661 and MCB-1518130) and the Madison Metropolitan Sewerage District. Additional

funding from the Chilean National Commission for Scientific and Technological Research

(CONICYT) as a fellowship to Pamela Camejo is also acknowledged. The U.S. Environmental

Protection Agency, through its Office of Research and Development, partially funded and

collaborated in the research described herein. Any opinions expressed in this paper are those of

the authors and do not necessarily reflect the views of the agency; therefore, no official

endorsement should be inferred. Any mention of trade names or commercial products does not

constitute endorsement or recommendation for use.

FIGURES

Figure 1. Nutrient profiles of (A) nitrogenous compounds, (B) phosphorous and acetate, and (C)

oxygen concentration in a regular cycle of the lab-scale SBR during Stage 1. ANA: Anaerobic,

AER: Micro-aerobic, SET: Settling.

Figure 2. Comparison of the genome-wide average nucleotide identity and alignment percentage,

of *Nitrospira*-like genomes. The heatmap shows the average nucleotide identity (red upper section

of matrix) and the percentage of the two genomes that aligned (yellow lower section).

Figure 3. RAxML phylogenetic tree of a concatenated alignment of 37 marker genes (nucleotide sequence) data set with the root placed on the branch leading to *Nitrobacter winogradskyi*. The numbers at the nodes of both trees show support values derived from 100 RAxML bootstrap.

Figure 4. (A) Normalized frequency of metagenomic reads mapping to the genome of Comammox, AOB, AOA, NOB and Anammox-related organisms in samples from Stage 1 and 2 of the lab-scale SBR. (B) Relative abundance of reads mapping to genomes of *Nitrospira*-related bacteria in Stage 1 sample, including the draft genomes retrieved in this study.

Figure 5. Genomic differences between *Nitrospira sp.* UW-LDO-01 and *Ca*. Nitrospira nitrosa. (A) Venn diagram of ortholog clusters shared between the two draft genomes; (B) distribution of COGs functional classes in the fraction of orthologs shared by the two genomes; (C) distribution of COGs functional classes in ortholog clusters found in only one of the genomes (genome-specific fraction).

TABLES

Table 1. Inventory of transcriptional regulators with implications on adaptive metabolism, from complete and draft genomes of *Nitrospira*. Grey and white rectangles represent presence and absence of each gene, respectively.

Transcriptional Regulator	Gene	Comammox					NOB				Function
		Nitrospira sp. UW-LDO-01	*Ca.* Nitrospira nitrosa	*Ca.* Nitrospira nitrificans	*Ca.* Nitrospira inopinata	*Nitrospira* sp. Ga0074138	*Nitrospira* sp. UW-LDO-02	*Nitrospira moscoviensis*	*Nitrospira defluvii*	*Nitrospira* sp. OL23	
Formate hydrogenlyase transcriptional activator	*fhlA*	■	■	■	■	■	■	■	■	■	FhlA binds to formate hydrogenlyase structural genes (formate dehydrogenase and group 4 hydrogenase complex) to activate transcription of their promoters [66].
Transcriptional activator protein NhaR	*nhaR*	■	■	■	■	■	■	■	□	■	NhaR regulates *nhaA*, a pH-dependent sodium-proton antiporter that responds to alkaline and saline conditions [76]. It is also responsible for *osmC* induction [67], required for resistance to organic peroxides and osmotic conditions and for long-term survival in stationary phase [77, 78]. NhaR also stimulates *pga* transcription, a set of genes responsible of poly-β-1,6-N-acetyl-D-glucosamine (PGA) synthesis [68]. PGA is involved in cell-cell adhesion and attachment, which stabilizes biofilm formation [79].
Hydrogen peroxide-inducible genes activator	*oxyR*	■	■	■	■	■	■	□	■	■	OxyR is required for the induction of a hydrogen peroxide-inducible regulon in response to elevated levels of hydrogen peroxide [69].
Chemotaxis regulator CheZ	*cheZ*	■	■	■	■	■	□	■	■	■	CheZ is a component of the chemotaxis signal-transduction pathway [80]. It controls the phosphorylation of CheY, a protein involved in the cell excitation response. Absence of CheZ results in non-chemotactic cells or long stimuli response latencies, demonstrating its critical importance during response to stimuli [81].
Fumarate and nitrate reductase regulatory protein	*fnr*	■	■	■	□	□	■	■	■	■	Fnr is an oxygen-responsive regulator required for the expression of a number of genes involved in anaerobic metabolism [73, 82, 83], including fumarate reductase, nitrate and nitrite reductase, and cytochrome oxidase genes [84]

 ASSOCIATED CONTENT

 Supplementary Material. Supplementary Figures and Tables supporting the information

 presented in this manuscript.

 REFERENCES

504 1.   Daims, H.; Lebedeva, E. V.; Pjevac, P.; Han, P.; Herbold, C.; Albertsen, M.; Jehmlich,

 N.; Palatinszky, M.; Vierheilig, J.; Bulaev, A.; Kirkegaard, R. H.; von Bergen, M.; Rattei, T.;

 Bendinger, B.; Nielsen, P. H.; Wagner, M., Complete nitrification by Nitrospira bacteria.

 Nature **2015,** *528*, (7583), 504-9.

 2.   van Kessel, M. A. H. J.; Speth, D. R.; Albertsen, M.; Nielsen, P. H.; Op den Camp, H.

 J. M.; Kartal, B.; Jetten, M. S. M.; Lucker, S., Complete nitrification by a single

 microorganism. *Nature* **2015,** *528*, (7583), 555-+.

 3.   Palomo, A.; Jane Fowler, S.; Gulay, A.; Rasmussen, S.; Sicheritz-Ponten, T.; Smets, B.

 F., Metagenomic analysis of rapid gravity sand filter microbial communities suggests novel

 physiology of Nitrospira spp. *ISME J* **2016,** *10*, (11), 2569-2581.

 4.   Pjevac, P.; Schauberger, C.; Poghosyan, L.; Herbold, C. W.; van Kessel, M. A. H. J.;

 Daebeler, A.; Steinberger, M.; Jetten, M. S. M.; Lücker, S.; Wagner, M.; Daims, H., Amo-

 targeted polymerase chain reaction primers for the specific detection and quantification of

 comammox Nitrospira in the environment. *bioRxiv* **2017**.

 5.   Okabe, S.; Satoh, H.; Watanabe, Y., In situ analysis of nitrifying biofilms as determined

 by in situ hybridization and the use of microelectrodes. *Appl Environ Microb* **1999,** *65*, (7),

 3182-3191.

6.   Daims, H.; Nielsen, J. L.; Nielsen, P. H.; Schleifer, K. H.; Wagner, M., In situ

characterization of Nitrospira-like nitrite-oxidizing bacteria active in wastewater treatment

plants. *Appl Environ Microbiol* **2001,** *67*, (11), 5273-84.

7.   Juretschko, S.; Timmermann, G.; Schmid, M.; Schleifer, K. H.; Pommerening-Roser,

A.; Koops, H. P.; Wagner, M., Combined molecular and conventional analyses of nitrifying

bacterium diversity in activated sludge: Nitrosococcus mobilis and Nitrospira-like bacteria as

dominant populations. *Appl Environ Microbiol* **1998,** *64*, (8), 3042-51.

8.   Zilles, J. L.; Peccia, J.; Kim, M. W.; Hung, C. H.; Noguera, D. R., Involvement of

Rhodocyclus-related organisms in phosphorus removal in full-scale wastewater treatment

plants. *Appl Environ Microb* **2002,** *68*, (6), 2763-2769.

9.   Park, H. D.; Whang, L. M.; Reusser, S. R.; Noguera, D. R., Taking advantage of

aerated-anoxic operation in a full-scale University of Cape Town process. *Water Environment*

Research **2006,** *78*, (6), 637-642.

10.   Camejo, P. Y.; Owen, B. R.; Martirano, J.; Ma, J.; Kapoor, V.; Santo Domingo, J.;

McMahon, K. D.; Noguera, D. R., Candidatus Accumulibacter phosphatis clades enriched

under cyclic anaerobic and microaerobic conditions simultaneously use different electron

acceptors. *Water Research* **2016,** *102*, 125-137.

11.   APHA In *Standard methods for the examination of water and wastewater*, Washington,

DC, 2005; Andrew D. Eaton, L. S. C., Eugene W. Rice, Arnold E. Greenberg, Ed. Washington,

DC, 2005.

12.   Peng, Y.; Leung, H. C. M.; Yiu, S. M.; Chin, F. Y. L., IDBA-UD: a de novo assembler

for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics*

2012, *28*, (11), 1420-1428.

13.   Karst, S. M.; Kirkegaard, R. H.; Albertsen, M., mmgenome: a toolbox for reproducible

genome extraction from metagenomes. *bioRxiv* **2016**.

14.   Albertsen, M.; Karst, S. M.; Ziegler, A. S.; Kirkegaard, R. H.; Nielsen, P. H., Back to

Basics--The Influence of DNA Extraction and Primer Choice on Phylogenetic Analysis of

Activated Sludge Communities. *PLoS One* **2015,** *10*, (7), e0132783.

15.   Boetzer, M.; Henkel, C. V.; Jansen, H. J.; Butler, D.; Pirovano, W., Scaffolding pre-

assembled contigs using SSPACE. *Bioinformatics* **2011,** *27*, (4), 578-579.

16.   Parks, D. H.; Imelfort, M.; Skennerton, C. T.; Hugenholtz, P.; Tyson, G. W., CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. *Genome Res* **2015,** *25*, (7), 1043-55.

17.   Hanson, N. W.; Konwar, K. M.; Hawley, A. K.; Altman, T.; Karp, P. D.; Hallam, S. J.,

Metabolic pathways for the whole community. *BMC Genomics* **2014,** *15*, 619.

18.   Richter, M.; Rossello-Mora, R., Shifting the genomic gold standard for the prokaryotic

species definition. *Proceedings of the National Academy of Sciences of the United States of*

America **2009,** *106*, (45), 19126-19131.

19.   Darling, A. E.; Jospin, G.; Lowe, E.; Matsen, F. A. t.; Bik, H. M.; Eisen, J. A., PhyloSift:

phylogenetic analysis of genomes and metagenomes. *PeerJ* **2014,** *2*, e243.

20.   Stamatakis, A., RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses

with thousands of taxa and mixed models. *Bioinformatics* **2006,** *22*, (21), 2688-90.

21.   Clark, K.; Karsch-Mizrachi, I.; Lipman, D. J.; Ostell, J.; Sayers, E. W., GenBank.

Nucleic Acids Research **2016,** *44*, (D1), D67-D72.

22.   Wright, E. S., DECIPHER: harnessing local sequence context to improve protein

multiple sequence alignment. *Bmc Bioinformatics* **2015,** *16*.

23.   Tamura, K.; Stecher, G.; Peterson, D.; Filipski, A.; Kumar, S., MEGA6: Molecular

Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution* **2013,** *30*, (12),

2725-9.

24.   Muller, J.; Muller, K., TREEGRAPH: automated drawing of complex tree figures using

an extensible tree description format. *Molecular Ecology Notes* **2004,** *4*, (4), 786-788.

25.   Li, L.; Stoeckert, C. J.; Roos, D. S., OrthoMCL: Identification of ortholog groups for

eukaryotic genomes. *Genome Research* **2003,** *13*, (9), 2178-2189.

26.   Huerta-Cepas, J.; Szklarczyk, D.; Forslund, K.; Cook, H.; Heller, D.; Walter, M. C.;

Rattei, T.; Mende, D. R.; Sunagawa, S.; Kuhn, M.; Jensen, L. J.; von Mering, C.; Bork, P.,

eggNOG 4.5: a hierarchical orthology framework with improved functional annotations for

eukaryotic, prokaryotic and viral sequences. *Nucleic Acids Research* **2016,** *44*, (D1), D286-

D293.

27.   Bin, Z.; Bin, X.; Zhigang, Q.; Zhiqiang, C.; Junwen, L.; Taishi, G.; Wenci, Z.; Jingfeng,

W., Denitrifying capability and community dynamics of glycogen accumulating organisms

during sludge granulation in an anaerobic-aerobic sequencing batch reactor. *Sci Rep* **2015,** *5*,

12904.

28.   Konstantinidis, K. T.; Tiedje, J. M., Genomic insights that advance the species

definition for prokaryotes. *Proceedings of the National Academy of Sciences of the United*

States of America **2005,** *102*, (7), 2567-2572.

29.	Wu, D. Y.; Jospin, G.; Eisen, J. A., Systematic Identification of Gene Families for Use

as "Markers" for Phylogenetic and Phylogeny-Driven Ecological Studies of Bacteria and

Archaea and Their Major Subgroups. *Plos One* **2013,** *8*, (10).

30.	Keluskar, R.; Desai, A., Evaluation of hydroxylamine oxidoreductase as a functional

and phylogenetic marker to differentiate Nitrosomonas spp. *Journal of Basic Microbiology*

2014, *54*, (4), 261-268.

31.	Junier, P.; Molina, V.; Dorador, C.; Hadas, O.; Kim, O. S.; Junier, T.; Witzel, K. P.;

Imhoff, J. F., Phylogenetic and functional marker genes to study ammonia-oxidizing

microorganisms (AOM) in the environment. *Applied Microbiology and Biotechnology* **2010,**

85, (3), 425-440.

32.	Purkhold, U.; Wagner, M.; Timmermann, G.; Pommerening-Roser, A.; Koops, H. P.,

16S rRNA and amoA-based phylogeny of 12 novel betaproteobacterial ammonia-oxidizing

isolates: extension of the dataset and proposal of a new lineage within the nitrosomonads.

International Journal of Systematic and Evolutionary Microbiology **2003,** *53*, 1485-1494.

33.	Pester, M.; Rattei, T.; Flechl, S.; Grongroft, A.; Richter, A.; Overmann, J.; Reinhold-

Hurek, B.; Loy, A.; Wagner, M., amoA-based consensus phylogeny of ammonia-oxidizing

archaea and deep sequencing of amoA genes from soils of four different geographic regions.

Environmental Microbiology **2012,** *14*, (2), 525-539.

34.	Fitzgerald, C. M.; Camejo, P.; Oshlag, J. Z.; Noguera, D. R., Ammonia-oxidizing

microbial communities in reactors with efficient nitrification at low-dissolved oxygen. *Water*

Res **2015,** *70*, 38-51.

35.   Davey, M. E.; Duncan, M. J., Enhanced biofilm formation and loss of capsule synthesis:

deletion of a putative glycosyltransferase in Porphyromonas gingivalis. *J Bacteriol* **2006,** *188*,

(15), 5510-23.

36.   Li, J. Y.; Wang, N., The gpsX gene encoding a glycosyltransferase is important for

polysaccharide production and required for full virulence in Xanthomonas citri subsp citri. *Bmc*

Microbiology **2012,** *12*.

37.   Tao, F.; Swarup, S.; Zhang, L. H., Quorum sensing modulation of a putative

glycosyltransferase gene cluster essential for Xanthomonas campestris biofilm formation.

Environ Microbiol **2010,** *12*, (12), 3159-70.

38.   Theilacker, C.; Sava, I.; Sanchez-Carballo, P.; Bao, Y.; Kropec, A.; Grohmann, E.;

Holst, O.; Huebner, J., Deletion of the glycosyltransferase bgsB of Enterococcus faecalis leads

to a complete loss of glycolipids from the cell membrane and to impaired biofilm formation.

BMC Microbiol **2011,** *11*, 67.

39.   Wallden, K.; Rivera-Calzada, A.; Waksman, G., Type IV secretion systems: versatility

and diversity in function. *Cell Microbiol* **2010,** *12*, (9), 1203-12.

40.   Klausen, M.; Heydorn, A.; Ragas, P.; Lambertsen, L.; Aaes-Jorgensen, A.; Molin, S.;

Tolker-Nielsen, T., Biofilm formation by Pseudomonas aeruginosa wild type, flagella and type

IV pili mutants. *Mol Microbiol* **2003,** *48*, (6), 1511-24.

41.   Zechner, E. L.; Lang, S.; Schildbach, J. F., Assembly and mechanisms of bacterial type

IV secretion machines. *Philos Trans R Soc Lond B Biol Sci* **2012,** *367*, (1592), 1073-87.

42.   Pascoe, B.; Meric, G.; Murray, S.; Yahara, K.; Mageiros, L.; Bowen, R.; Jones, N. H.;

Jeeves, R. E.; Lappin-Scott, H. M.; Asakura, H.; Sheppard, S. K., Enhanced biofilm formation

and multi-host transmission evolve from divergent genetic backgrounds in Campylobacter

jejuni. *Environ Microbiol* **2015,** *17*, (11), 4779-89.

43.   Bronnec, V.; Turonova, H.; Bouju, A.; Cruveiller, S.; Rodrigues, R.; Demnerova, K.;

Tresse, O.; Haddad, N.; Zagorec, M., Adhesion, Biofilm Formation, and Genomic Features of

Campylobacter jejuni Bf, an Atypical Strain Able to Grow under Aerobic Conditions. *Front*

Microbiol **2016,** *7*, 1002.

44.   Wong, E. H. J.; Ng, C. G.; Chua, E. G.; Tay, A. C. Y.; Peters, F.; Marshall, B. J.; Ho,

B.; Goh, K. L.; Vadivelu, J.; Loke, M. F., Comparative Genomics Revealed Multiple

Helicobacter pylori Genes Associated with Biofilm Formation In Vitro. *Plos One* **2016,** *11*,

(11).

45.   Chipasa, K. B.; Medrzycka, K., Behavior of lipids in biological wastewater treatment

processes. *J Ind Microbiol Biotechnol* **2006,** *33*, (8), 635-45.

46.   Koch, H.; Lucker, S.; Albertsen, M.; Kitzinger, K.; Herbold, C.; Spieck, E.; Nielsen, P.

H.; Wagner, M.; Daims, H., Expanded metabolic versatility of ubiquitous nitrite-oxidizing

bacteria from the genus Nitrospira. *Proc Natl Acad Sci U S A* **2015,** *112*, (36), 11371-6.

47.   Nowka, B.; Off, S.; Daims, H.; Spieck, E., Improved isolation strategies allowed the

phenotypic differentiation of two Nitrospira strains from widespread phylogenetic lineages.

FEMS Microbiol Ecol **2015,** *91*, (3).

48.   Deboer, W.; Laanbroek, H. J., Ureolytic Nitrification at Low Ph by Nitrosospira Spec.

Arch Microbiol **1989,** *152*, (2), 178-181.

49.   Koops, H. P.; Bottcher, B.; Moller, U. C.; Pommereningroser, A.; Stehr, G.,

Classification of 8 New Species of Ammonia-Oxidizing Bacteria - Nitrosomonas-Communis

Sp-Nov, Nitrosomonas-Ureae Sp-Nov, Nitrosomonas-Aestuarii Sp-Nov, Nitrosomonas-
Marina Sp-Nov, Nitrosomonas-Nitrosa Sp-Nov, Nitrosomonas-Eutropha Sp-Nov,
Nitrosomonas-Oligotropha Sp-Nov and Nitrosomonas-Halophila Sp-Nov. *J Gen Microbiol*
1991, *137*, 1689-1699.

50.   Koper, T. E.; El-Sheikh, A. F.; Norton, J. M.; Klotz, M. G., Urease-encoding genes in
ammonia-oxidizing bacteria. *Appl Environ Microb* **2004,** *70*, (4), 2342-2348.

51.   Palatinszky, M.; Herbold, C.; Jehmlich, N.; Pogoda, M.; Han, P.; von Bergen, M.;
Lagkouvardos, I.; Karst, S. M.; Galushko, A.; Koch, H.; Berry, D.; Daims, H.; Wagner, M.,
Cyanate as an energy source for nitrifiers. *Nature* **2015,** *524*, (7563), 105-+.

52.   Qian, M. W.; Eaton, J. W.; Wolff, S. P., Cyanate-mediated inhibition of neutrophil
myeloperoxidase activity. *Biochemical Journal* **1997,** *326*, 159-166.

53.   Purcarea, C.; Ahuja, A.; Lu, T.; Kovari, L.; Guy, H. I.; Evans, D. R., Aquifex aeolicus
aspartate transcarbamoylase, an enzyme specialized for the efficient utilization of unstable
carbamoyl phosphate at elevated temperature. *Journal of Biological Chemistry* **2003,** *278*, (52),
52924-52934.

54.   Ubalua, A. O., Cyanogenic Glycosides and the fate of cyanide in soil. *Australian*
Journal of Crop Science **2010,** *4*, (4), 223-237.

55.   Widner, B.; Mulholland, M. R.; Mopper, K., Chromatographic Determination of
Nanomolar Cyanate Concentrations in Estuarine and Sea Waters by Precolumn Fluorescence
Derivatization. *Analytical Chemistry* **2013,** *85*, (14), 6661-6666.

56.   Ushiki, N.; Fujitani, H.; Aoi, Y.; Tsuneda, S., Isolation of Nitrospira belonging to

Sublineage II from a Wastewater Treatment Plant. *Microbes and Environments* **2013,** *28*, (3),

346-353.

57.   Gruber-Dorninger, C.; Pester, M.; Kitzinger, K.; Savio, D. F.; Loy, A.; Rattei, T.;

Wagner, M.; Daims, H., Functionally relevant diversity of closely related Nitrospira in

activated sludge. *Isme Journal* **2015,** *9*, (3), 643-655.

58.   Koch, H.; Galushko, A.; Albertsen, M.; Schintlmeister, A.; Gruber-Dorninger, C.;

Lucker, S.; Pelletier, E.; Le Paslier, D.; Spieck, E.; Richter, A.; Nielsen, P. H.; Wagner, M.;

Daims, H., Growth of nitrite-oxidizing bacteria by aerobic hydrogen oxidation. *Science* **2014,**

345, (6200), 1052-1054.

59.   Silva, P. J.; de Castro, B.; Hagen, W. R., On the prosthetic groups of the NiFe

sulfhydrogenase from Pyrococcus furiosus: topology, structure, and temperature-dependent

redox chemistry. *J Biol Inorg Chem* **1999,** *4*, (3), 284-91.

60.   Andrews, S. C.; Berks, B. C.; McClay, J.; Ambler, A.; Quail, M. A.; Golby, P.; Guest,

J. R., A 12-cistron Escherichia coli operon (hyf) encoding a putative proton-translocating

formate hydrogenlyase system. *Microbiology* **1997,** *143 (Pt 11)*, 3633-47.

61.   Ferry, J. G., Co Dehydrogenase. *Annual Review of Microbiology* **1995,** *49*, 305-333.

62.   Sant'Anna, F. H.; Lebedinsky, A. V.; Sokolova, T. G.; Robb, F. T.; Gonzalez, J. M.,

Analysis of three genomes within the thermophilic bacterial species Caldanaerobacter

subterraneus with a focus on carbon monoxide dehydrogenase evolution and hydrolase

diversity. *Bmc Genomics* **2015,** *16*.

63.   Stover, C. K.; Pham, X. Q.; Erwin, A. L.; Mizoguchi, S. D.; Warrener, P.; Hickey, M.

J.; Brinkman, F. S. L.; Hufnagle, W. O.; Kowalik, D. J.; Lagrou, M.; Garber, R. L.; Goltry, L.;

Tolentino, E.; Westbrock-Wadman, S.; Yuan, Y.; Brody, L. L.; Coulter, S. N.; Folger, K. R.;

Kas, A.; Larbig, K.; Lim, R.; Smith, K.; Spencer, D.; Wong, G. K. S.; Wu, Z.; Paulsen, I. T.;

Reizer, J.; Saier, M. H.; Hancock, R. E. W.; Lory, S.; Olson, M. V., Complete genome sequence

of Pseudomonas aeruginosa PAO1, an opportunistic pathogen. *Nature* **2000,** *406*, (6799), 959-

964.

64.   Gioia, J.; Qin, X.; Jiang, H. Y.; Clinkenbeard, K.; Lo, R.; Liu, Y. M.; Fox, G. E.;

Yerrapragada, S.; McLeod, M. P.; McNeill, T. Z.; Hemphill, L.; Sodergren, E.; Wang, Q. Y.;

Muzny, D. M.; Homsi, F. J.; Weinstock, G. A.; Highlander, S. K., The genome sequence of

Mannheimia haemolytica A1: Insights into virulence, natural competence, and Pasteurellaceae

phylogeny. *Journal of Bacteriology* **2006,** *188*, (20), 7257-7266.

65.   Brinkrolf, K.; Brune, I.; Tauch, A., The transcriptional regulatory network of the amino

acid producer Corynebacterium glutamicum. *J Biotechnol* **2007,** *129*, (2), 191-211.

66.   Schlensog, V.; Lutz, S.; Bock, A., Purification and DNA-binding properties of FHLA,

the transcriptional activator of the formate hydrogenlyase system from Escherichia coli. *J Biol*

Chem **1994,** *269*, (30), 19590-6.

67.   Toesca, I.; Perard, C.; Bouvier, J.; Gutierrez, C.; Conter, A., The transcriptional

activator NhaR is responsible for the osmotic induction of osmC(p1), a promoter of the stress-

inducible gene osmC in Escherichia coli. *Microbiology-Sgm* **2001,** *147*, 2795-2803.

68.   Goller, C.; Wang, X.; Itoh, Y.; Romeo, T., The cation-responsive protein NhaR of

Escherichia coli activates pgaABCD transcription, required for production of the biofilm

adhesin poly-beta-1,6-N-acetyl-D-glucosamine. *J Bacteriol* **2006,** *188*, (23), 8022-32.

69. Storz, G.; Tartaglia, L. A., OxyR: a regulator of antioxidant genes. *J Nutr* **1992,** *122*,

(3 Suppl), 627-30.

70. Bueno, E.; Mesa, S.; Bedmar, E. J.; Richardson, D. J.; Delgado, M. J., Bacterial

adaptation of respiration from oxic to microoxic and anoxic conditions: redox control. *Antioxid*

Redox Signal **2012,** *16*, (8), 819-52.

71. Green, J.; Crack, J. C.; Thomson, A. J.; LeBrun, N. E., Bacterial sensors of oxygen.

Curr Opin Microbiol **2009,** *12*, (2), 145-51.

72. Green, J.; Scott, C.; Guest, J. R., Functional versatility in the CRP-FNR superfamily of

transcription factors: FNR and FLP. *Adv Microb Physiol* **2001,** *44*, 1-34.

73. Reents, H.; Munch, R.; Dammeyer, T.; Jahn, D.; Hartig, E., The Fnr regulon of Bacillus

subtilis. *J Bacteriol* **2006,** *188*, (3), 1103-12.

74. Salmon, K.; Hung, S. P.; Mekjian, K.; Baldi, P.; Hatfield, G. W.; Gunsalus, R. P.,

Global gene expression profiling in Escherichia coli K12. The effects of oxygen availability

and FNR. *J Biol Chem* **2003,** *278*, (32), 29837-55.

75. Park, H. D.; Noguera, D. R., Nitrospira community composition in nitrifying reactors

operated with two different dissolved oxygen levels. *J Microbiol Biotechnol* **2008,** *18*, (8),

1470-4.

76. Padan, E.; Maisler, N.; Taglicht, D.; Karpel, R.; Schuldiner, S., Deletion of ant in

Escherichia coli reveals its function in adaptation to high salinity and an alternative Na+/H+

antiporter system(s). *J Biol Chem* **1989,** *264*, (34), 20297-302.

77.   Conter, A.; Gangneux, C.; Suzanne, M.; Gutierrez, C., Survival of Escherichia coli

during long-term starvation: effects of aeration, NaCl, and the rpoS and osmC gene products.

Research in Microbiology **2001,** *152*, (1), 17-26.

78.   Mongkolsuk, S.; Praituan, W.; Loprasert, S.; Fuangthong, M.; Chamnongpol, S.,

Identification and characterization of a new organic hydroperoxide resistance (ohr) gene with

a novel pattern of oxidative stress regulation from Xanthomonas campestris pv. phaseoli. *J*

Bacteriol **1998,** *180*, (10), 2636-43.

79.   Itoh, Y.; Wang, X.; Hinnebusch, B. J.; Preston, J. F., 3rd; Romeo, T., Depolymerization

of beta-1,6-N-acetyl-D-glucosamine disrupts the integrity of diverse bacterial biofilms. *J*

Bacteriol **2005,** *187*, (1), 382-7.

80.   Bren, A.; Welch, M.; Blat, Y.; Eisenbach, M., Signal termination in bacterial

chemotaxis: CheZ mediates dephosphorylation of free rather than switch-bound CheY. *Proc*

Natl Acad Sci U S A **1996,** *93*, (19), 10090-3.

81.   Khan, S.; Amoyaw, K.; Spudich, J. L.; Reid, G. P.; Trentham, D. R., Bacterial

chemoreceptor signaling probed by flash photorelease of a caged serine. *Biophys J* **1992,** *62*,

(1), 67-8.

82.   Unden, G.; Schirawski, J., The oxygen-responsive transcriptional regulator FNR of

Escherichia coli: the search for signals and reactions. *Mol Microbiol* **1997,** *25*, (2), 205-10.

83.   Spiro, S.; Guest, J. R., FNR and its role in oxygen-regulated gene expression in

Escherichia coli. *FEMS Microbiol Rev* **1990,** *6*, (4), 399-428.

84.   Li, B.; Wing, H.; Lee, D.; Wu, H. C.; Busby, S., Transcription activation by Escherichia

coli FNR protein: similarities to, and differences from, the CRP paradigm. *Nucleic Acids Res*

1998, *26*, (9), 2075-81.