1

Morphological and transcriptomic evidence for ammonium induction of sexual reproduction in *Thalassiosira pseudonana* and other centric diatoms

Eric R. Moore[1], Briana S. Bullington[1], Alexandra J. Weisberg[2], Yuan Jiang[3], Jeff Chang[2], Kimberly H. Halsey[1]*

[1]Department of Microbiology, Oregon State University, Corvallis, Oregon 97331, USA

[2]Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon 97331, USA

[3]Department of Statistics, Oregon State University, Corvallis, Oregon 97331, USA

* Corresponding Author: halseyk@science.oregonstate.edu

Abstract:

The reproductive strategy of diatoms includes asexual and sexual phases, but in many species,

including the model centric diatom *Thalassiosira pseudonana*, sexual reproduction has never

been observed. Furthermore, the environmental factors that trigger sexual reproduction in

diatoms are not understood. Although genome sequences of a few diatoms are available, little is

known about the molecular basis for sexual reproduction. Here we show that ammonium reliably

induces the key sexual morphologies, including oogonia, auxospores, and spermatogonia, in two

strains of *T. pseudonana*, *T. weissflogii*, and *Cyclotella cryptica*. RNA sequencing revealed 1,274

genes whose expression patterns changed when *T. pseudonana* was induced into sexual

reproduction by ammonium. Some of the induced genes are linked to meiosis or encode flagellar

structures of heterokont and cryptophyte algae. The identification of ammonium as an

environmental trigger suggests an unexpected link between diatom bloom dynamics and

strategies for enhancing population genetic diversity.

Introduction:

Diatoms are protists that form massive annual spring and fall blooms in aquatic environments and are estimated to be responsible for about half of photosynthesis in the global oceans [1]. This predictable annual bloom dynamic fuels higher trophic levels and initiates delivery of carbon into the deep ocean biome. Diatoms have complex life history strategies that are presumed to have contributed to their rapid genetic diversification into ~200,000 species [2] that are distributed between the two major diatom groups: centrics and pennates [3]. A defining characteristic of all diatoms is their restrictive and bipartite silica cell wall that causes them to progressively shrink during asexual cell division. At a critically small cell size and under certain conditions, auxosporulation restitutes cell size and prevents clonal death [4-6]. The entire lifecycles of only a few diatoms have been described and rarely have sexual events been captured in the environment [7-9].

So far, all centric diatoms appear to share the process of oogamous sexual reproduction (Fig 1). The average cell size of a population of asexually dividing diatoms decreases as a result of differential thecae inheritance. At a critically small size, cells become eligible to differentiate into male and female cells. Meiosis in the male spermatogonangium produces multinucleate spermatogonia that divide into individual haploid spermatocytes. Meiosis in the female oogonia produces a single functional haploid nucleus that is fertilized by a flagellated spermatocyte through an opening in the oogonia thecae. Fertilized oogonia expand into a large auxospore where new, large thecae are formed for the new, enlarged initial cell. Auxosporulation can also occur asexually, but it is considered an ancillary pathway for cell size restitution in diatom species that have a sexual path for reproduction [5].

The environmental factors that trigger formation of sexual cells and sexual reproduction

in centric diatoms are not well understood [10, 11], but sexualization appears to be strongly

associated with conditions causing synchronous sexuality in cells experiencing growth stress

[12]. Besides the size threshold requirement, previous observations indicate that sexualization is

possible when active growth has ceased, causing cell cycle arrest [13, 14] and cell densities are

sufficient to permit successful fertilization of the oogonia by the spermatocyte [15]. Light

interruption with an extended dark period [13], changing salinities, and nutrient shifts [16], have

sometimes been successful in inducing sexual reproduction, probably by causing cell cycle

arrest. Recently, pheromones produced by the pennate diatom, *Seminavis robusta,* have been

identified that cause cell cycle arrest and induce the sexual pathway [17]. However, we are aware

of no method that reliably causes induction of all of the sexual stages of centric diatoms shown

in figure 1.

The ecological importance of diatoms, combined with their potential uses in materials

chemistry, drug delivery, biosensing [18, 19], and bioenergy [20, 21], prompted genome

sequencing of *T. pseudonana* CCMP1335 (a 'centric' diatom collected from the North Atlantic

Ocean) and *Phaeodactulum tricornutum* (a 'pennate' diatom), which have become model

organisms for experimental studies [22, 23]. However, sexual morphologies have never been

observed in either of these species or in the vast majority of diatoms [10]. The inability to

reliably control the sexual cycle in centric diatoms has severely hindered studies to understand

the silica deposition process, as well as the genetic regulation, ecology, and evolution of sex [10,

24, 25]. Both of the model diatoms were thought to have repurposed their extant genetic toolkits

and lost the need and ability for a sexual lifestyle [10, 11, 26].

Here we show that two strains of *T. pseudonana* and two other centric species, *T.*

weisflogii and *Cyclotella cryptica,* can be reliably induced into the sexual reproductive pathway

when cells are below the critical size threshold and exposed to ammonium during the stationary

phase of growth. Ammonium induced oogonia, auxospore, and spermatocyte formation in each

of these species. Induction of sexuality was further supported by RNA sequencing (RNAseq)

which revealed 1,274 genes whose expression patterns changed when *T. pseudonana* became

sexualized by ammonium. Meiosis genes and genes associated with flagellar structures of

heterokont and cryptophyte algae were differentially expressed in ammonium-induced cells

compared to nitrate grown cells. We anticipate that this discovery will open opportunities to

study the evolution of diatom lifecycles and facilitate expansion of diatom breeding to explore

functional genetics for molecular ecology, nanotechnology and biofuels applications.

Results and Discussion:

Ammonium triggers sexual morphologies

We observed *T. pseudonana* CCMP1335 cell morphologies consistent with sexual

reproduction when cells were propagated in artificial seawater medium supplemented with

ammonium. The proportion of cells that differentiated into sexual cell types was dependent on

ammonium concentration, with up to 39% of the population identified as oogonia or auxospores

in cultures supplemented with 800 μM NH_4Cl (Fig 2A). Oogonia and auxospores were first

observed at the onset of stationary phase and reached maximum population proportions in late

stationary phase (Fig 2A). Ammonium also induced oogonia and auxospore production in *T.*

pseudonana CCMP1015 (collected from the North Pacific Ocean), *T. weissflogii*, and *Cyclotella*

cryptica (Fig S1). A few oogonia and auxospores were observed in nitrate grown cultures with

no added ammonium (Fig 2A and S1). However, with the exception of nitrate grown *C. cryptica*

cultures, which generated oogonia and auxospores constituting 11% of the total population,

oogonia and auxospores were only a small percentage of the total population in nitrate-grown *T.*

pseudonana and *T. weissflogii.* Even though auxospores can have diameters 3-4 times that of

asexual cells, such small population proportions do not lead to discernable shifts in cell size

distributions obtained by particle size analysis (e.g., Coulter counter, a commonly used method

to assess population size). We initially observed auxospores when performing visual inspections

of our cultures. For the data reported here, oogonia and auxospores were quantified by manually

counting the cell types using a hemocytometer. We suspect that reliance on laboratory

instruments such as particle counters and flow cytometers in place of microscopic analysis is one

reason that sexual morphologies in these well-studied diatom species have gone undetected until

now. Laboratory stock cultures are typically maintained in media with low concentrations of

nitrogen, especially when ammonium is supplied as the nitrogen source because it has been

considered to be toxic to diatoms in high concentrations [27]. Therefore, it may not be surprising

that sexual cells have gone un-noticed due to the low rates of sexual induction in the presence of

low ammonium concentrations (Fig. 2A).

Cells differentiated into oogonia and auxospores regardless of whether ammonium was

supplied at inoculation or at the onset of stationary phase (Fig. 1B). Thus, it appears that

stationary phase and ammonium availability are key factors that trigger formation of sexual cells

in centric diatoms. Resource depletion can arrest the cell cycle [14], and the presence of

ammonium at the onset of stationary phase appears to activate the sexual cycle. Auxospore

formation was observed in *Cyclotella meneghiniana* during stationary phase [28], and other

protists initiate sex under stress in response to nutrient depletion or oxidative DNA damage [29].

Ammonium can inhibit photosynthesis [27]; however, diatoms, including *T. pseudonana*, can

acclimate to millimolar ammonium concentrations [30]. It is possible that high ammonium

concentrations intensify the stress condition required for the sexual pathway. Nevertheless,

ammonium consistently caused formation of ten-fold more sexualized cells than the same

concentrations of nitrate (Fig 2 and S1).

Our results showing that ammonium induced formation of sexual cells in several centric

diatom species suggests that it may serve as a key environmental factor regulating the sexual

lifecycle across centric diatoms. Ammonium is typically present in very low concentrations in

aquatic ecosystems. However, ammonium reached 12.6 mM in a eutrophic lake where the centric

diatom, *A. subarctica,* was observed undergoing sexual reproduction [8]. Clearly, ammonium

was not the growth-limiting nutrient under those conditions or in our laboratory cultures (Fig 2).

Pseudo-nitzschia auxospore formation was positively correlated with ammonium, which was

measured to be 14 μM during a major bloom event off the coast of Washington [7]. Thus, the

formation of sexual cells appears to be triggered by the presence of ammonium while at least one

other growth factor becomes limiting, such as light (discussed below), phosphorous, silica [7],

vitamins, or trace elements.

Cell differentiation in *T. pseudonana* was induced irrespective of growth rate in

exponential phase, light intensity, or light regime. However, the growth parameters did affect the

proportion of differentiated cells. Oogonia and auxospores were only 0.5% of the population

when grown under very low light (5 μE) with 200 μM NH_4Cl and increased to 39% when grown

under moderate light (100 μE) and 800 μM NH_4Cl (Table 1). The proportions of oogonia and

auxospores increased with light up to moderate intensities (70-100 μE) but decreased at high

(220 µE) intensities (Table 1), suggesting that photon flux has an important role in meeting the

energetic demands of sexual reproduction. Other work showed sexualization was more prevalent

at light <50 µE [31] or with the addition of a dark period [13, 32, 33]. Likely, the optimum light

intensity or need for a dark period to precede sexual induction [34] is species-specific and linked

to adaptive life histories [5].

Visualization of sexual morphologies

Confocal, light, and scanning electron microscopy were used to document the cell

morphologies at various stages in the life history cycle of *T. pseudonana* (Fig 3 and S2). Oogonia

were elongated relative to asexually growing vegetative cells and exhibited a bent morphology

and swelling of the plasma membrane at the junction of the hypotheca and epitheca (Figs 3B, 3C

and S2B-D). In oogonia, cellular contents became localized to the ends of the cell resulting in an

apparent empty space near the area of membrane swelling where fertilization likely occurs [5].

T. pseudonana spermatogonium harbored at least eight nuclei (Fig 3D), suggesting that a

depauperating mitosis preceded meiosis [11, 35]. Sperm released were very small, about 1 µm,

and flagellated (Figs 3E and S2E, S2F), but they often became entangled with other cells and

debris [8, 36, 37]. Sperm cells attached to oogonia at the junction of the thecae for fertilization

(Figs 3F, 3G and S2I, S2J) as shown in *T. punctigera* [38]. Also similar to *T. punctigera*, flagella

were not visible at that stage, possibly because flagella are abandoned upon attachment to the

oogonia [36].

Oogonia developed into auxospores and these conspicuous cell morphologies were

always observed in cultures induced by ammonium. Auxospores were larger than vegetative

cells and oogonia, ranging from about 6 to 20 μm in diameter, with most being 10-15 μm in

diameter (Figs 3H, 3I and S2G, S2H, S2L). Auxospores were spherical, with most of the cellular

contents localized to one side (Fig 3H, 3I and S2G, S2H, S2L) and sometimes showing slight

distention where the mother valve was shed (Fig 3I), as described in *Stephanodiscus niagarae*

[39]. Thecae remained attached in some cases, especially on smaller cells. Oogonia, auxospores,

and spermatogonia in the other species studied displayed similar morphologies to those observed

in *T. pseudonana* and other centric diatoms (Figs 3L-O, S2M-P and 4A) [11, 16, 28, 33, 35, 37,

38, 40, 41].

Changes in DNA content in *T. pseudonana* cells induced into sexuality by ammonium

were observed using flow cytometry-based analysis. Fluorescence-activated cell sorting (FACS)

analysis showed that as the culture progressed from exponential into stationary growth phases,

the diploid population (Fig 4B) expanded to include DNA fluorescence intensities that were

consistent with the presence of spermatogonangia and spermatogonia containing multiple

gametes (Fig 4C). In late stationary phase, a new population was observed that had DNA

fluorescence signals consistent with haploid sperm cells with little to no chlorophyll [13] (Fig

4D).

Cell size restitution via auxosporulation produced initial cells that were considerably

larger than the parent cells that were maintained in nitrate and already at or below the critical

size threshold for induction into sexuality. We serially transferred ammonium-induced cultures

with high proportions of auxospores to generate *T. pseudonana*, *T. weissflogii*, and *C. cryptica*

cultures that were enriched in initial cells. The average cell diameters of the resulting cultures

were larger relative to stock cultures maintained in nitrate (Fig 4E and S3). The *T. pseudonana*

initial cells were 7-12 μm, the largest size reported for this species (Figs 3J, 3K, S2K).

We identified oogonia, male gametes, auxospores, and initial cells in cultures of the

model centric diatom, *T. pseudonana* providing new evidence for sexuality in this species that

was previously assumed to be asexual [10]. Although cell enlargement through

asexual/apomictic mechanisms has been recorded in other species [42-44], the presence of all

sexual cell types, and the expression of meiotic genes (discussed below), suggest apomixis is not

the mechanism being used by *T. pseudonana* for cell enlargement. Furthermore, apomixis

typically occurs in species that also undergo sexual reproduction [5]. Only spermatogenesis had

previously been reported in *T. weissflogii* [13, 45], but we have now also documented induction

of oogonia and auxospores by ammonium and subsequent formation of initial cells in this

species. A major challenge in visualizing the morphological characteristics of these species is

their smaller cell sizes compared to other species for which morphological details have been

documented. Now that we have determined a reliable method for inducing the sexual

morphologies, future studies will dissect additional details associated with the sexual pathways

in these, and perhaps other species inducible by ammonium, to determine their variation from

other centric diatoms. For example, the presence of auxospore scales, precise timing of

fertilization and meiotic activity, repeated auxosporulation, and polyspermy events (e.g., [38]).

Gene expression analysis of ammonium induced sexual morphologies

We used RNAseq to identify genes that were differentially expressed in conditions that

triggered cell differentiation into sexual morphologies. We compared the transcriptomes of *T.*

pseudonana harvested in exponential (EXP), stationary (STA), and late stationary phases (L-

STA). Cells were grown in 100 μM NaNO$_3$ or, to capture a dose-dependent change in gene

expression, either 100 or 800 μM NH$_4$Cl (Fig S4). We identified genes that were significantly

differentially expressed in multiple pairwise comparisons of growth phases and nitrogen sources

(Tables S1-S11). Next, we examined the statistical interactions of pairwise condition

comparisons to identify genes with significantly greater or lesser magnitude changes in

expression between growth stages in the presence of ammonium relative to 100 μM NaNO$_3$ (Fig

5A and S5).

This conservative approach yielded a total of 1,274 genes in the four analyses of

statistical interactions (Supplemental Tables S12-S15). A total of 89 of the genes have an

annotated function (Fig. 5A; Table S16). The set of 89 genes includes four meiotic genes (*mcm2,*

mcm8, mcm9, and mlh1, Fig 4B and S6) that were also up-regulated in pennate diatoms during

sexual reproduction [26]. The Mcm family of DNA helicases function in DNA replication, with

mcm8, mcm9, and *mlh1*, having roles in double stranded break repair [46-48]. Mcm8 was one of

the four genes related to meiosis that were upregulated in the pennate diatom, *Seminavis robusta*

during treatment with a sex-inducing pheromone [17]. Eight genes in our list are homologous to

yeast genes involved in meiosis [49] (Fig 4B, Table S17), including genes that regulate the

meiotic anaphase promoting complex (*cdc16*, *cdc23*, *ama1*) [50, 51] and *rad54*, a motor protein

that regulates branch migration of Holliday junctions during homologous recombination [52].

Expression of genes encoding other RAD proteins (*rad50 and rad 51*) increased in pennate

diatoms induced into meiosis [17, 26].

Of three 'sexually induced genes' that were up-regulated in *T. weissflogii* at the initiation

of gametogenesis [45] and associated with sperm flagella mastigonemes [53], one, *sig3*, was

significantly up-regulated in stationary phase compared to exponential phase (Fig 4B). In

addition, a gene encoding an intraflagellar transport protein (IFT88) was also up-regulated in

ammonium induced cells during stationary phase (Fig. 4B). An IFT system is required for

flagellar assembly [54] and five genes encoding IFT particle proteins, including IFT88, and a

kinesin-associated protein involved in anterograde transport were found in the *T. pseudonana*

genome [55]. The genes encoding Sig3 and IFT88 are unique to flagellar structures, and their

differential expression in ammonium induced *T. pseudonana* compared to the nitrate-grown

control treatments provide additional evidence that ammonium induced spermatogenesis in this

species.

The MYB factor and bZIP families of proteins are transcriptional regulators that control a

variety of cell processes including stress responses, development, and differentiation in plants

[56]. Expression of two genes having the characteristic R2R3 MYB DNA binding domains

common in plants, *myb24 and myb16,* was generally lower in ammonium induced cultures

compared to the nitrate grown controls (Fig 4B). In *Arabidopsis thaliana*, hormonal activation of

myb24 is required for stamen development and male fertility [57]. Whether *myb24, myb16,* and

bzip2 play roles in regulating gamete development or sex differentiation in diatoms remains to be

determined.

The 1,274 genes provide new avenues to understand the evolution of sexuality in the

Heterokont eukaryotic lineage. Diatoms emerged ~ 200 Mya, about 800 My after a eukaryotic

heterotroph engulfed a red alga in the secondary endosymbiosis event that gave rise to the SAR

eukaryotic supergroup [58]. Of 171 diatom genes of red algal origin [58], 17 were identified as

differentially expressed in conditions that induced sexual reproduction (Table S18). None of

these genes are annotated in the *T. pseudonana* genome, but in red algae they are predicted to

function in transport and plastid-targeted processes [23].

Conclusions:

That some of the most well studied centric diatoms were never observed undergoing sexual reproduction was a mystery. Possibly even more elusive was the ability to reliably control or induce the sexual pathway of centric diatoms in the laboratory [10] despite a myriad of efforts that ranged from sweet-talk to torture. Factors that have limited progress in this field center on the problem that even under 'favorable environmental conditions' that result in the sexual lifecycle, only a fraction of cells undergo sexual reproduction (Fig 2). Thus, capturing the sexual event requires near constant visual observation because (a) only cells that have become sufficiently small and reach the critical size threshold can undergo sexual reproduction [59], (b) only a fraction of those size-eligible cells may undergo sexual reproduction [8, 15, 16, 31], (c) there has been a lack of understanding about what constitutes conditions that are 'favorable' for triggering diatom sex [5, 7], and (d) morphological changes indicative of sex may not be recognized by untrained scientists [60].

Our results provide strong evidence that *T. pseudonana* is a sexual organism, expressing the major morphologies associated with the sexual pathway that result in enlarged initial cells. Furthermore, the sexual pathway was reliably induced in *T. pseudonana*, and two other centric diatom species by exposure of size-eligible cells to ammonium. Ammonium triggered formation of sexual cells in a dose-dependent manner and significant changes in expression of genes involved in meiosis, spermatocyte flagellar structures and assembly, and sex differentiation. RNAseq analysis revealed many more genes with unknown functions that were expressed under conditions of sexual differentiation. Other genes involved in sex are likely to have been missed by our analysis because their changes in expression were masked by the mixed population of asexual and sexual cells, or they were not captured in the coarse time-resolution of sampling

used in this study. Nevertheless, our discoveries resolve two persistent mysteries that have

plagued diatom researchers. Furthermore, the RNAseq data provide a subset of genes that can be

used to study the molecular ecology of diatoms.

The ecology of centric diatom sexual reproduction that can be inferred from our findings

appears best described as synchronous sexuality [12] triggered by ammonium in cells

experiencing growth stress. Asexual cell cycle arrest appears to be prerequisite to activation of

the diatom sexual life cycle [13, 14, 28, 29]. In the environment, diatoms bloom following

elevated nutrient concentrations driven by vertical mixing, coastal upwelling, or river inputs and

the bloom reaches its peak biomass when essential nutrients are depleted. Within a week, the

bulk of a bloom can be consumed by heterotrophic protists [61] that excrete ammonium to

maintain homeostatic elemental composition [62]. We propose that ammonium released by

grazers at bloom climax may be a principal ecological trigger for sexual morphologies in centric

diatoms. Synchronization of sexuality at the onset of resource depletion (stationary phase)

increases the chances for successful fertilization because cell density is at its maximum [12].

Environmental concentrations of ammonium in the environment rarely reach the concentrations

used in this study to demonstrate the dose response effect on sexuality. Other methods that have

sometimes successfully triggered sexual reproduction in other species are similarly unusual

compared to environmental conditions. For example, the magnitude of the salinity shifts used to

induce sexual reproduction in *Skeletonema marioni* in the laboratory do not occur in the Baltic

sea [16]. Nevertheless, pulses of ammonium, shifts in salinity, and other environmental

fluctuations do occur in aquatic ecosystems, and provided the other conditions for sexuality are

met (e.g., cell size threshold, stress, population density), are likely to induce sexuality in at least

a small fraction of a population. The presence of ammonium and the onset of stationary phase

also point to involvement of another growth factor whose depletion triggers sexual reproduction.

The specific collection of factors that lead to sexual reproduction in diatoms in the environment

is not yet known and neither is whether ammonium is a direct or indirect trigger of sexuality [4].

Nevertheless, this work suggests an intriguing ecological role for ammonium in the mechanisms

underlying sexuality in centric diatoms and will certainly be a valuable tool to control sexuality

in the laboratory.

The identification of ammonium as a reliable inducer of sexuality in *T. pseudonana* and

other centric diatoms has the potential to shift perspectives on diatom ecology, open avenues for

the experimental investigation of diatom reproductive mechanisms, and provide tools for genetic

manipulation of centric diatoms that have not heretofore been available. Diatom blooms have a

global impact but the factors that control these blooms and their demise are complex and a

consensus has not been reached about these processes. Our evidence suggests that induction of

sexuality may play a vital role in diatom bloom conclusion and the production of genetic

diversity that seeds future blooms [63]. Our analysis suggests an involvement of genes of red-

algal origin, providing new lines of evolutionary enquiry. Interest in diatoms for biotechnological

applications is high due to their uses in biofuels, materials chemistry and medicine. Our work

will likely propel this exploration by enabling improved breeding and genetic modification to

control and understand unique diatom traits.

<u>Methods</u>

Stock cultures of *T. pseudonana* (CCMP1335) were maintained in f/2 medium [64] with

200 μM NaNO$_3$ under continuous sub-saturating light at 18°C. Sexual cells were quantified in

triplicate cultures of *T. pseudonana* (CCMP1335 and CCMP1015)*, T. weissflogii* (CCMP1336)

and *C. cryptica* (CCMP332) (all obtained from NCMA) grown in f/2 amended with $NaNO_3$ or

NH_4Cl and grown at 18^oC under 50 μE continuous light, or under variable light intensities/cycles

as shown in Table 1. Cell populations were quantified using a Coulter counter (Beckman-

Coulter, Indianapolis, Indiana). Oogonia and auxospores were counted using a hemocytometer.

We found that a modified f/2 medium yielded better cell images using light and confocal

microscopy. This medium contained 0.939 mM KCl, 0.802 mM NO_3^-, 1 mM NH_4Cl, 0.05 mM

glycine, 0.01 mM methionine, 0.078 mM pyruvate, 0.84 μM pantothenate, 0.985 μM 4-amino-5-

hydroxymethyl-2-methylpyrimidine, 0.3 μM thiamine, 0.002 μM biotin, 0.117 μM $FeCl_3*6H_2O$,

0.009 μM $MnCl_2*4H_2O$, 0.0008 μM $ZnSO_4*7H_2O$, 0.0005 μM $CoCl_2*6H_2O$, 0.0003 μM

$Na_2MoO_4*2H_2O$, 0.001 μM Na_2SeO_3, and 0.001 μM $NiCl_2*6H_2O$, and sparged with filter-

sterilized carbon dioxide and air for 8 hours and overnight respectively. To view DNA, 1 ml live

samples were stained with 5 μl 1.62 μM Hoescht 33342 (0.2 μm filtered) for 10 min. For

scanning electron microscopy (SEM), 1 ml samples were diluted 1:3 with sterile f/2 and syringe

filtered onto 13 mm 0.2 μm polycarbonate filters using a Swinnex filter unit. The filter was

washed with 4 ml f/2 containing 0.5% gluteraldehyde and left submerged for at least 24 hours,

followed by a series of 4 ml washes: 0.2 μm filtered 80%, 60%, 40%, 20% and 0% f/2, followed

by 20%, 40%, 60% 80% and 100% ethanol, before critical point drying. SEM imaging was done

at the Oregon State University Electron Microscope facility.

For flow cytometry 1 ml culture samples were fixed with 1 μl gluteraldehyde (50%) and

stained with 10 μl Sybr green mix (1:25 dilution Sybr green in 0.01M Tris-EDTA, pH 8.0) for 30

min. Samples were run on a FACScan flow cytometer (Becton Dickinson, Franklin Lakes, New

Jersey). Settings were FL1=582 and FL3=450 for unstained cells and FL1=450 and FL3=450 for

stained cells.

For RNAseq analysis, $1.61 \times 10^8 - 1.10 \times 10^9$ cells from triplicate independent cultures

were filtered onto 0.8 μm 47 mm polycarbonate filters during exponential, stationary and late

stationary phases and flash frozen in liquid nitrogen. RNA was extracted using a Qiagen RNeasy

midi kit according to modified manufacturer's instructions [65]. Silica beads (0.5 mm) were

added to the cells and lysis buffer and vortexed until homogeneous before being filtered through

Qiashredder columns to remove large particles. Eluted RNA was subjected to off column RNase

free DNase I treatment and secondary purification according to manufacturer's

recommendations. Total RNA was prepared and sequenced as a 150 bp single end library on an

Illumina HiSeq 3000 at the Center for Genome Research and Biocomputing at Oregon State

University. Sequencing data/interaction analyses were conducted using the Ballgown pipeline

[66]. Sequencing reads were trimmed to remove sequencing adapters using BBDuk v. with the

parameters " ktrim=r k=23 mink=9 hdist=1 minlength=100 tpe tbo" [67]. Reads aligned to the *T.*

pseudonana reference genome (NCBI accession GCA_000149405.2) using HISAT2 v. 2.0.4

with the parameters " --min-intronlen 20 --max-intronlen 1500 --rna-strandness F --dta-cufflinks"

[68]. Transcripts were assembled for each dataset and merged using Stringtie v 1.2.4 [69].

Pairwise differential expression analyses for genes were performed using the "stattest" function

in Ballgown version 2.2.0 [70]. Interaction effects were tested by comparing the models with

(timepoint + treatment + timepoint * treatment) and without (timepoint + treatment) the

interaction term using the custom model option in the "stattest" function.

For construction of the phylogenetic tree, 18s rRNA sequences were obtained from the

Silva database and aligned using Muscle v3.8.31 (default settings) [71]. A genome editor

(BioEdit) was used to manually trim off overhanging sequence. The tree was built using

RAxML-HPC v8.0.26 using the GTRCAT model, "-f a" option, and 1000 bootstrap replicates

[72]. A visual representation was created using the TreeDyn [73] tool through LIRMM

(phylogeny.fr) [74].

Acknowledgements: The authors thank Kelsey McBeain for assistance with confocal

microscopy imaging and the Electron Microscope Facility at Oregon State University for

scanning electron micrographs.

REFERENCES:

1.   Nelson DM, Tréguer P, Brzezinski MA, Leynaert A, Quéguiner B. Production and dissolution of biogenic silica in the ocean: Revised global estimates, comparison with regional data and relationship to biogenic sedimentation Biogeochem Cycles. 1995;9(3):359-72.

2.   Mann DG. The species concept in diatoms. Phycologia. 1999;38:437-95.

3.   Simonsen R. The diatom system: Ideas on phylogeny. Bacillaria. 1979;2:9-71.

4.   Lewis WM. The diatom sex clock and its evolutionary importance. Amer Naturalist. 1984;123:73-80.

5.   Chepurnov VA, Mann DG, Sabbe K, Vyverman W. Experimental studies on sexual reproduction in diatoms. In: Jeon KW, editor. A Survey of Cell Biology. International Review of Cytology. 237. London, UK: Elsevier Academic Press; 2004. p. 91-154.

6.   Drebes. Sexuality. In: Werner D, editor. The Biology of Diatoms. Botanical Monographs. 13. Oxford: Blackwell Scientific Publications; 1977. p. 250-83.

7.   Holtermann KE, Bates SS, Trainer VL, Odell A, Armbrust EV. Mass sexual reproduction in the toxigenic diatoms *Pseudo-Nitzscia australis* and *P. pungens* (Bacillariophyceaea) on the Washington coast, USA. J Phycol. 2010;46:41-52.

8.   Jewson DH. Size reduction, reproductive strategy and the life cycle of a centric diatom. Philos T R Soc B. 1992;336:191-213.

9.   Koester JA, Brawley SH, Karp-Boss L, Mann DG. Sexual reproduction in the marine centric diatom *Ditylum brightwellii* (Bacillariophyta). Eur J Phycol. 2008;42(4):351-66.

10.   Chepurnov VA, Mann DG, von Dassow P, Vanormelingen P, Gillard J, Inze D, et al. In search of new tractable diatoms for experimental biology. BioEssays. 2008;30:692-702.

11. Von Dassow P, Montresor M. Unveiling the mysteries of phytoplankton life cycles: patterns and opportunities behind complexity. J Plankton Res. 2011;33(1):3-12.

12. Edlund MB, Stoermer EF. Ecological, evolutionary, and systematic significance of diatom life histories. J Phycol. 1997;33:897-918.

13. Armbrust EV, Chisholm SW, Olson RJ. Role of light and the cell cycle on the induction of spermatogenesis in a centric diatom. J Phycol. 1990;26(3):470-8.

14. Huysman MJJ, Vyverman W, De Veylder L. Molecular regulation of the diatom cell cycle. J Exp Bot. 2014;65(10):2573-84.

15. Assmy P, Henjes J, Smetacek V, Montresor M. Auxospore formation in the silica-sinking oceanic diatom *Fragilariopsis kerguelensis* (Bacillariophyceae). J Phycol. 2006;42:1002-6.

16. Godhe A, Kremp A, Montresor M. Genetic and microscopic evidence for sexual reproduction in the centric diatom *Skeletonema marinoi*. Protist. 2014;165(4):401-16.

17. Moeys S, Frenkel J, Lembke C, Gillard JTF, Devos V, Van den Berge K, et al. A sex-inducing pheromone triggers cell cycle arrest and mate attraction in the diatom Seminavis robusta. Sci Reports. 2016;6:19252.

18. Ezzati J, Dolatabadi N, de la Guardia M. Applications of diatoms and silica nanotechnology in biosensing, drug and gene delivery, and formation of complex metal nanostructures. J Analyt Chem. 2011;30(9):1538-48.

19. Kroth P. Molecular biology and the biotechnological potential of diatoms. Adv Exp Med Biol. 2007;616:23-33.

20. d'Ippolito G, Sardo A, Paris D, Vella FM, Adelfi MG, Botte P, et al. Potential of lipid metabolism in marine diatoms for biofuel production. Biotechnology for Biofuels. 2015;8:28-38.

21. Hildebrand D, Smith SR, Traller JC, Abbriano RM. The place of diatoms in the biofuels industry. Biofuels. 2012;3(2):221-40.

22. Armbrust EV, Berges JA, Bowler C, Green BR, Martinez D, Putnam NH, et al. The genome of the diatom *Thalassiosira pseudonana*: Ecology, evolution, and metabolism. Science. 2004;306:79-86.

23. Bowler CB, Allen AE, Badger JH, Grimwood J, Jabbari K, Kuo A, et al. The *Phaeodactylum* genome reveals the evolutionary history of diatom genomes. Nature. 2008;456(7219):239-44.

24. Chepurnov VA, Chaerle P, Roef L, Van Meirhaeghe A, Vanhoutte K. Classical breeding in diatoms: Scientific background and practical perspectives. Diatom Biology. Cellular Origin, Life in Extreme Habitats and Astrobiology. 19: Springer; 2011. p. 167-94.

25. Apt KE, Kroth-Pancic PG, Grossman A. Stable nuclear transformation of the diatom *Phaeodactylum tricornutum*. Mol Gen Genet. 1996;252(5):572-9.

26. Patil S, Moeys S, Von Dassow P, Huysman MJJ, Mapleson D, De Veylder L, et al. Identification of the meitic toolkit in diatoms and exploration of meiosis-specific *SPO11* and *RAD51* homologs in the sexual species *Pseudo-nitzchia multistriata* and *Seminavis robusta*. BMC Genomics. 2015;2015(930).

27. Yoshiyama K, Sharp JH. Phytoplankton response to nutrient enrichment in an urbanized estuary: Apparent inhibition of primary production by overeutrophication. Limnol Oceanogr. 2006;51(1, part 2):424-34.

28. Hoops HJ, Floyd GL. Ultrastructure of the centric diatom, *Cyclotella meneghiniana*: vegetative cell and auxospore development. Phycologia. 1979;18:424-35.

29. Bernstein H, Bernstein C. Evolutionary origin of recombination during meiosis. BioScience. 2010;60(7):498-505.

30. Collos Y, Harrison PJ. Acclimation and toxicity of high ammonium concentrations to unicellular alage. Mar Poll Bull. 2014;80(1-2):8-23.

31. Mouget J-L, Gastineau R, Davidovich O, Gaudin P, Davidovich NA. Light is a key factor in triggering sexual reproduction in the pennate diatom *Haslea ostrearia*. FEMS Microbiol Ecol. 2009;69(2):194-201.

32. Vaulot D, Chisholm SW. A simple model for the growth of phytoplankton populations in light/dark cycles. J Plankton Res. 1987;9:345-66.

33. Mills KE, Kaczmarska I. Autogamic reproductive behavior and sex cell structure in *Thalassiosira angulata* (Bacillariophyta). Botanica Mar. 2006;49:417-30.

34. Gillard J, Devos V, Huysman MJJ, De Veylder L, D'Hondt S, Martens C, et al. Physiological and transcriptomic evidence for a close coupling between chloroplast ontogeny and cell cycle progression in the pennate diatom *Seminavis robusta*. Plant Physiol. 2008;148:1394-411.

35. Kaczmarska I, Poulíčková A, Sato S, Edlund MB, Idei M, Watanabe T, et al. Proposals for a terminology for diatom sexual reproduction, auxospores and resting stages. J Diatom Res. 2013;28(3):263-94.

36. French III FW, Hargraves PE. Spore formation in the life cycles of the diatoms *Chaetoceros diadema* and *Leptocylindrus danicus*. J Phycol. 1985;21:477-83.

37. Idei M, Osada K, Sato S, Toyoda K, Nagumo T, Mann DG. Gametogenesis and auxospore development in *Actinocyclus* (Bacillariophyta). PLOS One. 2012;7(8):e41890.

38. Chepurnov VA, Mann DG, Von Dassow P, Armbrust EV, Sabbe K, Dasseville R, et al. Oogamous reproduction, with two-step auxosporulation, in the centric diatom *Thalassiosira punctigera* (Bacillariophyta). J Phycol. 2006;42:845-58.

39. Edlund MB, Stoermer EF. Sexual reproduction in *Stephanodiscus niagarae* (Bacillariophyta). J Phycol. 1991;27:780-93.

40. Idei M, Sato S, Nagasato C, Motomura T, Toyoda K, Nagumo T, et al. Spermatogenesis and auxospore structure in the multipolar centric diatom Hydrosera. J Phycol. 2015;51(1):144-58.

41. Shirokawa Y, Shimada M. Sex allocation pattern of the diatom *Cyclotella meneghiniana*. Proceedings of the Royal Society B: Biological Sciences. 2013;280(1761):20130503.

42. Nagai S, Hori Y, Manabe T, Imai I. Restoration of cell size by vegetative cell enlargement in *Coscinodiscus wailesii* (Bacillariophyceae). Phycologia. 1995;34(6):533-5.

43. Sabbe K. Apomixis in *Achnanthes* (Bacillariophyceae); development of a model system for diatom reproductive biology. Eur J Phycol. 2004;39:327-41.

44. Gallagher JC. Cell Enlargement in *Skeletonema costatum* (Bacillariophyceae). J Phycol. 1983;19:539-42.

45. Armbrust EV. Identification of a new gene family expressed during the onset of sexual reproduction in the centric diatom *Thalassiosira weissflogii*. Appl Environ Micriobiol. 1999;65:3121-8.

46. Lutzmann M, Grey C, Traver S, Ganier O, Maya-Mendoza A, Ranisavljevic N, et al. MCM8- and MCM9-deficient mice reveal gametogenesis defects and genome instability due to impaired homologous recombination. Mol Cell. 2012;47(4):523-34.

47. Park J, Long DT, Lee KY, Abbas T, Shibata E, Negishi M, et al. The MCM8-MCM9 complex promotes RAD51 recruitment at DNA damage sites to facilitate homologous recombination. Mol Cell Biol. 2013;33(8):1632-44.

48. Baker SM, Plug AW, Prolla TA, Bronner CE, Harris AC, Yao X, et al. Involvement of mouse Mlh1 in DNA mismatch repair and meiotic crossing over. Nat Genet. 1996;13(3):336-42.

49.   Butler G, Rasmussen MD, Lin MF, Santos MAS, Sakthikumar S, Munro CA, et al. Evolution of pathogenicity and sexual reproduction in eight *Candida* genomes. Nature. 2009;459:657-62.

50.   Cooper KF, Mallory JJ, Egeland DB, Jarnik M, Strich R. Ama1p is a meiosis-specific regulator of the anaphase promoting complex/cyclosome in yeast. PNAS. 2000;97(26):14548-53.

51.   Zachariae W, Shevchenko A, Andrews PD, Ciosk R, Galova M, Stark MJR, et al. Mass spectrometric analysis of the anaphase-promoting complex from yeast: Identificaiton of a subunit related to cullins. Science. 1998;279(5354):1216-9.

52.   Mazin AV, Mazina OM, Bugreev DV, Rossi MJ. Rad54, the motor of homologous recombination. DNA Repair. 2010;9(3):286-302.

53.   Honda D, Shono T, Kimura K, Fujita S, Iseki M, Makino Y, et al. Homologs of the sexually induced gene 1 (*sig1*) product constitute the Stramenopile mastigonemes. Protist. 2007;158(1):77-88.

54.   Rosenbaum JL, Witman GB. Intraflagellar transport. Nat Rev Mol Cell Biol. 2002;3:813-25.

55.   Merchant SS, Prochnik SE, Vallon O, Harris EH, Karpowicz SJ, Witman GB, et al. The *Chlamydomonas* genome reveals the evolution of key animal and plant functions. Science. 2007;318(5848):245-50. doi: 10.1126/science.1143609.

56.   Ambawat S, Sharma P, Yadav NR, Yadav RC. MYB transcription factor genes as regulators for plant responses: an overview. Physiol Mol Biol Plants. 2013;19(3):307-21.

57.   Cheng H, Song S, Xiao LX, Soo HM, cheng Z, Xie D, et al. Gibberellin acts through jasmonate to control the expression of *MYB21, MYB24,* and *MYB57* to promote stamen filament growth in *Arabidopsis*. Plos Genet. 2009;5. doi: 10.1371/journal.pgen.1000440.

58.   Archibald JM. The puzzle of plastid evolution. Curr Biol. 2009;19(2):R81-R8.

59.   Geitler L. Der Formwechsel der pennaten Diatomeen. Arch Protistenk. 1932;78(1):1-226.

60.	Mann DG. Why didn't Lund see sex in Asterionella? A discussion of the diatom life cycle in nature. In: Round FE, editor. Algae and the Aquatic Environment, contributions in honour of JWG Lund. Bristol: Biopress; 1988. p. 384-412.

61.	Tillmann U. Interactions between planktonic microalgae and protozoan grazers. J Euk Microbiol. 2004;51:156-68.

62.	Sterner RW, Elser JJ. Ecological stoichiometry: the biology of elements from molecules to the biosphere. Princeton, New Jersey, USA: Princeton University Press; 2002.

63.	Chen G, Rynearson TA. Genetically distinct populations of a diatom co-exist during the North Atlantic spring bloom. Limnol Oceanogr. 2016;61(6):2165-79.

64.	Guillard RRL. Culture of phytoplankton for feeding marine invertebrates. In: Smith WL, Chanley MH, editors. Culture of Marine Invertebrate Animals. New York: Plenum Press; 1975. p. 26-60.

65.	Dyhrman ST, Jenkins BD, Rynearson TA, Saito MA, Mercier ML, Alexander H, et al. The transcriptome and proteome of the diatom *Thalassiosira pseudonana* reveal a diverse phosphorus stress response. PLOS One. 2012;7(3):e33768.

66.	Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nat Prot. 2016;11:1650-67.

67.	Bushnell B. BBtools 2016.

68.	Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. Nat Methods. 2015;12:357-60.

69.	Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotech. 2015;33(290-295):290-5.

70.   Frazee AC, Pertea GM, Jaffe AE, Langmead B, Salzberg SL, Leek JT. Ballgown bridges the gap between transcriptome assembly and expression analysis. Nat Biotech. 2015;33(3):243-6.

71.   Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. BMC Bioinformatics. 2004;19(5):113.

72.   Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. Bioinformatics. 2006;22:2688-90.

73.   Chevenet F, Brun C, Bañuls AL, Jacq B, Christen R. TreeDyn: towards dynamic graphics and annotations for analyses of trees. BMC Bioinformatics. 2006;7:439.

74.   Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, et al. Phylogeny.fr: robust phylogenetic analysis for the non-specialist. Nucleic Acids Res. 2008;1(36):W465-W9.

Table 1. Effects of growth parameters on induction of sexual reproduction in *T. pseudonana* CCMP1335, *T. weissflogii* and *C. cryptica*. Oogonia and auxospores always appeared in stationary phase. The percentage of the total population (at least 300 cells counted per replicate) differentiated into oogonia or auxospores when grown in nitrate or ammonium is shown for the day they were at their maximum number; data are mean values ± s.d., biological replicates n=3.

Species	Light intensity (μE)	Light regime	Nitrogen concentration (μM)	Specific growth rate (d^{-1})	Oogonia and auxospores in NO_3^- (%)	Oogonia and auxospores in NH_4^+ (%)
T. pseudonana	5	24 h	200	0.28	0.21 ± 0.22	0.52 ± 0.52
	50	24 h	200	0.89	3.67 ± 0.55	7.67 ± 0.50
	50	12 h/12 h L/D	200	0.80	0.67 ± 0.27	2.32 ± 0.52
	220	24 h	200	1.2	0.26 ± 0.26	1.30 ± 0.91
	70	24 h	800	0.98	3.01 ± 0.52	38.9 ± 0.04
C. cryptica	100	24 h	800	0.38	11.1 ± 2.27	37.6 ± 3.71
T. weissflogii	100	24 h	800	0.52	4.50 ± 2.36	39.8 ± 5.65

1
2
3
Fig 1. The life cycle of a centric diatom. The average cell size of a population of asexually
dividing diatoms decreases as a result of differential thecae inheritance. At a critically small size,
cells can initiate sexual reproduction and differentiate into male and female cells. Meiosis in the
male spermatogonangium produces multinucleate spermatogonia that divide into individual
haploid spermatocytes. Meiosis in the female oogonia produces a single functional haploid
nucleus that is fertilized by a flagellated spermatocyte through an opening in the oogonia thecae.
Fertilized oogonia expand into a large auxospore where new, large thecae are formed for the new
initial cell.

Fig 2. Ammonium induces sexual morphologies in *T. pseudonana* CCMP1335. (**A**)
Proportion of sexual cells (oogonia and auxospores) relative to the total population in cultures of
T. pseudonana grown in the presence of NH_4Cl or $NaNO_3$; n=3 independent cultures, average of
300 cells counted per replicate. Oogonia and auxospores were only observed beginning in
stationary phase, data are mean values, error bars are s.d.. Inset: corresponding growth curve
linking the onset of stationary phase with first appearance of sexual cells on day six. (**B**) Sexual
cells were observed in cultures with NH_4Cl present at inoculation (blue hatched and solid blue
bars) or following NH_4Cl addition at the onset of stationary phase (yellow bars). Legend shows
concentration of nitrogen source provided at inoculation and concentration of nitrogen source
added at the time of the second dosing. Two control treatments were supplied 200 µM nitrogen
source at inoculation only. Inset: corresponding growth curve showing the onset stationary phase
and timing of 2nd nitrogen addition; n=3 independent cultures, average of 281 cells counted per
replicate, data are mean values, error bars are s.d..

Fig 3. The life cycle stages of *T. pseudonana* (A-K), T. weissflogii (L) and C. cryptica (M-O) imaged using scanning electron microscopy (SEM), light (LM), and confocal microscopy (CFM). **A:** Two vegetative cells (LM, CCMP1335). **B:** Oogonium displaying separation of thecae (arrowhead) and putative pycnotic nucleus indicated by the arrow (CFM, CCMP1015). **C:** Oogonium sharply bending at the thecae junction. Arrowhead indicates protrusion of the plasma membrane (CFM, CCMP1335). Oogonia images are representative of 38 total images. **D:** Spermatogonium containing multiple spermatocytes seen as individual red (DNA stained) clusters (CFM, CCMP1335); representative of 8 images. **E:** Motile spermatocytes (in red, arrow) with moving flagella (arrowheads, CFM, CCMP1335, representative of 10 images). **F-G:** SEM images of spermatocytes (arrowhead) attached to early oogonia (SEM, CCMP1335, representative of 20 images). **H,I:** Auxospores; representative of 60 images in CCMP1015 (H, LM) and CCMP1335 (I, CFM) showing bulging where mother valve was attached (arrowhead). Two nuclei are visible in red following non-cytokinetic mitosis. **J:** Small parental cell (arrow) with initial cells produced by sexual reproduction to the left (partial valve view) and right (girdle view) indicated by the arrowheads (SEM, CCMP1335). **K:** 7 x 12 µm initial cell (LM); j and k representative of 12 images of CCMP1335. **L:** *T. weissflogii* auxospore (LM); representative of 12 similar images. **M:** *C. cryptica* spermatogonium (upper left) and vegetative cell (lower right). CFM shows stained DNA (red, arrow) and multiple nuclei in the spermatogonium. Arrowheads indicate chlorophyll autofluorescence (green). Oogonium (**N**, representative of 6 images) and auxospore (**O**, representative of 4 similar images) of *C. cryptica* (LM). Confocal microscopy images (b-e, i, m) show chlorophyll autofluorescence (green) and Hoescht 33342 stained DNA (red). Scale bars: **A**: 5 µm; **B-E**: 5 µm; **F**: 2 µm; **G**: 1 µm; **H-O**: 10 µm.

53
54
Fig 4. Evidence for meiosis and initial cells. **A:** 18s rRNA phylogeny of diatoms including
penates (pink rectangle), centrics (blue rectangle). Highlighted in red are the four strains induced
into sexual reproduction in this study. Species for which some evidence already exists for sexual
reproduction are starred [9, 13, 16, 28, 33, 38]. **B-D:** Changes in DNA and chlorophyll
fluorescence in exponential (EXP), stationary (STA) and late stationary (L-STA) growth phases
of *T. pseudonana* induced by ammonium; 30,000 events recorded, representative of two
biological replicates. **E:** Coulter Counter distributions of cell diameter for *T. pseudonana*
cultures in exponential phase of growth and maintained in $NaNO_3$ (red) and after six successive
transfers to medium with ammonium (blue), single replicates of cultures with cell densities of 2.4
x 10^6 ml^{-1} ($NaNO_3$) and 2.3 x 10^6 ml^{-1} (ammonium). Dashed lines are the mode for each peak.
65

Fig 5. Transcriptomic evidence for sexual reproduction in *T. pseudonana*. **A:** Heat map of 89 genes having annotated functions that were differentially expressed during differentiation and sexual reproduction in *T. pseudonana* CCMP1335. Color indicates normalized expression value (FPKM) for each nitrogen treatment (control = 100 μM NO_3^-; 100NH4 = 100 μM NH_4^+; 800NH4 = 800 μM NH_4^+) and growth phase (EXP, STA, L-STA). **B:** FPKM values of select genes across growth phases for each nitrogen treatment.

Supporting Information:

S1 Table. Pairwise comparison, Stationary phase 800 µM Ammonium vs. exponential

phase 800 µM Ammonium.

S2 Table. Pairwise comparison, Late-stationary phase 800 µM Ammonium vs. exponential

phase 800 µM Ammonium.

S3 Table. Pairwise comparison, Exponential phase 100 µM Ammonium vs. Late stationary

phase 100 µM Ammonium.

S4 Table. Pairwise comparison, Stationary phase 800 µM Ammonium vs. Late stationary

phase 800 µM Ammonium.

S5 Table. Pairwise comparison, Exponential phase nitrate vs. Exponential phase 800 µM

Ammonium.

S6 Table. Pairwise comparison, Stationary phase nitrate vs. Stationary phase 800 µM

Ammonium.

S7 Table. Pairwise comparison, Late Stationary phase nitrate vs. Late Stationary phase 800

µM Ammonium.

S8 Table. Pairwise comparison, Exponential phase nitrate vs. Exponential phase 100 μM

Ammonium.

S9 Table. Pairwise comparison, Late Stationary phase nitrate vs. Late Stationary phase 100

μM Ammonium.

S10 Table. Pairwise comparison, Exponential phase 100 μM Ammonium vs. Exponential

phase 800 μM Ammonium.

S11 Table. Pairwise comparison, Late Stationary phase 100 μM Ammonium vs. Late

Stationary phase 800 μM Ammonium.

S12 Table. Interaction analysis, Exponential - Stationary phases (800 μM Ammonium v.

Nitrate control).

S13 Table. Interaction analysis, Exponential - Late Stationary phases (800 μM Ammonium

v. Nitrate control).

S14 Table. Interaction analysis, Exponential - Late Stationary phases (100 μM Ammonium

v. Nitrate control).

S15 Table. Interaction analysis, Stationary - Late Stationary phases (800 μM Ammonium v.

Nitrate control).

S16 Table. Genes that were identified as differentially expressed in conditions that

triggered cell differentiation and sexual reproduction and have annotated functions.

S17 Table. Genes that were identified as differentially expressed in conditions that

triggered cell differentiation and sexual reproduction and are homologous to yeast genes

involved in meiosis.

S18 Table. Genes that were identified as differentially expressed in conditions that

triggered cell differentiation and sexual reproduction and are of red algal origin.