1 **Short tandem repeats with massive variation and functional consequences**
2 **across strains of *Arabidopsis thaliana*.**
3

4 Maximilian O. Press[1], Rajiv C. McCoy[1], Ashley N. Hall[2], Joshua Akey[1], and Christine
5 Queitsch[1]*
6
7 1: University of Washington Department of Genome Sciences, Seattle, WA
8 2: University of Washington Department of Molecular and Cellular Biology, Seattle, WA
9 *: to whom correspondence should be addressed: queitsch@uw.edu

11 Keywords: Short tandem repeat, microsatellite, quantitative genetics, selection,
12 *Arabidopsis thaliana*.

Abstract

Short tandem repeat (STR) mutations may be responsible for more than half of the

mutations in eukaryotic coding DNA, yet STR variation is rarely examined as a

contributor to complex traits. We assess the scope of this contribution across a

collection of 96 strains of *Arabidopsis thaliana* by massively parallel STR genotyping.

95% of examined STRs are polymorphic, and the median STR has six alleles. Modest

STR expansions are found in most strains, some of which have evident functional

effects. For instance, three of six intronic STR expansions are associated with intron

retention. We infer selective constraint on STRs, and find the strongest signatures of

purifying selection on coding STRs. Lastly, we detect dozens of novel STR-phenotype

associations that could not be detected with SNPs, and validate some experimentally.

Our results demonstrate that STRs comprise a large unascertained reservoir of

functionally relevant genomic variation.

Introduction

Mutation rates vary within a genome by several orders of magnitude[1], from ~10^{-8}-10^{-9} for

substitutions to 10^{-3}-10^{-4} on average for short tandem repeats (STRs)[2–4]. STR mutations

occur through addition or subtraction of repeat units. Given the prevalence of STR loci

in eukaryotic genomes, we would expect more *de novo* STR mutations than single

nucleotide substitutions in the human genome per generation[4], even in coding regions[5].

Thus, while the overall mutation rate is under strong control from natural selection,

some loci experience many more mutations than others[6]. The existence of such highly

mutable loci contradicts the commonly-used infinite sites model[7], which assumes that

no locus mutates more than once in a population; it further poses complications for

quantitative genetic models that assume contributions from many independent loci [8].

In spite of the large effect that STRs can have on complex traits and diseases in model

organisms and humans[9–11], their variation is largely not considered in genotype-

phenotype association studies due to technical obstacles. However, STR genotyping

methods of sufficient accuracy, throughput, and cost-effectiveness to ascertain STR

alleles at high throughput have recently become available[12–14]. Studies leveraging these

methods suggest considerable contributions of STRs to heritable phenotypic

variation[12,15].

From an evolutionary perspective, the high mutation rate and clear phenotypic effects of

STRs are speculated to provide readily accessible evolutionary paths for rapid

adaptation[16–18]. On larger time scales, the presence of STRs with high mutation rates in

coding regions appears to be maintained by selection[19–21]. These observations

independently argue for important functional roles of STRs.

In the present study, we apply massively parallel STR genotyping to a diverse panel of

well-characterized *A. thaliana* strains. We use these data to generate and test

hypotheses about the functional effects of STR variation, combining observations of

gene disruption by STR expansion, inferences about STR conservation, and phenotypic

association analyses with follow-up experiments. Based on our results, we argue that

STRs must be included in any comprehensive account of phenotypically relevant

genomic variation.

Results

STR genotyping reveals complex allele frequency spectra.

We targeted 2,050 STR loci for genotyping with molecular inversion probes (MIPs)[12]

across a core collection of 96 *A. thaliana* strains (Methods). These loci were all less

than 200bp in size and had nucleotide purity of at least 90%, encompassing nearly all

gene-associated STRs and ~40% of intergenic STRs (Fig. 1a). We used comparisons

with the Col-0 reference genome, PCR analysis of selected STRs, and dideoxy

sequencing to estimate that MIP STR genotype calls were ~95% accurate, and

inaccurate calls were generally only one to two units away from the correct copy

number (Supplementary Note, Supplementary Fig. 1-4, Supplementary Table 1).

1
Figure 1. STRs in *A. thaliana* show a complex allele frequency distribution and geographic
differentiation. (a): Distribution and ascertainment of STR loci. "All": all STRs matching the
definition of STRs for this study, *e.g.* <=180 bp length in TAIR10, >=89% purity in TAIR10, 2-10
bp nucleotide motif. "Targeted": the 2050 STRs targeted for MIP capture. "Typed": STRs
successfully genotyped in the Col-0 genome in a MIPSTR assay. Numbers above bars indicate
the proportion of targeted STRs in the relevant category that were successfully genotyped. (b):
The distribution of allele counts across all genotyped STRs. (c): The distribution of major allele
frequencies (frequency of the most frequent allele at each locus) across genotyped STRs. (d):
Principal component analysis (PCA) reveals substantial geographic structure according to STR
variation. PC1 and PC2 correspond, respectively, to 5.2% and 4.0% of total STR allele variance.
12

95% of STRs were polymorphic. Most STRs were highly multiallelic (Fig. 1b; mean=6.4

alleles, median=6 alleles), and this variation was mostly unascertained by the 1001

Genomes resource for *A. thaliana* (Supplementary Fig. 3a). Coding STRs were only

slightly less polymorphic (mean=4.5 alleles, median=4 alleles), though whether this

difference is due to selection or to mutation rate variation is unclear. 45% of STRs had a

major allele with frequency < 0.5, confusing the familiar concepts of major and minor

alleles, which have provided a common framework for detecting genotype associations

(Fig. 1c). Specifically, the Col-0 reference strain carries the major STR allele at only

48% of STR loci. Moreover, rarefaction analysis implied that more STR alleles at these

loci are expected with further sampling of *A. thaliana* strains (Supplementary Fig. 4c).

Principal component analysis of STR variation revealed genetic structure corresponding

to Eurasian geography (Fig. 1d, Supplementary Fig. 5), consistent with previous

observations that population structure is associated with geography in *A. thaliana*[22,23].

This result, which agrees with previous observations from a much larger set of genome-

wide single nucleotide polymorphism markers, shows that a comparatively small panel

of STRs suffices to capture detailed population structure in *A. thaliana*. Overall, we find

that STRs are extremely allele-rich, and their variation reflects expected population

structure.

Novel STR expansions are associated with splice disruptions.

We next examined the frequency and functional consequences of STR expansions in *A.*

thaliana. STR expansions, extreme high-copy-number variants of comparatively short

STRs, are widely recognized as contributing to human disease[24] and other

phenotypes[25]. While large (>150 bp) expansions are difficult to infer, we detected

modest STR expansions with a simple heuristic comparing the longest allele observed

at each locus to the median allele (Fig. 2a). We found expansions in 64 of 96 *A. thaliana*

strains, each carrying at least one expanded STR allele from one of 28 expansion-prone

STRs (9 coding, 6 intronic, 8 UTR, 5 intergenic). Most expansions were found in

multiple strains (Fig. 2b), although expansion frequency was likely underestimated due

to a higher rate of missing data at these loci. STR expansions causing human disease

can be as small as 25 copies[24], whereas we detected expansions up to ~50 copies.

1
Figure 2. Inferring and assessing the functional effects of modest STR expansions. (a): The
distribution of expansion scores across STRs, where the expansion score is computed as
(max(STR length) – median(STR length)) / median STR length. We called any STRs with a score
greater than 2 a modest expansion (indicated). (b): Distribution of allele frequencies of the 28
expanded STR alleles. (c, d): Distribution of STR copy number of the intronic STR in the *NTM1*
gene and the 3' UTR STR in the *MEE36* gene. (e): RT-PCR demonstrates intron retention in
NTM1 mRNA in the Mr-0 strain, which carries the STR expansion, yielding an aberrant 437-bp
product. (f): *MEE36* transcript abundances measured by qRT-PCR and normalized relative to
UBC21 transcript levels. For each strain, two independent biological replicates are shown as
points. Transcript levels are expressed relative to Col-0 levels (set to 1). *: STR genotype
corrected by dideoxy sequencing. Strains and order are the same between (e) and (f).

We assayed the effects of STR expansions on expression of associated genes. The

most dramatic expansions (with large relative copy number increase) affected an

intronic STR in the *NTM1* gene (Fig. 2c; five other expansions also resided in introns)

and a STR in the 3' UTR of the *MEE36* gene (Fig. 2d). These genes, respectively, have

roles in cell proliferation and embryonic development. Intronic STR mutations can

disrupt splicing, causing altered gene function[26] and human disease[27]. Therefore, we

assayed effects on splicing of all six expanded intronic STRs. In three cases, the

expanded allele was associated with partial or full retention of its intron (Supplementary

Fig. 6, Supplementary Note), including the major *NTM1* splice form in the Mr-0 strain

(Fig. 2e). We confirmed intron retentions by dideoxy sequencing of cDNA

(Supplementary Fig. 7a). Retention of the *NTM1* intron is predicted to lead to a

nonsense mutation truncating most of the NTM1 protein (Supplementary Fig. 7b). For

the other two retentions, more complex and STR allele-specific mRNA species were

formed (Supplementary Fig. 6, Supplemental Text). The *MEE36* STR expansion alleles

were associated with dramatically reduced *MEE36* transcript levels (Fig. 2f), possibly

due to the STR expansion altering transcript processing[28]. These examples emphasize

the potential for previously unascertained STR variation to modify gene function.

Moreover, considering STR allele frequency distributions, as we did here by focusing on

outliers, enables predictions about STR functional effects.

Signatures of functional constraint on STR variation.

Using our observed STR allele frequency distributions, we next attempted to infer

selective processes acting on STRs. Previous models for evaluating functional

constraint on STRs are few[29], though there is consensus that selection affects STR

variation[17,19,29,30]. Naively, we would expect that coding STRs should show increased

constraint (lower variation). Consistent with this expectation, we observed that most

invariant STRs are coding (53 of 84 invariant STRs genotyped across at least 70

strains; odds ratio = 4.5, p = $5*10^{-11}$, Fisher's Exact Test). However, methods of inferring

selection by allele counting are confounded by population structure and by mutation

rate, which varies widely across STRs in this (Fig. 3a) and other studies[3,31].

Therefore, to account for mutation rate and population structure, we used support vector

regression (SVR) to predict STR variability across these 96 strains, using well-

established correlates of STR variability (*e.g.* STR unit number and STR purity; Online

Methods)[3,32,33]. Selection was defined as deviation from expected variation of a neutral

STR among strains. We trained SVRs on the set of intergenic STRs, which should

experience minimal selection relative to STRs associated with genes (Supplementary

Fig. 8-10). We used bootstrap aggregation of SVR models to compute a putative

constraint score for each STR by comparing its observed variability to the expected

distribution from bootstrapped SVR models (Fig. 3b, Supplementary Note,

Supplementary File 1).

Figure 3. Detecting functionally constrained STRs. (a): The distribution of $\hat{\theta}$ (estimated

mutation rate[7]) across all genotyped STR loci. (b): Distribution of "selection scores" across all
STRs, separated by locus category. Vertical lines indicate 2.5% and 97.5% quantiles of the
distribution of intergenic STRs, which are used as thresholds for putative constraint and
hypervariability respectively. (c): Constrained or hypervariable STRs separated by locus
category. White boxes indicate the expected numbers for each bar, based on number of STRs in
each locus category and number of STRs under different types of selection. (d): *BIN4* intron STR
is constrained relative to similar STRs. Allele frequency spectra are normalized by subtracting
the median copy number (9 for the *BIN4* STR). All pure STRs with TA/AT motifs and a median
copy number between 7 and 12 are included in the "similar STRs" distribution. (e): Lack of
association between near-expansion *CMT2* STR alleles and previously described nonsense
mutations. (f): Neighbor-joining tree of a 10KB region of *A. thaliana* chromosome 4
encompassing the *CMT2* gene across 81 strains with available data. Tip labels in red carry an
adaptive nonsense mutation early in the first exon of *CMT2*, as noted previously[41]. Red bars
drawn on tips of the tree indicate the length of the *CMT2* intronic STR (as a proportion of its
maximum length, 36.5 units). The bars are omitted for tips with missing STR data. (g):
Constraint score from (b) plotted with respect to nearest TSS. (h): Constraint score from (b)
plotted with respect STR annotations.

According to constraint scores, 132 STRs were less variable than expected, suggesting

purifying selection on these loci (Fig. 3c). Among these, 63 STRs were coding (OR =

2.4, p = 3.7*10^{-6}, Fisher's Exact Test); this enrichment of coding STRs among

constrained STRs agrees with our naïve analysis of invariant STRs above. Examples of

constrained coding STRs included STRs encoding homologous polylysines in three

different histone H2B proteins. Generally, coding STRs showing purifying selection

encoded roughly half as many polyserines and twice as many acidic homopolymers as

expected from the *A. thaliana* proteome (Supplementary Table 3)[34]. Although many

more coding STRs are probably functionally constrained, our power to detect such

constraints is limited by the size of the dataset. We also observed high conservation of

some STRs in non-coding regions, though this is less interpretable given the ambiguous

relationship between sequence conservation and regulatory function in *A. thaliana*[35].

The most constrained intronic STR, in the *BIN4* gene, which is required for

endoreduplication and normal development[36], shows a restricted allele frequency

spectrum compared to similar STRs (Fig. 3d).

Hypervariable coding STRs (showing more alleles than expected) were too few for

statistical arguments, but nonetheless showed several notable patterns (Supplementary

Note, Supplementary Fig. 11). For example, 3/24 hypervariable coding STRs encoded

polyserines in F-box proteins, suggesting that STR variation may serve as a mechanism

of diversification in this protein family, which shows dramatically increased family size

and sequence divergence in some plant lineages[37,38]. One non-coding hypervariable

STR was associated with the *Chromomethylase 2* (*CMT2*) gene, which is under

selection in *A. thaliana*[39]. Specifically, CMT2 nonsense mutations in some populations

are associated with temperature seasonality. We considered whether the extreme

CMT2 STR alleles might be associated with these nonsense mutations. Instead, these

extreme alleles exclusively occurred in strains with full-length CMT2 (Fig. 3e). Strains

with the common *CMT2* nonsense mutation form a tight clade in the *CMT2* sequence

tree, whereas the *CMT2* STR length fluctuates rapidly throughout the tree and appears

to converge on longer alleles independently in different clades (Fig. 3f). These

convergent changes are consistent with a model in which the *CMT2* STR is a target of

selection.

We further assessed whether STR conservation can be attributed to *cis*-regulatory

function. The abundance of STRs in eukaryotic promoters[40], and their associations with

gene expression (in *cis*)[15,41], have suggested that STRs affect transcription, possibly by

altering nucleosome positioning[41]. We examined whether STRs near transcription start

sites (TSSs) showed signatures of functional constraint (Fig. 3g). We found little

evidence for reduced STR variation near TSSs, suggesting that *cis*-regulatory effects do

not generally constrain STR variation. Moreover, we found no relationship between

putative constraint and whether STRs reside in accessible chromatin sites marking

regulatory DNA (Fig. 3h).

STRs yield numerous novel genotype-phenotype associations.

We next addressed whether STR genotypes contribute new information for explaining

phenotypic variation. It is often assumed that STR phenotype associations should be

captured through linkage to common single nucleotide polymorphisms (SNPs)[42]. This

assumption is inconsistent with available data. Human STR data[42,43] and simulation

studies[44] indicate decreased linkage disequilibrium (LD) between STRs and SNPs,

making it uncertain whether SNP markers can meaningfully tag STR genotypes. Indeed,

we found little evidence of LD between STR and SNPs in *A. thaliana* (Fig. 4a),

Moreover, the observed LD around STR loci declined with increasing STR allele

number (Supplementary Fig. 12), consistent with an expected higher mutation rate at

multiallelic loci[7,44]. This result suggests that STR-phenotype associations need to be

directly tested rather than relying on linkage to SNPs. *A. thaliana* offers extensive high-

quality phenotype data for our inbred strains, which have been previously used for SNP-

based association studies[45]. We tested each polymorphic STR for associations across

the 96 strains with each of 105 published phenotypes. For the subset of 32 strains for

which RNA-seq data were available[46], we also tested for associations between STR

genotypes and expression of genes within 1MB (*i.e.*, eQTLs; Supplementary Note).

Although our power to detect eQTLs was limited by sample size, we detected 12

associations. The strongest association was between a STR residing in long noncoding

RNA gene *AT4G07030* and expression of the nearby stress-responsive gene *AtCPL1*

(Supplementary Fig 13, Supplementary Table 4).

We next focused on organismal phenotypes. Certain STRs showed associations with

multiple phenotypes, and flowering time phenotypes were particularly correlated with

one another (Fig. 4c,d). Similar to these patterns, SNPs have also shown associations

with multiple phenotypes, and correlated flowering time phenotypes are among the

strongest associations in the same strains[45]. As in previous association studies using

STRs[15], some inflation was apparent in test p-values compared to expectations, though

the same tests using permuted STR genotypes showed negligible inflation (Fig. 4b,

Supplementary Fig 14). Negligible inflation with permuted genotypes has been used

previously to exclude confounding from population structure[15], which we will also

presume here (Supplementary Note). We found 137 associations between 64 STRs and

25 phenotypes at stringent genome-wide significance levels (Methods; Supplementary

Table 5). Given the low LD observed between STRs and other variants, STRs are likely

to be the causal variants, rather than merely tagging them. Our analysis found plausible

candidate genes, such as *COL9*, which acts in flowering time pathways and contains a

flowering time-associated STR, and *RABA4B*, which acts in the salicylic acid defense

response[47] and contains a STR associated with lesion formation.

Figure 4. Diverse associations of STRs with quantitative phenotypes. (a): Multiallelic LD[64]

estimates for STR and SNP loci. Lowess lines for each category are plotted. All values of $r^2 < 0.05$

are omitted from lowess calculation for visualization purposes. (b): Quantile-quantile plot of p-

values from tests of association between STRs and germination rate after 28 days of storage.

(c): An example association between an STR (33085) and a phenotype (flowering time in long

days after 4 weeks vernalization) in *A. thaliana* strains. Median of each distribution is indicated

by a bar proportional in width to the number of observations. (d): Heatmap showing pairwise

associations between STRs and phenotypes, summarized by the p-value from a linear mixed

model, fitting STR allele as a fixed effect and kinship as a random effect. Both rows and columns

are clustered, though the row dendrogram was omitted for clarity. STRs with genotype

1 information in fewer than 25 strains are not displayed. Flowering time phenotypes are boxed in
2 black.
3

We evaluated whether these associations might have been found using SNP-based
analyses, and whether STR effects are large enough to be meaningful. We found that
STR effects on phenotype are largely not accounted for by nearby SNP variation.
Considering the strongest association for each STR, only 18 of the 64 STRs were near
potentially confounding SNP variants, and most associations (14/18) were robust to
adjustment for nearby SNP genotypes[45] (Supplementary Table 6, Supplementary Note).
One notable exception was an STR closely linked to a well-known deletion of the *RPS5*
gene in a hypervariable region of chromosome 1[48] that causes resistance to bacterial
infection[49]. *RPS5* status is under balancing selection in *A. thaliana*[48]. In this case, the
association and the linkage are apparently strong enough (the STR is ~4KB upstream of
the deletion) that this STR tags *RPS5*'s effect on infection. The observation of a STR
tagging a hypervariable region leads us to speculate that STR variation holds
information about genomic regions with complex mutational histories.

To assess the STR contribution to the variance of a specific trait, we performed a naïve
variance decomposition of the long-day flowering phenotype into SNP and STR
components, as represented by the loci showing associations with this trait. Our results
suggested that STRs potentially contribute as much or more variance than SNPs to this
phenotype (Supplementary Note, Supplementary Table 7). Estimated effect sizes for
STR variants on this phenotype were similar to those of large-effect SNP variants[45]
(Supplementary Fig 15).

Finally, we used mutant analysis to evaluate the two strongest flowering time
associations, a coding STR in *AGL65* and an intronic STR in the uncharacterized gene
AT4G01390; neither locus had been associated with flowering time phenotypes. We
found that disruptions of both STR-associated genes had modest early flowering
phenotypes (by ~2 days and ~1 rosette leaf, $p < 0.05$ for each in linear mixed models;
Supplementary Fig 16), supporting the robustness of our STR-phenotype associations.

Taken together, our study suggests that STRs contribute substantially to phenotypic

variation.

Discussion

Our results imply that STRs contribute substantially to trait heritability in *A. thaliana.* Far

from being "junk DNA", STRs are apparently constrained by functional requirements,

STR variation can disrupt gene function, and is associated with phenotypic variation.

Considering that STR variation is represented only poorly through linkage to nearby

SNPs, STRs likely contribute to the missing heritability of complex traits.

STRs are particularly relevant to the phenotypic variance due to *de novo* mutations

(V_m). Estimates of V_m from model organisms are on the order of 1%, but may vary

substantially from trait to trait[50]. STRs are good candidates for a substantial proportion

of this variance, given their high mutation rate, residence in functional regions, and their

functional constraint observed here. In previous work[51], we showed apparent copy

number conservation of a STR in spite of a high mutation rate. In this case, deviation

from the conserved copy number produced aberrant phenotypes. Our observation that

constrained STRs are common suggests that STRs are a likely source of deleterious *de*

novo mutations which are removed by selection.

The extent to which STRs affect phenotype is only partially captured in this study.

Specifically, we assayed two STRs shown to cause phenotypic variation in prior

transgenic *A. thaliana* studies[25,52], but these STR loci did not show strong signatures of

phenotypic association or of selection. This lack of ascertainment suggests that many

more functionally important STRs exist in *A. thaliana* than we can detect with the

analyses presented here. For example, the polyQ-encoding STR in the *ELF3* gene

causes dramatic variation in developmental phenotypes[52], yet we find no statistical

associations between this locus and phenotype across our 96 strains. In this case the

lack of phenotype association is expected; *ELF3* STR alleles interact epistatically with

several loci[53] and associations are thus difficult to detect. Indeed, we have argued that

STRs are more likely than less mutable classes of genomic variation to exhibit

epistasis[9]. In consequence, we expect that the associations described in the present

study are an underestimate of STR effects on phenotype. Moreover, our data are

constrained by MIP technology, which limits the size and composition of STR alleles

that we can ascertain (Figure 1a, Supplementary Figures 1-2).

Considering next a mechanistic perspective, the association we observe between

intronic STR expansions and splice disruptions may be an important mechanism by

which STRs contribute to phenotypic variation. In humans, unascertained diversity of

splice forms contributes substantially to disease[54], and this diversity is larger than

commonly appreciated[55]. We demonstrate that this mechanism is common at least for

expansions; future work should evaluate how tolerant introns are to different

magnitudes of STR variation, as these effects on protein function may prove to be both

large and of relatively high frequency. The phenotypic contributions of loci with high

mutation rate remain underappreciated, specifically in cases where such loci are difficult

to ascertain with high-throughput sequencing. The results presented here argue that

STRs are likely to play a substantial role in phenotypic variation and heritability.

Accounting for the heterogeneity of different classes of genomic variation, and

specifically variation in mutation rate, will advance our understanding of the genotype-

phenotype map and the trajectory of molecular evolution.

METHODS

ONLINE METHODS

Probe design

We used TRF[56] (parameters: matching weight 2, mismatching penalty 5, indel penalty 5,

match probability 0.8, indel probability 0.1, score $\geq$ 40 and maximum period 10) to

identify STRs in the TAIR8 build of the *Arabidopsis thaliana* genome, identifying 7826

putative STR loci under 200bp (Supplementary File 2). We restricted further analysis to

the 2409 loci with repeat purity >=89%. We chose 2307 STRs from among these,

prioritizing STRs in coding regions, introns, or untranslated regions (UTRs), higher STR

unit purity, and expected variability (VARscore[32]). We designed[57] molecular inversion

probes (MIPs) targeting these STR loci in 180bp capture regions with 8bp degenerate

tags in the common MIP backbone. For this purpose, we converted STR coordinates

to the TAIR10 build and used the TAIR10 build as a reference genome. We used single

nucleotide variants (SNVs) in 10 diverse *Arabidopsis thaliana* strains[58] to avoid

polymorphic sites in designing MIP targeting arms. We filtered out MIPs predicted to

behave poorly (MIPGEN logistic score < 0.7), discarded MIPs targeting duplicate

regions, and substituted MIPs designed around SNVs as appropriate. We attempted to

re-design filtered MIPs with 200bp capture regions using otherwise identical criteria.

This yielded a final set of 2050 STR-targeting MIP probes (Supplementary File 3).

MIP and library preparation

These 2050 probes were ordered from Integrated DNA Technologies as desalted DNAs

at the 0.2 picomole scale and resuspended in Tris-EDTA pH 8.0 (TE) to a concentration

of 2 µM and stored at 4° C. We pooled and diluted probes to a final stock

concentration of 1 nM. We phosphorylated probes as described previously[59]. We

performed DNA preparation from whole aerial tissue of adult *A. thaliana* plants. We

prepared MIP libraries essentially as described previously[12,59] using 100 ng *A. thaliana*

genomic DNA for each of 96 *A. thaliana* strains.

Sequencing

We sequenced pooled capture libraries essentially as previously described[12] on

NextSeq and MiSeq instruments collecting a 250bp forward read sequencing the

ligation arm and captured target sequence, an 8bp index read for library

demultiplexing, and a 50bp reverse read sequencing the extension arm and

degenerate tag for single-molecule deconvolution. In each run, 10% of the sequenced

library pool consisted of high-complexity whole-genome library to increase sequence

complexity. For statistics and further details of data acquired for each library see

Supplementary Tables 8 and 9.

STR annotation

We annotated STRs according to Araport11[60], classifying all STRs as coding, intronic,

intergenic, or UTR-localized, and indicating whether each STR overlapped with

transposable element sequence. To identify regulatory DNA, we used the union of

seven distinct DNaseI-seq experiments[61] covering pooled or isolated tissue types. For

additional details, see the Supplementary Note.

Sequence analysis

Sequences were demultiplexed and output into FASTQ format using BCL2FASTQ

v2.17 (Illumina, San Diego). We performed genotype calling essentially as described

previously[12], with certain modifications (Supplementary Note, Supplementary Table 1).

Note that our *A. thaliana* strains are inbred, and more stringent filters and data

processing would be necessary to account for heterozygosity. For information about

comparison with the Bur-0 genome, see the Supplementary Note. Updated scripts

implementing the MIPSTR analysis pipeline used in this study are available at

https://osf.io/mv2at/?view_only=d51e180ac6324d2c92028b2bad1aef67.

Statistical analysis and data processing

We performed all statistical analysis and data exploration using R v3.2.1[62]. For plant

experiments, we fit mixed-effects models using Gaussian (flowering) or binomial

generalized linear models using experiment and position as random effects and

genotype as a fixed effect.

STR expansion inference

We inferred STR expansions where the maximum copy number of an STR is at least

three times larger than the median copy number of that STR. Various alleles of STR

expansions were inspected manually in BAM files. Selected cases were dideoxy-

sequenced and analyzed as described.

Plant material and growth conditions

Plants were grown on Sunshine soil #4 under long days (16h light : 8h dark) at 22°C

under cool-white fluorescent light. T-DNA insertion mutants were obtained from

ABRC[63] (Supplementary Table 11). For flowering time experiments, plants were grown

in 36-pot or 72-pot flats; days to flowering (DTF) and rosette leaf number at flowering

(RLN) were recorded when inflorescences were 1cm high. Results are combined

across at least three experiments.

Gene expression and splicing analysis

We grew bulk seedlings of indicated strains on soil for 10 days, harvested at Zeitgeber

time 12 (ZT12), froze samples immediately in liquid nitrogen, and stored samples at -

80°C until further processing. We extracted RNA from plant tissue using the SV RNA

Isolation kit (including DNase step; Promega, Madison, WI), and subsequently treated it

with a second DNAse treatment using the Turbo DNA-free kit (Ambion, Carlsbad, CA).

We performed cDNA synthesis on ~500ng RNA for each sample with oligo-dT adaptors

using the RevertAid kit (ThermoFisher, Carlsbad, CA). We performed PCR analysis of

cDNA with indicated primers (Supplementary Table 10) and ~25 ng cDNA with the

following protocol: denaturation at 95° 5 minutes, then 30 cycles of 95° 30 seconds,

55° 30 seconds, 72° 90 seconds, ending with a final extension step for 5 minutes at

72°. We gel-purified and sequenced electrophoretically distinguishable splice variants

associated with STR expansions. Each RT-PCR experiment was performed at least

twice with different biological replicates.

Population genetic analyses

For PCA, STRs with missing data across the 96 strains were omitted, leaving 987 STRs

with allele calls for every strain. $\hat{\theta}$ was estimated using the approximation $\hat{\theta} = \frac{1}{8\bar{X}^2} -$

0.5,[7] where $\bar{X}$ is the average frequency of all STR alleles at a locus. We computed

multiallelic linkage disequilibrium estimates for SNP-SNP and SNP-STR locus pairs

using MCLD[64]. We downloaded array SNP data for the same lines (TAIR9 coordinates)

from http://bergelson.uchicago.edu/wp-

content/uploads/2015/04/call_method_75.tar.gz[65]. For each locus, both SNP and STR,

we computed linkage disequilibrium scores with 150 surrounding loci. To facilitate

comparison, we computed lowess estimates of linkage only for those locus pairs in the

plotted distance window in each case, and only for locus pairs with $r^2 > 0.05$.

Inference of conservation

STRs typed across 70 *A. thaliana* strains or fewer were dropped from this analysis, as

the estimates of their variability were unlikely to be accurate, leaving 1825 STRs. We

measured STR variation as the base-10 logarithm of the standard deviation of STR

copy number (Supplementary Note). We used bootstrap aggregation ("bagging") to

describe a distribution of predictions as follows. An ensemble of 1000 support vector

regression (SVR, fit using the *ksvm()* function in the kernlab package[66]) models was

used to predict expected neutral variation of each STR as quantified by each measure

(Supplementary Note). We used this distribution of bootstrapped predictions for

intergenic STRs to compute putative conservation scores (Z-scores) for each STR.

Scores below the 2.5% (Z < -3.46) and above the 97.5% (Z > 3.65) quantiles of

intergenic STRs were considered to be putatively constrained and hypervariable

respectively.

eQTL inference

We downloaded normalized transcriptome data for *A. thaliana* strains from NCBI GEO

GSE80744[46]. We used the Matrix eQTL package[67] to detect associations, fitting also 10

principal components from SNP genotypes to correct for population structure.

Following precedent[15], we fitted additive models assuming that STR effects on

expression would be a function of STR copy number. We used Kruskal-Wallis rank-

sum tests to test the null hypothesis of no association following correction.

Genotype-phenotype associations

We downloaded phenotype data from https://github.com/Gregor-Mendel-

Institute/atpolydb/blob/master/miscellaneous_data/phenotype_published_raw.tsv. We

followed precedent[45] in log-transforming certain phenotypes. In all analyses we treated

STRs as factorial variables (to avoid linearity assumptions) in a linear mixed-effect

model analysis to fit STR allele effects on phenotype as fixed effects while modeling

the identity-by-state kinship matrix between strains (computed from SNP data) as a

correlation structure for strain random effects on phenotype. We performed this

modeling using the *lmekin()* function from the *coxme* R package[68]. We repeated every

analysis using permuted STR genotypes as a negative control to evaluate p-value

inflation, and discarded traits showing such inflation. We used $p < 10^{-6}$ as a genome-

wide significance threshold commensurate to the size of the *A. thaliana* genome and

the data at hand. For flowering time phenotypes we used a more stringent $p < 10^{-10}$

threshold, as these phenotypes showed somewhat shifted p-value distributions (which

were nonetheless inconsistent with inflation, according to negative controls). We

identified potentially confounding SNP associations using the

https://gwas.gmi.oeaw.ac.at/#/study/1/phenotypes resource, using the criterion that a

SNP association must have a $p <= 10^{-4}$ and be within roughly 100KB of the STR to be

considered. We fit models including SNPs as fixed effects as before, and performed

model selection using AICc[69]. Additional details about association analyses are in the

Supplementary Note.

Data and code availability

MIP sequencing data are available in FASTQ format at the Sequence Read Archive,

under project number PRJNA388228. Analysis scripts are provided at

https://osf.io/5jm2c/?view_only=324129c85b3448a8bd6086263345c7b0, along with

data sufficient to reproduce analyses.

ACKNOWLEDGMENTS

We thank Keisha Carlson, Alberto Rivera, and members of the Queitsch lab for

technical assistance and important conversations. We thank Evan Boyle, Choli Lee,

Matthew Snyder, and Jay Shendure for assistance and advice concerning MIP design

and use. We thank the Dunham Lab and the Fields Lab for access to and help with

sequencing instruments. We thank UW Genome Sciences Information Technology for

References

1. Acuna-Hidalgo, R., Veltman, J. A. & Hoischen, A. New insights into the generation and role of de novo mutations in health and disease. *Genome Biol.* **17,** 241 (2016).

2. Sun, J. X. *et al.* A direct characterization of human mutation based on microsatellites. *Nat. Genet.* **44,** 1161–5 (2012).

3. Gymrek, M., Willems, T., Erlich, Y. & Reich, D. E. A framework to interpret short tandem repeat variation in humans. *bioRxiv* 92734 (2016). doi:10.1101/092734

4. Willems, T. *et al.* Population-Scale Sequencing Data Enable Precise Estimates of Y-STR Mutation Rates. *Am. J. Hum. Genet.* **98,** 919–933 (2016).

5. Payseur, B. A., Jing, P. & Haasl, R. J. A Genomic Portrait of Human Microsatellite Variation. *Mol. Biol. Evol.* **28,** 303–312 (2011).

6. Harpak, A., Bhaskar, A. & Pritchard, J. K. Mutation Rate Variation is a Primary Determinant of the Distribution of Allele Frequencies in Humans. *PLOS Genet.* **12,** e1006489 (2016).

7. Haasl, R. J. & Payseur, B. A. The Number of Alleles at a Microsatellite Defines the Allele Frequency Spectrum and Facilitates Fast Accurate Estimation of θ. *Mol. Biol. Evol.* **27,** 2702–2715 (2010).

8. Yang, J. *et al.* Common SNPs explain a large proportion of the heritability for human height. *Nat. Genet.* **42,** 565–9 (2010).

9. Press, M. O., Carlson, K. D. & Queitsch, C. The overdue promise of short tandem repeat variation for heritability. *Trends Genet.* **30,** 504–512 (2014).

10. Fondon, J. W., Hammock, E. A. D., Hannan, A. J. & King, D. G. Simple sequence repeats: genetic modulators of brain function and behavior. *Trends Neurosci.* **31,** 328–34 (2008).

11.   Hannan, A. J. Tandem repeat polymorphisms: modulators of disease susceptibility and candidates for 'missing heritability'. *Trends Genet.* **26,** 59–65 (2010).

12.   Carlson, K. D. *et al.* MIPSTR: A method for multiplex genotyping of germline and somatic STR variation across many individuals. *Genome Res.* **125,** 750–761 (2015).

13.   Willems, T., Zielinski, D., Gordon, A., Gymrek, M. & Erlich, Y. Genome-wide profiling of heritable and de novo STR variations. *bioRxiv* 77727 (2016). doi:10.1101/077727

14.   Highnam, G. *et al.* Accurate human microsatellite genotypes from high-throughput resequencing data using informed error profiles. *Nucleic Acids Res.* **41,** e32 (2013).

15.   Gymrek, M. *et al.* Abundant contribution of short tandem repeats to gene expression variation in humans. *Nat. Genet.* **48,** 22–29 (2015).

16.   Moxon, E. R., Rainey, P. B., Nowak, M. A. & Lenski, R. E. Adaptive evolution of highly mutable loci in pathogenic bacteria. *Curr. Biol.* **4,** 24–33 (1994).

17.   King, D. G. Indirect selection of implicit mutation protocols. *Ann. N. Y. Acad. Sci.* **1267,** 45–52 (2012).

18.   Gemayel, R., Vinces, M. D., Legendre, M. & Verstrepen, K. J. Variable tandem repeats accelerate evolution of coding and regulatory sequences. *Annu. Rev. Genet.* **44,** 445–77 (2010).

19.   Mularoni, L., Ledda, A., Toll-Riera, M. & Albà, M. M. Natural selection drives the accumulation of amino acid tandem repeats in human proteins. *Genome Res.* **20,** 745–54 (2010).

20.   Yu, F. *et al.* Positive selection of a pre-expansion CAG repeat of the human SCA2 gene. *PLoS Genet.* **1,** e41 (2005).

21. Sawaya, S. M., Lennon, D., Buschiazzo, E., Gemmell, N. & Minin, V. N. Measuring microsatellite conservation in mammalian evolution with a phylogenetic birth-death model. *Genome Biol. Evol.* **4,** 636–47 (2012).

22. Nordborg, M. *et al.* The pattern of polymorphism in Arabidopsis thaliana. *PLoS Biol.* **3,** e196 (2005).

23. Alonso-Blanco, C. *et al.* 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. *Cell* **0,** (2016).

24. Usdin, K. The biological effects of simple tandem repeats: Lessons from the repeat expansion diseases. *Genome Res.* **18,** 1011–1019 (2008).

25. Sureshkumar, S. *et al.* A genetic defect caused by a triplet repeat expansion in Arabidopsis thaliana. *Science* **323,** 1060–3 (2009).

26. Li, Y.-C., Korol, A. B., Fahima, T. & Nevo, E. Microsatellites Within Genes: Structure, Function, and Evolution. *Mol. Biol. Evol.* **21,** 991–1007 (2004).

27. Ranum, L. P. W. & Day, J. W. Dominantly inherited, non-coding microsatellite expansion disorders. *Curr. Opin. Genet. Dev.* **12,** 266–271 (2002).

28. Jackson, R. J. Cytoplasmic regulation of mRNA function: The importance of the 3′ untranslated region. *Cell* **74,** 9–14 (1993).

29. Haasl, R. J. & Payseur, B. A. Microsatellites as targets of natural selection. *Mol. Biol. Evol.* **30,** 285–98 (2013).

30. Huntley, M. A. & Clark, A. G. Evolutionary Analysis of Amino Acid Repeats across the Genomes of 12 Drosophila Species. *Mol. Biol. Evol.* **24,** 2598–2609 (2007).

31.   Schlötterer, C., Kauer, M. & Dieringer, D. Allele Excess at Neutrally Evolving Microsatellites and the Implications for Tests of Neutrality. *Proc. Biol. Sci.* **271,** 869–874 (2004).

32.   Legendre, M., Pochet, N., Pak, T. & Verstrepen, K. J. Sequence-based estimation of minisatellite and microsatellite repeat variability. *Genome Res.* **17,** 1787–96 (2007).

33.   Eckert, K. A. & Hile, S. E. Every microsatellite is different: Intrinsic DNA features dictate mutagenesis of common microsatellites present in the human genome. *Mol. Carcinog.* **48,** 379–88 (2009).

34.   Karlin, S., Brocchieri, L., Bergman, A., Mrazek, J. & Gentles, A. J. Amino acid runs in eukaryotic proteomes and disease associations. *Proc. Natl. Acad. Sci. U. S. A.* **99,** 333–8 (2002).

35.   Alexandre, C. M. *et al.* Regulatory DNA in A. thaliana can tolerate high levels of sequence divergence. *bioRxiv* 104323 (2017). doi:10.1101/104323

36.   Breuer, C. *et al.* BIN4, a Novel Component of the Plant DNA Topoisomerase VI Complex, Is Required for Endoreduplication in Arabidopsis. *Plant Cell* **19,** 3655–3668 (2007).

37.   Xu, G., Ma, H., Nei, M. & Kong, H. Evolution of F-box genes in plants: Different modes of sequence divergence and their relationships with functional diversification. *Proc. Natl. Acad. Sci.* **106,** 835–840 (2009).

38.   Clark, R. M. *et al.* Common Sequence Polymorphisms Shaping Genetic Diversity in Arabidopsis thaliana. *Science* **317,** 338–342 (2007).

39.   Shen, X. *et al.* Natural CMT2 Variation Is Associated With Genome-Wide Methylation Changes and Temperature Seasonality. *PLOS Genet.* **10,** e1004842 (2014).

40.   Sawaya, S. *et al.* Microsatellite tandem repeats are abundant in human promoters and are

associated with regulatory elements. *PloS One* **8,** e54710 (2013).

41.   Vinces, M. D., Legendre, M., Caldara, M., Hagihara, M. & Verstrepen, K. J. Unstable

tandem repeats in promoters confer transcriptional evolvability. *Science* **324,** 1213–6 (2009).

42.   Payseur, B. A., Place, M. & Weber, J. L. Linkage disequilibrium between STRPs and

SNPs across the human genome. *Am. J. Hum. Genet.* **82,** 1039–50 (2008).

43.   Willems, T. F., Gymrek, M., Highnam, G., Mittelman, D. & Erlich, Y. The landscape of

human STR variation. *Genome Res.* gr.177774.114– (2014). doi:10.1101/gr.177774.114

44.   Sawaya, S., Jones, M. & Keller, M. *Linkage disequilibrium between single nucleotide*

polymorphisms and hypermutable loci. 20909 (Cold Spring Harbor Labs Journals, 2015).

45.   Atwell, S. *et al.* Genome-wide association study of 107 phenotypes in Arabidopsis

thaliana inbred lines. *Nature* **465,** 627–31 (2010).

46.   Kawakatsu, T. *et al.* Epigenomic Diversity in a Global Collection of Arabidopsis thaliana

Accessions. *Cell* **166,** 492–505 (2016).

47.   Antignani, V. *et al.* Recruitment of PLANT U-BOX13 and the PI4Kβ1/β2

Phosphatidylinositol-4 Kinases by the Small GTPase RabA4B Plays Important Roles during

Salicylic Acid-Mediated Plant Defense Signaling in Arabidopsis. *Plant Cell* **27,** 243–261

(2015).

48.   Tian, D., Araki, H., Stahl, E., Bergelson, J. & Kreitman, M. Signature of balancing

selection in Arabidopsis. *Proc. Natl. Acad. Sci.* **99,** 11525–11530 (2002).

49.   Karasov, T. L. *et al.* The long-term maintenance of a resistance polymorphism through

diffuse interactions. *Nature* **512,** 436–440 (2014).

50.   Lynch, M. The rate of polygenic mutation. *Genet. Res.* **51,** 137–148 (1988).

51.   Rival, P. *et al.* The conserved PFT1 tandem repeat is crucial for proper flowering in

Arabidopsis thaliana. *Genetics* **198,** 747–754 (2014).

52.   Undurraga, S. F. *et al.* Background-dependent effects of polyglutamine variation in the

Arabidopsis thaliana gene ELF3. *Proc. Natl. Acad. Sci. U. S. A.* **109,** 19363–19367 (2012).

53.   Press, M. O. & Queitsch, C. Variability in a Short Tandem Repeat Mediates Complex

Epistatic Interactions in Arabidopsis thaliana. *Genetics* genetics.116.193359 (2016).

doi:10.1534/genetics.116.193359

54.   Cummings, B. B. *et al.* Improving genetic diagnosis in Mendelian disease with

transcriptome sequencing. *Sci. Transl. Med.* **9,** eaal5209 (2017).

55.   Nellore, A. *et al.* Human splicing diversity and the extent of unannotated splice junctions

across human RNA-seq samples on the Sequence Read Archive. *Genome Biol.* **17,** 266

(2016).

56.   Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids*

Res. **27,** 573–580 (1999).

57.   Boyle, E. A., O'Roak, B. J., Martin, B. K., Kumar, A. & Shendure, J. MIPgen: optimized

modeling and design of molecular inversion probes for targeted resequencing. *Bioinformatics*

30, 2670–2672 (2014).

58.   Gan, X. *et al.* Multiple reference genomes and transcriptomes for Arabidopsis thaliana.

Nature **477,** 419–23 (2011).

59.   Hiatt, J. B., Pritchard, C. C., Salipante, S. J., O'Roak, B. J. & Shendure, J. Single

molecule molecular inversion probes for targeted, high-accuracy detection of low-frequency

variation. *Genome Res.* **23,** 843–54 (2013).

60.   Cheng, C.-Y. *et al.* Araport11: a complete reannotation of the Arabidopsis thaliana

reference genome. *Plant J. Cell Mol. Biol.* **89,** 789–804 (2017).

61.   Sullivan, A. M. *et al.* Mapping and Dynamics of Regulatory DNA and Transcription

Factor Networks in A. thaliana. *Cell Rep.* **8,** 2015–30 (2014).

62.   R Core Team. *R: A Language and Environment for Statistical Computing.* (R Foundation

for Statistical Computing, 2016).

63.   Alonso, J. M. *et al.* Genome-wide insertional mutagenesis of Arabidopsis thaliana.

Science **301,** 653–7 (2003).

64.   Zaykin, D. V., Pudovkin, A. & Weir, B. S. Correlation-Based Inference for Linkage

Disequilibrium With Multiple Alleles. *Genetics* **180,** 533–545 (2008).

65.   Horton, M. W. *et al.* Genome-wide patterns of genetic variation in worldwide

Arabidopsis thaliana accessions from the RegMap panel. *Nat. Genet.* **44,** 212–216 (2012).

66.   Karatzoglou, A., Smola, A. & Hornik, K. *kernlab: Kernel-Based Machine Learning Lab*.

(2016).

67.   Shabalin, A. A. Matrix eQTL: ultra fast eQTL analysis via large matrix operations.

Bioinformatics **28,** 1353–1358 (2012).

68.   Therneau, T. M. *coxme: Mixed Effects Cox Models*. (2015).

69.   Hurvich, C. M. & Tsai, C.-L. Regression and time series model selection in small

samples. *Biometrika* **76,** 297–307 (1989).