1

Using connectivity to identify climatic drivers of local adaptation

Stewart L. Macdonald*[1,2], John Llewelyn[1,2], & Ben L. Phillips[3,1]

1. Centre for Tropical Biodiversity and Climate Change, James Cook University, Townsville, Qld, 4811, Australia. stewart.macdonald@my.jcu.edu.au
2. CSIRO Land and Water Flagship, Townsville, Qld, Australia
3. School of Biosciences, University of Melbourne, Vic, Australia. phillipsb@unimelb.edu.au
* Author and address for correspondence

Running title: Identifying drivers of local adaptation

Statement of authorship: BLP and SLM designed research; SLM, JL, and BLP collected data; SLM & BLP performed analyses and wrote the manuscript; all authors contributed to revisions.

Keywords: local adaptation, climate, connectivity, gene flow.

Abstract

Despite being able to conclusively demonstrate local adaptation, we are still often

unable to objectively determine the climatic drivers of local adaptation. Given the

rapid rate of global change, understanding the climatic drivers of local adaptation is

vital. Not only will this tell us which climate axes matter most to population fitness,

but such knowledge is critical to inform management strategies such as translocation

and targeted gene flow. Targeted gene flow, for example, requires knowledge of

where in the landscape we can find populations with pre-adapted climate-relevant

traits; we cannot find these populations if we do not know the dominant climatic

drivers of local adaptation. While simple assessments of geographic trait variation are

useful, geographic variation — and its associations with environment — may

represent plastic, rather than evolved differences. Additionally, the vast number of

environment–trait combinations makes it difficult to determine which aspects of the

environment populations adapt to. Here we argue that by incorporating a measure of

landscape connectivity as a proxy for gene flow, we can differentiate between

environment–trait relationships that are under selection versus those that reflect

phenotypic plasticity. By doing so, we can rapidly shorten the list of environment–

trait combinations that may be of adaptive significance. We demonstrate this method

using data on geographic trait variation in a lizard species from Australia's Wet

Tropics rainforest.

Significance Statement

Despite 150 years of evolutionary research, we still do not have a good method for

determining the climatic drivers of local adaptation. Having such a method is critical

if we are to understand and mitigate the impact of climate change. Here we exploit the

fact that local adaptation is eroded by gene flow to develop a new method for

identifying the environmental drivers of local adaptation. Our method will allow

workers around the world, for the first time (and often using existing datasets), to

determine the climatic drivers of local adaptation in their system. As such, the method

has powerful implications not only for pure evolutionary research, but also for

management.

Introduction

It is only recently that we have begun to appreciate the speed with which evolution can happen; not only over relatively short timespans (e.g., 1, 2-4), but also at small spatial scales (5). Rapid local adaptation has been recorded in response to a wide suite of environmental drivers, including invasive species, and pollution (6). We expect climate to also be a major driver of local adaptation (e.g., 7, 8), and understanding the way in which species respond to climate is of increasing importance because anthropogenic climate change is proceeding at such a rate that there are concerns that many species will be unable to evolve rapidly enough to avoid extinction (9, 10).

Evolution typically optimizes phenotypes, but the optimum will vary through both time and space (11, 12), in turn leading to populations ('demes') that have, on average, higher fitness in their home environment than an immigrant would: local adaptation. While adaptive optima for traits almost always vary geographically, it does not follow that all geographic trait variation is due to local adaptation. Geographic trait variation can arise due to other factors, such as phenotypic plasticity (including developmental plasticity and maternal effects) and environmental factors (such as geographic variation in fitness-reducing parasites). These factors, and particularly phenotypic plasticity, can give the appearance of local adaptation, despite having no underlying heritable basis (10, 11), complicating our identification of climate-relevant adaptive variation.

To circumvent these issues, evolutionary biologists use experimental approaches to demonstrate local adaptation (12, 13). Experiments designed to detect local adaptation typically utilise one of two techniques: 1) reciprocal transplants, which are done *in*

situ, and are considered the gold standard for demonstrating local adaptation; or 2)

common garden experiments, which are usually done in the lab where it is easier to

control each environmental variable (12). Both of these techniques can be difficult,

for reasons of time, expense, logistics, or ethics. This difficulty increases as the

number of separate demes and environmental variables to be tested increases and as

the generation time of the organism increases (12). Additionally, although reciprocal

transplants will detect signs of local adaptation, they are not necessarily suited to

identifying the environmental drivers of that local adaptation (14). This is because *in*

situ reciprocal transplants necessarily encompass all the environmental variables that

differ between the transplant locations. Lab-based common garden approaches may,

in principle, be more suited to identifying environmental drivers (because the

environment may be under a degree of control), but in practice it often remains

impossible to identify the environmental drivers of trait variation seen in the wild.

Thus, the best experimental tools we have for studying local adaptation are

demanding in terms of time and cost, and are unsuitable for assigning environmental

drivers (such as climate variables) to adaptive variation. If we are looking for climate-

driven local adaptation, this is a problem: we want to know which climate variable or

variables are the main drivers of adaptation, and we urgently need this information for

many species.

By definition, local adaptation has a genetic basis and is consequently weakened by

gene flow (11, 15, 16). Demes with high levels of inward gene flow are therefore

likely to be less optimally adapted, causing an observable mismatch between optimal

and actual phenotypes. Some examples of this are birds dispersing and producing

clutch sizes that are not optimised for the habitat quality in which they are now

nesting (17), larval salamander colouration not matching streambed colouration due to

high levels of gene flow from nearby but predator-free streams (18), and stick insects

in smaller habitat patches having non-cryptic colouration when the surrounding

patches are larger and environmentally dissimilar (19). These observations of

"migrant load" suggest an alternative technique for identifying and assessing local

adaptation. First, we look across populations for correlations between the environment

(e.g., mean annual temperature) and traits (e.g., morphology, physiology). By

themselves, these correlations are not sufficient evidence of local adaptation — they

could also be caused by phenotypic plasticity. Second, knowing that local adaptation

is hindered by gene flow, we can look at whether gene flow diminishes the

environmental effect. In cases where data on gene flow are absent (which is often the

case), habitat connectivity can be used as a substitute for gene flow. Environment–

trait correlations that are strong, but which are also weakened by connectivity, are

indicative of environment–trait correlations that have arisen through local adaptation.

If we now collect data on a large number of environment–trait correlations, and their

interaction with connectivity, we can imagine several possible patterns emerging.

These possibilities are depicted in Figure 1. Each panel represents a possible

relationship between environment–trait correlations (environmental coefficients on

the x-axis) and the interaction between environment and connectivity (y-axis). Panel

A shows a set of environment–trait relationships that are strong, but that are not

influenced by connectivity (i.e., no environment–connectivity interaction). This

pattern is indicative of a system in which environment–trait correlations are

predominantly driven by plastic responses of traits to their environment (i.e., traits

always match the local environment, regardless of the level of inward gene flow).

Plastic responses are still of interest, and many species are likely to show such

plasticity as the climate changes (20). Panel B shows a system in which environment–

trait relationships are eroded by connectivity: increased connectivity diminishes the

correlation between the environment and the trait. In this situation, the interaction

between the environmental variable and connectivity is negative when the

environmental coefficient is positive (i.e., greater connectivity causes the

environmental coefficient to decrease towards zero; bottom-right quadrant), and

positive when the environmental coefficient is negative (i.e., greater connectivity

causes the environmental coefficient to increase towards zero; top-left quadrant). This

is the pattern we would expect if there is a genetic basis to the environment–trait

correlation, i.e., local adaptation. Panel C shows the situation where the effect of the

environment tends to be enhanced by connectivity. This pattern might arise in

organisms that are highly mobile and can actively move to their ideal environment,

thus avoiding the selective pressures that would lead to local adaptation.

Understanding how species respond to specific aspects of their environment is vital if

we are to have any hope of halting the current rapid loss of biodiversity. Climate

change is undoubtedly one of the biggest threats to global biodiversity (21, 22), and

conservation biologists are looking to a variety of techniques to assess and help

mitigate the impacts of climate change on vulnerable species (23-25). One technique

that is likely to see increasing use is targeted, or assisted gene flow [TGF; for review,

see (23, 26)]. This technique involves the spatial redistribution of long-standing

adaptations, and acts to increase genetic diversity in recipient populations, thereby

bolstering capacity for evolutionary adaptation (10, 23, 25, 26). When applying TGF

to help species adapt to climate change, we need to find an existing location that

matches the future climate at our recipient site, and then translocate animals from that

source location. It is a simple idea, but climate is multidimensional and species will

not be adapting equally to each climate axis: is a difference of 0.5°C in mean

temperature more important than a difference of 100mm in annual rainfall? The

answer depends upon which axes have the strongest influence on fitness.

Here we explore the idea of using connectivity to infer local adaptation. To do this we

develop a case study of a lizard species from northern Australia. We use this system

to examine the relationship, across sites, between traits and climatic variables. We

assess how habitat connectivity affects these relationships and use the interaction

between the environmental variable and connectivity to rank environment–trait

combinations. In doing so, we reveal a set of environment–trait relationships

dominated by local adaptation.

Methods

Study species and site selection

The Rainforest Sunskink (*Lampropholis coggeri*) is a small (snout–vent length up to

45 mm), diurnal scincid lizard restricted to the rainforests of the Wet Tropics region

of northeastern Australia (27). The rainforests of this region cover a wide range of

environmental conditions, spanning significant elevation (0–1600 m ASL),

precipitation (annual mean precipitation of 1432–8934 mm, not including input from

cloud stripping), and temperature (annual mean temperature of 16.3–25.8˚C)

gradients. This heliothermic skink is active year-round, often seen basking in patches

of sunlight on the rainforest floor. Lizards were captured by hand from sites that were

selected to maximize the environmental heterogeneity sampled (Fig. 2).

Morphological measurements were obtained from 532 skinks from 32 sites.

Physiological measurements were obtained from a smaller subset of these lizards: 259

skinks from 12 sites. Between 8–20 skinks were caught at each site on each collecting

trip. Following capture, skinks were transported to James Cook University (JCU) in

Townsville for trait measurement. All procedures involving lizards were approved by

the JCU animal ethics committee (projects A1976 and A1726).

198
Physiological trials

Physiological trials commenced within seven days of skinks being collected from the

field; skinks being used only for morphology were measured and released back at

their point of capture within seven days. The following measures were taken from

each skink (n = 259) during laboratory trials: critical thermal minimum (CTmin),

critical thermal maximum (CTmax), thermal-performance breadth for sprinting

(breadth80), maximum sprint speed (Rmax), temperature at which sprint speed is

optimized (Topt), active body temperature as measured in a thermal gradient

(Tactive), and desiccation rate (des) (see Table S1 for further details). Details of trait

measurement procedures are detailed elsewhere (see 28, 29).

<u>Morphological measurements</u>

The following measurements were taken from each skink (n = 532) using digital

calipers: head width (HeadW); head length (HeadL); interlimb length (ILimbL);

hindlimb length (HindLL). Left and right measurements were averaged to obtain one

measurement for that trait. We also recorded snout–vent length (SVL), total length,

and mass (see Table S1 for further details). All measurements were taken by one

person (SLM) to minimize observer bias. All morphological variables were log-

transformed prior to regression analyses.

<u>Climatic variables, and connectivity</u>

As our study aimed to assess adaptation to local climate, various temperature and

precipitation variables were extracted for each site (see Table S2 for details). We

considered both means and extremes. It is important to consider climatic extremes,

because temperature extremes may be increasing faster than mean temperatures (30),

and many species are likely to be particularly vulnerable to these extremes (31). Many

environmental variables are highly correlated (28), so only the less-derived variables

were used in analyses, specifically: annual mean precipitation (AMP); seasonality of

precipitation (Pcov); precipitation of the driest quarter (Pdry); annual mean

temperature (AMT); coefficient of variation of temperature (Tcov); average minimum

daily temperature (Tmin); average maximum daily temperature (Tmax); average

variance of daily maximum temperature (TmaxVar); and average variance of daily

Tmin (TminVar).

Our connectivity index was designed to capture the flux of individuals through a

location and is detailed in the Supplementary Material. Briefly, it is a measure of

habitat suitability for our focal skink species, averaged over space using a species-

specific estimate of dispersal potential. As our species is an obligate rainforest-

dweller, grid cells in the landscape that are rainforest and that are surrounded by

rainforest have high connectivity indices, while grid cells of rainforest surrounded by

non-rainforest matrix have low indices. See Table S2 for further details on all

variables, and Figure S1 for correlations between all variables.

<u>Analysis</u>

Our analysis aimed to assess: 1) the relationship, across sites, between each trait and

each environmental variable; and 2) how connectivity affected each of these

relationships (i.e., the interaction between connectivity and environment). To allow

comparison of coefficients across variables, and to make interaction effect-sizes

meaningful, all trait and environmental variables were standardized so they had a

mean of 0 and a standard deviation of 1. Linear models were fitted for each pair of

environment–trait variables, with all models including the effect of lizard body size

and sex, as well as the interaction between environment and connectivity:

$\text{trait}_i = A + B_{svl} \times \text{SVL}_i + B_{sex} \times \text{Sex}_i + B_{env} \times \text{Env} + B_{conn} \times \text{Conn} + B_{int} \times \text{Env} \times \text{Conn} + error_i$

Where:

trait_i = standardized trait value of interest for lizard i

A = intercept

B_{svl} = coefficient of SVL

SVL = lizard snout–vent length, to control for effect of body size

B_{sex} = coefficient of Sex

Sex = lizard sex (this species is sexually dimorphic in some morphological

traits)

B_{env} = coefficient of environmental variable

Env = environmental variable (e.g., annual mean temperature) at the lizard's

site

B_{conn} = coefficient of connectivity

Conn = connectivity index at the lizard's site

B_{int} = coefficient of interaction between Env and Conn

$error_i$ = deviation between expected value and trait value of lizard i

A score for ranking the strength of local adaptation was then calculated as:

270                              $$L = -B_{env} B_{int}$$

If the signs of the two coefficients (B_{env} and B_{int}) are opposite (which indicates an

environmental effect being eroded by high connectivity, i.e., evidence for local

adaptation), L will be positive. If the signs are the same (which indicates an

environmental effect being enhanced by high connectivity, a situation not consistent

with local adaptation), L will be negative. Thus, higher numbers on this scale equate

to stronger evidence for local adaptation in that environment–trait pair. This score

can, in theory, range from $-\infty$ to $+\infty$. Once many environment–trait combinations

have been assessed, the coefficients for all pairs can be plotted (see Fig. 1). As

described in the Introduction, in a system dominated by local adaptation, we expect to

see a negative relationship between B_{env} and B_{int} (Fig. 1B). All analyses were

conducted in R v3.2 (32).

Results

There was substantial variation in the effect of environment (B_{env}) and its interaction

with connectivity (B_{int}) across climate and trait variables, with B_{env} ranging from -1.64

to 1.37, and B_{int} ranging from -0.89 to 0.74 (Fig. 3). Despite this variation, a clear

pattern is evident, with most points in Figure 3 appearing in the top-left or bottom-

right quadrants: the quadrants in which the two coefficients have opposing signs, and

where we would expect points to fall if environment–trait relationships are due to

local adaptation. Across these environment–trait combinations there is a distinct

negative linear trend (slope= -0.36, p < 0.001). It is especially noteworthy that the

environment–trait pairs with the largest coefficients (i.e., those points furthest away

from the origin) are in the two quadrants indicative of local adaptation.

Overall, physiological traits showed substantially stronger environmental effects (i.e.,

larger values of B_{env}) than did morphological traits, with the largest environmental

effects being exhibited by CTmin (AMP: -1.64; Tmax: 1.31; Pdry: 1.28) and CTmax

(Pdry: 1.37). Physiological traits also showed stronger interactions between

environmental effects and connectivity, again with CTmin and CTmax showing the

largest interactions. These trends are apparent when we examine our index of local

adaptation, L. Figure 4 shows a heatmap of all environment–trait pairs, ranked via

reciprocal averaging according to the strength of their local adaptation index. The

environment–trait pairs that show the strongest signature of local adaptation appear at

the top-left in red. There is a rough divide, with most of the physiological traits on the

left and most of the morphological traits on the right. The exceptions are the

physiological traits Topt and Rmax, which appear at the far right of the figure.

The two environmental variables that produced the strongest effects (topmost rows in

Fig 4) were both precipitation related: annual mean precipitation (AMP) and

precipitation of the driest quarter (Pdry). In our system, AMP and Pdry are both

highly correlated with connectivity (see Fig. S1). This is expected, because our

connectivity index is largely a measure of where rainforest is, and the distribution of

rainforest in our study region is driven to a large degree by rainfall.

Discussion

Understanding environment–trait correlations will help us plan management

strategies, such as targeted gene flow (TGF), that can mitigate the impact of climate

change on vulnerable species. Numerous studies have looked for (and found)

environment–trait correlations (e.g., 18, 19, 33, 34-36), but the interpretation of these

associations is plagued with uncertainty: are they associations due to local adaptation,

or plasticity? By acknowledging that gene flow undermines adaptation, we can

incorporate connectivity (a proxy for the flux of genes) into our analysis, and in doing

so, separate correlations due to local adaptation from those due to plasticity.

<u>Local adaptation</u>

In the environment–trait combinations we assessed, physiological traits typically

showed a substantially stronger effect of environment than did morphological traits,

with the largest environmental effects shown in CTmax and CTmin (Figs. 3 & 4).

Physiological traits also generally showed stronger environment–connectivity

interactions, again with CTmin and CTmax showing the largest interactions. Overall,

physiological traits generally showed stronger evidence of local adaptation than did

morphological traits. This result is intuitive: we would expect an ectotherm's

physiological traits to be under strong selection from climate (37-39), but the fitness

link between morphology and climate is much less clear. Had we also included some

environmental variables that had a clearer bearing on morphology, we might have

detected stronger environment–trait relationships for morphology. For example,

skinks that occur in rockier habitats show various morphological adaptations to that

environment (40). Including a measure of rockiness in our set of environmental

variables, for example, might have allowed us to detect environmental correlations

with limb length.

Of the environmental variables used, our analysis suggests that precipitation is a very

strong driver of local adaptation, even in thermal traits that might not seem obviously

related to precipitation (e.g., CTmin, CTmax). Although this may seem a surprising

result, precipitation has been shown to directly affect growth rate, body temperature,

activity patterns, and thermoregulatory opportunities in lizards (38, 41-45). Wetter

areas also have higher thermal inertia (and so lower cyclical thermal fluctuations

(46)), and changed environmental variance in temperature potentially has a strong

influence on thermal limits (47). Additionally, Bonebrake and Mastrandrea (48) found

that changes in precipitation can significantly affect modeled fitness and performance

curves. Finally, comparative analyses also suggest that precipitation can influence

thermal traits in many species (38). Thus, although the mechanisms linking

precipitation to thermal limits are diffuse and poorly resolved, they do exist, and our

analyses suggest that precipitation is a strong driver of local adaptation at thermal

physiological traits.

Our analysis also suggests that temperature is an important driver of local adaptation

in this system, but that extremes of temperature (encapsulated in minimum and

maximum temperatures) are at least as strongly associated with local adaptation as is

mean temperature. Again, this result is intuitive (natural selection from climate is

likely stronger during extreme events than during normal daily temperatures) and

agrees with results of empirical studies (38). Finally, the environmental variables with

the weakest signals of local adaptation are Tcov (temperature seasonality), TminVar,

and TmaxVar (variance of minimum and maximum daily temperatures, respectively).

These variables represent predictable environmental variation occurring within an

individual's lifespan and so are variables to which we might expect individuals to

develop plastic responses, rather than adaptive responses (49-51).

System-wide signal of local adaptation

The clear negative linear trend displayed in Fig. 3 is precisely what we would expect

in a set of environment–trait combinations dominated by local adaptation. Migrant

load (the negative effect of the immigration of less-locally adapted individuals) scales

positively with immigration as well as with the strength of selection [see equation 5 in

Polechová, Barton and Marion (52)]. The reason for this is that, when the strength of

selection is moderately high, the environment will have a large effect on relevant

traits, and therefore any immigrants coming from differing environments will be

particularly maladapted and will therefore have a large and negative impact on the

local phenotype. Thus, we expect environment–trait combinations with strong local

adaptation to show strong effects of connectivity on the environment–trait correlation

(52).

We used long-term climatic averages and found strong evidence that local adaptation

dominates over plasticity in our system. If we had included different environmental

variables, such as the conditions the lizards had recently encountered, signals of

plasticity may have been more apparent. Clearly environmental variables that are

similar across generations should lead to local adaptation, while environmental

variables that fluctuate within generations should have a strong influence on

phenotypic plasticity

Phenotypic plasticity

The importance of accounting for phenotypic plasticity is exemplified in our dataset

by the relatively strong effect of precipitation of the driest quarter (Pdry) on the

temperature at which maximum sprint speed is achieved (Topt) and on maximum

sprint speed (Rmax) itself. On their own, these strong correlations might traditionally

be considered evidence for local adaptation. Our analysis, however, suggests that the

environmental effect is largely unaffected by connectivity, implying that variation in

these traits is due to plasticity. Other work (28) has shown little temporal variation in

Topt (within generations) despite clear geographic variation and this, together with

our results, suggests that this trait undergoes developmental plasticity, but is fixed in

adult lizards. In principle, this non-effect of connectivity could also arise due to

selection that is so strong that it maintains local adaptation despite high levels of gene

flow [i.e., immigrants are selected against so strongly that they do not contribute to

the effective population (11)]. The environment–trait relationships for Topt and Rmax

are, however, weaker than those for some other traits (e.g., CTmax and CTmin) that

show clear effects of connectivity, so extremely strong selection seems an unlikely

explanation for the pattern we see here.

Assumptions and limitations

Our analysis requires an index of population connectivity across the landscape,

something that can be calculated relatively easily for many species by using broad

scale habitat mapping datasets [e.g., vegetation mapping from DERM (53)]. Where

possible, these measures of connectivity should be calculated at a scale relevant to the

scale of dispersal of the species in question [as was ours, using dispersal rate data for

Lampropholis coggeri from Singhal and Moritz (54)]. For highly mobile species, such

as plants or insects that have wind-assisted dispersal and for which calculating

connectivity between populations may be difficult, it is possible our approach will not

work. But for many species of animals with relatively low vagility (i.e., the species

for which TGF is most needed), our approach should be generally applicable.

Our approach implicitly assumes that traits have a linear response to the environment

(at least at the environmental scale across which we are looking). In most instances,

this will be a reasonable assumption: it seems unlikely, for example, that a trait such

as desiccation resistance would be high in dry environments, low in moderately wet

environments, and then high again in very wet environments. The reason our

assumption of linear environment–trait relationships bears mentioning is that a

limitation of our technique arises when the connectivity index is strongly correlated

with one or more of the other environmental variables being used. In our system, for

example, AMP and Pdry are correlated with connectivity (Fig. S1). Where the

environment–connectivity correlation is very strong, the interaction term in our model

(Conn×Env) could be interpreted as a quadratic term for environment (i.e., Env^2). In

these cases, it is possible that a strong connectivity interaction is, in fact, pointing to a

non-linear environment–trait relationship. Thus, for environmental variables that

correlate with connectivity (and there will always be some), careful consideration

needs to be given to the possibility of a quadratic fitness function between

environment and trait. In our case, it remains possible, for example, that the strong

influence of precipitation on local adaptation in our system is spurious, and instead

reflects non-linear relationships between optimal trait values and precipitation. We

can, however, think of no obvious reason why thermal limits should respond

quadratically to precipitation, nor why desiccation rates and other physiological traits

should also do so. Thus we are inclined to accept the importance of this

environmental variable in driving local adaptation in our system.

Finally, our approach, by examining one environment–trait combination at a time,

may potentially miss relationships that only appear in multivariate analyses. For

example, if two environmental variables are negatively correlated but both have a

positive effect on a trait, it is possible that these countergradients can obscure the

univariate relationship. Similar problems are encountered when examining response

to selection over time (55), and with our approach may lead us to underestimate the

number of important environmental drivers of local adaptation. To minimize this

effect, care should be taken in future work to sample environmental spaces in such a

way as to minimize correlations between environmental variables. Such an aim can be

achieved by, for example, strategically exploiting latitudinal and altitudinal gradients.

<u>Conclusion</u>

There is increasing urgency to identify populations that will act as suitable sources for

targeted gene flow efforts in the face of climate change. To identify these populations,

we need to know which traits influence sensitivity to climate and are locally adapted.

Traditional approaches to unearthing local adaptation (reciprocal transplants and

common garden experiments) are time consuming, and often cannot attribute

adaptation to any particular environmental driver. Local adaptation is, however,

undermined by gene flow, and we should be able to use this fact to sort patterns of

local adaptation from patterns with other causes. Here we have demonstrated this

approach: using connectivity as a proxy for gene flow, and looking for its effect on

environment–trait correlations. Our analysis, using a species of lizard from Australia's

Wet Tropics rainforest, suggests the approach has merit: the results we achieve are

coherent and suggest local adaptation is the overwhelming signal in the set of

environment-trait correlations tested. As well as confirming a strong role for local

adaptation, we have effectively ranked environmental drivers of local adaptation,

finding evidence that precipitation and temperature are important environmental

variables with regard to local adaptation. Our analysis also suggests that some traits

exhibit strong plastic responses to the environment, particularly in response to

precipitation of the driest quarter and the seasonality of temperature and precipitation.

These specific results will likely apply to other species that are phylogenetically or

ecologically similar to our focal species, but the method has the potential to apply

much more broadly. Given the potential of this method to provide evidence of local

adaptation, and rapid ranking of the climatic drivers of local adaptation, assessment of

the method in a broader array of systems is warranted.

Acknowledgments

This work was funded by the Tropical Landscapes Joint Venture, the Wet Tropics

Management Authority, an Australian Postgraduate Award to SLM, and Australian

Research Council project DP130100318. All physiological data used in these analyses

were collected by Amberlee Hatcher at James Cook University. The authors are

eternally grateful for her patience and attention to detail while undertaking this task.

References

1.   Reznick DN & Ghalambor CK (2001) The population ecology of
488   contemporary adaptations: what empirical studies reveal about the conditions
489   that promote adaptive evolution. *Genetica* 112:183-198.

490   2.   Reznick DN, Shaw FH, Rodd FH, & Shaw RG (1997) Evaluation of the rate
491   of evolution in natural populations of guppies (*Poecilia reticulata*). *Science*
492   275(5308):1934–1937.

493   3.   Losos JB, Warheit KI, & Schoener TW (1997) Adaptive differentiation
494   following experimental island colonization in *Anolis* lizards. *Nature*
495   387(6628):70–73.

496   4.   Stuart YE, *et al*. (2014) Rapid evolution of a native species following invasion
497   by a congener. *Science* 346(6208):463–466.

498   5.   Richardson JL, Urban MC, Bolnick DI, & Skelly DK (2014) Microgeographic
499   adaptation and the spatial scale of evolution. *Trends Ecol Evol* 29(3):165–176.

500   6.   Stockwell CA, Hendry AP, & Kinnison MT (2003) Contemporary evolution
501   meets conservation biology. *Trends Ecol. Evol.* 18(2):94-101.

502   7.   Kelly MW, Sanford E, & Grosberg RK (2012) Limited potential for
503   adaptation to climate change in a broadly distributed marine crustacean.
504   *Proceedings of the Royal Society B-Biological Sciences* 279(1727):349–356.

505   8.   Olsson M & Uller T (2003) Thermal environment, survival and local
506   adaptation in the common frog, *Rana temporaria*. *Evol Ecol Res* 5(3):431–
507   437.

508   9.   Sinervo B, *et al*. (2010) Erosion of Lizard Diversity by Climate Change and
509   Altered Thermal Niches. *Science* 328(5980):894–899.

10. Hoffmann AA & Sgrò CM (2011) Climate change and evolutionary

adaptation. *Nature* 470(7335):479–485.

11. Kawecki TJ & Ebert D (2004) Conceptual issues in local adaptation. *Ecol Lett*

7(12):1225–1241.

12. Reznick DN & Travis J (1996) The empirical study of adaptation in natural

populations. *Adaptation*, (Academic Press, San Diego, CA), pp 243–289.

13. Blanquart F, Kaltz O, Nuismer SL, & Gandon S (2013) A practical guide to

measuring local adaptation. *Ecology Letters* 16:1195-1205.

14. Wright JW, Stanton ML, & Scherson R (2006) Local adaptation to serpentine

and non-serpentine soils in *Collinsia sparsiflora*. *Evol Ecol Res* 8(1):1–21.

15. García-Ramos G & Kirkpatrick M (1997) Genetic models of adaptation and

gene flow in peripheral populations. *Evolution* 51(1):21–28.

16. Wright S (1931) Evolution in Mendelian populations. *Genetics* 16(2):0097–

0159.

17. Dhondt AA, Adriaensen F, Matthysen E, & Kempenaers B (1990)

Nonadaptive Clutch Sizes in Tits. *Nature* 348(6303):723–725.

18. Storfer A, Cross J, Rush V, & Caruso J (1999) Adaptive coloration and gene

flow as a constraint to local adaptation in the streamside salamander,

Ambystoma barbouri. *Evolution* 53(3):889–898.

19. Sandoval CP (1994) The Effects of the Relative Geographic Scales of Gene

Flow and Selection on Morph Frequencies in the Walking-Stick *Timema*

christinae. *Evolution* 48(6):1866–1879.

20. Seebacher F, White CR, & Franklin CE (2015) Physiological plasticity

increases resilience of ectothermic animals to climate change. *Nature Climate*

Change 5(1):61–66.

21.   IPCC (2014) *Climate Change 2014: Synthesis Report*. *Contribution of*

Working Groups I, II and III to the Fifth Assessment Report of the

Intergovernmental Panel on Climate Change (IPCC, Geneva, Switzerland) p

151.

22.   Meehl GA, *et al*. (2007) Global Climate Projections. in *Climate Change 2007:*

The Physical Science Basis. *Contribution of Working Group I to the Fourth*

Assessment Report of the Intergovernmental Panel on Climate Change, eds

Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M,

& Miller HL (Cambridge University Press, Cambridge, United Kingdom and

New York, NY, USA).

23.   Aitken SN & Whitlock MC (2013) Assisted gene flow to facilitate local

adaptation to climate change. *Annu Rev Ecol Evol Syst* 44:367–388.

24.   Williams SE, Shoo LP, Isaac JL, Hoffmann AA, & Langham G (2008)

Towards an integrated framework for assessing the vulnerability of species to

climate change. *PLoS Biol* 6(12):2621–2626.

25.   Weeks AR, *et al*. (2011) Assessing the benefits and risks of translocations in

changing environments: a genetic perspective. *Evolutionary Applications*

4(6):709–725.

26.   Kelly EL & Phillips BL (2016) Targeted gene flow for conservation. *Cons*.

Biol. 30:259-267.

27.   Wilson SK & Swan G (2010) *A complete guide to reptiles of Australia* (Reed

New Holland, Sydney, N.S.W.) p 512 pp.

28.   Llewelyn J, Macdonald S, Hatcher A, Moritz C, & Phillips BL (2016)

Intraspecific variation in climate-relevant traits in a tropical rainforest skink.

Diversity and Distributions 22(10):1000-1012.

29.   Phillips BL, Llewelyn J, Hatcher A, Macdonald S, & Moritz C (2014) Do evolutionary constraints on thermal performance manifest at different organizational scales? *J. Evol. Biol.* 27:2687-2694.

30.   Easterling DR*, et al.* (2000) Climate extremes: Observations, modeling, and impacts. *Science* 289(5487):2068-2074.

31.   Parmesan C, Root TL, & Willig MR (2000) Impacts of extreme weather and climate on terrestrial biota. *B Am Meteorol Soc* 81(3):443–450.

32.   R Core Team (2015) R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, Vienna, Austria).

33.   Llewelyn J, Macdonald SL, Hatcher A, Moritz C, & Phillips BL (submitted) Intraspecific variation in climate-relevant traits in a tropical rainforest skink.

34.   Wegener JE, Gartner GEA, & Losos JB (2014) Lizard scales in an adaptive radiation: variation in scale number follows climatic and structural habitat diversity in *Anolis* lizards. *Biol J Linn Soc* 113(2):570–579.

35.   Klaczko J, Ingram T, & Losos J (2015) Genitals evolve faster than other traits in *Anolis* lizards. *Journal of Zoology* 295(1):44–48.

36.   Bogert CM (1949) Thermoregulation in Reptiles, a Factor in Evolution. *Evolution* 3(3):195–211.

37.   Hoffmann AA (2010) Physiological climatic limits in *Drosophila*: patterns and implications. *J Exp Biol* 213(6):870–880.

38.   Clusella-Trullas S, Blackburn TM, & Chown SL (2011) Climatic Predictors of Temperature Performance Curve Parameters in Ectotherms Imply Complex Responses to Climate Change. *Am Nat* 177(6):738–751.

39.   Angilletta MJ (2009) *Thermal adaptation: a theoretical and empirical*

synthesis (Oxford University Press, Oxford; New York) pp xii, 289 p. 281 col.

plate.

40.   Goodman BA, Miles DB, & Schwarzkopf L (2008) Life on the Rocks: Habitat

Use Drives Morphological and Performance Evolution in Lizards. *Ecology*

89(12):3462–3471.

41.   Huey RB & Webster TP (1976) Thermal Biology of *Anolis* Lizards in a

Complex Fauna: the *cristatellus* Group on Puerto-Rico. *Ecology* 57(5):985–

994.

42.   Crowley SR (1987) The Effect of Desiccation Upon the Preferred Body-

Temperature and Activity Level of the Lizard *Sceloporus indulatus*. *Copeia*

(1):25–32.

43.   Jones SM, Waldschmidt SR, & Potvin MA (1987) An Experimental

Manipulation of Food and Water - Growth and Time-Space Utilization of

Hatchling Lizards (*Sceloporus undulatus*). *Oecologia* 73(1):53–59.

44.   Lorenzon P, Clobert J, Oppliger A, & John-Alder H (1999) Effect of water

constraint on growth rate, activity and body temperature of yearling common

lizard (*Lacerta vivipara*). *Oecologia* 118(4):423–430.

45.   Stamps J & Tanaka S (1981) The Influence of Food and Water on Growth-

Rates in a Tropical Lizard (*Anolis aeneus*). *Ecology* 62(1):33–40.

46.   Myers VI & Heilman MD (1969) Thermal Infrared for Soil Temperature

Studies. *Photogramm Eng* 35(10):1024–1032.

47.   Martin TL & Huey RB (2008) Why "Suboptimal" is optimal: Jensen's

inequality and ectotherm thermal preferences. *Am Nat* 171(3):E102–E118.

48.   Bonebrake TC & Mastrandrea MD (2010) Tolerance adaptation and

precipitation changes complicate latitudinal patterns of climate change

impacts. *Proc Natl Acad Sci U S A* 107(28):12581–12586.

49.   Kingsolver JG & Huey RB (1998) Evolutionary analyses of morphological

and physiological plasticity in thermally variable environments. *Am Zool*

38(3):545–560.

50.   Gilchrist GW (1995) Specialists and Generalists in Changing Environments. I.

Fitness Landscapes of Thermal Sensitivity. *Am Nat* 146(2):252–270.

51.   Phillips BL*, et al*. (2016) Heat hardening in a tropical lizard: geographic

variation explained by the predictability and variance in environmental

temperatures. *Func. Ecol*. 30:1161-1168.

52.   Polechová J, Barton N, & Marion G (2009) Species' range: adaptation in space

and time. *Am Nat* 174(5):E186–E204.

53.   DERM (2011) Queensland Department of Environment and Resource

Management, regional ecosystem mapping version 7.0.

54.   Singhal S & Moritz C (2012) Strong selection against hybrids maintains a

narrow contact zone between morphologically cryptic lineages in a rainforest

lizard. *Evolution* 66(5):1474–1489.

55.   Merila J, Sheldon BC, & Kruuk LEB (2001) Explaining stasis:

microevolutionary studies in natural populations. *Genetica* 112:199-222.

56.   Jones, D.A., Wang, W. & Fawcett, R. (2009) High-quality spatial climate

data-sets for Australia. *Australian Meteorological and Oceanographic*

Journal, 58, 233–248.

57.   McMahon, J.P., Hutchinson, M.F., Nix, H.A. & Ord, K.D. (1995). *ANUCLIM*

631   *user's guide*. Centre for Resource and environmental Studies, Australian

632   National University, Canberra.

633   58.   Storlie, C.J., Phillips, B.L., VanDerWal, J.J. & Williams, S.E. (2013)

634   Improved spatial estimates of climate predict patchier species distributions.

635   *Diversity and Distributions*, **19**, 1106–1113.

636   59.   Graham, Catherine H., VanDerWal, Jeremy, Phillips, Steven J., Moritz, Craig,

637   and Williams, Stephen E. (2010). Dynamic refugia and species persistence:

638   tracking spatial shifts in habitat through time. *Ecography*, 33(6):1062-1069.

Figure Captions

Figure 1. Graphs showing the concepts illustrated by plotting a set of environment–trait coefficients (x-axis) and the corresponding environment–connectivity interaction coefficients (y-axis). A) Phenotypic plasticity is suggested when environment–trait relationships are strong, but are not influenced by connectivity. B) Local adaptation is suggested when increasing connectivity diminishes the correlation between the environment and the trait. C) The effect of the environment is enhanced by connectivity. This latter pattern might arise in organisms that are highly mobile and can actively move to their ideal environment, thus avoiding the selective pressures that would lead to local adaptation.

Figure 2. Map of the southern Australian Wet Tropics bioregion, showing the distribution of rainforest and the sampling locations.

Figure 3. Scatterplot showing the results of 99 linear models run to assess the relationship between each pair of environment–trait variables and the environment–connectivity interaction. Environment–trait coefficients are on the x-axis, and environment–connectivity interaction coefficients are on the y-axis. Local adaptation is suggested in environment–trait pairs that show a strong environmental effect that is eroded by increasing connectivity.

Figure 4. Heatmap showing the relative rankings of climate variables (rows) and morphological and physiological traits (columns). The matrix has been sorted (by reciprocal averaging) and coloured according to the strength of local adaptation, with higher values coloured red and being sorted to the top/left. See Tables S1 and S2 for explanations of the trait and environmental variables used.

L = local adaptation index: $-B_{env} B_{int}$

B_{env} = coefficient of the environmental variable.

B_{int} = coefficient of the interaction between the environment and connectivity terms.

674   **Figures**

676   **Figure 1**

678
679 **Figure 2**
680
681

Figure 3

	CTmin	CTmax	breadth80	Rmax	des	HeadW	Tactive	HindLL	HeadL	ILimbL	Topt
AMP	L = 0.7751	L = -0.6644	L = -0.2265	L = 0.1570	L = -0.4007	L = 0.0835	L = -0.0315	L = 0.0551	L = 0.0219	L = -0.0159	L = -0.1404
Pdry	L = 0.7763	L = -0.6289	L = -0.2683	L = 0.1961	L = -0.3532	L = 0.1395	L = -0.1134	L = 0.0613	L = 0.0279	L = -0.0150	L = -0.0771
Tmin	L = -0.5004	L = 0.7345	L = 0.2380	L = -0.3484	L = 0.2510	L = -0.0399	L = -0.0329	L = -0.0157	L = -0.0219	L = -0.0116	L = 0.0977
AMT	L = -0.2156	L = 0.6655	L = 0.4128	L = -0.1072	L = 0.2139	L = -0.0730	L = -0.0447	L = -0.0071	L = -0.0099	L = -0.0043	L = 0.1511
Tmax	L = -0.2173	L = 0.5316	L = 0.4846	L = 0.0206	L = 0.1275	L = -0.0864	L = -0.0813	L = -0.0020	L = -0.0041	L = 0.0002	L = 0.1438
Pcov	L = -0.5868	L = 0.2996	L = 0.0522	L = -0.2473	L = 0.1719	L = -0.1279	L = 0.0474	L = -0.0642	L = -0.0440	L = 0.0082	L = 0.1089
Tcov	L = -0.6766	L = 0.2366	L = -0.0206	L = -0.4338	L = 0.0219	L = -0.1579	L = -0.0219	L = 0.0397	L = 0.0083	L = 0.0253	L = -0.0104
TmaxVar	L = -0.7312	L = 0.3308	L = 0.0820	L = -0.4284	L = 0.0898	L = -0.0394	L = 0.0984	L = 0.0853	L = 0.0267	L = 0.0034	L = -0.2013
TminVar	L = -0.4241	L = -0.0721	L = -0.1174	L = -0.2979	L = -0.0289	L = -0.2144	L = -0.1083	L = -0.0172	L = 0.0095	L = 0.0409	L = 0.2357

688 **Figure 4**

689