Using connectivity to identify climatic drivers of local adaptation

Stewart L. Macdonald*[1,2], John Llewelyn[1,2], & Ben L. Phillips[3,1]

1. Centre for Tropical Biodiversity and Climate Change, James Cook University, Townsville, Qld, 4811, Australia. stewart.macdonald@my.jcu.edu.au
2. CSIRO Land and Water Flagship, Townsville, Qld, Australia
3. School of BioSciences, University of Melbourne, Vic, Australia. phillipsb@unimelb.edu.au
* Author and address for correspondence

Running title: Identifying drivers of local adaptation

Statement of authorship: BLP and SLM designed research; SLM, JL, and BLP collected data; SLM & BLP performed analyses and wrote the manuscript; all authors contributed to revisions.

Keywords: local adaptation, climate, connectivity, gene flow.

Revised manuscript, 24 August 2017

Abstract

Despite being able to conclusively demonstrate local adaptation, we are still often unable to objectively determine the climatic drivers of local adaptation. Given the rapid rate of global change, understanding the climatic drivers of local adaptation is vital. Not only will this tell us which climate axes matter most to population fitness, but such knowledge is critical to inform management strategies such as translocation and targeted gene flow. While simple assessments of geographic trait variation are useful, geographic variation (and its associations with environment) may represent plastic, rather than evolved, differences. Additionally, the vast number of trait–environment combinations makes it difficult to determine which aspects of the environment populations adapt to. Here we argue that by incorporating a measure of landscape connectivity as a proxy for gene flow, we can differentiate between trait–environment relationships underpinned by genetic differences versus those that reflect phenotypic plasticity. By doing so, we can rapidly shorten the list of trait–environment combinations that may be of adaptive significance. We demonstrate how this reasoning can be applied using data on geographic trait variation in a lizard species from Australia's Wet Tropics rainforest. Our analysis reveals an overwhelming signal of local adaptation for the traits and environmental variables we investigated. Our analysis also allows us to rank environmental variables by the degree to which they appear to be driving local adaptation. Although encouraging, methodological issues remain: we point to these issue in the hope that the community

57   can rapidly hone the methods we sketch here. The promise is a rapid and general

58   approach to identifying the environmental drivers of local adaptation.

Introduction

It is only recently that we have begun to appreciate the speed with which evolution

can happen; not only over relatively short timespans (e.g., 1, 2-4), but also at small

spatial scales (5). Rapid local adaptation has been recorded in response to a wide suite

of environmental drivers, including invasive species, and pollution (6). We expect

climate to also be a major driver of local adaptation (e.g., 7, 8), and understanding the

way in which species respond to climate is of increasing importance because

anthropogenic climate change is proceeding at such a rate that there are concerns that

many species will be unable to evolve rapidly enough to avoid extinction (9, 10).

Evolution typically optimizes phenotypes, but the optimum will vary through both

time and space (11, 12), in turn leading to populations ('demes') that have, on average,

higher fitness in their home environment than an immigrant would: local adaptation.

While adaptive optima for traits almost always vary geographically, it does not follow

that all geographic trait variation is due to local adaptation. Geographic trait variation

can arise due to other factors, such as phenotypic plasticity (including developmental

plasticity and maternal effects), neutral clines, and environmental factors (such as

geographic variation in fitness-reducing parasites). These factors can give the

appearance of local adaptation (10, 11), complicating our identification of climate-

relevant adaptive variation.

To circumvent these issues, evolutionary biologists use experimental approaches to

demonstrate local adaptation (12, 13). Experiments designed to detect local adaptation

typically utilise one of two techniques: 1) reciprocal transplants, which are done *in*

situ, and are considered the gold standard for demonstrating local adaptation; or 2)

common garden experiments, which are usually done in the lab where it is easier to

control each environmental variable (12). Both of these techniques can be difficult,

for reasons of time, expense, logistics, or ethics. This difficulty increases as the

number of separate demes and environmental variables to be tested increases and as

the generation time of the organism increases (12). Additionally, although reciprocal

transplants will detect signs of local adaptation, they are not necessarily suited to

identifying the environmental drivers of that local adaptation (14). This is because *in*

situ reciprocal transplants necessarily encompass all the environmental variables that

differ between the transplant locations. Lab-based common garden approaches may,

in principle, be more suited to identifying environmental drivers (because the

environment may be under a degree of control), but in practice it often remains

impossible to identify the environmental drivers of trait variation seen in the wild.

Thus, the best experimental tools we have for studying local adaptation are

demanding in terms of time and cost, and are unsuitable for assigning environmental

drivers (such as climate variables) to adaptive variation. If we are looking for climate-

driven local adaptation, this is a problem: we want to know which climate variable or

variables are the main drivers of adaptation, and we urgently need this information for

many species.

By definition, local adaptation has a genetic basis and is consequently weakened by

high levels of gene flow (11, 15, 16). Demes with excessive inward gene flow are

therefore likely to be less optimally adapted, causing an observable mismatch between

optimal and actual phenotypes. Some examples of this are birds dispersing and

producing clutch sizes that are not optimised for the habitat quality in which they are

now nesting (17), larval salamander colouration not matching streambed colouration

due to high levels of gene flow from nearby but predator-free streams (18), and stick

insects in smaller habitat patches having non-cryptic colouration when the

surrounding patches are larger and environmentally dissimilar (19). These

observations of "migrant load" suggest an alternative technique for identifying and

assessing local adaptation. First, we look across populations for relationships between

the environment (e.g., mean annual temperature) and traits (e.g., morphology,

physiology). By themselves, these relationships are not sufficient evidence of local

adaptation — they could also be caused by phenotypic plasticity. Second, knowing

that local adaptation is hindered by gene flow, we can look at whether gene flow

diminishes the environmental effect. With some caveats (discussed below), in cases

where data on gene flow are absent (which is often the case), habitat connectivity can

be used as a substitute for gene flow. Trait–environment relationships that are strong,

but which are also weakened by connectivity, are indicative of trait–environment

relationships that have a genetic basis. In a statistical model, this idea would be

represented as follows:

$$\text{Trait}_i = A + B_{env}{\times}\text{Env} + B_{conn}{\times}\text{Conn} + B_{int}{\times}\text{Env}{\times}\text{Conn} + error_i$$

Where:

trait_i = trait value for individual i

A = intercept

B_{env} = coefficient of the environmental variable

Env = environmental variable (e.g., annual mean temperature) at the

individual's site

B_{conn} = coefficient of connectivity

Conn = connectivity at the individual's site

B_{int} = coefficient of the interaction between Env and Conn

135    $error_i$ = deviation between expected value and trait value of individual i

136 Which, with slight rearrangement, can also be expressed as:

137      $Trait_i = A + (B_{env} + B_{int} \times Conn) \times Env + B_{conn} \times Conn + error_i$

138 showing that the slope of the relationship between the trait and the environment now

139 depends on the connectivity value. When the signs of B_{env} and B_{int} are in opposition,

140 then we have a situation in which the relationship between the trait and the

141 environment diminishes as connectivity increases.

143 If we now collect data on a large number of trait–environment relationships, and their

144 interaction with connectivity, we can imagine several possible patterns emerging.

145 These possibilities are depicted in Figure 1. Each panel represents a possible

146 relationship between trait–environment coefficients (along the x-axis) and the

147 interaction between environment and connectivity (y-axis). Panel A shows a set of

148 trait–environment relationships that vary in strength, but that are not influenced by

149 connectivity (i.e., no environment–connectivity interaction). This pattern is indicative

150 of a system in which trait–environment relationships are predominantly driven by

151 plastic responses of traits to their environment (i.e., traits always match the local

152 environment, regardless of the level of inward gene flow). Panel B shows a system in

153 which trait–environment relationships are eroded by connectivity: increased

154 connectivity diminishes the relationship between the environment and the trait. In this

155 situation, the interaction between the environmental variable and connectivity is

156 negative when the environmental coefficient is positive (i.e., greater connectivity

157 causes the environmental coefficient to decrease towards zero; bottom-right

158 quadrant), and positive when the environmental coefficient is negative (i.e., greater

159 connectivity causes the environmental coefficient to increase towards zero; top-left

quadrant). This is the pattern we would expect if there is a genetic basis to the trait–

environment relationship, such as is exhibited by local adaptation. Panel C shows the

situation where the effect of the environment tends to be enhanced by connectivity.

This pattern might arise in organisms that are highly mobile and can actively move to

their ideal environment, thus avoiding the selective pressures that would lead to local

adaptation.

Understanding how species respond to specific aspects of their environment is vital if

we are to have any hope of halting the current rapid loss of biodiversity. Climate

change is undoubtedly one of the biggest threats to global biodiversity (20, 21), and

conservation biologists are looking to a variety of techniques to assess and help

mitigate the impacts of climate change on vulnerable species (22-24). One technique

that is likely to see increasing use is targeted, or assisted, gene flow [TGF; for review,

see (22, 25)]. This technique involves the spatial redistribution of long-standing

adaptations, and acts to increase genetic diversity in recipient populations, thereby

bolstering capacity for evolutionary adaptation (10, 22, 24, 25). When applying TGF

to help species adapt to climate change, we need to find an existing location that

matches the future climate at our recipient site, and then translocate animals from that

source location. It is a simple idea, but climate is multidimensional and species will

not be adapting equally to each climate axis: is a difference of 0.5°C in mean

temperature more important than a difference of 100mm in annual rainfall? The

answer depends upon which aspects of climate (hereafter "climatic axes") have the

strongest influence on fitness.

Here we explore the idea of using connectivity to infer local adaptation. To do this we

develop a case study of a lizard species from northern Australia. We use this system

to examine the relationship, across sites, between traits and climatic variables. We

assess how habitat connectivity affects these relationships and use the interaction

between the environmental variable and connectivity to rank trait–environment

combinations. In doing so, we reveal a set of trait–environment relationships that

appears to be dominated by local adaptation.

Methods

<u>Study species and site selection</u>

The Rainforest Sunskink (*Lampropholis coggeri*) is a small (snout–vent length up to

45 mm), diurnal scincid lizard restricted to the rainforests of the Wet Tropics region

of northeastern Australia (26). The rainforests of this region cover a wide range of

environmental conditions, spanning significant elevation (0–1600 m ASL),

precipitation (annual mean precipitation of 1432–8934 mm, not including input from

cloud stripping), and temperature (annual mean temperature of 16.3–25.8˚C)

gradients. This heliothermic skink is active year-round, often seen basking in patches

of sunlight on the rainforest floor. Lizards were captured by hand from sites that were

selected to maximize the environmental heterogeneity sampled (Fig. 2).

Morphological measurements were obtained from 532 skinks from 32 sites.

Physiological measurements were obtained from a smaller subset of these lizards: 259

skinks from 12 sites. At each site, 8–20 skinks were caught per collecting trip.

Following capture, skinks were transported to James Cook University (JCU) in

Townsville for trait measurement. All procedures involving lizards were approved by

the JCU animal ethics committee (projects A1976 and A1726).

<u>Physiological trials</u>

Physiological trials commenced within seven days of skinks being collected from the

field; skinks being used only for morphology were measured and released back at

their point of capture within seven days. The following measures were taken from

each skink (n = 259) during laboratory trials: critical thermal minimum (CTmin),

critical thermal maximum (CTmax), thermal-performance breadth for sprinting

(breadth80), maximum sprint speed (Rmax), temperature at which sprint speed is

optimized (Topt), active body temperature as measured in a thermal gradient

(Tactive), and desiccation rate (des) (see Table S1 for further details). Details of trait

measurement procedures are detailed elsewhere (see 27, 28).

Morphological measurements

The following measurements were taken from each skink (n = 532) using digital

calipers: head width (HeadW); head length (HeadL); interlimb length (ILimbL);

hindlimb length (HindLL). Left and right measurements were averaged to obtain one

measurement for that trait. We also recorded snout–vent length (SVL), total length,

and mass (see Table S1 for further details). All measurements were taken by one

person (SLM) to minimize observer bias. All morphological variables were log-

transformed prior to regression analyses.

Climatic variables, and connectivity

Because our study aimed to assess adaptation to local climate, various temperature

and precipitation variables were extracted for each site (see Table S2 for details). We

considered both means and extremes. It is important to consider climatic extremes,

because temperature extremes may be increasing faster than mean temperatures (29),

and selection may often occur during extreme weather events (30). Many

environmental variables are highly correlated (27), so only the less-derived variables

were used in analyses, specifically: annual mean precipitation (AMP); seasonality of

precipitation (Pcov); precipitation of the driest quarter (Pdry); annual mean

temperature (AMT); coefficient of variation of temperature (Tcov); average minimum

daily temperature (Tmin); average maximum daily temperature (Tmax); average

variance of daily maximum temperature (TmaxVar); and average variance of daily

Tmin (TminVar).

Our connectivity index was designed to capture the flux of individuals through a

location and is detailed in (31). Briefly, it is a measure of habitat suitability for our

focal skink species, averaged over space using a species-specific estimate of dispersal

potential. This approach is reasonable for any species exhibiting diffusive dispersal,

and similar techniques (though different spatial-weighting functions) can be used for

species exhibiting non-diffusive dispersal. As our species is an obligate rainforest-

dweller, grid cells in the landscape that are rainforest and that are surrounded by

rainforest have high connectivity indices, while grid cells of rainforest surrounded by

non-rainforest matrix have low indices. See Table S2 for further details on all

variables, and Figure S1 for correlations between all variables.

<u>Analysis</u>

Our analysis aimed to assess: 1) the relationship, across sites, between each trait and

each environmental variable; and 2) how connectivity affected each of these

relationships (i.e., the interaction between connectivity and environment). To allow

comparison of coefficients across variables, and to make interaction effect-sizes

meaningful, all trait and environmental variables were standardized so they had a

mean of 0 and a standard deviation of 1. Linear models were fitted for each pair of

environment–trait variables, with all models including the effect of lizard body size

and sex, as well as the interaction between environment and connectivity:

$\text{trait}_i = A + B_{svl} \times \text{SVL}_i + B_{sex} \times \text{Sex}_i + B_{env} \times \text{Env} + B_{conn} \times \text{Conn} + B_{int} \times \text{Env} \times \text{Conn} + error_i$

Where:

trait_i = standardized trait value of interest for lizard i

A = intercept

B_{svl} = coefficient of SVL

SVL = lizard snout–vent length, to control for effect of body size

273   B_{sex} = coefficient of Sex

274   Sex = lizard sex (this species is sexually dimorphic in some morphological

275   traits)

276   B_{env} = coefficient of environmental variable

277   Env = environmental variable (e.g., annual mean temperature) at the lizard's

278   site

279   B_{conn} = coefficient of connectivity

280   Conn = connectivity index at the lizard's site

281   B_{int} = coefficient of interaction between Env and Conn

282   $error_i$ = deviation between expected value and trait value of lizard i

A score for ranking the strength of local adaptation (L) was then calculated as:

$$L = -B_{env} \times B_{int}$$

If the signs of the two coefficients (B_{env} and B_{int}) are opposite (which indicates an

trait–environmental relationship that is diminished by increasing connectivity, i.e.,

evidence for local adaptation), L will be positive. If the signs are the same (which

indicates an environmental effect being enhanced by increased connectivity, a

situation not consistent with local adaptation), L will be negative. Thus, higher

numbers on this scale equate to stronger evidence for local adaptation in that

environment–trait pair. This score can, in theory, range from $-\infty$ to $+\infty$. Once many

environment–trait combinations have been assessed, the coefficients for all pairs can

be plotted (see Fig. 1). As described in the Introduction, in a system dominated by

local adaptation, we expect to see a negative relationship between B_{env} and B_{int} (Fig.

1B). All analyses were conducted in R v3.2 (32).

Results

There was substantial variation in the effect of environment (B_{env}) and its interaction

with connectivity (B_{int}) across climate and trait variables, with B_{env} ranging from -1.8

to 1.61, and B_{int} ranging from -0.73 to 0.78 (Fig. 3). Despite this variation, a clear

pattern is evident, with most points in Figure 3 appearing in the top-left or bottom-

right quadrants: the quadrants in which the two coefficients have opposing signs, and

where we would expect points to fall if trait–environment relationships have a genetic

basis. Across these trait–environment combinations there is a distinct negative linear

trend (slope= -0.36, p < 0.001). It is especially noteworthy that the trait–environment

pairs with the largest coefficients are in the two quadrants indicative of local

adaptation.

Overall, physiological traits showed substantially stronger environmental effects (i.e.,

larger values of B_{env}) than did morphological traits, with the largest environmental

effects being exhibited by CTmin (AMP: -1.80; Tmax: 1.35; Pdry: -1.55) and CTmax

(Pdry: 1.61; AMP: 1.21). Physiological traits also showed stronger interactions

between environmental effects and connectivity, again with CTmin and CTmax

showing the largest interactions. These trends are apparent when we examine our

index of local adaptation, L. Figure 4 shows a heatmap of all trait–environment pairs,

ranked via reciprocal averaging according to the strength of their local adaptation

index. The trait–environment pairs that show the strongest signature of local

adaptation appear at the top-left in red. There is a rough divide, with most of the

physiological traits on the left and most of the morphological traits on the right. The

exceptions are the physiological traits Topt and Rmax, which appear at the far right of

the figure.

The two environmental variables that produced the strongest effects (topmost rows in

Fig 4) were both precipitation related: annual mean precipitation (AMP) and

precipitation of the driest quarter (Pdry). In our system, AMP and Pdry are both

highly correlated with connectivity (see Fig. S1). This is expected, because our

connectivity index is largely a measure of where rainforest is, and the distribution of

rainforest in our study region is driven to a large degree by rainfall.

330
331
Discussion

Understanding relationships between traits and the environment will help us plan

management strategies, such as targeted gene flow (TGF), that can mitigate the

impact of climate change on vulnerable species. Numerous studies have looked for

(and found) trait–environment relationships (e.g., 18, 19, 33, 34-36), but the

interpretation of these associations is plagued with uncertainty: are they associations

due to local adaptation, neutral clines, habitat choice, or plasticity? By acknowledging

that gene flow undermines adaptation, we can incorporate connectivity (a proxy for

the flux of genes) into our analysis, and in doing so, separate those relationships due

to fixed genetic differences, from those due to plasticity or habitat choice.

Local adaptation

In the trait–environment combinations we assessed, physiological traits typically

showed a substantially stronger effect of environment (B_{env}) than did morphological

traits, with the largest environmental effects shown in CTmax and CTmin (Figs. 3 &

4). Physiological traits also generally showed stronger environment–connectivity

interactions (B_{int}), again with CTmin and CTmax showing the largest interactions.

Overall, physiological traits generally showed stronger evidence of local adaptation

than did morphological traits. This result is intuitive: we would expect an ectotherm's

physiological traits to be under strong selection from climate (37-39), but the fitness

link between morphology and climate is much less clear. Had we also included some

environmental variables that had a clearer bearing on morphology, we might have

detected stronger trait–environment relationships for morphology. For example,

skinks that occur in rockier habitats show various morphological adaptations to that

environment (40). Including a measure of rockiness in our set of environmental

variables might have allowed us to detect a signal of local adaptation for limb length.

Here, however, our focus is on climatic aspects of the environment.

Of the environmental variables used, our analysis suggests that precipitation is a very

strong driver of local adaptation, even in thermal traits that might not seem obviously

related to precipitation (e.g., CTmin, CTmax). Although this may seem a surprising

result, precipitation has been shown to directly affect growth rate, body temperature,

activity patterns, and thermoregulatory opportunities in lizards (38, 41-45). Wetter

areas also have higher thermal inertia (and so lower cyclical thermal fluctuations

(46)), and changed environmental variance in temperature potentially has a strong

influence on thermal limits (47). Additionally, Bonebrake and Mastrandrea (48) found

that changes in precipitation can significantly affect modeled fitness and performance

curves. Finally, comparative analyses also suggest that precipitation can influence

thermal traits in many species (38). Thus, although the mechanisms linking

precipitation to thermal limits are diffuse and poorly resolved, they do exist, and our

analyses suggest that precipitation is a strong driver of local adaptation at thermal

physiological traits.

Our analysis also suggests that temperature is an important driver of local adaptation

in this system, but that extremes of temperature (encapsulated in minimum and

maximum temperatures) are at least as strongly associated with local adaptation as is

mean temperature. Again, this result is intuitive (natural selection from climate is

likely stronger during extreme events than during normal daily temperatures) and

agrees with results of empirical studies (38). Finally, the environmental variables with

the weakest signals of local adaptation are Tcov (temperature seasonality), TminVar,

and TmaxVar (variance of minimum and maximum daily temperatures, respectively).

These variables represent predictable environmental variation occurring within an

individual's lifespan and so are variables to which we might expect individuals to

develop plastic responses, rather than fixed differences; local adaptation to these

variables would likely be reflected in reaction norms, rather than point values for

traits. (49-51).

<u>System-wide signal of local adaptation</u>

The clear negative linear trend displayed in Fig. 3 is precisely what we would expect

in a set of trait–environment combinations dominated by local adaptation. Migrant

load (the negative effect of the immigration of less-locally adapted individuals) scales

positively with immigration as well as with the strength of selection [see equation 5 in

Polechová, Barton and Marion (52)]. The reason for this is that, when the strength of

selection is moderately high, the environment will have a large effect on relevant

traits, and therefore any immigrants coming from differing environments will be

particularly maladapted and will therefore have a large and negative impact on the

local phenotype. Thus, we expect trait–environment combinations with strong local

adaptation to show strong effects of connectivity on the trait–environment relationship

(52).

By setting up a statistical model in which the trait–environment relationship is altered

by connectivity, we have allowed the possibility that the trait–environment

relationship could be reversed as connectivity increases. Such an outcome is absurd

from a theoretical perspective. In practice, however, our interaction coefficients were

typically estimated to be around 0.36 times as strong as the main effect of

environment. In this situation, reversal would only happen when connectivity values

were more than 2.7 standard deviations beyond the mean (a situation that is

exceedingly rare). Thus, encouragingly, our system wide analysis consistently

provides parameter estimates that are theoretically sensible, despite there being no

constraint within the model for them to be so.

We used long-term climatic averages and found strong evidence that local adaptation

dominates over plasticity in our trait–environment set. If we had included different

environmental variables, such as the conditions the lizards had recently encountered,

signals of plasticity may have been more apparent. Clearly environmental variables

that are similar across generations should lead to local adaptation, while

environmental variables that fluctuate within generations should have a strong

influence on phenotypic plasticity.

Phenotypic plasticity

The importance of accounting for phenotypic plasticity is, however, exemplified in

our dataset by the relatively strong effect of precipitation of the driest quarter (Pdry)

on the temperature at which maximum sprint speed is achieved (Topt) and on

maximum sprint speed (Rmax) itself. On their own, these strong trait–environment

relationships might be interpreted as evidence for local adaptation. Our analysis,

however, suggests that the environmental effect is largely independent of

connectivity, implying that variation in these traits is due to plasticity rather than

genetic differentiation. Other work (27) has shown little temporal variation in Topt

(within generations) despite clear geographic variation and this, together with our

results, suggests that this trait undergoes developmental plasticity, but is fixed in adult

lizards. In principle, this non-effect of connectivity could also arise due to selection

that is so strong that it maintains local adaptation despite high levels of gene flow

[i.e., immigrants are selected against so strongly that they do not contribute to the

recipient population (11)]. The trait–environment relationships for Topt and Rmax

are, however, weaker than those for some other traits (e.g., CTmax and CTmin) that

show clear effects of connectivity, so extremely strong selection seems an unlikely

explanation for the pattern we see here.

The generally weak evidence for plasticity in our dataset should not be considered

weak evidence for plasticity in these traits. Indeed many of the physiological traits we

use (e.g., CTmax) are notoriously plastic, responding reversibly on timescales ranging

from hours to months (53, 54). That we do not see signals of plasticity in these traits

here reflects our choice of environmental variables: long-term climatic variables,

rather than short-term weather variables (such as the temperature in the week before

an animal was collected). We chose these long-term variables precisely because we

are interested in unearthing patterns of local adaptation, rather than patterns due to

rapid, reversible plasticity.

<u>Caveats and challenges</u>

Our intent here has been to point out the additional inference that can be drawn from

data on geographic trait variation if we account for the effect of gene flow on trait

differentiation. The idea that local adaptation is eroded by gene flow offers a novel

way to identify the environmental drivers of local adaptation. Such a capacity is of

fundamental interest, and is also sorely needed if we are to effectively manage the

impacts of climate change. The methods we use here are, however, embryonic, and in

the following we point out caveats and challenges for future work.

Gene flow and connectivity

Our approach requires a measure of gene flow across a landscape. Here we have used

environmental connectivity as a proxy for gene flow. We chose connectivity because

it can be calculated relatively easily for many species by using broad scale habitat

mapping datasets [e.g., vegetation mapping from DERM (55)]. Of course, these

measures of connectivity should be calculated at a scale relevant to the scale of

dispersal of the species in question [as ours was, using dispersal rate data for

Lampropholis coggeri from Singhal and Moritz (56)]. While connectivity measures

will often correlate with gene flow, e.g., (57)], a measure of gene flow, rather than the

flow of individuals, would be preferable. Such measures are increasingly becoming

available with the rise of landscape genomics tools (e.g., 58), but may still be cost-

prohibitive in many cases.

While there may be better measures of gene flow, our inference might also be

improved by taking into account landscape heterogeneity in the environment. Gene

flow, per se, does not erode local adaptation. Rather it is an influx of maladapted

genes that erodes local adaptation. Thus, a better index of this "migrant load" may

well be one in which connectivity is multiplied by a measure of environmental

heterogeneity, where connectivity and heterogeneity are calculated over the same

spatial scale (e.g., 31). An index such as this should, in principle be a better measure

of migrant load than our simple measure of connectivity. The cost, however, is that

this index would need to be calculated in a standardized manner for every

environmental variable under consideration.

Clearly connectivity is an imperfect measure of migrant load. By using it, we

implicitly assume that all migrants are equally maladapted and have equal fitness in

the recipient population. Nonetheless, connectivity should scale positively with

migrant load, and our analysis using simple connectivity generated a coherent and

intuitively sensible result. This is encouraging, suggesting that, in the absence of

precise estimates of migrant load, a readily calculable connectivity metric may suffice

to elucidate broad patterns.

Linear trait–environment relationships, and covariation with connectivity

Our method assumed that traits have a linear relationship to the environment (at least

at the environmental scale across which we are looking). In many instances, this will

be a reasonable null assumption: it seems unlikely, for example, that a trait such as

desiccation resistance would be high in dry environments, low in moderately wet

environments, and then high again in very wet environments. The assumption bears

particular mention, however, in the situation where the connectivity index is strongly

correlated with one or more of the other environmental variables being used. In our

system, for example, AMP and Pdry are correlated with connectivity (Fig. S1). Where

the environment–connectivity correlation is very strong, the interaction term in our

model (Conn×Env) could be interpreted as a quadratic term for environment (i.e.,

Env^2). In these cases, it is possible that a strong connectivity interaction is, in fact,

pointing to a non-linear trait–environment relationship. Thus, for environmental

variables that correlate with connectivity (and there will always be some), careful

consideration needs to be given to the possibility of a quadratic fitness function

between trait and environment. In our case, it remains possible, for example, that the

strong influence of precipitation on local adaptation in our system is spurious, and

instead reflects non-linear relationships between optimal trait values and precipitation.

We can, however, think of no obvious reason why thermal limits should respond

quadratically to precipitation, nor why desiccation rates and other physiological traits

should also do so. Thus, in our case, we are inclined to accept the importance of this

environmental variable in driving local adaptation in our system.

Covariation between explanatory variables

As in any multiple regression analysis, our capacity to make precise coefficient

estimates diminishes if there is substantial covariation between our explanatory

variables. If a sampling regime is being designed *de novo*, care should be taken to

sample sites in such a way that covariation between environmental variables

(including connectivity) is avoided as far as possible. Such an aim can be achieved by,

for example, strategically exploiting latitudinal and altitudinal gradients.

Multivariate traits and environments

Here we examined one trait–environment combination at a time. Doing so may

potentially miss relationships that only appear in multivariate analyses. For example,

if two environmental variables are negatively correlated but both have a positive

effect on a trait, it is possible that these countergradients can obscure the univariate

relationship. Similar problems are encountered when examining response to selection

over time (59) and, with our approach, may lead us to underestimate the number of

important environmental drivers of local adaptation. Analysis incorporating multiple

environmental predictors is possible, but such a model will rapidly become saturated

with parameters. To minimize the problem of countergradients, again, care should be

taken to sample environmental spaces in such a way as to minimize correlations

between environmental variables.

An additional analytical challenge is to treat traits as multivariate. Here we have

treated each measured trait as independent. In reality, however, traits covary and this

covariance can have both genetic and environmental origins (60). As a corollary,

selection acts on the multivariate trait, and causes populations to move in multivariate

trait space (61). Consequently, local adaptation perhaps should be measured in a

multivariate trait space rather than on a univariate basis. Such an aim, however,

requires considerable theoretical development and may well require substantially

more data. For now, however, we should be aware that we are collapsing our trait

space, and each of our measured traits is not independent. For example, in our system

there is a strong correlation between CTmin and CTmax, thus we should be aware

that these two traits should not get equal weighting when we use our traits to rank

environmental variables by their importance to local adaptation.

Neutral clines

Finally, our approach should allow us to identify when geographic variation is a result

of genetic variation. That is, it can weed out relationships that are driven by plasticity

or habitat choice. Covariation between genotype and environment will often be the

result of local adaptation, but can also arise for non-adaptive reasons, the most

obvious being trait clines caused by the historical spread of population (62). In

principle, and again, with careful attention to sample design (i.e., a sample design

which minimizes the covariation between space and environment), it should be

possible to separate spatial from environmental patterns.

<u>Conclusion</u>

There is increasing urgency to identify populations that will act as suitable sources for

targeted gene flow efforts in the face of climate change. To identify these populations,

we need to know which traits influence sensitivity to climate and are locally adapted.

Traditional approaches to unearthing local adaptation (reciprocal transplants and

common garden experiments) are time consuming, and often cannot attribute

adaptation to any particular environmental driver. Local adaptation is, however,

undermined by gene flow, and we should be able to use this fact to sort patterns of

local adaptation from patterns with other causes. Here we have demonstrated this

approach: using connectivity as a proxy for gene flow, and looking for its effect on

trait–environment relationships. Our analysis, using a species of lizard from

Australia's Wet Tropics rainforest, suggests the approach has merit: the results we

achieve are coherent and suggest local adaptation is the overwhelming signal in the

set of trait–environment relationships tested. As well as implying a strong role for

local adaptation, we have effectively ranked environmental drivers of local

adaptation, finding evidence that precipitation and temperature are important

environmental variables with regard to local adaptation in our system. Our analysis

also suggests that some traits exhibit strong plastic responses to the environment,

particularly in response to precipitation of the driest quarter and the seasonality of

temperature and precipitation. These specific results will likely apply to other species

that are phylogenetically or ecologically similar to our focal species, but the method

has the potential to apply much more broadly. Analytical and sampling challenges

remain, however, and we point to avenues whereby the method can be improved.

Given the potential of this method to provide evidence of local adaptation, and to

provide rapid ranking of the climatic drivers of local adaptation, assessment of the

method in a broader array of systems is warranted.

Acknowledgments

This work was funded by the Tropical Landscapes Joint Venture, the Wet Tropics

Management Authority, an Australian Postgraduate Award to SLM, and the

Australian Research Council (DP130100318, FT160100198). All physiological data

used in these analyses were collected by Amberlee Hatcher at James Cook University.

The authors are extremely grateful for her patience and attention to detail while

undertaking this task.

References

1.   Reznick DN & Ghalambor CK (2001) The population ecology of contemporary adaptations: what empirical studies reveal about the conditions that promote adaptive evolution. *Genetica* 112:183-198.

2.   Reznick DN, Shaw FH, Rodd FH, & Shaw RG (1997) Evaluation of the rate of evolution in natural populations of guppies (*Poecilia reticulata*). *Science* 275(5308):1934–1937.

3.   Losos JB, Warheit KI, & Schoener TW (1997) Adaptive differentiation following experimental island colonization in *Anolis* lizards. *Nature* 387(6628):70–73.

4.   Stuart YE*, et al.* (2014) Rapid evolution of a native species following invasion by a congener. *Science* 346(6208):463–466.

5.   Richardson JL, Urban MC, Bolnick DI, & Skelly DK (2014) Microgeographic adaptation and the spatial scale of evolution. *Trends Ecol Evol* 29(3):165–176.

6.   Stockwell CA, Hendry AP, & Kinnison MT (2003) Contemporary evolution meets conservation biology. *Trends Ecol. Evol.* 18(2):94-101.

7.   Kelly MW, Sanford E, & Grosberg RK (2012) Limited potential for adaptation to climate change in a broadly distributed marine crustacean. *Proceedings of the Royal Society B-Biological Sciences* 279(1727):349–356.

8.   Olsson M & Uller T (2003) Thermal environment, survival and local adaptation in the common frog, *Rana temporaria*. *Evol Ecol Res* 5(3):431–437.

9.   Sinervo B*, et al.* (2010) Erosion of Lizard Diversity by Climate Change and Altered Thermal Niches. *Science* 328(5980):894–899.

616   10.   Hoffmann AA & Sgrò CM (2011) Climate change and evolutionary

617   adaptation. *Nature* 470(7335):479–485.

618   11.   Kawecki TJ & Ebert D (2004) Conceptual issues in local adaptation. *Ecol Lett*

619   7(12):1225–1241.

620   12.   Reznick DN & Travis J (1996) The empirical study of adaptation in natural

621   populations. *Adaptation*, (Academic Press, San Diego, CA), pp 243–289.

622   13.   Blanquart F, Kaltz O, Nuismer SL, & Gandon S (2013) A practical guide to

623   measuring local adaptation. *Ecology Letters* 16:1195-1205.

624   14.   Wright JW, Stanton ML, & Scherson R (2006) Local adaptation to serpentine

625   and non-serpentine soils in *Collinsia sparsiflora*. *Evol Ecol Res* 8(1):1–21.

626   15.   García-Ramos G & Kirkpatrick M (1997) Genetic models of adaptation and

627   gene flow in peripheral populations. *Evolution* 51(1):21–28.

628   16.   Wright S (1931) Evolution in Mendelian populations. *Genetics* 16(2):0097–

629   0159.

630   17.   Dhondt AA, Adriaensen F, Matthysen E, & Kempenaers B (1990)

631   Nonadaptive Clutch Sizes in Tits. *Nature* 348(6303):723–725.

632   18.   Storfer A, Cross J, Rush V, & Caruso J (1999) Adaptive coloration and gene

633   flow as a constraint to local adaptation in the streamside salamander,

634   *Ambystoma barbouri*. *Evolution* 53(3):889–898.

635   19.   Sandoval CP (1994) The Effects of the Relative Geographic Scales of Gene

636   Flow and Selection on Morph Frequencies in the Walking-Stick *Timema*

637   *christinae*. *Evolution* 48(6):1866–1879.

638   20.   IPCC (2014) *Climate Change 2014: Synthesis Report. Contribution of*

639   *Working Groups I, II and III to the Fifth Assessment Report of the*

Intergovernmental Panel on Climate Change (IPCC, Geneva, Switzerland) p

151.

21.   Meehl GA*, et al.* (2007) Global Climate Projections. in *Climate Change 2007:*

The Physical Science Basis. Contribution of Working Group I to the Fourth

Assessment Report of the Intergovernmental Panel on Climate Change, eds

Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M,

& Miller HL (Cambridge University Press, Cambridge, United Kingdom and

New York, NY, USA).

22.   Aitken SN & Whitlock MC (2013) Assisted gene flow to facilitate local

adaptation to climate change. *Annu Rev Ecol Evol Syst* 44:367–388.

23.   Williams SE, Shoo LP, Isaac JL, Hoffmann AA, & Langham G (2008)

Towards an integrated framework for assessing the vulnerability of species to

climate change. *PLoS Biol* 6(12):2621–2626.

24.   Weeks AR*, et al.* (2011) Assessing the benefits and risks of translocations in

changing environments: a genetic perspective. *Evolutionary Applications*

4(6):709–725.

25.   Kelly EL & Phillips BL (2016) Targeted gene flow for conservation. *Cons.*

Biol. 30:259-267.

26.   Wilson SK & Swan G (2010) *A complete guide to reptiles of Australia* (Reed

New Holland, Sydney, N.S.W.) p 512 pp.

27.   Llewelyn J, Macdonald S, Hatcher A, Moritz C, & Phillips BL (2016)

Intraspecific variation in climate-relevant traits in a tropical rainforest skink.

Diversity and Distributions 22(10):1000-1012.

28.   Phillips BL, Llewelyn J, Hatcher A, Macdonald S, & Moritz C (2014) Do

evolutionary constraints on thermal performance manifest at different

organizational scales? *J. Evol. Biol.* 27:2687-2694.

29.   Easterling DR*, et al.* (2000) Climate extremes: Observations, modeling, and

impacts. *Science* 289(5487):2068-2074.

30.   Parmesan C, Root TL, & Willig MR (2000) Impacts of extreme weather and

climate on terrestrial biota. *B Am Meteorol Soc* 81(3):443–450.

31.   Macdonald SL, Llewelyn J, Moritz C, & Phillips BL (2017) Peripheral isolates

as sources of adaptive diversity under climate change. *Frontiers in Ecology*

and Evolution 5.

32.   R Core Team (2015) R: A Language and Environment for Statistical

Computing (R Foundation for Statistical Computing, Vienna, Austria).

33.   Llewelyn J, Macdonald SL, Hatcher A, Moritz C, & Phillips BL (submitted)

Intraspecific variation in climate-relevant traits in a tropical rainforest skink.

34.   Wegener JE, Gartner GEA, & Losos JB (2014) Lizard scales in an adaptive

radiation: variation in scale number follows climatic and structural habitat

diversity in *Anolis* lizards. *Biol J Linn Soc* 113(2):570–579.

35.   Klaczko J, Ingram T, & Losos J (2015) Genitals evolve faster than other traits

in *Anolis* lizards. *Journal of Zoology* 295(1):44–48.

36.   Bogert CM (1949) Thermoregulation in Reptiles, a Factor in Evolution.

Evolution 3(3):195–211.

37.   Hoffmann AA (2010) Physiological climatic limits in *Drosophila*: patterns

and implications. *J Exp Biol* 213(6):870–880.

38.   Clusella-Trullas S, Blackburn TM, & Chown SL (2011) Climatic Predictors of

Temperature Performance Curve Parameters in Ectotherms Imply Complex

Responses to Climate Change. *Am Nat* 177(6):738–751.

39.   Angilletta MJ (2009) *Thermal adaptation: a theoretical and empirical*

synthesis (Oxford University Press, Oxford; New York) pp xii, 289 p. 281 col.

plate.

40.   Goodman BA, Miles DB, & Schwarzkopf L (2008) Life on the Rocks: Habitat

Use Drives Morphological and Performance Evolution in Lizards. *Ecology*

89(12):3462–3471.

41.   Huey RB & Webster TP (1976) Thermal Biology of *Anolis* Lizards in a

Complex Fauna: the *cristatellus* Group on Puerto-Rico. *Ecology* 57(5):985–

994.

42.   Crowley SR (1987) The Effect of Desiccation Upon the Preferred Body-

Temperature and Activity Level of the Lizard *Sceloporus indulatus*. *Copeia*

(1):25–32.

43.   Jones SM, Waldschmidt SR, & Potvin MA (1987) An Experimental

Manipulation of Food and Water - Growth and Time-Space Utilization of

Hatchling Lizards (*Sceloporus undulatus*). *Oecologia* 73(1):53–59.

44.   Lorenzon P, Clobert J, Oppliger A, & John-Alder H (1999) Effect of water

constraint on growth rate, activity and body temperature of yearling common

lizard (*Lacerta vivipara*). *Oecologia* 118(4):423–430.

45.   Stamps J & Tanaka S (1981) The Influence of Food and Water on Growth-

Rates in a Tropical Lizard (*Anolis aeneus*). *Ecology* 62(1):33–40.

46.   Myers VI & Heilman MD (1969) Thermal Infrared for Soil Temperature

Studies. *Photogramm Eng* 35(10):1024–1032.

47. Martin TL & Huey RB (2008) Why "Suboptimal" is optimal: Jensen's

inequality and ectotherm thermal preferences. *Am Nat* 171(3):E102–E118.

48. Bonebrake TC & Mastrandrea MD (2010) Tolerance adaptation and

precipitation changes complicate latitudinal patterns of climate change

impacts. *Proc Natl Acad Sci U S A* 107(28):12581–12586.

49. Kingsolver JG & Huey RB (1998) Evolutionary analyses of morphological

and physiological plasticity in thermally variable environments. *Am Zool*

38(3):545–560.

50. Gilchrist GW (1995) Specialists and Generalists in Changing Environments. I.

Fitness Landscapes of Thermal Sensitivity. *Am Nat* 146(2):252–270.

51. Phillips BL*, et al.* (2016) Heat hardening in a tropical lizard: geographic

variation explained by the predictability and variance in environmental

temperatures. *Func. Ecol.* 30:1161-1168.

52. Polechová J, Barton N, & Marion G (2009) Species' range: adaptation in space

and time. *Am Nat* 174(5):E186–E204.

53. Seebacher F (2005) A review of thermoregulation and physiological

performance in reptiles: what is the role of phenotypic flexibility? *Journal of*

comparative physiology. B, Biochemical, systemic, and environmental

physiology 175(7):453-461.

54. Llewelyn J, Macdonald S, Moritz C, & Phillips BL (2017) Thermoregulatory

behaviour explains countergradient variation in the upper thermal limit of a

rainforest skink. *Oikos* 126(5):748-757.

55. DERM (2011) Queensland Department of Environment and Resource

Management, regional ecosystem mapping version 7.0.

56. Singhal S & Moritz C (2012) Strong selection against hybrids maintains a

narrow contact zone between morphologically cryptic lineages in a rainforest

lizard. *Evolution* 66(5):1474–1489.

57. Palmer SCF, Coulon A, & Travis JMJ (2011) Introducing a 'stochastic

movement simulator' for estimating habitat connectivity. *Methods in Ecology*

and Evolution 2(3):258-268.

58. Petkova D, Novembre J, & Stephens M (2016) Visualizing spatial population

structure with estimated effective migration surfaces. *Nature Genetics*

48(1):94-100.

59. Merila J, Sheldon BC, & Kruuk LEB (2001) Explaining stasis:

microevolutionary studies in natural populations. *Genetica* 112:199-222.

60. Lynch M & Walsh B (1998) *Genetics and Analysis of Quantitative Traits*

(Sinauer Associates, Sunderland, MA).

61. Blows MW (2007) A tale of two matrices: multivariate approaches in

evolutionary biology. *J. Evol. Biol.* 20(1):1-8.

62. Phillips BL, Brown GP, & Shine R (2010) Life-history evolution in range-

shifting populations. *Ecology* 91(6):1617-1627.

56. Jones, D.A., Wang, W. & Fawcett, R. (2009) High-quality spatial climate

data-sets for Australia. *Australian Meteorological and Oceanographic*

Journal, 58, 233–248.

57. McMahon, J.P., Hutchinson, M.F., Nix, H.A. & Ord, K.D. (1995). *ANUCLIM*

user's guide. Centre for Resource and environmental Studies, Australian

National University, Canberra.

58.   Storlie, C.J., Phillips, B.L., VanDerWal, J.J. & Williams, S.E. (2013)

759   Improved spatial estimates of climate predict patchier species distributions.

760   *Diversity and Distributions*, **19**, 1106–1113.

761   59.   Graham, Catherine H., VanDerWal, Jeremy, Phillips, Steven J., Moritz, Craig,

762   and Williams, Stephen E. (2010). Dynamic refugia and species persistence:

763   tracking spatial shifts in habitat through time. *Ecography*, 33(6):1062-1069.

Figure Captions

Figure 1. Graphs showing the concepts illustrated by plotting a set of trait–environment

coefficients (e.g., the coefficient from a linear model examining the effect of annual mean site

temperature on the sprint performance of organisms from that site: B_{env}, x-axis) and the

corresponding environment–connectivity interaction coefficients (B_{int}, y-axis). Broad grey

line represents the approximate area in which these points would fall. A) Phenotypic plasticity

is suggested when trait–environment relationships are strong, but are not influenced by

connectivity. B) Local adaptation is suggested when increasing connectivity diminishes the

relationship between the environment and the trait. C) The effect of the environment is

enhanced by connectivity. This latter pattern might arise in organisms that are highly mobile

and can actively move to their ideal environment, thus avoiding the selective pressures that

would lead to local adaptation.

Figure 2. Map of the southern Australian Wet Tropics bioregion, showing the distribution of

rainforest in green and the sampling locations as black dots.

Figure 3. Scatterplot showing the results of 99 linear models run to assess the relationship

between each trait–environment pair, and the environment–connectivity interaction. Trait–

environment coefficients (B_{env}) are on the x-axis, and environment–connectivity interaction

coefficients (B_{int}) are on the y-axis. Local adaptation is suggested when these two parameters

are opposite in sign: in trait–environment pairs in which a strong environmental effect is

eroded by increasing connectivity.

Figure 4. Heatmap showing the relative rankings of climate variables (rows) and

morphological and physiological traits (columns). The matrix has been sorted (by reciprocal

averaging) and coloured according to the strength of local adaptation, with higher values

coloured red and being sorted to the top/left. See Tables S1 and S2 for explanations of the

trait and environmental variables used.

L = local adaptation index: $-B_{env} \times B_{int}$

802   **Figures**

804   **Figure 1**

Figure 2

Figure 3

	CTmin	CTmax	breadth80	Rmax	des	HeadW	Tactive	HindLL	HeadL	ILimbL	Topt
AMP	L = 1.3955	L = 0.8062	L = 0.1132	L = 0.1802	L = 0.1076	L = 0.0044	L = -0.0046	L = 0.0005	L = 0.0001	L = -0.0001	L = -0.0032
Pdry	L = 1.2042	L = 1.0140	L = 0.2861	L = 0.1401	L = 0.1713	L = 0.0379	L = 0.0385	L = 0.0007	L = 0.0002	L = 0.0004	L = -0.0232
Tmin	L = 0.5787	L = 0.6494	L = 0.0894	L = 0.1409	L = 0.0673	L = 0.0081	L = 0.0038	L = 0.0004	L = 0.0015	L = -0.0002	L = 0.0069
AMT	L = 0.2890	L = 0.5014	L = 0.0619	L = 0.0630	L = 0.0597	L = 0.0121	L = 0.0059	L = 0.0003	L = 0.0007	L = -0.0000	L = 0.0044
Tmax	L = 0.2932	L = 0.4480	L = 0.0281	L = -0.0148	L = 0.0467	L = 0.0121	L = 0.0089	L = 0.0001	L = 0.0003	L = -0.0000	L = 0.0029
Pcov	L = 0.2155	L = 0.2200	L = 0.0290	L = -0.0148	L = 0.0435	L = 0.0237	L = 0.0068	L = -0.0009	L = 0.0006	L = 0.0001	L = -0.0176
Tcov	L = 0.0676	L = 0.1122	L = -0.0071	L = -0.0418	L = 0.0058	L = 0.0119	L = 0.0022	L = -0.0051	L = -0.0007	L = -0.0008	L = -0.0011
TmaxVar	L = -0.0164	L = 0.1009	L = 0.0162	L = -0.0496	L = 0.0139	L = 0.0056	L = -0.0229	L = -0.0121	L = -0.0026	L = -0.0000	L = -0.0519
TminVar	L = -0.0518	L = 0.0040	L = 0.0297	L = 0.0957	L = -0.0068	L = -0.0457	L = 0.0049	L = 0.0003	L = 0.0002	L = -0.0012	L = 0.0683

816 **Figure 4**

817